Genomics of Root System Architecture Adaptation in Sorghum under Nitrogen and Phosphorus Deficiency

Joshua Magu[1], Joel Masanga[2], Catherine Muui[3], Steven Runo[1*]

[1]Department of Biochemistry, Microbiology and Biotechnology, Kenyatta University, P O Box 43844-00100, Nairobi, Kenya

[2]Department of Biology, The Pennsylvania State University, 432 Science Drive, University Park, PA 16802, USA

[3]Department of Agricultural Science and Technology, Kenyatta University, P O Box 43844-00100, Nairobi, Kenya

Corresponding author email: runo.steve@ku.ac.ke

Abstract

Plant root systems play a crucial role in water and nutrient uptake, yet the genetic mechanisms underlying root architecture adaptation to environmental stress remain poorly understood, particularly in non-model crops. Here, we investigate the diversity of root system architecture (RSA) in *Sorghum bicolor*, a climate-resilient cereal, under nitrogen (N) and phosphorus (P) deficiency. We used a diverse sorghum reference set, employed crown root imaging technology to quantify RSA traits and performed a genome-wide association study (GWAS) to identify genetic factors driving these responses. Our analysis revealed significant changes in six RSA traits—average root diameter, surface area, volume, total root length, and root length diameter classes (RLDR1 and RLDR2)—under N and P deficiency. GWAS identified SNPs associated with these traits, pinpointing three candidate genes—*ILR3-like*, *bHLH*, and a *LEUNIG* homolog—with known roles in root growth regulation. These findings provide novel genetic

insights into sorghum root adaptation to nutrient limitations and offer potential targets for

breeding climate-resilient, resource-efficient crop varieties.

Summary

Plants absorb water and nutrients from the soil through their roots, yet for most crops, little is

known about how root shape and structure adapt to stressful conditions such as poor soil

fertility. In this study, we used a globally diverse collection of sorghum genotypes to investigate

how sorghum roots respond to nitrogen and phosphorus deficiency. Remarkably, some

genotypes showed strong root adaptations under these conditions, leading us to identify the

genetic factors driving these responses. Our findings not only improve our understanding of root

adaptation to nutrient stress but also highlight promising genetic targets for breeding more

nutrient-efficient crops.

Introduction

Sorghum (*Sorghum bicolor* L. Moench) is a globally important cereal crop used for food, feed,

and bioenergy. It can survive adverse conditions such as high temperatures and drought

(Trouche et al. 2014), making it a strategic crop for mitigating the impacts of climate change.

However, sorghum production in many regions is constrained by poor soil fertility. In particular,

nitrogen (N) and phosphorus (P) are critical determinants of yield (White and Brown 2010), yet

their availability is often limited due to depleted soils and the high cost or limited accessibility of

fertilizers (Sheahan and Barrett 2017). One way to address these nutrient limitations is to exploit

plants' natural ability to adapt their root system architecture (RSA) to optimize nutrient extraction

in resource-limited environments.

Considerable research has proposed optimal RSAs to help crops cope with

environmental stress. Lynch (2013) described the "steep, cheap, and deep" ideotype as an

optimized RSA for improving water and nitrogen acquisition in maize. This ideotype emphasizes

steep root angles, which enable roots to penetrate deeper soil layers and access subsoil water

reserves critical for drought resilience. A deeper root system also captures nitrate that tends to

leach beyond shallow root zones, enhancing nitrogen uptake. The "cheap" component focuses

on reducing the metabolic costs of root growth by favoring thinner axial roots and reduced

lateral branching, thereby conserving carbon resources. For optimal phosphorus acquisition,

Lynch and Brown (2012) proposed that shallow root angles promote efficient foraging in the

topsoil, where phosphorus is most abundant due to its low mobility. Increased lateral root

branching expands the soil volume explored, allowing roots to access more phosphorus. Longer

and denser root hairs further enhance the effective root surface area, improving phosphorus

uptake, particularly under phosphorus-deficient conditions. Furthermore, the formation of more

aerenchyma – air-filled cavities in roots – relative to cortical cells reduces the metabolic cost of

root growth, allowing plants to allocate more resources to nutrient and water acquisition.

While these ideotypes outline general principles of RSA optimization, crop species and

genotypes can vary widely in how they deploy such strategies under natural resource

constraints. Maize, for example, shows considerable genotype-dependent variation in RSA

under nitrogen stress, with increased lateral root branching, a higher number of nodal roots, and

greater root length density, particularly in the topsoil where nitrogen is concentrated (Wang et al.

2022). Maize tends to allocate resources toward forming more lateral roots and denser root

systems in upper soil horizons to enhance nitrogen uptake from shallow nutrient-rich zones. In

contrast, sorghum typically produces fewer lateral roots while maintaining deeper and thicker

axial roots under nitrogen-limited conditions (Liu et al. 2024). Sorghum roots generally grow at

steeper angles to explore deeper soil layers, with only moderate increases in lateral branching

(Lynch et al. 2024), reflecting a lower degree of lateral root growth than maize. Additionally,

sorghum often maintains a higher root-to-shoot ratio under nutrient stress, prioritizing structural

root stability and resource conservation (Lopez et al. 2023).

Given the importance of these adaptations, this study aimed to identify distinct RSA

changes in sorghum under nitrogen and phosphorus deprivation, as well as the genetic regions

that underpin these traits for possible integration into breeding programs targeting nutrient-

limited ecosystems. To achieve this objective, we combined three interrelated components: (i) a

diverse sorghum panel to capture a wide range of natural genetic variation, (ii) an efficient high-

throughput phenotyping system, and (iii) high-resolution genotypic data to identify SNPs

associated with RSA.

We hypothesized that sorghum genotypes would exhibit extensive genetic variation for

RSA traits, reflecting ecological adaptations that could be harnessed for crop improvement. To

test this hypothesis, we used 123 accessions of the sorghum reference set, representing all

major cultivated, wild, and intermediate sorghum races from diverse agroecological regions

across Africa, India, the Middle East, Europe, and North America (Casa et al. 2008). This panel

has proven effective for identifying new traits, including parasitic plants (*Striga*) resistance

(Kavuluko et al. 2021).

Numerous technologies are now available for root imaging and phenotyping, including

hydroponic root elutriation systems (Box et al. 1989), minirhizotron and coring methods (Box et

al. 1989; Phillips et al. 2000), non-destructive radar and electrical resistance tomography (Ducut

et al. 2022), and root crown phenotyping (York 2018). In this study, we used a modified root

crown phenotyping approach involving excavation of the upper root system, removal of the

growth substrate (vermiculite), and subsequent measurements. The integrated RhizoVision

Crown platform (Seethepalli et al. 2020) provided a reliable way to extract data from numerous

root systems. Similar platforms have been successfully used in studies of nutrient acquisition in

crops such as soybean (*Glycine max*), common bean (*Phaseolus vulgaris*), cowpea (*Vigna

unguiculata*), wheat (*Triticum aestivum*), and maize (York 2018).

With advances in high-throughput RNA sequencing (RNA-seq), quantitative trait loci

(QTL) can now be resolved at the gene level. In our study, we used genotyping-by-sequencing

(GBS) SNP data from the sorghum diversity panel (Morris et al. 2013). This approach has been

similarly applied in other cereals, including rice, sorghum (Gelli et al. 2014), maize (Yu et al.

2015), and wheat (Oono et al. 2013), to identify genes and novel transcripts involved in RSA

control. To achieve our objective of identifying genetic factors in sorghum that influence root

architecture under nutrient-limited conditions, we controlled N and P availability in the growth

medium and analyzed the resulting RSA changes, followed by association mapping of these

traits with genotype data. We describe these findings and their implications for breeding

sorghum varieties better adapted to low-nutrient environments.

Materials and Methods

Plant material

A subset comprising 123 accessions of the sorghum reference set [Dataset S1 and

(https://genebank.icrisat.org/Common/Viewer?ctg=Referenceset&ref=RS1)] was obtained from

the International Crops Research Institute for the Semi-Arid Tropics (ICRISAT) Nairobi, Kenya

and maintained at the Plant Transformation Laboratory (Kenyatta University, Nairobi, Kenya).

Genetic description of sorghum accessions

Sorghum distribution maps were generated using MapTool package (Lemmond, 1994) version 2

in R based on metadata available at https://www.morrislab.org/data (Lasky et al. 2015) . To

perform population genetic structure analysis, we retrieved SNP data from the repository of

Colorado State University (Morris et al. 2013), available at https://www.morrislab.org/data.

Population genetic structure was analyzed using phylogenetic and Bayesian clustering

methods.

For phylogenetic analysis, we first converted the retrieved SNP HapMap file into Variant

Call Format (VCF) using TASSEL (Trait Analysis by aSSociation, Evolution and Linkage)

version 5.0 (Bradbury et al. 2007). The resulting VCF file was then utilized to construct a

neighbor-joining (NJ) tree with the 'ape' package (Paradis et al. 2004) in R.

For Bayesian clustering, the SNP HapMap file was transformed into PLINK format using

TASSEL, then ADMIXTURE software employed to analyze population clusters, evaluating K

values ranging from 1 to 10. The optimal number of clusters was determined based on the K

value with the lowest cross-validation error. The ADMIXTURE results were visualized using

plots generated with the 'ggplot2' package (Villanueva and Chen 2019) in R.

Sorghum seedling establishment

Sorghum seedlings were sowed in pots measuring 13.5 cm × 11 cm × 11 cm containing

vermiculite. For each of the 123 sorghum accessions, 5 seedlings were selected and planted in

3 replicates and 3 treatments of watering with; Long Ashton media (Hudson 1967) (optimal N

and P), Long Ashton media minus N, and Long Ashton Media minus P. Pots were laid out in a

complete randomized block and maintained in the greenhouse for 14 days. Greenhouse

conditions were set and maintained at a light/ dark cycle of 12/12 h and a temperature of 28℃

day and 24℃ nights.

Root imaging and analysis of sorghum phenotypes

On the 14th day, fresh root samples were photographed using a Nikon DX VR 18–140 mm

camera at a resolution of 1400 dpi. The acquired images were processed and analyzed using

RhizoVision Explorer,

(https://academic.oup.com/aobpla/article/13/6/plab056/6366354?utm_source=chatgpt.com) an

open-source software designed for root phenotyping.

Images were firstly scaled and cropped to produce grayscale versions, followed by the

application of a global image threshold to distinguish roots (foreground) from the background

using the software's 'Broken Roots' mode. The pixel-to-centimeter conversion was calibrated to

41.02 pixels per millimeter. Image thresholding was adjusted to achieve clear root boundaries

while minimizing external noise. Non-root artifacts smaller than 2 mm² were filtered out to

prevent interference with the analysis. Edge smoothing was enabled and set to level 2 to reduce

false root appearances during skeletonization, thereby enhancing the delineation of root

features. The root pruning threshold was configured to remove false root segments where tips

are shorter than the diameter of the primary root, ensuring accurate feature extraction.

Consequently, RhizoVision Explorer output 13 RSA traits, grouped by size (total root

length, perimeter, volume, surface area, and network area), distribution (average, maximum,

and median diameters), and extent (number of branching points, number of root tips, and

branching frequency). These parameters are detailed in Table 1 and illustrated in Figure 1. Raw

data is provided in Dataset S2.

Data analysis

Data of the above-mentioned parameters was subjected to analysis of variance (ANOVA).

Means and standard deviations of the means were generated and compared across the three

study conditions for all the genotypes. The Tukey's HSD test at a 95% confidence interval

(P<0.05) was used to separate the means and to ascertain whether there is a significant

difference in the data set. Pearson correlation coefficients, principal component analysis (PCA)

were implemented in R Software version 3.5.10. Root system architecture classification of

sorghum under the three conditions was achieved following the procedure as described by

(Bodner et al. 2013) based on PCA and computed with FactoMineR (Lê et al. 2008).

Heritability of each trait under the tree conditions was estimated according to (Hill et al., 1998).

$$h^2 = \frac{\sigma_G^2}{\sigma_G^2 + \sigma_{G \times E}^2 \frac{1}{n} + \sigma^2 \frac{1}{n \times r}}$$

Where σ_G^2 represent the genotypic variance, $\sigma_{G \times E}^2$ denotes variance of genotype by environment

interaction, σ^2 error of variance, n number of environments and r number of replicates.

Genome wide association studies (GWAS)

Genome-wide association studies (GWAS) were conducted on RSA phenotypes under three

conditions: optimal, nitrogen-deficient (-N), and phosphorus-deficient (-P). 32,955 single

nucleotide polymorphisms (SNPs) were used to perform a multivariate analysis with the

Genome Association and Prediction Integration Tool (GAPIT) Version 2 (Tang et al. 2016),

employing two models: the Fixed and Random Model Circulating Probability Unification

(FarmCPU) (Kusmec and Schnable 2018) and the Multi-Locus Mixed Model (MLMM) (Segura et

al. 2012).

Markers were filtered to retain those with a minor allele frequency (MAF) of at least 0.05.

To correct for multiple testing, we applied a 5% false discovery rate (FDR) implemented in R.

The results were visualized using quantile-quantile (QQ) plots and Manhattan plots.

Significant SNPs associated with the phenotypic traits were mapped onto the *Sorghum*

bicolor genome using Phytozome v3.1.1's JBrowse tool (Goodstein et al. 2012). Candidate

genes of interest were selected based on their proximity to the SNPs and their potential

involvement in the regulation of root development.

Results

Description of sorghum root system architecture

To extract root data from various sorghum genotypes, we used the RhizoVision platform, which

provided 13 root traits. These traits are illustrated in Figure 1 and described in more detail in

Table S1, with further illustrations in Figure 2. The traits can be grouped as follows: (i) Size:

Total root length (TRL), Perimeter (P), Surface Area (SA), Network Area (NA), and Volume (V);

(ii) Extent: Branch Points (BP), Branching Frequency (BF), and Number of Root Tips (NRT); (iii)

Distribution: Average diameter (AD), Median diameter (MD), and Maximum diameter (MD).

Figure 1. RhizoVision analysis of root system architecture showing a skeletonized sorghum root image used to extract measurements. The upper panel shows measurement for network area (left panel), root length, root tip and branching point (middle panel) and root length diameter range 1 (RLDR1) and RLDR2 (right panel). The lower panel shows a small root region that is amplified to show root diameter (RD), and root radius (RR) in the left panel, root area and root perimeter (middle panel) and root volume and root surface area (right panel). Scale of skeletonized root crown = 140 mm.

Genetic variation of sorghum root system architecture and response to N and P nutrients

deficiency

To examine genetic variation in the sorghum root system we used a subset of 123 genotypes of

the sorghum diversity panel with a global representation (Figure S1a). The panel was highly

diverse and comprised all major sorghum races. Phylogenetic analysis using Neighbor joining

grouped the genotypes into 7 clades corresponding to races: bicolor, caudatum, durra, guinea,

Guinea subraces, kafir and wild genotypes (Figure S1b). These same groupings were reflected

in Bayesian hierarchical clustering (Figure S1c) at the most optimal K (K=7). Additionally, the

sorghum genotypes represented populations originating in diverse agro-ecological regions

across Africa, Asia, America and Europe.

To determine if there were distinct differences in RSA that could be used to cluster all

123 genotypes used in the study, we computed Euclidean distances for genotypes across

treatments. Under optimum nutrient availability, hierarchical clustering (HC) grouped the

genotypes into 5 clusters that represented different root types (Figure 2). Cluster I represented

by IS19685 had the most root area while cluster V represented by IS23574 had the lowest root

area (Figure 2). Under nutrient constraint (N and P deficiency) HC grouped the genotypes into 2

clusters – representing genotypes with large (cluster I) or small (cluster II) root area (Figure S2).

The 5 clusters under optimal conditions could point to high inter-genotypic variation in root

structure while the two structures are indicative of specific inter genotypic differences of roots in

response to N and P deficiency. Overall, nutrient availability significantly reshaped the

relationships among these genotypes, causing them to cluster differently under stress

conditions.

Figure 2. Variation of sorghum root architecture: Dendrogram showing hierarchical clustering of root traits from a dataset of 123 sorghum genotypes. Clustering was performed using the complete method and Euclidean distance. The colored boxes highlight five distinct

root-type clusters, each represented with crown images corresponding to the genotypes within the cluster.

Partitioning of genotypic variation of sorghum root system architecture

Next, we examined how the variation of RSA was partitioned among the genotypes using

principal component analysis (PCA) for each treatment condition: optimal, minus N and minus

P.

Under optimal conditions (Figure 3A), PCA1 and PCA2 accounted for 88.4% of the total

variation (PCA1 = 56.9% and PCA2 = 31.5%). Variation in PCA1 was attributed to six traits: BF,

RLDR1, NRTP, NBP, P, and TRL. Five of these traits clustered together, indicating a strong

correlation between them. PCA2 comprised seven traits: RDL2, SA, NA, V, MaxD, AD, and MD.

SA and NA overlapped, suggesting a high correlation between them. Distribution-related traits

(AD, MaxD, and MD) formed a separate cluster with significant variance, indicating a lower

correlation and weaker influence relative to other traits (Figure 3A).

In N deficiency (Figure 3B), PCA1 accounted for 62.6% of the variation and PCA2

captured 28%, totaling 90.6% of the variance (Figure 3B). Under P deficiency (Figure 3C), the

total variance explained by PCA1 and PCA2 was 89.4% (Figure 3B). Both conditions exhibited

greater variance compared to the optimal N and P nutrient conditions (88.4%), suggesting

distinct changes due to nitrogen and phosphorus removal. Notably, distribution traits (AD,

MaxD, and MD) shifted to PCA3 under both N- and P-deficient treatments. This relocation

indicates a significant change in their behavior and relationship with other traits, emphasizing a

stronger negative or opposite correlation compared to their interactions under optimal N and P

conditions.

In all cases, dots representing different races did not form clusters but were widely

dispersed within the PCAs, indicating high internal diversity. However, larger dots representing

races with high influence on variance aligned with specific PCAs. Dura and wild genotypes, for

example, showed strong associations with traits represented in PCA1 and PCA2, while kafir,

guinea, and bicolor aligned were partitioned PCA3, which did not contribute to root variation.

These findings suggest that dura, caudatum, and wild sorghum genotypes exhibit strong

associations with root traits while bicolor, kafir, and guinea in PC3 and PC4 have less

association with root trait variation. This implies that some genotypes exhibit greater resilience

under nitrogen-limited conditions, making them potential candidates for breeding and further

investigation.

Figure 3. Partitioning genotypic variation in sorghum root system architecture (RSA) using biplot analysis of 123 genotypes under different treatments: (A) Under optimal conditions, variation was mainly explained by two principal components (PC1 = 23.02% (56.9), PC2 = 19% (31.5)). Distribution traits were weakly related to BF and other traits. (B) Under nitrogen-deficient conditions, variation shifted, with PCs showing a different contribution across experimental runs. (C) Under phosphorus-deficient conditions, distribution traits shifted similarly to the third quadrant. Colored dots represent sorghum races of the genotypes analyzed. Traits measured are shown on the key.

Interactions of sorghum root system architecture traits

To determine levels of correlations among various RSA traits under all experimental conditions,

we performed a Person's correlation analysis. This analysis revealed a strong positive

correlation between primary root length (P) and TRL (r = 0.99), as well as between RLDR1 and

the number BPs (r = 0.91). Additionally, AD and MD were highly correlated (r = 0.99). Size-

related traits; MaxD, AD, and MD exhibited negative correlations with most other traits,

particularly BF (Figure 4A).

Under minus N conditions (Figure 4B), we observed a strong positive correlation

between SA and NRT (r = 0.99). Positive correlations (Figure 4C) were also noted between P

and TRL, AD, and MD, as well as between traits describing extent NBP and size (P and TRL).

Conversely, a negative correlation was found between traits describing extent (BP) and

distribution (MD, AD, MaxD). These trends were consistent under phosphorus-depleted

conditions. Specifically, size traits (SA and NA), SA and V, SA and mass (MA), and AD were

positively correlated, while negative correlations were observed with distribution traits such as

AD, MaxD, BF, NBP, and number of root tips (NRT). These findings highlight the inverse

relationship between size and distribution in RSA traits.

Figure 4: Pearson's correlation analysis between root system architecture (RSA) traits under: (A) Optimal (B) Minus nitrogen and (C) Minus phosphorus. Dark blue to dark brown indicates highly positive to highly negative correlations. Traits measured are shown on the key.

Key
AD = Average Diameter
BD = Branching Frequency
BF = Branching Frequency
MaxD = Maximum diameter
MD = Median Diameter
NA = Network Area
NPB = Number of Branching Points
NRTP = Total Number of Root Tips
P = Perimeter
RLDR_1 = Root Length Diameter Range_1
RLDR_2 = Root Length Diameter Range_2
SA = Surface Area
V = Volume

To further determine the correlation of RSA traits, we used the Gaussian graphical model

described in (Bhushan et al. 2019). In this model, a set of items or variables, are depicted by

circles, and a set of lines that visualize relationships between the items or variables; where line

thickness represents the strength of the relationships between items while the absence of a line

implies no or very weak relationships. The Gaussian graphical model also allows for

examination of the conditional dependencies between studied traits. Greater values for

outdegree and betweenness indicate greater centrality in a network, suggesting that a trait has

influence on other traits. Results showed that the network area had the highest betweenness

connecting other traits under optimal nutrients, minus N and minus P conditions (Figure 5 A-C).

Results also showed that BF weakly connected with NA, MD, P and RLDR1. Expectedly, traits

linked to distribution (MaxD, AD, MD) clustered together and had weaker connection to traits

associated with size (TRL, P, NA) and extent (NBP, NRT, BF) this was also observed in roots

under N deficiency (figure B). Together, these results suggest that NA has the highest

conditional dependency, signifying its great influence on other traits.

Figure 5. Gaussian graphical model showing strength of correlation between traits: A. Optimal nutrients, B, nitrogen deficiency and C. Phosphorus deficiency. Red and green lines show negative and positive correlations, respectively.

Key
AD = Average Diameter
BD = Branching Frequency
BF = Branching Frequency
MaxD = Maximum diameter
MD = Median Diameter
NA = Network Area
NPB = Number of Branching Points
NRTP = Total Number of Root Tips
P = Perimeter
RLDR_1 = Root Length Diameter Range_1
RLDR_2 = Root Length Diameter Range_2
SA = Surface Area
V = Volume

Impact of nutrient stress on sorghum root architecture

To assess the response of RSA traits under N and P deficiency, we conducted a Multivariate

Analysis of Variance (MANOVA). The analysis revealed significant differences in six RSA traits

when conditions shifted from optimal to N and P deficiency.

Specifically, root length traits—RLDR1, RLDR2, and TRL—increased markedly under P

deficiency, while remaining unchanged under N deficiency. Conversely the RSA traits indicative

of secondary growth—AD, SA and V—decreased under both N and P deficiency, with a more

pronounced reduction observed under P deficiency (Figure 6).

These findings suggest that root elongation traits are enhanced under phosphorus

constraint, whereas root secondary growth diminish under both N and P limitations.

Figure 6: Impact of nutrient stress on sorghum root architecture
The traits that had statistical differences between treatments were total root length (TRL), root length diameter range 1 (RLDR1), and root length diameter range 2 (RLDR2), Surface area, volume and average diameter. Box plots show the distribution of each trait across the three nutrient conditions. Significant differences between treatments are indicated by asterisks (*$p < 0.05$, and ***$p < 0.001$).

Genome wide association studies to identify genetic factors underpinning root system

architecture traits in sorghum

To investigate the genetic factors contributing to the observed RSA responses under N and P

stress, we conducted GWAS on all 13 measured traits. These analyses employed both the

FarmCPU and MLMM frameworks within the GAPIT pipeline. Compared to traditional single-

locus mixed linear models (MLM), these multi-locus models offer greater statistical power and

improved resolution for dissecting complex traits, increasing the likelihood of identifying

associations that might otherwise go undetected. FarmCPU achieves this by iteratively

separating marker testing from kinship correction, treating significant markers as fixed effects

while modeling random background variation, thus reducing false positives (Liu et al. 2016).

And, MLMM extends MLM by incorporating significant markers stepwise as cofactors, allowing

simultaneous modeling of multiple loci and capturing more complex genetic architectures

(Segura et al. 2012). Application of these approaches enabled the identification of a total of 25

significant SNPs which were detected under N and P deficiency using RSA traits of V, SA,

RLDR2, and NBP (N deficiency) and RLDR2 (P deficiency).

The most prominent SNP was S1_26205071 identified 11 times using V, SA, NBP, and

RLDR2 under N deficiency and RLDL2 under P deficiency. This SNP did not fall on a gene

region, but its highly significant P value and association with multiple traits in two models

pointed to its importance in shaping the root architecture. Therefore, we characterized the

phenotypic variation related to each haplotype of the SNP change. We determined that 92 and 8

accessions harbored the C and T haplotypes, respectively, whereas the remaining 37

accessions lacked genotype information. Under all environments, the accessions carrying the T

allele showed significantly higher values of the RSA traits SA, RLDR2, and NRT than those

carrying the C allele ($p \leq 0.05$), indicating that the T haplotype had a positive effect on RSA

(Figure 5C).

Our analysis further identified several SNPs located on annotated genes with functions

and phenotypes directly linked to RSA. For instance, S7_2952840 (p = 6.96E-08) was mapped

to a gene annotated as *ILR3-like*. *ILR3* (also known as *bHLH105*) encodes a basic helix-loop-

helix (bHLH) transcription factor involved in iron and auxin signaling—two pathways known to

influence RSA (Tissot et al. 2019). Likewise, S10_11497177 (p = 3.08E-08) was found

upstream of the *bHLH18*; is associated with stress response and hormone signaling pathways,

including ethylene and gibberellins. These pathways regulate cell differentiation and patterning,

affecting traits such as root branching and primary root length (Qin et al. 2022). Additionally,

SNP 54899232 (p = 8.77E-09) was positioned in the exon of a gene annotated as a

transcription corepressor of *LEUNIG*. Prior studies have shown that the *LEUNIG_HOMOLOG*

(*LUH*) plays a critical role in root elongation. For example, *luh-1* mutants in Arabidopsis exhibit

significantly shorter roots compared to wild-type plants (Sitaraman et al. 2008). The discovery of

these root-associated genes highlights the robustness of GWAS analysis in uncovering the

genetic mechanisms driving RSA trait adaptation under N and P deficiency.

Figure 7: Genome-wide association studies (GWAS) identifying genetic factors influencing root system architecture (RSA) traits in sorghum. (A) Fixed and Random Model Circulating Probability Unification (FarmCPU) model circular Manhattan plot highlighting significant single nucleotide polymorphisms (SNPs) associated with root system architecture (RSA) traits, including volume (V), root length diameter 2 (RLD2), number of branch points (NBP), and surface area (SA). (B) Corresponding Quantile-quantile (QQ) plot comparing observed and expected p-values to assess model accuracy and false positives. (C) Boxplot illustrating the allelic effect of the most significant SNP (S1_26205071) on SA and

RLD2. Favorable alleles increase trait expression, while unfavorable alleles decrease it. (D) Representative root phenotypes comparing "C" and "T" allele genotypes under optimal, nitrogen-deficient, and phosphorus-deficient conditions. Genotypes with the "C" allele exhibited significantly higher SA and RLD2. Scale bar = 140 mm.

Table 1. Sorghum single nucleotide polymorphisms (SNPs) showing significant genome-wide
associations with root system architecture (RSA) traits (FDR correction at $\propto 0.05$). SNPs within
genes are presented with their annotations in order of chromosomal location. AF: allele
frequency; AdjP: P-value after FDR adjustment. *represent SNPs identified under nitrogen
deficiency, †represent SNPs identified under phosphorus deficiency. Letters after SNP positions
represent GWAS model used: a□=□BLINK, b□=□FarmCPU, c□=□MLMM.
354

Chr	Position	AF	AdjP	Associated Trait	Gene ID	Annotation
1	26205071[*c]	0.16	6.18E-08	Surface area	N/A	N/A
1	26205071[*c]	0.16	1.67E-07	Volume	N/A	N/A
1	26205071[*b]	0.16	4.05E-08	Surface area	N/A	N/A
1	26205071[†c]	0.16	1.89E-13	RLDL2	N/A	N/A
1	26205071[*c]	0.16	5.43E-08	Surface area	N/A	N/A
1	26205071[*c]	0.16	5.43E-08	Surface area	N/A	N/A
1	26205071[*c]	0.16	1.42E-09	RLDL2	N/A	N/A
1	26423820[*c]	0.20	1.05E-07	Volume	N/A	N/A
1	26731398[*c]	0.13	1.78E-07	Surface area	N/A	N/A
1	26731398[*c]	0.13	1.72E-08	Volume	N/A	N/A
1	26731398[*b]	0.13	1.19E-07	Surface area	N/A	N/A
1	26731398[*b]	0.13	3.70E-11	Volume	N/A	N/A
1	26731398[*c]	0.13	3.29E-08	Volume	N/A	N/A
1	27129528[*c]	0.10	2.20E-08	Volume	XM_002464586.2	CCRP 4 homolog
1	26205071[*a]	0.16	1.10E-10	Surface area	N/A	N/A
1	26423820[*a]	0.20	4.68E-08	Volume	N/A	N/A
1	26205071[*a]	0.16	1.56E-12	RLDL2	N/A	N/A
1	20102651[*b]	0.40	8.66E-11	Volume	N/A	N/A
2	63695738[†c]	0.42	2.03E-08	RLDL2	N/A	N/A
2	63695779[†b]	0.41	5.28E-09	RLDL2	N/A	N/A
3	15123886[*b]	0.10	1.97E-08	RLDL2	XP_002455526	Uncharacterised protein
3	54899232[†c]	0.08	8.77E-09	RLDL2	XP_02131268.1	*LEUNIG*
3	67879725[*a]	0.12	6.48E-11	Volume	XP_002456657.1	Cell Division Control Protein 6

4	12760823[†b]	0.28	6.82E-11	RLDL2	N/A	N/A
	5702394[*b]	0.09	8.74E-08	NBP	XP_002451690.2	Uncharacterised protein
4	58775820[*b]	0.16	3.12E-09	NBP	XP_021314639.1	*MAPK14-like*
5	17754110*b	0.04	3.38E-08	NBP	XP_002449381.2	1-aminocyclopropane-1-carboxylate oxidase 1
5	59062428[*b]	0.12	2.63E-08	Volume	N/A	N/A
6	55153514[†b]	0.08	6.48E-08	RLDL2	XP_021318054.1	E3 ubiquitin-protein ligase RNF14
6	56342501[*c]	0.03	2.55E-08	N NBP	XP_021318339.1	probable LRR receptor-like serine/threonine-protein kinase/hypothetical protein
6	56929524[*b]	0.03	4.41E-09	Volume	SORBI_3006G223101	Hypothetical protein
6	4077580[*a]	0.05	1.16E-09	NBP	XP_021318130.1	Uncharacterised protein
6	45806791[*b]	0.27	1.29E-10	Volume	XP_002446521.1	Transcription factor bHLH18
7	2952840[*c]	0.07	6.96E-08	NBP	XP_0213120712.1	ILR3
7	64170934[*b]	0.43	3.75E-13	Volume	N/A	N/A
8	15675606[*b]	0.05	3.96E-09	Volume	N/A	N/A
9	49587358[†b]	0.35	1.35E-11	NBP	N/A	N/A
9	49587358[*a]	0.35	1.44E-09	NBP	N/A	N/A
10	9879684[*b]	0.16	5.00E-09	NBP	XP_002438177.2	peroxygenase 4
10	17059423[*a]	0.35	2.99E-09	Volume	N/A	N/A
10	11497177[†b]	0.46	3.08E-08	Volume	N/A	N/A

Discussion

Our study sought to address the following questions: (i) what are the inter genotypic RSA

variation and associated adaptation to N and P deficiency? (ii) how do RSA adopt to low N and

P deficiency? and (ii) which genetic factors underpin RSA trait adaptability? The following

sections provide an in-depth discussion of these questions, exploring the implications of our

findings and their relevance to sorghum improvement.

What are the inter genotypic RSA variation and associated adaptation to N and P

deficiency? Data from neighbor joining phylogenetic analysis grouped our sample size of 123

sorghum genotypes into all representative races of sorghum. The results were further confirmed

with HBC which grouped the races into 7 clusters. Expectedly, genetic diversity of the panel was

further enhanced by the broad range from where the sorghum genotypes were sourced – owing

to agroecological adaptations.

We found that all RSA traits varied significantly among the genotypes. A HC dendrogram

grouped the genotypes into five clusters, each representing distinct root types. However, when

N and P were removed, only two clusters emerged. This reduction in the number of clusters

suggests that the differences in root architecture are primarily driven by N and P deprivation. It

also indicates that most genotypes exhibit similar responses to N and P deficiency. Genotypes

that maintain RSA traits under these conditions may possess specialized mechanisms for N and

P uptake or utilization, making them unique even under low nitrogen availability. These

genotypes are of particular interest for breeding programs focused on improving N and P use

efficiency or enhancing tolerance to nutrient stress.

PCA analysis revealed that NA, SA and V accounted for most phenotypic variations,

implying that these traits account for most of the diversity observed between the sorghum

genotypes tested. This observation was also highlighted in network analysis which showed NA

and BF to have the most significant effect. There are studies that both support and contrast

these findings depending on the species, environment, and focus of the study. (Lynch 2011)

emphasizes that root surface area and volume are critical for resource acquisition and account

for significant variation in root functionality, particularly under stress conditions like low N or P

availability. Similarly, (York et al. 2013) showed that traits like root network area and volume

explain much of the variation in resource acquisition, especially under water-limited conditions.

(Postma and Lynch 2011) showed that root length and branching density can account for more

variation in nutrient acquisition than surface area or volume, depending on the soil nutrient

distribution.

For both PCA, and correlation analysis, extent traits (MaxD, AD, and MD) on PC 2

showed a weak correlation with other traits – especially distribution traits. This negative

correlation highlights trade-offs among RSA traits as suggested by York et al., (2013). The

authors argue that thicker roots (higher root diameter) tend to be less branched or have lower

branching frequency, particularly in resource-constrained environments. This observation aligns

with the idea that plants optimize root traits for specific soil conditions—favoring fewer but

thicker roots for penetrating compacted soils or thinner, more branched roots for nutrient

exploration in less compacted soils.

How do RSA adopt to low N and P deficiency? Results showed that both N and P

deficiency led to enhanced root length and diminished secondary growth. Enhanced root growth

under P deficiency is consistent with results obtained in rice (Wissuwa 2003; Shimizu et al.

2004; Yi et al. 2005) and in maize (Mollier and Pellerin 1999); but in contrast to shorter root

growth observed in Arabidopsis grown in low P conditions. The varying effects of P deficiency

on root development may be attributed to species and genotypic variations, as Reviewed in (Niu

et al. 2013). For this study, the increased elongation of lateral results in a more branched root

system with a greater efficiency for P uptake.

Root elongation under low P was also accompanied by reduced secondary growth. This

is consistent with the notion that reduced root secondary growth decreases maintenance and

construction costs, allowing greater root elongation and soil exploration, thereby improving P

acquisition and plant growth under P stress. The afore hypothesis was advanced by (Strock et

al.2017) and verified by computer simulations and root measurements in beans.

Which genetic factors underpin RSA trait adaptability? GWAS results revealed a

significant SNP that colocalized with SA and RLDR2. Further analysis of phenotypic variation

linked to the different haplotypes of S1_26205071 (C allele vs. T allele) demonstrated the

phenotypic impact of this SNP where C Allele accessions exhibited significantly higher values of

SA and RLDR2. The haplotype analysis confirms that this SNP is not just statistically significant

but biologically relevant in shaping important root traits, making it a strong candidate for further

validation and incorporation into breeding strategies aimed at improving root system

architecture. Although no gene was associated with the SNP, this SNP could be linked to a

causative variant in the vicinity, functioning through linkage disequilibrium. Additional fine-

mapping or functional studies, such as transcriptome analysis, could reveal specific genes

affected by this SNP.

Our GWAS analysis revealed significant associations with additional SNPs located in

gene-coding regions, with three noteworthy genes identified: (i) The *ILR3-like* gene belongs to

the basic helix-loop-helix (*bHLH*) family of transcription factors, which are known to play pivotal

roles in iron homeostasis, nutrient acquisition, and root development. In *Arabidopsis thaliana*,

the *ILR3* gene regulates iron deficiency responses, influencing root architecture by modulating

root hair development and lateral root growth (Tissot et al. 2019). Given its involvement in iron

regulation, *ILR3-like* may mediate crosstalk between nitrogen signaling and other nutrients,

such as iron. Under conditions of nutrient imbalance, such as nitrogen deficiency, *ILR3-like*

might trigger structural changes in roots, enhancing branching and improving nutrient foraging

efficiency. (ii) *bHLH18*, another member of the bHLH transcription factor family, is associated

with diverse plant growth and developmental processes (Qin et al. 2022), including root

development, stress responses, and hormonal signaling pathways (e.g., auxin and gibberellin).

The association between *bHLH18* and root volume suggests that this gene may influence root

growth by regulating cell proliferation, expansion, or overall biomass accumulation in roots. (iii)

Transcriptional Repressor of *LEUNIG* (*LUH*) likely acts as a positive regulator of root elongation,

influencing cell division and expansion processes in the root meristem or elongation zone

(Sitaraman et al. 2008). This hypothesis is further supported by the identification of a SNP

linked to LUH using root length diameter range 2 (RLDR2) as a trait, suggesting its role in

promoting elongation under specific root system adaptations.

The strong associations of these genes with RSA traits suggest they may function in the

same or complementary pathways that modulate root structure and function. Further

experimental validation—such as expression analysis, mutant studies, and co-expression

network analysis—will provide critical insights into their roles and interactions, offering new

avenues for understanding and improving root system adaptations to nutrient stress.

In conclusion, our study underscores the vast genetic diversity within the sorghum gene

pool, making it a valuable resource for identifying alleles for germplasm improvement.

Integrating high-throughput root architecture phenotyping with freely accessible genomic data

enabled us to pinpoint genetic regions associated with RSA adaptation under nutrient-limiting

conditions and to infer, plausibly the biological processes involved. This knowledge may inform

crop breeding strategies designed to improve resilience to challenges associated with soil

nutrient deficiency.

Author contributions

SR and second author JM conceived and designed the study. First author JM performed the

experiments and all analysis guided by SR and CM. SR, and first author JM, wrote the

manuscript. All authors read and approved the final manuscript.

Data availability statement

The data that support the findings were derived from the following resources available in the

public domain: [https://www.morrislab.org/data and

https://genebank.icrisat.org/Common/Viewer?ctg=Referenceset&ref=RS1]. The data that

supports the findings of this study are available in the supplementary material of this article.

Dataset S1 contain the sorghum accessions metadata and Dataset S2 contain the root system

architecture (RSA). Data of sorghum accessions used in the study. Full results from the GWAS,

heritability estimates, PCA analysis, and all associated scripts used in the study have been

uploaded to the Genetics Society of America's Figshare portal.

Acknowledgements

This work was funded by the National Research Fund grant to S.R. contract number

KU/DVCR/NRF/VOL1/27 under the Multi Disciplinary Research Grants. The authors thank and

acknowledge the International Crops Research Institute for the Semi-Arid Tropics (ICRISAT)

through Dr. Eric Manyasa for availing the sorghum germplasm used in this project.

Supplemental information

The following Supporting Information is available for this article:

Figure S1. Genetic diversity of sorghum accessions used for root system architecture analysis.

Table S1: Description of root system architecture (RSA) traits extracted from RhizoVision

analysis.

Figure S2. Sorghum root clusters generated under different nutrient conditions.

Figure S3. Multivariate detection of Single Nucleotide Polymorphisms (SNPs) associated with

Root System Architecture (RSA) using multiple traits and models

Dataset S1: Sorghum accessions metadata.

Dataset S2. Root system architecture (RSA) data of sorghum accessions used in the study.

References

Bhushan N, Mohnert F, Sloot D, Jans L, Albers C, Steg L. 2019. Using a Gaussian Graphical
Model to Explore Relationships Between Items and Variables in Environmental Psychology
Research. Front Psychol. 10:1050. doi:10.3389/fpsyg.2019.01050.

Bodner G, Leitner D, Nakhforoosh A, Sobotik M, Moder K, Kaul H-P. 2013. A statistical
approach to root system classification. Front Plant Sci. 4:292. doi:10.3389/fpls.2013.00292.

Box JE, Smucker AJM, Ritchie JT. 1989. Minirhizotron Installation Techniques for Investigating
Root Responses to Drought and Oxygen Stresses. Soil Sci Soc Am J. 53(1):115–118.
doi:10.2136/sssaj1989.03615995005300010021x.

Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES. 2007. TASSEL:
software for association mapping of complex traits in diverse samples. Bioinformatics.
23(19):2633–2635. doi:10.1093/bioinformatics/btm308.

Casa AM, Pressoir G, Brown PJ, Mitchell SE, Rooney WL, Tuinstra MR, Franks CD, Kresovich
S. 2008. Community Resources and Strategies for Association Mapping in Sorghum. Crop Sci.
48(1):30–40. doi:10.2135/cropsci2007.02.0080.

Ducut JD, Alipio M, Go PJ, II RC, Vicerra RR, Bandala A, Dadios E. 2022. A Review of
Electrical Resistivity Tomography Applications in Underground Imaging and Object Detection.
Displays. 73:102208. doi:10.1016/j.displa.2022.102208.

Freschet GT, Roumet C, Comas LH, Weemstra M, Bengough AG, Rewald B, Bardgett RD,
Deyn GBD, Johnson D, Klimešová J, et al. 2021. Root traits as drivers of plant and ecosystem
functioning: current understanding, pitfalls and future research needs. N Phytol. 232(3):1123–
1158. doi:10.1111/nph.17072.

Gelli M, Duo Y, Konda AR, Zhang C, Holding D, Dweikat I. 2014. Identification of differentially
expressed genes between sorghum genotypes with contrasting nitrogen stress tolerance by
genome-wide transcriptional profiling. BMC Genom. 15(1):179. doi:10.1186/1471-2164-15-179.

Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten
U, Putnam N, et al. 2012. Phytozome: a comparative platform for green plant genomics. Nucleic
Acids Res. 40(Database issue):D1178–D1186. doi:10.1093/nar/gkr944.

Hudson JP. 1967. Sand and Water Culture Methods Used in the Study of Plant Nutrition. :104--
104.

Kavuluko J, Kibe M, Sugut I, Kibet W, Masanga J, Mutinda S, Wamalwa M, Magomere T,
Odeny D, Runo S. 2021. GWAS provides biological insights into mechanisms of the parasitic
plant (Striga) resistance in sorghum. BMC Plant Biol. 21(1):392. doi:10.1186/s12870-021-
03155-7.

Kusmec A, Schnable PS. 2018. FarmCPUpp: Efficient large-scale genomewide association
studies. Plant Direct. 2(4):e00053. doi:10.1002/pld3.53.

Lasky JR, Upadhyaya HD, Ramu P, Deshpande S, Hash CT, Bonnette J, Juenger TE, Hyma K,
Acharya C, Mitchell SE, et al. 2015. Genome-environment associations in sorghum landraces
predict adaptive traits. Sci Adv. 1(6):e1400218. doi:10.1126/sciadv.1400218.

Lê S, Josse J, Husson F. 2008. FactoMineR□: An R Package for Multivariate Analysis. J Stat
Softw. 25(1). doi:10.18637/jss.v025.i01.

Liu Chunjuan, Gu W, Li B, Feng Y, Liu Chang, Shi X, Zhou Y. 2024. Screening key sorghum
germplasms for low-nitrogen tolerance at the seedling stage and identifying from the carbon and
nitrogen metabolism. Front Plant Sci. 15:1340509. doi:10.3389/fpls.2024.1340509.

Liu X, Huang M, Fan B, Buckler ES, Zhang Z. 2016. Iterative Usage of Fixed and Random
Effect Models for Powerful and Efficient Genome-Wide Association Studies. Listgarten J, editor.
Plos Genet. 12(2):e1005767. doi:10.1371/journal.pgen.1005767.

Lopez G, Ahmadi SH, Amelung W, Athmann M, Ewert F, Gaiser T, Gocke MI, Kautz T, Postma
J, Rachmilevitch S, et al. 2023. Nutrient deficiency effects on root architecture and root-to-shoot
ratio in arable crops. Front Plant Sci. 13:1067498. doi:10.3389/fpls.2022.1067498.

Lynch JP. 2011. Root Phenes for Enhanced Soil Exploration and Phosphorus Acquisition: Tools
for Future Crops. Plant Physiol. 156(3):1041–1049. doi:10.1104/pp.111.175414.

Lynch JP. 2013. Steep, cheap and deep: an ideotype to optimize water and N acquisition by
maize root systems. Ann Bot. 112(2):347–357. doi:10.1093/aob/mcs293.

Lynch JP, Galindo-Castañeda T, Schneider HM, Sidhu JS, Rangarajan H, York LM. 2024. Root
phenotypes for improved nitrogen capture. Plant Soil. 502(1–2):31–85. doi:10.1007/s11104-
023-06301-2.

Mollier A, Pellerin S. 1999. Maize root system growth and development as influenced by
phosphorus deficiency. J Exp Bot. 50(333):487–497. doi:10.1093/jxb/50.333.487.

Morris GP, Ramu P, Deshpande SP, Hash CT, Shah T, Upadhyaya HD, Riera-Lizarazu O,
Brown PJ, Acharya CB, Mitchell SE, et al. 2013. Population genomic and genome-wide
association studies of agroclimatic traits in sorghum. Proc National Acad Sci. 110(2):453–458.
doi:10.1073/pnas.1215985110.

Niu YF, Chai RS, Jin GL, Wang H, Tang CX, Zhang YS. 2013. Responses of root architecture
development to low phosphorus availability: a review. Ann Bot. 112(2):391–408.
doi:10.1093/aob/mcs285.

Oono Y, Kobayashi F, Kawahara Y, Yazawa T, Handa H, Itoh T, Matsumoto T. 2013.
Characterisation of the wheat (triticum aestivum L.) transcriptome by de novo assembly for the
discovery of phosphate deficiency-responsive genes: gene expression in Pi-stressed wheat.
BMC Genom. 14(1):77–77. doi:10.1186/1471-2164-14-77.

Paradis E, Claude J, Strimmer K. 2004. APE: Analyses of Phylogenetics and Evolution in R
language. Bioinformatics. 20(2):289–290. doi:10.1093/bioinformatics/btg412.

Phillips DL, Johnson MG, Tingey DT, Biggart C, Nowak RS, Newsom JC. 2000. Minirhizotron
installation in sandy, rocky soils with minimal soil disturbance. Soil Sci Soc Am J. 64(2):761–
764. doi:10.2136/sssaj2000.642761x.

Postma JA, Lynch JP. 2011. Root Cortical Aerenchyma Enhances the Growth of Maize on Soils
with Suboptimal Availability of Nitrogen, Phosphorus, and Potassium. Plant Physiol.
156(3):1190–1201. doi:10.1104/pp.111.175489.

Qin T, Kazim A, Wang Y, Richard D, Yao P, Bi Z, Liu Y, Sun C, Bai J. 2022. Root-Related
Genes in Crops and Their Application under Drought Stress Resistance—A Review. Int J Mol
Sci. 23(19):11477. doi:10.3390/ijms231911477.

Saengwilai P, Nord EA, Chimungu JG, Brown KM, Lynch JP. 2014. Root Cortical Aerenchyma
Enhances Nitrogen Acquisition from Low-Nitrogen Soils in Maize. Plant Physiol. 166(2):726–
735. doi:10.1104/pp.114.241711.

Seethepalli A, Guo H, Liu X, Griffiths M, Almtarfi H, Li Z, Liu S, Zare A, Fritschi FB, Blancaflor
EB, et al. 2020. RhizoVision Crown: An Integrated Hardware and Software Platform for Root
Crown Phenotyping. Plant Phenomics. 2020:3074916. doi:10.34133/2020/3074916.

Segura V, Vilhjálmsson BJ, Platt A, Korte A, Seren Ü, Long Q, Nordborg M. 2012. An efficient
multi-locus mixed-model approach for genome-wide association studies in structured
populations. Nat Genet. 44(7):825–830. doi:10.1038/ng.2314.

Sheahan M, Barrett CB. 2017. Ten striking facts about agricultural input use in Sub-Saharan
Africa. Food Polic. 67:12–25. doi:10.1016/j.foodpol.2016.09.010.

Shimizu A, Yanagihara S, Kawasaki S, Ikehashi H. 2004. Phosphorus deficiency-induced root
elongation and its QTL in rice (Oryza sativa L.). Theor Appl Genet. 109(7):1361–1368.
doi:10.1007/s00122-004-1751-4.

Sitaraman J, Bui M, Liu Z. 2008. LEUNIG_HOMOLOG and LEUNIG Perform Partially
Redundant Functions during Arabidopsis Embryo and Floral Development. Plant Physiol.
147(2):672–681. doi:10.1104/pp.108.115923.

Strock CF, Riva LM de la, Lynch JP. 2017. Reduction in Root Secondary Growth as a Strategy
for Phosphorus Acquisition. Plant Physiol. 176(1):691–703. doi:10.1104/pp.17.01583.

Tang Y, Liu X, Wang J, Li M, Wang Q, Tian F, Su Z, Pan Y, Liu D, Lipka AE, et al. 2016. GAPIT
Version 2: An Enhanced Integrated Tool for Genomic Association and Prediction. Plant
Genome. 9(2):0. doi:10.3835/plantgenome2015.11.0120.

Tissot N, Robe K, Gao F, Grant-Grant S, Boucherez J, Bellegarde F, Maghiaoui A, Marcelin R,
Izquierdo E, Benhamed M, et al. 2019. Transcriptional integration of the responses to iron
availability in Arabidopsis by the bHLH factor ILR3. N Phytol. 223(3):1433–1446.
doi:10.1111/nph.15753.

Trouche G, Bastianelli D, Hamadou TVC, Chantereau J, Rami J-F, Pot D. 2014. Exploring the
variability of a photoperiod-insensitive sorghum genetic panel for stem composition and related
traits in temperate environments. Field Crop Res. 166:72–81. doi:10.1016/j.fcr.2014.06.008.

Villanueva RAM, Chen ZJ. 2019. ggplot2: Elegant Graphics for Data Analysis (2nd ed.). Meas
Interdiscip Res Perspectives. 17(3):160–167. doi:10.1080/15366367.2019.1565254.

Wang H, Wu Y, An T, Chen Y. 2022. Lateral root elongation enhances nitrogen-use efficiency in
maize genotypes at the seedling stage. J Sci Food Agric. 102(12):5389–5398.
doi:10.1002/jsfa.11892.

White PJ, Brown PH. 2010. Plant nutrition for sustainable development and global health. Ann
Bot. 105(7):1073–1080. doi:10.1093/aob/mcq085.

Wissuwa M. 2003. How Do Plants Achieve Tolerance to Phosphorus Deficiency? Small Causes
with Big Effects. Plant Physiol. 133(4):1947–1958. doi:10.1104/pp.103.029306.

Yi K, Wu Z, Zhou J, Du L, Guo L, Wu Y, Wu P. 2005. OsPTF1, a Novel Transcription Factor
Involved in Tolerance to Phosphate Deficiency in Rice. Plant Physiol. 138(4):2087–2096.
doi:10.1104/pp.105.063115.

York LM. 2018. Phenotyping Crop Root Crowns: General Guidance and Specific Protocols for
Maize, Wheat, and Soybean. Methods Mol Biol (Clifton, NJ). 1761:23–32. doi:10.1007/978-1-
4939-7747-5_2.

York LM, Nord EA, Lynch JP. 2013. Integration of root phenes for soil resource acquisition.
Front Plant Sci. 4:355. doi:10.3389/fpls.2013.00355.

Yu P, Hochholdinger F, Li C. 2015. Root-type-specific plasticity in response to localized high
nitrate supply in maize (Zea mays). Ann Bot. 116(5):751–762. doi:10.1093/aob/mcv127.

Supporting information

Genomics of Root System Architecture Adaptation in Sorghum under Nitrogen and

Phosphorus Deficiency

Joshua Magu[1], Joel Masanga[2], Catherin Muui[3], Steven Runo[1*]

7
8
9 The following Supporting Information is available for this article:
10
Figure S1. Genetic diversity of sorghum accessions used for root system architecture analysis.

Table S1: Description of root system architecture (RSA) traits extracted from RhizoVision

analysis.

Figure S2. Sorghum root clusters generated under different nutrient conditions.

Figure S3. Multivariate detection of Single Nucleotide Polymorphisms (SNPs) associated with

Root System Architecture (RSA) using multiple traits and models

Figure S1. Genetic diversity of sorghum accessions used for root system architecture analysis.

(A) Map showing the collection locations of the sorghum accessions. (B) A neighbor-joining (NJ)

tree illustrating the genetic structure and relationships among sorghum races. (C) An admixture

plot depicting the clustering patterns of sorghum races at the optimal K value (K = 7). The key

shows color codes for sorghum races on the map and NJ tree. The key provides color codes

for sorghum races shown on the map and the NJ tree.

Table S1: Description of root system architecture (RSA) traits extracted from RhizoVision analysis
(Seethepalli et al., 2020).
28

Trait	Abbreviation	Category	Description
Total root length	TRL	Size	Computed by determining the Euclidean distance of the medial axis pixel in the skeletonized image.
Perimeter	P	Size	Total Euclidean distance of contour pixels in the segmented image
Average diameter	AD	Distribution	For each pixel in the skeletonized image, the distance to the nearest non-root pixel is computed. This distance is used as radius to fit a circle. The diameter of the circle at each pixel is noted as the root diameter. The list of diameter from all the medial axis is used to determine the average median and maximum diameter for the entire root crown.
median diameter	MD	Distribution	
Maximum diameter	MaxD	Distribution	
Branch points	NBP	Extent	Computed by counting the total number of branch pixels in the skeletonized image with topology.
Branching frequency	BF	Extent	The number of branch points divided by the total root length.
Number of Root Tips	NRTP	Extent	Count of total number of tip pixels in the skeletonized image.
Volume	V	Size	Using the radii determined earlier, the sum of all cross-sectional areas across all the medial axis pixels is noted as the volume and the sum of the perimeter across all the medial axis pixels is noted as the surface area.
Surface area	SA	Size	
Network Area	NA	Size	Total number of pixels in the segmented image.

Figure S2. Sorghum root clusters generated under different nutrient conditions. The average silhouette width is plotted against the number of clusters (k) to determine the optimal number of clusters for each condition: Top (Optimum): Clustering under optimal conditions shows the highest average silhouette width at k = 5, indicating five distinct RSA clusters. Middle (N-Dep): Clustering under nitrogen deficiency (N-Dep) shows the highest silhouette width at k = 2, suggesting two major RSA clusters. Bottom (P-Dep): Clustering under phosphorus deficiency (P-Dep) indicates an optimal cluster number of k = 2 based on the highest silhouette width.

Figure S3. Multivariate detection of Single Nucleotide Polymorphisms (SNPs) associated with Root System Architecture (RSA) using multiple traits and models. **A**. Manhattan Plots: The plots highlight the most prominent SNP, S1_26205071, identified using the Fixed and Random Model Circulating Probability Unification (FarmCPU) model based on four RSA traits: root length diameter range 1 (RLDR1), volume (V), surface area (SA), and number of branching points (NBP). Additionally, SNP S1_26731398 was detected in both the Multi-Locus Mixed Model (MLMM) and Bayesian-information and Linkage-disequilibrium Iteratively Nested Keyway (BLINK) models using the RLDR1 trait. **B**. Quantile-Quantile (Q-Q) Plots: **Bi**: Q-Q plot for the multivariate analysis performed using the FarmCPU model. **Bii**: Combined Q-Q plot for results obtained from the FarmCPU, BLINK, and MLMM models.