Soma to neuron communication links stress adaptation to stress avoidance behavior

Julia Witrado[1], Ella Gundrum[1], Maria Victoria Veroli[1], A.J. Gilsrud[2], Mahdi Eskandarisani[3], Ioannis K. Zervantonakis[3], Steven Mullett[4,5], Stacy Gelhaus[4,5], and Todd Lamitina[1,2*]

[1]Division of Child Neurology, Department of Pediatrics, Children's Hospital of Pittsburgh, University of Pittsburgh Medical Center, Pittsburgh, PA, USA

[2]Center for Neuroscience at the University of Pittsburgh Graduate Training Program

[3]Department of Bioengineering, University of Pittsburgh, Pittsburgh, PA, USA

[4]Department of Pharmacology and Chemical Biology, University of Pittsburgh, Pittsburgh, PA 15261

[5]Health Sciences Mass Spectrometry Core, University of Pittsburgh, Pittsburgh, PA 15261

*Address correspondence to Todd Lamitina, Ph.D., Children's Hospital of Pittsburgh of the University of Pittsburgh Medical Center, Departments of Pediatrics and Cell Biology, 4401 Penn Avenue, Rangos 7122, Pittsburgh, PA 15224, phone - (412) 692-9437; fax – (412) 641-1844; email - stl52@pitt.edu

Abstract

In multicellular organisms, signaling from the nervous system to the peripheral tissues can activate physiological responses to stress. Here, we show that inter-tissue stress communication can also function in reverse, i.e. from the peripheral tissue to the nervous system. *osm-8* mutants, which activate the physiological osmotic stress response in the *C. elegans* skin, also exhibit defective osmotic avoidance (Osm) behavior, via a direct and specific effect on ASH osmosensory neuron excitability. Both *osm-8* and the Patched-related gene *ptr-23*, mutations in which suppress all *osm-8* phenotypes, function in the hypodermal lysosomes to regulate both physiology and behavior. Unbiased lipidomics shows that *osm-8* leads to a *ptr-23*-dependent elevation of the lysosome specific lipid bis(monoacylglycero)phosphate (BMP) and expansion of the pool of hypodermal lysosomes. Just as genetic activation of the osmotic stress response by loss of *osm-8* in the hypodermis causes an Osm phenotype, acute physiological exposure to osmotic stress also confers a reversible Osm phenotype. Behavioral and genetic plasticity requires biosynthesis of the compatible solute glycerol, a key physiological output of the organismal osmotic stress response. However, *ptr-23* is only required for *osm-8* induced behavioral plasticity and not physiological plasticity. Instead, both genetic and physiologically induced Osm phenotypes require the unusual non-neuronal lysosomal V-ATPase subunit *vha-5*, which is also critical for organismal osmotic stress survival. Together, these data reveal that genetic or physiological activation of stress signaling from the skin elicits lysosome-associated signals that modulate organismal neurophysiology to attenuate a sensory neuron circuit. Such 'body-brain' interoceptive communication may allow organisms to better match neuronal decision-making with organismal physiological state.

Introduction

Sensory neuron control of somatic stress responses. Cellular responses to alterations in homeostasis, such as changes in temperature, oxygen, and osmolarity, are commonly referred to as cellular stress responses. This is because cells contain autonomous mechanisms for sensing, signaling, and adapting to the particular challenges posed by each individual stressor. However, in an organismal setting, these autonomous cellular stress response pathways become subservient to systemic control mechanisms. For example, in mice, genetic or pharmacological activation of autophagy in the hypothalamic POMC neurons leads to activation of autophagy in peripheral tissues such as the liver and adipose cells (MARTINEZ-LOPEZ *et al.* 2016). In *C. elegans*, genetic activation of the mitochondrial or endoplasmic reticulum stress response in neurons leads to activation of these same stress response pathways in peripheral intestinal cells (TAYLOR AND DILLIN 2013; BERENDZEN *et al.* 2016). Similarly, optogenetic activation of temperature-sensing neurons can activate the heat shock transcription factor in the germline (TATUM *et al.* 2015). While this neuron-to-soma control of stress response pathways is now well established, what is not clear is whether or not information flows in the reverse direction, i.e. from soma to neurons. Some studies suggest that food and/or pathogen based stress signaling from epithelial cells in *C. elegans* can signal to neurons to modify physiological states, such as longevity, sleep, and learned pathogen avoidance (SINNER *et al.* 2021; SAVINI *et al.* 2022; WU *et al.* 2025). Whether or not abiotic factors, such as osmolarity, also modify behavior through changes in physiological state is less well understood.

Osmotic behavior and physiology in *C. elegans*. The nematode *C. elegans* exhibits two robust responses to increased environmental osmolarity, i.e. hypertonic stress. First, *C. elegans* exhibit behavioral avoidance to hypertonicity (CULOTTI AND RUSSELL 1978).

This response is driven by a well-described neuronal avoidance circuit made up of the amphid and phasmid sensory neurons ASH and PHB which signal to the command interneuron AVA to initiate backwards motion and reorientation away from the stressor (KAPLAN AND HORVITZ 1993; GAT *et al.* 2023). Second, when worms are unable to avoid hypertonicity, they activate physiological adaptation pathways, primarily in the hypodermis (LAMITINA *et al.* 2004). This includes transcriptional up- and down-regulation of >300 osmotic response genes, such as the glycerol biosynthetic enzyme *gpdh-1* (ROHLFING *et al.* 2010) via the activity of hypodermal molecular signaling pathways (URSO *et al.* 2020; URSO *et al.* 2023). Glycerol accumulation closely tracks extracellular solute levels which allows *C. elegans* to maintain cell and tissue volume without raising cytoplasmic ionic strength (LAMITINA *et al.* 2004). Currently, the connection between osmotic avoidance behavior and peripheral tissue osmotic stress responses are not clear. ASH osmo-sensing does not appear to be required for hypertonicity induced *gpdh-1* upregulation or glycerol accumulation in the somatic tissues (LEE *et al.* 2016). Whether or not somatic stress signaling modifies ASH osmotic avoidance behavior has not been investigated.

***osm-8* and *ptr-23*.** Upregulation of the glycerol biosynthetic enzyme *gpdh*-1 in the epithelial tissues is a robust and specific marker of the physiological response to hypertonic stress in *C. elegans* (LAMITINA *et al.* 2006). *gpdh-1* upregulation by hypertonicity is positively regulated by the O-GlcNAc transferase *ogt-1* and the polyadenylation and cleavage factor complex subunits such as Symplekin/*symk-1* (URSO *et al.* 2020; URSO *et al.* 2023). Unexpectedly, *gpdh-1* is also under extensive negative regulation, as inhibition of >100 genes cause significant upregulation of *gpdh-1* under non-stressful isotonic conditions (LAMITINA *et al.* 2006). These negative regulators result in a physiological state change, as mutant animals become completely resistant to

dehydration when exposed to extreme hypertonicity (>500mM NaCl) and accumulate the osmolyte glycerol at close to molar levels (SOLOMON *et al.* 2004; WHEELER AND THOMAS 2006). One of these negative regulators is *osm-8*, a 331 residue protein with a signal sequence and S/T-rich repeats (ROHLFING *et al.* 2011). Loss of *osm-8* leads to upregulation of *gpdh-1*, as well as most of the ORG transcription program, and accumulation of glycerol. As a result, >90% of *osm-8* mutants retain mobility in >500mM NaCl, a condition in which 0% of wild type animals are motile. Through a genetic suppressor screen, we found that all of these physiological phenotypes require the Nieman-Pick C1 (NPC1)/patched-related (ptr) receptor *ptr-23* (ROHLFING et al. 2011). Tissue-specific and temporal rescue experiments reveal that *osm-8* is required in the hypodermal epithelial cells during the L4 stage for these phenotypes. If and how *osm-8* and *ptr-23* regulate neuronal osmotic avoidance behavior has not been determined.

Here, we show that, in addition to its role in regulating peripheral physiology, *osm-8* also regulates neuronal osmotic avoidance behavior by reducing the sensitivity of the ASH neuron to hypertonic stimuli. *osm-8* behavioral defects are completely dependent on *ptr-23*. Furthermore, the expression of *osm-8* and *ptr-23* in the hypodermis is necessary and sufficient to regulate neuronal behavior. Endogenously tagged alleles show that OSM-8 and PTR-23 co-localize to the hypodermal lysosomes. Unbiased lipidomic analysis shows that the lysosome-specific lipid bis(monoacylglycero)phosphate (BMP) exhibits *ptr-23*-dependent upregulation in *osm-8* mutants, which results in an expansion of the pool of lysosomes in the hypodermis. Acute adaptation of wild type worms to mild hypertonicity also induces a reversible osmotic avoidance defective phenotype like *osm-8*. The acquired Osm phenotype is dependent on both glycerol biosynthesis enzymes *gpdh-1* and *gpdh-2* and the lysosomal V-ATPase subunit *vha-5*, which is also required for *osm-8* behavioral phenotypes.

Together, these data suggest that the *osm-8-ptr-23-vha-5* pathway that regulates peripheral stress physiology in the skin also modulates ASH-driven behavioral sensitivity to osmotic stress, thus synchronizing behavioral and physiological states at the organismal level.

Results

osm-8 and _ptr-23_ regulate osmotic stress resistance and osmotic behavioral avoidance. The single _osm-8_ allele, _n1518_, was previously shown to constitutively upregulate the glycerol biosynthesis gene _gpdh-1_, contain elevated glycerol levels, and exhibit an osmotic stress resistance (Osr) phenotype to 500 mM NaCl, a normally lethal hypertonic stress to naïve wild type animals (ROHLFING _et al._ 2011). All of these phenotypes are dependent on the function of the patched-related receptor _ptr-23_. To create definitive null alleles for these genes, we generated deletion alleles for both _osm-8_ and _ptr-23_ using CRISPR/Cas9 (Fig. 1A). Similar to the originally reported alleles, the _osm-8(dr170)_ allele was also Osr while the _ptr-23(dr180)_ allele exhibited no Osr phenotype on its own but fully suppressed the Osr phenotype of _osm-8(d170)_ (Fig. 1B). Mutations in neuron-expressed Osm genes, such as _osm-6_, do not exhibit an Osr phenotype (Fig. 1B). These new alleles represent definitive null alleles for both _osm-8_ and _ptr-23_ that phenocopy previously described mutants and were used for all subsequent studies, unless otherwise indicated.

While _osm-8_ is annotated to exhibit Osmotic Avoidance (Osm) behavioral defects (www.wormbase.org), these phenotypes have never been described or published. When _C. elegans_ encounter hypertonic environments, they first attempt to avoid the stimuli through a well characterized behavioral avoidance response (CULOTTI AND RUSSELL 1978; BARGMANN _et al._ 1990; KAPLAN AND HORVITZ 1993; TROEMEL _et al._ 1995). This avoidance response circuit is primarily driven through detection of osmotic stress via the polymodal sensory neuron ASH. In addition to hypertonicity, ASH also directs avoidance of the volatile odorant 1-octanol via mechanisms that are distinct from those involved in osmosensing (HART _et al._ 1999). _osm-8_ mutants exhibited a strong Osm phenotype similar to that of other Osm mutants, such as _osm-6_ (Fig. 1C). Like the Osr phenotype,

the Osm phenotype of *osm-8*, was also suppressed by *ptr-23(dr180)* or *ptr-23(RNAi)*

(Fig. 1C,D). *ptr-23(RNAi)* did not suppress the *osm-6* Osm phenotype (Fig. 1D), showing

that *ptr-23* is a specific suppressor of *osm-8* physiological and behavioral phenotypes.

Many Osm mutants inhibit sensory cilia formation in chemosensory neurons, which

leads to characteristic morphological, functional, and behavioral phenotypes. To test if

osm-8 also alters sensory cilia formation, we first investigated the morphology or

function of the ASH osmosensory neuron. Unlike *osm-6*, *osm-8* ASH neurons were able

to endocytose the extracellular dye DiI, indicating the presence of functional cilia in *osm-*

8 (Fig. 2A). Next, we determined if the ASH neuron was generally dysfunctional or

specifically dysfunctional towards hypertonicity by examining behavioral responses to

another noxious chemical detected by ASH – 1-octanol. *osm-8* mutants exhibit a

significantly shorter octanol response latency compared to *osm-6*, indicating that the

general ability of ASH to respond to stimuli and the downstream neurons mediating

avoidance behavior is not strongly affected in *osm-8* (Fig. 2B). Consistent with this, we

also found that *osm-8* animals exhibit normal reversal response rates upon direct

stimulation of ASH by the light activated channel channelrhodopsin (Fig. 2C). Finally, we

compared the behaviors of *osm-6* and *osm-8* in the Osm ring assay (Video S1-S3;

(CULOTTI AND RUSSELL 1978)). *osm-6* exhibits relatively few ring encounters because

they tend to cross the ring very early in the assay (Fig. 2D,E). In contrast, *osm-8* mutants

have significantly more ring encounters and do not cross the ring until much later in the

assay as compared to *osm-6* (Fig. 2D,E). Together, these data show that *osm-8* is

required for osmotic avoidance behavior and that this phenotype is not due to defects in

ASH neuron development or gross abnormalities in the ASH avoidance circuit. However,

the kinetics of avoidance in *osm-8* was significantly delayed compared to *osm-6*,

suggesting they regulate avoidance through different mechanisms.

The normal responses to 1-octanol and ChR2 stimulation suggest that the behavioral defect in *osm-8* mutants is due to altered ASH osmosensing. If this is the case, ASH neurons should exhibit less neuronal activation in response to hypertonic stimuli than wild type. To test this, we used a microfluidics-based imaging approach and an ASH-specific GCaMP6s reporter to directly measure the calcium responses in the ASH neuron (CHRONIS *et al.* 2007; PECHUK *et al.* 2022). As a control, we also measured the response of *osm-6* mutants. As previously reported, we found that *osm-6* exhibited no ASH response to a hypertonic stimulus (PHILBROOK *et al.* 2024)(Fig. 3A-C). In contrast, *osm-8* mutants did respond to hypertonicity, but their response was significantly lower than wild type, although still significantly stronger than *osm-6* mutants (Fig.3A-C). The Osm phenotype of *osm-8* mutants was unaffected by the GCaMP transgene (Fig. S1). These data again show that *osm-6* and *osm-8* exhibit substantial phenotypic differences at the level of ASH activation and that the basis of the *osm-8* behavioral defect is likely due in part to the reduced response of ASH to hypertonic stimuli.

There are at least two classes of Osm mutants. Class 1 mutants, such as *osm-6*, are required for the formation of sensory cilia, while Class 2 mutants, such as the TRPV channel *osm-9* (COLBERT *et al.* 1997), have normal cilia but are nonetheless functionally defective in the ability to detect all ASH stimuli. Both Class 1 and Class 2 genes are expressed in ASH, as well as other ciliated sensory neurons. When we examined the single cell gene expression map for the *C. elegans* nervous system (TAYLOR *et al.* 2021), we found that the expression of both Class 1 and Class 2 Osm genes were dramatically enriched in neurons and absent from the hypodermal cells (Fig. 4A). Consistent with previous studies using overexpression reporter transgenes (ROHLFING *et al.* 2011), expression of both *osm-8* and *ptr-23* mRNA exhibit the inverse pattern - they were not detected in sensory neurons, including ASH, but were highly enriched in the hypodermis

(Fig. 4A). These expression data suggest the possibility that *osm-8* and *ptr-23* may alter neuronal osmosensing via their function outside of the nervous system.

Hypodermal function of *osm-8* and *ptr-23* is necessary and sufficient for both physiological stress resistance and neuronal stress avoidance behavior. To test if *osm-8* and *ptr-23* act outside of the nervous system to control behavior, we took two approaches. First, we utilized tissue-specific rescue constructs to restore *osm-8* or *ptr-23* expression solely in the hypodermis and examined their ability to rescue *osm-8* or *osm-8;ptr-23* Osm behavioral phenotypes. In *osm-8* mutants, rescue of *osm-8* using either its endogenous promoter or the hypodermal cell specific *dpy-7* promoter fully rescued both the Osr and Osm phenotypes (Fig. 4B,C). Likewise, in *osm-8;ptr-23* mutants, expression of *ptr-23* from either its endogenous promoter or the hypodermal cell specific *dpy-7* promoter restored the Osr and Osm phenotypes (Fig. 4B,C). Therefore, expression of both *ptr-23* and *osm-8* in the hypodermis is sufficient for rescue of neuronal behavioral defects.

Second, we utilized tissue specific RNAi strains in which genes could be inhibited in either the hypodermis or the intestine (WATTS *et al.* 2020). Since *osm-8* does not produce a strong or consistent RNAi phenotype (ROHLFING *et al.* 2011), we engineered the *osm-8* deletion into each of these strains using CRISPR/Cas9 and performed *ptr-23(RNAi)*. In a global RNAi background, *ptr-23(RNAi)* fully suppressed the Osm and Osr phenotypes of *osm-8* (Fig. 4D,E). Likewise, when RNAi was restricted to the hypodermis, *ptr-23(RNAi)* also fully suppressed the Osm and Osr phenotypes of the *osm-8* mutant (Fig. 4D,E). However, in an intestine-restricted RNAi strain, *ptr-23(RNAi)* had no effect of any *osm-8* phenotypes (Fig. 4D,E). Taken together, the rescue and tissue specific RNAi data show that *osm-8* and *ptr-23* expression outside of the nervous

system in the hypodermis is necessary and sufficient to regulate the physiological Osr and the behavioral Osm phenotypes.

OSM-8 and PTR-23 are co-localized to hypodermal lysosomes and regulate lysosome abundance and volume. We previously generated a rescuing *osm-8::GFP* transgene and failed to observe any fluorescent signal (ROHLFING *et al.* 2011). Since *osm-8* contains a canonical N-terminal signal sequence, this led us to speculate that OSM-8::GFP might be secreted into a cellular compartment that is recalcitrant to GFP folding and/or fluorescence (EVANS *et al.* 2023). To better understand how *osm-8* and *ptr-23* might be regulating behavioral and physiological phenotypes, we generated endogenously tagged alleles using environmentally stable, monomeric fluorophores. Both *osm-8:mNeonGreen::AID*::3XFLAG* and *ptr-23:mScarlett:: AID*::3XFLAG* alleles (referred to as *osm-8::mNG* and *ptr-23:mSc*) were functional, since *osm-8::mNG* did not exhibit an Osr phenotype and *ptr-23:mSc* did not suppress the *osm-8(dr170)* Osr phenotype (Fig. S2). We observed that both OSM-8::mNG and PTR-23::mSc were only expressed in the hypodermis where they exhibited strong co-localization during the L4 stage of development (Fig. 5A-C). However, in adults, OSM-8::mNG was no longer detectable, while PTR-23::mSc was still present (Fig. 5D-F). An identical pattern was observed for both *osm-8:mSc* and *osm-8::sfGFP* CRISPR alleles (Fig. S3). During the L4 stage, PTR-23::mSc co-localized with the lysosomal marker SCAV-3::sfGFP (Fig. 5H-J) while OSM-8::mNG co-localized with the lysosomal marker NUC-1::mCh (Fig. 5K-M). However, PTR-23::mSc was only observed in the lysosomes, suggesting that *osm-8-* dependent signaling occurs through *ptr-23* function in the lysosome.

Given the association of OSM-8 and PTR-23 with the lysosome, which is a major regulator of lipid metabolism (EBNER *et al.* 2025), and the established roles of Patched and Patched-related genes in the transport of hydrophobic molecules including lipids

(Zhong *et al.* 2014) , we hypothesized that *osm-8* and *ptr-23* may regulate lipid homeostasis. Therefore, we performed unbiased liquid chromatography-high resolution mass spectrometry (LC-HRMS) to identify *osm-8* induced, *ptr-23*-dependent changes in relative lipid abundance. We identified 2,708 distinct lipid species among all 3 genotypes (wild type, *osm-8(dr170), osm-8(dr170); ptr-23(dr180)*). Notably, we found that the lysosome/endosome-specific lipid bis(monoacylglycero)phosphate (also known BMP) was upregulated in *osm-8* mutants but restored to wild type levels in *osm-8; ptr-23* mutants (Fig. 6A,B,C,D). *osm-8* upregulated primarily polyunsaturated BMP species, while monounsaturated species were unaffected (Fig. 6C), indicating some level of specificity in the upregulation of BMP lipids by *osm-8*. Interestingly, the proposed precursor for BMP synthesis, phosphatidylglycerol (PG) (Chen *et al.* 2023; Medoh *et al.* 2023; Bulfon *et al.* 2024; Singh *et al.* 2024), did not increase in *osm-8* but was reduced in *osm-8; ptr-23* (Fig. 6E). Triglycerides, which would be predicted to be elevated in *osm-8* mutants due to the extreme levels of glycerol accumulation, were actually reduced in *osm-8* and this was partially reversed in *osm-8; ptr-23* (Fig. 6F). Similar to previous studies, elevation of BMP by *osm-8* also led to increased cholesterol storage, which is known to mainly occur in the lysosome (McCauliff *et al.* 2019). Increased *osm-8* cholesterol levels were partially reversed in *osm-8; ptr-23* animals (Fig. 6G). Together, these lipidomic data show that *osm-8* and *ptr-23* play key roles in the homeostasis of the lysosomal lipid BMP and accumulation of cholesterol.

The elevation in BMP lipids and cholesterol levels observed in *osm-8* suggests that lysosome homeostasis may also be altered in *osm-8*. To test this, we examined lysosomal abundance in the hypodermis of *osm-8*, *ptr-23*, and *osm-8; ptr-23* mutants using the lysosomal marker *nuc-1:mCherry*. We observed a significant increase in the number of hypodermal lysosomes in *osm-8* (Fig. 7A). Automated image analysis

revealed that the increase in hypodermal lysosomes seen in *osm-8* was completely

eliminated in *osm-8; ptr-23* animals (Fig. 7B). *ptr-23* alone had no effect on the size of

the lysosome pool. The increase in the number of lysosomes did not seem to be due to

increased autophagy as the abundance of the autophagosomal marker *lgg-*

1:gfp:mCherry was not altered by *osm-8* (Fig. 7C). We conclude that *osm-8* and *ptr-23*

regulate lysosomal abundance through an autophagy-independent mechanism.

Acute physiological adaptation of wild type animals to mild hypertonic stress

induces a reversible Osm phenotype that requires glycerol biosynthesis and the

lysosomal ATPase subunit *vha-5*. *osm-8* mutants genetically activate the same gene

expression program seen in wild type animals exposed to non-lethal adaptive levels of

hypertonicity (ROHLFING *et al.* 2010) and the expression of many, but not all, of these

genes depends on *ptr-23* (ROHLFING *et al.* 2011). We hypothesized that the osmotic

avoidance behavioral phenotypes observed in *osm-8* might be replicated in wild type

animals by acutely exposing them to mild hypertonic stress. To test this prediction, we

exposed wild type animals to 250 mM NaCl for 1, 3, 6, or 24 hours and then performed

Osm behavioral ring assays. While non-adapted animals are not Osm, we observed that

wild type animals started exhibiting Osm behavior after only 3 hours of adaptation and

this increased by 24 hours (Fig. 8A). The acquired Osm behavior in wild type animals

was reversed by shifting animals back to 51 mM NaCl and was back to baseline levels

after 6 hours (Fig. 8A). Acquired Osm behavior was dose-dependent and could be

induced by exposing animal to 150mM NaCl and above (Fig. S4). These data show that

the Osm behavior is a physiologically plastic phenotypic trait that is constitutively

activated in *osm-8* mutants.

One major physiological consequence of osmotic adaptation in *C. elegans* is

accumulation of the organic osmolyte glycerol (LAMITINA *et al.* 2004). Genetic deletion of

the glycerol-3-phosphate dehydrogenase enzymes *gpdh-1* and *gpdh-2* reduces the accumulation of glycerol by ~50% (LAMITINA *et al.* 2006). To test if glycerol is required for adaptation induced Osm behavior, we adapted *gpdh-1*, *gpdh-2*, and *gpdh-1; gpdh-2* knockouts and analyzed their Osm behavior. We found that Osm behavior was significantly reduced, but not eliminated, in the *gpdh-1; gpdh-2* double mutants, possibly due to the incomplete suppression of glycerol production in this background (Figure 8B). Likewise, the Osm phenotype of *osm-8* was significantly reduced in a *gpdh-1; gpdh-2* mutant background (Fig. 8C). This suggests that both adaptation induced and *osm-8-*induced Osm behavior depends in part on the production of glycerol.

We hypothesized that adaptation induced Osm behavior may require *ptr-23*, since it is required for *osm-8*-induced Osm behavior. However, acquired Osm behavior was not affected by loss of *ptr-23* (Fig. S5). Therefore, we re-examined our prior list of *osm-8* suppressors (ROHLFING *et al.* 2011) and tested if any were required for adaptation induced Osm behavior. We found that the V0 'a' subunit of the V-ATPase *vha-5* was required for both *osm-8* and wild type acquired Osm behavior (6 hour adaptation; Fig. 8D). After 24 hours of exposure to 250mM NaCl, but not 51mM NaCl, all *vha-5(RNAi)* animals died, showing that *vha-5* is also required for hypertonic stress survival (Fig. S6). *vha-5* is only expressed in the hypodermis and excretory cell suggesting that, like *ptr-23*, *vha-5* regulates osmotic avoidance behavior from outside the nervous system (OKA *et al.* 2001; LIEGEOIS *et al.* 2006; LIEGEOIS *et al.* 2007).

Previous studies showed that *vha-5* has both V-ATPase-dependent and -independent functions. The E830Q mutation specifically interferes with the V-ATPase-independent functions in the hypodermis but does not affect the V-ATPase-dependent function in the excretory cell (LIEGEOIS *et al.* 2006). We tested whether *vha-5^{E830Q}* was required for *osm-8* phenotypes. We found that *vha-5(+)* but not *vha-5^{E830Q}* exhibited an Osr and Osm

phenotype in the *osm-8* mutant (Fig. S7A,B). Similarly, *vha-5(+)* exhibited an adaptation-induced Osm phenotype but *vha-5^{E830Q}* did not (Fig. 8E). Therefore, the hypodermal-specific V-ATPase independent functions of *vha-5* mediate the effects of *osm-8* on osmotic stress resistance and behavior. Based on our data, we conclude that *osm-8* and *ptr-23* comprise one branch of a multipronged osmotic stress response pathway, which converges on V-ATPase-independent regulation of the V0 subunit *vha-5* and that signaling from the hypodermis via *vha-5* is required to suppress neuronal osmotic avoidance behavior and promote organismal osmotic stress survival (Fig. 8F).

Discussion

Peripheral stress resistance pathways drive changes in behavior. Our data show
that a behavioral avoidance neuronal circuit can be over-ridden by the activation of
stress response pathways in peripheral cells. Genetic or physiological activation of
osmotic stress response pathways in the barrier hypodermal cells was sufficient to
directly inhibit the osmotic sensitivity of the ASH sensory neuron. This effect was specific
for the osmotic avoidance modality of the ASH circuit, since octanol response latency
was unaffected. Interestingly, the time course of physiological adaptation and reversal
closely matches the accumulation kinetics of the organic osmolyte glycerol and the
glycerol biosynthetic enzymes *gpdh-1* and *gpdh-2* were required for physiological
adaptation. This suggests an elegant feedback model in which the activation of a
physiological response in the peripheral skin tissues produces a metabolite(s) that both
blunts the ability of a sensory neuron to detect and avoid threatening environments and
enhances organismal survival. Such interoceptive 'body-brain' axis signaling between
the peripheral digestive system and the brain for the regulation of feeding behavior is
well documented (STOVER *et al.* 2023). However, our studies of the osmotic stress
response and *osm-8* signaling show that such interoceptive signaling is more pervasive
than previously appreciated and also takes place outside of the gustatory system.

***osm-8-ptr-23* regulation of lysosomes may play a key role in communicating
physiological state to the nervous system.** We showed that *osm-8* and *ptr-23* play
essential roles in coordinating stress physiology with behavior. Surprisingly, we find that
both genes carry out these functions from the hypodermis. In the case of OSM-8, we find
that its accumulation during the L4 stage and disappearance in the adult precisely
matches the temporal rescue requirements that we previously defined (ROHLFING *et al.*
2011). While OSM-8 and PTR-23 are localized to lysosomes, regulate lysosome

abundance, and control the abundance of lysosome-specific lipids such as BMP, the role of the lysosome in osmotic physiology remains enigmatic. A previous genome wide RNAi screen identified several lysosomal genes as being required for hyperosmotic survival, although *vha-5* was not among the genes identified (CHOE AND STRANGE 2008). Another recent study shows that mutation of the cuticular collagen *dpy-7*, which causes an Osr phenotype similar to *osm-8*, exhibits upregulation of some lysosomal genes (FONG *et al.* 2024). One possibility is that lysosomes generate signaling molecules which lead to activation of the osmotic stress response and production of secretory factors that attenuate osmotic avoidance behavior. Such a paradigm was recently described for the role of intestinal lysosomes during aging (SAVINI *et al.* 2022). Another possibility is that the increase in lysosomal abundance and BMP lipids is a downstream consequence of the activation of glycerol accumulation by *ptr-23*-dependent signaling upon loss of *osm-8*. BMP is the major phospholipid constituent of intralumenal membranes of the late endosome/lysosome, where it regulates sorting and degradation of lipids or lipid cargo (KOBAYASHI *et al.* 1998). BMP is composed of two monoacylglycerols linked by a phosphate. Elevation of glycerol could drive increased flux through the BMP biosynthesis pathway, resulting in the expansion of BMP enriched lysosomal membranes. The synthesis mechanism of BMP is not well understood, although several enzymes were recently suggested to catalyze different steps in its biosynthesis (CHEN *et al.* 2023; MEDOH *et al.* 2023; BULFON *et al.* 2024; SINGH *et al.* 2024). Whether or not these genes play important roles in the generation of BMP by the *osm-8/ptr-23* pathway or contribute to the osmotic stress response will require further study.

The V-ATPase proton pump plays a key role in lysosomal physiology by generating the acidic environment required for many aspects of lysosomal function. Like *ptr-23*, the V-ATPase subunit *vha-5* is required for the *osm-8* phenotypes. We were unable to find a

consistent role for other Vha genes in *osm-8* phenotypes due to the strong lethal

phenotype that was observed in most Vha knockdowns except for *vha-5*. This suggests

the possibility that the lysosomal functions of *osm-8*, *ptr-23,* and *vha-5* may not be pH-

dependent. *vha-5* has well established V-ATPase-independent roles in *C. elegans*

(LIEGEOIS *et al.* 2006). Indeed, our data show that these non-canonical functions of *vha-*

5 are likely critical since both the Osm and Osr phenotypes of *osm-8* could be

suppressed by wild type *vha-5* but not the *vha-5(E830Q)* mutant which preserves

canonical V-ATPase functions (LIEGEOIS *et al.* 2006). Non-canonical *vha-5* functions are

not well understood but *vha-5* is associated with the secretion of proteins from the

hypodermis and serves as a marker of specialized apical plasma membrane structures

called meisosomes (LIEGEOIS *et al.* 2006; AGGAD *et al.* 2023). Whether or not the

meisosomal functions of *vha-5* are required cannot yet be determined since mutations

that specifically affect this function(s) are not known.

ptr-23 is part of the expanded Patched Receptor (*ptc*) and Patched-related (*ptr*) gene

family which includes >20 genes in *C. elegans* (ZHONG *et al.* 2014). *ptr* genes have

established roles in molting and apical extracellular matrix development, as well as the

formation of unicellular tube structures (COHEN *et al.* 2021). While Patched and Patched-

related proteins are thought to be hydrophobic transporters due to their overall similarity

with bacterial resistance-nodulation division (RND) transporters, their specific transport

substrates are mostly unknown. One exception is the Patched homolog NPC1, which

encodes a lysosomal cholesterol transporter associated with inherited metabolic disorder

Niemann-Pick C1 Disease (SAHA *et al.* 2020). Mutations in NPC1 cause lysosomal

accumulation of cholesterol and elevation of the lipid BMP (MCCAULIFF *et al.* 2019). This

causes NPC1 patients to develop a progressive form of ataxia in adolescence and

adulthood. Interestingly, we find that mutations in *osm-8* mimic the lipid phenotypes of

NPC1 mutations by raising BMP and cholesterol levels. Conversely, we find that mutation of the NPC1-related gene *ptr-23* reverses the *osm-8* induced changes in lipid composition by lowering cholesterol and BMP. Thus, even though PTR-23 localizes to the lysosome and has sequence similarity to the NPC1 gene, it is unlikely to function as a cholesterol transporter. Consistent with this, many of the key residues required for cholesterol transport in NPC1 are not conserved in PTR-23 (SAHA *et al.* 2020). Further analysis of the Osr/Osm mutant genetic pathway should help to better understand how OSM-8 and PTR-23 contribute to these dramatic lipidomic and phenotypic changes.

Physiological signals that drive changes in the nervous system from the hypodermal lysosome. The precise molecular mechanism by which hypodermal osmotic stress signaling represses the osmotic behavioral response remains unclear. Our Ca2+ imaging data suggest hypodermal stress signaling directly inhibits the ASH osmosensory neuron response to hypertonicity. This may be due to the accumulation of the osmolyte glycerol in both osm-8 mutants and wild type adapted animals, which might reduce the osmotic gradient experienced by ASH sensory machinery. Indeed, adaptation-induced behavioral changes precisely track the accumulation and secretion kinetics of whole animal glycerol accumulation (LAMITINA *et al.* 2004). Consistent with this model, we find that both *osm-8* and wild type adaptation induced behavioral changes are reduced but not eliminated in animals lacking the biosynthetic enzymes *gpdh-1* and *gpdh-2*, likely due to the fact that these mutants reduce but do not eliminate accumulation of glycerol under hypertonic conditions (LAMITINA *et al.* 2006). Other mechanisms could also contribute to the observed behavioral plasticity. For example, a previous study demonstrated that lysosomes generate lipid signals in non-neuronal somatic cells that alter nervous system function and behavior (SAVINI *et al.* 2022). Given the significant changes in lipid content in *osm-8* mutants, it is possible that such lipid

signals may also be involved here. Yet another possibility is that the secretion of several hypodermal expressed neuropeptides, which are upregulated by both *osm-8* and osmotic stress (ROHLFING *et al.* 2010), may communicate with neurons to modify their responses to osmotic stimuli. These neuropeptides are similarly upregulated by infection and were previously shown to regulate infection-induced behavior (SINNER *et al.* 2021). Future studies will be aimed at testing each of these models.

Surprisingly, we found that while loss of *ptr-23* is required for the behavioral adaptation induced by loss of *osm-8*, it is not required for physiological behavioral adaptation induced by high salt. Previously, we showed that even in the absence of *ptr-23*, hypertonicity could still upregulate *gpdh-1* (ROHLFING *et al.* 2011). These data suggests that there is a parallel pathway(s) that regulates the peripheral osmotic stress response and can communicate with neurons. The observation that *vha-5(RNAi)* blocks adaptation induced behavioral changes suggests that this pathway may function in parallel to *ptr-23* and that both pathways converge on *vha-5*. Candidates for such a pathway include the related secreted proteins *osm-7* and *osm-11* (WHEELER AND THOMAS 2006; KOMATSU *et al.* 2008), which may act independently of *ptr-23* (VEROLI AND LAMITINA 2023). Unlike *osm-8*, *osm-7* and *osm-11* are also reported to alter octanol response latency (SINGH *et al.* 2011). This suggests that *osm-7* and *osm-11* might modify the ASH avoidance circuit through a different mechanism from *osm-8*. Our findings here show that *osm-8* mutants reduce the response of ASH to a hypertonic stimulus. Whether or not *osm-7* and/or *osm-11* also inhibit ASH responses to hypertonicity is not known. Future studies investigating ASH responses in these mutants will be important to test these models.

In conclusion, our data reveals a non-gustatory 'body-brain' axis in *C. elegans* that links peripheral physiological state with behavioral plasticity. The sensation and integration of internal physiological state by the nervous system, a process called interoception, is an

emerging but poorly understood process (KHALSA *et al.* 2018). By activating a

physiological stress response pathway without altering external environmental input, the

osm-8-ptr-23 pathway in *C. elegans* represents a primitive but tractable interoceptive

state in which to define the interoceptive signals that arise from the peripheral tissue, the

interoreceptors on neurons, and the mechanisms by which these signals modulate the

nervous system. Such mechanisms could provide insights into a wide variety of

conditions associated with disruptions in the body-brain axis, such as depression (AVERY

et al. 2014), autism (HATFIELD *et al.* 2019), post-traumatic stress syndrome (SIMMONS *et

al.* 2009), and chronic pain (SUAREZ-ROCA *et al.* 2019).

Materials and methods

***C. elegans* strains and culture**: All strains were cultured at 20ºC. A table of all strains used in these studies can be found in Table 1. Wormbase was utilized as the source for all genomic sequence data (STERNBERG *et al.* 2024).

Osmotic stress resistance, adaptation, and osmotic avoidance assays: All assays were performed with Day 1 adults (24 hours post L4 stage). For Osr assays, four replicates of 20 animals were placed in a glass depression slide containing 200μl of M9 with 550 mM NaCl. The number of animals exhibiting movement after 10 minutes was counted.

Osm behavioral assay were performed as previously described (CULOTTI AND RUSSELL 1978), with slight modifications. We printed a ring of 4M NaCl, 0.1% light green SF yellowish dye, pH 7.0 onto standard 3cm NGM plates using a 12mm Falcon snap cap as a template. The cap was gently placed on the agar surface and 40μl of 4M NaCl solution was evenly pipeted around the edge of the cap. The cap was then removed and the 4M NaCl solution was allowed to soak into the plate for 5 minutes prior to the start of the assay. For each assay, 10 day 1 adult animals were placed in S-buffer for ~1 minute to remove bacteria and then placed in the center of the ring using an eyelash pick. Eight replicates of 10 animals were performed for each genotype and/or condition. After 30 minutes, each plate was scored for the number of animals that crossed the ring. Animals that paralyzed on the ring were censored. To prevent animals from re-entering the ring after crossing, we placed a 10ul spot of OP50 or *empty vector(RNAi)* HT115 bacteria on the outer edge of the plate to which worms navigated and dwelled once they crossed the ring. The presence of food did not result in wild type animals crossing the ring, even after up to 1 hour.

For the analysis of ring encounters and ring dwell time, we collected video (2 frames per second) of the whole 3cm plate over the course of the 30 minute Osm assay using the Wormlab imaging system (N=5-6 animals per video, 4-6 videos per genotype, 18-21 animals total per genotype. Animals were picked into 5µl of S-buffer in the center of the ring to prevent engagement with the ring prior to starting the video recording. After each plate was recorded, worms were tracked using WormLab software throughout the course of the 30 minutes. In order to differentiate an encounter from a crossing, two concentric circular labels were created in WormLab, where one label denoted the inner boundary of the ring and the other label denoted the outer boundary. The sizes of these labels were saved and applied to each video. A worm crossing the first label was denoted as an 'encounter', and the same worm crossing the second label was denoted as an escape. The number of encounters a worm underwent was counted, and the time before which the worm escaped the outer label was recorded in seconds. Once a worm crossed the ring, it was no longer considered throughout the remainder of the video. If a worm did not cross the ring after 30 minutes, the time in the ring was denoted as 1800 seconds. Animals that paralyzed inside or on the ring were censored from the analysis.

Octanol response assay - The 1-octanol avoidance assays were performed as previously published with some modifications (TROEMEL *et al.* 1995). The morning before conducting the assay, 4µl of OP50 were seeded as 10 separate spots on five 60mm NGM plates per strain and allowed to dry for ~30 minutes at room temperature. Ten day 1 animals were placed individually onto each spot and left to acclimate to the new plate conditions for approximately 30 minutes. During that time, 1-octanol was diluted to 10% in 100% ethanol. 50µl of diluted 1-octanol was aliquoted into separate PCR tubes for the assays. The 1-octanol aliquots were kept sealed and twelve feet away from the

nematodes when assessing 1-octanol avoidance. For the experiments, a fresh horsehair was parafilmed onto the tip of a Q-tip. Then the horsehair was saturated in the diluted 1-octanol, avoiding the parafilm. Then, the horsehair was placed in front of the forward-moving animal. The 1-octanol avoidance response time was recorded upon initiation of backward locomotion. For the subsequent two biological replicates per animal, the horsehair was re-saturated in diluted 1-octanol following the recording of avoidance time to ensure the presence of the odorant throughout the assay.

Osmotic Survival Assays - Day 1 adults placed onto *empty vector(RNAi)* or *vha-5(RNAi)* plates and incubated for ~65-72 hr at 20°C. Progeny L4 stage animals were then placed onto either the control or high salt RNAi plates containing 250mM NaCl. 24 hours later, day 1 adults were tested for survival by tapping the plate and by tapping the worm's body. Animals that moved forward or backward one full body length were scored as alive.

Imaging and microscopy: L4 stage or day 1 adult animals were immobilized in 50mM levamisole. For images in Figure 4, widefield *nuc-1:mCherry* images were collected as a Z-series (0.27µm step size) on a Leica DMI4000 equipped using a 63x/1.4N.A. lens and a DFC365 digital camera. For quantification of lysosome number, three non-overlapping 25.69µm x 25.69µm x 8.64µm sections centered on the *nuc-1:mCherry* signal in the hypodermis were cropped from each master image. Each cropped section was deconvolved using identical standard settings in imaging software (Leica AF). Puncta in the 3D deconvolved cropped Z-stacks were then analyzed using Aivia software. Training took place on one wildtype image stack and equivalent threshold settings were then applied to all image stacks for all genotypes to generate object counts.

Microfluidics and Ca^{2+} imaging: Custom microfluidic devices were based on a previous design with modification (dual layers to facilitate sample loading/removal and a narrower 50μm diameter worm trap tapered to a 15μm opening) (CHRONIS *et al.* 2007). Silicone wafer molds were printed using maskless photolithography at the University of Pittsburgh Nanofabrication Core Facility. Following silianization of the molds, PDMS elastomer (1:10) was mixed, degassed, poured onto silicon wafers, and incubated at 65ºC for 2-4 hours. The polymerized PDMS was removed from the wafer, cut into individual devices, and 1mm access ports were punched. Chips were sealed to cleaned glass coverslips with a 1 minute plasma treatment of the contact surfaces under vacuum and then applying gentle pressure to seal the PDMS to the coverslip. Sealed devices were baked at 80ºC for 30 minutes to enhance bonding.

Devices were connected to four fluid reservoirs via a direct PVC tubing connection. Stimulus flow was controlled through a valve control system (Valvebank 4, Automate Scientific). Stimuli were diluted in filtered Milli-Q water. Images were acquired with a Leica DMi8 inverted microscope with a 40x dry objective (0.6NA) and a Leica K5 sCMOS camera at 10Hz with 4x4 binning (100ms exposure time). Day 1 adult animals grown without starvation for at least 2 generations were picked onto a clean NGM plate to remove bacteria, aspirated from the clean plate in 0.3mM levamisole to minimize movement in the device, and loaded into the device trap. Prior to imaging, each animal was allowed to equilibrate in the device with flow for 3 minutes without illumination. Each animal was then subjected to 20 seconds of control, 20 seconds of stimulus, and 20 seconds of control solution (600 frames). Data were collected over at least two days using the same solutions with some animals from each genotype being collected each day. Only a single ASH neuron was recorded from each animal.

For GCaMP analysis, images stacks were first aligned using the 'Template Matching' plugin in Fiji. An ROI was manually drawn around the ASH soma and a matching ROI was placed outside the worm for measuring background fluorescence. Mean neuron soma and background fluorescence measurements were used for analysis using custom Python scripts. First, the background subtracted fluorescence was calculated for each neuron. Next, frame numbers were converted to time (ie 10 frames per second). F0 was then calculated for each neuron as the average background subtracted fluorescence during the 15-20 second period. The delta F/F0 was then calculated for each neuron using the formula (F-F0) / F0 where F is the background-subtracted F at each time point. To correct for GCaMP photobleaching, an exponential decay function was fit to the deltaF/F0 data for each neuron during the 10-20 second and 50-60 second periods and this curve was subtracted from the deltaF/F0 data for each neuron. Finally, the corrected deltaF/F0 values for each neuron were converted to percentages and this data was used for time plots, heat maps, and peak value plots. For all time data, the first 10 seconds of data is omitted since this period often contained sample positioning and focusing artifacts. Peak %deltaF/F0 was calculated as the maximum change in fluorescence in the 20-40 second stimulus period. All code is available at:

https://github.com/LamitinaLab/Witrado_et_al_2025.git

Transgene rescue experiments: *osm-8* and *ptr-23* genomic fragments containing the promoter (~2Kb), coding sequence, and 3' UTR (~1Kb) was PCR amplified (Q5 polymerase, New England Biolabs) and cloned into the pMinT2.0 vector. To derive clones with hypodermal-specific expression, we replaced the *osm-8* or *ptr-23* promoters with a *dpy-7* promoter using recombination-based cloning (High-fidelity assembly, NEB).

The sequence of all plasmids was verified by whole-plasmid DNA sequencing (Plasmidsaurus).

To generate *osm-8* rescue lines, we microinjected *osm-8(dr170)* with a mix containing pBluescript II (80ng/µl), pCFJ90/*myo-2p::mCherry* (2ng/µl), and the *osm-8* rescue plasmid (20ng/µl or 2ng/µl for the *dpy-7p* plasmid, which was toxic at higher concentrations). To generate *ptr-23* rescue lines, we injected *osm-8(dr170); ptr-23(dr180)* with a mix containing pBluescript II (80ng/µl), pCFJ90/*myo-2p::mCherry* (2ng/µl), and the *ptr-23* rescue plasmid (20ng/µl or 2ng/µl for the *dpy-7*p plasmid, which was toxic at higher concentrations). Stable extrachromosomal array lines that segregated *myo-2p::mCherry* in the progeny were selected for analysis.

CRISPR/Cas9 genomic editing: The *osm-8* deletion allele *dr170* and the *ptr-23* deletion allele *dr180* were generated using previously described CRISPR/Cas9 editing methods (VEROLI AND LAMITINA 2023). mNeonGreen and mScarlett tags were amplified from pJW2171 and pJW2098 (ASHLEY *et al.* 2021) and inserted into the endogenous *osm-8* or *ptr-23* genes using our standard CRISPR/Cas9 methods (URSO *et al.* 2023). While these alleles contain an AID* degron tag, we did not observe a reduction in either mNeongreen or mScarlett signals in the presence of 1mM K-NAA and a globally expressed TIR1 (*wrdSi58*), likely due to the AID* degron tag being located in the lysosome lumen, which would be inaccessible to the nucleocytoplasmic TIR1. All alleles were verified by DNA sequencing. A table of all primers used in this study can be found in Table 2.

Lipidomics – For each replicate, N2, *osm-8(dr170)*, and *osm-8(dr170); ptr-23(dr180)* animals were seeded on 3x15 cm NGM plate with a lawn of OP50 as starved L1s. Animals were grown at 20ºC until animals were day 1 adults. We collected a total of six

replicates for each genotype for analysis. Worms were washed off plates using M9 buffer, pelleted in a swinging bucket centrifuge at 2,000xg for 1 minute, and then washed three times. Worm pellets were resuspended in 10ml of M9 buffer and incubated at 20ºC with rocking for 15 minutes to evacuate OP50 bacteria from the intestine. Worms were pelleted and the density of worms per microliter was calculated for each strain. For each sample, ~5,000 worms were pipeted into 2.0ml tubes containing a MP Bio Matrix A (garnet/ceramic sphere) and frozen at -80ºC prior to processing.

Metabolic quenching, lysis, and lipid extraction was performed by adding 0.5mL ice cold phosphate buffered saline. Samples were homogenized in MP Biomatrix A tubes at 60 Hz for 1 minute. Uncleared supernatant (400 μL) was transferred to a clean glass tube containing 10 μL of LipidSplash, deuterated internal standards (Avanti Polar Lipids. Alabaster, AL) and subjected to Folch extraction. Samples were rested on ice for 10 minutes before phase separation via centrifugation at 2500 x g for 15 minutes. The organic phase (0.7 mL) was transferred to a clean glass vial and dried to completion under N2. Samples were resuspended in 100 μL of 1:1 acetonitrile:isopropanol and 3μL was injected for online LC-HRMS analysis.

Briefly, samples were injected via a Thermo Vanquish UHPLC and separated over a reversed phase Thermo Accucore C-18 column (2.1×100mm, 5μm particle size) maintained at 55°C.   For the 30 minute LC gradient, the mobile phase consisted of the following: solvent A (50:50 H2O:ACN 10 mM ammonium acetate/0.1% acetic acid) and solvent B (90:10 IPA:ACN 10 mM ammonium acetate/0.1% acetic acid).   The gradient started at 30% B and increased to 43%B over 2 minutes followed by an increase to 55%B in 0.1 minute. The organic increased to 65%B over the next 10 minutes, followed by an increase to 85%B over 6 minutes. For column washing the gradient increased to 100%B over 2 minutes and was held for 5 minutes followed by 5 minutes of equilibration

at 30%B. The Thermo ID-X tribrid mass spectrometer was operated in both positive and negative ESI mode. A data-dependent MS2 method was used with scanning in Full MS mode from 200 to 1500 m/z at 120,000 resolution with an AGC target of 5e4 for triggering MS2 fragmentation using stepped HCD collision energies at 20, 40, and 60% in the Orbitrap at 15,000 resolution. Source ionization settings were 3.5kV and 2.4kV spray voltage, respectively for positive and negative mode. Source gas parameters were 35 sheath gas, 5 auxiliary gas at 300°C, and 1 sweep gas. Calibration was performed prior to analysis using the PierceTM FlexMix Ion Calibration Solutions (Thermo Fisher Scientific). Internal standard peak areas were then extracted manually using Quan Browser (Thermo Fisher Xcalibur ver. 2.7), normalized to weight and internal standard peak area, then graphed using GraphPad PRISM (ver 9.0).   Untargeted differential comparisons were performed using LipidSearch 4.2 (Thermo Fisher) to generate a ranked list of significant lipid compounds at the class and species-specific levels.

Statistical Analysis: Comparisons of means were analyzed with either a two-tailed Students t-test (2 groups) or ANOVA (3 or more groups) using either the Dunnett's, Sidak, or Tukey's post-test analysis as indicated in GraphPad Prism 9 (GraphPad Software, Inc., La Jolla, CA) or Python 3 libraries. p-values of <0.05 were considered significant.

Data Availability Statement: Strains and plasmids are available upon request. The authors affirm that all data necessary for confirming the conclusions of the article are present within the article, figures, and tables. All raw data for all figures is available via FigShare.

Acknowledgements

This work was supported by NIH R01GM135577 to T.L. and NIHS10OD023402 to S.G.. Some mutant strains were provided by the CGC, which was funded by the NIH Office of Research Infrastructure Programs (P40 OD010440). We thank Jun Chen for work performed in the University of Pittsburgh Nanofabrication and Characterization Core Facility (RRID:SCR_05124). NFCF services and instruments used in this project were graciously supported, in part, by the University of Pittsburgh. We also thank Piali Sengupta, Ph.D. and Jamie Yeon, Ph.D. (Brandeis University) for guidance and advice on Ca^{2+} imaging.

Conflict of Interest Statement

The authors have no conflicts of interest to declare

Table 1. Strains used in this study

Strain	Genotype	Source
N2	+/+	CGC
MT3571	osm-8(n1518)	CGC
OG2201	osm-8(dr170)	This study
PR811	osm-6(p811)	(CULOTTI AND RUSSELL 1978)
MAH215	sqIs11 [lgg-1p:lgg-1::mCherry::gfp::lgg-1 3'UTR; rol-6]	(CHANG et al. 2017)
OG1117	ogt-1(jah01)	(KONZMAN et al. 2022)
OG2199	qxIs722; ptr-23(dr169) [CRISPR ptr-23 mScarlet]	This study
OG2180	osm-8(dr163) [CRISPR insertion of mNeonGreen at C-terminus of endogenous osm-8 genomic locus]	This study
OG2192	osm-8(dr165) [osm-8:mSc CRISPR]; qxIs722 [dpy-7p::dpy-7::sfGFP]	This study
OG2200	osm-8(dr163); qxIs257 [ced-1p:nuc-1::mCherry]	This study
OG2210	ptr-23(dr169) [ptr-23:mSc]; qxIs430 [scav-3::gfp]	This study
OG2215	ptr-23(dr179) [CRISPR ptr-23 del.]; osm-8(n1518); qxIs257	This study
OG2216	osm-8(n1518); qxIs257	This study
OG2217	ptr-23(dr180); qxIs257	This study
OG2255	osm-8(dr170); drEx524 [osm-8p::osm-8 genomic DNA (10ng/ul); myo-2p::mCherry (2ng/ul); pBSII (80ng/ul)]	This study
OG2256	osm-8(dr170); ptr-23(dr180); drEx525 [ptr-23p::ptr-23 genomic DNA (10ng/ul); myo-2p::mCherry (2ng/ul); pBSII (80ng/ul)]	This study
OG2257	frSi17 [mtl- 2p::rde-1 3'UTR]; frIs7 [nlp-29p::GFP + col-12p::DsRed]; rde-1(ne300); osm-8(dr187) [CRISPR osm-8 deletion into IG1839]	This study
OG2258	frSi21 [col-62p::rde-1 3'UTR]; frIs7 [nlp-29p::GFP + col-12p::DsRed]; rde-1(ne300); osm-8(dr188) [CRISPR osm-8 deletion into IG1846]	This study
OG2263	osm-8(dr170); drEx526 [dpy-7p::osm-8 genomic DNA (1ng/ul), myo-2p::RFP(2ng/ul), pBSII(100ng/ul)]	This study
OG2264	osm-8(dr170); ptr-23(dr180); drEx527 [exdpy-7p::ptr-23 genomic DNA(1ng/ul), myo-2p::RFP(2ng/ul), pBSII (100ng/ul)]	This study
OG2267	osm-8(dr190) [CRISPR sfGFP]	This study
OG2271	ptr-23(dr180) [ptr-23 deletion]	This study
OG2272	ptr-23(dr180); osm-8(dr170)	This study

OG2274	*gpdh-2(dr195)*	This study
OG2275	*gpdh-1(dr154)*	(VEROLI AND LAMITINA 2023)
OG2276	*gpdh-1(dr154); gpdh-2(dr195)*	This study
OG2290	*osm-6(p811); etYIs1*	This study
OG2292	*osm-8(dr206) etYIs1*	This study
MOS88	*him-5(e1490); etYIs1[[gpa-13p::FLPase + sra-6p::FRT::mCherry::StopCodon::FRT::GCaMP6s]*	(PECHUK *et al.* 2022)
XW5399	*unc-76(e911); qxIs257 [ced-1p::nuc-1::mCherry + unc-76(+)]*	(LI *et al.* 2016)
XW8056	*unc-76(e911); qxIs430 [scav-3 GFP]*	(LI *et al.* 2016)
XW18042	*qxIs722 [dpy-7p::dpy-7::SfGFP]*	(MIAO *et al.* 2020)

Table 2. Oligos used in this study

Primer	Sequence	Purpose
OG108	TTCGTCATTTCCGAAGCTATCATCA	*osm-8* genomic rescue sense
OG109	ATCGGCGTTGGAGCAACATA	*osm-8* genomic rescue antisense
OG2174	TTCTTAGCAGGCGGAGAAGC	*osm-8* C-terminal gRNA
OG2177	GTTCACTACACCATCTCCTCCACA	*osm-8* C-terminal fluorescent reporter genotyping, sense
OG2178	gcaagaggagcagcttctcaacctc	*osm-8* C-terminal fluorescent reporter genotyping, antisense
OG2241	gtattttgcactacgggtcg	*gpdh-2* N-terminal gRNA
OG2252	GCATTTGTGCAGGTGCATCT	*osm-8* N-terminal gRNA
OG2276	tatttccaagcagaaaaagcaaaatgtttgacttcttggctggaggggaggctggacgacttcttggaggtggcggtggatcgggaggag	*osm-8:mScarlett:AID:3XFLAG* C-terminal repair template primer, sense
OG2277	gttcacaaataatttaaagcgaaaaaaataaatTTAGCCTCCGGACGATGCTCCTGAGGCTCC	*osm-8:mScarlett:AID:3XFLAG C-terminal repair template primer, antisense*
OG2278	GAGTGTGGTCACTGATCGAC	*ptr-23* C-terminal gRNA
OG2279	gttcagccggaaaagagtgtggtcactgatcgacctgaaatcggtggcggtggatcgggaggagg	*ptr-23:mSc:AID:3XFLAG* C-terminal repair template primer, sense
OG2280	AAGTTAAAAAGAAATATAATTAATGGTTTTTTTTAGCCTCCGGACGATGCTCCTGAGGC	*ptr-23:mSc:AID:3XFLAG* C-terminal repair template primer, antisense
OG2281	CTTCATGGTCTCGTGTTCTTACC	*ptr-23:mSc:AID:3XFLAG* genotyping primer outer sense
OG2282	cttacctgctatcttgattcttttg	*ptr-23:mSc:AID:3XFLAG* genotyping primer inner sense
OG2283	CATGGATCCGCCTCCTCCTGAAC	*ptr-23:mSc:AID:3XFLAG* genotyping primer outer antisense
OG2284	TCCTCCTGAACCACCGCCGCCAC	*ptr-23:mSc:AID:3XFLAG* genotyping primer inner antisense
OG2285	gactacaaagaccatgacggtg	*ptr-23:mSc:AID:3XFLAG* genotyping primer outer sense

OG2286	gacggtgattataaagatcatg	*ptr-23:mSc:AID:3XFLAG* genotyping primer inner sense
OG2287	GGCAAAATGAAACTCAAGAAGAT	*ptr-23* deletion screening primer, outer antisense
OG2288	CTCAAGAAGATGTCGGAAAATCAC	*ptr-23* deletion screening primer, inner antisense
OG2289	ACCTGCAGCCGCTTCGACGA	*ptr-23* N-terminal gRNA
OG2290	ccacattggcgccgccacctgcagccgcttcgacgaaaaaaccatta attatatttcttttaacttctc	*ptr-23* deletion CRISPR HDR template
OG2291	ataagtttcgttcagaagtcctg	*ptr-23* deletion screening primer outer sense
OG2292	cctggattaagttgaaatacttc	*ptr-23* deletion screening primer inner sense
OG2302	actaatacatatattttttgattcattcaatattttgacgacttcttggataaattt atttttttcgcttt	*osm-8* deletion CRISPR HDR template
OG2366	ttccagataaatgccagccaaaatgttg	plasmid PCR for *dpy-7p:osm-8* promoter replacement, sense
OG2367	ttggtagggaatcatggttttacctcctg	plasmid PCR for *dpy-7p:osm-8* promoter replacement, antisense
OG2368	aaaccatga ttccctaccaattgaaaattc	*dpy-7* promoter primer for osm-8 promoter replacement, sense
OG2369	tggctggcatttatctggaacaaaatgtaagaatattc	*dpy-7* promoter primer for osm-8 promoter replacement, antisense
OG2370	ttccagataaatggacgggagcatcgctc	*osm-8* plasmid PCR for *dpy-7p:ptr-23* promoter replacement, sense
OG2371	tcacctaaattgcgtttaaaccttcctgcag	*osm-8* plasmid PCR for *dpy-7p:ptr-23* promoter replacement, antisense
OG2372	tttaaacgcaatttaggtgacactatagaag	*dpy-7* promoter primer for *ptr-23* promoter replacement, sense
OG2373	tcccgtccatttatctggaacaaaatgtaagaatattc	*dpy-7* promoter primer for *ptr-23* promoter replacement, antisense
OG2374	GGAGAAGCTGGACGACTTCT	*osm-8* C-terminal gRNA
OG2375	acttcttagcaggcggagaagctggacgacttcttatgagtaaaggaga agaactttcactg	*osm-8:sfGFP* repair primer, sense
OG2376	ttcacaaataatttaaagcgaaaaaaataaatttatcctttgtatagttcatc catgccatgtg	*osm-8:sfGFP* repair primer, antisense
OG2377	tttccagtagtgcaaataaatttaag	*osm-8:sfGFP* screening primer antisense

OG2378	tcaacaaaatactccaattggcgatg	*osm-8:sfGFP* screening primer sense
OG2379	AATGATATTTACATGTGTTC	*gpdh-2* C-terminal gRNA
OG2384	TTCTTCATCAACAAAATCGAAAACTAGAATTTCCT TAAATATCATTATAGGTTTCGCAAATTTCAATTAG	*gpdh-2* deletion repair oligo
OG2385	CAACTGCGCGCAGTGAATGCTGAG	*gpdh-2* deletion screening primer, sense
OG2386	GCTAGAACGAGGCATTGCACAGTGC	*gpdh-2* deletion screening primer, antisense

Figure legends

Figure 1 – *osm-8* physiological and behavioral phenotypes depend on *ptr-23*. A) Diagram of the *osm-8* and *ptr-23* alleles generated in this study. A detailed list of these and other strains can be found in Table 1. The *osm-8(n1518)* allele was characterized in a prior study (ROHLFING *et al.* 2011). B) Osmotic stress resistance (OSR) phenotype of the indicated genotypes. N=8 replicates per genotype (10 animals per replicate, N=80 per genotype). ****-p<0.0001, One-way ANOVA with Kruskal-Wallis post hoc test. Individual data points are shown along with the mean ± S.D. C) Osmotic avoidance phenotype of the indicated genotypes. N=8 replicates per genotype (10 animals per replicate, N=80 per genotype). ***-p<0.001, 'ns' – p>0.05, One-way ANOVA with Kruskal-Wallis post hoc test. Individual data points are shown along with the mean ± S.D. D) Osmotic avoidance phenotype of the indicated genotypes. N=8 replicates per genotype (10 animals per replicate, N=80 per genotype). **-p<0.01, 'ns' – p>0.05, One-way ANOVA with Kruskal-Wallis post hoc test. Individual data points are shown along with the mean ± S.D.

Figure 2 –*osm-8* mutants have normal ASH osmosensory neuron development and exhibit a distinct behavioral phenotype from the ciliary mutant *osm-6*. A) DiI staining of day 1 adults of the indicated genotypes. B) Latency to reversal response with 10% octanol. N= 98-136 individual trials per genotype. ****-p<0.0001, One-way ANOVA with Tukey's multiple comparison test. Individual data points are shown along with the mean ± S.D. C) Proportion of animals reversing in response to ChR2 stimulation in the presence of all-trans retinal. Data shown are the mean +/- S.D. of 9-12 replicates of 5-10 animals each. *-p<0.05, 'n.s.' – not significant, One-way ANOVA with Tukey post-hoc testing. D) Number of encounters with the hypertonic ring in the osmotic avoidance

assay per individual animal. N=18-21 animals per genotype. ****-p<0.0001, One-way

ANOVA with Tukey's multiple comparison test. Individual data points are shown along

with the mean ± S.D. E) Time inside the hypertonic ring in the osmotic avoidance assay

per individual animal. N=18-21 animals per genotype. ****-p<0.0001, One-way ANOVA

with Tukey's multiple comparison test. Individual data points are shown along with the

mean ± S.D.

Figure 3 – ASH neuronal response to hypertonicity is reduced in *osm-8* mutants.

A) Mean +/- SE of wild type, *osm-8*, and *osm-6* GCaMP6s fluorescence in ASH neurons

in response to a 20 second exposure to 2M NaCl. N=17, 15, and 12 animals per

genotype. Gray shading indicates the period of exposure to 2M NaCl. B) Normalized

color coded GCaMP6s calcium responses of ASH to 2M NaCl. Heatmaps show the

calcium levels of individual neurons for each genotype. The stimulus is applied from 10-

30 seconds. The number of neurons per group is shown at the bottom left corner of each

heatmap. C) Mean +/- SEM of the maximum peak %deltaF/F0 during the stimulus

period. *** - p<0.001, One-way ANOVA with Tukey post-hoc testing.

Figure 4 – *osm-8* and *ptr-23* hypodermal expression is necessary and sufficient to

regulate osmotic avoidance behavior. A) Predicted expression of *osm-8*, *ptr-23*, and

other Class 1 & 2 Osm mutants across sensory neurons and hypodermal cells adapted

from the *C. elegans* neuronal gene expression map and network (CeNGEN) (TAYLOR *et*

al. 2021)   B) Osmotic stress resistance (OSR) phenotype of the indicated genotypes

with *osm-8* or *ptr-23* rescue under either their endogenous promoter or the hypodermal

specific promoter *dpy-7p*. N=4 replicates per genotype (20 animals per replicate, N=80

per genotype). ***-p<0.001, One-way ANOVA with Kruskal-Wallis post hoc test.

Individual data points are shown along with the mean ± S.D. C) Osmotic avoidance

phenotype of the indicated genotypes with *osm-8* or *ptr-23* rescue under either their

endogenous promoter or the hypodermal specific promoter *dpy-7p*. N=8 replicates per

genotype (10 animals per replicate, N=80 per genotype). ****-p<0.0001, *** – p<0.001,

**-p<0.01, One-way ANOVA with Kruskal-Wallis post hoc test. Individual data points are

shown along with the mean ± S.D. D) Osmotic stress resistance (OSR) phenotype of

the indicated genotypes with empty vector or *ptr-23* RNAi. N=4 replicates per genotype

(20 animals per replicate, N=80 per genotype). ***-p<0.001, **-p<0.01, One-way ANOVA

with Kruskal-Wallis post hoc test. Individual data points are shown along with the mean

± S.D.. E) Osmotic avoidance phenotype of the indicated genotypes with empty vector or

ptr-23 RNAi. N=8 replicates per genotype (10 animals per replicate, N=80 per genotype).

****-p<0.0001, *-p<0.05, 'ns'-p>0.05, One-way ANOVA with Kruskal-Wallis post hoc test.

Individual data points are shown along with the mean ± S.D.

Figure 5 – OSM-8 and PTR-23 co-localize in hypodermal lysosomes. A) Wide-field

fluorescence images of the hypodermis in L4 stage animals expressing A) *ptr-23::mSc*,

B) *osm-8::mNG*, and C) the merged overlay. Scale bar = 10µm. Wide-field fluorescence

images of the hypodermis in day 1 adult animals expressing D) *ptr-23::mSc*, E) *osm-8::mNG*, and F) the merged overlay the indicated endogenously tagged alleles. Scale

bar = 10µm. Wide field fluorescence images of Day 1 adults expressing H) the

lysosomal marker *scav-3::sfGFP*, I) endogenously tagged *ptr-23:mSc*, and J) the

merged overlay. Scale bar=10µm. Deconvolved wide field image of L4 stage animals

expressing K) *osm-8::mNG*, L) the lysosomal marker *nuc-1:mCh*, and M) the merged

overlay. Scale bar= 10µm. Dotted lines indicate the seam cell boundary, which lack

OSM-8 expression.

Figure 6 – The lysosome-specific lipid bis(monoacylglyerol)phosphate (BMP) and cholesterol are upregulated in *osm-8* mutants in a *ptr-23* dependent manner. A) Heatmap of the normalized $\log_2$ fold change for each indicated lipid class. Each column represents an individual replicate of 5,000 worms. AcCa=Acyl Carnitine, AEA=N-Acylethanolamine, BisMePE=Bis-methyl phosphatidylethanolamine, BMP=Bis(monoacylglyerol)phosphate, Cer=Ceramide, CerPE=Ceramide PE, Ch=Cholesterol, CL=Cardiolipins, Co=CoenzymeQ, DG=Diglycerides, Hex1Cer=Hexosylceramide, LPC=LysoPC, LPE=LysoPE, MG=Monoglycerides, OAHFA=OAcyl-(gamma-hydroxy)FA, PA=Glycerophosphatidic acid, PC=Glycerophosphocholines, PE=Glycerophosphoethanolamines, PFAA=Primary fatty acid amines, PG=Glycerophosphoglycerols, PI=Glycerophosphoinositols, PS=Glycerophosphoserines, SM=Sphingomyelins, TG=Triacylglycerols. B) Volcano plot of the $\log_2$ fold change versus $\log_{10}$ p-value for all identified lipid species in *osm-8(dr170)* compared to wild type (top) or *osm-8(dr170); ptr-23(dr180)* compared to wild type (bottom). BMP lipids, which are strongly upregulated in *osm-8(dr170)*, are highlighted in red. C) Normalized fold change for all BMP lipid species. Data shown are mean ± S.D. Lipids are organized as poly-unsaturated (>1 acyl chain double bonds) or mono-unsaturated (1 or 0 acyl chain double bonds). D) Normalized $\log_2$ fold change for BMP lipid group in wild type, *osm-8(dr170)* and *osm-8(dr170); ptr-23(dr180)*. N=6 samples per genotype. Horizontal line indicates the mean. ****-$p<0.0001$, One-way ANOVA with Tukey post hoc test. E) Normalized $\log_2$ fold change for phosphoglycerol lipid group in wild type, *osm-8(dr170)* and *osm-8(dr170); ptr-23(dr180)*. N=6 samples per genotype. Horizontal line indicates the mean. ****-$p<0.0001$, **-$p<0.01$, One-way ANOVA with Tukey post hoc test. F) Normalized $\log_2$ fold change for triglyceride lipid group in wild

type, *osm-8(dr170)* and *osm-8(dr170); ptr-23(dr180)*. N=6 samples per genotype. Horizontal line indicates the mean. ****-p<0.0001, ***-p<0.001, One-way ANOVA with Tukey post hoc test. G) Normalized log$_2$ fold change for phosphoglycerol lipid group in wild type, *osm-8(dr170)* and *osm-8(dr170); ptr-23(dr180)*. N=6 samples per genotype. Horizontal line indicates the mean. ****-p<0.0001, ***-p<0.001, **-p<0.01, One-way ANOVA with Tukey post hoc test.

Figure 7 – *osm-8* expands the pool of hypodermal lysosomes through a *ptr-23*-dependent, autophagy-independent mechanism. A) Deconvolved flattened Z-stacks of hypodermal *nuc-1:mCh* lysosomal fluorescence in day 1 adult animals of the indicated genotype. Scale bar = 5µm. B) Quantification of the number of *nuc-1:mCh* puncta in each genotype. Individual data points are shown along with the mean ± S.D.. N=18-21 measurements from 6-7 individual animals per genotype. ****-p<0.0001, 'ns'-not significant, One-way ANOVA with Tukey post hoc multiple comparison testing. C) Widefield images of *lgg-1::gfp:mCh* autophagy marker in day 1 adults of the indicated genotype. Individual green puncta represent autophagosomes while red puncta represent autolysosomes. Scale bar = 5µm

Figure 8 – Mild physiological adaptation to osmotic stress drives a reversible Osm phenotype in wild type animals that is dependent on the V-ATPase subunit *vha-5*.
A) Adaptation-induced osmotic avoidance behavior in day 1 wild type animals. Black symbols represent animals that were grown on 50mM NaCl and then shifted to 250mM NaCl for the indicated amount of time prior to performance of a standard Osm assay. Red symbols indicate animals that were grown on 250mM NaCl and then shifted back to 50mM NaCl for the indicated amount of time prior to the performance of a standard Osm

assay. N=8 replicates per time point for adaptation and reversal (10 animals per replicate, N=80 per genotype). Data points are the mean ± S.D. ****-p<0.0001. ***-p<0.001, 'ns'-not significant versus t=0. Two-way ANOVA with Tukey post hoc test. B) Osmotic adaptation osmotic avoidance assay in wild type, *gpdh-1(dr154)*, *gpdh-2(dr195)*, and *gpdh-1(dr154); gpdh-2(dr195)* day 1 adult animals. N=8 replicates per genotype and per time point (10 animals per replicate, N=80 per genotype for each time point). **-p<0.01, ***-p<0.001, ****-p<0.0001, One-way ANOVA with Tukey post hoc test. Individual data points are shown along with the mean ± S.D. C) Osmotic avoidance assay in wild type, *osm-8(dr170)*, and *osm-8(dr170); gpdh-1(dr154); gpdh-2(dr195)* day 1 adult animals. N=8 replicates per genotype and per time point (10 animals per replicate, N=80 per genotype for each time point). 'ns'-not significant, One-way ANOVA with Tukey post hoc test. Individual data points are shown along with the mean ± S.D. D) Osmotic adaptation osmotic avoidance assay in wild type, *vha-5(RNAi), osm-8(dr170),* and *osm-8(dr170); vha-5(RNAi)* day 1 adult animals. N=8 replicates per genotype and per time point (10 animals per replicate, N=80 per genotype for each time point). ****-p<0.0001, 'ns'-not significant, One-way ANOVA with Tukey post hoc test. Individual data points are shown along with the mean ± S.D. E) Osmotic adaptation assay in vha-5(mc38) day 1 adult null mutant with either *vha-5(+)* wild type rescue or *vha-5^{E830Q}* rescue. N=8 replicates per genotype and per time point (10 animals per replicate, N=80 per genotype for each time point). ****-p<0.0001, 'ns'-not significant, One-way ANOVA with Tukey post hoc test. Individual data points are shown along with the mean ± S.D. F) working model for the role of *osm-8*, *ptr-23*, and *vha-5* in the regulation of peripheral physiology and behavior.

Supplemental Figures

Figure S1 – Osmotic avoidance behavior in wild type and *osm-8* mutant animals with or without the *etYIs1* GCaMP6s transgene. N=8 replicates per genotype (10 animals per replicate, N=80 per genotype). ****-p<0.0001, 'ns'-not significant, One-way ANOVA with Tukey post hoc test. Individual data points are shown along with the mean ± S.D.

Figure S2 – Osmotic stress resistance phenotype of tagged *osm-8* and *ptr-23* alleles. N=4 replicates per genotype (20 animals per replicate, N=80 per genotype). ****-p<0.0001, 'n.s.' – not significant, One-way ANOVA with Tukey posthoc testing.

Figure S3 – Localization of endogenously tagged *osm-8* alleles. A) *osm-8(dr165) [mScarlet* CRISPR allele] B) *dpy-7::sfGFP* C) merge. Scale bar = 10 microns. D) *osm-8(dr190) [sfGFP* CRISPR allele] in L4 or E) day 1 adult animals. Scale bar = 10 microns.

Figure S4 – Dose response of osmotic adaptation-induced Osm behavior in wild type animals. Adaptation period was 24 hours. N=8 replicates per genotype and per time point (10 animals per replicate, N=80 per genotype for each concentration). ****-p<0.0001, 'ns'-not significant, One-way ANOVA with Tukey post hoc test. Individual data points are shown along with the mean ± S.D.

Figure S5 – Adaptation induced Osm behavior in wild type and *ptr-23(dr180)* mutants. N=8 replicates per genotype and per time point (10 animals per replicate, N=80 per genotype for each time point). ****-p<0.0001, 'ns'-not significant, One-way ANOVA with Tukey post hoc test. Individual data points are shown along with the mean ± S.D.

Figure S6 - Survival of *vha-5(RNAi)* day 1 adults in either wild type or *ptr-23(dr180)* after 24 hour exposure to NGM plates with either 50mM NaCl or 250mM NaCl. N=4 replicates per condition (20-25 animals per replicate, 80-100 animals per condition per genotype). ****-p<0.0001, 'ns'-not significant, One-way ANOVA with Tukey post hoc testing. Individual data points are shown along with the mean ± S.D.

Figure S7 – Non-canonical V-ATPase independent functions of *vha-5* are required for *osm-8* phenotypes. A) OSR phenotype of wild type or *osm-8* mutants in the *vha-5* null mutant *mc38* with rescue from either a *vha-5(+)* wild type or *vha-5E830Q* mutant transgene. N=4 replicates per genotype (20 animals per replicate, N=80 per genotype). ****- p<0.0001, 'n.s.' – not significant, One-way ANOVA with Tukey post hoc testing. B) Osm behavior in wild type or *osm-8* mutants in the *vha-5* null mutant *mc38* with rescue from either a *vha-5(+)* wild type or *vha-5E830Q* mutant transgene. N=8 replicates per genotype and per time point (10 animals per replicate, N=80 per genotype for each time point). ****-p<0.0001, 'ns'-not significant, One-way ANOVA with Tukey post hoc test. Individual data points are shown along with the mean ± S.D.

Video S1. Wild type osmotic avoidance behavior. Ring assay with 4M NaCl. Spot of OP50 *E. coli* in upper right quadrant provides both a motivating stimulus to cross the ring and a 'sink' to retain worms that cross the barrier to prevent them from re-entering the ring. Note that wild type animals exhibit chemotaxis towards the OP50 stimulus even when inside the ring, consistent with an odorant-based attractive mechanism.

Video S2. *osm-6(p811)* osmotic avoidance behavior. Ring assay with 4M NaCl. Spot of OP50 *E. coli* in upper right quadrant provides both a motivating stimulus to cross the ring

and a 'sink' to retain worms that cross the barrier to prevent them from re-entering the ring. Note that *osm-6* mutants do not appear to chemotaxis towards the OP50 stimulus, consistent with their general defect in both soluble and odorant-based sensory neuron function.

Video S3. *osm-8(dr170)* osmotic avoidance behavior. Ring assay with 4M NaCl. Spot of OP50 *E. coli* in upper right quadrant provides both a motivating stimulus to cross the ring and a 'sink' to retain worms that cross the barrier to prevent them from re-entering the ring. Note that unlike *osm-6*, *osm-8* mutants chemotaxis towards the OP50 stimulus, consistent with the hypothesis that *osm-8* mutants do not cause a general defect in chemosensory function.

Aggad, D., N. Brouilly, S. Omi, C. L. Essmann, B. Dehapiot *et al.*, 2023 Meisosomes, folded membrane microdomains between the apical extracellular matrix and epidermis. Elife 12.

Ashley, G. E., T. Duong, M. T. Levenson, M. A. Q. Martinez, L. C. Johnson *et al.*, 2021 An expanded auxin-inducible degron toolkit for Caenorhabditis elegans. Genetics 217.

Avery, J. A., W. C. Drevets, S. E. Moseman, J. Bodurka, J. C. Barcalow *et al.*, 2014 Major depressive disorder is associated with abnormal interoceptive activity and functional connectivity in the insula. Biol Psychiatry 76**:** 258-266.

Bargmann, C. I., J. H. Thomas and H. R. Horvitz, 1990 Chemosensory cell function in the behavior and development of Caenorhabditis elegans. Cold Spring Harb Symp Quant Biol 55**:** 529-538.

Berendzen, K. M., J. Durieux, L. W. Shao, Y. Tian, H. E. Kim *et al.*, 2016 Neuroendocrine Coordination of Mitochondrial Stress Signaling and Proteostasis. Cell 166**:** 1553-1563 e1510.

Bulfon, D., J. Breithofer, G. F. Grabner, N. Fawzy, A. Pirchheim *et al.*, 2024 Functionally overlapping intra- and extralysosomal pathways promote bis(monoacylglycero)phosphate synthesis in mammalian cells. Nat Commun 15**:** 9937.

Chang, J. T., C. Kumsta, A. B. Hellman, L. M. Adams and M. Hansen, 2017 Spatiotemporal regulation of autophagy during Caenorhabditis elegans aging. Elife 6.

Chen, J., A. Cazenave-Gassiot, Y. Xu, P. Piroli, R. Hwang, Jr. *et al.*, 2023 Lysosomal phospholipase A2 contributes to the biosynthesis of the atypical late endosome lipid bis(monoacylglycero)phosphate. Commun Biol 6**:** 210.

Choe, K. P., and K. Strange, 2008 Genome-wide RNAi screen and in vivo protein aggregation reporters identify degradation of damaged proteins as an essential hypertonic stress response. Am J Physiol Cell Physiol 295**:** C1488-1498.

Chronis, N., M. Zimmer and C. I. Bargmann, 2007 Microfluidics for in vivo imaging of neuronal and behavioral activity in Caenorhabditis elegans. Nat Methods 4**:** 727-731.

Cohen, J. D., C. E. Cadena Del Castillo, N. D. Serra, A. Kaech, A. Spang *et al.*, 2021 The Caenorhabditis elegans Patched domain protein PTR-4 is required for proper organization of the precuticular apical extracellular matrix. Genetics 219.

Colbert, H. A., T. L. Smith and C. I. Bargmann, 1997 OSM-9, a novel protein with structural similarity to channels, is required for olfaction, mechanosensation, and olfactory adaptation in Caenorhabditis elegans. J Neurosci 17**:** 8259-8269.

Culotti, J. G., and R. L. Russell, 1978 Osmotic avoidance defective mutants of the nematode Caenorhabditis elegans. Genetics 90**:** 243-256.

Ebner, M., F. Frohlich and V. Haucke, 2025 Mechanisms and functions of lysosomal lipid homeostasis. Cell Chem Biol 32**:** 392-407.

Evans, D. C. S., A. B. Khamas, L. Marcussen, K. S. Rasmussen, J. K. Klitgaard *et al.*, 2023 GFP fusions of Sec-routed extracellular proteins in Staphylococcus aureus reveal surface-associated coagulase in biofilms. Microb Cell 10**:** 145-156.

Fong, A., M. Rodriguez and K. P. Choe, 2024 Increased expression of metabolism and lysosome-associated genes in a C. elegans dpy-7 cuticle furrow mutant. MicroPubl Biol 2024.

Gat, A., V. Pechuk, S. Peedikayil-Kurien, S. Karimi, G. Goldman *et al.*, 2023 Integration of spatially opposing cues by a single interneuron guides decision-making in C. elegans. Cell Rep 42**:** 113075.

Hart, A. C., J. Kass, J. E. Shapiro and J. M. Kaplan, 1999 Distinct signaling pathways mediate touch and osmosensory responses in a polymodal sensory neuron. J Neurosci 19**:** 1952-1958.

Hatfield, T. R., R. F. Brown, M. J. Giummarra and B. Lenggenhager, 2019 Autism spectrum disorder and interoception: Abnormalities in global integration? Autism 23**:** 212-222.

Kaplan, J. M., and H. R. Horvitz, 1993 A dual mechanosensory and chemosensory neuron in Caenorhabditis elegans. Proc Natl Acad Sci U S A 90**:** 2227-2231.

Khalsa, S. S., R. Adolphs, O. G. Cameron, H. D. Critchley, P. W. Davenport *et al.*, 2018 Interoception and Mental Health: A Roadmap. Biol Psychiatry Cogn Neurosci Neuroimaging 3**:** 501-513.

Kobayashi, T., E. Stang, K. S. Fang, P. de Moerloose, R. G. Parton *et al.*, 1998 A lipid associated with the antiphospholipid syndrome regulates endosome structure and function. Nature 392**:** 193-197.

Komatsu, H., M. Y. Chao, J. Larkins-Ford, M. E. Corkins, G. A. Somers *et al.*, 2008 OSM-11 facilitates LIN-12 Notch signaling during Caenorhabditis elegans vulval development. PLoS Biol 6**:** e196.

Konzman, D., T. Fukushige, M. Dagnachew, M. Krause and J. A. Hanover, 2022 O-GlcNAc transferase plays a non-catalytic role in C. elegans male fertility. PLoS Genet 18**:** e1010273.

Lamitina, S. T., R. Morrison, G. W. Moeckel and K. Strange, 2004 Adaptation of the nematode Caenorhabditis elegans to extreme osmotic stress. Am J Physiol Cell Physiol 286**:** C785-791.

Lamitina, T., C. G. Huang and K. Strange, 2006 Genome-wide RNAi screening identifies protein damage as a regulator of osmoprotective gene expression. Proc Natl Acad Sci U S A 103**:** 12173-12178.

Lee, E. C., H. Kim, J. Ditano, D. Manion, B. L. King *et al.*, 2016 Abnormal Osmotic Avoidance Behavior in C. elegans Is Associated with Increased Hypertonic Stress Resistance and Improved Proteostasis. PLoS One 11**:** e0154156.

Li, Y., B. Chen, W. Zou, X. Wang, Y. Wu *et al.*, 2016 The lysosomal membrane protein SCAV-3 maintains lysosome integrity and adult longevity. J Cell Biol 215**:** 167-185.

Liegeois, S., A. Benedetto, J. M. Garnier, Y. Schwab and M. Labouesse, 2006 The V0-ATPase mediates apical secretion of exosomes containing Hedgehog-related proteins in Caenorhabditis elegans. J Cell Biol 173**:** 949-961.

Liegeois, S., A. Benedetto, G. Michaux, G. Belliard and M. Labouesse, 2007 Genes required for osmoregulation and apical secretion in Caenorhabditis elegans. Genetics 175**:** 709-724.

Martinez-Lopez, N., M. Garcia-Macia, S. Sahu, D. Athonvarangkul, E. Liebling *et al.*, 2016 Autophagy in the CNS and Periphery Coordinate Lipophagy and Lipolysis in the Brown Adipose Tissue and Liver. Cell Metab 23**:** 113-127.

McCauliff, L. A., A. Langan, R. Li, O. Ilnytska, D. Bose *et al.*, 2019 Intracellular cholesterol trafficking is dependent upon NPC2 interaction with lysobisphosphatidic acid. Elife 8.

Medoh, U. N., A. Hims, J. Y. Chen, A. Ghoochani, K. Nyame *et al.*, 2023 The Batten disease gene product CLN5 is the lysosomal bis(monoacylglycero)phosphate synthase. Science 381**:** 1182-1189.

Miao, R., M. Li, Q. Zhang, C. Yang and X. Wang, 2020 An ECM-to-Nucleus Signaling Pathway Activates Lysosomes for C. elegans Larval Development. Dev Cell 52: 21-37 e25.

Oka, T., T. Toyomura, K. Honjo, Y. Wada and M. Futai, 2001 Four subunit a isoforms of Caenorhabditis elegans vacuolar H+-ATPase. Cell-specific expression during development. J Biol Chem 276: 33079-33085.

Pechuk, V., G. Goldman, Y. Salzberg, A. H. Chaubey, R. A. Bola et al., 2022 Reprogramming the topology of the nociceptive circuit in C. elegans reshapes sexual behavior. Curr Biol 32: 4372-4385 e4377.

Philbrook, A., M. P. O'Donnell, L. Grunenkovaite and P. Sengupta, 2024 Cilia structure and intraflagellar transport differentially regulate sensory response dynamics within and between C. elegans chemosensory neurons. PLoS Biol 22: e3002892.

Rohlfing, A. K., Y. Miteva, S. Hannenhalli and T. Lamitina, 2010 Genetic and physiological activation of osmosensitive gene expression mimics transcriptional signatures of pathogen infection in C. elegans. PLoS One 5: e9010.

Rohlfing, A. K., Y. Miteva, L. Moronetti, L. He and T. Lamitina, 2011 The Caenorhabditis elegans mucin-like protein OSM-8 negatively regulates osmosensitive physiology via the transmembrane protein PTR-23. PLoS Genet 7: e1001267.

Saha, P., J. L. Shumate, J. G. Caldwell, N. Elghobashi-Meinhardt, A. Lu et al., 2020 Inter-domain dynamics drive cholesterol transport by NPC1 and NPC1L1 proteins. Elife 9.

Savini, M., A. Folick, Y. T. Lee, F. Jin, A. Cuevas et al., 2022 Lysosome lipid signalling from the periphery to neurons regulates longevity. Nat Cell Biol 24: 906-916.

Simmons, A., I. A. Strigo, S. C. Matthews, M. P. Paulus and M. B. Stein, 2009 Initial evidence of a failure to activate right anterior insula during affective set shifting in posttraumatic stress disorder. Psychosom Med 71: 373-377.

Singh, K., M. Y. Chao, G. A. Somers, H. Komatsu, M. E. Corkins et al., 2011 C. elegans Notch signaling regulates adult chemosensory response and larval molting quiescence. Curr Biol 21: 825-834.

Singh, S., U. E. Dransfeld, Y. A. Ambaw, J. Lopez-Scarim, R. V. Farese, Jr. et al., 2024 PLD3 and PLD4 synthesize S,S-BMP, a key phospholipid enabling lipid degradation in lysosomes. Cell 187: 6820-6834 e6824.

Sinner, M. P., F. Masurat, J. J. Ewbank, N. Pujol and H. Bringmann, 2021 Innate Immunity Promotes Sleep through Epidermal Antimicrobial Peptides. Curr Biol 31: 564-577 e512.

Solomon, A., S. Bandhakavi, S. Jabbar, R. Shah, G. J. Beitel et al., 2004 Caenorhabditis elegans OSR-1 regulates behavioral and physiological responses to hyperosmotic environments. Genetics 167: 161-170.

Sternberg, P. W., K. Van Auken, Q. Wang, A. Wright, K. Yook et al., 2024 WormBase 2024: status and transitioning to Alliance infrastructure. Genetics 227.

Stover, P. J., M. S. Field, M. L. Andermann, R. L. Bailey, R. L. Batterham et al., 2023 Neurobiology of eating behavior, nutrition, and health. J Intern Med 294: 582-604.

Suarez-Roca, H., R. Y. Klinger, M. V. Podgoreanu, R. R. Ji, M. I. Sigurdsson et al., 2019 Contribution of Baroreceptor Function to Pain Perception and Perioperative Outcomes. Anesthesiology 130: 634-650.

Tatum, M. C., F. K. Ooi, M. R. Chikka, L. Chauve, L. A. Martinez-Velazquez et al., 2015 Neuronal serotonin release triggers the heat shock response in C. elegans in the absence of temperature increase. Curr Biol 25: 163-174.

Taylor, R. C., and A. Dillin, 2013 XBP-1 is a cell-nonautonomous regulator of stress resistance and longevity. Cell 153**:** 1435-1447.

Taylor, S. R., G. Santpere, A. Weinreb, A. Barrett, M. B. Reilly *et al.*, 2021 Molecular topography of an entire nervous system. Cell 184**:** 4329-4347 e4323.

Troemel, E. R., J. H. Chou, N. D. Dwyer, H. A. Colbert and C. I. Bargmann, 1995 Divergent seven transmembrane receptors are candidate chemosensory receptors in C. elegans. Cell 83**:** 207-218.

Urso, S., M. Comly, J. A. Hanover and T. Lamitina, 2020 The O-GlcNAc transferase OGT is a conserved and essential regulator of the cellular and organismal response to hypertonic stress. BioRxiv.

Urso, S. J., A. Sathaseevan, W. Brent Derry and T. Lamitina, 2023 Regulation of the hypertonic stress response by the 3' mRNA cleavage and polyadenylation complex. Genetics 224.

Veroli, M. V., and T. Lamitina, 2023 Endogenous gpdh-1 transcriptional reporters as new tools for the study of the osmotic stress response. MicroPubl Biol 2023.

Watts, J. S., H. F. Harrison, S. Omi, Q. Guenthers, J. Dalelio *et al.*, 2020 New Strains for Tissue-Specific RNAi Studies in Caenorhabditis elegans. G3 (Bethesda) 10**:** 4167-4176.

Wheeler, J. M., and J. H. Thomas, 2006 Identification of a novel gene family involved in osmotic stress response in Caenorhabditis elegans. Genetics 174**:** 1327-1336.

Wu, Y. C., I. Beets, B. W. Fox, D. Fajardo Palomino, L. Chen *et al.*, 2025 Intercellular sphingolipid signaling mediates aversive learning in C. elegans. Curr Biol.

Zhong, Y., L. J. Gu, X. G. Sun, S. H. Yang and X. H. Zhang, 2014 Comprehensive analysis of patched domain-containing genes reveals a unique evolutionary pattern. Genet Mol Res 13**:** 7318-7331.