Joint impact on thermotolerance of *Saccharomyces* species divergence in mitochondrial and nuclear genomes

Jun-Ting Johnson Wang[1,2], Ping Ling Priscilla Ng[1,3], Maceo E. Powers[1,4], Rachel B. Brem[1*]

[1]Department of Plant and Microbial Biology, University of California, Berkeley, Berkeley, CA, 94720

[2]Graduate Group in Comparative Biochemistry, University of California, Berkeley, Berkeley, CA, 94720

[3]Current address: BillionToOne, Menlo Park, CA, 94025

[4]Current address: Laboratory of Viral Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, 20892

*To whom correspondence should be addressed (rbrem@berkeley.edu)

Keywords: Trait variation, evolution, mitochondrial divergence, positive epistasis

Abstract

Many traits of interest in biology evolved long ago and are fixed in a particular species, distinguishing it from other sister taxa. Elucidating the mechanisms underlying such ancient traits has been a central goal for evolutionary biologists. The yeast *Saccharomyces cerevisiae* is unique among its relatives for its ability to thrive at high temperature, via a genetic architecture that remains incompletely understood. We sought to understand the contribution of species variation in mitochondrial DNA to yeast thermotolerance. We used mitochondrial transgenesis to show that *S. cerevisiae* mitotypes were sufficient for a partial boost to thermotolerance and respiration in the *S. paradoxus* background. These mitochondrial alleles worked best when the background also harbored a pro-thermotolerance nuclear genotype, attesting to positive epistasis between the two genomes. The benefits of *S. cerevisiae* alleles in terms of respiration and growth at high temperature came at the cost of worse performance in cooler conditions. Together, our results establish this system as a case in which mitoalleles have fitness benefits and work well in multiple backgrounds; given the high mutation rate of the mitochondrial compartment, they were also likely extra easy for the ancestral population to acquire. We thus propose a broader model in which mitochondrial variants may prove to be especially good candidates in the search for genetic mechanisms of the adaptive process.

Introduction

A central aim in evolutionary genetics is to figure out how organisms acquire new traits. In the study of within-species variation, a rich literature has provided insights into the mechanisms of local adaptation in the wild (Elena 2017; Kraemer and Boynton 2017; Sork 2017; Rees et al. 2020; Bazzicalupo 2022). Yet many characters of basic and applied interest evolved long ago and are fixed in a particular species, distinguishing it from unaffected relatives. Against a backdrop of years of genomic and experimental pursuit (Allen Orr 2001; Nikolov and Tsiantis 2015; Weiss and Brem 2019), as a field we still know surprisingly little about how evolution engineers and refines phenotypes over long timescales, with the widest knowledge gap in the case of polygenic architectures. Shedding light on these processes, by dissecting the genetics of ancient traits, remains a key challenge for the field.

Sequence variants in mitotype, the DNA of the mitochondrial organelle, have emerged as an important part of the genetic architecture of naturally varying traits. Mitotype variants have been directly implicated in trait innovation, both independently and in concert with nuclear genomes (Melvin and Ballard 2006; Tranah 2011; Houtkooper et al. 2013; Sullivan and Chandel 2014; Sun et al. 2019; Cao et al. 2022; Meng et al. 2022; Oppong et al. 2022; Quéméneur et al. 2022; Parra et al. 2023; Moran et al. 2024). Though small in length, the mitochondrial genome has the potential for an outsize influence on trait variation, given its increased mutation rate and large mutational target of the multi-copy mitochondrial genome relative to nuclear chromosomes (Pakendorf and Stoneking 2005; Chou and Leu 2015; Ferreira and Rodriguez 2024).

Saccharomyces budding yeasts diverged from a common ancestor ~20 million years ago (Kellis et al. 2003) and have served for decades as a model for the study of interspecific trait variation, including in metabolism, genome content, cell cycle, and reproductive isolation (Hou et al. 2014; Roop et al. 2016; Bozdag et al. 2021; Fredericks et al. 2021; Lupo et al. 2021; Marsit et al. 2021; Smukowski Heil et al. 2021; Swamy et al. 2022; Crandall et al. 2023; Forejt and Jansa 2023; Peris et al. 2023). A long-standing subset of this field has focused on species-unique traits driven by mitochondrial genomes (Sulo et al. 2003; Wolff et al. 2014; De Chiara et al. 2020; Hewitt et al. 2020; Hénault et al. 2022), with particularly detailed mechanistic validation in studies of speciation (Lee et al. 2008; Chou et al. 2010; Mário et al. 2015; Jhuang et al. 2017). Such findings from ancient traits serve as a complement to studies of within-species mitotype

variation in yeast (Wolters et al. 2015; Leducq et al. 2017; Wolters et al. 2018; Vijayraghavan et
al. 2019; Vijayraghavan et al. 2019; Nguyen et al. 2020). A key model for ancient trait variation
in the *Saccharomyces* system is temperature response, as the best-studied yeast species, *S.*
cerevisiae, is unique within the *Saccharomyces* clade for its ability to thrive at high temperature
(Sweeney et al. 2004; Gonçalves et al. 2011; Salvadó et al. 2011), whereas others in the clade
exhibit cryotolerance (Paget et al. 2014; Pinto et al. 2025). A growing literature has implicated
nuclear loci in species differences in thermotolerance (Weiss et al. 2018; AlZaben et al. 2021;
Walunjkar et al. 2024) and cold tolerance (Paget et al. 2014; Pinto et al. 2025). The contribution
of mitochondrial genome variation to this trait has also been suggested, based on analyses of
interspecies hybrids (Baker et al. 2019; Li et al. 2019; Hewitt et al. 2020); exactly what the
divergent mitotypes of *Saccharomyces* species do phenotypically, and how they exert their
effects, remains incompletely understood.

We set out to deepen our understanding of the phenotypic impact of mitochondrial variation
between *Saccharomyces* species, both in isolation and in conjunction with nuclear loci. We
anticipated that our findings would further illuminate the mechanisms of thermotolerance and
their evolutionary history. We designed an approach that used *S. paradoxus*, a relatively
thermosensitive species sister to *S. cerevisiae,* as a model for the common ancestor on whose
background traits would have arisen as the two species diverged. We manipulated the nuclear
and mitochondrial genome in this system to investigate their contributions to metabolism and
thermotolerance.

79 Results

80 *S. cerevisiae* mitotypes are sufficient to heighten respiratory behaviors in *S. paradoxus*

Previous studies have used yeast interspecies hybrids to establish an association between *S.*
cerevisiae mitochondrial alleles and thermotolerance (Baker et al. 2019; Li et al. 2019; Hewitt et
al. 2020), which parallels the advantage of *S. cerevisiae* purebreds at high temperature, relative
to other species in the clade. We reasoned that investigating mitotype genetics in purebred *S.*
paradoxus could help shed additional light on the mechanisms of evolutionary innovation along
the *S. cerevisiae* lineage. For this purpose, we developed a series of cybrids (cytoplasmic
hybrids), in which we introduced each mitochondrial genome in turn from a suite of *S. cerevisiae*
isolates from different geographical locations, which all grew well in standard conditions and at
high temperature (Figure S1), into a tester strain of *S. paradoxus*, the English oak tree isolate
Z1 (Figure 1). Since they lack the endogenous *S. paradoxus* mitotype, these strains
represented an experimental resource for the study of the phenotypic impact of *S. cerevisiae*
mitochondrial alleles in an isogenic background, and they were viable and stable as expected
(Sulo et al. 2003; Mário et al. 2015).

Before using our cybrids to address thermotolerance *per se*, we first investigated the hypothesis
that respiration, as a proximal function of the mitochondrion and its genome, would be perturbed
by species variation in mitotype. Wild-type *S. cerevisiae* strains as a rule grow better than those
of *S. paradoxus* on glycerol, a non-fermentable carbon source (Figure S2), and in interspecies
hybrid backgrounds, *S. cerevisiae* mitotypes promote growth in this condition (Li et al. 2019). To
explore the impact of *S. cerevisiae* mitochondrial DNA alleles on respiratory growth in a
purebred context, we subjected our cybrids to the liquid growth assay in glycerol media. The
results revealed robust improvement in glycerol growth attributable to *S. cerevisiae* mitotypes in
the *S. paradoxus* background, for three of our *S. cerevisiae* donors (Figure 2A). The fourth
cybrid, harboring mitochondrial DNA from the *S. cerevisiae* vineyard isolate Y55, had a defect in
growth on glycerol (Figure 2A), which echoes previously reported incompatibilities of this
mitotype in other backgrounds (Wolters et al. 2018). Together, these data suggested that pro-

respiratory effects could be a prevalent characteristic of mitochondrial genomes from across the
S. cerevisiae population.

As a more direct measure of metabolic effects driven by *S. cerevisiae* mitochondria DNA, we
established an assay of reduction of the colorimetric substrate 1-(4,5-dimethylthiazol-2-yl)-3,5-
diphenyltetrazolium bromide (MTT) in glucose medium. This signal serves as a reporter of
activities of the electron transport chain, particularly succinate dehydrogenase (Slater et al.
1963; Sánchez and Königsberg 2006; Jain et al. 2018) (though see (Shoemaker et al. 2004)
and (Peng et al. 2005)), and thus is informative as a marker of respiration. A first set of
observations revealed no detectable impact of mitotype on MTT reduction during culture in
liquid glucose medium at 28°C, in our cybrids in the Z1 *S. paradoxus* background (Figure 3A).
We reasoned that the metabolic effect of *S. cerevisiae* mitotypes might be better resolved at
higher temperatures, in which yeast cells respire more avidly (Rikhvanov et al. 2001).
Consistent with this notion, in culture at 39°C in glucose medium, we observed increased MTT
reduction in all cybrids except for that harboring *S. cerevisiae* Y55 mitochondrial DNA, relative
to wild-type Z1 *S. paradoxus* with its endogenous mitochondrial genome (Figure 3B). Together,
these results established that in the *S. paradoxus* background, *S. cerevisiae* mitotypes can be
sufficient for increased growth and metabolic behaviors that are signatures of respiration.

S. cerevisiae* mitotypes boost thermotolerance in *S. paradoxus

We now turned to high-temperature growth proper, as an additional phenotype that we
hypothesized would be modulated by mitotype in our yeast species comparison. To test the
latter prediction, we measured carrying capacity during growth in liquid glucose medium at
39°C, for each of our cybrid strains harboring *S. cerevisiae* mitochondrial DNA in Z1 *S.*
paradoxus. Results revealed a 10% to 70% improvement in accumulated biomass in this assay,
relative to wild-type *S. paradoxus,* in all cybrids except the one for which *S. cerevisiae* Y55 was
the mitochondrial donor (Figure 2A). Thus, *S. cerevisiae* mitotypes were sufficient to confer
thermotolerance benefits, in some cases sizeable ones, in *S. paradoxus*. After 24 hours of
incubation at 39°C, only *S. cerevisiae* wild-type cells retained viability, whereas cultures of all
strains in the *S. paradoxus* background harbored very few viable cells (Figure S3). We conclude
that in our cybrids, *S. cerevisiae* mitochondrial DNAs enable additional cell divisions during
high-temperature exposure, though failures of factors encoded by the *S. paradoxus* nuclear
genome ultimately prevent further growth and kill cells outright.

Thermotolerance phenotypes as they varied across our panel of strains correlated tightly with
the respiratory behaviors we had documented in assays of growth in the non-fermentable
carbon source glycerol (Figures 2A and 2B) and of MTT reduction (Figure 3). That is, among our
strains, those performing best in respiratory contexts were the most thermotolerant, and the
worst in respiratory assays grew the worst at high temperature. These results are most
consistent with a causal relationship between the traits, whereby the *S. cerevisiae* mitotypes
that foster better thermotolerance do it by improving respiration—a metabolic underpinning for
the capacity to grow under high-temperature challenge.

Antagonistic pleiotropy by *S. cerevisiae* mitotypes in temperate conditions and ethanol

Previous studies using interspecies hybrids have reported a defect in cold temperatures
attributable to *S. cerevisiae* mitotypes relative to those of *S. paradoxus* or other relatives (Baker
et al. 2019; Li et al. 2019; Hewitt et al. 2020). To test whether such effects would be
recapitulated in purebred backgrounds, for each of our Z1 *S. paradoxus* derivatives carrying *S.*
cerevisiae mitochondrial DNA, we assayed the carrying capacity after culture 23°C in liquid
glucose medium, for comparison against a 28°C control and the high-temperature condition
(39°C), also with glucose as a carbon source. The results revealed compromised carrying

capacity at the lower temperatures in the cybrids that performed the best at high temperature
(Figure 2A). This set up an almost perfect anticorrelation between the growth patterns of high
and moderate temperatures across strains (Figure 2C), mirroring the observation we had made
with respiratory behaviors (Figures 2 and 3). Interestingly, a similar principle held true in assays
using ethanol as the carbon source, where again the least thermotolerant strains grew the best
(Figures 2A and 2D). Together, our growth and respiration data make clear that *S. cerevisiae*
mitotypes have specialized to heat stress at the expense of performance at lower temperatures
and during ethanol utilization.

161 Positive epistasis between mitochondria and nuclear genomes in terms of
162 thermotolerance

During the evolution of *S. cerevisiae* as a species, the pro-thermotolerance mitotypes we study
here would have arisen alongside changes in nuclear genes which could modify their effects.
We sought to shed light on these relationships by modeling them in the *S. paradoxus*
background. To this end, we introduced *S. cerevisiae* mitochondrial genomes into a derivative of
S. paradoxus Z1 that also harbored nuclear alleles from the DBVPG1373 Dutch soil strain of *S.*
cerevisiae, at eight loci which contribute to thermotolerance (*AFG2, APC1, CEP3, DYN1, ESP1,*
MYO1, SCC2, and *TAF2*, most of which encode cell division factors; (AlZaben et al. 2021)). In
assays of carrying capacity at 39°C, the combination of *S. cerevisiae* alleles in the nuclear and
mitochondrial genomes together conferred a two-fold improvement relative to wild-type *S.*
paradoxus (Figure 4, squares). No such effect was detectable in 28°C cultures (Figure S4). This
represented an increase in cell divisions early in the treatment, as no cybrids harboring *S.*
cerevisiae nuclear alleles maintained viability after 24 hours at 39°C (Figure S5), paralleling
results from cybrids in the wild-type *S. paradoxus* nuclear background (Figure S3). Advantages
in carrying capacity at 39°C manifested regardless of the origin of the mitotype: mitochondrial
DNA from each of the four *S. cerevisiae* strain donors yielded identical high-temperature growth
in *S. paradoxus* in the presence of the nuclear thermotolerance genes from *S. cerevisiae,*
including the Y55 mitotype that had compromised the phenotype in wild-type *S. paradoxus*
(Figures 2A and 4). Thus, the latter incompatibility was resolved by *S. cerevisiae* alleles at our
focal nuclear genes.

Also evident from our nuclear-mitochondrial transgenic strain panel was a pattern of positive
epistasis. That is, the impact of each *S. cerevisiae* mitotype on thermotolerance was greater in
the *S. paradoxus* derivative also harboring nuclear alleles from *S. cerevisiae* than in otherwise
wild-type *S. paradoxus* (Figure 4; compare circles and squares). No such dependence was
detectable at 28°C (Figure S4). Thus, mitochondrial DNA from distinct *S. cerevisiae* donors,
including Y55, all achieved peak performance at high temperature in the presence of *S.*
cerevisiae nuclear genes. This effect did not reach significance for the *S. cerevisiae* mitotype
that promoted thermotolerance most strongly on its own in the *S. paradoxus* background, from
DBVPG1373; it benefited the least from the addition of DBVPG1373 alleles of nuclear genes
(Figure 4, magenta). Relative to this transgenic, strains harboring other *S. cerevisiae* mitotypes,
which had less impact on their own, achieved the same carrying capacity at 39°C in the
presence of DBVPG1373 nuclear alleles (Figure 4, blue, green, and orange). These results are
most consistent with a ceiling beyond which any *S. cerevisiae* alleles of the mitochondrial
genome, even when combined with *S. cerevisiae* alleles of our eight focal nuclear genes, are
unable to improve thermotolerance, in the face of defects accruing from other *S. paradoxus* loci.
Nonetheless, our data make clear that pro-thermotolerance alleles of mitochondrial DNAs and
nuclear genes from *S. cerevisiae* together contribute to the phenotype in a manner that exceeds
the sum of their parts.

Discussion

A key goal of comparative genetics is to understand how evolution builds traits over long
timescales. In this work, we have leveraged tractable *Saccharomyces* as a testbed for the study
of interspecies trait genetics and evolution focused on the mitochondrial genome. Given the
evidence that *S. cerevisiae* thermotolerance is the product of positive selection (Weiss et al.
2018; Abrams et al. 2021), it serves as a useful model of the evolutionary genetics of adaptation
over deep divergences. Our findings complement previous studies of mitotype effects in this
system in interspecies hybrids (Baker et al. 2019; Li et al. 2019; Hewitt et al. 2020) by tracing
the phenotypes conferred by *S. cerevisiae* mitochondrial alleles in purebred backgrounds,
including their genetic dependencies.

Our results shed new light on the genetic architecture and evolution of thermotolerance. Our
finding of a 70% boost in the trait by an *S. cerevisiae* mitotype outstrips the effect of any one of
the nuclear loci previously reported in this system (Weiss et al. 2018). That said, not all *S.*
cerevisiae mitotypes had a benefit of this magnitude, as might be expected given the extensive
precedent for polymorphism among mitochondrial genomes and their effects on human disease
(Wallace 2015) as well as traits in animals (Tsai and St. John 2016; Brand et al. 2024; Serrano
et al. 2024) and plants (Dubrovina and Kiselev 2016; Adhikari et al. 2019; Jo et al. 2019). Some
of the *S. cerevisiae* mitotypes we study here could be ancient, if each was part of a soft
selective sweep in the ancestral population when the trait arose, of the kind reported recently in
a plant selfing system (Bergero et al. 2019). Perhaps more plausibly, *S. cerevisiae* mitochondrial
polymorphisms could be the product of recent mutations on a background that underwent a
hard sweep long ago.

Our data also make clear that alongside largely beneficial marginal effects, all *S. cerevisiae*
mitotypes are even more advantageous at high temperature in the context of pro-
thermotolerance nuclear alleles. To date, such evidence for positive epistasis has been at a
premium in the literature describing adaptive loci from the wild (Nelson and Grishin 2018;
Syenina et al. 2020; Stern et al. 2022). This may be in part owing to a heavier emphasis in the
field on the evolutionarily difficult repacking of a given protein interior to achieve fitness
(Weinreich et al. 2005; Starr and Thornton 2016), which could be more constrained than
combinations of adaptive alleles of multiple proteins. But even the unlinked nuclear genes
underlying S. *cerevisiae* thermotolerance do not exhibit positive epistasis among themselves
(though their combinations do boost the trait; AlZaben et al. 2021). As such, our data point to
mitoalleles, during the evolution of *S. cerevisiae* thermotolerance, as candidate cases of
especially easily accessible (Liao et al. 2022) and highly beneficial mutations collected by the
ancestral population on the trajectory toward its final phenotype, the so-called low-hanging fruit
(Lenski 2017) of the adaptive process.

Though our data leave open the precise mechanism for these benefits, we propose that *S.*
cerevisiae mitotypes produce extra energy and metabolites (Malecki et al. 2020) during thermal
stress via enhanced respiration, relative to the putative ancestral behaviors of mitochondria from
S. paradoxus. If so, an excess of basic building blocks would support the capacity of *S.*
cerevisiae nuclear genes to confer their effects—those we study here govern housekeeping
functions, namely cell division and transcription/translation. Indeed, evidence for epistasis
between mitochondrial and nuclear genomes is rife in non-adaptive contexts, including
reproductive barriers and aging and disease (Ellison and Burton 2008; Arnqvist et al. 2010;
Tranah 2011; Joseph et al. 2013; Lovell et al. 2015; Roux et al. 2016; Loewen and Ganetzky
2018; Wolters et al. 2018; Sujkowski et al. 2019; Nguyen et al. 2020; Bushel et al. 2022; Biot-
Pelletier et al. 2023; Nguyen et al. 2023; Ibrahim et al. 2024 Sep 3). For *S. cerevisiae*
thermotolerance, the emerging model from our work and that of others is that evolution can use
mitonuclear interactions for gains of function, as traits are built from variants in both genome
compartments.

Likewise, the growth defects we have seen by cybrids at lower temperatures dovetail with results from studies of *S. cerevisiae* nuclear thermotolerance loci and those using interspecies backgrounds (Baker et al. 2019; Li et al. 2019; Hewitt et al. 2020; AlZaben et al. 2021). As a general rule, evolutionary tradeoffs are routine in the adaptation literature, from prokaryotes (Abram et al. 2021; Backman et al. 2024; Dalldorf et al. 2024) to animals (Ou et al. 2020; Ackermans 2023; Metzler et al. 2023) and plants (Gao et al. 2022; He et al. 2022; Kurepa and Smalle 2023). For yeast in particular, the exact mechanism of antagonistic pleiotropy exhibited by thermotolerance loci in cooler conditions remains unclear. Exciting recent work has revealed that *S. cerevisiae* proteins, including those encoded by mitochondria, are more thermostable than those of sister species, establishing the likely biochemical underpinning of the trait (Walunjkar et al. 2024). But it may not follow that the *S. cerevisiae* alleles of the loci we study here compromise protein structure *per se* at low temperatures (Yang et al. 2015; Cui et al. 2024). Amino acid changes unique to *S. cerevisiae* could well have tuned the temperature optima of many enzymes, as appears to be the case for the thermotolerance gene *DYN1* (Hong et al. 2016). Further work will be required to pinpoint exactly what biochemical functionality *S. cerevisiae* gave up as it acquired pro-thermotolerance alleles. Even as they stand, however, the genetics of this system help tell a story of a complex balancing act by evolution along the *S. cerevisiae* lineage, as nuclear and mitochondrial variants worked together to convert a thermosensitive ancestor to a high-temperature specialist.

Deta availability statement

The authors affirm that all data necessary for confirming the conclusion of the article are present within the article, figures, and tables. Strains are available upon request.

Acknowledgements

The authors thank members of the Brem lab, Nick Hoeffner (Stanford University), and Jun-Yi Leu (Academia Sinica, Taiwan) for experimental support and helpful discussions. This work was supported by NIH 2R01GM120430 to R.B.B.

Conflicts of interest

No conflicts of interest.

Figure captions

Figure 1. Experimental design and isolates for genetic dissection of *Saccharomyces* species variation in thermotolerance. (A) Wild-type Z1 *S. paradoxus* (purple) is used as the background for a cybrid (right) harboring the endogenous nuclear genome (N) and the mitochondrial genome from a *S. cerevisiae* donor (orange). **(B)** Strains used in this work: Sp, *S. paradoxus* Z1; Sc1 to Sc4, *S. cerevisiae* DBVPG1373, YPS128, DBVPG6044 and Y55, respectively.

Figure 2. Condition-dependent growth and tradeoffs across *Saccharomyces* species and cybrids. (A) In the main plot, each cell reports cell density, as a mean across replicates ($n \geq 3$), after 24 hours of growth in liquid culture in the respective condition, with glucose as the carbon source unless otherwise indicated. Each column reports results from wild-type *S. paradoxus* (Sp) or an isogenic cybrid harboring the mitochondrial genome from a *S. cerevisiae* donor (Scmt) labeled as in Figure 1. Colors report row-normalized z-score. In the inset, data are as in the main figure with the addition of wild-type *S. cerevisiae* DBVPG1373 (Sc). **(B-D)** Shown are reanalyses of cell density measurements across strains from the cultures and conditions in **(A)**, with respective Spearman coefficients of -0.8, 1, and 0.67. In a given panel, each dot reports median across replicates ($n \geq 3$), and error bars report standard deviations. Strain labels are as in **(A)**. *, two-tailed Mann-Whitney $p < 0.05$. Raw cell density measurements, Spearman correlation coefficients, and statistical analyses are reported in Table S2.

Figure 3. *S. cerevisiae* mitotypes boost respiration in the *S. paradoxus* nuclear background at 39°C. In each panel, in the main plot, the *y*-axis reports measurements of the reduction of 1-(4,5-dimethylthiazol-2-yl)-3,5-diphenyltetrazolium bromide after 24 hours of growth of a given strain in liquid culture with glucose as the carbon source at one temperature, normalized to the respective measurement wild-type *S. paradoxus*. In a given column, each point reports results from one biological replicate and the bar height reports the average. The *x*-axis denotes strains: wild-type *S. paradoxus* (Sp) or cybrids the mitochondrial genome from a *S. cerevisiae* donor (Scmt) labeled as in Figure 1. In the inset, data are as in the main figure with the addition of wild-type *S. cerevisiae* DBVPG1373 (Sc). **(A),** 28°C; **(B)**, 39°C. *, one-sample, two-tailed Wilcoxon $p < 0.05$. Raw data and statistical analyses are reported in Table S2.

Figure 4. Epistasis between nuclear and mitochondrial *Saccharomyces* species variants that impact thermotolerance. Each panel reports results from wild-type *S. paradoxus* Z1 (Sp) and isogenic derivatives the mitochondrial genome from a *S. cerevisiae* donor (Scmt) labeled as in Figure 1; eight nuclear thermotolerance genes from *S. cerevisiae* DBVPG1373 (8X); or both. The *y*-axis reports cell density after 24 hours of growth in liquid culture at 39°C, with glucose as the carbon source. In a given column, transparent points report replicates ($n \geq 6$) and the bold bordered point reports their average. *, two-way ANOVA $p < 0.01$ for mito-nuclear interaction. Raw cell density measurements and statistical analyses are reported in Table S2.

Supplementary figure captions

Figure S1. Stark growth differences between *S. cerevisiae* and *S. paradoxus* at high temperature. In a given panel, the *y*-axis reports cell density after 24 hours of growth in liquid culture of a wild-type homozygous diploid *Saccharomyces* strain in one temperature condition, with glucose as the carbon source. Points reports technical replicates (n = 3) and bar heights report means. Sc1, Sc2, Sc3 and Sc4 denote *S. cerevisiae* DBVPG1373, YPS128, DBVPG6044 and Y55, respectively. Sp1, Sp2, Sp3 and Sp4 denote *S. paradoxus* KPN3828, UFRJ50816, A4 and Z1, respectively. **(A)** 28°C; **(B)** 39°C. *, two-tailed Mann-Whitney $p < 0.05$. Raw cell density measurements and statistical analyses are reported in Table S2.

Figure S2. Wild-type strain and species variation in growth on glycerol. In a given panel, each bar reports results from one wild-type *S. cerevisiae* or *S. paradoxus* strain grown in liquid culture with glycerol as the carbon source from (Warringer et al. 2011). The *y*-axis reports relative growth rate **(A)** and efficiency **(B)**. *, two-talied Mann-Whitney $p < 0.05$.

Figure S3. Chronic 39°C treatment is lethal to *S. paradoxus* and to its cybrids. Shown are results of plating of the indicated strains after 24 hours of culture in one temperature condition, with glucose as the carbon source, followed by incubation of plates at 28°C for two days. Strains were wild-type *S. cerevisiae* (Sc), wild-type *S. paradoxus* (Sp), or a cybrid in the *S. paradoxus* background harboring the mitochondrial genome from a *S. cerevisiae* donor (Scmt) labeled as in Figure 1 of the main text. Raw cell density measurements and statistical analyses are reported in Table S2.

Figure S4. No detectable epistasis between nuclear and mitochondrial *Saccharomyces* species variants that impact 28°C growth. Labels are as in Figure 4. The *y*-axis reports cell density after 24 hours of growth in liquid culture at 28°C, with glucose as the carbon source. In a given column, transparent points report replicates ($n \geq 6$) and the bold bordered point reports their average. ns, two-way ANOVA $p > 0.05$ for mito-nuclear interaction. Raw cell density measurements and statistical analyses are reported in Table S2.

Figure S5. Chronic 39°C treatment is lethal to *S. paradoxus* harboring *S. cerevisiae* nuclear thermotolerance loci and to its cybrids. Labels are as in **Figure S3** except that the strains were wild-type *S. cerevisiae* (Sc), wild-type *S. paradoxus* (Sp), *S. paradoxus* harboring eight nuclear loci from *S. cerevisiae* (8X), or a cybrid in the 8X background harboring the mitochondrial genome from a *S. cerevisiae* donor (Scmt) labeled as in Figure 1 of the main text. Raw cell density measurements and statistical analyses are reported in Table S2.

Supplementary table captions

Table S1. Strains used in this work.

Table S2. Growth data and statistics. D1373, YPS128, D6044 and Y55 indicate *S. cerevisiae* strains DBVPG1373, YPS128, DBVPG6044 and Y55, respectively; Sp, *S. paradoxus* Z1. The first sheet contains all OD_{600} measurements of each strain at 39°C in standard glucose medium. The second sheet reports the statistical analysis of the OD_{600} measurements of the cybrids at 39°C compared to wild-type *S. paradoxus* or the *S. paradoxus* harboring *S. cerevisiae* nuclear loci (8X) by Mann-Whitney U test. The third sheet contains all OD_{600} measurements of each strain at 28°C in standard glucose medium. The fourth sheet contains all OD_{600} measurements of each strain at 28°C in medium with glycerol as a carbon source. The fifth sheet contains all OD_{600} measurements of each strain at 28°C in medium with ethanol as a carbon source. The sixth sheet contains all OD_{600} measurements of each strain at 23°C in standard glucose medium. The seventh sheet reports the *p*-values of two-way ANOVA of the impact of the interaction between *S. cerevisiae* mitotypes and the 8 *S. cerevisiae* loci in the *S. paradoxus* background on high temperature growth (Figure 4). The eighth sheet reports the raw readouts of the MTT assay at 28°C and 39°C. The ninth sheet reports the *p*-values of the normalized MTT measurements by one-sample Wilcoxon with a hypothesized value of 1. The tenth sheet reports the Spearman correlation coefficients and respective *p*-values of Figures 2B, 2C and 2D. The eleventh sheet reports the OD_{600} measurements of wild-type *S. cerevisiae* and *S. paradoxus* strains at 28°C and 39°C in standard glucose medium with the statistical analysis (Figure S1). The twelfth sheet reports the *p*-values of glycerol growth efficiency and rate of the wild-type *S. cerevisiae* and *S. paradoxus* strains by two-tailed Mann-Whitney U test (Warringer et al. 2011).

References

Abram F, Arcari T, Guerreiro D, O'Byrne CP. 2021. Evolutionary trade-offs between growth and survival: The delicate balance between reproductive success and longevity in bacteria. In: Advances in Microbial Physiology. Vol. 79. Elsevier. p. 133–162. https://linkinghub.elsevier.com/retrieve/pii/S006529112100014X.

Abrams MB, Dubin CA, AlZaben F, Bravo J, Joubert PM, Weiss CV, Brem RB. 2021. Population and comparative genetics of thermotolerance divergence between yeast species. G3. 11(7):jkab139. doi:10.1093/g3journal/jkab139.

Ackermans NL. 2023. Neurobiological tradeoffs of headbutting bovids. Trends in Neurosciences. 46(11):898–900. doi:10.1016/j.tins.2023.08.004.

Adhikari B, Caruso CM, Case AL. 2019. Beyond balancing selection: frequent mitochondrial recombination contributes to high-female frequencies in gynodioecious *Lobelia siphilitica* (Campanulaceae). New Phytologist. 224(3):1381–1393. doi:10.1111/nph.16136.

Allen Orr H. 2001. The genetics of species differences. Trends in Ecology & Evolution. 16(7):343–350. doi:10.1016/S0169-5347(01)02167-X.

AlZaben F, Chuong JN, Abrams MB, Brem RB. 2021. Joint effects of genes underlying a temperature specialization tradeoff in yeast. PLoS Genet. 17(9):e1009793. doi:10.1371/journal.pgen.1009793.

Amine AAA, Liao C-W, Hsu P-C, Opoc FJG, Leu J-Y. 2021. Experimental evolution improves mitochondrial genome quality control in Saccharomyces cerevisiae and extends its replicative lifespan. Current Biology. 31(16):3663-3670.e4. doi:10.1016/j.cub.2021.06.026.

Arnqvist G, Dowling DK, Eady P, Gay L, Tregenza T, Tuda M, Hosken DJ. 2010. Genetic architecture of metabolic rate: environment specific epistasis between mitochondrial and nuclear genes in an insect. Evolution. 64(12):3354–3363. doi:10.1111/j.1558-5646.2010.01135.x.

Backman T, Burbano HA, Karasov TL. 2024. Tradeoffs and constraints on the evolution of tailocins. Trends in Microbiology. 32(11):1084–1095. doi:10.1016/j.tim.2024.04.001.

Baker EP, Peris D, Moriarty RV, Li XC, Fay JC, Hittinger CT. 2019. Mitochondrial DNA and temperature tolerance in lager yeasts. Sci Adv. 5(1):eaav1869. doi:10.1126/sciadv.aav1869.

Bazzicalupo A. 2022. Local adaptation in fungi. FEMS Microbiology Reviews. 46(6):fuac026. doi:10.1093/femsre/fuac026.

Bergero R, Levsen N, Wolff K, Charlesworth D. 2019. Arms races with mitochondrial genome soft sweeps in a gynodioecious plant, Plantago lanceolata. :78254 bytes. doi:10.5061/DRYAD.TH757VG. https://datadryad.org/dataset/doi:10.5061/dryad.th757vg.

Biot-Pelletier D, Bettinazzi S, Gagnon-Arsenault I, Dubé AK, Bédard C, Nguyen THM, Fiumera HL, Breton S, Landry CR. 2023. Evolutionary Trajectories are Contingent on Mitonuclear Interactions. Molecular Biology and Evolution. 40(4):msad061. doi:10.1093/molbev/msad061.

Bozdag GO, Ono J, Denton JA, Karakoc E, Hunter N, Leu J-Y, Greig D. 2021. Breaking a
species barrier by enabling hybrid recombination. Current Biology. 31(4):R180–R181.
doi:10.1016/j.cub.2020.12.038.

Brand JA, Garcia-Gonzalez F, Dowling DK, Wong BBM. 2024. Mitochondrial genetic variation as
a potential mediator of intraspecific behavioural diversity. Trends in Ecology & Evolution.
39(2):199–212. doi:10.1016/j.tree.2023.09.009.

Bushel PR, Ward J, Burkholder A, Li J, Anchang B. 2022. Mitochondrial-nuclear epistasis
underlying phenotypic variation in breast cancer pathology. Sci Rep. 12(1):1393.
doi:10.1038/s41598-022-05148-4.

Cao Y, Vergnes L, Wang Y-C, Pan C, Chella Krishnan K, Moore TM, Rosa-Garrido M, Kimball
TH, Zhou Z, Charugundla S, et al. 2022. Sex differences in heart mitochondria regulate diastolic
dysfunction. Nat Commun. 13(1):3850. doi:10.1038/s41467-022-31544-5.

Chou J-Y, Hung Y-S, Lin K-H, Lee H-Y, Leu J-Y. 2010. Multiple Molecular Mechanisms Cause
Reproductive Isolation between Three Yeast Species. PLoS Biol. 8(7):e1000432.
doi:10.1371/journal.pbio.1000432.

Chou J-Y, Leu J-Y. 2015. The Red Queen in mitochondria: cyto-nuclear co-evolution, hybrid
breakdown and human disease. Front Genet. 6. doi:10.3389/fgene.2015.00187.
http://www.frontiersin.org/Evolutionary_and_Population_Genetics/10.3389/fgene.2015.00187/ab
stract.

Crandall JG, Fisher KJ, Sato TK, Hittinger CT. 2023. Ploidy evolution in a wild yeast is linked to
an interaction between cell type and metabolism. PLoS Biol. 21(11):e3001909.
doi:10.1371/journal.pbio.3001909.

Cubillos FA, Louis EJ, Liti G. 2009. Generation of a large set of genetically tractable haploid and
diploid *Saccharomyces*   strains. FEMS Yeast Research. 9(8):1217–1225. doi:10.1111/j.1567-
1364.2009.00583.x.

Cui S, Furukawa R, Akanuma S. 2024. Insights into the low-temperature adaptation of an
enzyme as studied through ancestral sequence reconstruction. doi:10.1101/2024.09.05.611558.
http://biorxiv.org/lookup/doi/10.1101/2024.09.05.611558.

Dalldorf C, Rychel K, Szubin R, Hefner Y, Patel A, Zielinski DC, Palsson BO. 2024. The
hallmarks of a tradeoff in transcriptomes that balances stress and growth functions. mSystems.
9(7):e00305-24. doi:10.1128/msystems.00305-24.

De Chiara M, Friedrich A, Barré B, Breitenbach M, Schacherer J, Liti G. 2020. Discordant
evolution of mitochondrial and nuclear yeast genomes at population level. BMC Biol. 18(1):49.
doi:10.1186/s12915-020-00786-4.

Dubrovina AS, Kiselev KV. 2016. Age-associated alterations in the somatic mutation and DNA
methylation levels in plants. Plant Biol J. 18(2):185–196. doi:10.1111/plb.12375.

Elena SF. 2017. Local adaptation of plant viruses: lessons from experimental evolution.
Molecular Ecology. 26(7):1711–1719. doi:10.1111/mec.13836.

Ellison CK, Burton RS. 2008. Genotype-dependent variation of mitochondrial transcriptional
profiles in interpopulation hybrids. Proc Natl Acad Sci USA. 105(41):15831–15836.
doi:10.1073/pnas.0804253105.

Ferreira T, Rodriguez S. 2024. Mitochondrial DNA: Inherent Complexities Relevant to Genetic
Analyses. Genes. 15(5):617. doi:10.3390/genes15050617.

Forejt J, Jansa P. 2023. Meiotic Recognition of Evolutionarily Diverged Homologs:
Chromosomal Hybrid Sterility Revisited. Molecular Biology and Evolution. 40(4):msad083.
doi:10.1093/molbev/msad083.

Fredericks LR, Lee MD, Crabtree AM, Boyer JM, Kizer EA, Taggart NT, Roslund CR, Hunter SS,
Kennedy CB, Willmore CG, et al. 2021. The Species-Specific Acquisition and Diversification of a
K1-like Family of Killer Toxins in Budding Yeasts of the Saccharomycotina. PLoS Genet.
17(2):e1009341. doi:10.1371/journal.pgen.1009341.

Gao H, Cui J, Liu S, Wang S, Lian Y, Bai Y, Zhu T, Wu H, Wang Y, Yang S, et al. 2022. Natural
variations of ZmSRO1d modulate the trade-off between drought resistance and yield by
affecting ZmRBOHC-mediated stomatal ROS production in maize. Molecular Plant.
15(10):1558–1574. doi:10.1016/j.molp.2022.08.009.

Gonçalves P, Valério E, Correia C, De Almeida JMGCF, Sampaio JP. 2011. Evidence for
Divergent Evolution of Growth Temperature Preference in Sympatric Saccharomyces Species.
PLoS ONE. 6(6):e20739. doi:10.1371/journal.pone.0020739.

He Z, Webster S, He SY. 2022. Growth–defense trade-offs in plants. Current Biology.
32(12):R634–R639. doi:10.1016/j.cub.2022.04.070.

Hénault M, Marsit S, Charron G, Landry CR. 2022. Hybridization drives mitochondrial DNA
degeneration and metabolic shift in a species with biparental mitochondrial inheritance. Genome
Res. 32(11–12):2043–2056. doi:10.1101/gr.276885.122.

Hewitt SK, Duangrattanalert K, Burgis T, Zeef LAH, Naseeb S, Delneri D. 2020. Plasticity of
Mitochondrial DNA Inheritance and its Impact on Nuclear Gene Transcription in Yeast Hybrids.
Microorganisms. 8(4):494. doi:10.3390/microorganisms8040494.

Hoffman GA, Garrison TR, Dohlman HG. 2002. Analysis of RGS Proteins in Saccharomyces
cerevisiae. In: Methods in Enzymology. Vol. 344. Elsevier. p. 617–631.
https://linkinghub.elsevier.com/retrieve/pii/S0076687902447441.

Hong W, Takshak A, Osunbayo O, Kunwar A, Vershinin M. 2016. The Effect of Temperature on
Microtubule-Based Transport by Cytoplasmic Dynein and Kinesin-1 Motors. Biophysical Journal.
111(6):1287–1294. doi:10.1016/j.bpj.2016.08.006.

Hou J, Friedrich A, de Montigny J, Schacherer J. 2014. Chromosomal Rearrangements as a
Major Mechanism in the Onset of Reproductive Isolation in Saccharomyces cerevisiae. Current
Biology. 24(10):1153–1159. doi:10.1016/j.cub.2014.03.063.

Houtkooper RH, Mouchiroud L, Ryu D, Moullan N, Katsyuba E, Knott G, Williams RW, Auwerx
J. 2013. Mitonuclear protein imbalance as a conserved longevity mechanism. Nature.
497(7450):451–457. doi:10.1038/nature12188.

Ibrahim R, Bahilo Martinez M, Dobson AJ. 2024 Sep 3. Rapamycin's lifespan effect is
modulated by mito-nuclear epistasis in *Drosophila*. Aging Cell.:e14328. doi:10.1111/acel.14328.

Jain AK, Singh D, Dubey K, Maurya R, Mittal S, Pandey AK. 2018. Models and Methods for In
Vitro Toxicity. In: In Vitro Toxicology. Elsevier. p. 45–65.
https://linkinghub.elsevier.com/retrieve/pii/B9780128046678000031.

Jhuang H, Lee H, Leu J. 2017. Mitochondrial–nuclear co-evolution leads to hybrid
incompatibility through pentatricopeptide repeat proteins. EMBO Reports. 18(1):87–101.
doi:10.15252/embr.201643311.

Jo YD, Lee H-Y, Ro N-Y, Kim SH, Kim J-B, Kang B-C, Kang S-Y. 2019. Mitotypes Based on
Structural Variation of Mitochondrial Genomes Imply Relationships With Morphological
Phenotypes and Cytoplasmic Male Sterility in Peppers. Front Plant Sci. 10:1343.
doi:10.3389/fpls.2019.01343.

Joseph B, Corwin JA, Li B, Atwell S, Kliebenstein DJ. 2013. Cytoplasmic genetic variation and
extensive cytonuclear interactions influence natural variation in the metabolome. eLife.
2:e00776. doi:10.7554/eLife.00776.

Kellis M, Patterson N, Endrizzi M, Birren B, Lander ES. 2003. Sequencing and comparison of
yeast species to identify genes and regulatory elements. Nature. 423(6937):241–254.
doi:10.1038/nature01644.

Kraemer SA, Boynton PJ. 2017. Evidence for microbial local adaptation in nature. Molecular
Ecology. 26(7):1860–1876. doi:10.1111/mec.13958.

Kurepa J, Smalle JA. 2023. Plant Hormone Modularity and the Survival-Reproduction Trade-Off.
Biology. 12(8):1143. doi:10.3390/biology12081143.

Leducq J-B, Henault M, Charron G, Nielly-Thibault L, Terrat Y, Fiumera HL, Shapiro BJ, Landry
CR. 2017. Mitochondrial Recombination and Introgression during Speciation by Hybridization.
Molecular Biology and Evolution. 34(8):1947–1959. doi:10.1093/molbev/msx139.

Lee H-Y, Chou J-Y, Cheong L, Chang N-H, Yang S-Y, Leu J-Y. 2008. Incompatibility of Nuclear
and Mitochondrial Genomes Causes Hybrid Sterility between Two Yeast Species. Cell.
135(6):1065–1073. doi:10.1016/j.cell.2008.10.047.

Lenski RE. 2017. Experimental evolution and the dynamics of adaptation and genome evolution
in microbial populations. The ISME Journal. 11(10):2181–2194. doi:10.1038/ismej.2017.69.

Li XC, Peris D, Hittinger CT, Sia EA, Fay JC. 2019. Mitochondria-encoded genes contribute to
evolution of heat and cold tolerance in yeast. Sci Adv. 5(1):eaav1848.
doi:10.1126/sciadv.aav1848.

Liao S, Chen L, Song Z, He H. 2022. The fate of damaged mitochondrial DNA in the cell.
Biochimica et Biophysica Acta (BBA) - Molecular Cell Research. 1869(5):119233.
doi:10.1016/j.bbamcr.2022.119233.

Liti G, Carter DM, Moses AM, Warringer J, Parts L, James SA, Davey RP, Roberts IN, Burt A,
Koufopanou V, et al. 2009. Population genomics of domestic and wild yeasts. Nature.
458(7236):337–341. doi:10.1038/nature07743.

Loewen CA, Ganetzky B. 2018. Mito-Nuclear Interactions Affecting Lifespan and
Neurodegeneration in a *Drosophila* Model of Leigh Syndrome. Genetics. 208(4):1535–1552.
doi:10.1534/genetics.118.300818.

Lovell JT, Mullen JL, Lowry DB, Awole K, Richards JH, Sen S, Verslues PE, Juenger TE, McKay
JK. 2015. Exploiting Differential Gene Expression and Epistasis to Discover Candidate Genes
for Drought-Associated QTLs in *Arabidopsis thaliana*. Plant Cell. 27(4):969–983.
doi:10.1105/tpc.15.00122.

Lupo O, Krieger G, Jonas F, Barkai N. 2021. Accumulation of *cis* - and *trans* -regulatory
variations is associated with phenotypic divergence of a complex trait between yeast species.
G3 Genes|Genomes|Genetics. 11(2):jkab016. doi:10.1093/g3journal/jkab016.

Malecki M, Kamrad S, Ralser M, Bähler J. 2020. Mitochondrial respiration is required to provide
amino acids during fermentative proliferation of fission yeast. EMBO Reports. 21(11):e50845.
doi:10.15252/embr.202050845.

Mário Š, Poláková S, Jatzová K, Sulo P. 2015. Post-zygotic sterility and cytonuclear
compatibility limits in S. cerevisiae xenomitochondrial cybrids. Front Genet. 5.
doi:10.3389/fgene.2014.00454.
http://journal.frontiersin.org/article/10.3389/fgene.2014.00454/abstract.

Marsit S, Hénault M, Charron G, Fijarczyk A, Landry CR. 2021. The neutral rate of whole-
genome duplication varies among yeast species and their hybrids. Nat Commun. 12(1):3126.
doi:10.1038/s41467-021-23231-8.

Melvin RG, Ballard JWO. 2006. Intraspecific variation in survival and mitochondrial oxidative
phosphorylation in wild-caught *Drosophila simulans*. Aging Cell. 5(3):225–233.
doi:10.1111/j.1474-9726.2006.00211.x.

Meng X, Li L, Pascual J, Rahikainen M, Yi C, Jost R, He C, Fournier-Level A, Borevitz J,
Kangasjärvi S, et al. 2022. GWAS on multiple traits identifies mitochondrial ACONITASE3 as
important for acclimation to submergence stress. Plant Physiology. 188(4):2039–2058.
doi:10.1093/plphys/kiac011.

Metzler S, Kirchner J, Grasse AV, Cremer S. 2023. Trade-offs between immunity and
competitive ability in fighting ant males. BMC Ecol Evo. 23(1):37. doi:10.1186/s12862-023-
02137-7.

Moran BM, Payne CY, Powell DL, Iverson ENK, Donny AE, Banerjee SM, Langdon QK, Gunn
TR, Rodriguez-Soto RA, Madero A, et al. 2024. A lethal mitonuclear incompatibility in complex I
of natural hybrids. Nature. 626(7997):119–127. doi:10.1038/s41586-023-06895-8.

Nelson ED, Grishin NV. 2018. Inference of epistatic effects in a key mitochondrial protein. Phys
Rev E. 97(6):062404. doi:10.1103/PhysRevE.97.062404.

Nguyen THM, Sondhi S, Ziesel A, Paliwal S, Fiumera HL. 2020. Mitochondrial-nuclear
coadaptation revealed through mtDNA replacements in Saccharomyces cerevisiae. BMC Evol
Biol. 20(1):128. doi:10.1186/s12862-020-01685-6.

Nguyen THM, Tinz-Burdick A, Lenhardt M, Geertz M, Ramirez F, Schwartz M, Toledano M,
Bonney B, Gaebler B, Liu W, et al. 2023. Mapping mitonuclear epistasis using a novel
recombinant yeast population. PLoS Genet. 19(3):e1010401.
doi:10.1371/journal.pgen.1010401.

Nikolov LA, Tsiantis M. 2015. Interspecies Gene Transfer as a Method for Understanding the
Genetic Basis for Evolutionary Change: Progress, Pitfalls, and Prospects. Front Plant Sci.
6:1135. doi:10.3389/fpls.2015.01135.

Oppong RF, Terracciano A, Picard M, Qian Y, Butler TJ, Tanaka T, Moore AZ, Simonsick EM,
Opsahl-Ong K, Coletta C, et al. 2022. Personality traits are consistently associated with blood
mitochondrial DNA copy number estimated from genome sequences in two genetic cohort
studies. eLife. 11:e77806. doi:10.7554/eLife.77806.

Ou Q, Vannier J, Yang X, Chen A, Mai H, Shu D, Han J, Fu D, Wang R, Mayer G. 2020.
Evolutionary trade-off in reproduction of Cambrian arthropods. Sci Adv. 6(18):eaaz3376.
doi:10.1126/sciadv.aaz3376.

Paget CM, Schwartz J-M, Delneri D. 2014. Environmental systems biology of cold-tolerant
phenotype in Saccharomyces species adapted to grow at different temperatures. Mol Ecol.
23(21):5241–5257. doi:10.1111/mec.12930.

Pakendorf B, Stoneking M. 2005. MITOCHONDRIAL DNA AND HUMAN EVOLUTION. Annu
Rev Genom Hum Genet. 6(1):165–183. doi:10.1146/annurev.genom.6.080604.162249.

Parra M, Libkind D, Hittinger CT, Álvarez L, Bellora N. 2023. Assembly and comparative
genome analysis of a Patagonian *Aureobasidium pullulans* isolate reveals unexpected
intraspecific variation. Yeast. 40(5–6):197–213. doi:10.1002/yea.3853.

Peng L, Wang B, Ren P. 2005. Reduction of MTT by flavonoids in the absence of cells. Colloids
and Surfaces B: Biointerfaces. 45(2):108–111. doi:10.1016/j.colsurfb.2005.07.014.

Peris D, Ubbelohde EJ, Kuang MC, Kominek J, Langdon QK, Adams M, Koshalek JA, Hulfachor
AB, Opulente DA, Hall DJ, et al. 2023. Macroevolutionary diversity of traits and genomes in the
model yeast genus Saccharomyces. Nat Commun. 14(1):690. doi:10.1038/s41467-023-36139-
2.

Pinto J, Balarezo-Cisneros LN, Delneri D. 2025. Exploring adaptation routes to cold
temperatures in the Saccharomyces genus. PLoS Genet. 21(2):e1011199.
doi:10.1371/journal.pgen.1011199.

Piskur J, Rozpedowska E, Polakova S, Merico A, Compagno C. 2006. How did Saccharomyces
evolve to become a good brewer? Trends in Genetics. 22(4):183–186.
doi:10.1016/j.tig.2006.02.002.

Quéméneur J-B, Danion M, Cabon J, Collet S, Zambonino-Infante J-L, Salin K. 2022. The
relationships between growth rate and mitochondrial metabolism varies over time. Sci Rep.
12(1):16066. doi:10.1038/s41598-022-20428-9.

Rees JS, Castellano S, Andrés AM. 2020. The Genomics of Human Local Adaptation. Trends in
Genetics. 36(6):415–428. doi:10.1016/j.tig.2020.03.006.

Reuß O, Vik Å, Kolter R, Morschhäuser J. 2004. The SAT1 flipper, an optimized tool for gene
disruption in Candida albicans. Gene. 341:119–127. doi:10.1016/j.gene.2004.06.021.

Rikhvanov EG, Varakina NN, Rusaleva TM, Rachenko EI, Kiseleva VA, Voinikov VK. 2001. [No
title found]. Microbiology. 70(4):462–465. doi:10.1023/A:1010442429489.

Roop JI, Chang KC, Brem RB. 2016. Polygenic evolution of a sugar specialization trade-off in
yeast. Nature. 530(7590):336–339. doi:10.1038/nature16938.

Roux F, Mary-Huard T, Barillot E, Wenes E, Botran L, Durand S, Villoutreix R, Martin-Magniette
M-L, Camilleri C, Budar F. 2016. Cytonuclear interactions affect adaptive traits of the annual
plant *Arabidopsis thaliana* in the field. Proc Natl Acad Sci USA. 113(13):3687–3692.
doi:10.1073/pnas.1520687113.

Salvadó Z, Arroyo-López FN, Guillamón JM, Salazar G, Querol A, Barrio E. 2011. Temperature
Adaptation Markedly Determines Evolution within the Genus *Saccharomyces*. Appl Environ
Microbiol. 77(7):2292–2302. doi:10.1128/AEM.01861-10.

Sánchez NS, Königsberg M. 2006. Using yeast to easily determine mitochondrial functionality
with 1-(4,5-dimethylthiazol-2-yl)-3,5-diphenyltetrazolium bromide (MTT) assay. Biochem
Molecular Bio Educ. 34(3):209–212. doi:10.1002/bmb.2006.49403403209.

Serrano IM, Hirose M, Valentine CC, Roesner S, Schmidt E, Pratt G, Williams L, Salk J, Ibrahim
S, Sudmant PH. 2024. Mitochondrial haplotype and mito-nuclear matching drive somatic
mutation and selection throughout ageing. Nat Ecol Evol. 8(5):1021–1034. doi:10.1038/s41559-
024-02338-3.

Shoemaker M, Cohen I, Campbell M. 2004. Reduction of MTT by aqueous herbal extracts in the
absence of cells. Journal of Ethnopharmacology. 93(2–3):381–384.
doi:10.1016/j.jep.2004.04.011.

Slater TF, Sawyer B, Sträuli U. 1963. Studies on succinate-tetrazolium reductase systems.
Biochimica et Biophysica Acta. 77:383–393. doi:10.1016/0006-3002(63)90513-4.

Smukowski Heil C, Patterson K, Hickey AS-M, Alcantara E, Dunham MJ. 2021. Transposable
Element Mobilization in Interspecific Yeast Hybrids. Genome Biology and Evolution.
13(3):evab033. doi:10.1093/gbe/evab033.

Sork VL. 2017. Genomic Studies of Local Adaptation in Natural Plant Populations. Journal of
Heredity. 109(1):3–15. doi:10.1093/jhered/esx091.

Starr TN, Thornton JW. 2016. Epistasis in protein evolution. Protein Science. 25(7):1204–1218.
doi:10.1002/pro.2897.

Stern DB, Anderson NW, Diaz JA, Lee CE. 2022. Genome-wide signatures of synergistic
epistasis during parallel adaptation in a Baltic Sea copepod. Nat Commun. 13(1):4024.
doi:10.1038/s41467-022-31622-8.

Sujkowski A, Spierer AN, Rajagopalan T, Bazzell B, Safdar M, Imsirovic D, Arking R, Rand DM,
Wessells R. 2019. Mito-nuclear interactions modify Drosophila exercise performance.
Mitochondrion. 47:188–205. doi:10.1016/j.mito.2018.11.005.

Sullivan LB, Chandel NS. 2014. Mitochondrial reactive oxygen species and cancer. Cancer
Metab. 2(1):17. doi:10.1186/2049-3002-2-17.

Sulo P, Pirek M, Oltesova A, Marinoni G, Pikur J. 2003. The efficiency of functional
mitochondrial replacement in species has directional character. FEMS Yeast Research. 4(1):97–
104. doi:10.1016/S1567-1356(03)00109-0.

Sun J, Duan X, Hoffmann AA, Liu Y, Garvin MR, Chen L, Hu G, Zhou J, Huang H, Xue X, et al.
2019. Mitochondrial variation in small brown planthoppers linked to multiple traits and probably
reflecting a complex evolutionary trajectory. Molecular Ecology. 28(14):3306–3323.
doi:10.1111/mec.15148.

Swamy KBS, Lee H-Y, Ladra C, Liu C-FJ, Chao J-C, Chen Y-Y, Leu J-Y. 2022. Proteotoxicity
caused by perturbed protein complexes underlies hybrid incompatibility in yeast. Nat Commun.
13(1):4394. doi:10.1038/s41467-022-32107-4.

Sweeney J, Kuehne H, Sniegowski P. 2004. Sympatric natural and populations have different
thermal growth profiles. FEMS Yeast Research. 4(4–5):521–525. doi:10.1016/S1567-
1356(03)00171-5.

Syenina A, Vijaykrishna D, Gan ES, Tan HC, Choy MM, Siriphanitchakorn T, Cheng C,
Vasudevan SG, Ooi EE. 2020. Positive epistasis between viral polymerase and the 3´
untranslated region of its genome reveals the epidemiologic fitness of dengue virus. Proc Natl
Acad Sci USA. 117(20):11038–11047. doi:10.1073/pnas.1919287117.

Tranah GJ. 2011. Mitochondrial–nuclear epistasis: Implications for human aging and longevity.
Ageing Research Reviews. 10(2):238–252. doi:10.1016/j.arr.2010.06.003.

Tsai T, St. John JC. 2016. The role of mitochondrial DNA copy number, variants, and haplotypes
in farm animal developmental outcome. Domestic Animal Endocrinology. 56:S133–S146.
doi:10.1016/j.domaniend.2016.03.005.

Vijayraghavan S, Kozmin SG, Strope PK, Skelly DA, Lin Z, Kennell J, Magwene PM, Dietrich
FS, McCusker JH. 2019. Mitochondrial Genome Variation Affects Multiple Respiration and
Nonrespiration Phenotypes in *Saccharomyces cerevisiae*. Genetics. 211(2):773–786.
doi:10.1534/genetics.118.301546.

Wallace DC. 2015. Mitochondrial DNA Variation in Human Radiation and Disease. Cell.
163(1):33–38. doi:10.1016/j.cell.2015.08.067.

Walunjkar N, Lai TY, Akhter N, Miller JH, Bettinger JQ, Marcus E, Phizicky EM, Ghaemmaghami
S, Fay JC. 2024. Pervasive divergence in protein thermostability is mediated by both structural

changes and cellular environments. doi:10.1101/2024.12.09.627561.
http://biorxiv.org/lookup/doi/10.1101/2024.12.09.627561.

Warringer J, Zörgö E, Cubillos FA, Zia A, Gjuvsland A, Simpson JT, Forsmark A, Durbin R,
Omholt SW, Louis EJ, et al. 2011. Trait Variation in Yeast Is Defined by Population History. PLoS
Genet. 7(6):e1002111. doi:10.1371/journal.pgen.1002111.

Weinreich DM, Watson RA, Chao L. 2005. Perspective: Sign epistasis and genetic constraint on
evolutionary trajectories. Evolution. 59(6):1165–1174.

Weiss CV, Brem RB. 2019. Dissecting Trait Variation across Species Barriers. Trends in
Ecology & Evolution. 34(12):1131–1136. doi:10.1016/j.tree.2019.07.013.

Weiss CV, Roop JI, Hackley RK, Chuong JN, Grigoriev IV, Arkin AP, Skerker JM, Brem RB.
2018. Genetic dissection of interspecific differences in yeast thermotolerance. Nat Genet.
50(11):1501–1504. doi:10.1038/s41588-018-0243-4.

Wolff JN, Ladoukakis ED, Enríquez JA, Dowling DK. 2014. Mitonuclear interactions:
evolutionary consequences over multiple biological scales. Philos Trans R Soc Lond B Biol Sci.
369(1646):20130443. doi:10.1098/rstb.2013.0443.

Wolters JF, Charron G, Gaspary A, Landry CR, Fiumera AC, Fiumera HL. 2018. Mitochondrial
Recombination Reveals Mito–Mito Epistasis in Yeast. Genetics. 209(1):307–319.
doi:10.1534/genetics.117.300660.

Wolters JF, Chiu K, Fiumera HL. 2015. Population structure of mitochondrial genomes in
Saccharomyces cerevisiae. BMC Genomics. 16(1):451. doi:10.1186/s12864-015-1664-4.

Yang L-L, Tang S-K, Huang Y, Zhi X-Y. 2015. Low Temperature Adaptation Is Not the Opposite
Process of High Temperature Adaptation in Terms of Changes in Amino Acid Composition.
Genome Biol Evol. 7(12):3426–3433. doi:10.1093/gbe/evv232.

Materials and methods

Strain construction

Strains used in this study are listed in Table S1. For phenotyping in Figure S1, we used wild-type, homozygous diploid *S. cerevisiae*, DBVPG1373, YPS128, DBVPG6044 and Y55, and *S. paradoxus* strains, Z1, UFRJ50816, A4 and KPN3828, from the Saccharomyces Genome Resequencing Project (SGRP) collection (Table S1) (Cubillos et al. 2009). To make haploid strains for the recipients for *S. cerevisiae* mitochondrial DNA, we first generated prototrophic versions of diploid *S. paradoxus* Z1 and the *S. paradoxus* harboring the 8 *S. cerevisiae* loci, designated as 8X (AlZaben et al. 2021). For this, the *HO* loci of the diploid wild-type Z1 and 8X strains were knocked out using the hygromycin resistance gene as the selection marker using the transformation protocol from (Reuß et al. 2004), with selection on 200 μg/ml hygromycin and PCR and sequence confirmation. The resulting *hoΔ*::Hyg diploid strains were induced to sporulate essentially as in (Lee et al. 2008), and mating types of the resulting haploids were checked via halo assay following the protocol in (Hoffman et al. 2002), using JRY02375 and JRY02176 as the mating testers. Next, for haploid MATα *S. paradoxus* Z1 and its descendant 8X, we generated a petite version of each (lacking mitochondrial DNA, ρ^0), using the protocol adopted from (Amine et al. 2021).

Finally, we used wild-type, MATα haploid, G418-resistant, hygromycin-resistant *S. cerevisiae* strains (DBVPG1373, YPS128, DBVPG6044 and Y55) mitochondrial DNA from the SGRP collection (Table S1) (Cubillos et al. 2009) as donors of mitochondrial DNA for cybrids in our *S. paradoxus* wild-type and 8X backgrounds. Our method for cybrid construction used the cytoduction protocol of (Nguyen et al. 2020) as follows. We first constructed a heterokaryon-competent *S. cerevisiae* strain harboring each mitotype of interest, referred to as a carrier. For this, a single colony of each donor in turn, and a single colony of the ρ^0 *kar1*-d15 mutant (JRY5450, a generous gift from Dr. Jasper Rine), were each inoculated into a separate 3 mL of liquid YPD (1% yeast extract, 2% peptone, 2% glucose) and incubated at 28°C overnight with shaking at 250 rpm. We mixed 100 μL of the two, spotted on a YPD plate, and incubated at 28°C for 4 hours. The mixed cellular patch was inoculated into 3 mL of YPD and incubated at 28°C for 2 hours with shaking at 250 rpm to promote cell division. After a 2-hour incubation, 10^{-5} OD_{600} of cells were spread on a YPG plate (1% yeast extract, 2% peptone, 2% glycerol) to select for mitochondrial genome retention. Colonies on the YPG plates were transferred to a YPD plate with 200 μg/mL hygromycin or 200 μg/mL G418 to check for hygromycin and G418 sensitivity; we earmarked and preserved drug-sensitive, respiration-competent strains as those in which the donor mitochondrial DNA had successfully been transferred to the JRY5450 nuclear background. We next used each carrier strain in turn to transfer the *S. cerevisiae* mitochondrial DNA of interest into the haploid *S. paradoxus* Z1 ρ^0 strain and, separately, its descendant, the 8X ρ^0 strain. This cytoduction proceeded as above except that the carrier and the *S. paradoxus* strain were mated and the progeny were selected on YPG alone; afterward we subjected them to the halo assay, to eliminate the carrier-*S. paradoxus* diploid hybrids, and only those strains that behaved as haploid MATα were retained as cybrids.

Growth assays

Measurements of carrying capacity at 39°C were done as described (AlZaben et al. 2021) with some modifications. To make sure the cells were respiratory-competent before the assay, strains were streaked on YPG plates and were allowed to grow at 28°C for 4 days to form colonies. Three colonies were inoculated into 15 mL of liquid YPG and incubated at 28°C for overnight with shaking at 250 rpm. The cultures would reach 0.4-0.8 OD_{600} after a 16-hour growth. The cultures were then back-diluted into 5 mL of YPD, YPE (1% yeast extract, 2%

peptone, 2% ethanol), or YPG to achieve 0.1 OD_{600}/mL, and then incubated at 23°C, 28°C or
39°C for 24 hours. This procedure, from streaking on solid agar plates to biomass
measurement, was repeated at least twice for each strain with three technical repeats in each
biological replicate. We collected the readouts of carrying capacity from all the replicates across
all days for a given strain and compared each mitochondrial transgenic to *S. paradoxus* with a
two-tailed Mann-Whitney U test. For epistatic analysis, a two-way ANOVA test was carried out to
examine the impact of the genotype-mitotype interaction on thermotolerance. All the statistical
analyses were done with scipy.stats in Python 3.7.

MTT assay

To measure respiration efficiency, pre-culture and treatment were as in Growth assays, above,
and after 24 hours we carried out the MTT assay as in (Sánchez and Königsberg 2006).
Independent experiments from growth to MTT measurement were performed at least six times.
The readouts were normalized to that of wild-type *S. paradoxus* at either 28°C or 39°C. Two-
tailed one-sample Wilcoxon test was performed using the normalized data with a hypothesized
value of 1 with scipy.stats in Python 3.7.

Growths of *S. cerevisiae* and *S. paradoxus* in glycerol

Data are from (Liti et al. 2009) and Mann-Whitney U test was performed to compare the growths
of the two species in glycerol with scipy.stats in Python 3.7.

Fig 1

(A)

S. paradoxus
S. cerevisiae

(B)

Sp
Sc1
Sc4
Sc2
Sc3

Oak tree
Soil
Wine
Soil
Grapes

Fig 2

(A)

(B)

(C)

(D)

Fig 3

(A)

(B)

Fig 4