Using single-cell genomics to explore transcriptional divergence and *cis*-regulatory dynamics of duplicated genes

Xiang Li[1], Xuan Zhang[1], Robert J. Schmitz[1]

[1]Department of Genetics, University of Georgia, Athens, GA, 30602, USA.

Address for correspondence: schmitz@uga.edu

Abstract

Gene duplication is a major source of evolutionary innovation, enabling the emergence of novel expression patterns and functions. Leveraging single-cell genomics, we investigated the transcriptional dynamics and *cis*-regulatory evolution of duplicated genes in cultivated soybean (*Glycine max*), a species that has undergone two rounds of whole-genome duplication. Our analysis revealed extensive transcriptional divergence among these duplicated gene pairs, including dosage balanced, paralogue dominance, divergence, specialization, and widespread mono-expressed pattern. These expression divergences are correlated with sequence variation in their associated accessible chromatin regions (ACRs), where *cis*-regulatory elements reside. Moreover, different duplication mechanisms, including whole-genome duplication and small-scale duplication, likely give rise to distinct types of *cis*-regulatory variants, leading to varying effects on transcriptional divergence. To further explore the evolution of *cis*-regulatory elements and their impact on gene expression, we focused on gene sets in which duplicated genes are derived from two rounds of whole-genome duplication and share a common ancestral gene, as *cis*-regulatory elements are typically co-duplicated with genes during whole-genome duplication. Most ACRs were retained after duplication whereas a subset likely arose *de novo*. Both mutations in duplicated elements and the emergence of *de novo* evolved elements contribute to widespread

23   expression divergence within these gene sets. We also characterized the evolution of cell-type-specific

24   expression and cell-type-specific ACRs among these sets. Collectively, our findings highlight the

25   important role of *cis*-regulatory evolution in shaping transcriptional divergence and facilitating the

26   retention of duplicated genes, offering a high-resolution view through the lens of single-cell genomics.

Introduction

Gene duplication is a crucial driver of evolution, providing the raw materials for new gene functions

and expression patterns (Lynch and Conery 2000; Birchler and Yang 2022). It arises through small-scale

duplication, such as tandem or proximal duplications, or through whole-genome duplication that

replicates entire genomic segments simultaneously (Zhang 2003; Kuzmin et al. 2022). These mechanisms

generate duplicated genes with varying evolutionary rates, essentiality, and functions (Freeling 2009;

Qiao et al. 2019; Shi et al. 2020; Cai and Des Marais 2024; Benoit et al. 2025). In plants, approximately

65% of annotated genes have duplicates, mostly derived from whole-genome duplication, consistent

with the prevalence of palaeopolyploidy in land plants (Wendel 2000; Blanc and Wolfe 2004; Panchy et

al. 2016; Tiley et al. 2016). The retention of these duplicates has been instrumental in shaping plant

genomes and contributing to important agricultural traits (Panchy et al. 2016; Salman-Minkov et al.

2016; Benoit et al. 2025).

Following gene duplication, one or both copies may be subject to relaxed selective constraints and,

in some instances, acquire new functions that contribute to adaptation (neofunctionalization) or

partition ancestral functions (subfunctionalization). However, functional divergence is not always

necessary for their retention and typically unfolds over extended evolutionary timescales (Panchy et al.

2016; Birchler and Yang 2022). Multiple mechanisms have been proposed to account for the

dispensability of retained duplicates (Innan and Kondrashov 2010; Panchy et al. 2016; Qiu et al. 2020).

Moreover, factors such as selection, genetic drift and genetic redundancy (or genetic buffering) can also

influence gene retention (Panchy et al. 2016; Iohannes and Jackson 2023). On the other hand, this

redundancy invariably complicates genotype-phenotype associations and limits the transferability of

genetic findings across closely related species (Iohannes and Jackson 2023; Benoit et al. 2025). Despite

their evolutionary and functional significance, the dynamics of gene duplication and diversification over

short evolutionary timescales remain insufficiently explored (Benoit et al. 2025).

Divergent expression among duplicate genes is a widespread phenomenon and provides crucial

insight into the evolution of duplicated genes. In Arabidopsis, 70% of duplicated gene pairs exhibit

significant transcriptional differences (Ganko et al. 2007). Similarly, approximately 50% of soybean

duplicated genes, derived from whole-genome duplication, show transcriptional divergence (Roulin et al.

2013). This divergence is likely driven by genetic variants within regulatory regions (Wray et al. 2003),

particularly *cis*-regulatory elements that control temporospatial expression patterns (Wittkopp and Kalay

2012; Schmitz et al. 2022). In cotton, divergence in *cis*-regulatory sequences between its diploid

progenitors contributes to transcriptional divergence in 40% of the duplicated genes (Chaudhary et al.

2009). Similar patterns have been observed in Arabidopsis (Haberer et al. 2004), soybean (Fang et al.

2023), strawberry (Fang et al. 2024) and additional cotton studies (Han et al. 2022; Hu et al. 2024).

Despite these valuable insights, a key underexplored question remains: how do changes in gene

expression and *cis*-regulatory control influence the retention of duplicated genes? Specifically, what

types of transcriptional divergence and genetic variations in *cis*-regulatory regions persist in duplicated

genes to ensure their retention over relatively short evolutionary timescales?

Soybean (*Glycine max*) has gone through two rounds of whole-genome duplication, approximately

59 and 13 million years ago (Schmutz et al. 2010; Koenen et al. 2021; Zhuang et al. 2022), which make it

an excellent model for exploring this question. Gene pairs resulting from both whole-genome and small-

scale duplications have been identified in soybean (Qiao et al. 2019). Previous studies have also explored

the evolutionary fates of duplicated genes in soybean, highlighting divergence in gene expression and

the role of *cis*-regulatory control (Du et al. 2012; Zhao et al. 2017; Fang et al. 2023). However, these

studies primarily assessed transcriptional divergence at the tissue level, often relying on differential

expression as the sole metric (Fang et al. 2023), which likely oversimplifies "divergence" and overlooks

differences at the cellular level. For instance, even when two duplicated genes exhibit similar overall

expression levels within a given tissue, they may display distinct expression patterns across different cell

types within that tissue. The recently generated single-cell assay for transposase-accessible chromatin

sequencing (scATAC-seq) and single-nucleus RNA sequencing (snRNA-seq) data in soybean (Zhang et al.

2024) provide a valuable opportunity to study transcriptional and *cis*-regulatory sequence divergence of

duplicated genes at a single-cell resolution. snRNA-seq enables precise measurement of gene expression

at the cellular level, whereas scATAC-seq profiles chromatin accessibility, offering deeper insights into

how *cis*-regulatory activity, particularly cell-type-specific ones, contributes to gene expression

divergence.

In this study, we investigated duplicated genes by integrating single-cell expression and chromatin

accessibility profiles to examine transcriptional and *cis*-regulatory sequence divergence. This approach

provides a more comprehensive understanding of transcriptional divergence and its correlation with *cis*-

regulatory divergence, shedding light on how these processes are intertwined with the distinct fates of

duplicated genes arising from different duplication mechanisms. Given that *cis*-regulatory elements are

typically duplicated along with genes during whole genome duplication, we further examined their

evolution and impacts on expression divergence by focusing on gene sets in which genes are derived

from two rounds of whole-genome duplications and share the same ancestral gene. Our analysis

revealed that most *cis*-regulatory elements were retained after whole-genome duplication, whereas a

subset evolved *de novo*. Mutations within these duplicated elements and the emergence of novel ones

collectively drive transcriptional divergence. Lastly, we explored the evolution of cell-type-specific

expression and cell-type-specific *cis*-regulatory sequences among these gene sets. Overall, our findings

underscore the power of single-cell genomics in revealing the divergence of expression and chromatin

accessibility in retained duplicated genes, offering key insights into the evolution of *cis*-regulatory

elements and gene duplication.

Results

Single-cell transcriptomics reveal the expression relationships of retained duplicated genes

A previous study identified 73,983 duplicated gene pairs in soybean, classifying them into two main

categories: whole-genome duplication and small-scale duplication gene pairs (Qiao et al. 2019). The

small-scale duplicated gene pairs were further subdivided into four types based on their duplication

mechanisms: tandem duplication (closely adjacent duplicated genes on the same chromosome),

proximal duplication (duplicated genes located nearby on the same chromosome but separated by up to

10 other genes), transposed duplication (duplicated genes transposed to distant genomic positions), and

dispersed duplication (duplicated genes scattered across the genome, neither neighboring nor colinear)

(**Fig. 1a and supplementary table S1**) (Qiao et al. 2019). Importantly, individual genes can undergo

multiple duplication events via distinct duplication mechanisms. For instance, a gene duplicated through

whole-genome duplication may later experience tandem duplication. Among all identified duplicated

gene pairs, whole-genome duplication derived gene pairs represented the largest proportion (47.1%),

followed by dispersed duplication (40.9%) (**Fig. 1b and supplementary table S2**). The remaining 12% of

duplicated gene pairs arose from tandem, proximal, and transposed duplications, with transposed

duplication being the most prevalent among them (**Fig. 1b and supplementary table S2**).

The expression dynamics of duplicated gene pairs play a crucial role in their retention (Thompson

et al. 2016; Birchler and Yang 2022; Iohannes and Jackson 2023). To investigate this, we integrated the

gene pair information with the single-cell gene expression profiles from seven tissues - root, hypocotyl,

nodule, globular stage seed, heart stage seed, cotyledon stage seed and early maturation stage seed

(Zhang et al. 2024). Building on the methods developed by (Benoit et al. 2025), we refined the approach

to assess gene expression at the level of individual cells within each tissue, rather than solely across

different tissues (Benoit et al. 2025). This cell-level resolution allows for a more precise and

comprehensive examination of expression relationships of duplicated genes, providing insight into the

expression-based mechanisms that contribute to their retention (Coate et al. 2020). In addition to the

four previously described biologically meaningful expression patterns (Benoit et al. 2025) – (1) dosage

balanced, where duplicates retain similar expression profiles and levels across cell types within a tissue

(**Fig. 1c**); (2) paralogue dominance, where duplicates share similar expression profiles but exhibit

divergent expression levels (**Fig. 1d**); (3) specialization, where duplicates display distinct expression

profiles, with one acquiring cell-type-specific expression (**Fig. 1e**); (4) divergence, where duplicates differ

in both expression profiles and levels – we introduced a fifth category (**Fig. 1f**): (5) mono-expression, in

which only one gene in the pair is expressed whereas the other remains inactive in a given tissue (**Fig.**

1g). Additionally, gene pairs in which neither gene is expressed in the examined tissues are classified as

non-expression, whereas those that do not fit any of the above categories are grouped as "others" (**Fig.**

1h). Among the five biologically meaningful expression patterns mentioned above, the mono-expression

category consistently contains the most gene pairs across different tissue types (**Fig. 2a and**

supplementary table S3). This is followed by dosage balanced, divergence, paralogue dominance, and

specialization patterns, except in the hypocotyl, where divergence gene pairs outnumber dosage

balanced ones, and in the early maturation seed stage, where paralogue dominance exceeds divergence

(**Fig. 2a and supplementary table S3**).

When examining the expression pattern of gene pairs based on their duplication mechanisms, we

found that those derived from whole-genome duplication and dispersed duplication exhibited similar

patterns, whereas gene pairs from tandem, proximal, and transposed duplications showed greater

similarity to each other (**Fig. 2b and supplementary fig. S1 and table S3**). The unpredictable, seemingly

random nature of dispersed duplication events (Ganko et al. 2007; Qiao et al. 2019) may explain why the

expression pattern of gene pairs derived from this mechanism differ from those derived from other

small-scale duplications. Based on this distinction, we focused on gene pairs arising from tandem,

proximal, and transposed duplications—collectively referred to as regional duplications due to their

tendency to remain within specific chromosomal regions—to compare their expression patterns with

those of gene pairs derived from whole-genome duplication.

First, the proportion of expressed gene pairs—where at least one gene is active in a given tissue—

varied by duplication mechanisms. For instance, in early nodule tissue, over 60% of gene pairs from

regional duplications (excluding transposed duplication) were not expressed, whereas only 37% of gene

pairs derived from whole-genome duplication remained inactive (**supplementary table S3**). Second,

among expressed gene pairs, most regional duplications exhibited mono-expression (**Fig. 2b and**

supplementary fig. S1). Specifically, up to 72% of expressed gene pairs from regional duplications

showed mono-expression in early nodule tissue, compared to 43% of those from whole-genome

duplication (**supplementary table S3**). Lastly, a few gene pairs from regional duplications displayed other

biologically meaningful expression patterns, such as dosage balanced, paralogue dominance,

specialization, or divergence expression (**Fig. 2b and supplementary fig. S1**). For example, only 11% of

expressed gene pairs from proximal duplication exhibited these patterns in early nodule tissue,

compared to approximately 20% of expressed gene pairs from whole-genome duplication

(**supplementary table S3**). These results suggested that gene pairs exhibiting correlated expression

patterns, such as dosage balanced, paralogue dominance and specialization expression, are more likely

to originate from whole-genome duplication than from regional duplications. Similar trends have been

previously reported in Arabidopsis, where duplicated genes originating from large-scale duplications and

retained in duplicated segments exhibit more correlated expression patterns than those arising from

small-scale duplications or those not located within duplicated segments (Casneuf et al. 2006).

By assigning an expression pattern, including non-expressed, "others", mono-expressed, dosage

balanced, paralogue dominance, specialization and divergence, to each gene pair across the seven

examined tissue types, we were able to analyze their expression dynamics across tissues. In total, we

identified 1,344 distinct expression patterns (**supplementary table S4**). To simplify the results, we

focused on the top 25 expression patterns across tissue types, each comprising at least 0.5% of the gene

pairs (**Fig. 2c and supplementary table S4**). The most prevalent expression patterns involved gene pairs

exhibiting mono-expression in at least one tissue and non-expression in the remaining tissues (**Fig. 2c**).

Together, these patterns accounted for more than 20% of the gene pairs analyzed (**supplementary table**

S4). Additionally, in gene pairs exhibiting a mono-expression across tissues, the same gene within the

pair was always expressed across different tissue types (**supplementary fig. S2 and table S5**). A previous

study also found that differential expression within gene pairs tend to occur in the same direction across

tissues (Roulin et al. 2013). The non-expression in all examined tissues was the second most common

expression pattern across tissue types (**Fig. 2c**), which was expected given that our study included only

seven tissue types. A lot of gene pairs also exhibited mono-expression across seven tissues, ranking the

third most common expression pattern (**Fig. 2c**). The remaining top 25 expression patterns across tissue

types often involved "others" expression pattern along with biologically meaningful expression patterns

such as mono-expression, dosage balanced and divergence (**Fig. 2c**). Notably, dosage balanced

expression was the most frequently observed biologically meaningful pattern when combined with

"others" expression pattern (**Fig. 2c**). Together, the single-cell expression profiles revealed a wide variety

of expression patterns exhibited by duplicated gene pairs both within and across tissue types.

Correlations between genetic variants within ACRs and expression patterns of duplicated gene pairs

Cis-regulatory variants are thought to drive transcriptional divergence in duplicated gene pairs and

contribute to their retention (Iohannes and Jackson 2023); to explore this, we analyzed Accessible

Chromatin Regions (ACRs), where *cis*-regulatory elements reside, from scATAC-seq data in each of the

seven tissue types corresponding to the snRNA-seq data (Zhang et al. 2024). For each gene pair, we

associated ACRs with duplicated gene based on their proximity and counted the number of ACRs

associated with each gene separately. In some cases, both genes had at least one associated ACR,

whereas in others, only one gene was linked to ACR whereas the other one had no associated ACRs. The

latter one was more frequently observed in gene pairs exhibiting mono-expression than in those

exhibiting other biologically meaningful expression patterns (**Fig. 3a and supplementary fig. S3**). Gene

pairs where only one gene had associated ACR also showed a higher proportion of mono-expression in

the respective tissue compared to those where both genes had ACR (**supplementary table S6**). These

findings suggest that the presence of *cis*-regulatory elements in only one duplicated gene, may be a key

factor, although not the sole mechanism, driving mono-expression in duplicated gene pairs.

To assess genetic variation in ACRs associated with duplicated gene pairs, we used a BLAST-based

approach following the method of (Mendieta et al. 2024). For each gene pair, the ACR sequence

associated with one duplicated gene was used as the query, whereas the reference was defined as the

genomic region spanning from the upstream to the downstream gene of the corresponding duplicate

(**Fig. 3b**). This approach accounts for the possibility that the ACR linked to the corresponding duplicate

may be positioned between it and either its upstream or downstream gene (**Fig. 3b**). Conserved ACRs

were defined based on BLASTed ratio, calculated as the length of the BLASTed sequence in the reference

region divided by the length of the query ACR. ACRs with BLASTed ratios greater than 0.1 were

considered conserved (**Fig. 3c**). We found that gene pairs exhibiting dosage balanced had the highest

percentage of conserved ACRs, whereas divergence gene pairs had the lowest (**Fig. 3d, supplementary**

fig. S3 and table S7). Dosage balanced gene pairs exhibit the most similar expression profiles and levels

(**Fig. 1c**), whereas divergence gene pairs show great divergence (**Fig. 1f**). Similarly, gene pairs displaying

paralogue dominance - characterized by similar expression profiles but varied expression level (**Fig. 1d**) -

also had a high percentage of conserved ACRs (**Fig. 3d, supplementary fig. S3 and table S7**). These

results suggest that proportion of conserved ACRs correlates with the degree of expression similarity

between duplicated genes.

Further analysis of genetic variation between conserved ACRs and their corresponding BLASTed

sequences - including average BLASTed ratio and average mismatch rate (calculated as the ratio of

mismatches between the BLASTed sequence and the ACR query to the BLASTed sequence length) -

revealed that ACRs and their BLASTed sequences were more divergent in gene pairs with divergence

expression pattern compared to those exhibiting other expression patterns (**Fig. 3e, supplementary fig.**

S3, table S7). Interestingly, gene pairs exhibiting mono-expression had a higher percentage of conserved

ACRs, greater BLASTed ratio, and lower mismatch rate than those with divergence expression pattern

(**Fig. 3d-3e, supplementary fig. S3, table S7**), despite mono-expression being a more extreme form of

divergent expression. One possible explanation is that these measures were based on conserved ACRs,

indicating the presence of both ACRs and their corresponding BLASTed sequences in the reference

region. However, a higher percentage of unBLASTed ACRs - those without a corresponding sequence in

the reference region - was found in mono-expressed gene pairs compared to those with other

biologically meaningful expression patterns (**Fig. 3f, supplementary fig. S3, table S7**). This suggests that

ACRs associated with mono-expressed gene pairs are more likely to be unique, lacking counterparts in

their associated duplicated genes. This uniqueness may result from the *de novo* emergence or loss of *cis-*

regulatory elements, which could play a key role in driving mono-expression. Overall, our analysis

highlights the correlation between expression divergence and ACR sequence variation, considering both

the presence or absence of corresponding ACRs and the sequence divergence between conserved ACR

and its corresponding counterpart.

When further examining these BLAST results in the context of duplication mechanisms, distinct

patterns emerged among gene pairs derived from different duplication mechanisms (**supplementary fig.**

S4 and table S8). Gene pairs originating from tandem duplication exhibited the highest percentage of

conserved ACRs (**supplementary fig. S4**). This observation may be influenced by a methodological

limitation of our BLAST approach, which tends to map ACRs back to themselves for tandem-duplication-

associated ACRs. Specifically, the reference region of one gene encompasses both the tandemly

duplicated gene and its adjacent upstream or downstream gene, often the gene itself, as tandemly

duplicated genes are positioned next to each other (**Fig. 3b**). Tandemly duplicated genes can sometimes

exhibit similar expression profiles, with increased copy number leading to higher overall expression

levels, suggesting that either the shared *cis*-regulatory elements regulate both copies, or that *cis*-

regulatory elements were co-duplicated along with the genes. However, relatively fewer tandem-

duplicated gene pairs exhibited dosage balanced expression pattern (**Fig. 2b and supplementary fig. S1**),

indicating that genetic variation in ACRs may be disproportionately masked or skewed by the current

method. Alternatively, the transcriptional activities of these genes could be impacted by positional

differences of *cis*-regulatory elements, the dependence of expression on DNA folding or competition for

transcription factors at duplicated *cis*-regulatory elements.

For gene pairs derived from other duplications—particularly whole-genome duplication, proximal

duplication, and transposed duplication—ACRs and their corresponding BLASTed sequences displayed

similar levels of divergence (**supplementary fig. S4 and table S8**), despite differing gene expression

patterns (**Fig. 2b and supplementary fig. S1**). This pattern may be explained by the difference in the ratio

of unBLASTed ACRs. Gene pairs derived from whole-genome duplication had a much lower percentage of

unBLASTed ACRs compared to those from proximal and transposed duplications (**supplementary fig. S4**

and table S8). *Cis*-regulatory elements of duplicated genes derived from whole-genome duplication were

more likely to be entirely duplicated (Shoemaker et al. 2006), whereas the variation in the presence or

absence of duplicated *cis*-regulatory elements was more likely common in gene pairs derived from other

duplication mechanisms. Together, different duplication mechanisms likely give rise to different types of

cis-regulatory variants, leading to varying effects on transcriptional divergence.

The evolution of duplicated *cis*-regulatory regions and their impact on expression dynamics and gene

retention following two rounds of whole-genome duplications

Soybean is a paleopolyploid species that has undergone two rounds of whole-genome duplication,

theoretically resulting in four duplicated genes originating from a single ancestral gene. Although

extensive gene loss followed these duplications (Schmutz et al. 2010; Du et al. 2012; Zhao et al. 2017),

the retained complete gene sets provide a valuable opportunity to examine the evolution of *cis*-

regulatory elements - duplicated alongside their associated genes (Li and Schmitz 2025) - and their

impact on gene expression and retention. Specifically, we focused on gene sets containing four

duplicated genes derived from whole-genome duplication and sharing a common ancestral gene across

all bean-like (papilionoid) legume species.

A total of 2,383 gene sets, comprising 9,532 genes, were identified for analysis (**supplementary**

table S9). Within individual tissue, these duplicated gene sets showed transcriptional divergence, both in

the number of genes expressed and in the expression correlation among the expressed genes - an

indicator of expression similarity (**Fig. 4a and 4b, supplementary fig. S5**). For instance, in cotyledon stage

seed, ~ 40% of gene sets contained at least one duplicated gene being not expressed (**Fig. 4a and**

supplementary fig. S5). Given that a coexpression value of 0.9 had been established as the threshold for

defining co-expression for duplicated gene pairs (Benoit et al. 2025), most gene sets with more than two

expressed genes had mean coexpression value below this threshold, indicating their low expression

correlations (**Fig. 4b and supplementary fig. S5**). Based on these observations, we categorized gene sets

into mono-, dual-, tri-, and quad-expressed groups, and further classified them (except mono-expressed)

as correlated if their mean coexpression value exceeded 0.9, and uncorrelated otherwise (**Fig. 4c and**

supplementary table S10). Among these, the quad-uncorrelated category contained the largest number

of gene sets (**Fig. 4c, supplementary fig. S5 and table S10**). Additionally, quad-expressed gene sets

displayed higher expression levels than those in other categories, with the difference most pronounced

when compared to mono-expressed gene sets, which consistently exhibited the lowest expression levels

(**supplementary fig. S6**). Across tissues, a variety of expression pattern combinations were observed,

with quad-uncorrelated expression across all seven examined tissues being the most prevalent (**Fig. 4d,**

supplementary tables S11 and S12). Collectively, these results reveal widespread transcriptional

divergence among the four duplicated genes within the set, reflected in variation in expression presence

or absence, expression correlation, and overall expression level.

To investigate genetic variation in ACRs underlying transcriptional divergence within duplicated

gene sets, we BLASTed each ACR associated with a duplicated gene against the reference regions of the

other three duplicated genes in the same set. To account for varying levels of sequence divergence

among ACRs arising from the first and second rounds of whole-genome duplication, we assigned each

ACR a conservation score, defined as the number of times it was conserved (BLASTed ratio > 0.1) across

the three BLASTs. Based on this metric, ACRs were classified into three categories: unique ACRs, which

did not BLAST to any regions near the other duplicated genes (conservation score = 0); two-copy

conserved ACRs, which appeared to be conserved only between the two genes duplicated from the most

recent whole-genome duplication (conservation score = 1/3); and multiple-copy conserved ACRs, which

were conserved across three or four duplicated genes from both rounds of whole-genome duplications

(conservation score = 1 or 2/3, with the latter suggesting possible loss of ACR in one duplicate) (**Fig. 5a**).

For each gene set, we then calculated the average BLASTed ratio and mismatch rate for two-copy and

multi-copy conserved ACRs to quantify ACR sequence divergence.

The proportion of ACRs associated with gene set that were two-copy or multi-copy conserved was

generally similar, and both were substantially higher than the proportion of unique ACRs—a pattern

consistent across gene sets with varying expression profiles (**Fig. 5b, supplementary fig. S7 and table**

S13). These findings suggest that most ACRs associated with duplicated gene sets are conserved

elements, likely retained from the first or second rounds of whole-genome duplication, whereas only a

small fraction are unique ACRs, potentially representing *de novo* elements that evolved at a low rate.

Notably, multi-copy conserved ACRs showed a lower average BLASTed ratio and higher average mismatch

rate than two-copy conserved ACRs (**Fig. 5c, supplementary fig. S8, and table S13**), indicating greater

sequence divergence in multi-copy conserved ACRs. This pattern supports a temporal model of

regulatory sequence divergence, where ACRs gradually accumulate sequence divergence over time

following duplication.

To examine the relationship between genetic variants associated with ACRs and expression

divergence within duplicated gene sets, we compared the proportions of different ACR categories and

their BLAST sequence similarity across gene sets with varying expression patterns. First, in gene sets with

correlated expression—particularly those showing tri- or quad-correlated patterns —the proportion of

multi-copy conserved ACRs was higher than that of two-copy conserved ACRs (**Fig. 5b, supplementary**

fig. S7 and table S13), suggesting that multi-copy conserved ACRs may play a more prominent role in

maintaining coordinated expression, especially in sets with more than two co-expressed genes. Second,

sequence variation among ACRs was associated with expression divergence in corresponding gene

duplicates. For example, in cotyledon tissue, tri- or quad-uncorrelated groups exhibited a lower average

BLASTed ratio in multi-copy conserved ACRs compared to correlated groups, as highlighted by red

squares in **Fig. 5c**. A similar pattern was observed for two-copy conserved ACRs in some other tissues

and when using mismatch rate as an alternative measure of sequence divergence (**supplementary fig. S8**

and table S13). These findings suggested that genetic variation in either multi-copy or two-copy

conserved ACRs can contribute to expression divergence. Lastly, we found no consistent patterns in the

proportions of different ACR types, average BLASTed ratios or average mismatch rates across gene sets

grouped by the number of expressed genes (**Fig. 5b – 5c, supplementary fig. S8 and table S13**),

indicating that regulation of gene expression presence or absence may be more case-dependent or

influenced by mechanisms beyond *cis*-regulatory elements.

The evolution of cell-type-specific expression and cell-type-specific ACRs

Some genes exhibiting cell-type-specific expression are essential for determining or maintaining

cell identity, with *cis*-regulatory elements controlling at which cell types, during what developmental

stages and at what levels these genes are expressed (Wittkopp and Kalay 2012; Schmitz et al. 2022;

Marand et al. 2023). This specificity can differ among duplicated gene sets - some retain expression in

the same cell type across all four copies (**Fig. 6a**), whereas others exhibit varied cell-type-specific

expression (**Fig. 6b - 6c**). These patterns, in turn, enable the identification of candidate *cis*-regulatory

regions responsible for expression specificity and provides insights into the mechanisms driving its

evolution.

Here, we continued to focus on gene sets originating from two rounds of whole-genome

duplication that share a common ancestral origin, exclusively selecting sets in which all four genes are

expressed and at least one exhibit cell-type-specific expression. This filtering step allowed us to focus on

changes in cell-type-specific expression, rather than the confounding effects of loss of expression in

some duplicates. To assess the evolutionary dynamics of cell-type-specific expression, we clustered

duplicated genes based on their expression contributions and then counted the number of shared

clusters within each set, which reflects the degree of conservation in cell-type-specific expression (**Fig.**

6). Only a few sets retained a single shared cluster (**Fig. 7, supplementary fig. S9 and table S14**), implying

the potential conservation of cell-type-specific expression from the ancestral gene (**Fig. 6a**), whereas

most sets had multiple clusters (**Fig. 7, supplementary fig. S9 and table S14**), particularly in uncorrelated

sets, likely reflecting subfunctionalization of ancestral gene expression or the emergence of novel cell-

type-specific expression (**Fig. 6b -6c**).

To explore the role of *cis*-regulatory elements in shaping these expression patterns, we examined

cell-type-specific ACRs (ctACRs) associated with these gene sets (Zhang et al., 2024). Gene sets that

retained a single shared cluster always contained at least one conserved ctACR; notably, in some tissue

types, no gene sets composed exclusively of unique ctACRs contained a single shared cluster (**Fig. 7,**

supplementary fig. S10). In contrast, gene sets with multiple shared clusters showed a variety of ctACR

configurations—some with only conserved ctACRs, some with only unique ctACRs, and others with a mix

of both (**Fig. 7, supplementary fig. S10**). These patterns suggest that changes in cell-type-specific

expression can arise from either sequence divergence within conserved ctACRs or from the emergence

of *de novo* evolved ctACRs. Moreover, multi-copy conserved ctACRs in gene sets with a single shared

cluster exhibited much higher average BLASTed ratios and lower average mismatch rates compared to

those in gene sets with multiple shared clusters (**supplementary fig. S11 and table S15**), highlighting the

important role of multi-copy conserved ctACRs in maintaining conserved cell-type-specific expression

within the set.

Discussion

This study investigates the expression patterns of duplicated gene pairs and gene sets across

diverse cell types and tissues using single-cell genomic data, offering a comprehensive understanding of

the diversity and prevalence of their transcriptional profiles. Further analysis of ACRs associated with

these duplicated gene pairs or sets revealed correlations between the sequence variation in ACRs and

expression divergence. Additionally, different duplication mechanisms likely give rise to distinct types of

cis-regulatory variants, resulting in varying effects on transcriptional divergence. To gain deeper insights

into the evolution of *cis*-regulatory elements, we analyzed the expression patterns and associated ACRs

of four genes derived from whole-genome duplication that share a common evolutionary origin. Most

ACRs associated with these gene sets were retained from the first or second whole-genome duplication

events and have accumulated mutations over time, whereas a small subset of these ACRs likely evolved

de novo. The mutations in conserved ACRs and the emergence of novel ACRs collectively contribute to

widespread expression divergence—reflected in the number of genes expressed, expression level, and

expression correlation. Furthermore, we examined the evolution of cell-type-specific expression and

ctACRs, revealing dynamics of cell-type-specific gene expression within duplicated gene sets. These

changes were driven by mutations in conserved ctACRs as well as the emergence of novel regulatory

elements, highlighting the regulatory complexity underlying cell-type-specific expression patterns.

Single-cell genomics: a key to understanding transcriptional divergence and genetic variants in chromatin

accessibility

The transcriptional divergence of duplicated genes has been extensively studied in polyploid plants

(Du et al. 2012; Roulin et al. 2013; Zhao et al. 2017; Bird et al. 2020; Wang et al. 2021; Han et al. 2022;

Fang et al. 2023; Hu et al. 2024). However, due to technological limitations, most previous studies have

relied on bulk RNA-seq, assessing divergence primarily through differential expression analyses at tissue

level, with only few focusing on the single-cell level (Coate et al. 2020). Yet, such tissue-level differential

expression—whether high or low across entire tissues—can result from diverse underlying patterns,

including paralog dominance, specialization, or divergence (**Fig. 1**), which cannot be resolved by using

bulk RNA-seq. In contrast, single-cell genomics enables gene expression analysis at the cellular level,

providing a more detailed view of expression divergence, including shifts in expression across specific cell

types, the presence or absence of expression in particular tissues, and altered expression patterns across

tissue types (**Fig. 1 and 2**). Additionally, this improved resolution of transcriptional divergence,

particularly across different cell types within individual tissues, allows for a deeper understanding of its

relationship with ACR genetic variants (**Fig. 3**).

Admittedly, bulk ATAC-seq, which identifies ACRs within tissues, is sufficient to uncover broad

correlations. Combining bulk ATAC-seq with snRNA-seq could potentially pinpoint candidate regulatory

regions underlying specific transcriptional patterns, such as conserved ACRs with high BLASTed ratio or

low mismatch rate associated with gene pairs exhibiting dosage balanced expression or unique ACRs

linked to gene pairs with mono- or divergence expression patterns. However, integrating scATAC-seq with

snRNA-seq provides a more powerful approach to explore the evolution of cell-type-specific expression

and ctACRs. We did observe the correlations between genetic similarity in ctACRs and expression

similarity of their associated gene sets (**supplementary fig. S11**) and the contributions of both mutations

in conserved ctACRs and emergence of novel ACRs to the dynamics of the cell-type-specific expression

within the sets (**Fig. 7 and supplementary fig. S10**). Importantly, these results offer a valuable resource

to identify *cis*-regulatory elements responsible for the conservation (**e.g. Fig. 6a**) or shifts (**Fig. 6b - 6c**) in

cell-type-specific expression in specific case studies in the future.

The evolution of *cis*-regulatory elements

The duplication of *cis*-regulatory elements serves as a key source for the emergence of new

regulatory elements (McDonald and Reed 2024; Li and Schmitz 2025). Similar to protein-coding

sequences of duplicated genes, which often experience reduced selective constraints (Kondrashov et al.

2002), duplicated *cis*-regulatory regions may also undergo relaxed selection, allowing mutations to

accumulate through genetic drift and potentially driving phenotypic novelty. Moreover, this pattern may

be more pronounced in whole-genome duplications than small-scale duplications. Duplicates arising

from whole-genome duplication tend to retain shared *cis*-regulatory elements (Shoemaker et al. 2006),

although this is not necessarily the case for those evolved from small-scale duplications (Arsovski et al.

2015). For example, a higher number of conserved non-coding regions are more likely to be retained in

duplicated genes arising from whole-genome duplication (Schnable et al. 2011). Our observation of a

higher percentage of ACRs that can be BLASTed to corresponding duplicated regions in whole-genome

duplication-derived genes compared to small-scale duplication-derived genes (**supplementary fig. S4**),

also underscores this distinction. These differences, in turn, can influence expression pattern divergence,

with more pronounced divergence seen in duplicated genes from small-scale duplications, especially

those arising from regional duplications (**Fig. 2 and supplementary fig. S1**).

On the other hand, the evolution of *cis*-regulatory elements can involve modifications to

preexisting elements through mutations, deletions, insertions, or the *de novo* emergence of regulatory

sequences from previously non-regulatory regions (Katikaneni and Lowe 2025; Li and Schmitz 2025). Our

analysis of proportional of different categories of ACRs associated with the gene sets with four whole-

genome duplication derived genes with shared common ancestor (**Fig. 5a**), potentially provides insights

into the relative contributions of these mechanisms to *cis*-regulatory sequence evolution. Most ACRs

associated with these sets were retained from the first or second rounds of whole-genome duplication

whereas only a small subset of ACRs evolved *de novo* (**Fig. 5b and supplementary figs. S7 and S8**)

suggesting that both modifications of existing sequences and *de novo* emergence of *cis*-regulatory

elements exist but the latter one probably occur in a lower rate. Additionally, the obvious lower average

BLASTed ratio and higher average mismatch rate in multi-copy conserved ACRs than those in two-copy

conserved ACRs (**Fig. 5c**), suggest that mutations in ACRs have accumulated over time.

The retention of duplicated genes and the involvement of *cis*-regulatory control

Three canonical outcomes of duplicated genes over long evolutionary time include gene loss,

subfunctionalization and neofunctionalization (Ohno 1970; Lynch and Conery 2000; Birchler and Yang

2022; Kuzmin et al. 2022; Gout et al. 2023). However, before duplicated genes can undergo

subfunctionalization or neofunctionalization, they must first be maintained without being selected

against—a process that can persist for an extended period, though in some situations

subfunctionalization and neofunctionalization can arise quickly (Birchler et al. 2005; Birchler and Yang

2022). The prevailing consensus in the field suggests that the prolonged maintenance phase is primarily

driven by selective constraints on gene dosage (Iohannes and Jackson 2023). Duplicated genes can

impose negative fitness effects due to increased expression dosage, which can lead to stoichiometric

imbalances (Birchler et al. 2005; Freeling and Thomas 2006; Veitia et al. 2008; Veitia et al. 2013; Birchler

and Yang 2022). This issue is particularly critical for genes involved in protein-protein interactions, as

proper stoichiometries among interacting partners are essential for maintaining functional complexes

(Defoort et al. 2019; Birchler and Yang 2022; Iohannes and Jackson 2023). A key distinction between

whole-genome duplication and small-scale duplication lies in dosage balance: whereas whole-genome

duplication preserves the relative dosage of all interacting partners, small-scale duplication disrupts this

balance because only a subset of genes is duplicated (Ohno 1970; Defoort et al. 2019; Kuzmin et al.

2022). The greater prevalence of similar expression patterns—including dosage balanced, paralogue

dominant, and specialization expression—and the reduced incidence of mono-expression in gene pairs

derived from whole-genome duplication compared to those from small-scale duplication, particularly

regional ones, (**Fig. 2b and supplementary fig. S1**) likely reflects the impact of dosage-balance selection.

Although the initial similarity in expression among gene pairs derived from whole-genome duplication

may simply result from the simultaneous duplication of *cis*-regulatory elements alongside the genes, the

continued maintenance of these regulatory elements suggests the potential influence of selective

pressures.

On the other hand, the process of maintaining dosage balance is complex and can involve several

mechanisms: hypofunctionalization, where both duplicates reduce their expression; dosage balance,

where the duplicates remain reduced but similar expressions to preserve the original gene's function; or

compensatory drift, where one duplicate decreases in expression while the other increases to maintain

functionality of the progenitor gene; and these processes can occur gradually or bypass certain stages

(Birchler et al. 2005; Qian et al. 2010; Thompson et al. 2016; Veitia 2017; Birchler and Yang 2022;

Iohannes and Jackson 2023). These transcriptional dynamics correspond with the widespread expression

divergence we observed among four-gene sets derived (**Fig. 4 and supplementary fig. S5**), suggesting

that these genes are likely in the midst of dosage-balance processes and may ultimately evolve toward

subfunctionalization, neofunctionalization, or loss.

Our analyses of single-cell genomic data also reflect an important additional factor to consider in

gene retention - the influence of cell-type specificity. Duplicated genes may be in dosage balance or

compensatory drift in certain cell types, but not in other cell types (Birchler and Yang 2022; Iohannes and

Jackson 2023). This is reflected in our observation that different combinations of expression patterns

occur across tissues within the gene pairs (**Fig. 2c**) or gene sets (**Fig. 4d**). Another example is that the

gene set exhibiting consistent phloem-specific expression in the globular stage seed tissue but different

cell-type-specific expression in early-stage nodule tissue (**supplementary fig. S12**). Such variability

suggests that gene retention may be driven by expression in specific tissues while being dispensable in

others. This highlights the importance of understanding the context dependence of this phenomenon,

particularly through the lens of single-cell genomics.

In summary, understanding the evolution of paralogs is crucial, as many crops - including Solanum,

grasses, and legumes - have similarly complex gene families that undoubtedly influence agriculturally

important traits. Gaining insights into the retention of gene duplications and their underlying

mechanisms can inform strategies for fine-tuning gene dosage, facilitating the engineering of

quantitative variation, and overcoming redundancy barriers in functional studies.

Materials and Methods

Duplicated gene pairs and gene sets in cultivated soybean

Duplicated gene pairs in the soybean genome (Williams 82), were categorized based on their duplication mechanisms - whole-genome duplication, tandem duplication, proximal duplication, transposed duplication and dispersed duplication (**Fig. 1**) - and obtained from the Plant Duplication Gene Database (http://pdgd.njau.edu.cn:8080) (Qiao et al. 2019). Gene pairs located in scaffolds were excluded from further analyses. To identify gene sets containing four whole-genome duplication derived genes, which share a common ancestral gene across all bean-like (papilionoid) legume species, we overlapped the legume gene family dataset (https://www.soybase.org/) with duplicated genes originating from whole-genome duplication. Specifically, we first filtered the legume gene families to retain only those containing four duplicated genes in soybean, and then further refined the selection by ensuring that these four genes were derived from whole-genome duplication based on the gene pair information obtained earlier.

Expression relationships of gene pairs or gene sets

Raw gene counts from seven tissues - root, hypocotyl, nodule, globular stage seed, heart stage seed, cotyledon stage seed and early maturation stage seed - were obtained from snRNA-seq (Zhang et al. 2024). Data from each tissue was analyzed separately. Only genes with two replicates per cell type within the same tissue were included in subsequent analyses. For each tissue, we calculated the Spearman correlation between cell type replicates and excluded cell types with low correlation (0.75 or blow). Lowly expressed genes were filtered using "filterByExpr" function, and the remaining gene counts were normalized using "cpm" function in edgeR (Chen et al. 2025). Genes with zero expression across all cell types within a tissue were classified as "non-expressed" and excluded from further analysis. The

remaining genes were used to construct coexpression networks (Benoit et al. 2025) to assess whether

duplicated gene pairs exhibit similar expression profiles across cell types within a tissue.

To quantify expression relationships between duplicated genes within the same pair, we calculated

the absolute mean and standard deviation (SD) of log2-transformed fold change across cell types within

the same tissue (Benoit et al. 2025). Following (Benoit et al. 2025), we classified duplicated gene pairs

into six categories based on coexpression and variation in expression levels, introducing "mono-

expression" as an additional category:

1. Dosage-balanced: coexpression > 0.9, mean $\log_2$ fold change <1, SD $\log_2$ fold change <1

2. Paralogue dominance: coexpression > 0.9, mean $\log_2$ fold change >= 1, SD $\log_2$ fold change <1

3. Specialized: coexpression > 0.9, mean $\log_2$ fold change >= 1, SD $\log_2$ fold change >=1

4. Diverged: coexpression < 0.5, mean $\log_2$ fold change >= 1, SD $\log_2$ fold change >=1

5. Mono-expression: coexpression = NA.

6. Others: expressed gene pairs that did not fall into any of the above categories.

To investigate expression relationships among gene sets, we summarized the metric of

coexpression for all six gene pairs within each gene set. Gene sets were first grouped based on the

number of genes expressed. For sets with at least two expressed genes, we further classified them as

either correlated or uncorrelated based on their average pairwise coexpression values. Gene sets with a

mean coexpression value > 0.9 were designated as correlated, while those below this threshold were

categorized as uncorrelated.

Identification of cell-type-specific genes in each tissue in cultivated soybean

To identify cell-type-specific genes, we performed differential expression analysis for each cell type

within its respective tissue. Gene counts were aggregated across nuclei of the same cell type and

compared to a reference panel, which uniformly sampled gene counts from other cell types within the

same tissue, using edgeR (Chen et al. 2025). Genes with a $\log_2$ fold change > 1 and an adjusted p-value <

0.05 were classified as highly expressed in a given cell type. A gene was defined as cell-type-specific if it

was classified as highly expressed in at least one cell type within its tissue.

To investigate the evolution of cell-type-specific expression among gene sets, we first filtered for

gene sets in which four duplicated genes were expressed and at least one duplicate exhibited cell-type-

specific expression in the relevant tissue. For each gene, we calculated the proportion of expression

attributable to each cell type by dividing the expression level of that cell type by the total expression

level within the tissue. The resulting matrix of cell-type contribution proportions was used for clustering

analysis. Pairwise distances between genes were computed as 1 minus the Spearman correlation

coefficient, and agglomerative hierarchical clustering was performed using the average linkage method

in R. The optimal number of clusters was determined using the gap statistic method with 50

permutations and a maximum of 60 clusters using cluster package. Genes were then assigned to clusters

by cutting the dendrogram at the selected value of k. The number of clusters observed within each gene

set reflects the conservation of cell-type-specific expression among the duplicated genes: a single shared

cluster indicates conserved expression across all four genes (e.g. **Fig. 6a**), whereas two or more clusters

indicate divergence in cell-type-specific expression among the duplicates (e.g. **Fig. 6b - 6c**).

ACR comparisons between the two duplicated genes within each pair

ACRs identified in cultivated soybean was obtained from (Zhang et al. 2024). The gene closest to

each ACR was defined as the gene-associated ACR, and this association was determined using bedtools

(Quinlan and Hall 2010). BLAST analysis was performed to examine the sequence divergence of ACRs

associated with the two genes within the same duplication gene pair. The sequence of the ACR

associated with one duplicated gene was used as the query, while the sequence of the two adjacent

genes from the other duplicated gene served as the reference (**Fig. 3b**). These sequences were isolated

using the "getfasta" function in bedtools (Quinlan and Hall 2010). Then sequence comparisons were

performed using BLASTn with the following parameters "-task blastn-short -evalue 1e-3 -

max_target_seqs 4-word_size 7 -gapopen 5 -gapextend 2 -penalty -1 -reward 1 -dust no -outfmt 6"

following (Mendieta et al. 2024). To align ACR sequence comparisons with the expression relationships

of duplicated genes in each tissue, we identified ACRs associated with each gene in each tissue

separately and conducted BLAST analyses independently.

ACRs were classified as conserved if more than 10% (approximately 50 bp) were BLASTed to the

reference region (**Fig. 3c**). The mean percentage of conserved ACRs were calculated for each gene pair by

dividing the number of conserved ACRs by the total number of ACRs associated with that pair and then

averaged across gene pairs with the same expression pattern (**Fig. 3d**). The BLASTed ratio was calculated

by dividing the length of the BLASTed sequence in the reference region by the length of the query ACR.

The mismatch rate was calculated as the number of mismatches between the ACR and the BLASTed

sequence, divided by the length of the BLASTed sequence in the reference region. The percentage of

unBLASTed ACRs was calculated by dividing the number of ACRs could not be BLASTed to the reference

region by the total number of ACRs associated with the gene pair. The average BLASTed ratio, mismatch

rate, and percentage of unBLASTed ACRs across all the ACRs associated with the gene pair were

calculated to represent the level of sequence divergence of ACRs for each gene pair.

When measuring ACR sequence divergence across duplicated genes with the gene sets, we

classified the ACRs based on conservation score, defined as the number of times an ACR was conserved

(with a BLASTed ratio > 0.1) across the three BLASTs. Based on this metric, ACRs were categorized as:

unique ACRs, two-copy conserved ACRs and multi-copy conserved ACRs (**Fig. 5a**). We examined the

compositions of these ACRs for ACRs associated with each gene set. For each gene set, we calculated the

average BLASTed ratio and mismatch rate for two-copy conserved ACRs, and multiple-copy conserved

ACRs associated with each gene set. When comparing the ACR explaining cell-type-specific expression,

we specifically focused on cell-type-specific ACRs identified in (Zhang et al. 2024).

During the preparation of this manuscript, we used ChatGPT to refine code and improve readability

and language. All content was subsequently reviewed and edited by the authors, who take full

responsibility for the final version.

Acknowledgement

600   We sincerely appreciate the insightful feedback from Dr. Jeffrey Doyle from Cornell University, Dr. Scott

601   Jackson from University of Georgia, Dr. Steven Cannon from USDA-ARS, and the members of the Schmitz

602   lab on this manuscript, particularly Cullan Meyer, Kevin Sun, Dr. Hao Zhang, Dr. Shan Liang and Dr. Mark

603   Minow. This research was supported by the United Soybean Board (2432-201-0102) and the National

604   Science Foundation (IOS-1856627) to R.J.S.

Reference

Arsovski AA, Pradinuk J, Guo XQ, Wang S, Adams KL. 2015. Evolution of Cis-Regulatory Elements
and Regulatory Networks in Duplicated Genes of Arabidopsis. *Plant Physiol*. 169:2982–
2991.

Benoit M, Jenike KM, Satterlee JW, Ramakrishnan S, Gentile I, Hendelman A, Passalacqua MJ,
Suresh H, Shohat H, Robitaille GM, et al. 2025. Solanum pan-genetics reveals paralogues
as contingencies in crop engineering. *Nature*:1–11.

Birchler JA, Riddle NC, Auger DL, Veitia RA. 2005. Dosage balance in gene regulation: biological
implications. *Trends Genet*. 21:219–226.

Birchler JA, Yang H. 2022. The multiple fates of gene duplications: Deletion, hypofunctionalization,
subfunctionalization, neofunctionalization, dosage balance constraints, and neutral
variation. *Plant Cell* 34:2466–2474.

Bird KA, Niederhuth CE, Ou S, Gehan M, Pires JC, Xiong Z, VanBuren R, Edger PP. 2020. Replaying
the evolutionary tape to investigate subgenome dominance in allopolyploid Brassica napus.
New Phytol.

Blanc G, Wolfe KH. 2004. Widespread paleopolyploidy in model plant species inferred from age
distributions of duplicate genes. *Plant Cell* 16:1667–1678.

Cai H, Des Marais DL. 2024. Expression variability following gene duplication facilitates gene
retention. *bioRxiv* [Internet]:2024.11.06.622370. Available from:
https://www.biorxiv.org/content/10.1101/2024.11.06.622370v2.abstract

Casneuf T, De Bodt S, Raes J, Maere S, Van de Peer Y. 2006. Nonrandom divergence of gene
expression following gene and genome duplications in the flowering plant Arabidopsis
thaliana. *Genome Biol*. 7:R13.

Chaudhary B, Flagel L, Stupar RM, Udall JA, Verma N, Springer NM, Wendel JF. 2009. Reciprocal
silencing, transcriptional bias and functional divergence of homeologs in polyploid cotton
(gossypium). *Genetics* 182:503–517.

Chen Y, Chen L, Lun ATL, Baldoni PL, Smyth GK. 2025. edgeR v4: powerful differential analysis of
sequencing data with expanded functionality and improved support for small counts and
larger datasets. *Nucleic Acids Res*. 53:gkaf018.

Coate JE, Farmer AD, Schiefelbein JW, Doyle JJ. 2020. Expression partitioning of duplicate genes at
single cell resolution in Arabidopsis roots. *Front. Genet.* 11:596150.

Defoort J, Van de Peer Y, Carretero-Paulet L. 2019. The evolution of gene duplicates in angiosperms
and the impact of protein-protein interactions and the mechanism of duplication. *Genome*
Biol. Evol. 11:2292–2305.

Du J, Tian Z, Sui Y, Zhao M, Song Q, Cannon SB, Cregan P, Ma J. 2012. Pericentromeric effects shape
the patterns of divergence, retention, and expression of duplicated genes in the
paleopolyploid soybean. *Plant Cell* 24:21–32.

Fang C, Jiang N, Teresi SJ, Platts AE, Agarwal G, Niederhuth C, Edger PP, Jiang J. 2024. Dynamics of
accessible chromatin regions and subgenome dominance in octoploid strawberry. *Nat.*
Commun. 15:1–14.

Fang C, Yang M, Tang Y, Zhang L, Zhao H, Ni H, Chen Q, Meng F, Jiang J. 2023. Dynamics of *cis*-
regulatory sequences and transcriptional divergence of duplicated genes in soybean. *Proc.*
Natl. Acad. Sci. U. S. A. 120:e2303836120.

Freeling M. 2009. Bias in plant gene content following different sorts of duplication: tandem, whole-
genome, segmental, or by transposition. *Annu. Rev. Plant Biol.* 60:433–453.

Freeling M, Thomas BC. 2006. Gene-balanced duplications, like tetraploidy, provide predictable
drive to increase morphological complexity. *Genome Res.* 16:805–814.

Ganko EW, Meyers BC, Vision TJ. 2007. Divergence in expression between duplicated genes in
Arabidopsis. *Mol. Biol. Evol.* 24:2298–2309.

Gout J-F, Hao Y, Johri P, Arnaiz O, Doak TG, Bhullar S, Couloux A, Guérin F, Malinsky S, Potekhin A, et
al. 2023. Dynamics of gene loss following ancient whole-genome duplication in the cryptic
Paramecium complex. *Mol. Biol. Evol.* 40:msad107.

Haberer G, Hindemitt T, Meyers BC, Mayer KFX. 2004. Transcriptional Similarities, Dissimilarities,
and Conservation of cis-Elements in Duplicated Genes of Arabidopsis. *Plant Physiol.*
136:3009–3022.

Han J, Lopez-Arredondo D, Yu G, Wang Y, Wang B, Wall SB, Zhang X, Fang H, Barragán-Rosillo AC,
Pan X, et al. 2022. Genome-wide chromatin accessibility analysis unveils open chromatin
convergent evolution during polyploidization in cotton. *Proc. Natl. Acad. Sci. U. S. A.*
119:e2209743119.

Hu G, Grover CE, Vera DL, Lung P-Y, Girimurugan SB, Miller ER, Conover JL, Ou S, Xiong X, Zhu D, et
al. 2024. Evolutionary Dynamics of Chromatin Structure and Duplicate Gene Expression in
Diploid and Allopolyploid Cotton. *Mol. Biol. Evol.* [Internet] 41. Available from:
http://dx.doi.org/10.1093/molbev/msae095

Innan H, Kondrashov F. 2010. The evolution of gene duplications: classifying and distinguishing
between models. *Nat. Rev. Genet.* 11:97–108.

Iohannes SD, Jackson D. 2023. Tackling redundancy: genetic mechanisms underlying paralog
compensation in plants. *New Phytol.* 240:1381–1389.

Katikaneni A, Lowe CB. 2025. Novelty versus innovation of gene regulatory elements in human
evolution and disease. *Curr. Opin. Genet. Dev.* 90:102279.

Koenen EJM, Ojeda DI, Bakker FT, Wieringa JJ, Kidner C, Hardy OJ, Pennington RT, Herendeen PS,
Bruneau A, Hughes CE. 2021. The origin of the legumes is a complex paleopolyploid
phylogenomic tangle closely associated with the Cretaceous-Paleogene (K-Pg) mass
extinction event. *Syst. Biol.* 70:508–526.

Kondrashov FA, Rogozin IB, Wolf YI, Koonin EV. 2002. Selection in the evolution of gene
duplications. *Genome Biol.* 3:RESEARCH0008.

Kuzmin E, Taylor JS, Boone C. 2022. Retention of duplicated genes in evolution. *Trends Genet.*
38:59–72.

Li X, Schmitz RJ. 2025. Cis-regulatory dynamics in plant domestication. *Trends Genet.* [Internet] 0.
Available from: https://www.cell.com/action/showAbstract?pii=S0168952525000460

Lynch M, Conery JS. 2000. The evolutionary fate and consequences of duplicate genes. *Science*
290:1151–1155.

Marand AP, Eveland AL, Kaufmann K, Springer NM. 2023. cis-Regulatory Elements in Plant
Development, Adaptation, and Evolution. *Annu. Rev. Plant Biol.* 74:111–137.

McDonald JMC, Reed RD. 2024. Beyond modular enhancers: new questions in cis-regulatory
evolution. *Trends Ecol. Evol.* [Internet] 0. Available from:
http://www.cell.com/article/S0169534724001708/abstract

Mendieta JP, Tu X, Jiang D, Yan H, Zhang X, Marand AP, Zhong S, Schmitz RJ. 2024. Investigating the
cis-regulatory basis of C3 and C4 photosynthesis in grasses at single-cell resolution. *Proc.*
Natl. Acad. Sci. U. S. A. 121:e2402781121.

Ohno S. 1970. Evolution by gene duplication. 1970th ed. Berlin, Germany: Springer Available from:
https://link.springer.com/book/10.1007/978-3-642-86659-3

Panchy N, Lehti-Shiu MD, Shiu S-H. 2016. Evolution of gene duplication in Plants1[OPEN]. *Plant*
Physiol. 171:2294–2316.

Qian W, Liao B-Y, Chang AY-F, Zhang J. 2010. Maintenance of duplicate genes and their functional
redundancy by reduced expression. *Trends Genet.* 26:425–430.

Qiao X, Li Q, Yin H, Qi K, Li L, Wang R, Zhang S, Paterson AH. 2019. Gene duplication and evolution
in recurring polyploidization-diploidization cycles in plants. *Genome Biol.* 20:38.

Qiu Y, Van Tay Y, Ruan Y, Adams KL. 2020. Divergence of duplicated genes by repeated partitioning
of splice forms and subcellular localization. *New Phytol.* 225:1011–1022.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics 26:841–842.

Roulin A, Auer PL, Libault M, Schlueter J, Farmer A, May G, Stacey G, Doerge RW, Jackson SA. 2013.
The fate of duplicated genes in a polyploid plant genome. *Plant J.* 73:143–153.

Salman-Minkov A, Sabath N, Mayrose I. 2016. Whole-genome duplication as a key factor in crop
domestication. *Nat Plants* 2:16115.

Schmitz RJ, Grotewold E, Stam M. 2022. Cis-regulatory sequences in plants: Their importance,
discovery, and future challenges. *Plant Cell* 34:718–741.

Schmutz J, Cannon SB, Schlueter J, Ma J, Mitros T, Nelson W, Hyten DL, Song Q, Thelen JJ, Cheng J,
et al. 2010. Genome sequence of the palaeopolyploid soybean. *Nature* 465:120–120.

Schnable JC, Pedersen BS, Subramaniam S, Freeling M. 2011. Dose-sensitivity, conserved non-
coding sequences, and duplicate gene retention through multiple tetraploidies in the
grasses. *Front. Plant Sci.* 2:2.

Shi T, Rahmani RS, Gugger PF, Wang M, Li H, Zhang Y, Li Z, Wang Q, Van de Peer Y, Marchal K, et al.
2020. Distinct Expression and Methylation Patterns for Genes with Different Fates following
a Single Whole-Genome Duplication in Flowering Plants. *Mol. Biol. Evol.* 37:2394–2413.

Shoemaker RC, Schlueter J, Doyle JJ. 2006. Paleopolyploidy and gene duplication in soybean and
other legumes. *Curr. Opin. Plant Biol.* 9:104–109.

Thompson A, Zakon HH, Kirkpatrick M. 2016. Compensatory drift and the evolutionary dynamics of
dosage-sensitive duplicate genes. *Genetics* 202:765–774.

Tiley GP, Ané C, Burleigh JG. 2016. Evaluating and characterizing ancient whole-genome
duplications in plants with gene count data. *Genome Biol. Evol.* 8:1023–1037.

Veitia RA. 2017. Gene duplicates: Agents of robustness or fragility? *Trends Genet.* 33:377–379.

Veitia RA, Bottani S, Birchler JA. 2008. Cellular reactions to gene dosage imbalance: genomic,
transcriptomic and proteomic effects. *Trends Genet.* 24:390–397.

Veitia RA, Bottani S, Birchler JA. 2013. Gene dosage effects: nonlinearities, genetic interactions,
and dosage compensation. *Trends Genet.* 29:385–393.

Wang L, Jia G, Jiang X, Cao S, Chen Z, Song Q. 2021. Altered chromatin architecture and gene
expression during polyploidization and domestication of soybean. *Plant Cell* [Internet].
Available from: https://academic.oup.com/plcell/article/33/5/1430/6175070

Wendel JF. 2000. Genome evolution in polyploids. *Plant Mol. Biol.* 42:225–249.

Wittkopp PJ, Kalay G. 2012. Cis-regulatory elements: molecular mechanisms and evolutionary
processes underlying divergence. *Nature Reviews Genetics* 13:59–69.

Wray GA, Hahn MW, Abouheif E, Balhoff JP, Pizer M, Rockman MV, Romano LA. 2003. The evolution
of transcriptional regulation in eukaryotes. *Mol. Biol. Evol.* 20:1377–1419.

Zhang J. 2003. Evolution by gene duplication: an update. *Trends Ecol. Evol.* 18:292–298.

Zhang X, Luo Z, Marand AP, Yan H, Jang H, Bang S, Mendieta JP, Minow MAA, Schmitz RJ. 2024. A
spatially resolved multi-omic single-cell atlas of soybean development. *Cell* 188:550-
567.e19.

Zhao M, Zhang B, Lisch D, Ma J. 2017. Patterns and Consequences of Subgenome Differentiation
Provide Insights into the Nature of Paleopolyploidy in Plants. *Plant Cell* 29:2974–2994.

Zhuang Y, Wang X, Li X, Hu J, Fan L, Landis JB, Cannon SB, Grimwood J, Schmutz J, Jackson SA, et al.
2022. Phylogenomics of the genus Glycine sheds light on polyploid evolution and life-
strategy transition. *Nat. Plants* 8:233–244.

Fig. 1. Different duplication mechanisms and distinct expression patterns.

(a) Illustration of different gene duplication mechanisms, including whole-genome duplication (WGD),

dispersed duplication (DSD), proximal duplication (PD), tandem duplication (TD), and transposed

duplication (TRD); (b) Proportional contributions of each duplication mechanism to the duplicated gene

pairs identified in soybean; (c-h) Representative examples of duplicated gene pairs exhibiting distinct

expression patterns across cell types within early nodule tissue: (c) dosage balanced, (d) paralogue

dominance, (e) specialization, (f) divergence, (g) mono-expressed and (h) others. In each panel, blue and

red dots/lines represent the two genes within a duplicated pair, with two biological replicates included

for each gene.

Fig. 2. Distributions of expression patterns within and across tissues of duplicated gene pairs derived

from distinct duplication mechanisms.

(a) Number of duplicated gene pairs classified by distinct expression patterns within a tissue, including

mono-expressed, dosage balanced, paralogue dominance, specialization, divergence and others (as

described in Fig.1), as well as non-expressed category. (b) Relationship between gene duplication

mechanisms and expression patterns. The left panel categorizes genes based on duplication

mechanisms: whole-genome duplication (WGD), tandem duplication (TD), proximal duplication (PD),

dispersed duplication (DSD), and transposed duplication (TRD). The right panel represents gene

expression patterns. Flow width indicates the relative frequency of each transition, highlighting how

different duplication mechanisms contribute to expression variation. (c) Number of duplicated gene pairs

classified by a combination of expression patterns across seven examined tissues.

Fig. 3. Genetic variation within ACRs associated with gene pairs with different expression patterns.

(a) Number of ACRs associated with each duplicated gene within gene pairs exhibiting different

expression patterns in the cotyledon stage seed. Within each gene pair, the gene associated with a

higher number of ACRs is classified into the large count group, whereas the one with fewer ACRs is

classified into the small count group. (b) Illustration of the query and reference region for a BLAST: an

ACR sequence associated with duplicated gene1 as a query, whereas the genomic region spanning from

the upstream to the downstream gene of duplicated gene2 as a reference region. (c) Number of ACRs

BLASTed to the reference region in the cotyledon stage seed. ACRs with a BLASTed ratio greater than 0.1

are classified as conserved. (d) Mean percentage of conserved ACRs (black bars) across gene pairs with

different expression patterns in the cotyledon stage seed. (e) Mean BLASTed ratio of conserved ACRs

across gene pairs with different expression patterns in the cotyledon stage seed. The size of dots

represents the mean mismatch rate of conserved ACRs across gene pairs with different expression

patterns in the cotyledon stage seed. The color of dots represents different expression patterns. (f) Mean

percentage of ACRs without a corresponding sequence in the reference region (unBLASTed ACRs) across

gene pairs with different expression patterns in the cotyledon stage seed.

Fig. 4. Expression divergence of duplicated genes within the sets.

(a) Number of duplicated gene sets classified by the number of genes expressed with each set in cotyledon stage seed. (b) Mean coexpression among expressed genes within gene sets, grouped by the number of genes expressed per set in cotyledon stage seed. (c) Number of duplicated gene sets classified by distinct expression patterns in cotyledon stage seed, including mono-expressed, dual-uncorrelated, dual-correlated, tri-uncorrelated, tri-correlated, quad-uncorrelated, quad-correlated and non-expressed. (d) Number of duplicated gene sets classified by a combination of expression patterns across seven examined tissues.

Fig. 5. Proportions and sequence divergence of ACRs with different levels of conservation across gene

sets with distinct expression patterns.

(a) Classifications of ACRs based on their conservation score: unique ACR, two-copy conserved ACR and

multi-copy conserved ACR. (b) Proportions of each ACR category identified in gene sets with different

expression patterns in cotyledon stage seed. (c) Mean BLASTed ratio of two-copy conserved or multi-

copy conserved ACRs across gene sets with different expression patterns in cotyledon stage seed.

Comparisons where correlated groups had higher mean BLASTed ratios than uncorrelated ones are

highlighted with red squares.

Fig. 6. The dynamics of cell-type-specific expression within gene sets in cotyledon stage seed.

(a) Gene set with a single shared cluster: all four duplicated genes exhibited specific expression in the hilum epidermis and embryo epidermis. (b) Gene set with two shared clusters: the two genes on the left formed a cluster with expression specific to the hilum epidermis, while the two on the right formed a separate cluster with expression in both the hilum epidermis and hilum parenchyma. (c) Gene set with three shared clusters: the two genes on the left clustered together and were expressed in the hilum epidermis and hilum parenchyma; the third gene formed a second cluster and was expressed in the tracheid-bar and hilum parenchyma; the fourth gene formed a third cluster and was expressed in the tracheid-bar, hilum epidermis, and hilum parenchyma. Cell-type annotations for cotyledon stage seeds were shown on the right. Arrows indicated the cells where each gene was specifically expressed, with different arrow colors denoting distinct cell types in each panel. WGD: whole-genome duplication; SC: seed coat; Emb: embryo; End: endosperm.

Fig. 7. Heatmap of proportions of cell-type-specific ACRs (ctACRs) with different levels of conservation

within each gene set with at least one gene exhibiting cell-type-specific expression in root tissue.

824   Each row represents a gene set in which all four genes are expressed and at least one exhibits cell-type-

825   specific expression in the root. The three columns display the proportions of two-copy conserved,

826   multiple-copy conserved and unique ACRs associated with each gene set. Color intensity ranges from

827   blue (0%) to red (100%). The bar alongside the heatmap indicates the number of shared clusters

828   identified for each gene set.