Development of a digital analysis system for a novel 3D culture-based colony

formation to detect malignantly transformed cells in human cell-based therapeutic

products

Shinji Kusakawa[1,*], Tingshu Yang[1], Rumi Sawada[1], Yoji Sato[2], Satoshi Yasuda[1,**]

[1] Division of Cell-Based Therapeutic Products, National Institute of Health Sciences,

Kawasaki, Japan

[2] Division of Drugs, National Institute of Health Sciences, Kawasaki, Japan

[*] Correspondence: Shinji Kusakawa, Division of Cell-Based Therapeutic Products,

National Institute of Health Sciences, 3-25-26 Tonomachi, Kawasaki Ward, Kawasaki

City, Kanagawa 210-9501, Japan.

Phone: +81-44-270-6530

E-mail addresses: kusakawa@nihs.go.jp

^{**} Correspondence: Satoshi Yasuda, Division of Cell-Based Therapeutic Products,

National Institute of Health Sciences, 3-25-26 Tonomachi, Kawasaki Ward, Kawasaki

City, Kanagawa 210-9501, Japan.

Phone: +81-44-270-6525

E-mail addresses: yasuda@nihs.go.jp

Abstract

The presence of malignantly transformed cells in human cell-based therapeutic products (hCTPs) is a significant safety concern. Although such cellular impurities in hCTPs can be assessed by detecting anchorage-independent growth using conventional soft agar colony formation (SACF) assays, the sensitivity of these assays is often insufficient. To overcome this limitation, we previously developed a novel tumorigenicity-associated testing method, the digital SACF (D-SACF) assay, which combines a partitioned culture of test cells to concentrate target cells with colony detection via image analysis. However, conventional soft agar culture involves complicated operations, such as preparing multilayered culture media and temperature control, and further technical optimization is required for the widespread adoption of the D-SACF assay. In this study, we focused on a new culture system incorporating a three-dimensional (3D) culture method using a liquid medium containing the low-molecular-weight agar polymer LA717 in low-adhesion culture vessels. We initially confirmed conditions for the efficient high-density 3D culture of normal cells using LA717-supplemented medium in low-adhesion 96-well plates. Using human mesenchymal stem/stromal cells (MSCs) as a normal cell model and HeLa cells as a transformed cell model, we demonstrated that the new 3D culture system effectively maintained the dispersion of MSCs and

prevented their aggregation, while transformed HeLa cells exhibited robust anchorage

independence, thereby establishing the new liquid/low-molecular-weight agar colony

formation (LACF) method as an alternative to SACF. Finally, by systematizing the

digital analysis system for the LACF assay (D-LACF assay), which streamlines the

overall workflow from the performance evaluation of the test method to product testing

and result interpretation, the limitations of the conventional soft agar-based D-SACF

assay were addressed and its practicality and utility were enhanced. This *in vitro*

evaluation system is expected to provide a promising approach for improving the

quality and safety of hCTPs.

Keywords:

Tumorigenicity, malignantly transformed cells**,** human cell-based therapeutic products**,**

3D culture, *in vitro* assay

Introduction

In a three-dimensional (3D) culture environment without an adhesion substrate, normal cells typically undergo anoikis, a type of programmed cell death triggered by the loss of cell–matrix interactions [1,2]. In contrast, malignantly transformed cells exhibit anchorage-independent growth, allowing for their survival, proliferation, and colony formation under such conditions. This distinct phenotype is a hallmark of malignant transformation, and serves as a critical parameter for assessing cellular tumorigenicity [3]. The soft agar colony formation (SACF) assay is regarded as the gold standard for the *in vitro* assessment of cellular anchorage-independent growth [4–6]. Over the years, this assay has been refined to fulfill specific experimental needs, resulting in various modifications that enhance its practicality and sensitivity [7]. For example, the conventional SACF assay is insufficiently sensitive to evaluate the presence of transformed cells as cellular impurities in human cell-based therapeutic products (hCTPs), which consist of large numbers of cells [8]. To overcome this problem, we previously established the digital SACF (D-SACF) assay as an innovative test method that integrates soft agar culture, partitioned culture in multi-well plates, and image analysis to enable the digital detection of colonies derived from trace amounts of transformed cells within a population of normal cells [9]. Using an imaging cytometer

for high-throughput image acquisition and automated analysis, this approach allows the

efficient screening of large numbers of cells with high sensitivity to detect mixed

transformed cells. In a previous study, the detection limit of the D-SACF assay, when

evaluating its performance using human mesenchymal stem/stromal cells (MSCs) as the

product model and HeLa cells as the transformed cell model, was recorded as one HeLa

cell in 1×10^7 MSCs, i.e., 0.00001% of the total number of cells. Furthermore, recent

multisite studies have confirmed the high utility and feasibility of this test system [10].

Despite these advantages, the D-SACF assay presents practical challenges. As the

number of wells and plates increases, preparing multilayered agar media with different

concentrations and maintaining the required temperature becomes challenging. This

procedural difficulty imposes a significant barrier to the feasibility and throughput for

the large-scale application of the assay, underscoring the need for a more streamlined

and optimized approach for the D-SACF assay.

To address these issues, we focused on a novel 3D culture system that utilizes a

commercially available low-molecular-weight agar polymer LA717 (developed by

Nissan Chemical Corporation, Tokyo, Japan)-supplemented medium combined with

low-attachment culture vessels. Previous studies have demonstrated that using liquid

culture media containing LA717 and cultivating cells in low-adhesion containers

enables the even dispersion of cells throughout the culture medium, preventing adhesion

and aggregation of neighboring cells while maintaining contact between daughter cells

after division [11,12]. Thus, this system provides an easier means of establishing a 3D

environment than conventional soft agar preparation, allows for a high-density cell

culture, and may serve as a platform for evaluating anchorage-independent cell growth.

In this study, we used human MSCs as a model of normal cells and HeLa cells as

a   model   of   transformed   cells   to   determine   whether   the   novel

liquid/low-molecular-weight agar colony formation (LACF) approach utilizing LA717

could effectively detect colony formation of malignantly transformed cells through

anchorage-independent growth within the normal cell population. To overcome the

limitations of the soft agar-based D-SACF assay and further improve its practicality and

throughput, we optimized and systematized the protocol, ultimately establishing a

digital analysis system for the LACF assay (D-LACF assay). Herein, we demonstrate

how the D-LACF assay allows for the detection of malignantly transformed cells as

tumorigenic impurities in hCTPs with high sensitivity and reproducibility, providing a

promising new approach for the quality assessment of hCTPs.

Materials and Methods

Cells

All cell cultures were maintained in a humidified atmosphere with 5% CO_2 and 95% air

at 37°C. iCell Human iPSC-derived Mesenchymal Stem Cells (Lot 105305, Fujifilm

Cellular Dynamics, Madison, WI, USA) were cultured using the human Mesenchymal

Stem Cell Growth BulletKit Medium (Lonza, Basel, Switzerland). A human cervical

cancer cell line, wild-type HeLa (CCL-2, Lot 61647128), was obtained from the

American Type Culture Collection (Manassas, VA, USA). We obtained plasmids

pC13N-iCAG.copGFP, pZT-C13-R1, and pZT-C13-L1 described by Cerbini *et al*. [13]

from Addgene (Watertown, MA, USA) and introduced them into HeLa cells by

electroporation. This approach enabled transcription activator-like effector nuclease

(TALEN)-mediated green fluorescent protein (GFP) knock-in at the second intron of

CLYBLE on chromosome 13, which is recognized as a safe harbor site. A stable

GFP-expressing HeLa (HeLa-GFP) cell line was established for use in this study.

Wild-type HeLa (HeLa-WT) and HeLa-GFP cells were cultured in Minimum Essential

Medium (MEM; Sigma-Aldrich, St. Louis, MO, USA) supplemented with 10% fetal

bovine serum (FBS, Sigma-Aldrich), 1% GlutaMAX (Thermo Fisher Scientific,

Waltham, MA, USA), and 1% penicillin-streptomycin (Nacalai Tesque, Kyoto, Japan).

MSCs and HeLa cells pre-labeled with CellVue Claret (CVC) dye (Sigma-Aldrich)

were used in specific experiments following the manufacturer's instructions.

Novel LACF assay

We adopted a 3D culture method using a medium containing 0.03% (w/v) aqueous

solution of LA717, which was supplied as a 1% stock solution under the trade name

SphereMax (Nissan Chemical Corporation, Tokyo, Japan), based on previous research

[11] by Abe-Fukasawa *et al*. This culture method facilitates straightforward 3D cell

culture with a standard cell culture medium augmented by a fixed amount of LA717

(manufacturer's recommended concentration: 0.03%) in low-adhesion culture vessels.

In this study, cells were prepared in a prewarmed medium made up of a basal growth

medium containing GlutaMAX-supplemented Dulbecco's Modified Eagle Medium

(Thermo Fisher Scientific), prewarmed dissolved LA717, 10% FBS, 1% MEM

Non-Essential Amino Acid solution (Thermo Fisher Scientific), and 1%

penicillin-streptomycin to achieve a final concentration of 0.03–0.06% LA717 (w/v).

The cell suspensions were adjusted to a density of 15,000–600,000 cells/mL and

dispensed into 96-well flat-bottom low-adhesion plates (PhenoPlate 384-well

ULA-coated microplates, Revvity, Tokyo, Japan) at 100–400 μL (containing

10,000–60,000 cells) per well. These plates were incubated in a humidified atmosphere

with 5% CO_2 at 37°C for approximately 3 weeks (19–23 days) without any media

changes or addition.

Preparation of co-cultured cells: normal and reference transformed cells

In this study, MSCs were used as the normal cell model, whereas HeLa (-WT or -GFP)

cells served as the reference transformed cell model. Samples were prepared by mixing

these cells at a defined ratio, and the colony formation rate of individual transformed

cells was evaluated under 3D culture conditions. To each well of a 96-well plate, 200

µL of medium supplemented with 0.03% LA717 was added, along with 20,000 MSCs

and 0.125 to 1 HeLa cell. We employed a method in which the reference transformed

cell suspension was initially prepared at a minimum concentration of 100 cells/mL

using a procedure described later to minimize errors associated with repeated dilution

steps. The required volume was then directly taken from this suspension and added to

the MSC suspension, which had been preloaded into a reservoir. After thoroughly

mixing, the suspensions were seeded into multiple wells. Additionally, when sampling a

fixed volume from a low-concentration cell suspension, we considered the Poisson

distribution of the actual cell count along with its 95% confidence interval (CI). The CI

for the Poisson distribution was calculated using the poisson.test function in R (version

3.3.3, R Foundation for Statistical Computing, Vienna, Austria). For instance, when

spiking a suspension expected to contain 100 cells, executing "poisson.test (100,

conf.level = 0.95)" in R yields a 95% CI ranging from 81–122 cells (indicating a

rounded-down or rounded-up value).

Cell counting

The concentration of viable cells in the cell suspension was measured using

NucleoCounter NC-202 Consistent Cell Counter (ChemoMetec, Cambridge, MA, USA)

or Countess 3 FL Automated Cell Counter (Thermo Fisher Scientific). Based on these

results, serial dilutions were performed to prepare suspensions of appropriate

concentrations for each test. When preparing low-concentration HeLa cell suspensions

at 100 cells/mL, a portion of the diluted suspension was used to reconfirm the

concentration using the following method: 1.2 mL of the suspension was collected and

incubated at 37°C for 15 min with 1 µg/mL Hoechst 33342 (Dojindo Laboratories,

Kumamoto, Japan), 1 µg/mL propidium iodide (Dojindo Laboratories), and 1 µg/mL

Calcein-AM (Dojindo Laboratories) (not added when counting HeLa-GFP), and then

dispensed into 96 wells of a 384-well plate (Corning, Corning, NY, USA) at 10 µL each.

After centrifugation of the 384-well plate at $300 \times g$ for 1 min, the presence or absence

of viable cells in each well was analyzed by fluorescence imaging using the confocal

imaging cytometer CellVoyager CQ1 (Yokogawa Electric, Musashino, Japan). Based

on the number of negative wells (k) without cells out of the 96 wells, the average cell

count (λ) per 10 µL was calculated using the formula $\lambda = -ln(k)$, similar to

determining the average copy number per partition in droplet digital PCR (ddPCR) [14].

The final concentration of the suspension was thus determined.

Cell staining

After 3 weeks of incubation, the cells were stained with various reagents under the

following conditions. The LACF assay medium (containing 0.03% LA717) with each

staining dye was carefully added to each well of the plate at 20 µL. The plate was then

incubated at 37°C and 5% CO2 for at least 1 h. The final concentrations of each staining

reagent after addition were as follows: Hoechst 33342 at 1 µg/mL, Calcein-AM at 1 µM,

LysoTracker Red (Thermo Fisher Scientific) at 50 nM, or MitoTracker Deep Red

(Thermo Fisher Scientific) at 50 nM. The images of each well were captured using an

imaging cytometer. In this study, we used two different imaging devices: the

CellVoyager CQ1 (Yokogawa Electric) and the Cell3iMager duos2 (SCREEN Holdings,

Kyoto, Japan). The former was primarily used for acquiring fluorescent images for

analysis, whereas the latter was mainly employed to obtain clear brightfield images of

the entire well. When fixing samples stained with reagents other than Calcein-AM, 15

µL of 16% paraformaldehyde (PFA) solution (Muto Pure Chemicals, Tokyo, Japan)

prewarmed to 37°C was added to each well (with a final PFA concentration of

approximately 1%). The samples were kept in the dark and allowed to stand overnight

at 37°C. Subsequently, the samples were stored in the dark at room temperature. Fixed

samples were imaged within 2 weeks of this procedure.

Image analysis

An imaging cytometer was used to capture images of each well. Excitation light and

fluorescence filters corresponding to the fluorescent proteins expressed in the cells or

the added fluorescent reagents were selected and multichannel images of the entire well,

including bright-field images, were obtained. Furthermore, multiple Z-stack images

were acquired, and images focused on the cell colonies were selected. Automated image

analysis was conducted for each well using an application integrated into the analyzer,

employing an analysis script to segment colony features based on size, circularity, and

fluorescence intensity criteria. In this study, through image analysis, the cell survival

status was determined based on the expression of fluorescent proteins or the staining

reaction of cells to fluorescent reagents. HeLa cell-derived colonies were then defined

and detected based on their size, shape, and circularity. However, in some cases, debris

and other objects were detected as false-positive objects in the image analysis; therefore,

the presence or absence of colonies in each well was ultimately confirmed by visual

inspection based on bright-field images.

Calculation of the colony-forming efficiency (CFE) of reference transformed cells

Calculations were performed to determine the CFE (x) of reference transformed cells

based on the following assumptions:

1. Cells are assumed to be individually dispersed in the suspension without

aggregation.

2. Transformed cells seeded into a fixed number of wells (n) are distributed across the

wells following a Poisson distribution.

3. Under 3D culture conditions, a single transformed cell in a well of a 96-well plate

proliferates and forms a colony with a specific probability ($0 \leq x \leq 1$).

4. When multiple transformed cells are seeded into the same well, they each form

colonies independently with the same probability (x), without influencing one

another.

Using these assumptions, the relationship among the number of fractionated wells

(n), the average number of transformed cells per well (λ), the total number of wells with

colonies (y), and the probability of colony formation (x) is described by the following

equation (see Supplementary Figures 1 and 2 for detailed mathematical derivation and

application examples):

$$y = \sum_{k=1}^{n\lambda} (n \cdot \frac{\lambda^k \, e^{-\lambda}}{k!}) \cdot (1 - (1 - x)^k)$$

To determine x, the upper limit of k was estimated based on the actual number of

fractionated wells (n) and the average number of transformed cells per well (λ). As

shown in Supplementary Figure 1, the value of k was set to a level that ensures the

number of wells with colonies is not zero, regardless of the CFE, based on the seeding

conditions (particularly n and λ) of the reference transformed cells. Calculations for x

were performed using the Goal Seek function (What-If Analysis) in Microsoft Excel

(version 16.96; Microsoft, Redmond, WA, USA) to obtain approximate results.

Statistical analysis

Statistical analyses were performed using Prism 10 software (version 10.3.0; GraphPad

Software, San Diego, CA, USA). Statistical significance was defined as $P < 0.05$. The

effects of culture conditions and independent test trials on the mean CFE of HeLa-GFP

cells under different culture conditions were evaluated by one-way repeated measures

analysis of variance (ANOVA). CFE values are presented as the mean ± standard

deviation or with their corresponding 95% CI.

Results

LA717 enables stable and dispersed 3D MSC cultures at a high cell density

To evaluate the effect of LA717 on the high-density 3D culture of MSCs, cells were

seeded at various densities (10,000–60,000 cells/well) in low-adhesion culture plates. In

the absence of LA717, MSCs exhibited rapid aggregation, forming dense cell clusters

within the first day of culture, regardless of the initial seeding density (Figure 1, left

panel). This aggregation persisted and became more pronounced over time, especially in

the wells with higher cell densities. In contrast, the addition of 0.03% LA717 to the

culture medium significantly suppressed MSC aggregation. The cells remained largely

dispersed across all seeding densities even after 14 days of culture (Figure 1, right

panel). Although slight aggregation was observed at higher cell densities (50,000 and

60,000 cells/well), these aggregates were transient and shrank over time, indicating that

LA717 effectively promoted a stable dispersed state of MSCs under high-density

conditions.

Figure 1. LA717 prevents cell aggregation in high-density three-dimensional (3D) cultures of

human mesenchymal stem/stromal cells (MSCs). Representative whole-well images showing the time

course of MSC cultures over 14 days in low-adhesion 96-well plates under two different 3D culture

conditions. Insets show magnified views of the boxed areas to highlight cell aggregation or distribution.

MSCs were seeded at densities ranging from 10,000 to 60,000 cells/well in Dulbecco's Modified Eagle

Medium (DMEM) supplemented with 10% fetal bovine serum (FBS; left) or DMEM supplemented with

10% FBS and 0.03% LA717 (right; developed by Nissan Chemical Corporation, Tokyo, Japan). In the

absence of LA717, MSCs aggregated rapidly, particularly at higher seeding densities, and formed dense

clusters by day 1 that persisted throughout the culture period. In contrast, the addition of 0.03% LA717

effectively suppressed cell aggregation, maintaining a dispersed distribution of MSCs across all densities

over the entire 14-day period. All images were acquired using the Cell3iMager duos2 (SCREEN Holdings,

Kyoto, Japan). The inner diameter of each well was 6.4 mm.

Culture conditions to optimize high-density 3D MSC culture

In this study, we investigated the effects of LA717 concentration and medium volume

on the aggregation of MSCs in cultures in 3D using high-density cultures of 60,000

cells/well. MSCs cultured without LA717 showed significant aggregation regardless of

the medium volume. However, the addition of LA717 effectively suppressed

aggregation in a concentration-dependent manner, with higher concentrations of LA717

(0.04–0.06%) showing more pronounced effects (Figure 2). Furthermore, large medium

volumes (300–400 μL) were associated with a greater decrease in MSC aggregation,

even at low LA717 concentrations (0.03–0.04%). These results suggest that both the

LA717 concentration and medium volume play crucial roles in preventing MSC

aggregation in high-density 3D culture.

Figure 2. LA717 suppresses MSC aggregation in a concentration- and volume-dependent manner in 3D culture. Representative whole-well images showing the effects of LA717 concentration and culture medium volume on MSC aggregation after 14 days of 3D culture. Insets show magnified views of the boxed areas to highlight cell aggregation or distribution. MSCs (60,000 cells/well) were cultured in low-adhesion 96-well plates with DMEM + 10% FBS containing varying concentrations of LA717 (0–0.06%) and medium volumes (100, 200, 300, and 400 µL). In the absence of LA717, MSCs aggregated extensively regardless of the medium volume. The addition of LA717 progressively suppressed aggregation in a concentration-dependent manner, with higher concentrations (≥0.04%) and

greater medium volumes (≥300 µL) showing improved dispersion. These findings suggested that both

LA717 concentration and medium volume are important parameters for achieving a stable, high-density

3D culture of MSCs. All images were acquired using the Cell3iMager duos2 (SCREEN Holdings). The

inner diameter of each well was 6.4 mm.

A single HeLa cell co-cultured with MSCs in LA717-supplemented medium

proliferated in an anchorage-independent manner and formed a colony

Next, we examined the behavior of MSCs and a single HeLa-GFP cell co-cultured in a

3D environment with medium supplemented with 0.03% LA717 (200 µL per well). All

cells were pre-labeled with the CVC dye for fluorescence imaging (Figure 3).

Bright-field and CVC fluorescence (pseudo-colored in magenta) images demonstrated

that MSCs remained dispersed and showed no anchorage-independent growth under this

3D culture condition. Minor initial aggregation of MSCs was observed, which resolved

over time. In contrast, the single HeLa-GFP cell exhibited anchorage-independent

growth, doubled within 24 h, and continued to proliferate, ultimately forming a distinct

colony. GFP fluorescence highlighted the dynamic HeLa cell colony formation process

over 15 days. However, the fluorescence of the CVC dye in colonies derived from

HeLa-GFP cells almost disappeared, likely due to the continuous dilution of the

fluorescent label as the cells divided. These results suggest that the 3D culture system

utilizing LA717 effectively distinguishes the proliferative capacity of abnormally

transformed cells, such as HeLa cells, from that of normal MSCs. The images in Figure

3 show a section at each time point when time-lapse imaging (360 h, imaging at 1 h

intervals) was performed, and the video showing all the images for each time point

stitched together is shown in the Supplementary Movie.

Figure 3. Transformed HeLa-green fluorescent protein (GFP) cells exhibit anchorage-independent proliferation and colony formation in a 3D culture environment with 0.03% LA717, unlike surrounding MSCs. Representative images extracted from time-lapse imaging performed every hour for 360 h (15 days), showing the behavior of a single HeLa-GFP cell co-cultured with 20,000 MSCs in 200

µL of LA717 (0.03%)-supplemented medium in a low-adhesion 96-well plate. All cells were pre-labeled

with CellVue Claret (CVC; pseudo-colored magenta) for fluorescence visualization. Bright-field (BF),

GFP, CVC, and merged images at six time points revealed that, while MSCs remained dispersed and did

not proliferate, the single HeLa-GFP cell consistently proliferated in an anchorage-independent manner,

ultimately forming a distinct colony. The GFP signal expanded over time, highlighting the dynamic

growth of the transformed cells. A stitched movie of all time points is provided in the Supplementary

Movie. Images were acquired using a CellVoyager CQ1 (Yokogawa Electric). The inner diameter of each

well was 6.4 mm.

Single HeLa cells consistently formed colonies with a steady efficiency across

different 3D culture conditions in the presence of LA717

We previously established a method for calculating the CFE of reference transformed

cells, which serves as an indicator of the detection performance of transformed cells in

the D-SACF assay (an approach that allows the efficient analysis of a large number of

cells). Conventionally, the CFE is determined based on the total number of counted

colonies obtained from multi-well image analysis. However, in this study, we developed

and implemented a new approach whereby the CFE is calculated based on the total

number of wells containing at least one colony rather than the number of counted

colonies. Using this method, we assessed the CFE of HeLa-GFP cells under various

culture conditions.

First, we evaluated the CFE of HeLa-GFP cells co-cultured with MSCs in a

medium containing LA717 at varying concentrations (0.03–0.06%). We seeded cells

into a total of 240 wells (60 wells × 4 plates) under conditions where each well had 200

µL of medium with 20,000 MSCs and an average of 0.5 HeLa-GFP cells (i.e., $\lambda = 0.5$

cells/well). The cells were then co-cultured in a 3D environment, and colony analysis

was performed after 3 weeks. The results of three independent runs for calculating the

CFE of HeLa-GFP cells at four different concentrations of LA717 (0.03%, 0.04%,

0.05%, and 0.06%) are shown in Figure 4a. The data were analyzed using one-way

repeated measures ANOVA, which revealed no significant effects of LA717

concentration on the CFE of HeLa-GFP cells. These findings indicate that the CFE of

HeLa-GFP cells remained stable regardless of the LA717 concentration. This stability

suggests that LA717 did not influence the anchorage-independent growth capacity of

HeLa-GFP cells within the tested concentration range.

Next, to evaluate the impact of seeding conditions on the CFE of HeLa-GFP cells,

we conducted three independent runs under fixed culture conditions: 200 µL of the

medium containing 0.03% LA717 and 20,000 MSCs per well. We varied the seeding

ratio of HeLa-GFP cells (λ = 0–1.0 cells/well) and the total number of seeded wells

(60–840 wells). After 3 weeks of culture, the CFE was analyzed under each condition.

The results showed that the CFE of HeLa-GFP cells remained consistent across all

tested seeding conditions (Figure 4b). Statistical analysis using one-way repeated

measures ANOVA revealed no significant effect of the HeLa-GFP seeding ratio on the

CFE. These findings indicate that under the examined conditions, the evaluation of CFE

of reference transformed cells is robust and reliable. Importantly, under control

conditions (λ = 0; MSCs only, 60 wells), no colony formation was observed, confirming

the specificity of colony detection for HeLa-GFP cells.

Figure 4. The colony-forming efficiency (CFE) of HeLa-GFP cells co-cultured with MSCs in

LA717-supplemented medium under various culture conditions. (a) The CFE of HeLa-GFP cells was

evaluated under four different concentrations of LA717 (0.03%, 0.04%, 0.05%, and 0.06%) in 200 µL of

culture medium per well. HeLa-GFP cells (0.5 cells/well on average) were co-cultured with 20,000 MSCs

per well in a 96-well plate (n = 240 wells per condition) for 3 weeks. CFE was calculated based on the

number of wells containing colonies derived from HeLa-GFP cells, as determined by image analysis. Bar

graphs represent the mean and standard deviation (SD) values from three independent experiments, and

the colored lines show the individual results from each experiment. **(b)** The CFE of HeLa-GFP cells was

evaluated under varying cell input amounts (λ = 0.125, 0.25, 0.5, and 1.0) in 200 µL of medium

containing 0.03% LA717 and 20,000 MSCs per well. HeLa-GFP cells were seeded into 840, 420, 240,

and 120 wells, respectively, and cultured for 3 weeks. The CFE was calculated from the number of wells

containing colonies detected by image analysis. Bars represent the mean and SD values of three

independent experiments, and colored lines indicate the individual results. No colony formation was

observed under control conditions (λ = 0; i.e., MSCs only, 60 wells). Detailed numerical data for each

condition in **(a)** and **(b)** are provided in Supplementary Tables 1 and 2, respectively. The equation used to

calculate CFE in **(a)** and **(b)** is described in Supplementary Figures 1 and 2, respectively. LA, LA717.

398   Supplementary Tables 1 and 2 present the CFE of the reference transformed cells

399 under each culture condition, calculated using two distinct methods: (1) based on the

400 total number of wells containing at least one colony, as applied in this study, and (2)

401 based on the conventional method of counting the total number of colonies. The results

402 from both methods were consistent, indicating that the CFE of the reference

403 transformed cells remained nearly identical regardless of the calculation method

404 employed.

406 ***Improved visualization and detection of colonies using fluorescence staining in***

407 ***LA717-based 3D culture***

408 To enhance the efficiency and accuracy of colony detection through the image analysis

409 of whole-well images in a 96-well plate, we explored the application of fluorescence

410 staining in cultured samples. Based on our previous studies, we evaluated nuclear,

411 mitochondrial, and lysosomal fluorescent labeling for this purpose. Figure 5a shows

412 representative images of a cell colony observed after co-culturing 0.5 HeLa-GFP cells

413 with 20,000 MSCs per well in a medium supplemented with 0.03% LA717 for 3 weeks.

414 The presence of GFP fluorescence in the cell aggregates identified in the bright-field

415 images confirmed that these colonies originated from HeLa-GFP cells. The addition of

fluorescent dyes specific to the nucleus, mitochondria, and lysosomes during cell

culture enabled the organelle staining of live cells, leading to clear fluorescent labeling

of HeLa-GFP-derived colonies. Among the dyes used, LysoTracker and MitoTracker

are selective for live cells, which indicated that the colonies consisted of viable cells.

Figure 5b shows colonies formed by HeLa-WT cells under the same culture conditions.

Live staining with Calcein-AM resulted in strong fluorescence signals, confirming that

the colonies were composed of viable cells. Based on the findings from HeLa-GFP cells,

the colonies observed under these conditions were likely derived from HeLa-WT cells.

Fluorescent staining has proven useful for image-based colony detection. To

enhance its applicability, we further investigated fixation methods for stained cells. As

shown in Figure 5c, careful temperature management during fixation allowed

fluorescence signals from LysoTracker and Hoechst staining (left images show live

colony staining before fixation) to be preserved for at least 14 days (right images show

the fixed colony stained with LysoTracker and Hoechst dyes). This approach increases

the flexibility of the analytical process by enabling imaging-based colony evaluation

over an extended period, thereby improving the practicality of the method.

These fluorescent labeling techniques enabled clear visualization of colonies

derived from transformed cells in the LA717-based 3D culture environment.

Consequently, similar to the previous D-SACF assay, detecting transformed

cell-derived colonies and evaluating the CFE of reference cells through image analysis

became readily achievable.

Figure 5. Fluorescence-based detection of colonies formed by single HeLa cells co-cultured with

MSCs in LA717-supplemented medium. Representative images of colonies formed by a single

HeLa-GFP or wild-type HeLa cell co-cultured with 20,000 MSCs per well for 3 weeks in 0.03%

LA717-supplemented medium (200 µL/well). Fluorescent staining markedly enhanced the accuracy of

colony detection in image analysis. Images were acquired using the CellVoyager CQ1 confocal imaging

cytometer (Yokogawa Electric). **(a)** A colony derived from a single HeLa-GFP cell. Live staining was

performed by adding Hoechst 33342 (nucleus), MitoTracker Deep Red (mitochondria), or LysoTracker

Red (lysosomes) to the culture medium and incubating the cells for a defined period. The GFP signal

confirmed that the colony originated from HeLa-GFP cells, which was also clearly visualized with each

dye. **(b)** A colony derived from a single wild-type HeLa cell. Live staining with Hoechst 33342 and

Calcein-AM revealed strong fluorescence signals, indicating colony viability. **(c)** (Left) A wild-type live

HeLa colony stained with Hoechst 33342 and LysoTracker Red. (Right) The same colony was imaged 14

days after fixation with 1% paraformaldehyde (PFA). Fluorescence signals from Hoechst and

LysoTracker staining were retained, indicating that this dye combination is compatible with fixation and

long-term storage, allowing flexible scheduling of image acquisition and analysis. The inner diameter of

each well was 6.4 mm.

Discussion

A previous study conducted by the developers of LA717 [11] demonstrated that

culturing cells in low-attachment vessels with LA717-supplemented medium ensures

uniform cell dispersion, prevents adhesion and aggregation, and maintains contact

between newly divided cells while minimizing medium convection. These properties

allow cells and spheroids to stay in place and form the foundation of a new 3D culture

system. Building on this, we showed that MSCs, which typically form aggregates at

high densities, remained stably dispersed without proliferating in the LA717-containing

medium, supporting its use for scalable and uniform MSC culture. In contrast, HeLa

cells exhibited anchorage-independent growth from single cells and consistently formed

colonies under the same conditions. Fluorescent labeling enhanced the visualization of

colonies from transformed cells, making image-based colony detection easier and

supporting CFE evaluation with reference cells such as HeLa cells. The LACF assay,

developed using this 3D culture system with LA717, demonstrated practical advantages

over the SACF assay in terms of operability and analysis (Supplementary Table 3). We

further established the D-LACF assay to digitally measure the proportion of

transformed cells in hCTPs by detecting single-cell-derived colonies (Figure 6).

Quality control (QC) of the assay itself, using MSCs as a cell product and HeLa

cells as the reference, requires a 3-week co-culture period, followed by staining, fixation,

imaging, and analysis, which is typically completed within the same day or a few days.

Based on this workflow, we apply product testing under the same conditions. In our

prior D-SACF study, we developed a method to calculate the required sample size (i.e.,

the number of partitioned wells) to detect at least one colony at a predefined

false-negative rate (FNR), based on the CFE of the reference cells [10]. This approach

allows for the theoretical estimation of the assay scale needed to achieve a target

detection sensitivity and FNR for QC samples containing transformed cells at a known

frequency. It also enables the prediction of the probability of detecting at least one

colony when using QC samples, given the pre-calculated sample size. In product testing,

applying the same seeding conditions and number of wells helps determine whether

transformed cells comparable to the reference are present and if their proportion exceeds

the predefined detection limit—information that can be considered a quality attribute of

the product. This method is equally applicable to the D-LACF assay, supporting rational,

quantitative test design for assessing transformed cells in hCTPs under defined

conditions. By routinely monitoring assay performance, especially through system

suitability testing with reference cells, the D-LACF assay can be used for lot-release

testing of cell products. Its improved feasibility and scalability over the D-SACF assay

support large-scale screening of malignantly transformed cells in hCTPs.

Figure 6. Schematic overview of the digital liquid/low-molecular-weight agar colony formation

(D-LACF) assay. The D-LACF assay can be used to detect transformed cells by combining LA717-based

3D culture with image□based digital analysis. Section A illustrates the evaluation of the assay

performance using quality control (QC) samples: (1) assay design, (2) assay execution including split

culture, 3D growth in LA717-supplemented medium, fluorescent labeling, and high-throughput colony

detection, and (3) determination of the sample size based on the observed detection performance. Section

B presents product testing conducted under the same culture conditions but with the predetermined

sample size (number of wells) calculated in Section A; each well can then be classified as positive (P) or

negative (N) for transformed cell-derived colonies. This figure was created with BioRender.com.

The growth in low attachment assay (GILA) is another *in vitro* 3D culture system

for transformation assessment using only standard medium and low-adhesion vessels

[15,16]. It reportedly detects gene mutation-induced transformation with sensitivity

comparable to SACF assay [17]. The possibility of multiple assay options is

advantageous as it enables the selection of the most appropriate method based on

various requirements. The standard GILA protocol recommends seeding several

hundred to several thousand cells per well in a 96-well plate. However, in this study,

when MSCs were seeded at 10,000 cells/well in a conventional culture medium,

pronounced cell aggregation was observed. In contrast, the LA717-supplemented

medium effectively suppressed MSC aggregation under the same seeding conditions.

These results suggest that for quality assessment of hCTPs composed of a large number

of cells—particularly for detecting a trace amount of transformed cells—GILA may not

suitable for this purpose due to its limitations under high-density culture conditions.

Instead, the LACF assay, which enables high-density culture by modifying the culture

medium composition, is currently the most practical and effective method for such

applications.

The D-SACF/D-LACF assay share core principles with ddPCR and is structured

around three main components: (1) Fractionation of test cells into multi-well plates for

3D culture to enrich target transformed cells and improve the signal-to-noise ratio for

image analysis; (2) Use of high-throughput imaging systems for efficient multi-well

analysis; and (3) Digital detection of colony presence or absence in each well to

estimate the proportion of transformed cells. Similar to ddPCR, increasing the number

of partitions theoretically improves the detection of rare transformed cells, enhancing

sensitivity and lowering the limit of detection (LOD). However, in the D-LACF assay,

each well must receive a fixed number of total cells (including both normal and

transformed cells), followed by 3D culture and colony assessment. This imposes

constraints on optimal culture volume and cell density per well. Although switching

from 96- to 384-well plates could improve resolution and throughput, challenges such

as reduced seeding volume, higher risk of medium evaporation, and variations in cell

growth might outweigh these benefits. Nonetheless, optimizing culture formats,

including plate types, remains a crucial area of research for future assay automation. A

key difference from ddPCR is that while a target sequence in ddPCR is almost always

amplified, making the partition positive, not all transformed cells in D-SACF/D-LACF

necessarily form colonies. Thus, estimating their presence ratio based solely on colony

detection is difficult. This reflects a key challenge in detecting transformed cells with

unknown properties in a product. To address this, the D-SACF/D-LACF assay uses a

known transformed cell line, such as HeLa, as a reference and positive control. Its CFE

is pre-estimated and used as a performance indicator for the assay.

In this study, we improved part of the performance evaluation process from the

d-SACF assay. While assessing the CFE of reference cells, a key performance indicator,

we focused on two operational challenges: the difficulty of preparing low-concentration

cell suspensions and the CFE variability potentially caused by the cell spiking process.

To improve the accuracy and consistency of the CFE assessment, we optimized the

protocol for these steps. We refined the counting process by staining cells in 384-well

plates and using image-based counting, which partially automates the workflow. We

also incorporated Poisson distribution modeling and its 95% CI to account for sampling

error during spiking, enabling a more precise definition of the expected range in

performance metrics and enhancing analytical accuracy. Additionally, we developed a

simplified way for calculating CFE. Since tracking cell-to-colony conversion precisely

is impractical, we initially calculated the CFE as the ratio of the estimated number of

seeded cells to the total number of colonies. However, image analysis is often

complicated by overlapping or touching colonies, making exact counting difficult. To

address this, we developed an alternative approach: estimating CFE based on the

number of wells containing at least one colony. This method requires only binary

detection of colonies (presence or absence), significantly simplifying image analysis.

CFE values obtained with both methods were consistent and stable across various

conditions, including different LA717 concentrations and HeLa-GFP seeding densities.

These results demonstrate the robustness and reliability of the proposed approach,

which we have adopted as the standard.

Colony imaging and analysis in the D-LACF assays can be performed using a

standard fluorescence imaging system with multi-well scanning capabilities. In addition

to previously used systems (IN Cell Analyzer 2000 [Cytiva], Cellomics ArrayScan

[Thermo Fisher Scientific], and CellVoyager CQ1 [Yokogawa Electric]), this study

utilized the Cell3iMager duos2 (SCREEN Holdings), which demonstrated comparable

feasibility. To facilitate the widespread implementation of the D-LACF assays across

laboratories, the minimum requirements for an appropriate imaging system are outlined

in Supplementary Table 4. Image analysis remains a limiting factor. False positives can

arise from fluorescent debris or cell aggregates, which are common with dyes like

Hoechst 33342 or LysoTracker, or from insufficient mixing of the suspension. We

addressed this by pre-labeling cell membranes with CVC, which allowed for the

distinction between aggregates and proliferative colonies. Aggregates showed stable

fluorescence but no growth; true colonies exhibited decreasing signal due to division.

Incorporating metrics like total or mean fluorescence intensity could improve

identification accuracy. False negatives are actual colonies that go unnoticed. Our

current protocol focuses on decreasing false negatives, even if it causes more false

positives. Therefore, we use permissive image analysis settings, followed by manual

review of fluorescence and bright-field images to exclude artifacts through a two-step

verification process. To further enhance accuracy and scalability, advanced techniques

using more specific fluorescent dyes or artificial intelligence (AI)-based label-free

analysis will be important. Such methods could decrease both false positives and false

negatives, making the D-LACF assay more robust and dependable.

In product cell testing, the ideal outcome is that no colonies are detected. However,

if colony-like structures are observed and their status as true or false positives is

uncertain, the LACF assay offers a complementary option that the conventional SACF

assay lacks. Because the LA717-based 3D culture system uses a liquid medium, suspect

cell aggregates can be easily isolated and retrieved with a micropipette. The collected

cells can then be sub-cultured to determine whether they proliferate, allowing for further

characterization to establish whether they are truly malignant. Moreover, a detailed

analysis of any unexpectedly tumorigenic cells may provide new opportunities to

investigate the mechanisms of cellular transformation and develop strategies for its

suppression, thereby widening the research applications of this novel 3D culture

platform.

To facilitate broader adoption and practical use of the D-LACF assay, a systematic,

stepwise approach—including inter-laboratory validation and standardization—is

essential. Each participating facility should initially verify the assay's reproducibility

and feasibility using a standardized core protocol. We recommend referring to

Supplementary Table 5, which complements Figure 6 and highlights key considerations

and practical tips for implementing the D-LACF assay across laboratories. Overall,

these efforts will help ensure the assay's reliability and support its international

standardization.

Method validation should then be conducted according to ICH Q2(R1) guidelines

[19] . Since the D-LACF assay acts as a "limit test" within the "purity testing" category

of QC, validation should mainly emphasize specificity and the LOD. In this study, we

developed the D-LACF platform and demonstrated its feasibility using model cells: an

iPSC-derived MSC line and HeLa cells. However, to assess specificity, it is necessary

to evaluate additional tumor-derived cell lines and confirm that their colony formation

can be detected. Although previous work showed that several cancer cell lines can grow

in LA717-supplemented 3D culture [11], their CFEs have not been systematically

characterized.

As a preliminary study, we evaluated the performance of the D-LACF assay using

HEK-293 cells, a tumorigenic cell line different from HeLa (Supplementary Figure 3).

Although the observed CFE of HEK-293 cells was lower than that of HeLa, consistent

colony formation was still detected. These findings suggest the potential usefulness of

the D-LACF assay for detecting various transformed cells. Further studies with different

tumorigenic cell lines will be needed to assess their colony-forming capabilities, along

with quantitative evaluations of spike-in concentrations and detection accuracy.

In the earlier D-SACF study, multiple MSC lots showed no lot-dependent effect

on HeLa colony formation [10]. In this study, the iPSC-derived MSC line was chosen

for its high proliferative capacity and consistent stock availability. Both D-SACF and

D-LACF assays, which are based on similar principles, consistently showed that MSCs

have no inhibitory effects on colony formation by HeLa cells. However, donor

variability among MSCs has not been systematically investigated. Future studies should

incorporate additional MSC lots, including clinical-grade cell batches, to assess assay

robustness. Additionally, the applicability of the D-LACF assay to hCTP models

beyond MSCs—such as neural or cardiac cell-based products—should be explored. For

such validation, harmonizing reagents, cell sources, culture conditions, and assay design

is critical. Inter-laboratory studies will be essential to confirm the reproducibility of key

metrics, such as CFE.

In summary, the D-LACF workflow developed here can serve as a platform for

designing QC protocols for various hCTPs. A key benefit of the assay is its

flexibility—test parameters such as seeding density, well number, and LA717

concentration can be adjusted based on product characteristics. The D-LACF assay is

anticipated to improve the quality and safety of hCTPs, supporting the development of

more reliable therapies.

Conclusion

In this study, we established the D-LACF assay as a practical and sensitive platform for

detecting trace amounts of malignantly transformed cells in hCTPs. By combining a

novel LA717-based 3D culture system with digital image analysis and refined

evaluation protocols, the D-LACF assay offers improved feasibility, accuracy, and

scalability compared with conventional methods. These findings provide a foundation

for the broader implementation of the D-LACF assay in quality testing frameworks and

contribute to enhancing the safety and reliability of future hCTPs. Moving forward,

continued optimization and inter-laboratory validation of this assay will be crucial for

its practical implementation and international standardization as a QC tool for advanced

cell-based therapies.

Declaration of Competing Interests

The authors have no commercial, proprietary, or financial interest in the products,

methodologies, or companies described in this article.

Funding

This work was supported by JSPS KAKENHI Grant Number JP18K12137 to SK, by

the Japan Agency for Medical Research and Development (Grant Numbers

JP18mk0104118 and JP24mk0121279 to YS and SY, respectively), and by the Japanese

Ministry of Health, Labour and Welfare (MHLW) Grants-in-Aid for Scientific Research

(23KC5002 to SY).

Author Contributions

Study conception and design: SK and SY. Data acquisition: SK and TY. Analysis and

interpretation of data: All authors. Manuscript drafting and revision: SK and SY. All the

authors have approved the final manuscript.

Acknowledgments

We thank Keiichiro Otsuka and Taito Nishino from Nissan Chemical Corporation for

their help and support in establishing the 3D culture method using LA717. We thank

Atsune Kamada and Prof. Dr. Mitsutoshi Satoh of Meiji Pharmaceutical University for

generating the preliminary data that laid the foundation for this study. We also express

our gratitude to Takashi Yoshiura and Takayoshi Matsubara from Yokogawa Electric

for their dedicated efforts in supporting and coordinating image acquisition and analysis

with CellVoyager CQ1 (Yokogawa Electric). Additionally, we thank Yuki Mori and

Kazuo Onishi from SCREEN Holdings for their valuable contributions to image

acquisition and analysis using Cell3iMager duos2 (SCREEN Holdings).

Supplementary material

Supplementary materials associated with this article can be found in the online version

at doi:

References

[1] Frisch SM, Ruoslahti E. Integrins and anoikis. Curr Opin Cell Biol 1997;9:701–6.

https://doi.org/10.1016/s0955-0674(97)80124-x.

[2] Frisch S, Francis H. Disruption of epithelial cell-matrix interactions induces

apoptosis. J Cell Biol 1994;124:619–26. https://doi.org/10.1083/jcb.124.4.619.

[3] Larco JE de, Todaro GJ. Growth factors from murine sarcoma virus-transformed

cells. Proc Natl Acad Sci U S A 1978;75:4001–5.

https://doi.org/10.1073/pnas.75.8.4001.

[4] Roberts AB, Anzano MA, Wakefield LM, Roche NS, Stern DF, Sporn MB. Type

beta transforming growth factor: A bifunctional regulator of cellular growth. Proc Natl

Acad Sci U S A 1985;82:119–23. https://doi.org/10.1073/pnas.82.1.119.

[5] Puck TT, Marcus PI, Cieciura SJ. Clonal growth of mammalian cells in vitro;

growth characteristics of colonies from single HeLa cells with and without a feeder

layer. J Exp Med 1956;103:273–83. https://doi.org/10.1084/jem.103.2.273.

[6] Du F, Zhao X, Fan D. Soft Agar Colony Formation Assay as a Hallmark of

Carcinogenesis. Bio Protoc 2017;7:e2351. https://doi.org/10.21769/bioprotoc.2351.

[7] Borowicz S, Scoyk MV, Avasarala S, Rathinam MKK, Tauler J, Bikkavilli RK, et al.

The   Soft   Agar   Colony   Formation   Assay.   J   Vis   Exp   2014:e51998.

https://doi.org/10.3791/51998.

[8] Kusakawa S, Machida K, Yasuda S, Takada N, Kuroda T, Sawada R, et al.

Characterization of in vivo tumorigenicity tests using severe immunodeficient

NOD/Shi-scid IL2Rγnull mice for detection of tumorigenic cellular impurities in human

cell-processed   therapeutic   products.   Regen   Ther   2015;1:30–7.

https://doi.org/10.1016/j.reth.2014.12.001.

[9] Kusakawa S, Yasuda S, Kuroda T, Kawamata S, Sato Y. Ultra-sensitive detection of

tumorigenic cellular impurities in human cell-processed therapeutic products by digital

analysis   of   soft   agar   colony   formation.   Sci   Rep   2015;5:17892.

https://doi.org/10.1038/srep17892.

[10] Bando K, Kusakawa S, Adachi H, Yamamoto M, Iwata M, Kitanaka A, et al.

Protocol improvement and multisite validation of a digital soft agar colony formation

assay for tumorigenic transformed cells intermingled in cell therapy products. Cytotherapy 2024;26:769–77. https://doi.org/10.1016/j.jcyt.2024.03.005.

[11] Abe-Fukasawa N, Otsuka K, Aihara A, Itasaki N, Nishino T. Novel 3D Liquid Cell Culture Method for Anchorage-independent Cell Growth, Cell Imaging and Automated Drug Screening. Sci Rep 2018;8:3627. https://doi.org/10.1038/s41598-018-21950-5.

[12] Matsuzaka Y, Kusakawa S, Uesawa Y, Sato Y, Satoh M. Deep Learning-Based In Vitro Detection Method for Cellular Impurities in Human Cell-Processed Therapeutic Products. Appl Sci 2021;11:9755. https://doi.org/10.3390/app11209755.

[13] Cerbini T, Funahashi R, Luo Y, Liu C, Park K, Rao M, et al. Transcription Activator-Like Effector Nuclease (TALEN)-Mediated CLYBL Targeting Enables Enhanced Transgene Expression and One-Step Generation of Dual Reporter Human Induced Pluripotent Stem Cell (iPSC) and Neural Stem Cell (NSC) Lines. PLoS One 2015;10:e0116032. https://doi.org/10.1371/journal.pone.0116032.

[14] Huggett JF, Foy CA, Benes V, Emslie K, Garson JA, Haynes R, et al. The digital MIQE Guidelines: Minimum Information for Publication of Quantitative Digital PCR

Experiments. Clin Chem 2013;59:892–902.

https://doi.org/10.1373/clinchem.2013.206375.

[15] Izar B, Rotem A. GILA, a Replacement for the Soft☐Agar Assay that Permits

High☐Throughput Drug and Genetic Screens for Cellular Transformation. Curr Protoc

Mol Biology 2016;116:28.8.1-28.8.12. https://doi.org/10.1002/cpmb.26.

[16] Rotem A, Janzer A, Izar B, Ji Z, Doench JG, Garraway LA, et al. Alternative to the

soft-agar assay that permits high-throughput drug and genetic screens for cellular

transformation. Proc Natl Acad Sci U S A 2015;112:5708–13.

https://doi.org/10.1073/pnas.1505979112.

[17] Lemmens M, Fischer B, Zogg M, Rodrigues L, Kerr G, Rio-Espinola A del, et al.

Evaluation of two in vitro assays for tumorigenicity assessment of CRISPR-Cas9

genome-edited cells. Mol Ther - Methods Clin Dev 2021;23:241–53.

https://doi.org/10.1016/j.omtm.2021.09.004.

[18] Hays A, Wissel M, Colletti K, Soon R, Azadeh M, Smith J, et al.

Recommendations for Method Development and Validation of qPCR and dPCR Assays

in Support of Cell and Gene Therapy Drug Development. AAPS J 2024;26:24.

https://doi.org/10.1208/s12248-023-00880-9.

[19] ICH Harmonised Tripartitte Guideline - Validation of Analytical Procedures:Text

and Methodology Q2(R1),

https://database.ich.org/sites/default/files/Q2%28R1%29%20Guideline.pdf 2005

[accessed 26.6.25].