Nickel binding to a split ATCUN motif in c-Src SH3 domain facilitates crystallization

Xander Calicdan [a], Oriana S. Fisher [b], Byung Hak Ha [c], Titus J. Boggon [c,d]
and Amy L. Stiegler [3,*]

[a] Department of Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT, USA
[b] Department of Molecular Biology and Biochemistry, Wesleyan University, Middletown, CT, USA
[c] Department of Pharmacology, Yale University, New Haven, CT, USA
[d] Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, CT, USA
[*] Correspondence e-mail: amy.stiegler@yale.edu

ABSTRACT

Numerous X-ray crystal structures of the c-Src SH3 domain have provided a large sampling of atomic-level information for this important signaling domain. Multiple crystal forms have been reported, with variable crystal lattice contacts and chemical crystallization conditions. Here, we report a unique crystal structure of Src SH3 domain in trigonal space group $H3_2$ to 1.45 Å resolution. Crystal packing and anomalous scattering reveal that this crystal form is mediated by two ordered nickel ions provided by the crystallization buffer. Nickel coordination occurs in a 2:2 stoichiometry which dimerizes two SH3 domain monomers across a pseudo-twofold rotation axis, and involves the native N-terminal c-Src SH3 amino acid sequence, a surface-exposed histidine residue, and ordered water molecules. This provides an example of a split-ATCUN (amino terminal copper and nickel) motif that contrasts previous Src SH3 crystal forms containing monomeric ATCUN motifs and may provide an alternative mode of protein N-terminal metal binding.

Keywords: SH3 domain; c-Src tyrosine kinase; nickel coordination; crystal structure; anomalous signal

PDB References: 9OFX

1. INTRODUCTION

Cellular signaling proteins are frequently comprised of multiple modular domains with independent fold and function (Vogel et al., 2004). Among these protein domains, the Src homology 3 (SH3) domain is one of the most abundant (Kaneko et al., 2008). As its name implies, the SH3 domain was first identified as a conserved region shared among the amino acid sequences of cellular Src (c-Src) tyrosine kinase and the related cellular Abelson tyrosine kinase (c-Abl), phospholipase C, and the adapter protein Crk (Mayer et al., 1988). In total, 298 SH3 domains were recently reported in 221 human proteins (Mehrabipour et al., 2023). SH3 domains are also identified in all eukaryotes as well as in some prokaryotes and viruses (Mehrabipour et al., 2023). These domains typically mediate protein-protein interactions and affect enzyme regulation, substrate binding, protein localization, and motility, regulating pathways that impact many cellular processes including proliferation, migration, endocytosis, and actin cytoskeleton rearrangement (recently reviewed in (Kazemein Jasemi et al., 2024)).

The SH3 domain is a small protein fold comprised of approximately 60 amino acids. The first experimental three dimensional structures of SH3 domains were from the Src Family Kinases (SFKs) Fyn (Noble et al., 1993) and c-Src (Yu et al., 1992), and from alpha-spectrin (Musacchio et al., 1992). Together, these structures revealed a conserved SH3 fold comprising two small beta sheets with loops of variable lengths. This fold typically recognizes polyproline-II helical P-x-x-P motifs (or simply 'PxxP', where x is any amino acid) but can also bind other protein targets in atypical interactions (Chau et al., 2022; Kaneko et al., 2008; Saksela and Permi, 2012; Teyra et al., 2017). SH3 domain binding to polyproline motifs has been well studied by structural techniques including X-ray crystallography that have collectively helped resolve the binding site, specificity determinants, and bidirectionality of peptide binding (Feng et al., 1994; Lim et al., 1994; Musacchio et al., 1994; Yu et al., 1994; Yu et al., 1992).

The founding SH3 domain was identified in c-Src (Mayer et al., 1988) and has been shown to play a major role in autoinhibition of its kinase activity. In addition to the SH3 domain, SFKs include three other conserved domains: Src homology 1 (SH1) that contains the kinase domain, Src homology 2 (SH2) that binds directly to phosphorylated tyrosine residues, and a short Src homology (SH4) region at the N-terminus which is lipid modified (reviewed in (Boggon and Eck, 2004)). In the absence of activating signals, intradomain interactions maintain Src in an autoinhibited "closed" conformation; specifically, the SH2 domain binds an inhibitory

phosphotyrosine pY-527 in the C-terminal tail, while the SH3 domain binds the linker between SH2 and kinase domains at a non-proline helical-like sequence (Xu et al., 1997). In this conformation, the SH3 and SH2 domains impinge on the kinase domain and restrict it to its inactive conformation. In response to signaling, these SH3- and SH2-mediated autoinhibitory restraints are released, and the kinase domain adopts its active conformation. The SH3 domain of SFKs binds protein targets including proline enriched phosphatase (PEP) (Cloutier and Veillette, 1996), the adapter protein Crk-II (Anafi et al., 1996), the Nef protein of human immunodeficiency virus-1 (HIV-1) (Saksela et al., 1995), the 3BP1 RhoGAP (Cicchetti et al., 1992), choline kinase alpha (ChoKα) (Kall et al., 2019), and beta-arrestin (Luttrell et al., 1999).

The three-dimensional structure of the isolated SH3 domain of c-Src has been reported by numerous X-ray crystallographic and NMR studies (Mayer and Eck, 1995). Among the X-ray crystal structures available for Src SH3 in the Protein Data Bank, space group identity, unit cell dimensions, and/or lattice contacts are often replicated, presumably owing to the similarity in crystallization conditions or in the sequence of the recombinant protein. Here, we report an X-ray crystal structure of the c-Src SH3 domain in space group $H3_2$, which has not previously been observed. In the structure we find lattice contacts that are mediated by interchain nickel coordination. This coordination is achieved by the native protein sequence and demonstrates that native amino acids can serve as nickel coordinating residues. This crystal packing arrangement may widen the potential for future crystallography-based studies of the c-Src SH3 domain.

2. MATERIALS AND METHODS

2.1 Expression construct and expression

Human c-Src (UNIPROT ID: P12931) cDNA encoding the SH3 domain residues 85-143 was PCR-amplified using Platinum *Taq* HiFi polymerase (ThermoFisher) using a forward oligo containing an NdeI restriction site and encoding for a tobacco etch virus (TEV) protease recognition site, and a reverse oligo containing an XhoI restriction site. The PCR product and pET-28a(+) plasmid were separately doubly digested with NdeI/XhoI in CutSmart buffer (New England Biolabs), and ligated together using T4 DNA ligase (New England Biolabs) to yield an expression construct for N-terminal His$_6$-tagged protein with both thrombin and TEV protease recognition sequences. A positive clone was verified by whole-plasmid sequencing (Plasmidsaurus). See Table 1. Protein expression was carried out in Rosetta(DE3) *E. coli* cells transformed with the expression plasmid. Cells were cultured in the presence of 25 µg/ml kanamycin and 17 µg/ml chloramphenicol at 37°C shaking at 200 rpm in Miller's LB Broth (Fisher Scientific) to an OD$_{600}$ of 0.8, then cooled to 18°C prior to induction of protein expression with 0.2 mM Isopropyl β-D-1-thiogalactopyranoside (IPTG). Expression was conducted overnight (approximately 20 hours) at 18°C shaking at 200 rpm. Cells were harvested by centrifugation.

2.2 Protein purification

Cells were resuspended in lysis buffer (500 mM NaCl and 50 mM HEPES pH 7.5) and flash frozen in a dry ice/ethanol bath. The sample was thawed and lysozyme added to a final concentration of 0.1 mg/ml, followed by two more cycles of freeze/thaw. The sample was lysed by sonication for 3 cycles of 2 minutes using 10 second on/off pulses at 35 % with a half-inch probe (Q500, QSonica). DNase I (bovine pancrease, GoldBio) was added to 10 Units/ml and incubated 10 min on ice. The total lysate was clarified by centrifugation for 1 hr at 4°C at 32,928 x *g*. Clarified lysate was applied to 1 ml Ni-NTA agarose beads (Qiagen) in a sealed gravity flow column and incubated rocking for 1 hour at 4°C. The flow-through sample was collected, and beads washed with wash buffer (lysis buffer containing 20 mM imidazole). Recombinant Tobacco Etch Virus (rTEV) protease was added to the bead-bound His$_6$-tagged Src SH3 protein and incubated overnight at 4°C. The liberated untagged c-Src SH3 domain protein was then collected off the bead by washes with 1 ml aliquot of wash buffer and further purified by size exclusion chromatography (Superdex 75 Increase 10/300 GL) in buffer containing 150 mM NaCl, 20 mM Tris pH 7.5. The purified protein was concentrated in a spin concentrator (Amicon Ultra 3 kDa MWCO, Millipore Sigma) to a final concentration of 11 mg/ml. Protein concentration was determined by NanoDrop (Thermo Fisher)

A280 nm blanked with size exclusion buffer using an extinction coefficient of 16,960 M^{-1} cm^{-1} calculated using the primary amino acid sequence of the protein.

2.3 Protein crystallization

Purified c-Src SH3 domain protein at 11 mg/ml was crystallized in hanging drops by vapor diffusion at room temperature in 24-well VDX plates (Hampton Research) over 500 µl of reservoir solution. Crystallization drops contained 1 µl of protein with 1 µl of reservoir buffer containing 5 mM Nickel chloride, 1.7 M Ammonium Sulfate, 10 % glycerol, and 0.1 M HEPES pH 7.5. Crystals grew with morphology of rectangular prisms with typical edge dimension between 25 to 100 µm and appeared after 1 to 3 days. Crystals were harvested, washed in cryopreservative solution containing reservoir solution and 20% glycerol, and flash cooled in liquid nitrogen by manual plunging. See Table 2.

2.4 X-ray data collection and processing

X-ray diffraction data were collected at Brookhaven National Laboratory National Synchrotron Light Source II beamline 17-ID-2 (FMX) (Schneider et al., 2021). Crystal mounting, raster scanning, centering, and data collection were each performed automatically. Data were collected at wavelength 0.979315 Å. Autoprocessing was performed in the autoPROC pipeline (Vonrhein et al., 2011) which utilizes XDS for indexing and integration (Kabsch, 2010). Individual datasets from a single crystal contain 1200 images (4800 images in total) with 0.01 sec exposure and oscillation range of 0.2° per image for a total rotation range of 240°. Reflection data from 4 individual crystals were input to Aimless for scaling and merging (Evans and Murshudov, 2013). An anomalous signal was automatically detected by XDS in each of the 4 datasets; thus, Friedel pairs were treated separately. Both intensities ($I(+)$ and $I(-)$) and structure factors ($F(+)$ and $F(-)$) were output to MTZ. A resolution cutoff of 1.45 Å was determined based on a combination of values of $I/\sigma I$ > 1.5, CC½ > 0.5 and R_{pim} < 0.5, using the guidelines of (Evans and Murshudov, 2013). Crystals belong to space group $H3_2$ with unit cell dimensions a = 63.77 Å, b = 63.77 Å, c = 271.71 Å, α = β = 90°, γ = 120°. Phenix Xtriage analysis (Afonine et al., 2012) reveals that the diffraction data contain an off-origin native Patterson peak of height 31% due to translational non-crystallographic symmetry (tNCS). The free set for R_{free} calculation was randomly selected and flagged as 5% of reflections using Phenix reflection file editor. See Table 3.

2.5 Structure determination, model building and refinement

The crystal structure was determined by molecular replacement in Phaser (McCoy et al., 2007) using as the search model a previously published crystal structure of chicken c-Src SH3 domain truncated to residues 86-140 and lacking hydrogen atoms (PDB: 4JZ4 chain A (Bacarizo et al., 2014)). Translational NCS (tNCS) was detected automatically and applied during Phaser with NMOL equal to 2 and an inter molecule vector of (1, 0, 0.5). A molecular replacement solution containing 4 copies of the SH3 domain was found with top TFZ score of 10.9 and eLLG of 1070, initial R_{work} = 30.2% and R_{free} = 32.1%. Autobuilding was performed by Phenix autobuild (Terwilliger et al., 2008) to 227 residues in 4 chains and 169 waters with R_{work} = 25.2% and R_{free} = 26.53%. Manual model building was performed in Coot (Emsley and Cowtan, 2004) and refinement in Phenix (Afonine et al., 2012) using occupancy refinement and anisotropic B-factor refinement of protein and Nickel atoms. Waters were built automatically during autobuild and refinement, refined with isotropic B-factors, and were inspected and edited manually in Coot. Glycerol molecules from the cryopreservative solution were modeled manually and refined with isotropic B-factors. Nickel atoms were modeled into anomalous difference density. Secondary structure definitions are obtained by DSSP (Kabsch and Sander, 1983). See Table 4.

Phased anomalous difference maps were calculated in the Phenix utility phenix.find_peaks_holes (Afonine et al., 2012) using the protein model phases combined with the structure factors $F(+)$ and $F(-)$ from the diffraction data. Two strong anomalous peaks of heights 58.9 σ and 43.5 σ were used to guide modeling of two nickel atoms. These Ni^{2+} sites were validated in the CheckMyMetal server (Gucwa et al., 2023) and statistics given in Table 5. Refinement of the anomalous scattering factor f'' was calculated in Phenix to a value of 1.745 e- (analysis as per (Liu et al., 2013) and (El Omari et al., 2024)), which is consistent with the theoretical f'' value of 1.943 e- for Ni at wavelength 0.9793 Å (12,660 eV) obtained via X-ray absorption edge tabulations at http://www.bmsc.washington.edu/scatter/AS_index.html and (Sasaki, 1989). A third anomalous peak of 6.9 σ was present; however, a Ni atom was modeled and refined to occupancy of less than 0.2 with high B-factors and poor coordination. Therefore, this third nickel ion was omitted from the final model. Structures are illustrated with CCP4mg (McNicholas et al., 2011) and ChimeraX (Pettersen et al., 2021). All crystallography software was compiled by SBGrid (Morin et al., 2013).

3. RESULTS

3.1 Crystallization and structure determination

We set out to reproduce published crystallization conditions curated in the Protein Data Bank for the isolated c-Src SH3 domain protein for didactic purposes. Among those tested, crystallization conditions containing a cocktail of 1.4 - 1.8 M ammonium sulfate, HEPES pH 7.5, and 5 mM $NiCl_2$ in the reservoir solution yielded single crystals with rectangular prism morphology with edge length averaging between 25 - 100 μm despite heavy precipitate also being observed in the drops (Figure 1a, Table 2). We observed that when $NiCl_2$ was omitted from crystallization conditions, crystal growth was not achieved. Crystals were harvested from the drop, washed in reservoir solution, cryopreserved, and cryocooled in loops for X-ray data collection at the FMX beamline at NSLS-II. Datasets from four individual crystals were merged to achieve a multiplicity of 52.6. Consequently, anomalous signal improved and was detected automatically during data processing (Table 3).

The structure of the c-Src SH3 domain was determined to 1.45 Å resolution by molecular replacement (Table 4). Crystals belong to space group $H3_2$ with unit cell dimensions a = b = 63.77 Å, c = 271.71 Å, α = β = 90°, γ = 120° and four molecules per asymmetric unit. The $H3_2$ space group differs from the anticipated space groups $P2_12_12_1$ (PDB: 4jz4 and 4omo (Bacarizo et al., 2014) and 4rtz, unpublished)) or $P2_1$ (PDB: 4rtx, unpublished) of the previously reported c-Src SH3 crystals grown in similar $NiCl_2$-containing crystallization conditions. Examination of the Protein Data Bank reveals that no available crystal structure of the Src SH3 domain adopts the $H3_2$ space group or unit cell dimensions of our crystal form. Furthermore, interface search performed in the 'Protein interfaces, surfaces and assemblies' (PISA) server (Krissinel and Henrick, 2007) of crystal lattice contacts in the $H3_2$ crystal form reveals no common lattice contact in any available crystal structure of c-Src SH3 domain including the $P2_12_12_1$ or $P2_1$ crystal forms. Thus, this previously reported crystallization condition yielded a unique crystal lattice for c-Src SH3.

3.2 SH3 domain structure

We performed model building and refinement of the c-Src SH3 domain structure in space group $H3_2$. As our structure confirms, the c-Src SH3 domain structure is comprised of two antiparallel β-sheets formed by strands β1 - β2 - β5 and strands β3 - β4. These strands are connected by the long RT-loop (β1 - β2), n-Src loop (β2 - β3), distal loop (β3 - β4), and a 3/10 helix (β4 - β5) (Figure

1b and 1c, using nomenclature according to (Noble et al., 1993)). Chains A, B, and C are fully modeled from residues 85-143, while chain D lacks strong electron density for residues 115-119 (in the n-Src loop) which remain unmodelled. The four copies align closely with one another: upon superposition of residues Thr-88 through Ser-143, the structural alignment results in R.M.S.D. values of 0.38 - 0.99 Å over 54 Cα positions (Figure 1c, (Krissinel and Henrick, 2004)). Small positional changes in the RT and n-Src loops are the most pronounced differences between the modelled chains. Comparison of the four copies in the asymmetric unit also revealed differences in the N-terminal residues Gly-85, Val-86 and Thr-87. These variations in conformation are due to nickel binding which is discussed in detail below, and explain why the asymmetric unit is composed of four individual non-symmetric chains. Our c-Src SH3 structure also closely aligns with previously determined structures, with R.M.S.D. values ranging from 0.4 to 1.5 Å over 54 - 59 Cα positions for many of the available Src SH3 domain structures in the PDB (Figure S1, Dali server (Holm and Rosenstrom, 2010)). We therefore consider the unique lattice contacts in our $H3_2$ crystal form are not attributed to large conformational changes or structural rearrangements in the Src SH3 domain protein.

3.3 Nickel coordination in the asymmetric unit

Since a strong anomalous signal was detected in the X-ray diffraction data (Table 3), we computed phased anomalous difference maps which revealed two large anomalous peaks with heights of 58.9 σ and 43.5 σ (Figure 2a and Table 5) at locations where auto building had initially modeled water molecules; however, after refinement of these waters, the positive F_o-F_c difference map remained highly elevated (not shown), suggestive of the presence of a non-water atom (such as a metal) at these positions. As 5 mM $NiCl_2$ was required for crystal growth (Table 2) and nickel has a sufficiently high theoretical anomalous f'' atomic absorption value of 1.951 (Sasaki, 1989) electrons at the X-ray collection wavelength (0.98 Å), we modeled and refined nickel atoms at these anomalous peak positions (Figure 2b). The two nickel sites are located at the interface of Src SH3 chains A and B in a 2:2 stoichiometry. The nickel atoms bridge the A and B chains by coordination of the nitrogen and carboxyl oxygen atoms of the N-terminal residue Gly-85 of one chain and the imidazole epsilon nitrogen atom of His-125 on β3 in the neighboring chain. The crystallization pH value of 7.5 is favorable for metal coordination by histidine since its sidechain, with pKa value of 6.0, will be at least partially deprotonated. Octahedral metal coordination is completed by three water molecules (Figure 2b). The refined nickel-ligand bond distances (1.97 - 1.99 Å) are within the expected range (Table 5 and Figure S2). Both nickel sites are validated

by CheckMyMetal Binding Site Validation (Gucwa et al., 2023) and FindGeo (Andreini et al., 2012) servers (Table 5).

We further examined the 2:2 stoichiometry interaction between the Ni^{2+} ions and chains A and B of the SH3 domain. These chains are related by non-crystallographic twofold rotational symmetry and the interface buries a surface area of approximately 440 $Å^2$ per chain. In addition to the intrachain Ni^{2+} coordination, the interface includes hydrogen bonding between Gln-112 and Thr-87 and van der Waals interactions including Val-86 and Leu-127 near the nickel binding site (Figure 2c). There is also a small patch of van der Waals interactions involving Asn-115 and Thr-117 on the opposite side of the chains A/B interface (Figure S3). The nickel-mediated symmetrization aids in lattice formation by stabilizing the interactions between chains A and B in the asymmetric unit.

3.4 Role of nickel in lattice formation in $H3_2$ space group

We then more closely examined the other crystal contacts in this $H3_2$ c-Src SH3 crystal form. The four molecules in the asymmetric unit form two types of equivalent dimers: one between chains A/C and B/D which buries approximately 500 $Å^2$ per monomer, and a second between chains A/D and B/C which buries an average of 380 $Å^2$ per monomer (Figure 3a). Each of these dimer sets are related by non-crystallographic two-fold rotational symmetry. An interface search of the available structures in the Protein Data Bank for each of these dimers using the PISA server (Krissinel and Henrick, 2007) reveals no equivalent Src SH3 interface; thus, the lattice formation in the $H3_2$ space group is unique among available crystal structures of the Src SH3 domain. The binding of Ni^{2+} across the interface of chains A and B bridges these dimer pairs to help stabilize the $H3_2$ crystal lattice. Next, examination of the global crystal packing within the unit cell reveals that Ni coordination aides in asymmetric unit formation but is not involved in packing of the asymmetric units into the unit cell (Figure 3b). Taken together, these observations support the conclusion that nickel aids in formation of the asymmetric unit.

To further study this phenomenon, we attempted to detect nickel-induced dimer formation of the SH3 domain by comparing the elution profile from size exclusion chromatography and migration through semi-native SDS-PAGE (Nowakowski et al., 2014; Opdam et al., 2022) in the absence and presence of a five-fold molar excess of Ni^{2+} from $NiCl_2$; however, neither method yielded conclusive results. Therefore, whether nickel-induced dimerization occurs *in vitro* prior to crystallization or *in situ* in the crystallization dop during vapor diffusion remains unresolved.

3.5 A split amino-terminal copper and nickel (ATCUN) binding motif

The $H3_2$ crystal form of c-Src SH3 domain differs from those previously reported despite growing in similar crystallization conditions containing $NiCl_2$. To understand this discrepancy, we examined the protein constructs used for crystallization. We designed the expression plasmid to encode for a TEV protease recognition sequence (Glu-Asn-Leu-Tyr-Phe-Gln-Gly) occurring C-terminal to the His_6-tag which is absent from the parental pET-28 vector. After TEV proteolysis of the purified protein, which occurs between the Gln-Gly peptide bond of the recognition sequence, the resulting N-terminus bears the sequence Gly-Val, which is equivalent to Gly-85 and Val-86 in the native Src sequence (Table 1). This cloning scheme yields an all-native amino acid sequence following TEV proteolysis and facilitates the Ni^{2+} ion-mediated crystal form (Figure 2b and 2c).

We next compared the Ni^{2+} binding by c-Src SH3 in the $H3_2$ space group with the Ni^{2+} binding in the previously reported crystal structures of Src SH3 from similar $NiCl_2$-containing crystallization conditions and at similar pH values of 7.5 (PDB: 4jz4 and 4omo (Bacarizo et al., 2014), 4rtz (unpublished) and 4rtx (unpublished)). In these structures, Ni^{2+} ions are modelled and refined, and each chain of c-Src SH3 coordinates a single Ni^{2+} ion via the N-terminal residues Gly-Ser-His, which form a so-called 'amino terminal copper and nickel' (ATCUN) binding motif (Sankararamakrishnan et al., 2005). The ATCUN motif, consisting of sequence H_2N–x-x-His where x is any amino acid, is both a naturally occurring and engineered short peptide motif that is well established as a metal-coordinating moiety (reviewed in (Maiti et al., 2020)) that can chelate copper at very high affinity of 100 fM (Kotuniak and Bal, 2021). In these Src SH3 structures, the nickel ion is coordinated by the ATCUN motif of a single chain in distorted square planar geometry via four-nitrogen chelation from the backbone nitrogen atoms of Gly-81, Ser-82 and His-83 and an imidazole nitrogen from the sidechain of His-83 with bond distances ranging from 1.83 - 2.02 Å (Figure 4a and Figure S2). This Gly-Ser-His motif is present at the N-terminus of these c-Src SH3 domain proteins as the result of cloning artifacts and thrombin protease cleavage, and for c-Src SH3 is a non-native amino acid sequence. This differs from our interchain nickel coordination mediated by native amino acid sequences at the N-terminus Gly-85 and Val-86 and the sidechain of His-125 across the dimer interface (Figures 2 and S2). We therefore name this example a 'split' ATCUN motif.

Despite sharing a common mode of nickel binding by their N-terminal ATCUN motifs, the previous crystal structures differ in their N-terminal conformations and display flexibility in the positioning

of the Ni-bound ATCUN motifs with respect to the folded portion of the SH3 domain (Figure 4b). In addition, these Src SH3 crystals adopt different space groups ($P2_12_12_1$ and $P2_1$) and in the case of $P2_12_12_1$, two different unit cells with different packing interfaces. It has been postulated that this N-terminal flexibility contributes to different crystal packing among these earlier structures (Plaza-Garrido et al., 2020). In contrast to the $H3_2$ crystal form, the previous Ni-bound ATCUN motifs do not bridge monomers by direct nickel coordination, rather, the ATCUN motif promotes crystal packing by forming π-stacking interactions with one of two aromatic residues in neighboring symmetry-related chains, either Trp-118 (Figure 4c) or Phe-90 (Figure 4d and 4e). The fourth ATCUN-containing SH3 structure, PDB 4rtz (unpublished), is a co-crystal structure with a PxxP peptide. In this crystal form, the peptide contributes to crystal packing and the ATCUN-Ni participates in minor van der Waals interactions with a neighboring chain yet does not undergo π-stacking (not shown). Thus, flexibility in the N-terminal ATCUN motifs promote different crystal packing. The $H3_2$ crystal form represents further exploration of Src SH3 N-terminal mediated nickel coordination.

3.6 The Src SH3 $H3_2$ crystal form is not compatible with binding extended PxxP peptides

We next assessed whether the c-Src SH3 $H3_2$ crystal lattice is compatible with typical PxxP peptide binding, to ask if this crystal form might be amenable to in-crystal binding studies (Bacarizo and Camara-Artigas, 2013). SH3 domains bind typical PxxP-containing peptides via a binding pocket between the RT- and n-Src loops. This binding site can be broken up into three sub-pockets: two shallow Proline-binding (xP) pockets that accommodate the minimum PxxP binding motif and a third specificity site that binds residues in an extended peptide (Figure 5a) (Saksela and Permi, 2012). We superposed a co-crystal structure of the Src SH3 domain bound to the VSL12 peptide (PDB ID: 1rtz, unpublished, R.M.S.D. 1.1 - 1.4 Å over 56 equivalent Cα positions) onto each chain of our crystal structure and examined the positions of the modelled PxxP peptide position. Within the asymmetric unit, PxxP peptide binding to each of the four chains is compatible with the SH3 domain arrangement (Figure 5b). Next, we extended the asymmetric unit by generating symmetry mates using crystal symmetry and examined the symmetry-related lattice contacts. We observe two distinct lattice contact arrangements between chains in the asymmetric unit and symmetry-related molecules. First, chain A interacts with symmetry-related chain D (Figure 5c), and this lattice contact arrangement is shared between chain B and symmetry-related chain C (Figure S4a). When modeling a PxxP peptide bound to the SH3 domain, we observe that his lattice contact site sterically clashes with the predicted position of the PxxP peptide. Second, chain C interacts with symmetry-related chain C (Figure 5d) which is

shared between chain D and symmetry-related chain D (Figure S4b), but distinct from that observed for chains A and B (Figure 5c). In this arrangement, PxxP peptide binding to the typical binding site is also prevented by lattice contacts. We thus conclude that that standard PxxP peptide binding to the SH3 domain at the xP and specificity sites is incompatible with formation of the $H3_2$ crystal lattice since these sites in the SH3 domain are involved in crystal lattice contacts.

4. Conclusion and Discussion

In this study we describe a new crystal form for the c-Src SH3 domain protein that is dependent on interchain metal-mediated coordination and unique crystal contacts and expands the crystallization chemical space and lattice types for this important regulatory domain. This is achieved due to the native N-terminus of the isolated c-Src SH3 protein creating a previously unobserved binding site for Ni^{2+} ions. This finding offers potential benefits in biotechnology. For example, proteins in the environment of a crystal lattice are used as targets in drug discovery like crystal fragment-based screens (Davies and Tickle, 2012). The ability to obtain diverse lattice types for different protein targets affords a wider scope of available binding sites for fragment based screening (Patel et al., 2014).

Additionally, engineering of metal binding sites has emerged as an important aspect of metalloprotein studies. We introduce here the split-ATCUN motif as a derivative of the classic ATCUN motif that may be exploited by an expanded list of polypeptide targets. Previously, the engineered ATCUN motif has been extensively studied for its utility in many applications for both biology and chemistry (reviewed in (Maiti et al., 2020)). The classic monomeric ATCUN motif is comprised of the amino-terminal tripeptide consensus sequence (H_2N)-x-x-His (where x is any amino acid and histidine is at position 3), which is found in the naturally occurring ATCUN site found in proteins including human and bovine serum albumin where it is used for copper transport in the blood (Laussac and Sarkar, 1984). We provide evidence that in the absence of a histidine at position 3, nickel binding to N-terminal sequences may be achieved via a split-ATCUN motif in which the 'missing' histidine is provided by a second protein monomer, resulting in metal-mediated protein dimerization. While our observations are limited to crystallization space, it is possible that this nickel binding may also occur in solution. This observation also raises the possibility that copper could alternatively mediate split-ATCUN dimerization. Additionally, the SH3 domain fold has recently emerged as a useful structural template for an engineered metal binding, specifically as a miniaturized superoxide dismutase scaffold with catalytic activity (Hage et al., 2025). Together with our finding that the SH3 domain can engage in metal binding via a split-ATCUN motif, these results demonstrate an expanded role for the SH3 domain in metalloprotein studies.

In this example, metal binding facilitates protein crystallization. Multiple systematic approaches to improve the propensity of purified proteins to form well-ordered crystals for X-ray diffraction studies, also called "crystal lattice engineering," have been proposed. For example, protein modification tools including mutational surface entropy reduction (Goldschmidt et al., 2007) or

enzymatic lysine methylation (Walter et al., 2006) have been described as approaches to promote protein crystal lattice formation. Other approaches, sometimes termed synthetic (or mutational) symmetrization, involve engineered surface amino acid mutations to promote protein self-multimerization in a symmetric arrangement. This includes mutations to drive disulfide bond formation (Banatao et al., 2006; Forse et al., 2011), surface leucine zippers (Yamada et al., 2007), and metal coordination (Laganowsky et al., 2011; Salgado et al., 2010).

Here, we provide evidence of metal-mediated symmetrization in the absence of the additional step of creating synthetic (or mutational) sites. For the c-Src SH3 domain, this metal-mediated crystallization instead uses native amino acid residues: an N-terminal glycine residue that is the result of engineering of the protein expression construct, and a surface-exposed histidine in a second molecule. This raises the possibility of metal-mediated symmetrization in otherwise recalcitrant protein samples, as has been observed previously (for example, (Hegde et al., 2017; Trakhanov and Quiocho, 1995)). Thus, inclusion of metal ions during initial crystallization screening may help proteins self-associate into a crystal lattice. Common divalent metal salts include $NiCl_2$, $CdCl_2$, $CoCl_2$, and $CuSO_4$, which are readily accessible in many protein crystallography labs. Additionally, the absorption edge wavelengths of these common metal ions are readily achievable at standard synchrotron beamlines, which may allow for anomalous signal to be recorded and used for metal ion placement in the three-dimensional model as well as phasing for macromolecular structure solution (El Omari et al., 2024; Handing et al., 2018).

Lastly, our study demonstrates that continued studies of the SH3 domain can yield new results that expand the current knowledge of this important modular protein domain. Currently, a search of the term "SH3 domain" in the PDB yields over 6,000 entries for X-ray crystal structures, highlighting the abundance of structural information for this ubiquitous domain. SH3 domain structures were once referred to as the "darlings of the structural biology set" (Mayer and Eck, 1995) due to their prominence. Importantly, the function and activity of SH3 domains was first elucidated, in part, by structural biology studies which yielded three dimensional models of these domains. For example, three dimensional structures helped uncover the site of PxxP ligand binding, of specificity determinants, of autoinhibition within the Src family kinases, and atypical binding with non-PxxP binding partners to alternative sites on the SH3 domain (Teyra et al., 2017; Xu et al., 1997). Such sustained structural studies of SH3 domains have proven beneficial. The three dimensional structure of a Src family kinase SH3 domain was first observed in 1992-1993

(Noble et al., 1993; Yu et al., 1992), and Src SH3 continues to be a current target of structural studies (for example (Pakharukova et al., 2024)).

Acknowledgements

Anton Bennett and Sravan Perla are thanked for helpful discussions. This research used Center for BioMolecular Structure (CBMS) beamline 17ID-2 (FMX) of the National Synchrotron Light Source II, a U.S. Department of Energy (DOE) Office of Science User Facility operated for the DOE Office of Science by Brookhaven National Laboratory under Contract No. DE-SC0012704. CBMS is primarily supported by the National Institutes of Health, National Institute of General Medical Sciences (NIGMS) through a Center Core P30 Grant (P30GM133893), and by the DOE Office of Biological and Environmental Research (KP1607011). Data were collected using the beamtime obtained through NE-CAT BAG proposal #311401. We thank Alexei Soares for assistance with data collection. X.C. is a 2024 recipient of a Yale College Dean's Research Fellowship. O.S.F. is supported by NIH R35GM150644. This work is supported by NIH R01GM102262 to T.J.B.

Conflicts of interest. The authors declare no conflicts of interest.

Data availability. X-ray crystal structure model coordinates have been deposited in the RSCB Protein Data Bank (PDB) under accession code 9OFX. X-ray diffraction images are available online at SBGrid Data Bank (dataset 1158). The expression plasmid map and full plasmid sequence are provided as supplemental materials.

References

Afonine, P.V., Grosse-Kunstleve, R.W., Echols, N., Headd, J.J., Moriarty, N.W., Mustyakimov, M., Terwilliger, T.C., Urzhumtsev, A., Zwart, P.H., and Adams, P.D. (2012). Towards automated crystallographic structure refinement with phenix.refine. Acta Crystallogr D *68*, 352-367.

Anafi, M., Rosen, M.K., Gish, G.D., Kay, L.E., and Pawson, T. (1996). A potential SH3 domain-binding site in the Crk SH2 domain. J Biol Chem *271*, 21365-21374.

Andreini, C., Cavallaro, G., and Lorenzini, S. (2012). FindGeo: a tool for determining metal coordination geometry. Bioinformatics *28*, 1658-1660.

Bacarizo, J., and Camara-Artigas, A. (2013). Atomic resolution structures of the c-Src SH3 domain in complex with two high-affinity peptides from classes I and II. Acta Crystallogr D Biol Crystallogr *69*, 756-766.

Bacarizo, J., Martinez-Rodriguez, S., Martin-Garcia, J.M., Andujar-Sanchez, M., Ortiz-Salmeron, E., Neira, J.L., and Camara-Artigas, A. (2014). Electrostatic effects in the folding of the SH3 domain of the c-Src tyrosine kinase: pH-dependence in 3D-domain swapping and amyloid formation. PLoS One *9*, e113224.

Banatao, D.R., Cascio, D., Crowley, C.S., Fleissner, M.R., Tienson, H.L., and Yeates, T.O. (2006). An approach to crystallizing proteins by synthetic symmetrization. Proc Natl Acad Sci U S A *103*, 16230-16235.

Boggon, T.J., and Eck, M.J. (2004). Structure and regulation of Src family kinases. Oncogene *23*, 7918-7927.

Chau, J.E., Vish, K.J., Boggon, T.J., and Stiegler, A.L. (2022). SH3 domain regulation of RhoGAP activity: Crosstalk between p120RasGAP and DLC1 RhoGAP. Nature communications *13*, 4788.

Chen, V.B., Arendall, W.B., Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J., Murray, L.W., Richardson, J.S., and Richardson, D.C. (2010). MolProbity: all-atom structure validation for macromolecular crystallography. Acta Crystallogr D *66*, 12-21.

Cicchetti, P., Mayer, B.J., Thiel, G., and Baltimore, D. (1992). Identification of a protein that binds to the SH3 region of Abl and is similar to Bcr and GAP-rho. Science *257*, 803-806.

Cloutier, J.F., and Veillette, A. (1996). Association of inhibitory tyrosine protein kinase p50csk with protein tyrosine phosphatase PEP in T cells and other hemopoietic cells. EMBO J *15*, 4909-4918.

Davies, T.G., and Tickle, I.J. (2012). Fragment screening using X-ray crystallography. Top Curr Chem *317*, 33-59.

El Omari, K., Forsyth, I., Duman, R., Orr, C.M., Mykhaylyk, V., Mancini, E.J., and Wagner, A. (2024). Utilizing anomalous signals for element identification in macromolecular crystallography. Acta Crystallogr D Struct Biol.

Emsley, P., and Cowtan, K. (2004). Coot: model-building tools for molecular graphics. Acta crystallographica Section D, Biological crystallography *60*, 2126-2132.

Evans, P.R., and Murshudov, G.N. (2013). How good are my data and what is the resolution? Acta Crystallogr D Biol Crystallogr *69*, 1204-1214.

Feng, S., Chen, J.K., Yu, H., Simon, J.A., and Schreiber, S.L. (1994). Two binding orientations for peptides to the Src SH3 domain: development of a general model for SH3-ligand interactions. Science *266*, 1241-1247.

Forse, G.J., Ram, N., Banatao, D.R., Cascio, D., Sawaya, M.R., Klock, H.E., Lesley, S.A., and Yeates, T.O. (2011). Synthetic symmetrization in the crystallization and structure determination of CelA from. Protein Science *20*, 168-178.

Goldschmidt, L., Cooper, D.R., Derewenda, Z.S., and Eisenberg, D. (2007). Toward rational protein crystallization: A Web server for the design of crystallizable protein variants. Protein Sci *16*, 1569-1576.

Gucwa, M., Lenkiewicz, J., Zheng, H., Cymborowski, M., Cooper, D.R., Murzyn, K., and Minor, W. (2023). CMM-An enhanced platform for interactive validation of metal binding sites. Protein Sci *32*, e4525.

Hage, F.R., Schwan, M., Conde Gonzalez, M.R., Huber, J., Germer, S., Macri, M., Kopp, J., Sinning, I., and Thomas, F. (2025). Strand-Swapped SH3 Domain Dimer with Superoxide Dismutase Activity. ACS Cent Sci *11*, 157-166.

Handing, K.B., Niedzialkowska, E., Shabalin, I.G., Kuhn, M.L., Zheng, H., and Minor, W. (2018). Characterizing metal-binding sites in proteins with X-ray crystallography. Nat Protoc *13*, 1062-1090.

Hegde, R.P., Pavithra, G.C., Dey, D., Almo, S.C., Ramakumar, S., and Ramagopal, U.A. (2017). Can the propensity of protein crystallization be increased by using systematic screening with metals? Protein Science *26*, 1704-1713.

Holm, L., and Rosenstrom, P. (2010). Dali server: conservation mapping in 3D. Nucleic acids research *38*, W545-549.

Kabsch, W. (2010). Xds. Acta Crystallogr D *66*, 125-132.

Kabsch, W., and Sander, C. (1983). Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features. Biopolymers *22*, 2577-2637.

Kall, S.L., Whitlatch, K., Smithgall, T.E., and Lavie, A. (2019). Molecular basis for the interaction between human choline kinase alpha and the SH3 domain of the c-Src tyrosine kinase. Scientific reports *9*, 17121.

Kaneko, T., Li, L., and Li, S.S. (2008). The SH3 domain--a family of versatile peptide- and protein-recognition module. Front Biosci *13*, 4938-4952.

Kazemein Jasemi, N.S., Mehrabipour, M., Magdalena Estirado, E., Brunsveld, L., Dvorsky, R., and Ahmadian, M.R. (2024). Functional Classification and Interaction Selectivity Landscape of the Human SH3 Domain Superfamily. Cells *13*.

Kotuniak, R., and Bal, W. (2021). Kinetics of Cu(II) complexation by ATCUN/NTS and related peptides: a gold mine of novel ideas for copper biology. Dalton Trans *51*, 14-26.

Krissinel, E., and Henrick, K. (2004). Secondary-structure matching (SSM), a new tool for fast protein structure alignment in three dimensions. Acta Crystallogr D Biol Crystallogr *60*, 2256-2268.

Krissinel, E., and Henrick, K. (2007). Inference of macromolecular assemblies from crystalline state. Journal of molecular biology *372*, 774-797.

Laganowsky, A., Zhao, M., Soriaga, A.B., Sawaya, M.R., Cascio, D., and Yeates, T.O. (2011). An approach to crystallizing proteins by metal-mediated synthetic symmetrization. Protein Sci *20*, 1876-1890.

Laussac, J.P., and Sarkar, B. (1984). Characterization of the copper(II)- and nickel(II)-transport site of human serum albumin. Studies of copper(II) and nickel(II) binding to peptide 1-24 of human serum albumin by 13C and 1H NMR spectroscopy. Biochemistry *23*, 2832-2838.

Lim, W.A., Richards, F.M., and Fox, R.O. (1994). Structural determinants of peptide-binding orientation and of sequence specificity in SH3 domains. Nature *372*, 375-379.

Liu, Q., Liu, Q., and Hendrickson, W.A. (2013). Robust structural analysis of native biological macromolecules from multi-crystal anomalous diffraction data. Acta Crystallogr D Biol Crystallogr *69*, 1314-1332.

Luttrell, L.M., Ferguson, S.S., Daaka, Y., Miller, W.E., Maudsley, S., Della Rocca, G.J., Lin, F., Kawakatsu, H., Owada, K., Luttrell, D.K.*, et al.* (1999). Beta-arrestin-dependent formation of beta2 adrenergic receptor-Src protein kinase complexes. Science *283*, 655-661.

Maiti, B.K., Govil, N., Kundu, T., and Moura, J.J.G. (2020). Designed Metal-ATCUN Derivatives: Redox- and Non-redox-Based Applications Relevant for Chemistry, Biology, and Medicine. iScience *23*, 101792.

Mayer, B.J., and Eck, M.J. (1995). SH3 domains. Minding your p's and q's. Curr Biol *5*, 364-367.

Mayer, B.J., Hamaguchi, M., and Hanafusa, H. (1988). A novel viral oncogene with structural similarity to phospholipase C. Nature *332*, 272-275.

McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni, L.C., and Read, R.J. (2007). Phaser crystallographic software. J Appl Crystallogr *40*, 658-674.

McNicholas, S., Potterton, E., Wilson, K.S., and Noble, M.E. (2011). Presenting your structures: the CCP4mg molecular-graphics software. Acta Crystallogr D Biol Crystallogr *67*, 386-394.

Mehrabipour, M., Jasemi, N.S.K., Dvorsky, R., and Ahmadian, M.R. (2023). A Systematic Compilation of Human SH3 Domains: A Versatile Superfamily in Cellular Signaling. Cells *12*.

Morin, A., Eisenbraun, B., Key, J., Sanschagrin, P.C., Timony, M.A., Ottaviano, M., and Sliz, P. (2013). Collaboration gets the most out of software. Elife *2*, e01456.

Musacchio, A., Noble, M., Pauptit, R., Wierenga, R., and Saraste, M. (1992). Crystal structure of a Src-homology 3 (SH3) domain. Nature *359*, 851-855.

Musacchio, A., Saraste, M., and Wilmanns, M. (1994). High-resolution crystal structures of tyrosine kinase SH3 domains complexed with proline-rich peptides. Nat Struct Biol *1*, 546-551.

Noble, M.E., Musacchio, A., Saraste, M., Courtneidge, S.A., and Wierenga, R.K. (1993). Crystal structure of the SH3 domain in human Fyn; comparison of the three-dimensional structures of SH3 domains in tyrosine kinases and spectrin. EMBO J *12*, 2617-2624.

Nowakowski, A.B., Wobig, W.J., and Petering, D.H. (2014). Native SDS-PAGE: high resolution electrophoretic separation of proteins with retention of native properties including bound metal ions. Metallomics *6*, 1068-1078.

Opdam, L.V., Polanco, E.A., de Regt, B., Lambertina, N., Bakker, C., Bonnet, S., and Pandit, A. (2022). A screening method for binding synthetic metallo-complexes to haem proteins. Anal Biochem *653*, 114788.

Pakharukova, N., Thomas, B.N., Bansia, H., Li, L., Abzalimov, R.R., Kim, J., Kahsai, A.W., Pani, B., Bassford, D.K., Liu, S.*, et al.* (2024). Beta-arrestin 1 mediated Src activation via Src SH3 domain revealed by cryo-electron microscopy. bioRxiv.

Patel, D., Bauman, J.D., and Arnold, E. (2014). Advantages of crystallographic fragment screening: functional and mechanistic insights from a powerful platform for efficient drug discovery. Progress in biophysics and molecular biology *116*, 92-100.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., Morris, J.H., and Ferrin, T.E. (2021). UCSF ChimeraX: Structure visualization for researchers, educators, and developers. Protein Sci *30*, 70-82.

Plaza-Garrido, M., Salinas-Garcia, M.C., Martinez, J.C., and Camara-Artigas, A. (2020). The effect of an engineered ATCUN motif on the structure and biophysical properties of the SH3 domain of c-Src tyrosine kinase. J Biol Inorg Chem *25*, 621-634.

Saksela, K., Cheng, G., and Baltimore, D. (1995). Proline-rich (PxxP) motifs in HIV-1 Nef bind to SH3 domains of a subset of Src kinases and are required for the enhanced growth of Nef+ viruses but not for down-regulation of CD4. EMBO J *14*, 484-491.

Saksela, K., and Permi, P. (2012). SH3 domain ligand binding: What's the consensus and where's the specificity? FEBS Lett *586*, 2609-2614.

Salgado, E.N., Radford, R.J., and Tezcan, F.A. (2010). Metal-directed protein self-assembly. Acc Chem Res *43*, 661-672.

Sankararamakrishnan, R., Verma, S., and Kumar, S. (2005). ATCUN-like metal-binding motifs in proteins: identification and characterization by crystal structure and sequence analysis. Proteins *58*, 211-221.

Sasaki, S.a.N.L.f.H.E.P., Tsukuba, Ibaraki (Japan) (1989). Numerical Tables of Anomalous Scattering Factors Calculated by the Cromer and Liberman's Method.

Schneider, D.K., Shi, W., Andi, B., Jakoncic, J., Gao, Y., Bhogadi, D.K., Myers, S.F., Martins, B., Skinner, J.M., Aishima, J.*, et al.* (2021). FMX - the Frontier Microfocusing Macromolecular Crystallography Beamline at the National Synchrotron Light Source II. J Synchrotron Radiat *28*, 650-665.

Terwilliger, T.C., Grosse-Kunstleve, R.W., Afonine, P.V., Moriarty, N.W., Zwart, P.H., Hung, L.W., Read, R.J., and Adams, P.D. (2008). Iterative model building, structure refinement and density modification with the PHENIX AutoBuild wizard. Acta Crystallogr D *64*, 61-69.

Teyra, J., Huang, H., Jain, S., Guan, X., Dong, A., Liu, Y., Tempel, W., Min, J., Tong, Y., Kim, P.M.*, et al.* (2017). Comprehensive Analysis of the Human SH3 Domain Family Reveals a Wide Variety of Non-canonical Specificities. Structure *25*, 1598-1610 e1593.

Trakhanov, S., and Quiocho, F.A. (1995). Influence of divalent cations in protein crystallization. Protein Sci *4*, 1914-1919.

Vogel, C., Bashton, M., Kerrison, N.D., Chothia, C., and Teichmann, S.A. (2004). Structure, function and evolution of multidomain proteins. Current opinion in structural biology *14*, 208-216.

Vonrhein, C., Flensburg, C., Keller, P., Sharff, A., Smart, O., Paciorek, W., Womack, T., and Bricogne, G. (2011). Data processing and analysis with the autoPROC toolbox. Acta Crystallogr D Biol Crystallogr *67*, 293-302.

Walter, T.S., Meier, C., Assenberg, R., Au, K.F., Ren, J., Verma, A., Nettleship, J.E., Owens, R.J., Stuart, D.I., and Grimes, J.M. (2006). Lysine methylation as a routine rescue strategy for protein crystallization. Structure *14*, 1617-1622.

Xiao, R., Xi, X.D., Chen, Z., Chen, S.J., and Meng, G. (2013). Structural framework of c-Src activation by integrin beta3. Blood *121*, 700-706.

Xu, W., Harrison, S.C., and Eck, M.J. (1997). Three-dimensional structure of the tyrosine kinase c-Src. Nature *385*, 595-602.

Yamada, H., Tamada, T., Kosaka, M., Miyata, K., Fujiki, S., Tano, M., Moriya, M., Yamanishi, M., Honjo, E., Tada, H.*, et al.* (2007). 'Crystal lattice engineering,' an approach to engineer protein crystal contacts by creating intermolecular symmetry: crystallization and structure determination of a mutant human RNase 1 with a hydrophobic interface of leucines. Protein Sci *16*, 1389-1397.

Yu, H., Chen, J.K., Feng, S., Dalgarno, D.C., Brauer, A.W., and Schreiber, S.L. (1994). Structural basis for the binding of proline-rich peptides to SH3 domains. Cell *76*, 933-945.

Yu, H., Rosen, M.K., Shin, T.B., Seidel-Dugan, C., Brugge, J.S., and Schreiber, S.L. (1992). Solution structure of the SH3 domain of Src and identification of its ligand-binding site. Science *258*, 1665-1668.

Table 1. Macromolecule production information

Source organism	*Homo sapiens*
DNA source	cDNA
Expression vector	pET-28a(+)
Plasmid construction method	PCR and restriction-ligation
Forward primer (NheI and TEV site)	5' - GGATTCCATATGGAGAACCTGTACTTTCAA GGCGTGACCACCTTTGTGGCCC - 3'
Reverse primer (STOP codon and XhoI)	5' - CCGCTCGAGCTAGGAGGGCGCCACATAGTTG - 3'
Expression host	*E. coli* Rosetta(DE3)
Expression details	Induced with 0.2 mM IPTG at OD_{600} = 0.8, 18°C, 20 hr
Complete amino acid sequence of the protein produced	MGSSHHHHHHSSGLVPRGSHMENLYFQGVTTFVALYDY ESRTETDLSFKKGERLQIVNNTEGDWWLAHSLSTGQT GYIPSNYVAPS
Complete amino acid sequence of the protein after TEV proteolysis	85 - GVTTF VALYDYESRT ETDLSFKKGE RLQIVNNTEG DWWLAHSLST GQTGYIPSNY VAPS - 143

Table 2. Crystallization information

Method	Hanging drop vapor diffusion
Plate type	VDX™ Plate with sealant (Hampton Research)
Temperature (°C)	25º C (room temperature)
Protein concentration	1.8 mM, 11.9 mg/mL
Buffer composition of protein solution	150 mM NaCl, 20 mM Tris pH 7.5
Composition of reservoir solution	1.7 M Ammonium Sulfate, 5 mM $NiCl_2$, 10% glycerol, 0.1 M HEPES pH 7.5
Volume and ratio of drop	2 µL drop, 1:1 v/v protein to reservoir ratio
Volume of reservoir	500 µl in VDX plate (Hampton Research)
Composition of the cryopreservative	1.7 M Ammonium Sulfate, 5 mM $NiCl_2$, 20% glycerol, 0.1 M HEPES pH 7.5
Drop setting	Manual
Seeding	No

Table 3. Data collection and processing

Diffraction source	BNL NSLS-II 17-ID-2 FMX			
Wavelength (Å)	0.979315			
Detector	EIGER 16M			
Crystal	1	2	3	4
Temperature (K)	100	100	100	100
Crystal to detector distance (mm)	200	200	200	200
Total rotation range (°)	240	240	240	240
Rotation per image (°)	0.2	0.2	0.2	0.2
Exposure time per image (s)	0.01	0.01	0.01	0.01
Space group	$H3_2$	$H3_2$	$H3_2$	$H3_2$
Unit cell dimensions a (Å) b (Å) c (Å) α, β, γ (°)	63.760 63.760 271.669 90, 90, 120	63.976 63.976 272.214 90, 90, 120	63.976 63.976 272.816 90, 90, 120	63.867 63.867 272.628 90, 90, 120
Mosaicity (°)	0.09	0.06	0.07	0.06
Anomalous resolution limit (Å)[+]	2.63	2.40	2.16	2.21
Number of crystals merged	4			
Resolution range (Å) of merged data	31.95 - 1.45 (1.48 - 1.45)			
Mosaicity (°) of merged data	0.07			
Unit cell dimensions a, b, c (Å) α, β, γ (°)	63.77, 63.77, 271.71 90, 90, 120			
Anomalous resolution limit of merged data (Å)[+]	2.00			
Anomalous signal to noise Mn($dI/\sigma dI$))	1.3 (0.89)			
Number of reflections				
Total observations	2,030,203			
Unique (anomalous)	73,036			
Unique (non-anomalous)	38,628			
Completeness (%)	100 (100)			
Anomalous completeness (%)	100 (100)			
Redundancy	52.6 (46.8)			
Anomalous redundancy	27.2 (23.8)			
Mean $I/\sigma(I)$	22.9 (1.8)			
$CC_{1/2}$	1.00 (0.784)			
R_{pim}	0.012 (0.509)			
Wilson B-factor (Å^2)	23.07			

Values given in parentheses are for the highest resolution shell
[+] Anomalous signal detected in Aimless (Evans and Murshudov, 2013) where CC_{anom} < 0.15

Table 4. Structure refinement

Resolution range (Å)	31.89 - 1.45 (1.47 - 1.45)
Completeness (%)	99.9 (100)
No. of reflections, working set	73,036 (2,734)
No. of reflections, test set	3,450 (137)
Reflections, test set size (%)	4.72 (5.0)
Final R_{work}	16.25 (27.88)
Final R_{free}	19.61 (37.45)
No. of non-H atoms	
Protein	1,888
Metals (Ni^{2+})	2
Waters, other solvent	163, 54
Total	2,107
R.M.S. deviations from ideality[*]	
Bonds (Å)	0.005
Angles (°)	0.879
Average B-factors ($Å^2$)	
Protein	30.70
Metals (Ni^{2+})	24.62
Waters, other solvent	38.87, 58.29
Overall	32.07
MolProbity Statistics	
Ramachandran plot	
Favored regions (%)	98.19
Allowed regions (%)	1.81
Outliers (%)	0.00
Clashscore	2.63
Unmodelled residues (%)	2.12% (5 out of 236 total residues)
PDB Code	9OFX

Values given in parentheses are for the highest resolution shell.
* Phenix Geometry restraints library

Table 5. Nickel site information

Anomalous scatterer	Nickel	
Photon energy (eV)	12,660	
Theoretical f'' (e-)	1.943	
Refined f'' (e-)	1.745	
Nickel sites	Ni-1	Ni-2
Peak height (σ) [*]	58.9	43.5
CheckMyMetal[+] statistics		
B-factor (Å^2)	22.0	27.2
Valence	2.7	2.8
Geometry	Octahedral	Octahedral
R.M.S.D. Geometry[+,#]		
Angles (°)	4.3	6.0
Bonds (Å)	0.178	0.250
Metal-ligand distances (Å)		
Gly-85 N	1.97	1.97
Gly-85 O	1.97	1.98
His-125 N (Nε2)	1.99	1.99
Water	1.97 (1)	1.97 (4)
Water	1.98 (2)	1.97 (5)
Water	1.99 (3)	1.97 (6)

[*] Anomalous peak heights (σ) calculated in Phenix (phenix.find_peaks_holes)
[+] CheckMyMetal Server (Gucwa et al., 2023)
[#] FindGeo Server (Andreini et al., 2012)

Figure 1. Crystal structure of c-Src SH3 domain in $H3_2$ crystal form.
(*a*) Representative crystals of c-Src SH3 domain grown for this study. A scale bar representing 25 μM is included in each panel. The average crystal has edge dimension of 25 - 100 μm and fit within a 0.05-0.1 mm CryoLoop (Hampton Research).
(*b*) Ribbon diagram of the c-Src SH3 domain crystal structure from this study. Shown is chain A.
(*c*) Superposition of the structures of each chain of c-Src SH3 domain from this study. Shown is the Cα trace of each chain colored dark teal (chain A), orange (chain B), light teal (chain C) and tan (chain D). The positions of Thr-88 and Ser-143 are indicated with arrows. In (*b*) and (*c*), the secondary structure elements, loops, and amino- and carboxy-termini are named and labelled.

Figure 2. Nickel coordination by the c-Src SH3 domain.
(*a*) The asymmetric unit containing four chains of c-Src SH3 domain (Cα traces) and phased anomalous difference map contoured to 10 σ (pink) shows two strong peaks (peak 1 and peak 2) arranged between chains A (dark teal) and B (orange). The two views are related by a 90° rotation about the x-axis away from the reader.
(*b*) Final refined $2F_o$-F_c map (blue, contoured to 1.5 σ) and phased anomalous difference map (pink, contoured to 10 σ). Ni^{2+} atoms are drawn as grey spheres, and water molecules are red spheres.
(*c*) Details of nickel binding. Left panel: overview of chain A/chain B interface. Zoom-in: details of the interface including intrachain nickel coordination (black dashed lines) by Gly-85 and His-125.

Figure 3. Lattice contacts in the c-Src SH3 $H3_2$ crystal form.
(a) Left panel: asymmetric unit arrangement of four copies of the c-Src SH3 domain. Two types of dimers are depicted in the center and right subpanels.
(b) Expanded view of the unit cell, which is drawn as a box. The nickel sites are indicated as pink spheres. One copy of the asymmetric unit is drawn with protein ribbons and colored as in part (a). The other protein monomers are drawn as gray C-alpha traces. The orientation axes x (red), y (yellow) and z (blue) are indicated. The nickel sites bridge monomers A and B in the asymmetric unit but do not further aid in crystal packing between asymmetric units.

Figure 4. Nickel coordination by c-Src SH3 domains containing an ACTUN motif.
(a) The ATCUN motif Gly-Ser-His at the N-terminus of Src SH3 domain protein (PDB: 4jz4, (Bacarizo et al., 2014)) coordinates nickel in square planar geometry.
(b) Superposition of c-Src SH3 domains containing ATCUN and coordinating nickel. Included are PDBs 4jz4 (light blue, chains A and B, (Bacarizo et al., 2014)), 4rtz (light pink, $P2_12_12_1$, unpublished) and 4rtx (cyan, $P2_1$, unpublished). PDB 4omo (Bacarizo et al., 2014) is equivalent to 4jz4 and is not shown. The nickel sites are drawn as spheres and colored according to the corresponding PDB.
(c) Symmetry-related π-stacking interaction involving the nickel-binding ATCUN motif of chain A (light blue) and the sidechain of Tyr-90 in symmetry-related chain B (chain B_{symm}, gold). PDB: 4jz4 (Bacarizo et al., 2014).
(d) Symmetry-related π-stacking interaction involving the nickel-binding ATCUN motif of chain B (light blue) and the sidechain of Trp-118 in a symmetry-related chain A (chain A_{symm}, gold). PDB: 4jz4 (Bacarizo et al., 2014).
(e) Symmetry-related π-stacking interaction involving the nickel-binding ATCUN motif of one chain (cyan) and the sidechain of Tyr-90 in a second chain (gold). PDB: 4rtx (unpublished). For comparison, the orientation of the main SH3 domain molecule is equivalent in parts c, d, and e.

Figure 5. Modeling PxxP binding onto c-Src SH3 domains in $H3_2$ space group.

(a) Co-crystal structure of Src SH3 domain (teal, drawn in ribbon and semi-transparent surface) bound to the VSL12 peptide (in purple) (PDB ID: 4rtz, unpublished). Peptide binding sites on SH3 are highlighted in yellow and labeled.

(b) Asymmetric unit of c-Src SH3 $H3_2$ crystal structure. The PxxP peptide (purple surface) is modeled onto each chain by superposition of the complex from (a) with each individual chain of the asymmetric unit, with R.M.S.D. values ranging from 0.7 - 1.0 over 57 equivalent Cα positions.

(c) and (d) Modeling reveals that the predicted PxxP binding sites are blocked by symmetry-related molecules in the $H3_2$ crystal form. Residues in the symmetry mates (Chain D$_{symm}$ (c) and Chain C$_{symm}$ (d)) with predicted steric overlaps with the modeled PxxP peptide are colored red on the Cα trace (calculated in MolProbity (Chen et al., 2010)).

(c) c-Src SH3 chain A (teal ribbon) and its symmetry-related chain D (D$_{symm}$, tan Cα trace). Chain B shares a similar interface with a symmetry-related chain C (Figure S4a).

(d) c-Src SH3 chain C (light cyan ribbon) and symmetry-related chain C (C$_{symm}$, grey-blue). Chain D shares a similar interface with a symmetry-related chain D (Figure S4b).