Title

P-body factors Ddx6 and Ddx61 support development in mRNA-decay deficient *pnrc2* mutants

Running title

Pnrc2, P-bodies, and development

Authors

Gallagher, Thomas L.[1,2,‡]; Blatnik, Monica C.[1,2,3,*]; Austin, Clare C.[1,2]; Thompson, Kathryn G.[1,2];

Pvirre, Danielle M.[1]; Denniston, Ryan[2]; Morgan, Angelina[1]; Kearse, Michael G.[2,4]; Amacher,

Sharon L.[1,2,3,5‡]

1 Department of Molecular Genetics, The Ohio State University, Columbus, OH, 43210, USA

2 Center for RNA Biology, The Ohio State University, Columbus, OH, 43210, USA

3 Interdisciplinary Graduate Program in Molecular, Cellular and Developmental Biology, The

Ohio State University, Columbus, OH, 43210, USA

4 Department of Biological Chemistry and Pharmacology, The Ohio State University, Columbus,

OH, 43210, USA

5 Center for Muscle Health and Neuromuscular Disorders, The Ohio State University,

Columbus, OH, 43210, USA.

*Present address: Oberlin College of Arts and Sciences, 38 E. College St., Oberlin, OH, 44074,

USA.

‡Authors for correspondence (gallagher.184@osu.edu and amacher.6@osu.edu)

Key words

somitogenesis, oscillatory gene expression, P-bodies, translational regulation, deadenylation

Abstract

Somitogenesis is controlled by the segmentation clock, a molecular oscillator that

controls periodic gene expression in unsegmented mesoderm and is regulated by a negative

feedback loop driven by Hes/Her transcriptional repressors. In zebrafish, Pnrc2 is required for

decay of *her1* transcript and additional oscillatory gene transcripts. Despite accumulation of

numerous mRNAs including those encoding developmental regulators, overt embryonic

phenotypes are absent in *pnrc2* mutants. Our previous work suggested that accumulated

mRNAs are not translated in *pnrc2* mutants, though the underlying mechanism(s) was unknown.

We show here that many overexpressed transcripts in *pnrc2* mutants have shortened poly(A)

tails and are disengaged from ribosomes, and that deadenylation inhibition leads to somite

defects in *pnrc2* mutants. Transcripts encoding the P-body protein, Ddx61, are both

overexpressed and engaged with ribosomes in *pnrc2* mutants, leading to an increase in Ddx61

protein that may compensate against the negative effects of transcript accumulation. Co-

depletion of Ddx61 and Ddx6 in *pnrc2* mutants increases *her1* accumulation and leads to

pleiotropic defects. Together, our results show that multiple post-transcriptional mechanisms

ensure proper translation when mRNA decay is inhibited.

Summary statement

Pnrc2 promotes decay of tail-shortened, ribosome-disengaged mRNAs and co-regulates *her1*

mRNA decay with Ddx6 and Ddx61, P-body factors that sustain normal development in decay-

deficient *pnrc2* mutants.

Introduction

Vertebrate somitogenesis is a fundamental developmental process (Kageyama et al.,

2012) characterized by the anterior to posterior sequential segmentation of the embryonic

mesoderm into blocks of tissue called somites. This rhythmic process is controlled by a

molecular oscillator called the segmentation clock which operates in the presomitic mesoderm

(PSM) and is characterized by rapid cycles of mRNA and protein expression (Hubaud and

Pourquie, 2014; Oates et al., 2012; Pourquie, 2011). The segmentation clock is regulated by a

self-sustaining negative feedback loop, where a core oscillatory factor, a transcriptional

repressor, inhibits expression of downstream oscillatory genes, including itself.

For proper somite formation, rates of each oscillatory step, from transcription to

translation to decay, must be regulated to ensure correct somite size and number (Gomez et al.,

2008; Holley et al., 2000; Keynes and Stern 1988; Lewis, 2003; Matsuda et al., 2020; Palmeirim

et al., 1997; Schroter and Oates, 2010). Real-time segmentation clock reporters that

recapitulate clock dynamics in vivo must not only contain critical transcriptional regulatory

regions that drive oscillatory expression, but also contain features including 3′UTR sequences

and protein motifs that destabilize reporter mRNAs and proteins, respectively (Aulehla et al.,

2008; Delaune et al., 2012; Masamizu et al., 2006; Simsek et al., 2023; Yoshioka-Kobayashi et

al., 2020).

We have focused on post-transcriptional regulation of the segmentation clock (Blatnik et

al., 2023), motivated by our discovery that the zebrafish *pnrc2* gene promotes rapid decay of

oscillatory gene transcripts (Dill and Amacher, 2005; Gallagher et al., 2017; Tietz et al., 2020).

Transcriptional oscillations are normal in zebrafish *pnrc2* mutants, but *her1* mRNA (and other

oscillatory gene transcripts) accumulate due to mRNA decay defects (Gallagher et al., 2017).

Surprisingly, despite a 4-6-fold increase in *her1* and *dlc* oscillatory gene transcripts in *pnrc2*

mutants, oscillatory gene protein is expressed normally (Gallagher et al., 2017) and there is no

overt segmentation phenotype. Discordance between mRNA and protein levels might be caused

by a combination of factors: transcripts might be inefficiently translated due to deadenylation

and/or binding of translational repressors and/or localized to granules like P-bodies that are

sites of non-translating mRNAs (Luo et al., 2018; Ostareck et al., 2014).

To better understand the role of *pnrc2* during segmentation, we used RNA-Seq analysis

and identified over 1,700 transcripts that were significantly overexpressed at least 1.2-fold in

MZpnrc2 mutants. Because *pnrc2* is expressed broadly in embryos (Gallagher et al., 2017), we

anticipated that Pnrc2 likely functions in many tissues. GO analysis reveals enrichment for many

processes associated with development and mRNA regulation. Motif analysis shows that

overexpressed transcripts are enriched for well-known mRNA-destabilizing motifs. Several

oscillatory transcripts have shortened poly(A) tails in *MZpnrc2* mutants, suggesting that they are

inefficiently translated. Supporting this interpretation, polysome profiling of *MZpnrc2* mutants

shows that accumulated transcripts of several oscillatory genes are primarily disengaged from

ribosomes. *MZpnrc2* mutants are sensitive to deadenylation inhibition during segmentation, and

later to depletion of the P-body factors Ddx6 and Ddx61, suggesting that accumulated mRNAs

in *MZpnrc2* mutants are inefficiently translated when deadenylation and P-body functions are

intact, though whether these processes are connected remains unknown. Our results support

that discordant expression of oscillatory gene mRNA and protein in *MZpnrc2* mutants is due to

inefficient translation of accumulated transcripts. In the absence of *pnrc2* function, the P-body

factors Ddx6 and Ddx61 buffer against mRNA decay dysfunction and support embryonic

development, implicating a broader role for Pnrc2 and P-bodies during development across

many tissues.

Results:

Pnrc2-regulated transcripts are associated with key biological processes and are enriched for

mRNA destabilizing motifs

To globally identify transcripts that require *pnrc2* for proper expression, we performed

RNA-Seq analysis on wild-type (WT) and maternal-zygotic *pnrc2* (*MZpnrc2*) mutant embryos at

mid-segmentation stages. We detected 2,256 differentially expressed transcripts with a false

discovery rate of q < 0.05 (Fig. 1A, Tables S1-S3) using Kallisto to quantify transcript abundance

(Bray et al., 2016) and Sleuth for differential expression analysis (Pimentel et al., 2017). Most

differentially expressed transcripts were overexpressed (n = 1,769 transcripts), as expected for

an mRNA decay-deficient mutant. Among overexpressed transcripts, >99% are annotated as

protein-coding by Ensembl's biotype annotation (n = 1,759 transcripts), suggesting that Pnrc2

regulates decay of transcripts with full open reading frames rather than transcripts containing

premature termination codons that are subject to nonsense-mediated decay (NMD).

NMD can also be induced by specific features in transcripts with full open reading

frames. One such feature is the presence of an upstream open reading frame (uORF) (Mendell

et al., 2004; Hurt et al., 2013; Johnstone et al., 2016). Using stringent filtering criteria (see

methods), we identified uORF-containing transcripts using a list validated by ribosome profiling

(Johnstone et al., 2016) and find that overexpressed transcripts in *MZpnrc2* mutants are

enriched for uORFs relative to non-overexpressed transcripts (21.2% vs 15.7%, respectively, p

= 2.832×10^{-07} by a Fisher's Exact Test) (Table S3). These results indicate that Pnrc2 promotes

decay of some, but not all uORF-containing transcripts. NMD can also be induced by intron-

containing 3′UTRs (Schweingruber et al., 2013; Lykke-Andersen & Jensen, 2015), however, few

overexpressed transcripts in *MZpnrc2* mutants contain 3'UIs (n = 50 transcripts of 1769

overexpressed transcripts) when compared to a published list of 3'UI-containing transcripts

(Gangras et al., 2020). Finally, 3'UTR length can also influence mRNA stability (Schweingruber

et al., 2013; Lykke-Andersen & Jensen, 2015), so we compared the length distributions of

overexpressed, unaffected, and under-expressed transcript 3′UTRs (see methods). Although

overexpressed transcript 3′UTRs have a significantly longer length distribution than unaffected

transcript 3′UTRs, under-expressed transcripts have an even longer length distribution than

overexpressed and unaffected transcript 3′UTRs (Fig. 1B), suggesting that 3′UTR length is not a

primary mechanism by which Pnrc2 promotes transcript decay. To assess whether 3′UTR

lengths are different between WT and *MZpnrc2* mutant embryos, sequencing reads were

aligned using STAR (Dobin et al., 2013) and metagene plots were generated using deepTools

(Ramirez et al., 2016). Metagene plots of gene body read coverage for WT and *MZpnrc2*

mutants are very similar, including a steep drop in coverage downstream of transcription stop

sites (TESs), indicating that 3′UTR length is unaffected in *MZpnrc2* mutants (Fig. 1C).

Normalized read coverage bedgraphs for the oscillatory gene transcripts *her1*, *her7*, *dlc*, and

rhov, all of which are overexpressed in *MZpnrc2* mutants, similarly show no differences in read

coverage between genotypes (Fig. S1A-B). Read coverage downstream of TESs, although very

low relative to gene body coverage, is higher in both WT and *MZpnrc2* mutants for transcripts

that are overexpressed relative to non-overexpressed transcripts in *MZpnrc2* mutants (Fig. 1C).

These results suggest that transcripts that depend on Pnrc2 for decay may have higher rates of

Pnrc2-independent transcriptional read-through beyond their annotated TES (or alternative

poly(A) cleavage site usage).

141   Our previous work showed that the Pnrc2-regulated *her1* transcript contains two critical

destabilizing motifs in the 3′UTR, a Pumilio Response Element (PRE) and AU-rich Element

(ARE) (Tietz et al., 2020). To determine whether 3′UTRs of overexpressed transcripts in

MZpnrc2 mutants are enriched for PREs, AREs, and/or additional mRNA motifs, we performed

motif analysis using the motif discovery tool STREME (Bailey, 2021). Among all 8 statistically

significant enriched motifs (Fig. 1D, Tables S4-S5), the sequences UGUACAU and UGUAAAUA

match a PRE, a motif associated with transcript destabilization (Arvola et al., 2020; Bohn et al.,

2018; Galgano et al., 2008; Gerber et al., 2006; Hafner et al., 2010; Lu and Hall, 2011; Morris et

al., 2008; Murata et al., 1995; Wang et al., 2002; White et al., 2001; Zamore et al., 1997;

Zamore et al., 1999). The enriched motifs UAUUUAUU and UAUUAUUA match AREs, motifs

recognized by a broad group of ARE-binding proteins (ARE-BPs) that can both inhibit and

promote mRNA decay and/or translation (Briata et al., 2005; Garneau et al., 2007; Gratacos and

Brewer, 2010; Lykke-Andersen and Wagner, 2005; Sanduja et al., 2011). Additional enriched

motifs include GUUUGUUU that matches GU-rich elements (GREs), a motif associated with

mRNA decay (Rattenbacher et al., 2010), and CCCCCUCC that contains a destabilizing CCUC

motif (Vejner et al., 2019). Lastly, we find enrichment for two miRNA target sites, the motif

KACUUGA that matches the miR-26 seed target site UACUUGA (Crippa et al., 2016; Icli et al.,

2013; Klett et al., 2019; Watterston et al., 2019) and the motif AGCACUU that matches the miR-

430 target site GCACUU (Baia Amaral et al., 2024; Bazzini et al., 2012; Giraldez et al., 2006;

Liu et al., 2020; Rabani et al., 2017; Vejnar et al., 2019). Overall, all significantly enriched 3′UTR

motifs in overexpressed transcripts are associated with mRNA-destabilizing functions.

162   STREME analysis of unaffected transcript 3′UTRs reveals significant enrichment for two

motifs, none of which overlap with motifs enriched in the overexpressed motif analysis (Fig. 1E,

Tables S4-S5). One motif is a UA-dinucleotide repeat that in yeast is associated with enhanced

poly(A) site cleavage (Graber et al., 1999). The second enriched motif, UACWGUAU, matches

the miR-144 seed target sites ACAGUAU and UACAGUA (Fu et al., 2009; Su et al., 2014;

Kretov et al., 2024). miR-144 is an erythroid-specific miRNA that is not expressed at mid-

segmentation stages (Fu et al., 2009), which may explain why miR-144 motifs are enriched in

3′UTRs of unaffected transcripts. STREME analysis on under-expressed transcript 3′UTRs

showed no motif enrichment (Fig. 1F, Tables S4-S5). Altogether, 3′UTR motif analysis shows

that Pnrc2 influences decay of mRNAs enriched with destabilizing RNA motifs.

To explore possible functions of differentially expressed transcripts in *MZpnrc2* mutants,

we conducted GO enrichment analysis (Thomas et al., 2022). GO analysis of overexpressed

transcripts identified many significantly enriched GO terms (Figs 1G, S1C-E,Tables S4-S5)

including somite specification and negative regulation of neuron differentiation, processes that

rely upon *Hes/her* function for proper differentiation (Bonev et al., 2012; Hatakeyama et al.,

2004; Ishibashi et al., 1995; Ohtsuka et al., 1999; Soto et al., 2020; Tan et al., 2012). Additional

enriched terms are related to developmental pathways involving Fgf, Wnt, Notch, and Ephrin

signaling (Fig. S1D, Table S5). Terms associated with post-transcriptional regulation are also

enriched including 3′UTR-mediated destabilization, deadenylation-dependent decay, and P-

body assembly (Fig. 1G, Table S5). Overexpressed genes in these categories may represent

transcripts that depend on Pnrc2 for their decay or transcripts that compensate for loss of Pnrc2

function. Alternatively, overexpressed genes might be genes that are transcriptionally repressed

by Pnrc2. GO analysis of under-expressed transcripts identified significant enrichment for terms

including translation, peptide biosynthetic process, and eukaryotic translation elongation factor

complex (Fig. S1F-H). Under-expressed genes in these categories may represent transcripts

that are stabilized by Pnrc2 or genes that are downregulated as a compensatory mechanism to

prevent over-translation of accumulated transcripts in *MZpnrc2* mutants. Alternatively, under-

expressed genes might be genes that are transcriptionally promoted by Pnrc2.

mRNA expression patterns of many developmental genes are disrupted in pnrc2 mutants

We characterized mRNA expression patterns and validated mRNA expression levels of

select genes associated with segmentation, oscillatory expression, and/or Fgf or Wnt signaling

that were identified as overexpressed in *pnrc2* mutants via RNA-Seq analysis. For in situ

hybridization, coloration reactions for each probe were developed in parallel for identical

reaction times in WT and *MZpnrc2* mutants. For RT-qPCR, expression levels were quantified

using whole embryo extracts from mid-segmentation stage WT and *MZpnrc2* mutants. The

expression pattern of *rhov*, an oscillatory gene in zebrafish PSM (Krol et al., 2011), is restricted

in WT embryos but appears expanded in *MZpnrc2* mutants (Fig. 2A, A' vs B, B'), although it is

possible that the difference reflects signal detection limits rather than absence of expression.

RT-qPCR validates that *rhov* expression is significantly increased in *MZpnrc2* mutants (Fig. 2C).

Expression of *sp5l* (Fig. 2D, D' vs E, E', RT-qPCR in F), a gene whose ortholog *Sp5* oscillates in

mouse PSM (Krol et al., 2011; Matsuda et al., 2020), is similarly affected. Expression of somite

polarity genes *mespab* and *ripply2* also show expanded and elevated expression in *MZpnrc2*
mutants (Fig. 2G, G' vs H, H'' and J, J' vs K, K', RT-qPCR in I and L). Expression of *dact1*, a
modulator of Wnt/β-catenin signaling (Waxman et al., 2004) that oscillates in mouse and human
PSM cells (Krol et al., 2011; Matsuda et al., 2020) is also expanded and elevated in *MZpnrc2*
mutants (Fig. 2M, M' vs N, N', RT-qPCR in O). Expression of *spry1* and *spry4*, genes encoding
Fgf signaling antagonists whose orthologs oscillate in mouse and/or human PSM cells (Matsuda
et al., 2020) is also expanded and elevated (Fig. 2P, P' vs Q, Q' and S, S' vs T, T', RT-qPCR in
R and U). Additional genes known to oscillate in the zebrafish PSM (Krol et al., 2011) and
whose transcripts are overexpressed in *MZpnrc2* mutants were quantified by RT-qPCR (Fig.
S2). Of the 20 currently annotated genes known to oscillate in the zebrafish PSM (Krol et al.,
2011), 10 (half) are overexpressed, 8 are unaffected, and 2 are under-expressed in *MZpnrc2*
mutants (Figs 2 and S2, data not shown). Overall, results show that Pnrc2 promotes decay of
mRNAs that encode many developmental regulators; however, the lack of overt segmentation or
additional developmental phenotypes in *MZpnrc2* mutants suggests that Pnrc2-dependent
mRNAs are regulated translationally.
219
Overexpressed oscillatory gene transcripts in MZpnrc2 mutants have shortened poly(A) tails
Despite overexpression of many oscillatory and segmentation-related genes, *MZpnrc2*
mutants have no overt segmentation phenotype. Discordance between mRNA and protein
levels might be due to deadenylation of accumulated transcripts that leads to inefficient
translation. To examine poly(A) (pA) tail length, we performed pA tail length (PAT) assays and
find that although the predominant *her1* pA tail length is ~45 nts in both WT and *MZpnrc2*
mutants, the length distribution is reduced in *MZpnrc2* mutants, which tend to have shorter tails
compared to WT embryos as revealed by electrophoretic pixel intensity profiling of PAT products
(Figs 3A-B, S3A). Similar reduced length distributions are observed for oscillatory gene
transcripts *her7*, *dlc*, and *rhov* (Figs 3A-B, S3B-D). Interestingly, PAT profiling reveals a
reduction in products smaller than the predominant tail length in *MZpnrc2* mutants (Fig. 3B),
suggesting that accumulated transcripts in *MZpnrc2* mutants are initially deadenylated
efficiently, but further deadenylation is inefficient in the absence of Pnrc2 function.
Because most oscillatory gene transcripts with short pA tails in *MZpnrc2* mutants would
likely be transcripts that accumulate due to loss of Pnrc2 function, we quantified *her1* and
additional oscillatory gene transcripts in pA-enriched and pA-depleted extracts obtained by
oligo(dT) bead-based purification (see methods). The oscillatory gene transcripts *her1*, *her7*,
dlc, and *rhov* are all significantly increased in pA-depleted extracts, but not pA-enriched

extracts, derived from *MZpnrc2* mutant versus WT embryos (Fig. 3C). Transcripts with short pA

tails may not be efficiently captured using oligo(dT) beads, so we quantified oscillatory gene

transcripts using cDNA synthesized from total RNA extracts and primed with oligo(dT) to select

for polyadenylated mRNA (rather than priming with random primers as performed for all other

quantification experiments). Quantification of oligo(dT)-primed cDNA shows that oscillatory gene

transcripts *her1*, *her7*, *dlc*, and *rhov* are all significantly overexpressed in *MZpnrc2* mutants (Fig.

S3F-I) indicating that accumulated transcripts, although shortened, are not completely

deadenylated. Because *MZpnrc2* mutants lack segmentation phenotypes, we hypothesize that

accumulated tail-shortened oscillatory gene transcripts are inefficiently translated.

Overexpressed ddx61 transcripts in MZpnrc2 mutants are substantially polyadenylated

Accumulation of mRNA in *MZpnrc2* mutants might lead to localization to cytoplasmic

granules such as processing bodies (P-bodies) that are sites of translationally repressed

transcripts (Luo et al., 2018; Ostareck et al., 2014). In eukaryotic cells, P-body formation is

enhanced in cells that accumulate mRNA decay intermediates (Sheth and Parker, 2003;

Teixeira et al., 2005) as well as in cells treated with the translational inhibitor puromycin that

results in ribosomes dissociating from mRNAs (Eulalio et al., 2007; Zheng et al., 2008).

Transcripts of several genes associated with the enriched GO term "P-body assembly" are

overexpressed in *MZpnrc2* mutants (Fig. S3J-K, Table S5). These include *ddx6* and *ddx61* that

are orthologous to the highly conserved *DDX6/p54* DEAD-box helicase gene family that is

required for P-body formation (Luo et al., 2018; Ostareck et al., 2014). Both *ddx6* and *ddx61* are

expressed broadly based on single-cell RNA-Seq (Sur et al., 2023; Farrell et al., 2018). Of the

two zebrafish *ddx6* ohnologs (Singh et al., 2020), *ddx61* transcript levels are 16-fold higher than

ddx6 at mid-segmentation stages (White et al., 2017), indicating that *ddx61* is the predominantly

expressed ohnolog during segmentation. Upon profiling *ddx61* pA tail lengths, a broad tail length

distribution is observed in both WT and *MZpnrc2* mutants, although there is a shift in distribution

to shorter products in mutants (Figs 3A-B, S3E). Transcript profiling in pA-enriched and pA-

depleted extracts reveals that *ddx61* is significantly increased in both fractions (Fig. 3C).

Detection of increased *ddx61* transcript levels in pA-enriched extracts suggests that *ddx61*

overexpression in *MZpnrc2* mutants might lead to increased Ddx61 protein levels.

Accumulated oscillatory gene transcripts in MZpnrc2 mutants are disengaged from translational

machinery, whereas overexpressed ddx61 transcripts are translationally engaged

Because accumulated oscillatory gene transcripts in *MZpnrc2* mutants have reduced tail

lengths, we assessed engagement of *her1* and other oscillatory gene transcripts with

ribosomes. We conducted polysome profiling from WT and *MZpnrc2* mutants to compare

oscillatory gene transcript levels in actively translating versus non-translating fractions (see

methods). The overall polysome profiles are similar in WT and *MZpnrc2* mutants (Fig. 4A),

suggesting that global translation is normal in mutants; however, *her1* transcripts in *MZpnrc2*

mutants are 20-fold higher in unbound/subunit fractions, and not significantly different in

monosome- and polysome-bound fractions compared to WT *her1* levels (Fig. 4B). Similar

distributions were obtained for other oscillatory gene transcripts including *her7*, *dlc*, and *rhov*

(Fig. 4C-E). Together, these results support the hypothesis that accumulated oscillatory gene

transcripts in *MZpnrc2* mutants are not efficiently translated. Because *ddx61* mRNA is increased

in pA-enriched RNA extracts (Fig. 3C), we profiled *ddx61* mRNA abundance across fractions

and find that *ddx61* levels are 3-fold higher in *MZpnrc2* polysome fractions relative to WT

polysome fractions (Fig. 4F). To corroborate *ddx61* polysome profiling results, we quantified

protein levels in WT and *MZpnrc2* mutant extracts using an antibody that recognizes both Ddx6

and Ddx61. Antibodies that distinguish the two highly similar proteins are unavailable.

Immunoblot results show a significant increase in Ddx6/Ddx61 protein levels in *MZpnrc2*

mutants at mid-segmentation stages (Fig. 4G-H), corroborating polysome profiling. Overall, our

results show that *ddx61* gene products, unlike oscillatory genes, are overexpressed at both

mRNA and protein levels in *MZpnrc2* mutants.

MZpnrc2 mutants are sensitized to deadenylation inhibition

Because several oscillatory gene transcripts have shortened pA tails in *MZpnrc2*

mutants, we hypothesized that blocking deadenylation would alleviate their inefficient translation

and lead to segmentation and/or additional phenotypes. Broad ectopic expression of a CCR4-

NOT dominant negative deadenylase via mRNA injection into zebrafish embryos has previously

been shown to disrupt segmentation and prevent pA tail shortening of oscillatory gene

transcripts *her1* and *her7* (Fujino et al., 2018). Consistent with published work, we show that

Myc-Cnot7DN mRNA injection causes segmentation defects in WT embryos, with over 80% of

injectants exhibiting completely malformed segments (Fig. S4A-D). Because segmentation clock

gene transcripts are accumulated, but likely not over-translated, in *MZpnrc2* mutants, we

hypothesized that *MZpnrc2* mutants would be sensitized to deadenylation inhibition. To test this

idea, we pursued an inducible deadenylase inhibitor expression strategy by injecting a plasmid

containing a heat-shock inducible version of the deadenylase inhibitor. After inducing expression

of Myc-tagged Cnot7DN for 30 minutes at mid-segmentation stages, WT and *MZpnrc2* mutants

were scored for segmentation defects 6 hours post-heat shock (see methods). Mosaic

expression of Myc-Cnot7DN leads to abnormal segment formation in *MZpnrc2* mutants, but not

in WT embryos (Fig. 5A-E), consistent with the hypothesis that *MZpnrc2* mutants are sensitized

to deadenylase inhibition. Using anti-Myc immunohistochemistry on plasmid-injected embryos,

we confirmed that Myc-Cnot7DN expression was induced post-heat shock and was absent in

no-heat shock controls (Fig. 5F-G). No significant Myc-Cnot7DN expression pattern differences

were observed between genotypes (Fig. S4I-M), showing that *MZpnrc2* mutant sensitivity is not

due to differences in transgene induction. When comparing transgene expression patterns

between *MZpnrc2* mutants with and without somite defects, all *MZpnrc2* mutants with very high

Myc-Cnot7DN expression (i.e. embryos showing broad expression with little to no mosaicism)

displayed somite defects, though overall, expression pattern differences between *MZpnrc2*

mutants with and without somite defects did not reach significance (p = 0.056) (Fig. S4N).

MZpnrc2 mutants are sensitized to depletion of Ddx61 and Ddx6 function

Because *ddx61* is overexpressed at both mRNA and protein levels, we hypothesized

that depleting Ddx61 function in *MZpnrc2* mutants might lead to segmentation and/or additional

phenotypes. Using a highly efficient F0 CRISPR knockdown strategy (Hoshijima et al., 2019;

Klatt Shaw and Mokalled, 2021), we find that single *ddx61* crispants and single *ddx6* crispants

develop normally up to 5 days post-fertilization (dpf) and can be raised without significant losses

up to adulthood (data not shown), consistent with *ddx6* and *ddx61* genetic mutant analyses

(Pronobis et al., 2025). To estimate F0 CRISPR efficiency (Fig. S5A), we employed high

resolution melt analysis (HRMA) to detect CRISPR-induced mutations and confirmed that all

CRISPR-injected embryos were mutagenized (Fig. S5B-C). Because *ddx6* might compensate

for loss of *ddx61* function, we also targeted *ddx6*. In *ddx6*;*ddx61* crispants, *ddx6* and *ddx61*

mRNA levels are more than 50% reduced at 18 hpf compared to uninjected sibling controls (Fig.

S5D-E), likely because the premature termination codon-containing transcripts created by our

targeting strategy are degraded by NMD. Corresponding with reduced mRNA levels,

Ddx6/Ddx61 protein levels are reduced in *ddx6*;*ddx61* crispants at 18 hpf and are nearly absent

by 4 dpf (Fig. S5F). To assess whether *MZpnrc2* mutants are sensitized to Ddx6/Ddx61

depletion, we compared *ddx6*;*ddx61* crispant phenotypes in WT and *MZpnrc2* mutant

backgrounds over time. At mid-segmentation stages, most *ddx6;ddx61* crispants are

morphologically normal in WT and *MZpnrc2* mutant backgrounds (Fig. 6A-B, E-F, I). By 19 hpf,

~30% of *MZpnrc2 ddx6*;*ddx61* crispants exhibit morphological abnormalities, whereas

ddx6;*ddx61* crispants in the WT background (hereafter referred to as *ddx6*;*ddx61* crispants) as

well as uninjected controls for both genotypes are predominantly normal (Fig. 6C-D, G-H, J).

Defects worsen with time in *MZpnrc2 ddx6*;*ddx61* crispants, with nearly all crispants showing

abnormalities at 60 hpf when *ddx6*;*ddx61* crispants are mostly normal (Fig. S6A-J). By 84 hpf,

nearly half of *ddx6*;*ddx61* crispants show abnormal phenotypes similar to those observed earlier

in *MZpnrc2 ddx6;ddx61* crispants (Figs 7A-D, S6K). Similar proportions of abnormal phenotypes

are observed at 112 hpf and 120 hpf (Figs 7E-F, S6L) and correspond with strongly reduced

Ddx61/Ddx6 protein levels in *ddx6*;*ddx61* crispants (Figs 7G, S5F-G). Overall, these results

support that *MZpnrc2* mutants are sensitized to reduced Ddx6/Ddx61 function.

To determine whether *her1* expression is affected in *ddx6;ddx61* crispants, we

performed *her1* in situ hybridization at mid-segmentation stages. Although the *her1* expression

pattern in *ddx6*;*ddx61* crispants was indistinguishable from uninjected WT controls (Fig. 8A,A' vs

B,B'), *MZpnrc2 ddx6*;*ddx61* crispants showed an expanded misexpression phenotype compared

to control *MZpnrc2* mutants (Fig. 8C,C' vs D, D'). We quantified *her1* levels by RT-qPCR and

find that *her1* is more than 1.5-fold increased in *MZpnrc2 ddx6*;*ddx61* crispants relative to

control *MZpnrc2* mutants (Fig. 8E), but is unaffected in *ddx6*;*ddx61* crispants. In contrast, *her7*

expression is reduced, whereas *dlc* and *rhov* expression is unaffected in *MZpnrc2 ddx6*;*ddx61*

crispants (Fig. 8F-H). Overall, normal levels of oscillatory gene transcripts in *ddx6*;*ddx61*

crispants suggest that Ddx61 and Ddx6 are not required for mRNA decay when Pnrc2 is

functional. The synergistic effect of loss of *pnrc2*, *ddx6*, and *ddx61* shows that Ddx6 and Ddx61

partially compensate for loss of Pnrc2 function.

Discussion

In this work, we show that overexpressed transcripts in *MZpnrc2* mutants are primarily

protein-coding transcripts, whose sequences are enriched for features known to promote

transcript decay, including uORFs, long 3'UTRs, and mRNA destabilization sequences. Human

PNRC2 bridges NMD machinery by directly binding NMD factors Upf1 and Dcp1 (Cho et al.,

2009; Lai et al., 2012). Although depleting *upf1* via morpholino (MO) injection in zebrafish

embryos has no effect on *her1* expression, co-depletion of *upf1* and *pnrc2* via MO injection

enhances *her1* misexpression compared to depletion of *pnrc2* alone (Gallagher et al., 2017),

suggesting that Upf1 is important for the decay of *her1* and additional overexpressed transcripts

in *MZpnrc2* mutants.

Additional features of overexpressed transcripts in *MZpnrc2* mutants include enrichment

of mRNA destabilizing sequences, including PREs and AREs that are present in the oscillatory

gene transcripts *her1*, *her7*, and *dlc.* Among overexpressed transcripts, many correspond to
genes that are dynamically regulated, including oscillatory and/or developmental genes that
function in developmental signaling pathways. Remarkably, despite overexpression of >1,700
transcripts, segmentation and other developmental processes appear normal in *MZpnrc2*
mutants, although many fail to thrive as juveniles and adults. We previously suggested that
absence of segmentation defects is likely because accumulated mRNAs in *MZpnrc2* mutants
are not efficiently translated (Gallagher et al., 2017), though the mechanism was unknown. In
the current study, we explored mechanisms driving discordant expression of mRNA and protein
in *MZpnrc2* mutants by assessing the polyadenylation status of *her1* and other transcripts, their
level of engagement with translational machinery, and the impact of P-body factors Ddx6 and
Ddx61 on the *MZpnrc2* overexpression phenotype.

Inhibiting deadenylation in MZpnrc2 mutants leads to somite defects
Poly(A)-tail length analysis and polysome profiling suggests that many accumulated
oscillatory gene transcripts have reduced tail lengths in *MZpnrc2* mutants and are primarily
disengaged from ribosomes. These results are consistent with the idea that accumulated
oscillatory gene transcripts are not over-translated in *MZpnrc2* mutants. In support of this view,
inhibiting deadenylation leads to somite defects in *MZpnrc2* mutants under conditions that have
no effect in WT embryos, demonstrating that *MZpnrc2* mutants are sensitized to deadenylation
inhibition. We hypothesize that *her1* and additional oscillatory gene transcripts are over-
translated after inhibiting deadenylation in *MZpnrc2* mutants, which we will investigate in the
future using stably integrated inducible *Myc-cnot7DN* transgenic lines to quantify protein levels
and pA-tail length after deadenylation inhibition.

Depletion of RNA helicases Ddx6 and Ddx61 leads to developmental defects in MZpnrc2
mutants
We show here that Ddx6/Ddx61 RNA helicases are overexpressed at both mRNA and
protein levels in *MZpnrc2* mutants. Depletion of *ddx6* and *ddx61* function in *MZpnrc2* mutants
increases *her1* accumulation and leads to strong morphological phenotypes that are observed
at much lower frequency in WT crispants. This difference shows that *MZpnrc2* mutants are
sensitized to loss of *ddx6* and *ddx61* function. Future work will assess whether Pnrc2-regulated
transcripts are over-translated upon depletion of Ddx6 and Ddx61 function in *MZpnrc2*
crispants. It is possible that Ddx6/ Ddx61 promotes tail-shortening of accumulated mRNAs in

MZpnrc2 mutants, however a connection between deadenylation inhibition and Ddx6/Ddx61

depletion experiments has not been established.

Morphological defects are not observed in *ddx6*;*ddx61* crispants at 1 dpf despite

substantial Ddx6/Ddx61 protein depletion, potentially because maternally provided *ddx6/ddx61*

function is sufficient for normal early development. The absence of early phenotypes differs from

previous work that showed that *ddx6* and *ddx61* splice-blocking morpholino (MO) injection leads

to a curved tail phenotype in 24 hpf embryos (Zampedri et al., 2016), which could be due to

differences in knockdown approach and/or efficiency. Despite early differences, embryos

injected with *ddx6* and *ddx61* MOs survive to at least 5 dpf (Zampedri et al, 2016), similar to

what we observed for *ddx6*;*ddx61* crispants. More recently, genetic analysis of *ddx6* and *ddx61*

mutants showed that doubly homozygous *ddx6;ddx61* mutants are inviable by 1 dpf (Pronobis

et al., 2025). Although doubly homozygous *ddx6*;*ddx61* genetic mutants also have maternally

provided *ddx6/ddx61* function, we suspect that the differences between doubly homozygous

ddx6;*ddx61* genetic mutants versus *ddx6*;*ddx61* crispants and morphants is likely because

genetic mutants completely lack zygotic Ddx6/Ddx61 protein whereas crispants and morphants

likely have low levels of zygotic Ddx6/Ddx61 protein. Further analyses of genetic mutants will

clarify the role of Ddx6/Ddx61 function during early development.

In contrast to *ddx6*;*ddx61* crispants that appear largely normal up to 60 hpf, *MZpnrc2*

ddx6;ddx61 crispants show abnormalities as early as 19 hpf. We hypothesize that *MZpnrc2*

mutants require more Ddx6/Ddx61 function than WT embryos because of negative effects of

transcript accumulation. In fact, Ddx6/Ddx61 protein levels may be increased in *MZpnrc2*

mutants as a compensatory response that buffers against the negative effects of transcript

accumulation. The finding that *her1* mRNA levels in *MZpnrc2* mutants is further increased in

ddx6;*ddx61* crispants suggests that P-body factors promote *her1* mRNA decay in the absence

of Pnrc2. P-bodies are thought to primarily function as sites of mRNA storage rather than decay,

supported by observations that mRNA decay continues in the absence of P-bodies (Decker et

al., 2007; Eulalio et al., 2007; Stoecklin et al., 2006). Early reports of mRNA decay in P-bodies

are attributable to artifacts of the MS2 tethering system (Garcia & Parker, 2015; Heinrich et al.,

2017; Tutucci et al., 2018). It is also possible that Ddx6/Ddx61 promotes *her1* mRNA decay in a

P-body-independent manner, as both RNA helicases show diffuse cytoplasmic expression in

addition to enrichment in cytoplasmic foci (Zampedri et al., 2016). Future work will clarify the

extent to which *her1* mRNA is degraded in the cytoplasm versus in P-bodies and the

dependence of these activities on *pnrc2* function.

It is intriguing that among the 4 oscillatory gene transcripts we quantified, only *her1*
levels are increased in *MZpnrc2 ddx6*;*ddx61* crispants. It is possible that the *her1* transcript
contains features that promote Ddx6/Ddx61-dependent mRNA decay that are absent in *her7*,
dlc, and *rhov* transcripts. Alternatively, the level of Ddx6/Ddx61 depletion in *MZpnrc2* mutants
may not have been sufficient to detect accumulation of *her7*, *dlc*, and *rhov* oscillatory gene
transcripts. Future analysis using genetic mutants will help to distinguish these possibilities.
445

Conclusions
Pnrc2 promotes rapid mRNA decay during vertebrate segmentation, including the decay
of many dynamically expressed oscillatory gene transcripts. Overexpressed transcripts in
MZpnrc2 mutants are enriched for mRNA-destabilizing motifs that are associated with
translational repression and mRNA decay. *MZpnrc2* mutants develop normally despite
dysfunctional mRNA decay, likely because transcripts that depend on Pnrc2 for rapid decay
have shortened poly(A) tails that are inefficiently translated. Consistent with this view, inhibiting
deadenylation leads to somite defects in sensitized *MZpnrc2* mutants, potentially due to over-
translation of Pnrc2-regulated transcripts. Upregulation of *ddx61* at both mRNA and protein
levels suggests that compensatory mechanisms are employed in decay-deficient *pnrc2* mutants.
The synergistic effect of loss of *pnrc2*, *ddx6*, and *ddx61* leads to strong developmental
phenotypes, which may be due to over-translation of Pnrc2-regulated transcripts. Overall, our
results show that Ddx6 and Ddx61 are required for normal embryonic development in mRNA-
decay deficient *MZprnc2* mutants.
460

Materials and methods
Animal stocks and husbandry
All protocols using zebrafish were approved by the Ohio State University Institutional Animal
Care and Use Committee (protocol 2012A00000113). Adult zebrafish strains (*Danio rerio*) were
kept at 28.5°C on a 14 hour (h) light/10 h dark cycle and obtained by natural spawning or *in vitro*
fertilization (Westerfield, 2007). Embryos and larvae were staged according to Kimmel et
al.,1995. The *pnrc2*^{oz22} line has been described previously (Gallagher et al., 2017).
468

DNA extraction and genotyping
Individual embryos and adult fin tissue were lysed in 50 ul 1X ThermoPol Buffer (NEB) at 95°C
for 10 minutes, digested at 55°C for 1 hour in 1 mg/ml Proteinase K (BP1700, ThermoFisher),
followed by Proteinase K inactivation at 95°C for 10 minutes. Molecular identification of *pnrc2*^{oz22}

carriers by PCR amplification followed by NsiI-HF digestion to distinguish cleavable WT from

uncleavable mutant amplicons was performed as previously described (Tietz et al., 2020).

RNA-Seq sample collection and sequencing

Whole embryos at mid-segmentation stages (n = 50 embryos per biological replicate for each

genotype) were solubilized in Trizol for RNA extraction and purified following standard

procedures (Thermo Fisher). All genotypes were prepared in biological triplicate. 2 ug of total

RNA from each sample was processed at Novogene using a whole transcriptome sequencing

workflow with ribosomal RNA (rRNA) depletion that enables sequencing of RNA irrespective of

polyadenylation status. Each sample was depleted of rRNA using a Ribo-Zero Kit (Illumina),

followed by chemical fragmentation and 150 bp paired-end library synthesis using a NEBNext

Ultra II Directional RNA kit (NEB). Libraries were sequenced on the Illumina NovaSeq 6000

platform.

RNA-Seq data analysis

High-throughput sequencing reads were trimmed to remove adapter and primer sequences

using Cutadapt version 3.4 (Martin, 2011) and pseudoaligned to the zebrafish

GRCz11/danRer11 transcriptome reference using Kallisto (version 0.46.2), an RNA-Seq tool

that rapidly quantifies transcript abundance without the need for individual base alignment (Bray

et al., 2016). Cutadapt and Kallisto were installed and run in Python (version 2.7.5) using

standard procedures described in online guides available at

https://cutadapt.readthedocs.io/en/stable/guide.html and

https://pachterlab.github.io/kallisto/manual. Prior to running Kallisto, all sequence data files were

assessed using the quality control tool FastQC version 0.12.1 (Andrews, 2010) and no quality

issues were identified in relevant modules for RNA-Seq analysis (data not shown). Transcripts

with <5 estimated read counts in at least one sample were filtered out to reduce noise.

Differential gene expression analysis was performed using Sleuth version 0.30.1, a program

that uses variance decomposition to identify biological differences between RNA-Seq samples

(Pimentel et al., 2017). Sleuth was run in RStudio version 2023.06.2 Build 561 (Posit Software)

using standard procedures described in an online guide at

https://pachterlab.github.io/sleuth_walkthroughs/trapnell/analysis.html. Batch effects including

effects due to sequencing lane or biological replicate differences were tested and no effects

were observed (data not shown). A summary of the RNA-Seq processed data, Kallisto

normalized read counts, and Sleuth-generated differential expression results are included as

supplementary data (Tables S1-S3).

Genomic mapping and metagene plots

High throughput sequencing reads were trimmed to remove adapters using bbduk.sh v39.13

with k=50, mink=10, hdist=1, minoverlap=8, minlength=20, trimq=10, tbo and tpe options. Then,

trimmed reads were aligned to the GRCz11 zebrafish genome using STAR v2.7.10b with an

ensembl v113 GTF file. The --outWigType bedGraph option was used with STAR to allow

visualization of coverage for each gene (Fig. S1A-B). To generate metagene plots, various

modules from deepTools v3.5.6 were used. First, the bamCoverage module was used to

generate strand specific counts per million normalized bigWig files containing the coverage at

different genome positions for each sample. Then the computeMatrix module was used in scale-

regions mode to produce intermediate files containing coverage for each gene scaled to a fixed

length, as well as the coverage 1000 bases downstream of the genes. Two sets of

regions/genes were defined at the computeMatrix step, one set containing genes that

correspond with overexpressed transcripts in *MZpnrc2* mutants, and one set containing all other

genes that were expressed in *MZpnrc2* mutants (i.e. non-overexpressed genes). The

computeMatrixOperations utility was then used to combine the computeMatrix output for each

strand into a single file. plotProfile was then run with --averageType mean as well as the –

outFileNameData option in order to write the metagene coverage data to a file. In R v4.4.0, the

coverage data was then averaged within each experimental condition, and the coverage was

standardized so that the sum of the coverage for each condition/gene set combination summed

to one across all positions on the metagene plot. The standardized coverage was then plotted

using ggplot2 v3.5.2.

uORF enrichment analysis

To identify transcripts with functional uORFs, we used a published dataset of zebrafish

transcripts containing high-confidence uORFs from 24 hpf embryos (Johnstone et al., 2016).

Among the time points sampled in Johnstone et al (2016), the 24 hpf dataset was the closest to

mid-segmentation stages. Similar enrichment results were obtained using the 12 hpf dataset

(Johnstone et al., 2016) (data not shown). To select for uORFs with evidence of translation, we

followed methods in Gangras et al (2020) to filter the 24 hpf dataset (Johnstone et al., 2016)

using the following criteria: 5'UTR RPF RPKM ≥ 5, RNA-Seq RPKM≥ 5, RPF RPKM ≥ 5, and

ORF translation efficiency (RPF RPKM/RNA-Seq RPKM) >1. In addition to these criteria, the

functional uORF list was filtered to only include APPRIS primary protein coding transcript

isoforms that are functionally important (Rodriguez et al., 2022). Next, to identify functional

uORF-containing transcripts in our dataset, we filtered our expression dataset (Table S3) to only

include APPRIS primary protein coding transcript isoforms and compared this list with the

filtered functional uORF list. A one-sided Fisher's exact test was applied to evaluate the

enrichment of transcripts with uORFs among the transcripts that were significantly

overexpressed in *MZpnrc2* mutants (q < 0.05) versus transcripts that were non-overexpressed

in *MZpnrc2* mutants (Table S3). hpf = hours post-fertilization; RPF = ribosome-protected mRNA

fragment; RPKM = reads per kilobase mapped.

Motif analysis

Motif analysis was performed using 3'UTR input sequences corresponding to transcripts

identified from RNA-Seq analysis of *MZpnrc2* mutant embryos (Tables S3-S4) and the MEME

Suite tool STREME version 5.5.6 (Bailey et al., 2021). Annotated 3'UTR sequences from each

of the following datasets were analyzed using STREME: 1) significantly overexpressed

transcripts (q < 0.05; n = 1320), 2) unaffected transcripts (log2FC from -0.05 to 0.05; n = 1334),

and 3) significantly under-expressed transcripts (i.e. transcripts with decreased expression in

MZpnrc2 mutants relative to WT embryos; q < 0.05; n = 690) (Tables S3-S4). 3'UTR sequence

annotations were obtained using BioMart (Ensembl release 113, GRCz11 assembly) and those

less than 50 nts in length were removed to reduce annotation and length biases. Motif range

was set to 6-8 nts and shuffled input sequences were used as the control with Markov order set

to 2 (the default setting used by STREME). Sequence logos (Schneider and Stephens, 1990;

Schneider et al., 1986) were generated using probability matrix data from STREME and the

seqLogo package (Bembom and Ivanek, 2024) available with BioConductor version 3.19

(Gentleman et al., 2004; Huber et al., 2015) in RStudio version 2023.06.2 Build 561 (Posit

Software). Motif enrichment results are reported in Table S5.

GO analysis

GO analysis was performed using the PANTHER Overrepresentation Test (Released 2024-02-

26) that is available from the Gene Ontology Consortium at https://geneontology.org/ (Mi et al.,

2013; Thomas et al., 2022). The GO Ontology database version used was DOI:

10.5281/zenodo.10536401 (Released 2024-01-17). The input list contains gene IDs for which at

least one transcript was significantly overexpressed (q < 0.05) in *MZpnrc2* mutants and for

which a GO annotation was identified (n = 1619 uniquely annotated GO IDs) (Table S4), while

the reference list contains the full set of GO-annotated zebrafish genes in the database (n =

26,353). GO results are reported in Table S5. Statistical test type was a Fisher's Exact test

corrected by calculating the false discovery rate.

In situ hybridization

Whole mount in situ hybridization was performed as previously described (Broadbent and Read,

1999; Jowett, 1998) using digoxygenin (DIG)-labeled riboprobes. Riboprobe for *her1* was

previously described (Delaune et al., 2012; Dill and Amacher, 2005). Riboprobe for *rhov* was

synthesized from a plasmid containing the *rhov* cDNA (cb832) received from the Zebrafish

International Resource Center (ZIRC). *rhov*-containing plasmid was linearized with NotI-HF

(NEB) and transcribed with T7 RNA polymerase (Roche Life Science) to make DIG-labeled

antisense *rhov* riboprobe. Riboprobe templates for *sp5l*, *mespab*, *ripply2*, *dact1*, *spry1*, and

spry4 were generated by PCR amplification of coding sequence (primer sequences are listed in

Table S6). All DIG-labeled PCR-based probes were synthesized by transcribing PCR templates

with SP6 or T7 RNA polymerase (Roche Life Science). *dact1*, *sp5l*, and *spry4* produced

expression patterns consistent with direct submission in situ hybridization data (Thisse et al.,

2001; Thisse et al., 2005). Additionally, expression patterns for all PCR-based probes shown in

Fig. 2 match published expression patterns for WT embryos at mid-segmentation stages,

including *dact1* (Gillhouse et al., 2004), *mespab* (Cutty et al., 2012; Wanglar et al., 2014), *sp5l*

(Superina et al., 2014), *ripply2* (Junker et al., 2014; Kinoshita et al., 2018; Moreno et al., 2008;

Wanglar et al., 2014; Yabe et al., 2016), *spry1* (Komisarczuk et al., 2008), and *spry4* (Nguyen-

Chi et al., 2012). In situ hybridized embryos were mounted in Permount and imaged using an

Axiocam HRc digital camera with AxioPlan2 microscope and 10x objective (Zeiss).

RNA extraction and cDNA synthesis

Whole embryos at mid-segmentation stages (n = 15 embryos per biological replicate for each

genotype in Fig. 2; n = 10 embryos per biological replicate for each genotype and condition in

Fig. 7) were solubilized in Trizol for RNA extraction and purified following standard procedures

(Thermo Fisher). All genotypes and conditions were performed in biological triplicate. 500 ng of

total RNA was reverse transcribed with random primers and Superscript IV reverse

transcriptase according to the manufacturer's instructions (Thermo Fisher).

Quantitative RT-PCR analysis

Quantitative RT-PCR was performed using PowerUp SYBR Green Master Mix (Thermo Fisher)

and 4.5 ul cDNA (diluted 1:50 in Fig. 2, diluted 1:25 in Fig. 7) in 20 ul reactions, following

manufacturer's procedures. Negative controls lacking template were included for each primer

set. All reactions were subjected to thermal melting to confirm that each reaction gave single

peaks. Transcript levels were normalized to *mobk13* (*mob4*) (Gangras et al., 2020; Hu et al.,

2016; Tietz et al., 2020). Cycle thresholds (Ct) were determined using CFX Maestro version 2.3

software (Bio-Rad). The average Ct of 2 technical replicates per sample was used for all fold-

calculations except for calculations in Fig 4 for which the average Ct of 3 technical replicates per

sample was used to more accurately quantify RNA from low-abundance fractions. Changes in

mRNA expression were calculated by $\Delta\Delta Ct = \Delta Ct$ target $- \Delta Ct$ control. Relative changes in

mRNA expression levels are represented graphically as fold change, where relative mRNA fold

change = $2-\Delta\Delta Ct$. All graphs and statistics were generated using Prism 10 (GraphPad).

Poly(A) Tail-Length Assay

RNA for poly(A) tail-length assays was obtained by collecting whole embryos at mid-

segmentation stages (n = 18 embryos for each genotype) followed by solubilization in Trizol and

purification following standard manufacturer procedures (Thermo Fisher). Two biological

replicates were performed for each genotype and produced consistent results (data not shown).

Tail-lengths for *her1, her7, dlc, rhov,* and *ddx61* transcripts were assessed using the Poly(A)

Tail-Length Assay Kit following standard manufacturer procedures (Thermo Fisher). Minus RT

controls were performed for all samples and no amplification was detected as expected (data

not shown). Primers used for tail-length assay PCR are listed in Table S6. Gel images were

captured using an AlphaImager system (ProteinSimple) with exposure times set just below

saturation for each lane analyzed. Pixel intensity profiles were generated using the lane profile

analysis tool in AlphaImager HP version 3.4.0 (build 0728) and exported as png files for manual

tracing using the anchor point tool in Adobe Illustrator version 24.1.2. Peaks were identified

using the auto-detection option that identifies the upper and lower thresholds for peaks. Lane

profiles shown in Fig. 3B are scaled individually for each lane profile to better illustrate pixel

intensity distributions, whereas lane profiles in Fig. S3A-E are universally scaled between WT

(lane 2) and *MZpnrc2* (lane 3) profiles to better illustrate differences in pixel intensities between

lanes. Molecular weights were calculated using a molecular weight tool in AlphaImager that

utilizes known sizes of the molecular weight ladder to determine the size and range of PCR

products of interest.

Preparation and analysis of poly(A)-enriched and poly(A)-depleted extracts

RNA for comparison of expression levels in poly(A)-enriched versus poly(A)-depleted RNA

fractions was obtained from the same RNA sources as was used for RNA-Seq experiments.

Three biological replicates were obtained for both WT and *MZpnrc2* mutant embryos. From

each sample, 1 ug of total RNA was adjusted to a final volume of 50 ul in bead binding buffer

(20 mM Tri-HCl, pH 7.5, 1.0 M LiCl, and 2 mM EDTA) and incubated with Dynabeads

Oligo(dT)$_{25}$ following manufacturer procedures (Thermo Fisher). RNA bound to beads (poly(A)-

enriched RNA) was eluted in 10 mM Tris-HCl (pH 7.5). Unbound RNA (poly(A)-depleted RNA)

was precipitated by adding 0.1x volume 3M sodium acetate and 2x volume ice-cold ethanol,

followed by vortexing and incubation overnight at -80°C. After centrifugation at 4°C for 20

minutes at 12,000 x g, precipitated pellets from unbound RNA fractions were rinsed with ice-cold

70% ethanol and resuspended in 10 mM Tris-HCl (pH 7.5) at an equivalent volume used for

elution of bound RNA fractions. Each RNA fraction was reverse transcribed using random

primers and Superscript IV reverse transcriptase (RT) according to the manufacturer's

instructions (Thermo Fisher). As a control, 1 ug of total RNA from the same samples used for

bead-based fractionation were reverse transcribed in parallel. RT-qPCR analysis was performed

as described earlier using 4.5 ul of cDNA diluted 1:10.

Protein extraction and Immunoblot analysis

Protein extracts were prepared from zebrafish embryos or larvae at 1-5 days post-fertilization.

Individuals were collected and washed twice with rapid pipetting to disrupt yolks in 200 ul of ice-

cold Ringer's solution with protease inhibitors (116 mM NaCl, 2.9 mM KCl, 5.0 mM HEPES, pH

7.2, 100 µg/mL Soybean Trypsin Inhibitor [Sigma], 0.2 mg/mL PMSF, Complete Protease

Inhibitor Cocktail [Roche]). Embryo and larval suspensions were centrifuged at 4°C for 1 minute

at 300 x g for each wash. Cell lysis was performed by addition of 1 ul 4X LDS Buffer (Thermo

Fisher) for each embryo or larva collected (n = 20-65 individuals per extraction), followed by

homogenization with a plastic pestle, vortexing for 1 minute, and boiling for 10 minutes.

Homogenization, vortexing, and boiling were performed twice sequentially. Embryo or larval

equivalents were loaded on a 4-12% Novex Bis-Tris polyacrylamide gel (Thermo Fisher),

electrophoresed in NuPAGE MOPS SDS running buffer (Thermo Fisher) alongside a Novex

Sharp pre-stained protein standard (Thermo Fisher), and transferred to a PVDF membrane

using the Bio-Rad wet transfer system following standard manufacturer procedures. Immunoblot

was blocked in 5% non-fat skim milk in Tris-buffered saline (20 mM Tris, 150 mM NaCl, pH 7.5)

and subsequent steps were performed following standard procedures available online from Bio-

Rad in bulletin 6376 (https://www.bio-rad.com/). The immunoblot was co-incubated with DDX6

(A16270, AbClonal) and β-actin (sc-47778, Santa Cruz Biotechnology) primary antibodies

followed by co-incubation with AP-conjugated goat anti-rabbit IgG (H+L) secondary antibody

(ThermoFisher) to detect DDX6 primary antibody and goat anti-mouse IgG (H+L) secondary

antibody (Thermo Fisher) to detect β-actin primary antibody. In control experiments, immunoblot

antibody incubations were performed separately and showed no differences in DDX6 and β-

actin signal detection (data not shown). Primaries were diluted 1:1000 and secondaries were

diluted 1:10,000 in blocking solution with 0.1% Tween-20. Primary antibody incubations were

performed overnight at 4°C, secondary antibody incubations were performed for 1 hour at room

temperature. Chemiluminescent detection was performed using CDP-Star following the

manufacturer's procedure (Thermo Fisher) and imaged using the Sapphire FL Biomolecular

Imager (Azure Biosystems). Quantification of signal intensities was performed using lane profile

analysis in AlphaImager HP version 3.4.0, build 0728 (Protein Simple).

Plasmid construction & transgenesis

The heat shock construct *hsp70l:Myc-cnot7DN-SV40 pA* was assembled by first constructing

plasmid *pMA-Myc-cnot7DN* using the GeneArt Plasmid Construction Service (Thermo Fisher).

pMA-Myc-cnot7DN contains the coding sequence for a dominant negative form of zebrafish

Cnot7 (*Cnot7DN*) with an N-terminal C-myc epitope tag (Evan et al., 1985) and intervening

SGLRS flexible linker peptide. The *myc*-tagged *Cnot7DN* coding sequence was based on

plasmid M338 (*pCS2 + MT-CNOT7-D40A,E42E,C67E,L71E-sv40*) that has previously been

used to inhibit deadenylase function in zebrafish embryos (Fujino et al., 2018; Makino et al.,

2015; Mishima and Tomari, 2016; Mishima and Tomari, 2017). After sequence validation of

pMA-Myc-cnot7DN, the *Myc-cnot7DN* insert was cloned into *pBSKI2* (Thermes et al., 2002)

using standard restriction digestion-based cloning with the enzymes NheI-HF and BamHI-HF

(NEB). Plasmid and I-SceI enzyme were injected into the blastomere of 1-cell stage WT and

MZpnrc2 mutant embryos following standard transgenesis procedures (Thermes et al., 2002).

To determine a suitable dose, we injected 20 pg and 60 pg plasmid doses in the absence of

heat-shock induction and assessed morphology at mid-segmentation stages; the 20 pg dose

was well-tolerated (73 of 90 injected embryos had WT morphology), but the 60 pg dose caused

gross abnormalities (0 of 56 injected embryos had WT morphology).

Heat shock assay

Plasmid-injected embryos (were collected and raised to segmentation stages (10-12S), heat

shocked at 37□C for 30 minutes, and monitored for somite defects at 30 minute intervals post-

heat shock (pHS). Because *MZpnrc2* mutants are developmentally delayed relative to WT

embryos by ~2-3 hours, heat shock induction was performed on stage-matched embryos from

WT and *MZpnrc2* clutches based on somite number, requiring staggered heat shocks beginning

with WT embryos followed by heat shocks 2-3 hours later for *MZpnrc2* mutant embryos when

they reached the same somite stage. Live embryos at 6 hours post-heat shock were mounted in

3% methylcellulose and imaged at 20x magnification using AxioVision Software (Zeiss) on a

MZFLIII Fluorescence Stereo Microscope (Leica) with an AxioCam HRc digital camera (Zeiss)

and subsequently fixed in 4% PFA for anti-Myc immunohistochemistry.

Anti-Myc Immunohistochemistry

Embryos injected with *hsp70l:Myc-cnot7DN* plasmid that were heat shocked at ~14 hours post

fertilization (hpf) for 30 minutes were collected 6 hours post-heat shock (~22 hpf) alongside non-

heat shocked controls, fixed in 4% PFA in PBS overnight at 4$^-$C, and washed 5 x 5 minutes in

PBS with 0.1% Tween-20 (PBST). Embryos were then incubated in water for 5 minutes,

permeabilized with ice-cold acetone for 7 minutes, and incubated again in water for 5 minutes

followed by 3 x 5 minute PBST washes. Blocking was performed in PBST with 2% BSA, 2%

goat serum for 1 hour at room temperature, followed by incubation with 1:500 dilution

monoclonal anti-myc (clone 9E10, MA1-980, Thermo Fisher) in PBS with 2% BSA, 2% goat

serum, 1% DMSO and 0.1% Triton X-100 (PBDTx) overnight at 4$^-$C. Embryos were washed 8 x

15 minutes in PBST, blocked as described above, followed by incubation with 1:200 dilution

Alexa Fluor 546 goat anti-mouse secondary antibody (Thermo Fisher) in PBDTx overnight at

4$^-$C. Embryos were washed 8 x 15 minutes in PBST prior to imaging. Immuno-stained embryos

were mounted in Fluoromount-G and imaged at 20x magnification using AxioVision Software

(Zeiss) on an MZFLIII Fluorescence Stereo Microscope (Leica) with a mCherry filter set and

AxioCam HRc digital camera (Zeiss).

Polysome profiling

WT and *MZpnrc2* mutant embryos were raised to mid-segmentation stages and treated in 100

ug/ml cycloheximide for 5 minutes (n = 300 embryos per biological replicate for each genotype).

Embryos were triturated using a 200 ul pipette and washed to remove yolks as previously

described (Link et al., 2006) except that E2 embryo medium containing cycloheximide (15 mM

NaCl, 0.7 mM NaHCO$_3$, 0.5 mM KCl, 0.15 mM KH$_2$PO$_4$, 2.7 mM CaCl$_2$, 0.5 mM Na$_2$HPO$_4$, 1mM

MgSO$_4$, 100 ug/ml cycloheximide) was used in lieu of 0.5x Ginzburg Fish Ringer's buffer. Cell

pellets were resuspended in 500 ul of chilled lysis buffer (20 mM Tris HCl pH 7.5, 30 mM MgCl$_2$,

100 mM NaCl, 0.5 uM DTT, 100 ug/ml cycloheximide, 1 mg/ml heparin, 2.5 ul/ml NP-40) and

incubated on ice for 10 minutes. Lysates were centrifuged at 12,000 x g for 10 minutes at 4°C to

fractionate nuclei and debris (pellet) from cytoplasm (supernatant). After total RNA quantification

of cytoplasmic extracts on a Nanodrop Spectrophotometer (Thermo Fisher), a volume

corresponding to 50 ug of total RNA was transferred onto a 10-50% linear sucrose density

gradient per sample. Gradients were centrifuged at 35,000 rpm in a SW41 rotor for 2-3 hours at

4°C. Following centrifugation, gradients were fractionated into 12, 1 ml fractions using a

BioComp Piston Gradient Fractionator and A$_{260nm}$ profiles were recorded using a BioComp Triax

Flow Cell.

RNA extraction of polysome fractions

From each fraction, 400 ul was used for the following procedure. To each fraction, 2 ng

Luciferase Control RNA (Promega) was added as a normalization spike-in control, followed by

600 ul of Trizol and purification following standard procedures (Thermo Fisher). RNA pellets

were rinsed with ice-cold 70% ethanol, briefly dried, and resuspended in 20 ul nuclease-free

H$_2$O. For each fraction, 8 ul of RNA was reverse transcribed using random primers and

Superscript IV reverse transcriptase following the manufacturer's instructions (Thermo Fisher).

CRISPR-Cas9 mutagenesis of *ddx6* and *ddx61*

CRISPR RNA (crRNA) guide sequences were designed using Integrated DNA Technologies

(IDT) custom design tool

(https://www.idtdna.com/site/order/designtool/index/CRISPR_CUSTOM). crRNA sequences with

the highest on-target scores with no predicted off-target sites were chosen. The impact of

targeted mutations on gene function was maximized by targeting regions within exons encoding

the DEAD box helicase domain (see Fig. S5A-B). Reagents for CRISPR-Cas9 mutagenesis

were prepared as previously described for efficient mutagenesis (Klatt Shaw & Mokalled, 2021).

For double knockdown experiments, crRNAs targeting *ddx6* and *ddx61* (25 uM each, 50 uM

total) and trans-activating CRISPR RNA (tracrRNA) (IDT) (50 uM) were combined equally and

annealed to generate dual guide RNA duplexes (dgRNA), followed by 1:1 dilution in duplex

buffer (IDT) for a final concentration of 25 uM dgRNA. Annealed dgRNAs were next combined

equally with 25 uM Cas9-NLS (QB3 MacroLab, UC Berkeley) that had been diluted from a 40

uM stock solution using Cas9 dilution buffer B (IDT). Final injection solutions contained 0.1%

phenol red dye, 0.2 M potassium chloride, and 5 uM each of dgRNA and Cas9-NLS. Injection

solutions were incubated for 5 minutes at 37▯C to facilitate ribonucleoprotein complex (RNP)

formation. AB wild-type embryos were injected at the 1-cell stage with 2 nl of injection solution.

crRNA sequences are listed in Table S6. For single knockdown experiments, the same method

described for double knockdown experiments was employed except that twice the sgRNA dose

as double knockdown experiments was used to ensure that the total molar amount of gRNA

injected was equivalent between experiments (i.e 2.5 uM each of *ddx61* + *ddx6* gRNAs or 5 uM

ddx61 sgRNA or 5 uM *ddx6* gRNA).

Acknowledgements

We thank the Ohio State Zebrafish Facilities staff for excellent care, the Ohio Supercomputer

Center for computational resources, the Genomics Shared Resource at the Ohio State

Comprehensive Cancer Center for RNA analysis and DNA sequencing, and the Ohio State

Center for RNA Biology for support and resources. We thank Susan Cole, Kenneth Poss, Mira

Pronobis, and Guramrit Singh for advice and critical comments. We thank Anton Blatnik,

Geremy Lerma, Lauren Levesque, Benjamin Pastore, and Zhongxia Yi for advice.

Competing interests

The authors declare no competing or financial interests.

Funding

This work was supported by NIH grants R01GM117964 (S.L.A.), R35GM158157 (S.L.A.), and

R35GM146924 (M.G.K.), an OSU Dean's Enrichment Fellowship (M.C.B.), a Pelotonia

Fellowship (M.C.B.), an OSU Center for RNA Biology Undergraduate Fellowship (K.G.T.), a

Cellular, Molecular, and Biochemical Sciences T32 Training fellowship T32GM141955 (C.C.A.),

and NSF grant DBI-1560182 (A.M.).

Data and resource availability

RNA-Seq datasets and additional data files are available at GEO under accession GSE319255.

All additional data and resources are included in the article and supplementary information.

Author contributions

T.L.G., M.C.B., C.C.A, K.G.T, D.M.P, A.M., and S.L.A. performed experiments. T.L.G., M.C.B.,

R.D., M.G.K., and S.L.A. analyzed data. All authors contributed intellectually and discussed the

data and manuscript. T.L.G. wrote the manuscript and co-authors participated in editing.

References

Andrews, S. (2010). FastQC - A quality control tool for high throughput sequence data.

http://www.bioinformatics.babraham.ac.uk/projects/fastqc/. *Babraham Bioinformatics*.

Aulehla, A., Wiegraebe, W., Baubet, V., Wahl, M. B., Deng, C., Taketo, M., Lewandoski, M.

and Pourquié, O. (2008). A beta-catenin gradient links the clock and wavefront systems in

mouse embryo segmentation. *Nat Cell Biol* **10**, 186–193.

Arvola, R. M., Chang, C.-T., Buytendorp, J. P., Levdansky, Y., Valkov, E., Freddolino, L., &

Goldstrohm, A. C. (2020). Unique repression domains of Pumilio utilize deadenylation and

decapping factors to accelerate destruction of target mRNAs. *Nucleic Acids Research* **48**,

1843–1871.

Baia Amaral, D., Egidy, R., Perera, A. and Bazzini, A. A. (2024). miR-430 regulates zygotic

mRNA during zebrafish embryogenesis. *Genome Biol* **25**, 74.

Bailey, T. L. (2021). STREME: accurate and versatile sequence motif discovery. *Bioinformatics*

37, 2834–2840.

Bazzini, A. A., Lee, M. T. and Giraldez, A. J. (2012). Ribosome profiling shows that miR-430

reduces translation before causing mRNA decay in zebrafish. *Science* **336**, 233–237.

Bembom, O. and Ivanek, R. (2024). seqLogo: Sequence logos for DNA sequence alignments.

Benjamini, Y., Krieger, A. M. and Yekutieli, D. (2006). Adaptive linear step-up procedures that

control the false discovery rate. *Biometrika* **93**, 491–507.

Blatnik, M. C., Gallagher, T. L. and Amacher, S. L. (2023). Keeping development on time:

Insights into post-transcriptional mechanisms driving oscillatory gene expression during

vertebrate segmentation. *Wiley Interdiscip Rev RNA* **14**, e1751–e1751.

Bohn, J. A., Van Etten, J. L., Schagat, T. L., Bowman, B. M., McEachin, R. C., Freddolino,

L. and Goldstrohm, A. C. (2018). Identification of diverse target RNAs that are functionally

regulated by human Pumilio proteins. *Nucleic Acids Res* **46**, 362–386.

Bonev, B., Stanley, P. and Papalopulu, N. (2012). MicroRNA-9 Modulates Hes1 ultradian

oscillations by forming a double-negative feedback loop. *Cell Rep* **2**, 10–18.

Bonferroni, C. E. (1935). Il Calcolo delle Assicurazioni su Gruppi di Teste. In *Studi in Onore del*

Professore Salvatore Ortu Carboni.

Bonferroni, C. E. (1936). Teoria statistica delle classi e calcolo delle probabilità. *Pubblicazioni*
del R Istituto Superiore di Scienze Economiche e Commerciali di Firenze **8**, 3–62.

Bray, N. L., Pimentel, H., Melsted, P. and Pachter, L. (2016). Near-optimal probabilistic RNA-
seq quantification. *Nat Biotechnol* **34**, 525–527.

Briata, P., Forcales, S. V., Ponassi, M., Corte, G., Chen, C.-Y., Karin, M., Puri, P. L. and
Gherzi, R. (2005). p38-Dependent Phosphorylation of the mRNA Decay-Promoting Factor
KSRP Controls the Stability of Select Myogenic Transcripts. *Mol Cell* **20**, 891–903.

Causier, B., Li, Z., De Smet, R., Lloyd, J. P. B., Van de Peer, Y. and Davies, B. (2017).
Conservation of Nonsense-Mediated mRNA Decay Complex Components Throughout
Eukaryotic Evolution. *Sci Rep* **7**, 16692.

Cho, H., Kim, K. M. and Kim, Y. K. (2009). Human Proline-Rich Nuclear Receptor Coregulatory
Protein 2 Mediates an Interaction between mRNA Surveillance Machinery and Decapping
Complex. *Mol Cell* **33**, 75–86.

Cho, H., Kim, K. M., Han, S., Choe, J., Park, S. G., Choi, S. S. and Kim, Y. K. (2012).
Staufen1-Mediated mRNA Decay Functions in Adipogenesis. *Mol Cell* **46**, 495–506.

Cho, H., Han, S., Choe, J., Park, S. G., Choi, S. S. and Kim, Y. K. (2013). SMG5-PNRC2 is
functionally dominant compared with SMG5-SMG7 in mammalian nonsense-mediated
mRNA decay. *Nucleic Acids Res* **41**, 1319–1328.

Cho, H., Park, O. H., Park, J., Ryu, I., Kim, J., Ko, J. and Kim, Y. K. (2015). Glucocorticoid
receptor interacts with PNRC2 in a ligand-dependent manner to recruit UPF1 for rapid
mRNA degradation. *Proc Natl Acad Sci U S A* **112**, E1540–E1549.

Crippa, S., Nemir, M., Ounzain, S., Ibberson, M., Berthonneche, C., Sarre, A., Boisset, G.,
Maison, D., Harshman, K., Xenarios, I., Diviani, D., Schorderet D., Pedrazzini, T. (2016).
Comparative transcriptome profiling of the injured zebrafish and mouse hearts identifies
miRNA-dependent repair pathways. *Cardiovasc Res* **110**, 73–84.

Cutty, S. J., Fior, R., Henriques, P. M., Saúde, L. and Wardle, F. C. (2012). Identification and
expression analysis of two novel members of the Mesp family in zebrafish. *Int J Dev Biol* **56**,
285–294.

Decker, C. J., Teixeira, D. and Parker, R. (2007). Edc3p and a glutamine/asparagine-rich
domain of Lsm4p function in processing body assembly in Saccharomyces cerevisiae. *J Cell*
Biol **179**, 437–449.

Delaune, E. A., François, P., Shih, N. P. and Amacher, S. L. (2012). Single-cell-resolution
imaging of the impact of Notch signaling and mitosis on segmentation clock dynamics. *Dev*
Cell **23**, 995–1005.

Dill, K. K. and Amacher, S. L. (2005). tortuga refines Notch pathway gene expression in the
zebrafish presomitic mesoderm at the post-transcriptional level. *Dev Biol* **287**, 225–236.

Ditlev, J. A., Case, L. B. and Rosen, M. K. (2018). Who's In and Who's Out-Compositional
Control of Biomolecular Condensates. *J Mol Biol* **430**, 4666–4684.

**Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., & Gingeras, T. R.** (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics **29**, 15–21.

Dunn, O. J. (1961). Multiple Comparisons among Means. *J Am Stat Assoc* **56**, 52–64.

Eulalio, A., Behm-Ansmant, I., Schweizer, D. and Izaurralde, E. (2007). P-body formation is a
consequence, not the cause, of RNA-mediated gene silencing. *Mol Cell Biol* **27**, 3970–3981.

Farrell, J. A., Wang, Y., Riesenfeld, S. J., Shekhar, K., Regev, A., & Schier, A. F. (2018).
Single-cell reconstruction of developmental trajectories during zebrafish embryogenesis.
Science **360**, 6392.

**Fu, Y.-F., Du, T.-T., Dong, M., Zhu, K.-Y., Jing, C.-B., Zhang, Y., Wang, L., Fan, H.-B., Chen,
Y., Jin, Y., Yue, G.-P., Chen, S.-J., Chen, Z., Huang, Q.-H., Jing, Q., Deng, M., & Xi Liu, T.**
(2009). Mir-144 selectively regulates embryonic α-hemoglobin synthesis during primitive
erythropoiesis. *Blood* **113**, 1340–1349.

**Fujino, Y., Yamada, K., Sugaya, C., Ooka, Y., Ovara, H., Ban, H., Akama, K., Otosaka, S.,
Kinoshita, H., Yamasu, K., et al.,** (2018). Deadenylation by the CCR4-NOT complex
contributes to the turnover of *hairy*‐related mRNAs in the zebrafish segmentation clock.
FEBS Lett **592**, 3388–3398.

Galgano, A., Forrer, M., Jaskiewicz, L., Kanitz, A., Zavolan, M. and Gerber, A. P. (2008).
Comparative analysis of mRNA targets for human PUF-family proteins suggests extensive
interaction with the miRNA regulatory system. *PLoS One* **3**, e3164–e3164.

**Gallagher, T. L., Tietz, K. T., Morrow, Z. T., McCammon, J. M., Goldrich, M. L., Derr, N. L.
and Amacher, S. L.** (2017). Pnrc2 regulates 3′UTR-mediated decay of segmentation clock-
associated transcripts during zebrafish segmentation. *Dev Biol* **429**, 225–239.

**Gangras, P., Gallagher, T. L., Parthun, M. A., Yi, Z., Patton, R. D., Tietz, K. T., Deans, N. C.,
Bundschuh, R., Amacher, S. L. and Singh, G.** (2020). Zebrafish rbm8a and magoh
mutants reveal EJC developmental functions and new 3′UTR intron-containing NMD targets.
PLoS Genet **16**, e1008830–e1008830.

Garcia, J. F. and Parker, R. (2015). MS2 coat proteins bound to yeast mRNAs block 5' to 3′
degradation and trap mRNA decay products: implications for the localization of mRNAs by
MS2-MCP system. *RNA* **21**, 1393–1395.

Garneau, N. L., Wilusz, J. and Wilusz, C. J. (2007). The highways and byways of mRNA
decay. *Nat Rev Mol Cell Biol* **8**, 113–126.

Gentleman, R. C., Carey, V. J., Bates, D. M., Bolstad, B., Dettling, M., Dudoit, S., Ellis, B.,
Gautier, L., Ge, Y., Gentry, J., et al., (2004). Bioconductor: open software development for
computational biology and bioinformatics. *Genome Biol* **5**, R80–R80.

Gerber, A. P., Luschnig, S., Krasnow, M. A., Brown, P. O. and Herschlag, D. (2006).
Genome-wide identification of mRNAs associated with the translational regulator PUMILIO
in Drosophila melanogaster. *Proc Natl Acad Sci U S A* **103**, 4487–4492.

Gillhouse, M., Wagner Nyholm, M., Hikasa, H., Sokol, S. Y. and Grinblat, Y. (2004). Two
Frodo/Dapper homologs are expressed in the developing brain and mesoderm of zebrafish.
Developmental Dynamics **230**, 403–409.

Giraldez, A. J., Cinalli, R. M., Glasner, M. E., Enright, A. J., Thomson, J. M., Baskerville, S.,
Hammond, S. M., Bartel, D. P. and Schier, A. F. (2005). MicroRNAs Regulate Brain
Morphogenesis in Zebrafish. *Science (1979)* **308**, 833–838.

Giraldez, A. J., Mishima, Y., Rihel, J., Grocock, R. J., Van Dongen, S., Inoue, K., Enright, A.
J. and Schier, A. F. (2006). Zebrafish MiR-430 Promotes Deadenylation and Clearance of
Maternal mRNAs. *Science (1979)* **312**, 75–79.

Gomez, C., Özbudak, E. M., Wunderlich, J., Baumann, D., Lewis, J. and Pourquié, O.
(2008). Control of segment number in vertebrate embryos. *Nature* **454**, 335–339.

Graber, J. H., Cantor, C. R., Mohr, S. C., & Smith, T. F. (1999). In silico detection of control
signals: mRNA 3'-end-processing sequences in diverse species. *Proceedings of the*
National Academy of Sciences **96**, 14055–14060.

Gratacós, F. M. and Brewer, G. (2010). The role of AUF1 in regulated mRNA decay. *Wiley*
Interdiscip Rev RNA **1**, 457–473.

Hafner, M., Landthaler, M., Burger, L., Khorshid, M., Hausser, J., Berninger, P., Rothballer,
A., Ascano Jr, M., Jungkamp, A.-C., Munschauer, M., et al., (2010). Transcriptome-wide
identification of RNA-binding protein and microRNA target sites by PAR-CLIP. *Cell* **141**, 129–
141.

Hatakeyama, J., Bessho, Y., Katoh, K., Ookawara, S., Fujioka, M., Guillemot, F. and
Kageyama, R. (2004). *Hes*genes regulate size, shape and histogenesis of the nervous
system by control of the timing of neural stem cell differentiation. *Development* **131**, 5539–
5550.

He, F. and Jacobson, A. (2015). Nonsense-Mediated mRNA Decay: Degradation of Defective
Transcripts Is Only Part of the Story. *Annu Rev Genet* **49**, 339–366.

He, F. and Jacobson, A. (2023). Eukaryotic mRNA decapping factors: molecular mechanisms
and activity. *FEBS J* **290**, 5057–5085.

Heinrich, S., Sidler, C. L., Azzalin, C. M. and Weis, K. (2017). Stem-loop RNA labeling can
affect nuclear and cytoplasmic mRNA processing. *RNA* **23**, 134–141.

Holley, S. A., Geisler, R. and Nüsslein-Volhard, C. (2000). Control of *her1* expression during
zebrafish somitogenesis by a *Delta*-dependent oscillator and an independent wave-front
activity. *Genes Dev* **14**, 1678–1690.

Hoshijima, K., Jurynec, M. J., Klatt Shaw, D., Jacobi, A. M., Behlke, M. A. and Grunwald,
D. J. (2019). Highly Efficient CRISPR-Cas9-Based Methods for Generating Deletion
Mutations and F0 Embryos that Lack Gene Function in Zebrafish. *Dev Cell* **51**, 645-657.e4.

Hu, Y., Xie, S. and Yao, J. (2016). Identification of Novel Reference Genes Suitable for qRT-
PCR Normalization with Respect to the Zebrafish Developmental Stage. *PLoS One* **11**,
e0149277–e0149277.

Hubaud, A. and Pourquié, O. (2014). Signalling dynamics in vertebrate segmentation. *Nat Rev*
Mol Cell Biol **15**, 709–721.

Huber, W., Carey, V. J., Gentleman, R., Anders, S., Carlson, M., Carvalho, B. S., Bravo, H.
C., Davis, S., Gatto, L., Girke, T., et al., (2015). Orchestrating high-throughput genomic
analysis with Bioconductor. *Nat Methods* **12**, 115–121.

Hurt, J. A., Robertson, A. D., & Burge, C. B. (2013). Global analyses of UPF1 binding and
function reveal expanded scope of nonsense-mediated mRNA decay. *Genome Research*
23, 1636–1650.

Icli, B., Wara, A. K. M., Moslehi, J., Sun, X., Plovie, E., Cahill, M., Marchini, J. F., Schissler,
A., Padera, R. F., Shi, J., Cheng, H., Raghuram, S., Arany, Z., Liao, R., Croce, K.,
MacRae, C., Feinberg, M. (2013). MicroRNA-26a regulates pathological and physiological
angiogenesis by targeting BMP/SMAD1 signaling. *Circ Res* **113**, 1231–1241.

Icli, B., Dorbala, P. and Feinberg, M. W. (2014). An emerging role for the miR-26 family in
cardiovascular disease. *Trends Cardiovasc Med* **24**, 241–248.

Ishibashi, M., Ang, S. L., Shiota, K., Nakanishi, S., Kageyama, R. and Guillemot, F. (1995).
Targeted disruption of mammalian hairy and Enhancer of split homolog-1 (HES-1) leads to
up-regulation of neural helix-loop-helix factors, premature neurogenesis, and severe neural
tube defects. *Genes Dev* **9**, 3136–3148.

Johnstone, T. G., Bazzini, A. A., & Giraldez, A. J. (2016). Upstream ORFs are prevalent
translational repressors in vertebrates. *The EMBO Journal* **35**, 706–723.

Junker, J. P., Noël, E. S., Guryev, V., Peterson, K. A., Shah, G., Huisken, J., McMahon, A.
P., Berezikov, E., Bakkers, J. and van Oudenaarden, A. (2014). Genome-wide RNA
Tomography in the Zebrafish Embryo. *Cell* **159**, 662–675.

Karousis, E. D., Nasif, S. and Mühlemann, O. (2016). Nonsense-mediated mRNA decay:
novel mechanistic insights and biological impact. *Wiley Interdiscip Rev RNA* **7**, 661–682.

Kervestin, S. and Jacobson, A. (2012). NMD: a multifaceted response to premature
translational termination. *Nat Rev Mol Cell Biol* **13**, 700–712.

Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. and Schilling, T. F. (1995). Stages
of embryonic development of the zebrafish. *Developmental Dynamics* **203**, 253–310.

Klatt Shaw, D. and Mokalled, M. H. (2021). Efficient CRISPR/Cas9 mutagenesis for
neurobehavioral screening in adult zebrafish. *G3 (Bethesda)* **11**, jkab089.

Klett, H., Jürgensen, L., Most, P., Busch, M., Günther, F., Dobreva, G., Leuschner, F.,
Hassel, D., Busch, H. and Boerries, M. (2018). Delineating the Dynamic Transcriptome
Response of mRNA and microRNA during Zebrafish Heart Regeneration. *Biomolecules* **9**,
11.

Komisarczuk, A. Z., Topp, S., Stigloher, C., Kapsimali, M., Bally-Cuif, L. and Becker, T. S.
(2008). Enhancer detection and developmental expression of zebrafish *sprouty1*, a member
of the *fgf8* synexpression group. *Developmental Dynamics* **237**, 2594–2603.

Kretov, D. A., Folkes, L., Mora-Martin, A., Walawalkar, I. A., Imrat, Syedah, N., Vanuytsel,
K., Moxon, S., Murphy, G. J., & Cifuentes, D. (2024). The miR-144/Hmgn2 regulatory axis
orchestrates chromatin organization during erythropoiesis. *Nature Communications* **15**,
3821.

Krol, A. J., Roellig, D., Dequéant, M.-L., Tassy, O., Glynn, E., Hattem, G., Mushegian, A.,
Oates, A. C. and Pourquié, O. (2011). Evolutionary plasticity of segmentation clock
networks. *Development* **138**, 2783–2792.

Kusov, Y. Y., Shatirishvili, G., Dzagurov, G. and Gauss-Müller, V. (2001). A new G-tailing
method for the determination of the poly(A) tail length applied to hepatitis A virus RNA.
Nucleic Acids Res **29**, E57-e57.

Lai, T., Cho, H., Liu, Z., Bowler, M. W., Piao, S., Parker, R., Kim, Y. K. and Song, H. (2012).
Structural Basis of the PNRC2-Mediated Link between mRNA Surveillance and Decapping.
Structure **20**, 2025–2037.

Lewis, J. (2003). Autoinhibition with Transcriptional Delay. *Current Biology* **13**, 1398–1408.

Liu, Y., Zhu, Z., Ho, I. H. T., Shi, Y., Li, J., Wang, X., Chan, M. T. V and Cheng, C. H. K.
(2020). Genetic Deletion of miR-430 Disrupts Maternal-Zygotic Transition and Embryonic
Body Plan. *Front Genet* **11**, 853.

Lu, G. and Hall, T. M. T. (2011). Alternate modes of cognate RNA recognition by human
PUMILIO proteins. *Structure* **19**, 361–367.

Luo, Y., Na, Z. and Slavoff, S. A. (2018). P-Bodies: Composition, Properties, and Functions.
Biochemistry **57**, 2424–2431.

Lykke-Andersen, J. and Bennett, E. J. (2014). Protecting the proteome: Eukaryotic
cotranslational quality control pathways. *J Cell Biol* **204**, 467–476.

Lykke-Andersen, J. and Wagner, E. (2005). Recruitment and activation of mRNA decay
enzymes by two ARE-mediated decay activation domains in the proteins TTP and BRF-1.
Genes Dev **19**, 351–361.

Lykke-Andersen, S., & Jensen, T. H. (2015). Nonsense-mediated mRNA decay: an intricate
machinery that shapes transcriptomes. Nature Reviews Molecular Cell Biology 16, 665–677.

Makino, S., Mishima, Y., Inoue, K. and Inada, T. (2015). Roles of mRNA fate modulators Dhh1
and Pat1 in TNRC6-dependent gene silencing recapitulated in yeast. *J Biol Chem* **290**,
8331–8347.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet J* **17**, 10.

Martin, G. and Keller, W. (1998). Tailing and 3′-end labeling of RNA with yeast poly(A)
polymerase and various nucleotides. *RNA* **4**, 226-230.

Masamizu, Y., Ohtsuka, T., Takashima, Y., Nagahara, H., Takenaka, Y., Yoshikawa, K.,
Okamura, H. and Kageyama, R. (2006). Real-time imaging of the somite segmentation
clock: revelation of unstable oscillators in the individual presomitic mesoderm cells. *Proc*
Natl Acad Sci U S A **103**, 1313–1318.

Matsuda, M., Hayashi, H., Garcia-Ojalvo, J., Yoshioka-Kobayashi, K., Kageyama, R.,
Yamanaka, Y., Ikeya, M., Toguchida, J., Alev, C. and Ebisuya, M. (2020). Species-specific
segmentation clock periods are due to differential biochemical reaction speeds. *Science*
(1979) **369**, 1450–1455.

Mendell, J. T., Sharifi, N. A., Meyers, J. L., Martinez-Murillo, F., & Dietz, H. C. (2004).
Nonsense surveillance regulates expression of diverse classes of mammalian transcripts
and mutes genomic noise. *Nature Genetics* **36**, 1073–1078.

Mi, H., Muruganujan, A. and Thomas, P. D. (2013). PANTHER in 2013: modeling the evolution
of gene function, and other gene attributes, in the context of phylogenetic trees. *Nucleic*
Acids Res **41**, D377–D386.

Mishima, Y. and Tomari, Y. (2016). Codon Usage and 3′ UTR Length Determine Maternal
mRNA Stability in Zebrafish. *Mol Cell* **61**, 874–885.

Mishima, Y. and Tomari, Y. (2017). Pervasive yet nonuniform contributions of Dcp2 and Cnot7
to maternal mRNA clearance in zebrafish. *Genes to Cells* **22**, 670–678.

Moreno, T. A., Jappelli, R., Izpisúa Belmonte, J. C. and Kintner, C. (2008). Retinoic acid
regulation of the Mesp-Ripply feedback loop during vertebrate segmental patterning. *Dev*
Biol **315**, 317–330.

Morris, A. R., Mukherjee, N. and Keene, J. D. (2008). Ribonomic analysis of human Pum1
reveals cis-trans conservation across species despite evolution of diverse mRNA target sets.
Mol Cell Biol **28**, 4093–4103.

Mugridge, J. S., Ziemniak, M., Jemielity, J. and Gross, J. D. (2016). Structural basis of
mRNA-cap recognition by Dcp1-Dcp2. *Nat Struct Mol Biol* **23**, 987–994.

Murata, Y. and Wharton, R. P. (1995). Binding of pumilio to maternal hunchback mRNA is
required for posterior patterning in drosophila embryos. *Cell* **80**, 747–756.

Nguyen-Chi, M. E., Bryson-Richardson, R., Sonntag, C., Hall, T. E., Gibson, A., Sztal, T.,
Chua, W., Schilling, T. F. and Currie, P. D. (2012). Morphogenesis and cell fate
determination within the adaxial cell equivalence group of the zebrafish myotome. *PLoS*
Genet **8**, e1003014–e1003014.

Nicholson, P., Gkratsou, A., Josi, C., Colombo, M. and Mühlemann, O. (2018). Dissecting
the functions of SMG5, SMG7, and PNRC2 in nonsense-mediated mRNA decay of human
cells. *RNA* **24**, 557–573.

Oates, A. C., Morelli, L. G. and Ares, S. (2012). Patterning embryos with oscillations: structure,
function and dynamics of the vertebrate segmentation clock. *Development* **139**, 625–639.

Ohtsuka, T., Ishibashi, M., Gradwohl, G., Nakanishi, S., Guillemot, F. and Kageyama, R.
(1999). Hes1 and Hes5 as notch effectors in mammalian neuronal differentiation. *EMBO J*
18, 2196–2207.

Ostareck, D. H., Naarmann-de Vries, I. S. and Ostareck-Lederer, A. (2014). DDX6 and its
orthologs as modulators of cellular and viral RNA expression. *WIREs RNA* **5**, 659–678.

Palmeirim, I., Henrique, D., Ish-Horowicz, D. and Pourquié, O. (1997). Avian hairy Gene
Expression Identifies a Molecular Clock Linked to Vertebrate Segmentation and
Somitogenesis. *Cell* **91**, 639–648.

Pimentel, H., Bray, N. L., Puente, S., Melsted, P. and Pachter, L. (2017). Differential analysis
of RNA-seq incorporating quantification uncertainty. *Nat Methods* **14**, 687–690.

Pourquié, O. (2011). Vertebrate segmentation: from cyclic gene networks to scoliosis. *Cell* **145**,
650–663.

Presnyak, V. and Coller, J. (2013). The DHH1/RCKp54 family of helicases: an ancient family of
proteins that promote translational silencing. *Biochim Biophys Acta* **1829**, 817–823.

Pronobis, M. I., DeLuca, S., Lee, S., Garcia, L., Ou, J., Bursac, N., & Poss, K. D. (2025).
Ddx61-enriched condensates refine heart regeneration programs. *Nature Communications*
16, 8992.

Rabani, M., Pieper, L., Chew, G.-L. and Schier, A. F. (2017). A Massively Parallel Reporter
Assay of 3′ UTR Sequences Identifies In Vivo Rules for mRNA Degradation. *Mol Cell* **68**,
1083-1094.e5.

Raghavan, A., Ogilvie, R. L., Reilly, C., Abelson, M. L., Raghavan, S., Vasdewani, J.,
Krathwohl, M. and Bohjanen, P. R. (2002). Genome-wide analysis of mRNA decay in
resting and activated primary human T lymphocytes. *Nucleic Acids Res* **30**, 5529–5538.

Ramírez, F., Ryan, D. P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A. S., Heyne, S.,
Dündar, F., & Manke, T. (2016). deepTools2: a next generation web server for deep-
sequencing data analysis. *Nucleic Acids Research* **44**, W160–W165.

Rattenbacher, B., Beisang, D., Wiesner, D. L., Jeschke, J. C., von Hohenberg, M., St
Louis-Vlasova, I. A. and Bohjanen, P. R. (2010). Analysis of CUGBP1 targets identifies
GU-repeat sequences that mediate rapid mRNA decay. *Mol Cell Biol* **30**, 3970–3980.

Rodriguez, J. M., Pozo, F., Cerdán-Vélez, D., Di Domenico, T., Vázquez, J., & Tress, M. L.
(2022). APPRIS: selecting functionally important isoforms. *Nucleic Acids Research* **50**, D54–
D59.

Salkind, N. (2007). *Encyclopedia of Measurement and Statistics*. Sage Publications, Inc.

Sanduja, S., Blanco, F. F. and Dixon, D. A. (2011). The roles of TTP and BRF proteins in
regulated mRNA decay. *Wiley Interdiscip Rev RNA* **2**, 42–57.

Schneider, T. D. and Stephens, R. M. (1990). Sequence logos: a new way to display
consensus sequences. *Nucleic Acids Res* **18**, 6097–6100.

Schneider, T. D., Stormo, G. D., Gold, L. and Ehrenfeucht, A. (1986). Information content of
binding sites on nucleotide sequences. *J Mol Biol* **188**, 415–431.

Schoenberg, D. R. and Maquat, L. E. (2012). Regulation of cytoplasmic mRNA decay. *Nat Rev*
Genet **13**, 246–259.

Schröter, C. and Oates, A. C. (2010). Segment Number and Axial Identity in a Segmentation
Clock Period Mutant. *Current Biology* **20**, 1254–1258.

Schweingruber, C., Rufener, S. C., Zünd, D., Yamashita, A., & Mühlemann, O. (2013).
Nonsense-mediated mRNA decay — Mechanisms of substrate mRNA recognition and
degradation in mammalian cells. Biochimica et Biophysica Acta (BBA) - Gene Regulatory
Mechanisms 1829, 612–623.

Sheth, U. and Parker, R. (2003). Decapping and decay of messenger RNA occur in
cytoplasmic processing bodies. *Science* **300**, 805–808.

Sidák, Z. (1967). Rectangular Confidence Regions for the Means of Multivariate Normal
Distributions. *J Am Stat Assoc* **62**, 626–633.

Simsek, M. F., Chandel, A. S., Saparov, D., Zinani, O. Q. H., Clason, N. and Özbudak, E. M.
(2023). Periodic inhibition of Erk activity drives sequential somite segmentation. *Nature* **613**,
153–159.

Singh, P. P. and Isambert, H. (2020). OHNOLOGS v2: a comprehensive resource for the
genes retained from whole genome duplication in vertebrates. *Nucleic Acids Res* **48**, D724–
D730.

Soto, X., Biga, V., Kursawe, J., Lea, R., Doostdar, P., Thomas, R. and Papalopulu, N.
(2020). Dynamic properties of noise and Her6 levels are optimized by miR-9, allowing the
decoding of the Her6 oscillator. *EMBO J* **39**, e103558–e103558.

Stoecklin, G., Mayo, T. and Anderson, P. (2006). ARE-mRNA degradation requires the 5'-3'
decay pathway. *EMBO Rep* **7**, 72–77.

Su, Z., Si, W., Li, L., Zhou, B., Li, X., Xu, Y., Xu, C., Jia, H., & Wang, Q. K. (2014). MiR-144
regulates hematopoiesis and vascular development by targeting meis1 during zebrafish
development. *The International Journal of Biochemistry & Cell Biology* **49**, 53–63. **Superina,**
S., Borovina, A. and Ciruna, B. (2014). Analysis of maternal-zygotic ugdh mutants reveals
divergent roles for HSPGs in vertebrate embryogenesis and provides new insight into the
initiation of left–right asymmetry. *Dev Biol* **387**, 154–166.

Sur, A., Wang, Y., Capar, P., Margolin, G., Prochaska, M. K., & Farrell, J. A. (2023). Single-
cell analysis of shared signatures and transcriptional diversity during zebrafish development.
Developmental Cell **58**, 3028-3047.

Tan, S., Ohtsuka, T., González, A. and Kageyama, R. (2012). MicroRNA9 regulates neural
stem cell differentiation by controlling Hes1 expression dynamics in the developing brain.
Genes to Cells **17**, 952–961.

Teixeira, D., Sheth, U., Valencia-Sanchez, M. A., Brengues, M. and Parker, R. (2005).

Processing bodies require RNA for assembly and contain nontranslating mRNAs. *RNA* **11**,

371–382.

Thermes, V., Grabher, C., Ristoratore, F., Bourrat, F., Choulika, A., Wittbrodt, J. and Joly,

J.-S. (2002). I-SceI meganuclease mediates highly efficient transgenesis in fish. *Mech Dev*

118, 91–98.

Thisse, B. and Thisse, C. (2005). High Throughput Expression Analysis of ZF-Models

Consortium Clones. *ZFIN Direct Data Submission (http://zfin.org).*

Thisse, B., Pflumio, S., Furthauer, M., Loppin, B., Heyer, V., Degrave, A., Woehl, R., Lux,

A., Steffan, T., Charbonnier, X. Q., et al., (2001). Expression of the zebrafish genome

during embryogenesis. *ZFIN Direct Data Submission (http://zfin.org).*

Thomas, P. D., Ebert, D., Muruganujan, A., Mushayahama, T., Albou, L.-P. and Mi, H.

(2022). PANTHER: Making genome-scale phylogenetics accessible to all. *Protein Sci* **31**, 8–

22.

Tietz, K. T., Gallagher, T. L., Mannings, M. C., Morrow, Z. T., Derr, N. L. and Amacher, S. L.

(2020). Pumilio response and AU-rich elements drive rapid decay of Pnrc2-regulated cyclic

gene transcripts. *Dev Biol* **462**, 129–140.

Tutucci, E., Vera, M., Biswas, J., Garcia, J., Parker, R. and Singer, R. H. (2018). An improved

MS2 system for accurate reporting of the mRNA life cycle. *Nat Methods* **15**, 81–89.

Vejnar, C. E., Abdel Messih, M., Takacs, C. M., Yartseva, V., Oikonomou, P., Christiano, R.,

Stoeckius, M., Lau, S., Lee, M. T., Beaudoin, J.-D., et al., (2019). Genome wide analysis

of 3′ UTR sequence elements and proteins regulating mRNA stability during maternal-to-

zygotic transition in zebrafish. *Genome Res* **29**, 1100–1114.

Wang, X., McLachlan, J., Zamore, P. D. and Hall, T. M. T. (2002). Modular Recognition of RNA

by a Human Pumilio-Homology Domain. *Cell* **110**, 501–512.

Wanglar, C., Takahashi, J., Yabe, T. and Takada, S. (2014). Tbx protein level critical for clock-

mediated somite positioning is regulated through interaction between Tbx and Ripply. *PLoS*

One **9**, e107928–e107928.

Watterston, C., Zeng, L., Onabadejo, A. and Childs, S. J. (2019). MicroRNA26 attenuates

vascular smooth muscle maturation via endothelial BMP signalling. *PLoS Genet* **15**,

e1008163–e1008163.

Waxman, J. S. (2005). Regulation of the early expression patterns of the zebrafish Dishevelled-

interacting proteins Dapper1 and Dapper2. *Developmental Dynamics* **233**, 194–200.

Westerfield, M. (2007). The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish (Danio
rerio), 5th Edition. *University of Oregon Press, Eugene (Book)*.

White, E. K., Moore-Jarrett, T. and Ruley, H. E. (2001). PUM2, a novel murine Puf protein,
and its consensus RNA-binding site. *RNA* **7**, 1855-1866.

White, R. J., Collins, J. E., Sealy, I. M., Wali, N., Dooley, C. M., Digby, Z., Stemple, D. L.,
Murphy, D. N., Billis, K., Hourlier, T., et al., (2017). A high-resolution mRNA expression
time course of embryonic development in zebrafish. *Elife* **6**, e30860.

Yabe, T., Hoshijima, K., Yamamoto, T. and Takada, S. (2016). Quadruple zebrafish mutant
reveals different roles of Mesp genes in somite segmentation between mouse and zebrafish.
Development **143**, 2842–2852.

Yoshioka-Kobayashi, K., Matsumiya, M., Niino, Y., Isomura, A., Kori, H., Miyawaki, A. and
Kageyama, R. (2020). Coupling delay controls synchronized oscillation in the segmentation
clock. *Nature* **580**, 119–123.

Zamore, P. D., Williamson, J. R. and Lehmann, R. (1997). The Pumilio protein binds RNA
through a conserved domain that defines a new class of RNA-binding proteins. *RNA* **3**,
1421-1433.

Zamore, P. D., Bartel, D. P., Lehmann, R. and Williamson, J. R. (1998). The PUMILIO–RNA
Interaction: A Single RNA-Binding Domain Monomer Recognizes a Bipartite Target
Sequence. *Biochemistry* **38**, 596–604.

Zampedri, C., Tinoco-Cuellar, M., Carrillo-Rosas, S., Diaz-Tellez, A., Ramos-Balderas, J.
L., Pelegri, F. and Maldonado, E. (2016). Zebrafish P54 RNA helicases are cytoplasmic
granule residents that are required for development and stress resilience. *Biol Open* **5**,
1473–1484.

Zheng, D., Ezzeddine, N., Chen, C.-Y. A., Zhu, W., He, X. and Shyu, A.-B. (2008).
Deadenylation is prerequisite for P-body formation and mRNA decay in mammalian cells. *J*
Cell Biol **182**, 89–101.

1199

Figure legends

Fig. 1. Overexpressed transcripts in *MZpnrc2* mutants have a longer 3′UTR length
distribution than unaffected transcripts, are associated with many biological processes,
and are enriched for destabilizing motifs. (A) Volcano plot showing transcripts that are
overexpressed (q < 0.05, n = 1,769) (blue), under-expressed (i.e. transcripts with reduced
expression; q < 0.05, n = 856) (red) or not significantly differentially expressed (q > 0.05, n =
32,533) (black) in *MZpnrc2* mutant versus WT embryos at mid-segmentation stages. Key

segmentation-associated transcripts (Fig. 2) are indicated on the volcano plot. (B) Violin plots of
3′UTR lengths for overexpressed (q < 0.05, n = 1,333) (blue), under-expressed (q < 0.05, n =
685) (red), and unaffected (log2FC -0.05 to 0.05, n = 1,352) (gray) transcripts. (C) Metagene
plot showing normalized RNA-Seq read coverage in WT (brown) and *MZpnrc2* mutant (blue)
samples for transcripts that are overexpressed in *MZpnrc2* mutants (q < 0.05) and read
coverage in WT (gray) and *MZpnrc2* mutant (black) samples for transcripts that are non-
overexpressed in *MZpnrc2* mutants (i.e. all transcripts that were not significantly
overexpressed). (D-F) Enriched motifs in (D) overexpressed (n = 1320), (E) unaffected (n =
1334), and (F) under-expressed (n = 690) transcript 3′UTRs (E-value < 0.05) identified using
STREME. (G) Significantly enriched terms within the aspect biological process using GO
analysis of overexpressed genes in *MZpnrc2* mutants. Among the top 30 enriched terms (see
Fig. S1C), 15 terms are shown that contain gene sets ≥5 and ListHit values ≥4. FC = fold
change; kb = kilobase; TSS = transcription start site; TES = transcription end site; FDR = false
discovery rate; ListHit = number of overexpressed genes per GO term.
1221
Fig. 2. Expression patterns of transcripts associated with segmentation or other
developmental processes are disrupted in *MZpnrc2* mutants. (A-U) WT and *MZpnrc2*
mutants were raised to mid-segmentation stages (14-18 hpf) and probed for *rhov* (A, A' vs B, B',
RT-qPCR in C), *sp5l* (D, D' vs E, E', RT-qPCR in F), *mespab* (G, G' vs H, H', RT-qPCR in I),
ripply2 (J, J' vs K, K', RT-qPCR in L), *dact1* (M, M' vs N, N', RT-qPCR in O), *spry1* (P, P' vs Q,
Q', RT-qPCR in R), and *spry4* (S, S' vs T, T', RT-qPCR in U). Coloration reactions for each
probe were developed in parallel for identical reaction times. Representative images from each
genotype are shown. In situs were performed twice independently with consistent results (n =
15-20 embryos per probe per genotype for each experiment). RT-qPCR was performed
separately from in situ experiments using whole embryos (n = 15 embryos per bio-replicate per
genotype). Fold change values are the mean of three bio-replicate values derived from the
average of two technical replicates. P-values were calculated using an unpaired t-test. In situs
are shown in lateral view (A-T), magnified in dorsal view to the right of each embryo (A'-T').
Scale bars = 100 um (A-T); dorsal view = 50 um (A'-T'). hpf = hours post fertilization.
1236
Fig. 3. Overexpressed oscillatory gene transcripts and *ddx61* transcripts have shortened
poly(A) tails in *MZpnrc2* mutants. (A) Poly(A)-tail (pA-tail) lengths of transcripts from WT and
MZpnrc2 mutants at mid-segmentation stages were measured by G/I tailing-based RT-PCR
(Martin & Keller, 1998; Kusov et al., 2001), revealing a predominant pA-tail length of ~45 nts for

her1 and ~60 nts for *her7* and *rhov* in both WT and *MZpnrc2* mutants. A broad tail distribution

pattern is observed for *dlc* in WT embryos that becomes reduced in *MZpnrc2* mutants to a

predominant tail length of ~100 nts. In contrast, *ddx61* pA-tail lengths are broadly distributed in

both WT and *MZpnrc2* mutants, though the distribution is shifted to smaller tail lengths in

mutants. (B) Lane profile analysis reveals that the *MZpnrc2* mutant pA-tail length distribution is

reduced relative to WT for all transcripts profiled. Predominant peaks are demarcated with

hatched lines (see methods), with the relative proportion of tail products in hatched regions

relative to the entire plot (beginning at position A_0) indicated. All pA-tail length results were

consistent across two bio-replicates for each genotype. (C) Transcript quantification from WT

and *MZpnrc2* mutant extracts at mid-segmentation stages. Total RNA (n = 50 embryos per

extract per bio-replicate) was separated into pA-enriched and pA-depleted extracts by oligo-dT

bead-based purification (see methods). Fold change values are presented as the mean of 3 bio-

replicates with each bio-replicate value derived from the average of 2 technical replicates. Error

bars show standard deviation and p-values show results from unpaired t tests. hpf = hours post

fertilization; pA = poly(A) tail PCR; S = gene-specific PCR.

1256

Fig. 4. Accumulated oscillatory gene transcripts are primarily disengaged from

ribosomes, whereas accumulated *ddx61* transcripts are associated with ribosomes and

lead to higher Ddx6/Ddx61 protein levels in *MZpnrc2* mutants. (A) Representative 260 nm

absorbance profiles for WT and *MZpnrc2* mutant cytoplasmic extracts fractionated after

cycloheximide treatment and sucrose density gradient centrifugation (n = 300 embryos per

genotype at mid-segmentation stages) (see methods). Regions between dotted lines indicate

fractions containing mRNAs co-sedimenting in the following groups: ribosome/subunit-unbound

mRNAs (mRNP), 40S small ribosomal subunit (40S), 60S large ribosomal subunit (60S),

monosome-bound mRNAs (80S), and polysome-bound mRNAs (polysome). (B-F) Transcript

quantification from WT and *MZpnrc2* mutant extracts at mid-segmentation stages from

unbound/subunit (combined mRNP, 40S, and 60S fractions), monosome, and polysome

fractions. Fold change values are presented as the mean of 3 bio-replicates (n = 300 embryos

per genotype per bio-replicate) with each bio-replicate value derived from the average of 3

technical replicates. Error bars show standard deviation and q-values show adjusted p-values

from unpaired t tests adjusted with a two-stage linear step-up Benjamini, Krieger, and Yekutieli

false discovery rate (FDR) procedure (Benjamini et al., 2006). (G) Immunoblot showing

Ddx6/Ddx61 protein levels in extracts from WT and *MZpnrc2* mutants with β-actin as a loading

control (1 embryo equivalent loaded per lane at mid-segmentation stages). (H) Bar graph

showing relative Ddx6/Ddx61 protein levels in WT and *MZpnrc2* mutant extracts calculated

using pixel intensities from lane profile analysis of immunoblot in G. Signal intensities

normalized to β-actin. Fold change value is presented as the mean of 3 bio-replicates. Error bar

shows standard deviation and p-value was calculated from an unpaired t test. hpf = hours post

fertilization; nm = nanometers; mRNP = messenger ribonucleoprotein particle; S (panel A) =

Svedberg unit.

1281

Fig. 5. Induced expression of a dominant-negative deadenylase has no effect in WT

embryos but induces segmentation defects in *MZpnrc2* mutants. WT and *MZpnrc2* mutants

were injected with 20 pgs *hsp70l:Myc-cnot7DN* plasmid at the 1-cell stage and raised alongside

uninjected clutch-mate controls to ~14 hpf. Injected and uninjected controls were then heat

shocked at 37⬚C for 30 minutes and raised at standard temperature (28.5⬚C) for 6 hours post-

heat shock (pHS) with morphological assessment at hourly intervals (see methods). (A-D)

Representative embryos are shown for (A) uninjected WT embryos (n = 36), (B) *hsp70l:Myc-*

cnot7DN-injected WT embryos (n = 33), (C) uninjected *MZpnrc2* mutants (n = 62), and (D)

hsp70:Myc-cnot7DN-injected *MZpnrc2* mutants (n = 58). Arrow in D indicates an abnormal

segment. (E) Bar graph shows the proportion of embryos with normal and/or abnormal

segments, indicating that injected *MZpnrc2* mutants are the only group shown in A-D with

somite defects. (F-G) A subset of injected embryos was reserved as no heat-shock controls and

were negative for Myc-Cnot7DN expression by anti-Myc immunohistochemistry (n = 15) (F),

whereas mosaic expression was observed in injected embryos 6 hours pHS (n = 189) (G)

(representative images shown in F-G). P-value was calculated using a Fisher's Exact test

comparing the number of embryos with or without segment defects. Plasmid injection followed

by heat-shock induction was performed twice independently with consistent results. hpf = hours

post fertilization; pHS = post-heat shock; DN = dominant negative; scale bar = 300 um.

1300

Fig. 6. *MZpnrc2* mutant embryos with depleted Ddx6 and Ddx61 function first show

abnormal phenotypes at late segmentation stages. (A-H) Representative embryos are

shown at 16 hpf and 19 hpf, respectively, for (A, C) uninjected WT and (B, D) uninjected

MZpnrc2 mutants, (E, G) *ddx6;ddx61* crispants (on a WT background), (F, H) *MZpnrc2*

ddx6:*ddx61* crispants. (I-J) Bar graphs show the proportion of embryos with normal and

abnormal phenotypes at 16 hpf (I) and 19 hpf (J). Adjusted p-values were calculated using a

Fisher's Exact test comparing the number of embryos within each phenotype class (normal,

mildly abnormal, strongly abnormal) and corrected for multiple pairwise comparisons using a

Bonferroni adjustment (Bonferroni, 1935; Bonferroni, 1936; Dunn, 1961). Four knockdown

experiments were independently performed and produced consistent results. S = somite stage;

scale bar = 250 um.

1312

Fig. 7. *MZpnrc2* mutant larvae are sensitized to depletion of Ddx6 and Ddx61. (A-D)

Representative images are shown at 84 hpf for (A) uninjected WT, (B) uninjected *MZpnrc2*, (C)

ddx6;*ddx61* crispants (on a WT background), and (D) *MZpnrc2 ddx6*:*ddx61* crispants. (E) By

112 hpf, *ddx6;ddx61* crispants exhibit mild to severe abnormalities that occur at higher

frequency in *MZpnrc2* mutants (representative phenotypic classes shown in E with phenotypic

proportions in Fig. S6L). (F) Bar graph showing proportion of larvae at 120 hpf with normal and

abnormal phenotypes, indicating a synergistic effect from loss of *pnrc2*, *ddx6*, and *ddx61*.

Proportions were calculated from the following groups: uninjected WT, n = 112; *ddx6;ddx61*

crispant, n = 100; uninjected *MZpnrc2*, n = 81; *MZpnrc2 ddx6;ddx61* crispant, n = 47. Adjusted

p-values were calculated using a Fisher's Exact test comparing the number of embryos within

each phenotype class (normal, mildly abnormal, strongly abnormal) and corrected for multiple

pairwise comparisons using a Bonferroni adjustment (Bonferroni, 1935; Bonferroni, 1936; Dunn,

1961). Four independent knockdown experiments were performed and produced consistent

results. (C) Immunoblot showing increased Ddx6/Ddx61 protein levels in *MZpnrc2* mutants and

efficient depletion of Ddx61/Ddx6 protein in *ddx6;ddx61* crispants. One-half larval equivalents

were loaded in each lane with β-actin shown as a loading control. Immunoblot results were

consistent across two independent experiments. hpf = hours post-fertilization; dpf = days post-

fertilization; scale bars = 250 um.

1331

Fig. 8. Injection of *ddx6;ddx61* CRISPRs does not affect *her1* mRNA levels in WT

embryos, but increases *her1* accumulation in *MZpnrc2* mutants. WT and *MZpnrc2* mutants

were injected with *ddx6;ddx61* CRISPRs and raised to 18 hpf alongside uninjected clutch-mates

as controls. Uninjected and injected embryos were probed for *her1* expression by in situ

hybridization (n = 20 embryos per genotype per condition) and oscillatory gene transcript levels

were quantified by qRT-PCR (n = 10 embryos per genotype per condition; 5 bio-replicates per

condition with bio-replicate values derived from the average of 2 technical replicates each). (A-

D') Although *her1* expression in *ddx6;ddx61* crispants is indistinguishable from uninjected

clutch-mates (A, A' vs B, B'), *her1* expression appears expanded in *MZpnrc2 ddx6;ddx61*

crispants relative to uninjected *MZpnrc2* mutant clutch-mates (C, C' vs D, D'). Lateral views are

shown in A-D, magnified dorsal views are shown in A'-D'. (E-H) Bar graphs showing RT-qPCR

results with fold change values derived from the mean of 5 bio-replicates. Out of four quantified

oscillatory gene transcripts, only *her1* expression is significantly increased in *MZpnrc2*

ddx6;ddx61 crispants (E vs F-H). In contrast, *her7* expression is significantly reduced in

MZpnrc2 ddx6;ddx61 crispants compared to uninjected *MZpnrc2* mutants (F). Oscillatory gene

transcript expression is unaffected in *ddx6;ddx61* crispants on the WT background (E-H).

Adjusted p-values calculated using a one-way ANOVA with a Dunn-Šídák correction for multiple

comparisons (Šídák,1967; Abdi, 2007). Scale bars = 100 um (A-D); dorsal view = 50 um (A'-D').

Fig. 1

Fig. 2

WT   MZpnrc2

Fig. 3

A Tail Length PCR

B Tail Length Pixel Intensity Profiles

C Reverse Transcription Quantitative PCR

Fig. 4

Fig. 5

Uninjected control   *hsp70l:Myc-cnot7DN*

6 hours post-heat shock

A WT   36/36   B WT   33/33

C *MZpnrc2*   62/62   D *MZpnrc2*   33/58

E Segment Morphology 6 hrs pHS

p < 0.0001

% embryos

WT   *MZpnrc2*   WT   *MZpnrc2*

Uninjected   Injected

■ Normal segments
■ 1-2 abnormal segments

F anti-Myc

No heat shock control   15/15

G anti-Myc

6 hours post-heat shock   189/189

Fig. 6

Fig. 7

WT

MZpnrc2

Uninjected

A 84 hpf 90/104

B 80/107

ddx6;ddx61 crispant

C 39/106

D 85/100

Representative phenotypes in WT and MZpnrc2 mutant larvae

E **Phenotypic Classes**

Normal

112 hpf

Mildly abnormal

Strongly abnormal

F **120 hpf Phenotypes**

p-adj < 0.0004

p-adj < 0.0004

p-adj < 0.0004

p-adj < 0.0004

■ Normal
■ Mildly abnormal
■ Strongly abnormal

% embryos

Uninjected | ddx6;ddx61 crispant | Uninjected | ddx6;ddx61 crispant

WT | MZpnrc2

G **Immunoblot of 120 hpf extracts**

WT | MZpnrc2

Ddx6/Ddx61 — 54 kD

β-actin — 42 kD

ddx6;ddx61 CRISPRs: − + − +

Fig. 8

Fig. S1 (goes with Fig. 1)

C

Overexpressed - Biological Process

Top 30 Enriched GO Terms

D

Overexpressed - Molecular Function

Top 30 Enriched GO Terms

E

Overexpressed - Cellular Component

F

Under-expressed - Biological Process

G

Under-expressed - Molecular Function

H

Under-expressed - Cellular Component

Fig. S1 (goes with Fig. 1). (A-B) Oscillatory gene transcript 3′UTR lengths are similar in WT and *MZpnrc2* mutants. Normalized bedgraphs (read coverage density plots) from high throughput sequencing reads (see methods in main text) of WT and *MZpnrc2* mutants at mid-segmentation stages (n = 50 embryos for each genotype per biological replicate). Read coverage is individually scaled in A to better show read density distribution across 3′UTRs and downstream regions for each bio-replicate, whereas read coverage is singly scaled across bio-replicates in B to better show differences in read density that correspond with transcript accumulation in *MZpnrc2* mutants. (C-E) Overexpressed transcripts in *MZpnrc2* mutants at mid-segmentation stages are enriched for GO terms associated with developmental pathways and post-transcriptional mRNA regulation. Top 30 most-enriched terms within the aspects biological process (C), molecular function (D), and cellular component (E) using GO enrichment analysis of overexpressed genes in *MZpnrc2* mutant embryos. (F-H) Under-expressed transcripts in *MZpnrc2* mutants at mid-segmentation stages are enriched for GO terms associated with translation. Top 30 most-enriched terms within the aspect biological process (F), all enriched terms within the aspect molecular function (G), and top 30 most-enriched terms within the aspect cellular component (H) using GO enrichment analysis of under-expressed genes in *MZpnrc2* mutant embryos. FDR = false discovery rate; ListHit = number of overexpressed genes per GO term.

Fig. S2 (goes with Fig. 2)

Fig. S2 (goes with Fig. 2). Many genes known to oscillate in the zebrafish presomitic mesoderm are Pnrc2-regulated. (A-F) Bar graphs showing RT-qPCR validation of RNA-Seq results. Quantification was performed using cDNA synthesized from total RNA extracts from WT and *MZpnrc2* mutants at mid-segmentation stages (n = 15 embryos per biological replicate per genotype with 2-3 technical replicates per biological replicate). Fold change values are the mean of biological replicate values derived from the average of technical replicates. P-values were calculated using an unpaired t-test. Among the overexpressed oscillatory gene transcripts shown, 4 were overexpressed in RNA-Seq analysis with false discovery rates of q < 0.05, whereas 2 validated transcripts had larger q-values: *angptl7* (q = 0.12) and *rtn1a* (q = 0.13).

Fig. S3 (goes with Fig. 3)

Fig. S3 (goes with Fig. 3). (A-E) Accumulated oscillatory gene transcripts as well as *ddx61* transcripts have shortened poly(A) tails in *MZpnrc2* mutants, but are not completely deadenylated. Larger images of gels shown in Fig. 3 are shown in A-E, including molecular weight ladders for *her1* (A), *her7* (B), *dlc* (C), *rhov* (D), and *ddx61* (E) (note that for *dlc* and *ddx61*, gel images have been cropped and stitched so that the ladder is adjacent to profiled lanes for better size determination; stitched regions are denoted with a red line in C and E). Regions of interest for pixel intensity profiling are boxed in green, with blue lines denoting the width of each lane analyzed in pixel intensity plots shown below each gel. Predominant peaks are denoted with blue dots on each gel. Because *dlc* and *ddx61* lack a predominant peak in WT samples, a region corresponding to the peak in *MZpnrc2* mutants was identified by molecular weight and denoted with a blue dot in WT lanes. Gel images were captured under exposure times that were not saturated. Lane profiles of pixel intensities are scaled universally between lanes to better show differences in levels between WT and *MZpnrc2* mutants, although it is important to note that pA-tail reaction products are derived from end-point PCR where precise quantification is not possible. (F-I) Bar graphs showing transcript quantification using cDNA synthesized with oligo(dT) primers and total RNA extracts from WT and *MZpnrc2* mutants at mid-segmentation stages (n = 15 embryos per biological replicate, 2 technical replicates per biological replicate). Fold change values are the mean of biological replicate values derived from the average of two technical replicates. (J-K) *edc3* and *lsm14aa* transcripts that encode P-body associated factors are overexpressed in *MZpnrc2* mutants. Bar graphs showing transcript quantification using cDNA synthesized with random primers and total RNA extracts from WT and *MZpnrc2* mutants at mid-segmentation stages (n = 15 embryos per biological replicate per genotype, with 2 technical replicates per biological replicate). Fold change values are the mean of biological replicate values derived from the average of technical replicates. P-values were calculated using an unpaired t-test.

Fig. S4 (goes with Fig. 5)

Fig. S4 (goes with Fig. 5). (A-H) Segmentation defects arise in a dose-dependent manner in WT embryos after injection of mRNA encoding a dominant negative deadenylase. WT embryos were injected with *Myc-cnot7DN* mRNA at the 1-cell stage and raised to mid-segmentation stages. Representative images are shown for embryos injected with *Myc-cnot7DN* mRNA with normal segments (A), irregular segments (B), and no segments (C). (D) Bar graph shows the proportion of embryos with normal, irregular, or no segments (n = 108 uninjected embryos; n = 30 embryos injected with 200 pg *Myc-cnot7DN* mRNA; n = 58 embryos injected with 400 pg *Myc-cnot7DN* mRNA). (E-G) A subset of uninjected and injected embryos shown in A-D were probed for *her1* expression by in situ hybridization. Representative embryos are shown for uninjected (E) and *Myc-cnot7DN* mRNA-injected embryos (F-G). (H) Bar graph shows the proportion of embryos with normal and abnormal *her1* expression patterns. Proportions were calculated from a single experiment and were consistent with a second independent experiment (not shown). (I-N) Induction of Myc-Cnot7DN expression after injection of a heat-shock inducible construct encoding a Myc-tagged dominant negative deadenylase. WT and *MZpnrc2* mutant embryos were scored for Myc protein expression after performing anti-Myc immunohistochemistry on plasmid-injected embryos (see methods in main text). Individuals were scored as having low (i.e. substantial mosaicism), moderate, high, or very high (i.e. little to no mosaicism) Myc expression on the basis of anti-Myc staining pattern. Representative embryos are shown for embryos in each expression class (I-L). (M) Bar graph shows the proportion of embryos in each expression class comparing WT (n = 33) and *MZpnrc2* mutants (n = 58), showing that there is no difference in Myc-Cnot7DN expression when comparing WT and *MZpnrc2* mutant embryos (p = 0.913). (N) Bar graph shows the proportion of embryos in each expression class comparing *MZpnrc2* mutants with normal (n = 25) and abnormal (n = 33) segments, showing that all *MZpnrc2* mutants with very high Myc-Cnot7DN expression (i.e. embryos with broad expression with little to no mosaicism) have abnormal segments, though overall, expression differences between *MZpnrc2* mutants with and without somite defects did not reach significance (p = 0.056). Proportions were calculated by combining data from two independent experiments that showed consistent results. P-values were calculated using a Fisher's Exact test comparing the number of embryos within each phenotype class. Adjusted p-values that correct for multiple pairwise comparisons were calculated using a Bonferroni adjustment (Bonferroni, 1935; Bonferroni, 1936; Dunn, 1961). hpf = hours post fertilization; DN = dominant negative; scale bars = 200 um (A-C & I-L), 50 um (E-G).

Fig. S5 (goes with Figs. 6-7)

Fig. S5 (goes with Figs 6-7). CRISPRs targeting *ddx6* and *ddx61* efficiently reduce *ddx6* and *ddx61* mRNA and Ddx6/Ddx61 protein levels. (A) Cartoon schematics (exported from Ensembl) depict *ddx6* and *ddx61* gene structure with targeted regions denoted by red arrows, both of which lie just upstream of the DEAD-box helicase domain. (B-C) High resolution melt analysis (HRMA) difference plots showing CRISPR-induced mutagenesis at *ddx6* (B) and *ddx61* (C) targeted loci. CRISPR-injected embryo samples (blue) show strong deflections from uninjected embryos samples (red), indicating mutagenesis at the desired target loci. The presence of two distinct clusters (i.e. melt behavior patterns) in uninjected controls for *ddx61* (C) is due to a polymorphism outside of the targeted region within *ddx61* intron 3. (D-E) Bar graphs showing RT-qPCR results from 18 hpf embryos with fold change values derived from the mean of 5 biological replicates (biological replicate values derived from the average of 2 technical replicates each). Both *ddx6* and *ddx61* transcripts are significantly overexpressed in *MZpnrc2* embryos (D and E, blue vs orange bars), validating RNASeq results. Injection of *ddx6* and *ddx61* CRISPRs significantly decreases *ddx6* and *ddx61* transcript levels in both WT and *MZpnrc2* mutants at mid-segmentation (D and E, blue vs red bars and orange vs purple bars). Adjusted p-values calculated using a one-way ANOVA with a Dunn-Šídák correction for multiple comparisons (Šídák,1967; Abdi, 2007). (F) Immunoblot showing Ddx6/Ddx61 protein levels in WT uninjected and *ddx6*;*ddx61* crispant embryos over a 5 day period with β-actin as a loading control. (G) Full immunoblot image from Fig. 7G with molecular weight ladder indicating the expected sizes for Ddx6/Ddx61 and β-actin protein. Immunoblot results were consistent across two independent experiments. kD = kilodaltons.

Fig. S6 (goes with Fig. 7)

Fig. S6 (goes with Fig. 7). Abnormal phenotypes from depletion of Ddx6 and Ddx61 worsens over time in sensitized *MZpnrc2* mutants whereas WT embryos develop normally up to 60 hpf. (A-H) Representative embryos from the predominant class are shown at 42 hpf and 60 hpf, respectively, for (A, E) uninjected WT embryos and (B, F) uninjected *MZpnrc2* embryos, (C, G) *ddx6*;*ddx61* crispants (on the WT background), (D, H) *MZpnrc2 ddx6*:*ddx61* crispants. (I-L) Bar graphs show the proportion of embryos with normal and abnormal phenotypes at 42 hpf (I), 60 hpf (J), 84 hpf (K) and 112 hpf (L), indicating that *MZpnrc2 ddx6*;*ddx61* crispants are sensitized to Ddx6/Ddx61 depletion. Adjusted p-values were calculated using a Fisher's Exact test comparing the number of embryos within each phenotype class (normal, mildly abnormal, strongly abnormal) and corrected for multiple pairwise comparisons using a Bonferroni adjustment (Bonferroni, 1935; Bonferroni, 1936; Dunn, 1961). Representative larvae from predominant phenotypic classes at 84 hpf and 112 hpf are shown in Fig. 7A-E. hpf = hours post-fertilization; scale bars = 250 um.

Supplemental Methods

DNA extraction and high resolution melt analysis

Individual embryos were lysed in 50 ul 1X ThermoPol Buffer (NEB) at 95°C for 10 minutes, digested at 55°C for 1 hour using 50 ug Proteinase K (Thermo Fisher), followed by Proteinase K inactivation at 95°C for 10 minutes. CRISPR mutagenesis efficiency was analyzed by high resolution melt analysis (HRMA) on a CFX Duet instrument (Bio-Rad) using 1 ul of DNA extract as template in a 20 ul reaction with Precision Melt Mix following standard manufacturer procedures (Bio-Rad). gRNA and primer sequences are listed in Table S6. Melt analysis was performed using Precision Melt Analysis Software v1.2 (Bio-Rad).

RNA extraction and cDNA synthesis

Whole embryos at mid-segmentation stages (n = 15 embryos per biological replicate for each genotype in Figs S2-S3) were solubilized in Trizol for RNA extraction and purified following standard procedures (Thermo Fisher). All genotypes and conditions were performed in biological triplicate. 500 ng of total RNA was reverse transcribed using Superscript IV reverse transcriptase and random primers (Fig. S2) or oligo(dT) primers (Fig. S3) and according to the manufacturer's instructions (Thermo Fisher).

Quantitative RT-PCR analysis

Quantitative RT-PCR was performed using PowerUp SYBR Green Master Mix (Thermo Fisher) and 4.5 ul cDNA (diluted 1:50 in Fig. S2, diluted 1:25 in Fig. S3) in 20 ul reactions, following manufacturer's procedures. Negative controls lacking template were included for each primer set. All reactions were subjected to thermal melting to confirm that each reaction gave single peaks. Transcript levels were normalized to *mobk13* (*mob4*) (Hu et al., 2016; Gangras et al., 2020; Tietz et al., 2020). Cycle thresholds (Ct) were determined using Bio-Rad CFX manager software. The average Ct of 2 technical replicates per sample was used for calculations. Changes in mRNA expression were calculated by $\Delta\Delta Ct = \Delta Ct$ target $- \Delta Ct$ control. Relative changes in mRNA expression levels are represented graphically as fold change, where relative mRNA fold change = $2-\Delta\Delta Ct$. All graphs and statistics were generated using Prism 10 (GraphPad).

mRNA injection

The *Myc-cnot7DN* coding sequence from plasmid *pMA-Myc-cnot7DN* (see main methods) was PCR amplified and cloned into expression vector pCS2+ (Rupp et al., 1994; Turner and

Weintraub, 1994) using standard digestion-based cloning with the enzymes EcoRI-HF and XbaI (NEB) to generate the plasmid *pCS2-Myc-cnot7DN*. For deadenylation inhibition experiments, *Myc-cnot7DN* mRNA was synthesized using the SP6 mMessage Machine Kit (Thermo Fisher), diluted in 0.2M KCl with 0.1% phenol red, and injected into 1-cell stage embryos (400 pg mRNA per embryo). Primer sequences are listed in Table S6.

Microscopy and imaging

Live embryos were mounted in 3% methylcellulose and imaged at 63x magnification using AxioVision Software (Zeiss) on a MZFLIII Fluorescence Stereo Microscope (Leica) with an AxioCam HRc digital camera (Zeiss).

Protein extraction and Immunoblot analysis

See methods in main text.

Supplemental References

Rupp, R.A., Snider, L., Weintraub, H. (1994). Xenopus embryos regulate the nuclear localization of XMyoD. *Genes Dev* **8**, 1311–1323.

Turner, D. L. and Weintraub, H. (1994). Expression of achaete-scute homolog 3 in Xenopus embryos converts ectodermal cells to a neural fate. *Genes Dev* **8**, 1434–1447.