Title

Pnrc2 promotes rapid mRNA decay and coordinately supports early development with P-body

factors Ddx6 and Ddx61

Running title

Pnrc2, P-bodies, and development

Authors

Gallagher, Thomas L.[1,2,‡]; Blatnik, Monica C.[1,2,3,*]; Austin, Clare C.[1,2]; Thompson, Kathryn G.[1,2];

Pvirre, Danielle M.[1]; Morgan, Angelina[1]; Kearse, Michael G.[2,4]; Amacher, Sharon L.[1,2,3,5‡]

1 Department of Molecular Genetics, The Ohio State University, Columbus, OH, 43210, USA

2 Center for RNA Biology, The Ohio State University, Columbus, OH, 43210, USA

3 Interdisciplinary Graduate Program in Molecular, Cellular and Developmental Biology, The

Ohio State University, Columbus, OH, 43210, USA

4 Department of Biological Chemistry and Pharmacology, The Ohio State University, Columbus,

OH, 43210, USA

5 Center for Muscle Health and Neuromuscular Disorders, The Ohio State University,

Columbus, OH, 43210, USA.

*Present address: Oberlin College of Arts and Sciences, 38 E. College St., Oberlin, OH, 44074,

USA.

‡Authors for correspondence (gallagher.184@osu.edu and amacher.6@osu.edu)

Keywords

somitogenesis, oscillatory gene expression, P-bodies, translational regulation, deadenylation

Abstract

Somitogenesis, the sequential segmentation of vertebrate embryonic mesoderm, is

controlled by the segmentation clock, a molecular oscillator that controls periodic gene

expression in unsegmented mesoderm and is regulated by a negative feedback loop driven by

Hes/Her transcriptional repressors. In zebrafish, Pnrc2 is required for decay of *her1* transcript

and additional oscillatory gene transcripts. Despite accumulation of numerous mRNAs including

those encoding key developmental regulators, overt embryonic phenotypes are absent in *pnrc2*

mutants. Our previous work suggested that accumulated mRNAs are not translated in *pnrc2*

mutants, though the underlying mechanism(s) was unknown. We show here that many

overexpressed transcripts in *pnrc2* mutants have shortened poly(A) tails and are disengaged

from ribosomes, and that deadenylation inhibition leads to somite defects in *pnrc2* mutants. In

contrast, transcripts encoding the P-body protein, Ddx61, are both overexpressed and engaged

with ribosomes, leading to an increase in Ddx61 protein. Co-depletion of Ddx61 and its ohnolog

Ddx6 enhances *her1* accumulation and later leads to strong morphological defects in *pnrc2*

mutants. Together, our results show that multiple post-transcriptional mechanisms ensure

proper translation when mRNA decay is inhibited.

Summary statement

Pnrc2 promotes decay of tail-shortened, ribosome-disengaged mRNAs and co-regulates *her1*

mRNA decay with Ddx6 and Ddx61, P-body factors that sustain normal development in decay-

deficient *pnrc2* mutants.

Introduction

In vertebrates, somitogenesis is a fundamental developmental process (Kageyama et

al., 2012) characterized by the anterior to posterior sequential segmentation of the embryonic

mesoderm into blocks of tissue called somites, which later give rise to axial skeletal muscle,

vertebrae, and dermis (Christ et al., 1978; Christ et al, 1986; Keynes and Stern 1988). This

rhythmic and regular process is controlled by a molecular oscillator called the segmentation

clock which operates in the unsegmented presomitic mesoderm (PSM) and is characterized by

rapid cycles of mRNA and protein expression (Hubaud and Pourquie, 2014; Oates et al., 2012;

Pourquie, 2011). The segmentation clock is regulated by a self-sustaining negative feedback

loop, where a core oscillatory factor, a transcriptional repressor, inhibits expression of

downstream oscillatory genes, including itself. In all vertebrates examined to date,

transcriptional repressors of the *Hes*/*her* gene family are considered the conserved pacemakers

of the segmentation clock (Bessho et al., 2001; Oates and Ho 2002; Palmeirim et al., 1997).

For proper somite formation, rates of each oscillatory step, from transcription to

translation to decay, must be regulated to ensure correct somite size and number (Gomez et al.,

2008; Holley et al., 2000; Keynes and Stern 1988; Lewis, 2003; Matsuda et al., 2020; Palmeirim

et al., 1997; Schroter and Oates, 2010). Real-time in vivo segmentation clock reporters

designed to recapitulate clock dynamics must not only contain critical transcriptional regulatory

regions that drive oscillatory expression but must also contain features, typically 3'UTR

sequences and protein motifs, that destabilize reporter mRNAs and proteins, respectively

(Aulehla et al., 2008; Delaune et al., 2012; Masamizu et al., 2006; Simsek et al., 2023;

Yoshioka-Kobayashi et al., 2020).

We have focused on post-transcriptional regulation of the segmentation clock (reviewed

in Blatnik et al., 2023) which was originally motivated by our discovery that the zebrafish *pnrc2*

gene promotes rapid decay of oscillatory gene transcripts (Dill and Amacher, 2005; Gallagher et

al., 2017; Tietz et al., 2020). Pnrc2 (Proline-rich nuclear receptor coactivator 2) is related to the

EDC family of mRNA decapping enhancers and contains a large intrinsically disordered domain

and domains that facilitate interaction with the Dcp1a mRNA decapping factor and the Upf1

mRNA decay factor (Cho et al., 2009; Lai et al., 2012). Although Pnrc2 has been characterized

as a mRNA decay adaptor protein in cell culture, there is no evidence that it directly binds

mRNA. Transcriptional oscillations are normal in zebrafish *pnrc2* mutants, but *her1* mRNA (and

other oscillatory gene transcripts) accumulate due to mRNA decay defects (Dill and Amacher,

2005; Gallagher et al., 2017; Tietz et al., 2020). Surprisingly, despite a 4-6-fold increase in *her1*

and *dlc* oscillatory gene transcripts in *pnrc2* mutants, oscillatory gene protein is expressed

normally (Gallagher et al., 2017) and there is no overt segmentation phenotype. Discordance

between mRNA and protein levels might be caused by a combination of factors: transcripts

might be inefficiently translated due to deadenylation and/or binding of translational repressors,

as well as localized to cytoplasmic ribonucleoprotein granules like P-bodies that are sites of

non-translating deadenylated mRNAs (reviewed in: Luo et al., 2018; Ostareck et al., 2014). We

explore these possibilities using a combination of deep sequencing, molecular, and in vivo

approaches.

To better understand the role of *pnrc2* during segmentation, we used RNA-Seq analysis

and identified over 1700 transcripts that were significantly overexpressed at least 1.2-fold in

MZpnrc2 mutants. GO analysis reveals enrichment for many biological processes, including

terms associated with mRNA processing and decay. Motif analysis shows that overexpressed

transcripts are enriched for well-known RNA binding protein (RBP) and microRNA (miRNA)

target sites. Accumulated transcripts of several oscillatory genes have shortened poly(A) tails,

suggesting that they are inefficiently translated. Supporting this interpretation, polysome profiling

shows that several oscillatory gene transcripts that are overexpressed in *MZpnrc2* mutants have

normal levels of ribosome engagement. *MZpnrc2* mutants are sensitive to inhibition of

deadenylation and to depletion of the P-body factors Ddx6 and Ddx61, suggesting that

accumulated mRNAs in *pnrc2* mutants are not translated when deadenylation and P-body

functions are intact. Overall, our results show that discordant expression of oscillatory gene

mRNA and protein in *MZpnrc2* mutants is likely due to inefficient translation of accumulated

transcripts. In the absence of Pnrc2 function, the P-body factors Ddx6 and Ddx61 buffer against

mRNA decay dysfunction and support embryonic development, implicating a broader role for

Pnrc2 and P-bodies during development across many tissue types.

Results:

Pnrc2-regulated transcripts are associated with key biological processes and are enriched for

destabilizing motifs

During somitogenesis, *pnrc2* promotes decay of oscillatory gene transcripts (Gallagher

et al., 2017; Tietz et al., 2020). To globally identify transcripts that require *pnrc2* for proper

expression, we performed RNA-Seq analysis on wild-type and maternally-zygotic *pnrc2*

(*MZpnrc2*) mutant embryos at mid-segmentation stages. We detected 2,256 significantly

differentially expressed transcripts with a false discovery rate of q < 0.05 (Fig. 1A, Tables S1-S3)

using the program Kallisto to quantify transcript abundance (Bray et al., 2016) and Sleuth for

differential expression analysis (Pimentel et al., 2017). Most differentially expressed transcripts

were overexpressed (n = 1,769 transcripts), as expected for a mRNA decay-deficient mutant.

Among overexpressed transcripts, >99% are annotated as protein-coding by Ensembl's biotype

annotation (n = 1759 transcripts), suggesting that Pnrc2 regulates decay of transcripts with full

open reading frames rather than transcripts containing premature termination codons that are

subject to nonsense-mediated decay (NMD).

To explore characteristics of overexpressed transcripts in *MZpnrc2* mutants, we

conducted GO enrichment analysis (Thomas et al., 2022). GO analysis identified many

significantly enriched GO terms in the "biological process" aspect (Fig. 1B, Tables S4-S5)

including somite specification and neural development, processes that rely upon *Hes/her*

oscillations for proper differentiation (Bonev et al., 2012; Hatakeyama et al., 2004; Ishibashi et

al., 1995; Ohtsuka et al., 1999; Soto et al., 2020; Tan et al., 2012). Additional enriched terms in

the aspects biological process and molecular function are related to developmental pathways

involving Fgf, Wnt, Notch, and Ephrin signaling (Figs 1, S1, Table S5). Terms associated with

post-transcriptional regulation are also enriched including 3'UTR-mediated destabilization,

deadenylation-dependent decay, and P-body assembly (Fig. 1B, Table S5). Overexpressed

genes in these categories may represent transcripts that depend on Pnrc2 for their decay or

transcripts that are expressed to compensate for loss of Pnrc2 function.

Our previous work showed that the Pnrc2-regulated *her1* transcript contains two critical

destabilizing motifs in the 3'UTR, a Pumilio Response Element (PRE) and AU-rich Element

(ARE) (Tietz et al., 2020). To determine whether 3'UTRs of overexpressed transcripts identified

by RNA-Seq analysis of *MZpnrc2* mutants are enriched for PREs, AREs, and additional

destabilizing motifs, we performed motif analysis using the MEME Suite tool STREME (Simple,

Thorough, Rapid, Enriched Motif Elicitation) that finds motifs in large datasets (Bailey, 2021).

Among all 8 statistically significant enriched motifs (Tables S4-S5), the sequences UGUACAU

and UGUAAAUA (Fig. 1C) match the destabilizing PRE (Arvola et al., 2020; Bohn et al., 2018;

Galgano et al., 2008; Gerber et al., 2006; Hafner et al., 2010; Lu and Hall, 2011; Morris et al.,

2008; Murata et al., 1995; Wang et al., 2002; White et al., 2001; Zamore et al., 1997; Zamore et

al., 1999). The enriched motifs UAUUUAUU and UAUUAUUA (Fig. 1D) are AREs, motifs that

are recognized by a broad group of ARE-binding proteins (ARE-BPs) that can both inhibit and

promote mRNA decay and/or translation (Briata et al., 2005; Garneau et al., 2007; Gratacos and

Brewer, 2010; Lykke-Andersen and Wagner, 2005; Sanduja et al., 2011). Additional enriched

motifs (Fig. 1E) include GUUUGUUU that matches GU-rich elements (GREs), a motif that is

associated with mRNA decay (Rattenbacher et al., 2010), and the enriched sequence

CCC<u>CCUC</u>C that contains a destabilizing CCUC motif (Vejner et al., 2019). Lastly, we find

enrichment for miRNA target sites (Fig. 1F), including the motif K<u>ACUUGA</u> that matches the

miR-26 seed target site UACUUGA (Crippa et al., 2016; Icli et al., 2013; Klett et al., 2019;

Watterston et al., 2019) and the motif A<u>GCACUU</u> that matches the miR-430 target site

GCACUU (Baia Amaral et al., 2024; Bazzini et al., 2012; Giraldez et al., 2006; Liu et al., 2020;

Rabani et al., 2017; Vejnar et al., 2019). Enrichment of miRNA target sequences suggests that

there is potential cross-talk and/or interactions between Pnrc2- and miRNA-mediated decay.

Overall, we find that all significantly enriched 3'UTR motifs are known to destabilize transcripts.

Furthermore, STREME analysis of unaffected and under-expressed transcript 3'UTRs (see

methods) does not identify enrichment for destabilizing motifs, although unaffected transcript

3'UTRs are enriched for AU dinucleotide repeats (Tables S4-S5). Together, these results

suggest that Pnrc2 is important for decay of mRNAs with destabilizing 3'UTRs.

Loss of Pnrc2 disrupts mRNA expression patterns of many developmental genes

Expression patterns of the oscillatory gene transcripts *her1*, *her7*, and *dlc*, all of which

are associated with the GO term somite specification (Fig. 1B), are disrupted in *MZpnrc2*

mutants (Gallagher et al., 2017; Tietz et al., 2020). We used in situ hybridization to characterize

gene expression patterns and RT-qPCR to validate expression levels of select genes

overexpressed by RNA-Seq analysis and associated with segmentation, oscillatory expression,

and/or Fgf or Wnt signaling. For in situ hybridization, coloration reactions for each probe were

developed in both wild-type and *MZpnrc2* mutant embryos for identical reaction times in parallel.

For RT-qPCR, expression levels were quantified using whole embryo extracts from wild-type

and *MZpnrc2* mutant embryos at mid-segmentation stages. The expression pattern of *rhov*, an

oscillatory gene in zebrafish PSM (Krol et al., 2011), is restricted in wild-type embryos but

expanded in *MZpnrc2* mutants (Fig. 2A, A' vs B, B'). Quantification by RT-qPCR validates that

rhov expression is significantly increased in *MZpnrc2* mutants (Fig. 2C). Expression of *sp5l* (Fig.

2D, D' vs E, E', RT-qPCR in F), a gene whose ortholog *Sp5* oscillates in mouse PSM (Krol et al.,

2011; Matsuda et al., 2020), is similarly affected. Expression of the somite polarity genes

mespab and *ripply2 a*lso show expanded expression in *MZpnrc2* mutants (Fig. 2G, G' vs H, H''

and J, J' vs K, K', RT-qPCR in I and L). Expression of *dact1*, a modulator of Wnt/b-catenin

signaling (Waxman et al., 2004) that oscillates in mouse and human PSM cells (Krol et al., 2011;

Matsuda et al., 2020) is also expanded in *MZpnrc2* mutants (Fig. 2M, M' vs N, N', RT-qPCR in

O). Expression of *spry1* and *spry4*, genes encoding Fgf signaling antagonists whose orthologs

oscillate in mouse and/or human PSM cells (Matsuda et al., 2020) also show expanded

expression patterns (Fig. 2P, P' vs Q, Q' and S, S' vs T, T', RT-qPCR in R and U). Additional

genes known to oscillate in the zebrafish PSM (Krol et al., 2011) and whose transcripts are

overexpressed in *MZpnrc2* mutants were validated by RT-qPCR (Fig. S2). In total, of the 21

genes known to oscillate in the zebrafish PSM (Krol et al., 2011), 10 are Pnrc2-regulated.

Although 24 oscillatory genes were originally identified in microarray experiments (Krol et al.,

2011), annotations for 3 have been removed in GRCz11. Overall, our results show that Pnrc2

promotes decay of mRNAs that encode many developmental regulators; however, the lack of

overt segmentation or additional developmental phenotypes in *MZpnrc2* mutants suggests that

Pnrc2-dependent mRNAs are regulated translationally.

Overexpressed oscillatory gene transcripts in MZpnrc2 mutants have shortened poly(A) tails

Despite overexpression of many oscillatory and segmentation-related genes, *MZpnrc2*

mutants have no overt segmentation phenotype. Expression of *her1* and *dlc* mRNA is increased

4-6 fold in *MZpnrc2* mutants, but the corresponding oscillatory protein is expressed normally,

consistent with the lack of segmentation phenotypes (Gallagher et al., 2017; Tietz et al., 2020).

Discordance between mRNA and protein levels might be due to deadenylation of accumulated

transcripts that leads to inefficient translation. To examine poly(A) (pA) tail length, we performed

a G/I-tailing-based pA tail length (PAT) assay and find that although the predominant *her1* pA tail

length is ~45 nts in both wild-type and *MZpnrc2* mutant embryos, the length distribution is

reduced in *MZpnrc2* mutants, which tend to have shorter tails compared to wild-type embryos

(Fig. 3A-B). Because most *her1* transcripts with short pA tails in *MZpnrc2* mutants would likely

be transcripts that accumulate due to loss of Pnrc2 function, we quantified *her1* and other

oscillatory gene transcripts from extracts derived from pA-enriched and pA-depleted extracts

obtained by oligo(dT) bead-based purification (see methods for details). The oscillatory gene

transcripts *her1*, *her7*, *dlc*, and *rhov* are all significantly overexpressed in pA-depleted extracts,

in contrast to pA-enriched extracts that show no significant differences between wild-type and

MZpnrc2 mutant embryos (Fig. 3C-N). Transcripts with short pA tails may not be efficiently

captured using oligo(dT) beads, so we quantified oscillatory gene transcripts using cDNA

synthesized from total RNA extracts and primed with oligo(dT) to select for polyadenylated

mRNA (rather than priming with random primers as performed for all other quantification

experiments). Quantification of oligo(dT)-primed cDNA shows that oscillatory gene transcripts

her1, *her7*, *dlc*, and *rhov* are all significantly overexpressed in *MZpnrc2* mutants (Fig. S3)

indicating that accumulated transcripts, although shortened, are not completely deadenylated.

These results are consistent with *her1* PAT assay results showing that *her1* transcripts are not

completely deadenylated and instead tend to have short tails in *MZpnrc2* mutants (Fig. 3A-B).

Because *MZpnrc2* mutants lack segmentation phenotypes, we hypothesize that accumulated

tail-shortened oscillatory gene transcripts are inefficiently translated.

Overexpressed P-body associated ddx61 transcripts in MZpnrc2 mutants are substantially

polyadenylated

Accumulation of mRNA in *MZpnrc2* mutants might lead to localization to cytoplasmic

ribonucleoprotein (RNP) granules such as processing bodies (P-bodies) that are sites of

translationally repressed transcripts (Luo et al., 2018; Ostareck et al., 2014). In eukaryotic cells,

P-body formation is enhanced in cells that accumulate mRNA decay intermediates (Sheth and

Parker, 2003; Teixeira et al., 2005) as well as in cells treated with the translational inhibitor

puromycin that results in ribosomes dissociating from mRNAs (Eulalio et al., 2007; Zheng et al.,

2008). Transcripts of several genes associated with the enriched GO term "P-body assembly"

are overexpressed in *MZpnrc2* mutants (Tables S3 and S5). These include *ddx6* and *ddx61* that

are orthologous to the highly conserved *DDX6/p54* DEAD-box helicase gene family that is

required for P-body formation (reviewed in: Luo et al., 2018; Ostareck et al., 2014). Of the two

zebrafish *ddx6* ohnologs (Singh et al., 2020), *ddx61* expression is more than 16-fold higher than

ddx6 at mid-segmentation stages (White et al., 2017), indicating that *ddx61* is the predominantly

expressed ohnolog during segmentation. Upon profiling *ddx61* expression in pA-enriched and

pA-depleted extracts, we find that *ddx61* is significantly overexpressed in both fractions (Fig.

3O-Q). Detection of *ddx61* overexpression in pA-enriched extracts suggests that *ddx61*

overexpression in *MZpnrc2* mutants might lead to increased Ddx61 protein levels, in contrast to

oscillatory gene transcripts for which overexpression is detected only in pA-depleted extracts.

Accumulated oscillatory gene transcripts in MZpnrc2 mutants are disengaged from translational

machinery, whereas overexpressed ddx61 transcripts are translationally engaged

252   Because accumulated oscillatory gene transcripts in *MZpnrc2* mutants have reduced tail

lengths, we assessed engagement of *her1* and other oscillatory gene transcripts with

ribosomes. We conducted polysome profiling from wild-type and *MZpnrc2* mutant embryos at

mid-segmentation stages to compare oscillatory gene transcript levels in actively translating

versus non-translating fractions (see methods). The overall polysome profiles are similar in wild-

type and *MZpnrc2* mutant embryos (Fig. 4A), suggesting that global translation is normal in

mutants; however, *her1* transcripts in *MZpnrc2* mutants are 20-fold higher in unbound/subunit

fractions, and not significantly different in monosome- and polysome-bound fractions compared

to wild-type *her1* levels (Fig. 4B). Similar distributions were obtained for other oscillatory gene

transcripts including *her7*, *dlc*, and *rhov* (Fig. 4C-E). Together, these results support the

hypothesis that accumulated oscillatory gene transcripts in *MZpnrc2* mutants are not efficiently

translated. Because *ddx61* mRNA is overexpressed in pA-enriched RNA extracts (Fig. 3P), we

profiled *ddx61* mRNA abundance across fractions and find that *ddx61* levels are 3-fold higher in

MZpnrc2 polysome fractions relative to wild-type polysome fractions (Fig. 4F). To corroborate

polysome profiling results, we performed immunoblot detection of Ddx61 using wild-type and

MZpnrc2 mutant extracts. Ddx6 and Ddx61 cannot be discriminated because both have the

same molecular weight and antibodies that distinguish between the two highly similar proteins

are not available. Immunoblot results show a corresponding increase in Ddx6/Ddx61 protein

levels in *MZpnrc2* mutants at mid-segmentation stages (Fig. 4G-H), corroborating polysome

profiling. Overall, our results show that *ddx61* gene products, unlike oscillatory genes, are

overexpressed at both mRNA and protein levels in *MZpnrc2* mutants.

MZpnrc2 mutants are sensitized to deadenylation inhibition

Because several oscillatory gene transcripts have shortened pA tails in *MZpnrc2*

mutants, we hypothesized that blocking deadenylation would alleviate their inefficient translation

and lead to segmentation and/or additional phenotypes. Ectopic expression of a CCR4-NOT

dominant negative deadenylase via mRNA injection into zebrafish embryos has previously been

shown to disrupt segmentation and prevent pA tail shortening of oscillatory gene transcripts

including *her1* and *her7* (Fujino et al., 2018). Consistent with the published work, we confirmed

that *Myc-Cnot7DN* mRNA injection causes segmentation defects in wild-type embryos, with over

84% of injectants exhibiting segmentation defects (Fig. S4). To determine whether *MZpnrc2*

mutants are more sensitive to deadenylation inhibition than wild-type embryos, we induced

mosaic expression of Myc-tagged Cnot7DN at mid-segmentation stages in embryos injected

with a heat shock-inducible construct and scored them for segmentation defects 6 hours post-

heat shock (see methods for details). We find that mosaic expression of Myc-Cnot7DN leads to

abnormal segment formation in *MZpnrc2* mutants, but not in wild-type embryos (Fig. 5A-E),

consistent with the hypothesis that *MZpnrc2* mutants are sensitized to deadenylase inhibition.

Using anti-Myc immunohistochemistry on plasmid-injected embryos, we confirmed that *Myc-*

cnot7DN expression was mosaically induced post-heat shock and was absent in no-heat shock

controls (Fig. 5F-G).

MZpnrc2 mutants are sensitized to depletion of Ddx61 and Ddx6 function

Because *ddx61* is overexpressed at both mRNA and protein levels, depleting Ddx61

function in *MZpnrc2* mutants might lead to segmentation and/or additional phenotypes. Using a

highly efficient F0 CRISPR knockdown strategy (Hoshijima et al., 2019; Klatt Shaw and

Mokalled, 2021), we find that single *ddx61* crispants (as well as single *ddx6* crispants) develop

normally up to 5 days post-fertilization (dpf) and can be raised without significant losses up to

adulthood (data not shown). To estimate efficiency of our F0 CRISPR strategy (Fig. S5A), we

employed high resolution melt analysis (HRMA) to detect CRISPR-induced mutations and

confirmed that all CRISPR-injected embryos were mutagenized (Fig. S5B-C). Because *ddx6*

might compensate for loss of *ddx61* function, we also targeted *ddx6*. In *ddx6* and *ddx61* double

crispants, *ddx6* and *ddx61* mRNA levels are more than 50% reduced at 18 hpf compared to

uninjected sibling controls (Fig. S5D-E), likely because the premature termination codon-

containing transcripts created by our targeting strategy are degraded by NMD (reviewed in:

Causier et al., 2017; He and Jacobson, 2015; Karousis et al., 2016; Kervestin and Jacobson,

2012; Lykke-Andersen and Bennett, 2014; Schoenberg and Maquat, 2012). Corresponding with

reduced mRNA levels, Ddx6/Ddx61 protein levels are also reduced in *ddx6*;*ddx61* crispants

over time, becoming nearly absent by 5 dpf (Fig. S5F). Coinciding with reduced Ddx6/Ddx61

protein levels, we find that *ddx6;ddx61* crispant wild-type and *MZpnrc2* mutant embryos exhibit

morphological abnormalities beginning at 2 dpf that worsen with time. By 5 dpf, some

ddx6;*ddx61* crispant larvae exhibit strong morphological abnormalities, with *MZpnrc2* mutants

showing higher frequency and more severe abnormalities than wild-type larvae (Fig. 6A-B).

Corresponding with morphological defects, Ddx61/Ddx6 protein is strongly reduced in both

ddx6;*ddx61* crispant wild-type and *MZpnrc2* mutant embryos at 5 dpf (Figs 6C, S5G-H). The

lack of overt phenotypes prior to 2 dpf may be due to maternal contribution that compensates

for loss of zygotic function at early stages. Consistent with this possibility, both *ddx61* mRNA

and Ddx61 protein are abundant prior to zygotic genome activation according to the Zebrafish

Protein Profiling interactive tool (da Silva et al., 2024).

Morphological defects in *ddx6;dd61* crispant larvae might be preceded by earlier

molecular phenotypes. To determine whether *her1* expression is affected in *ddx6;ddx61*

crispants, we performed *her1* in situ hybridization on *ddx6*;*ddx61* crispant wild-type and

MZpnrc2 mutant embryos at mid-segmentation stages. Although the *her1* expression pattern in

ddx6;*ddx61* crispant wild-type embryos was indistinguishable from uninjected wild-type controls

(Fig. 7A,A' vs B,B'), *ddx6*;*ddx61* crispant *MZpnrc2* mutants showed an enhanced misexpression

phenotype compared to uninjected *MZpnrc2* mutant controls (Fig. 7C,C' vs D, D'). We

separately quantified *her1* levels by RT-qPCR and find that *her1* is increased more than 1.5-fold

in *ddx6*;*ddx61* crispant *MZpnrc2* mutants relative to control *MZpnrc2* mutants (Fig. 7E), but is

unaffected in *ddx6*;*ddx61* crispant wild-type embryos. In contrast, *her7* expression is modestly

reduced, whereas *dlc* and *rhov* expression is unaffected in *ddx6*;*ddx61* crispant *MZpnrc2*

mutants (Fig. 7F-H). Overall, normal levels of oscillatory gene transcripts in *ddx6*;*ddx61* crispant

wild-type embryos suggest that Ddx61 and Ddx6 are not required for mRNA decay when Pnrc2

is functional. The synergistic effect of loss of *pnrc2*, *ddx6*, and *ddx61* shows that Ddx6 and

Ddx61 partially compensate for loss of Pnrc2 function.

Discussion

In this work, we show that overexpressed transcripts in *MZpnrc2* mutants are enriched

for destabilizing sequences. Among overexpressed transcripts, many correspond to genes that

are dynamically regulated, including oscillatory and/or developmental genes that function in

Notch, Fgf, and Wnt signaling pathways. Remarkably, despite overexpression of >1700

transcripts, segmentation and other developmental processes appear normal in *MZpnrc2*

mutants, although many fail to thrive as juveniles and adults. We previously suggested that

absence of segmentation defects is likely because accumulated mRNAs in *MZpnrc2* mutants

are not efficiently translated (Gallagher et al., 2017), though the mechanism was unknown. In

the current study, we explored mechanisms driving discordant expression of mRNA and protein

in *MZpnrc2* mutants by assessing the polyadenylation status of *her1* and other transcripts, their

level of engagement with translational machinery, and the impact of P-body factors Ddx6 and

Ddx61 on the *MZpnrc2* overexpression phenotype.

Overexpressed transcripts in MZpnrc2 mutants are enriched for destabilizing motifs

Most differentially expressed transcripts in *MZpnrc2* mutants are overexpressed, which

we expected based on previous human cell-based studies that showed that PNRC2 functionally

interacts with mRNA decay machinery and promotes mRNA decay (Cho et al., 2009; Cho et al.,

2012; Cho et al., 2013; Cho et al., 2015; Lai et al., 2012; Mugridge et al., 2016; Nicholson et al.,

2018). Motif enrichment analysis of overexpressed transcripts in *MZpnrc2* mutants reveals that

PRE and ARE motifs are the most significantly enriched motifs, extending our previous findings

that the *her1*, *her7*, and *dlc* 3'UTRs each contain PRE and ARE motifs, and in the case of the

her1 3'UTR, terminally located PRE and ARE motifs drive rapid decay of reporter mRNA in vivo

(Tietz et al., 2020). In eukaryotes, transcriptome-wide profiling has shown that PREs and AREs

destabilize mRNA in human and mouse cell lines (Siegel et al., 2022; Schwanhausser et al.,

2011; Yang et al., 2003), mouse embryonic stem cells (Sharova et al., 2009), *Drosophila* cell

lines (Cairrao et al., 2009) and embryos (Burow et al., 2015), as well as in zebrafish embryos

(Rabani et al., 2017; Vejner et al., 2019). Pumilio (Pum) and ARE-binding proteins (ARE-BPs)

preferentially bind these motifs, respectively, and recruit deadenylation factors to initiate mRNA

decay (Arvola et al., 2020; Bulbrook et al., 2018; Enwerem et al., 2021; Joly et al., 2013;

Kadyrova et al., 2007; Weidmann et al., 2014). Because the poly(A) tail promotes translation

and mRNA stability, deadenylation leads to reduced translation and enhanced decay (Garneau

et al., 2007; Jackson et al., 2010).

In zebrafish, a reporter-based study found that PREs and AREs are associated with

rapid decay during the maternal-zygotic transition (MZT) (Rabani et al., 2017). Analysis of

endogenous transcripts revealed that AREs and CCUC motifs are among the most destabilizing

motifs during MZT (Vejner et al., 2019). We found that the C-rich motif CCC<u>CCUC</u>C that

contains a CCUC stretch is enriched in overexpressed transcripts in *MZpnrc2* mutants.

Similarly, C-rich elements are enriched in 3'UTRs of unstable mRNAs in human induced

pluripotent stem cells (Neff et al., 2012). These observations that C-rich sequences are

associated with transcript instability are somewhat surprising because C-rich sequences are

typically associated with mRNA stability through binding of poly(C)-binding proteins (PCBPs), a

family of multifunctional RBPs (Bomsztyk et al., 1997; Makeyev and Liebhaber, 2002; Ostareck-

Lederer et al., 1998). The role of C-rich sequences and their associated RBPs is likely context

and cell-type dependent, exemplified by the finding that zebrafish Pcbp2 preferentially binds

CCUC motifs and is associated with unstable mRNAs during MZT (Vejner et al., 2019). The GU-

rich sequence GUUUGUUU is also enriched in the 3'UTRs of overexpressed transcripts; this

sequence is found within the known destabilizing GU-rich element (GRE) U<u>GUUUGUUU</u>GU, a

3'UTR motif enriched in short-lived mRNAs in primary human T cells and associated with CUG-

binding proteins (Halees et al., 2011; Rhagavan et al., 2002; Vlasova et al., 2008).

Additional enriched motifs from our analysis include matches to miRNA seed target sites

for *miR-26* and *miR-430*. Our previous work showed that *her1* mRNA decay is independent of

Dicer-dependent miRNAs (Gallagher et al., 2017), consistent with the absence of either site in

the *her1* 3'UTR. Whether other Pnrc2-regulated transcripts utilize miRNA-based regulation for

their decay is unknown. miRNAs recruit the two major deadenylation complexes, Pan2-Pan3

and CCR4-NOT, to their target mRNAs by association with the miRNA-induced silencing

complex (Braun et al., 2011; Chekulaeva et al., 2011; Fabian et al., 2011). However,

mechanisms by which miRNAs reduce expression of target mRNAs are context dependent,

exemplified by studies using zebrafish embryos during MZT (Bazzini et al., 2012) and studies

using *Drosophila* S2 cells (Djuranovic et al., 2012) that showed that miRNA target mRNAs are

translationally repressed prior to deadenylation. Destabilizing 3'UTR motifs are associated with

translational repression, however distinguishing between direct effects that RBPs and miRNAs

bound to target mRNAs have on translational machinery from indirect effects due to

deadenylation and decay can be challenging. This is further complicated by the ability of some

members of the CCR4-NOT deadenylation complex to recruit translational repressors

independent of deadenylation (Cooke et al., 2010; Waghray et al., 2015).

Transcripts of many genes that regulate post-transcriptional processes depend on Pnrc2 for

normal expression levels

Among enriched GO terms identified by analysis of overexpressed transcripts in

MZpnrc2 mutants, some are related to post-transcriptional regulation including "3'UTR-mediated

mRNA destabilization, deadenylation-dependent decay" and "P-body assembly". Genes in the

first category are primarily *zfp36* family members, with five out of six *zfp36* orthologs

overexpressed in *MZpnrc2* mutant embryos (*zfp36l1a*, *zfp36l1l*, *zfp36l2*, *zgc:114130*, and

zgc:162730). Zfp36 members are CCCH-type zinc finger proteins that bind AREs and

destabilize target mRNAs by recruiting deadenylation factors (reviewed in: Blatnik et al., 2023;

Ciais et al., 2013; Hau et al., 2007). It is possible that *zfp36* overexpression in *MZpnrc2* mutant

embryos leads to an increase in Zfp36 protein levels that enhances deadenylation and reduces

excess translation of *pnrc2*-regulated transcripts, however it may simply be that *zfp36*

transcripts are *pnrc2*-dependent for their decay. Future analysis of Zfp36 protein levels in

MZpnrc2 mutant embryos coupled with genetic analysis of *zfp36* mutants will provide insight

into these possibilities.

Enrichment of the GO term "P-body assembly" raises the possibility that mechanisms

that sequester mRNAs from translational machinery are active in *MZpnrc2* mutants.

Overexpressed transcripts with this GO term include *edc3*, *lsm14aa, ddx6*, and *ddx61.* In yeast,

Edc3 is a decapping enhancer that activates decapping and promotes P-body formation

(reviewed in: Ditlev et al., 2018; He and Jacobson, 2023). LSm14/RAP55, similarly interacts

with decapping factors and promotes P-body formation (reviewed in: Marnef et al., 2009; Vidya

and Duchaine, 2022). The conserved Ddx6 family of helicases (also known as Rck/p54) are

decapping activators that are required for P-body formation and interact with Edc3, LSm14, and

other P-body factors (Luo et al., 2018; Ostareck et al., 2014). Zebrafish possess two Ddx6

helicases, Ddx6/P54a and Ddx61/P54b, both of which localize to cytoplasmic granules during

zebrafish embryonic development (Zampedri et al., 2016). Among P-body assembly genes, we

focused on *ddx6* and *ddx61* because transient knockdown of these genes leads to

morphological defects in several tissues including somites (Zampedri et al., 2016) and because

DDX6 overexpression in human pluripotent stem cells increases P-body number (Di Stefano et

al, 2019). Future analysis of *edc3* and *lsm14aa* will provide further insight into the role of P-

bodies during segmentation.

Depletion of RNA helicases Ddx6 and Ddx61 leads to developmental defects in MZpnrc2

mutants

In the current work, we show that Ddx6/Ddx61 RNA helicases are overexpressed at both

mRNA and protein levels in *MZpnrc2* mutants. Depletion of *ddx6* and *ddx61* function in *MZpnrc2*

mutants, but not wild-type embryos, enhances *her1* accumulation and leads to strong

morphological larval phenotypes that are also present at much lower frequency in wild-type

crispants. Higher rates of severe larval phenotypes in *ddx6;ddx61 MZpnrc2* crispants compared

to wild-type crispants suggest that Pnrc2-regulated transcripts are over-translated upon

depletion of Ddx6 and Ddx61 function. It is possible that Ddx6 and Ddx61 promote tail-

shortening of accumulated mRNAs in *pnrc2* mutants, similar to what has been shown for human

DDX6 that promotes deadenylation of inefficiently translated mRNAs in human cells (Weber and

Chang, 2024). We did not observe morphological defects in *ddx6;ddx61* crispant embryos at 1

dpf in either wild-type or *MZpnrc2* mutant backgrounds, likely due to maternal contribution that

compensates for loss of zygotic Ddx6 and Ddx61 function. The absence of an early phenotype

differs from previous work which showed that *ddx6* and *ddx61* morpholino (MO) injection leads

to somite and notochord defects (Zampedri et al., 2016), which we suspect may be due to MO-

induced toxicity and/or off-target effects caused by the high doses used (Schulte-Merker and

Stainier, 2014; Stainier et al., 2017). Future work using genetic mutants will help resolve this

discrepancy.

The finding that *her1* mRNA accumulation in *MZpnrc2* mutants is enhanced in

ddx6;ddx61 crispants suggests that P-body factors promote *her1* mRNA decay in the absence

of Pnrc2. P-bodies are thought to primarily function as sites of mRNA storage rather than decay,

supported by observations that mRNA decay continues even in the absence of P-bodies

(Decker et al., 2007; Eulalio et al., 2007; Stoecklin et al., 2006). Early reports of mRNA decay in

P-bodies are attributable to artifacts of the MS2 tethering system which inadvertently stabilizes

partially degraded reporter mRNAs that subsequently localize in P-bodies (Garcia & Parker,

2015; Heinrich et al., 2017; Tutucci et al., 2018). Recent work employing a new tethering

method to monitor mRNA dynamics suggests that some mRNAs are degraded rapidly within P-

bodies, but decay can be inadvertently inhibited due to stress induced by high laser intensities

frequently used during time-lapse imaging (Blake et al., 2024). An alternative explanation for

enhanced *her1* accumulation in *ddx6;ddx61* crispant *MZpnrc2* mutants is that Ddx6/Ddx61

promotes *her1* mRNA decay in a P-body-independent manner, as both RNA helicases show

diffuse cytoplasmic expression in addition to enrichment in cytoplasmic foci (Zampedri et al.,

2016). A recent human cell culture-based study found that DDX6 binds stalled ribosomes,

recruits the CCR4-NOT deadenylase complex, and triggers decay of inefficiently translated

mRNAs (Weber and Chang, 2024), an effect that presumably occurs outside of P-bodies. The

extent to which *her1* mRNA is degraded in the cytoplasm versus P-bodies as well as the

dependence of these activities on Pnrc2 function remains to be explored and future localization

studies will shed light on these possibilities.

Conclusions

Pnrc2 promotes rapid mRNA decay during vertebrate segmentation, including the decay

of many dynamically expressed oscillatory gene transcripts. Overexpressed transcripts in

MZpnrc2 mutants are enriched for destabilizing motifs that are associated with translational

repression and mRNA decay. *MZpnrc2* mutant embryos develop normally despite dysfunctional

mRNA decay and this is likely because transcripts that depend on Pnrc2 for their rapid decay

have shortened poly(A) tails that are inefficiently translated. Consistent with this view, inhibiting

deadenylation leads to somite and additional defects in sensitized *MZpnrc2* mutants, likely due

to over-translation of Pnrc2-regulated transcripts. Upregulation of *ddx61* at both mRNA and

protein levels suggests that compensatory mechanisms are employed in decay-deficient *pnrc2*

mutants. The synergistic effect of loss of *pnrc2*, *ddx6*, and *ddx61* leads to strong developmental

phenotypes, suggesting that accumulated mRNAs are translated in *pnrc2* mutants upon

depletion of the RNA helicases Ddx6 and Ddx61. Overall, our results show that Ddx6 and

Ddx61 regulate *her1* oscillatory expression and likely ensure proper protein levels in mRNA-

decay deficient mutants.

Materials and methods

Animal stocks and husbandry

All protocols using zebrafish were approved by the Ohio State University Institutional Animal

Care and Use Committee (protocol 2012A00000113-R4). Adult zebrafish strains (*Danio rerio*)

were kept at 28.5°C on a 14 hour (h) light/10 h dark cycle and obtained by natural spawning or

in vitro fertilization (Westerfield, 2007). Embryos and larvae were staged according to Kimmel et

al.,1995. The *pnrc2*^{oz22} line has been described previously (Gallagher et al., 2017).

DNA extraction and genotyping

Individual embryos and adult fin tissue were lysed in 50 ul 1X ThermoPol Buffer (NEB) at 95°C

for 10 minutes, digested at 55°C for 1 hour in 1 mg/ml Proteinase K (BP1700, ThermoFisher),

followed by Proteinase K inactivation at 95°C for 10 minutes. Molecular identification of *pnrc2*^{oz22}

carriers by PCR amplification followed by NsiI-HF digestion to distinguish cleavable wild-type

from uncleavable mutant amplicons was performed as previously described (Tietz et al., 2020).

RNA-Seq sample collection and sequencing

Whole embryos at mid-segmentation stages (n = 50 embryos per biological replicate for each

genotype) were solubilized in Trizol for RNA extraction and purified following standard

procedures (Thermo Fisher). Both wild-type and *MZpnrc2* mutant embryo samples were

prepared in biological triplicate. 2 ug of total RNA from each sample was processed at

Novogene using a whole transcriptome sequencing workflow with ribosomal RNA (rRNA)

depletion that enables sequencing of RNA irrespective of polyadenylation status. At Novogene,

each sample was depleted of rRNA using a Ribo-Zero Kit (Illumina), followed by chemical

fragmentation and 150 bp paired-end library synthesis using a NEBNext Ultra II Directional RNA

kit (NEB). Libraries were sequenced on the Illumina NovaSeq 6000 platform by Novogene.

RNA-Seq data analysis

High-throughput sequencing reads were trimmed to remove adapter and primer sequences

using Cutadapt version 3.4 (Martin, 2011) and pseudoaligned to the zebrafish

GRCz11/danRer11 transcriptome reference using Kallisto (version 0.46.2), an RNA-Seq tool

that rapidly quantifies transcript abundance without the need for individual base alignment (Bray

et al., 2016). Cutadapt and Kallisto were installed and run in Python (version 2.7.5) using

standard procedures described in online guides available at

https://cutadapt.readthedocs.io/en/stable/guide.html and

https://pachterlab.github.io/kallisto/manual. Prior to running Kallisto, all sequence data files were

assessed using the quality control tool FastQC version 0.12.1 (Andrews, 2010) and no quality

issues were identified in relevant modules for RNA-Seq analysis (data not shown). Transcripts

with <5 estimated read counts in at least one sample were filtered out to reduce noise.

Differential gene expression analysis was performed using Sleuth version 0.30.1, a program

that uses variance decomposition to identify biological differences between RNA-Seq samples

(Pimentel et al., 2017). Sleuth was run in RStudio version 2023.06.2 Build 561 (Posit Software)

using standard procedures described in an online guide at

https://pachterlab.github.io/sleuth_walkthroughs/trapnell/analysis.html. Batch effects including

effects due to sequencing lane or biological replicate differences were tested and no effects

were observed (data not shown). A summary of the RNA-Seq processed data, Kallisto

normalized read counts, and Sleuth-generated differential expression results are included as

supplementary data (Tables S1-S3).

Motif analysis

Motif analysis was performed using 3'UTR input sequences corresponding to transcripts

identified from RNA-Seq analysis of *MZpnrc2* mutant embryos (Tables S3-S4) and the MEME

Suite tool STREME version 5.5.6 (Bailey et al., 2021). Annotated 3'UTR sequences from each

of the following datasets were analyzed using STREME: 1) significantly overexpressed

transcripts (q < 0.05; n = 1320), 2) unaffected transcripts (log2FC from -0.05 to 0.05; n = 1334),

and 3) significantly under-expressed transcripts (i.e. transcripts with decreased expression in

MZpnrc2 mutants relative to wild-type embryos; q < 0.05; n = 690) (Tables S3-S4). 3'UTR

sequence annotations were obtained using BioMart (Ensembl release 113, GRCz11 assembly)

and those less than 50 nts in length were removed to reduce annotation and length biases.

Motif range was set to 6-8 nts and shuffled input sequences were used as the control with

Markov order set to 2 (the default setting used by STREME). Sequence logos (Schneider and

Stephens, 1990; Schneider et al., 1986) were generated using probability matrix data from

STREME and the seqLogo package (Bembom and Ivanek, 2024) available with BioConductor

version 3.19 (Gentleman et al., 2004; Huber et al., 2015) in RStudio version 2023.06.2 Build

561 (Posit Software). Motif enrichment results are reported in Table S5.

GO analysis

GO analysis was performed using the PANTHER Overrepresentation Test (Released 2024-02-

26) that is available from the Gene Ontology Consortium at https://geneontology.org/ (Mi et al.,

2013; Thomas et al., 2022). The GO Ontology database version used was DOI:

10.5281/zenodo.10536401 (Released 2024-01-17). The input list contains gene IDs for which at

least one transcript was significantly overexpressed (q < 0.05) in *MZpnrc2* mutants and for

which a GO annotation was identified (n = 1619 uniquely annotated GO IDs) (Table S4), while

the reference list contains the full set of GO-annotated zebrafish genes in the database (n =

26,353). GO results are reported in Table S5. Statistical test type was a Fisher's Exact test

corrected by calculating the false discovery rate.

In situ hybridization

Whole mount in situ hybridization was performed as previously described (Broadbent and Read,

1999; Jowett, 1998) using digoxygenin (DIG)-labeled riboprobes. Riboprobe for *her1* was

previously described (Delaune et al., 2012; Dill and Amacher, 2005). Riboprobe for *rhov* was

synthesized from a plasmid containing the *rhov* cDNA (cb832) received from the Zebrafish

International Resource Center (ZIRC). *rhov*-containing plasmid was linearized with NotI-HF

(NEB) and transcribed with T7 RNA polymerase (Roche Life Science) to make DIG-labeled

antisense *rhov* riboprobe. Riboprobe templates for *sp5l*, *mespab*, *ripply2*, *dact1*, *spry1*, and

spry4 were generated by PCR amplification of coding sequence (primer sequences are listed in

Table S6). All DIG-labeled PCR-based probes were synthesized by transcribing PCR templates

with SP6 or T7 RNA polymerase (Roche Life Science). *dact1*, *sp5l*, and *spry4* produced

expression patterns consistent with direct submission in situ hybridization data (Thisse et al.,

2001; Thisse et al., 2005). Additionally, expression patterns for all PCR-based probes shown in

Fig. 2 match published expression patterns for wild-type embryos at mid-segmentation stages,

including *dact1* (Gillhouse et al., 2004), *mespab* (Cutty et al., 2012; Wanglar et al., 2014), *sp5l*

(Superina et al., 2014), *ripply2* (Junker et al., 2014; Kinoshita et al., 2018; Moreno et al., 2008;

Wanglar et al., 2014; Yabe et al., 2016), *spry1* (Komisarczuk et al., 2008), and *spry4* (Nguyen-

Chi et al., 2012). In situ hybridized embryos were mounted in Permount and imaged using an

Axiocam HRc digital camera with AxioPlan2 microscope and 10x objective (Zeiss).

RNA extraction and cDNA synthesis

Whole embryos at mid-segmentation stages (n = 15 embryos per biological replicate for each

genotype in Fig. 2; n = 10 embryos per biological replicate for each genotype and condition in

Fig. 7) were solubilized in Trizol for RNA extraction and purified following standard procedures

(Thermo Fisher). All genotypes and conditions were performed in biological triplicate. 500 ng of

total RNA was reverse transcribed with random primers and Superscript IV reverse

transcriptase according to the manufacturer's instructions (Thermo Fisher).

Quantitative RT-PCR analysis

Quantitative RT-PCR was performed using PowerUp SYBR Green Master Mix (Thermo Fisher)

and 4.5 ul cDNA (diluted 1:50 in Fig. 2, diluted 1:25 in Fig. 7) in 20 ul reactions, following

manufacturer's procedures. Negative controls lacking template were included for each primer

set. All reactions were subjected to thermal melting to confirm that each reaction gave single

peaks. Transcript levels were normalized to *mobk13* (*mob4*) (Gangras et al., 2020; Hu et al.,

2016; Tietz et al., 2020). Cycle thresholds (Ct) were determined using CFX Maestro version 2.3

software (Bio-Rad). The average Ct of 2-3 technical replicates per sample was used for

calculations. Changes in mRNA expression were calculated by $\Delta\Delta Ct = \Delta Ct$ target $- \Delta Ct$ control.

Relative changes in mRNA expression levels are represented graphically as fold change, where

relative mRNA fold change = $2^{-\Delta\Delta Ct}$. All graphs and statistics were generated using Prism 10

(GraphPad).

Poly(A) Tail-Length Assay

RNA for poly(A) tail-length assays was obtained by collecting whole embryos at mid-

segmentation stages (n = 18 embryos for each genotype) followed by solubilization in Trizol and

purification following standard manufacturer procedures (Thermo Fisher). Tail-lengths for *her1*

transcripts were assessed using the Poly(A) Tail-Length Assay Kit following standard

manufacturer procedures (Thermo Fisher). Minus RT controls were performed for all samples

and no amplification was detected as expected (data not shown). Primers used for tail-length

assay PCR are listed in Table S6.

Preparation and analysis of poly(A)-enriched and poly(A)-depleted extracts

RNA for comparison of expression levels in poly(A)-enriched versus poly(A)-depleted RNA

fractions was obtained from the same RNA sources as was used for RNA-Seq experiments

described earlier. Three biological replicates were obtained for both wild-type and *MZpnrc2*

mutant embryos. From each sample, 1 ug of total RNA was adjusted to a final volume of 50 ul in

bead binding buffer (20 mM Tri-HCl, pH 7.5, 1.0 M LiCl, and 2 mM EDTA) and incubated with

Dynabeads Oligo(dT)$_{25}$ following manufacturer procedures (Thermo Fisher). RNA bound to

beads (poly(A)-enriched RNA) was eluted in 10 mM Tris-HCl (pH 7.5). Unbound RNA (poly(A)-

depleted RNA) was precipitated by adding 0.1x volume 3M sodium acetate and 2x volume ice-

cold ethanol, followed by vortexing and incubation overnight at -80°C. After centrifugation at 4°C

for 20 minutes at 12,000 x g, precipitated pellets from unbound RNA fractions were rinsed with

ice-cold 70% ethanol and resuspended in 10 mM Tris-HCl (pH 7.5) at an equivalent volume

used for elution of bound RNA fractions. Each RNA fraction was reverse transcribed using

random primers and Superscript IV reverse transcriptase (RT) according to the manufacturer's

instructions (Thermo Fisher). As a control, 1 ug of total RNA from the same samples used for

bead-based fractionation were reverse transcribed in parallel. RT-qPCR analysis was performed

as described earlier using 4.5 ul of cDNA diluted 1:10.

Protein extraction and Immunoblot analysis

Protein extracts were prepared from zebrafish embryos or larvae at 1-5 days post-fertilization.

Individuals were collected and washed twice with rapid pipetting to disrupt yolks in 200 ul of ice-

cold Ringer's solution with protease inhibitors (116 mM NaCl, 2.9 mM KCl, 5.0 mM HEPES, pH

7.2, 100 mg/mL Soybean Trypsin Inhibitor [Sigma], 0.2 mg/mL PMSF, Complete Protease

Inhibitor Cocktail [Roche]). Embryo and larval suspensions were centrifuged at 4°C for 1 minute

at 300 x g for each wash. Cell lysis was performed by addition of 1 ul 4X LDS Buffer (Thermo

Fisher) for each embryo or larva collected (n = 20-65 individuals per extraction), followed by

homogenization with a plastic pestle, vortexing for 1 minute, and boiling for 10 minutes.

Homogenization, vortexing, and boiling were performed twice sequentially. Embryo or larval

equivalents were loaded on a 4-12% Novex Bis-Tris polyacrylamide gel (Thermo Fisher),

electrophoresed in NuPAGE MOPS SDS running buffer (Thermo Fisher) alongside a Novex

Sharp pre-stained protein standard (Thermo Fisher), and transferred to a PVDF membrane

using the Bio-Rad wet transfer system following standard manufacturer procedures. Immunoblot

was blocked in 5% non-fat skim milk in Tris-buffered saline (20 mM Tris, 150 mM NaCl, pH 7.5)

and subsequent steps were performed following standard procedures available online from Bio-

Rad in bulletin 6376 (https://www.bio-rad.com/). The immunoblot was incubated with DDX6

antibody (A16270, AbClonal) followed by incubation with AP-conjugated goat anti-rabbit IgG

(H+L) secondary antibody (Thermo Fisher). Immunoblots were then incubated with β-actin (sc-

47778, Santa Cruz Biotechnology) antibody followed by incubation with AP-conjugated goat

anti-mouse IgG (H+L) secondary antibody (Thermo Fisher). Primaries were diluted 1:1000 and

secondaries were diluted 1:10,000 in blocking solution with 0.1% Tween-20. Primary antibody

incubations were performed overnight at 4°C, secondary antibody incubations were performed

for 1 hour at room temperature. Chemiluminescent detection was performed using CDP-Star

following the manufacturer's procedure (Thermo Fisher) and imaged using the Sapphire FL

Biomolecular Imager (Azure Biosystems). Quantification of signal intensities was performed

using lane profile analysis in Alpha Imager software (Protein Simple).

Plasmid construction & transgenesis

The heat shock construct *hsp70l:Myc-cnot7DN-SV40 pA* was assembled by first constructing

plasmid *pMA-Myc-cnot7DN* using the GeneArt Plasmid Construction Service (Thermo Fisher).

pMA-Myc-cnot7DN contains the coding sequence for a dominant negative form of zebrafish

Cnot7 (*Cnot7DN*) with an N-terminal C-myc epitope tag (Evan et al., 1985) and intervening

SGLRS flexible linker peptide. The *myc*-tagged *Cnot7DN* coding sequence was based on

plasmid M338 (*pCS2 + MT-CNOT7-D40A,E42E,C67E,L71E-sv40*) that has previously been

used to inhibit deadenylase function in zebrafish embryos (Fujino et al., 2018; Makino et al.,

2015; Mishima and Tomari, 2016; Mishima and Tomari, 2017). After sequence validation of

pMA-Myc-cnot7DN, the *Myc-cnot7DN* insert was cloned into *pBSKI2* (Thermes et al., 2002)

using standard restriction digestion-based cloning with the enzymes NheI-HF and BamHI-HF

(NEB). Plasmid and I-SceI enzyme were injected into the blastomere of 1-cell stage wild-type

and *MZpnrc2* mutant embryos following standard transgenesis procedures (Thermes et al.,

2002). To determine the optimal dose, we injected 20 pg and 60 pg plasmid doses in the

absence of heat-shock induction and assessed morphology at mid-segmentation stages; the 20

pg dose was well-tolerated (73 of 90 injected embryos had wild-type morphology), but the 60 pg

dose caused gross abnormalities (0 of 56 injected embryos had wild-type morphology).

Heat shock assay

Plasmid-injected embryos (described above) were collected and raised to segmentation stages

(10-12S), heat shocked at 37°C for 30 minutes, and monitored for somite defects at 30 minute

intervals post-heat shock (pHS). Live embryos at 6 hours post-heat shock were mounted in 3%

methylcellulose and imaged at 20x magnification using AxioVision Software (Zeiss) on a

MZFLIII Fluorescence Stereo Microscope (Leica) with an AxioCam HRc digital camera (Zeiss)

and subsequently fixed in 4% PFA for anti-Myc immunohistochemistry.

Anti-Myc Immunohistochemistry

Embryos injected with *hsp70l:Myc-cnot7DN* plasmid that were heat shocked at ~14 hours post

fertilization (hpf) for 30 minutes were collected 6 hours post-heat shock (~22 hpf) alongside non-

heat shocked controls, fixed in 4% PFA in PBS overnight at 4°C, and washed 5 x 5 minutes in

PBS with 0.1% Tween-20 (PBST). Embryos were then incubated in water for 5 minutes,

permeabilized with ice-cold acetone for 7 minutes, and incubated again in water for 5 minutes

followed by 3 x 5 minute PBST washes. Blocking was performed in PBST with 2% BSA, 2%

goat serum for 1 hour at room temperature, followed by incubation with 1:500 dilution

monoclonal anti-myc (clone 9E10, MA1-980, Thermo Fisher) in PBS with 2% BSA, 2% goat

serum, 1% DMSO and 0.1% Triton X-100 (PBDTx) overnight at 4°C. Embryos were washed 8 x

15 minutes in PBST, blocked as described above, followed by incubation with 1:200 dilution

Alexa Fluor 546 goat anti-mouse secondary antibody (Thermo Fisher) in PBDTx overnight at

4°C. Embryos were washed 8 x 15 minutes in PBST prior to imaging. Immuno-stained embryos

were mounted in Fluoromount-G and imaged at 20x magnification using AxioVision Software

(Zeiss) on an MZFLIII Fluorescence Stereo Microscope (Leica) with a mCherry filter set and

AxioCam HRc digital camera (Zeiss).

Polysome profiling

Wild-type and *MZpnrc2* mutant embryos were raised to mid-segmentation stages and treated in

100 ug/ml cycloheximide for 5 minutes (n = 300 embryos per biological replicate for each

genotype). Embryos were triturated using a 200 ul pipette and washed to remove yolks as

previously described (Link et al., 2006) except that E2 embryo medium containing

cycloheximide (15 mM NaCl, 0.7 mM $NaHCO_3$, 0.5 mM KCl, 0.15 mM KH_2PO_4, 2.7 mM $CaCl_2$,

0.5 mM Na_2HPO_4, 1mM $MgSO_4$, 100 ug/ml cycloheximide) was used in lieu of 0.5x Ginzburg

Fish Ringer's buffer. Cell pellets were resuspended in 500 ul of chilled lysis buffer (20 mM Tris

HCl pH 7.5, 30 mM $MgCl_2$, 100 mM NaCl, 0.5 uM DTT, 100 ug/ml cycloheximide, 1 mg/ml

heparin, 2.5 ul/ml NP-40) and incubated on ice for 10 minutes. Lysates were centrifuged at

12,000 x g for 10 minutes at 4°C to fractionate nuclei and debris (pellet) from cytoplasm

(supernatant). After total RNA quantification of cytoplasmic extracts on a Nanodrop

Spectrophotometer (Thermo Fisher), a volume corresponding to 50 ug of total RNA was

transferred onto a 10-50% linear sucrose density gradient per sample. Gradients were

centrifuged at 35,000 x g for 2-3 hours at 4°C. Following centrifugation, gradients were

fractionated into 12, 1 ml fractions using a BioComp Piston Gradient Fractionator and A_{260nm}

profiles were recorded using a BioComp Triax Flow Cell.

RNA extraction of polysome fractions

From each fraction, 400 ul was used for the following procedure. To each fraction, 2 ng

Luciferase Control RNA (Promega) was added as a normalization spike-in control, followed by

600 ul of Trizol and purification following standard procedures (Thermo Fisher). RNA pellets

were rinsed with ice-cold 70% ethanol, briefly dried, and resuspended in 20 ul nuclease-free

H_2O. For each fraction, 8 ul of RNA was reverse transcribed using random primers and

Superscript IV reverse transcriptase following the manufacturer's instructions (Thermo Fisher).

CRISPR-Cas9 mutagenesis of *ddx6* and *ddx61*

CRISPR RNA (crRNA) guide sequences were designed using Integrated DNA Technologies

(IDT) custom design tool

(https://www.idtdna.com/site/order/designtool/index/CRISPR_CUSTOM). crRNA sequences with

the highest on-target scores with no predicted off-target sites were chosen. The impact of

targeted mutations on gene function was maximized by targeting regions within exons encoding

the DEAD box helicase domain (see Fig. S5A-B). Reagents for CRISPR-Cas9 mutagenesis

were prepared as previously described for efficient mutagenesis (Shaw & Mokalled, 2001).

Briefly, crRNAs targeting *ddx6* and *ddx61* (25 uM each, 50 uM total) and trans-activating

CRISPR RNA (tracrRNA) (IDT) (50 uM) were combined equally and annealed to generate dual

guide RNA duplexes (dgRNA), followed by 1:1 dilution in duplex buffer (IDT) for a final

concentration of 25 uM dgRNA. Annealed dgRNAs were next combined equally with 25 uM

Cas9-NLS (QB3 MacroLab, UC Berkeley) that had been diluted from a 40 uM stock solution

using Cas9 dilution buffer B (IDT). Final injection solutions contained 0.1% phenol red dye, 0.2

M potassium chloride, and 5 uM each of dgRNA and Cas9-NLS. Injection solutions were

incubated for 5 minutes at 37°C to facilitate ribonucleoprotein complex (RNP) formation. AB

wild-type embryos were injected at the 1-cell stage with 2 nl of injection solution. crRNA

sequences are listed in Table S6.

Acknowledgements

We thank the Ohio State Zebrafish Facilities staff for excellent zebrafish care, the Ohio

Supercomputer Center for computational resources, the Genomics Shared Resource at the

Ohio State Comprehensive Cancer Center for RNA quality analysis and DNA sequencing, and

the Ohio State Center for RNA Biology for their support and resources. We thank Susan Cole,

Kenneth Poss, Mira Pronobis, and Guramrit Singh for advice and critical comments. We also

thank Anton Blatnik, Geremy Lerma, Ryan Denniston, Lauren Levesque, Benjamin Pastore, and

Zhongxia Yi for advice.

Competing interests

The authors declare no competing or financial interests.

Funding

This work was supported by NIH grants R01GM117964 (to S.L.A.) and R35GM146924 (to

M.G.K.), an OSU Dean's Enrichment Fellowship (to M.C.B.), a Pelotonia Fellowship (to M.C.B.),

an OSU Center for RNA Biology Undergraduate Fellowship (to K.G.T.), and a Cellular,

Molecular, and Biochemical Sciences T32 Training fellowship T32GM141955 (to C.C.A.).

Data and resource availability

RNA-Seq datasets will be made available with repository name and identifier at publication. All

additional data and details of resources can be found within the article and its supplementary

information.

Author contributions

T.L.G., M.C.B., C.C.A, K.G.T, D.M.P, A.M., and S.L.A. performed experiments. T.L.G., M.C.B.,

M.G.K., and S.L.A. analyzed data. All authors contributed intellectually and discussed the data

and manuscript. T.L.G. wrote the manuscript and all authors participated in the editing process.

References

Andrews, S. (2010). FastQC - A quality control tool for high throughput sequence data.

http://www.bioinformatics.babraham.ac.uk/projects/fastqc/. *Babraham Bioinformatics*.

Aulehla, A., Wiegraebe, W., Baubet, V., Wahl, M. B., Deng, C., Taketo, M., Lewandoski, M.

and Pourquié, O. (2008). A beta-catenin gradient links the clock and wavefront systems in

mouse embryo segmentation. *Nat Cell Biol* **10**, 186–193.

Baia Amaral, D., Egidy, R., Perera, A. and Bazzini, A. A. (2024). miR-430 regulates zygotic

mRNA during zebrafish embryogenesis. *Genome Biol* **25**, 74.

Bailey, T. L. (2021). STREME: accurate and versatile sequence motif discovery. *Bioinformatics*

37, 2834–2840.

Bazzini, A. A., Lee, M. T. and Giraldez, A. J. (2012). Ribosome profiling shows that miR-430

reduces translation before causing mRNA decay in zebrafish. *Science* **336**, 233–237.

Bembom, O. and Ivanek, R. (2024). seqLogo: Sequence logos for DNA sequence alignments.

Benjamini, Y., Krieger, A. M. and Yekutieli, D. (2006). Adaptive linear step-up procedures that

control the false discovery rate. *Biometrika* **93**, 491–507.

Bessho, Y., Sakata, R., Komatsu, S., Shiota, K., Yamada, S. and Kageyama, R. (2001).

Dynamic expression and essential functions of Hes7 in somite segmentation. *Genes Dev*

15, 2642–2647.

Blake, L. A., Watkins, L., Liu, Y., Inoue, T. and Wu, B. (2024). A Rapid Inducible RNA Decay

system reveals fast mRNA decay in P-bodies. *Nat Commun* **15**, 2720.

Blatnik, M. C., Gallagher, T. L. and Amacher, S. L. (2023). Keeping development on time:

Insights into post-transcriptional mechanisms driving oscillatory gene expression during

vertebrate segmentation. *Wiley Interdiscip Rev RNA* **14**, e1751–e1751.

Bohn, J. A., Van Etten, J. L., Schagat, T. L., Bowman, B. M., McEachin, R. C., Freddolino,

L. and Goldstrohm, A. C. (2018). Identification of diverse target RNAs that are functionally

regulated by human Pumilio proteins. *Nucleic Acids Res* **46**, 362–386.

Bomsztyk, K., Van Seuningen, I., Suzuki, H., Denisenko, O. and Ostrowski, J. (1997).

Diverse molecular interactions of the hnRNP K protein. *FEBS Lett* **403**, 113–115.

Bonev, B., Stanley, P. and Papalopulu, N. (2012). MicroRNA-9 Modulates Hes1 ultradian

oscillations by forming a double-negative feedback loop. *Cell Rep* **2**, 10–18.

Bonferroni, C. E. (1935). Il Calcolo delle Assicurazioni su Gruppi di Teste. In *Studi in Onore del*

Professore Salvatore Ortu Carboni.

Bonferroni, C. E. (1936). Teoria statistica delle classi e calcolo delle probabilità. *Pubblicazioni*

del R Istituto Superiore di Scienze Economiche e Commerciali di Firenze **8**, 3–62.

Braun, J. E., Huntzinger, E., Fauser, M. and Izaurralde, E. (2011). GW182 Proteins Directly Recruit Cytoplasmic Deadenylase Complexes to miRNA Targets. *Mol Cell* **44**, 120–133.

Bray, N. L., Pimentel, H., Melsted, P. and Pachter, L. (2016). Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* **34**, 525–527.

Briata, P., Forcales, S. V., Ponassi, M., Corte, G., Chen, C.-Y., Karin, M., Puri, P. L. and Gherzi, R. (2005). p38-Dependent Phosphorylation of the mRNA Decay-Promoting Factor KSRP Controls the Stability of Select Myogenic Transcripts. *Mol Cell* **20**, 891–903.

Bulbrook, D., Brazier, H., Mahajan, P., Kliszczak, M., Fedorov, O., Marchese, F. P., Aubareda, A., Chalk, R., Picaud, S., Strain-Damerell, C., et al. (2018). Tryptophan-Mediated Interactions between Tristetraprolin and the CNOT9 Subunit Are Required for CCR4-NOT Deadenylase Complex Recruitment. *J Mol Biol* **430**, 722–736.

Burow, D. A., Umeh-Garcia, M. C., True, M. B., Bakhaj, C. D., Ardell, D. H. and Cleary, M. D. (2015). Dynamic regulation of mRNA decay during neural development. *Neural Dev* **10**, 11.

Cairrao, F., Halees, A. S., Khabar, K. S. A., Morello, D. and Vanzo, N. (2009). AU-rich elements regulate Drosophila gene expression. *Mol Cell Biol* **29**, 2636–2643.

Causier, B., Li, Z., De Smet, R., Lloyd, J. P. B., Van de Peer, Y. and Davies, B. (2017). Conservation of Nonsense-Mediated mRNA Decay Complex Components Throughout Eukaryotic Evolution. *Sci Rep* **7**, 16692.

Chekulaeva, M., Mathys, H., Zipprich, J. T., Attig, J., Colic, M., Parker, R. and Filipowicz, W. (2011). miRNA repression involves GW182-mediated recruitment of CCR4-NOT through conserved W-containing motifs. *Nat Struct Mol Biol* **18**, 1218–1226.

Cho, H., Kim, K. M. and Kim, Y. K. (2009). Human Proline-Rich Nuclear Receptor Coregulatory Protein 2 Mediates an Interaction between mRNA Surveillance Machinery and Decapping Complex. *Mol Cell* **33**, 75–86.

Cho, H., Kim, K. M., Han, S., Choe, J., Park, S. G., Choi, S. S. and Kim, Y. K. (2012). Staufen1-Mediated mRNA Decay Functions in Adipogenesis. *Mol Cell* **46**, 495–506.

Cho, H., Han, S., Choe, J., Park, S. G., Choi, S. S. and Kim, Y. K. (2013). SMG5-PNRC2 is functionally dominant compared with SMG5-SMG7 in mammalian nonsense-mediated mRNA decay. *Nucleic Acids Res* **41**, 1319–1328.

Cho, H., Park, O. H., Park, J., Ryu, I., Kim, J., Ko, J. and Kim, Y. K. (2015). Glucocorticoid receptor interacts with PNRC2 in a ligand-dependent manner to recruit UPF1 for rapid mRNA degradation. *Proc Natl Acad Sci U S A* **112**, E1540–E1549.

Christ, B., Jacob, H. J. and Jacob, M. (1978). On the formation of the myotomes in avian
embryos. An experimental and scanning electron microscope study. *Experientia* **34**, 514–
516.

Christ, B., Jacob, M., Jacob, H.J., Brand, B., Wachtler, F. (1986). Myogenesis: A Problem of
Cell Distribution and Cell Interactions. In: Bellairs, R., Ede, D.A., Lash, J.W. (eds) Somites in
Developing Embryos. NATO ASI Series, vol 118. Springer, Boston, MA.

Ciais, D., Cherradi, N. and Feige, J.-J. (2013). Multiple functions of tristetraprolin/TIS11 RNA-
binding proteins in the regulation of mRNA biogenesis and degradation. *Cell Mol Life Sci* **70**,
2031–2044.

Cooke, A., Prigge, A. and Wickens, M. (2010). Translational repression by deadenylases. *J
Biol Chem* **285**, 28506–28513.

Crippa, S., Nemir, M., Ounzain, S., Ibberson, M., Berthonneche, C., Sarre, A., Boisset, G.,
Maison, D., Harshman, K., Xenarios, I., Diviani, D., Schorderet D., Pedrazzini, T. (2016).
Comparative transcriptome profiling of the injured zebrafish and mouse hearts identifies
miRNA-dependent repair pathways. *Cardiovasc Res* **110**, 73–84.

Cutty, S. J., Fior, R., Henriques, P. M., Saúde, L. and Wardle, F. C. (2012). Identification and
expression analysis of two novel members of the Mesp family in zebrafish. *Int J Dev Biol* **56**,
285–294.

da Silva Pescador, G., Baia Amaral, D., Varberg, J. M., Zhang, Y., Hao, Y., Florens, L. and
Bazzini, A. A. (2024). Protein profiling of zebrafish embryos unmasks regulatory layers
during early embryogenesis. *Cell Rep* **43**, 114769.

Decker, C. J., Teixeira, D. and Parker, R. (2007). Edc3p and a glutamine/asparagine-rich
domain of Lsm4p function in processing body assembly in Saccharomyces cerevisiae. *J Cell
Biol* **179**, 437–449.

Delaune, E. A., François, P., Shih, N. P. and Amacher, S. L. (2012). Single-cell-resolution
imaging of the impact of Notch signaling and mitosis on segmentation clock dynamics. *Dev
Cell* **23**, 995–1005.

Dill, K. K. and Amacher, S. L. (2005). tortuga refines Notch pathway gene expression in the
zebrafish presomitic mesoderm at the post-transcriptional level. *Dev Biol* **287**, 225–236.

Di Stefano, B., Luo, E.-C., Haggerty, C., Aigner, S., Charlton, J., Brumbaugh, J., Ji, F.,
Rabano Jiménez, I., Clowers, K. J., Huebner, A. J., et al. (2019). The RNA Helicase
DDX6 Controls Cellular Plasticity by Modulating P-Body Homeostasis. *Cell Stem Cell* **25**,
622-638.e13.

Ditlev, J. A., Case, L. B. and Rosen, M. K. (2018). Who's In and Who's Out-Compositional
Control of Biomolecular Condensates. *J Mol Biol* **430**, 4666–4684.

Djuranovic, S., Nahvi, A. and Green, R. (2012). miRNA-mediated gene silencing by
translational repression followed by mRNA deadenylation and decay. *Science* **336**, 237–240.

Dunn, O. J. (1961). Multiple Comparisons among Means. *J Am Stat Assoc* **56**, 52–64.

Enwerem, I. I. I., Elrod, N. D., Chang, C.-T., Lin, A., Ji, P., Bohn, J. A., Levdansky, Y.,
Wagner, E. J., Valkov, E. and Goldstrohm, A. C. (2021). Human Pumilio proteins directly
bind the CCR4-NOT deadenylase complex to regulate   the transcriptome. *RNA* **27**, 445–
464.

Eulalio, A., Behm-Ansmant, I., Schweizer, D. and Izaurralde, E. (2007). P-body formation is a
consequence, not the cause, of RNA-mediated gene silencing. *Mol Cell Biol* **27**, 3970–3981.

Fujino, Y., Yamada, K., Sugaya, C., Ooka, Y., Ovara, H., Ban, H., Akama, K., Otosaka, S.,
Kinoshita, H., Yamasu, K., et al. (2018). Deadenylation by the CCR4-NOT complex
contributes to the turnover of *hairy*-related mRNAs in the zebrafish segmentation clock.
FEBS Lett **592**, 3388–3398.

Galgano, A., Forrer, M., Jaskiewicz, L., Kanitz, A., Zavolan, M. and Gerber, A. P. (2008).
Comparative analysis of mRNA targets for human PUF-family proteins suggests extensive
interaction with the miRNA regulatory system. *PLoS One* **3**, e3164–e3164.

Gallagher, T. L., Tietz, K. T., Morrow, Z. T., McCammon, J. M., Goldrich, M. L., Derr, N. L.
and Amacher, S. L. (2017). Pnrc2 regulates 3'UTR-mediated decay of segmentation clock-
associated transcripts during zebrafish segmentation. *Dev Biol* **429**, 225–239.

Gangras, P., Gallagher, T. L., Parthun, M. A., Yi, Z., Patton, R. D., Tietz, K. T., Deans, N. C.,
Bundschuh, R., Amacher, S. L. and Singh, G. (2020). Zebrafish rbm8a and magoh
mutants reveal EJC developmental functions and new 3'UTR intron-containing NMD targets.
PLoS Genet **16**, e1008830–e1008830.

Garcia, J. F. and Parker, R. (2015). MS2 coat proteins bound to yeast mRNAs block 5' to 3'
degradation and trap mRNA decay products: implications for the localization of mRNAs by
MS2-MCP system. *RNA* **21**, 1393–1395.

Garneau, N. L., Wilusz, J. and Wilusz, C. J. (2007). The highways and byways of mRNA
decay. *Nat Rev Mol Cell Biol* **8**, 113–126.

Gentleman, R. C., Carey, V. J., Bates, D. M., Bolstad, B., Dettling, M., Dudoit, S., Ellis, B.,
Gautier, L., Ge, Y., Gentry, J., et al. (2004). Bioconductor: open software development for
computational biology and bioinformatics. *Genome Biol* **5**, R80–R80.

Gerber, A. P., Luschnig, S., Krasnow, M. A., Brown, P. O. and Herschlag, D. (2006).
Genome-wide identification of mRNAs associated with the translational regulator PUMILIO
in Drosophila melanogaster. *Proc Natl Acad Sci U S A* **103**, 4487–4492.

Gillhouse, M., Wagner Nyholm, M., Hikasa, H., Sokol, S. Y. and Grinblat, Y. (2004). Two
Frodo/Dapper homologs are expressed in the developing brain and mesoderm of zebrafish.
Developmental Dynamics **230**, 403–409.

Giraldez, A. J., Cinalli, R. M., Glasner, M. E., Enright, A. J., Thomson, J. M., Baskerville, S.,
Hammond, S. M., Bartel, D. P. and Schier, A. F. (2005). MicroRNAs Regulate Brain
Morphogenesis in Zebrafish. *Science (1979)* **308**, 833–838.

Giraldez, A. J., Mishima, Y., Rihel, J., Grocock, R. J., Van Dongen, S., Inoue, K., Enright, A.
J. and Schier, A. F. (2006). Zebrafish MiR-430 Promotes Deadenylation and Clearance of
Maternal mRNAs. *Science (1979)* **312**, 75–79.

Gomez, C., Özbudak, E. M., Wunderlich, J., Baumann, D., Lewis, J. and Pourquié, O.
(2008). Control of segment number in vertebrate embryos. *Nature* **454**, 335–339.

Gratacós, F. M. and Brewer, G. (2010). The role of AUF1 in regulated mRNA decay. *Wiley*
Interdiscip Rev RNA **1**, 457–473.

Hafner, M., Landthaler, M., Burger, L., Khorshid, M., Hausser, J., Berninger, P., Rothballer,
A., Ascano Jr, M., Jungkamp, A.-C., Munschauer, M., et al. (2010). Transcriptome-wide
identification of RNA-binding protein and microRNA target sites by PAR-CLIP. *Cell* **141**, 129–
141.

Halees, A. S., Hitti, E., Al-Saif, M., Mahmoud, L., Vlasova-St Louis, I. A., Beisang, D. J.,
Bohjanen, P. R. and Khabar, K. (2011). Global assessment of GU-rich regulatory content
and function in the human transcriptome. *RNA Biol* **8**, 681–691.

Hatakeyama, J., Bessho, Y., Katoh, K., Ookawara, S., Fujioka, M., Guillemot, F. and
Kageyama, R. (2004). *Hes*genes regulate size, shape and histogenesis of the nervous
system by control of the timing of neural stem cell differentiation. *Development* **131**, 5539–
5550.

Hau, H. H., Walsh, R. J., Ogilvie, R. L., Williams, D. A., Reilly, C. S. and Bohjanen, P. R.
(2006). Tristetraprolin recruits functional mRNA decay complexes to ARE sequences. *J Cell*
Biochem **100**, 1477–1492.

He, F. and Jacobson, A. (2015). Nonsense-Mediated mRNA Decay: Degradation of Defective
Transcripts Is Only Part of the Story. *Annu Rev Genet* **49**, 339–366.

He, F. and Jacobson, A. (2023). Eukaryotic mRNA decapping factors: molecular mechanisms
and activity. *FEBS J* **290**, 5057–5085.

Heinrich, S., Sidler, C. L., Azzalin, C. M. and Weis, K. (2017). Stem-loop RNA labeling can
affect nuclear and cytoplasmic mRNA processing. *RNA* **23**, 134–141.

Holley, S. A., Geisler, R. and Nüsslein-Volhard, C. (2000). Control of *her1* expression during
zebrafish somitogenesis by a *Delta*-dependent oscillator and an independent wave-front
activity. *Genes Dev* **14**, 1678–1690.

Hoshijima, K., Jurynec, M. J., Klatt Shaw, D., Jacobi, A. M., Behlke, M. A. and Grunwald,
D. J. (2019). Highly Efficient CRISPR-Cas9-Based Methods for Generating Deletion
Mutations and F0 Embryos that Lack Gene Function in Zebrafish. *Dev Cell* **51**, 645-657.e4.

Hu, Y., Xie, S. and Yao, J. (2016). Identification of Novel Reference Genes Suitable for qRT-
PCR Normalization with Respect to the Zebrafish Developmental Stage. *PLoS One* **11**,
e0149277–e0149277.

Hubaud, A. and Pourquié, O. (2014). Signalling dynamics in vertebrate segmentation. *Nat Rev*
Mol Cell Biol **15**, 709–721.

Huber, W., Carey, V. J., Gentleman, R., Anders, S., Carlson, M., Carvalho, B. S., Bravo, H.
C., Davis, S., Gatto, L., Girke, T., et al. (2015). Orchestrating high-throughput genomic
analysis with Bioconductor. *Nat Methods* **12**, 115–121.

Icli, B., Wara, A. K. M., Moslehi, J., Sun, X., Plovie, E., Cahill, M., Marchini, J. F., Schissler,
A., Padera, R. F., Shi, J., Cheng, H., Raghuram, S., Arany, Z., Liao, R., Croce, K.,
MacRae, C., Feinberg, M. (2013). MicroRNA-26a regulates pathological and physiological
angiogenesis by targeting BMP/SMAD1 signaling. *Circ Res* **113**, 1231–1241.

Icli, B., Dorbala, P. and Feinberg, M. W. (2014). An emerging role for the miR-26 family in
cardiovascular disease. *Trends Cardiovasc Med* **24**, 241–248.

Ishibashi, M., Ang, S. L., Shiota, K., Nakanishi, S., Kageyama, R. and Guillemot, F. (1995).
Targeted disruption of mammalian hairy and Enhancer of split homolog-1 (HES-1) leads to
up-regulation of neural helix-loop-helix factors, premature neurogenesis, and severe neural
tube defects. *Genes Dev* **9**, 3136–3148.

Jackson, R. J., Hellen, C. U. T. and Pestova, T. V (2010). The mechanism of eukaryotic
translation initiation and principles of its regulation. *Nat Rev Mol Cell Biol* **11**, 113–127.

Joly, W., Chartier, A., Rojas-Rios, P., Busseau, I. and Simonelig, M. (2013). The CCR4
deadenylase acts with Nanos and Pumilio in the fine-tuning of Mei-P26 expression to
promote germline stem cell self-renewal. *Stem Cell Reports* **1**, 411–424.

Junker, J. P., Noël, E. S., Guryev, V., Peterson, K. A., Shah, G., Huisken, J., McMahon, A.
P., Berezikov, E., Bakkers, J. and van Oudenaarden, A. (2014). Genome-wide RNA
Tomography in the Zebrafish Embryo. *Cell* **159**, 662–675.

Kadyrova, L. Y., Habara, Y., Lee, T. H. and Wharton, R. P. (2007). Translational control of
980   maternal *Cyclin B* mRNA by Nanos in the *Drosophila* germline. *Development* **134**, 1519–
981   1527.

982 **Karousis, E. D., Nasif, S. and Mühlemann, O.** (2016). Nonsense-mediated mRNA decay:
983   novel mechanistic insights and biological impact. *Wiley Interdiscip Rev RNA* **7**, 661–682.

984 **Kervestin, S. and Jacobson, A.** (2012). NMD: a multifaceted response to premature
985   translational termination. *Nat Rev Mol Cell Biol* **13**, 700–712.

986 **Keynes, R. J. and Stern, C. D.** (1988). Mechanisms of vertebrate segmentation. *Development*
987   **103**, 413–429.

988 **Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. and Schilling, T. F.** (1995). Stages
989   of embryonic development of the zebrafish. *Developmental Dynamics* **203**, 253–310.

990 **Klatt Shaw, D. and Mokalled, M. H.** (2021). Efficient CRISPR/Cas9 mutagenesis for
991   neurobehavioral screening in adult zebrafish. *G3 (Bethesda)* **11**, jkab089.

992 **Klett, H., Jürgensen, L., Most, P., Busch, M., Günther, F., Dobreva, G., Leuschner, F.,
993   Hassel, D., Busch, H. and Boerries, M.** (2018). Delineating the Dynamic Transcriptome
994   Response of mRNA and microRNA during Zebrafish Heart Regeneration. *Biomolecules* **9**,
995   11.

996 **Komisarczuk, A. Z., Topp, S., Stigloher, C., Kapsimali, M., Bally-Cuif, L. and Becker, T. S.**
997   (2008). Enhancer detection and developmental expression of zebrafish *sprouty1*, a member
998   of the *fgf8* synexpression group. *Developmental Dynamics* **237**, 2594–2603.

999 **Krol, A. J., Roellig, D., Dequéant, M.-L., Tassy, O., Glynn, E., Hattem, G., Mushegian, A.,
1000   Oates, A. C. and Pourquié, O.** (2011). Evolutionary plasticity of segmentation clock
1001   networks. *Development* **138**, 2783–2792.

1002 **Kusov, Y. Y., Shatirishvili, G., Dzagurov, G. and Gauss-Müller, V.** (2001). A new G-tailing
1003   method for the determination of the poly(A) tail length applied to hepatitis A virus RNA.
1004   *Nucleic Acids Res* **29**, E57-e57.

1005 **Lai, T., Cho, H., Liu, Z., Bowler, M. W., Piao, S., Parker, R., Kim, Y. K. and Song, H.** (2012).
1006   Structural Basis of the PNRC2-Mediated Link between mRNA Surveillance and Decapping.
1007   *Structure* **20**, 2025–2037.

1008 **Lewis, J.** (2003). Autoinhibition with Transcriptional Delay. *Current Biology* **13**, 1398–1408.

1009 **Liu, Y., Zhu, Z., Ho, I. H. T., Shi, Y., Li, J., Wang, X., Chan, M. T. V and Cheng, C. H. K.**
1010   (2020). Genetic Deletion of miR-430 Disrupts Maternal-Zygotic Transition and Embryonic
1011   Body Plan. *Front Genet* **11**, 853.

Lu, G. and Hall, T. M. T. (2011). Alternate modes of cognate RNA recognition by human
PUMILIO proteins. *Structure* **19**, 361–367.

Luo, Y., Na, Z. and Slavoff, S. A. (2018). P-Bodies: Composition, Properties, and Functions.
Biochemistry **57**, 2424–2431.

Lykke-Andersen, J. and Bennett, E. J. (2014). Protecting the proteome: Eukaryotic
cotranslational quality control pathways. *J Cell Biol* **204**, 467–476.

Lykke-Andersen, J. and Wagner, E. (2005). Recruitment and activation of mRNA decay
enzymes by two ARE-mediated decay activation domains in the proteins TTP and BRF-1.
Genes Dev **19**, 351–361.

Makeyev, A. V and Liebhaber, S. A. (2002). The poly(C)-binding proteins: a multiplicity of
functions and a search for mechanisms. *RNA* **8**, 265–278.

Makino, S., Mishima, Y., Inoue, K. and Inada, T. (2015). Roles of mRNA fate modulators Dhh1
and Pat1 in TNRC6-dependent gene silencing recapitulated in yeast. *J Biol Chem* **290**,
8331–8347.

Marnef, A., Sommerville, J. and Ladomery, M. R. (2009). RAP55: Insights into an
evolutionarily conserved protein family. *The International Journal of Biochemistry & Cell*
Biology **41**, 977–981.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet J* **17**, 10.

Martin, G. and Keller, W. (1998). Tailing and 3'-end labeling of RNA with yeast poly(A)
polymerase and various nucleotides. *RNA* **4**, 226-230.

Masamizu, Y., Ohtsuka, T., Takashima, Y., Nagahara, H., Takenaka, Y., Yoshikawa, K.,
Okamura, H. and Kageyama, R. (2006). Real-time imaging of the somite segmentation
clock: revelation of unstable oscillators in the individual presomitic mesoderm cells. *Proc*
Natl Acad Sci U S A **103**, 1313–1318.

Matsuda, M., Hayashi, H., Garcia-Ojalvo, J., Yoshioka-Kobayashi, K., Kageyama, R.,
Yamanaka, Y., Ikeya, M., Toguchida, J., Alev, C. and Ebisuya, M. (2020). Species-specific
segmentation clock periods are due to differential biochemical reaction speeds. *Science*
(1979) **369**, 1450–1455.

Mi, H., Muruganujan, A. and Thomas, P. D. (2013). PANTHER in 2013: modeling the evolution
of gene function, and other gene attributes, in the context of phylogenetic trees. *Nucleic*
Acids Res **41**, D377–D386.

Mishima, Y. and Tomari, Y. (2016). Codon Usage and 3′ UTR Length Determine Maternal
mRNA Stability in Zebrafish. *Mol Cell* **61**, 874–885.

Mishima, Y. and Tomari, Y. (2017). Pervasive yet nonuniform contributions of Dcp2 and Cnot7
to maternal mRNA clearance in zebrafish. *Genes to Cells* **22**, 670–678.

Moreno, T. A., Jappelli, R., Izpisúa Belmonte, J. C. and Kintner, C. (2008). Retinoic acid
regulation of the Mesp-Ripply feedback loop during vertebrate segmental patterning. *Dev*
Biol **315**, 317–330.

Morris, A. R., Mukherjee, N. and Keene, J. D. (2008). Ribonomic analysis of human Pum1
reveals cis-trans conservation across species despite evolution of diverse mRNA target sets.
Mol Cell Biol **28**, 4093–4103.

Mugridge, J. S., Ziemniak, M., Jemielity, J. and Gross, J. D. (2016). Structural basis of
mRNA-cap recognition by Dcp1-Dcp2. *Nat Struct Mol Biol* **23**, 987–994.

Murata, Y. and Wharton, R. P. (1995). Binding of pumilio to maternal hunchback mRNA is
required for posterior patterning in drosophila embryos. *Cell* **80**, 747–756.

Neff, A. T., Lee, J. Y., Wilusz, J., Tian, B. and Wilusz, C. J. (2012). Global analysis reveals
multiple pathways for unique regulation of mRNA decay in induced pluripotent stem cells.
Genome Res **22**, 1457–1467.

Nguyen-Chi, M. E., Bryson-Richardson, R., Sonntag, C., Hall, T. E., Gibson, A., Sztal, T.,
Chua, W., Schilling, T. F. and Currie, P. D. (2012). Morphogenesis and cell fate
determination within the adaxial cell equivalence group of the zebrafish myotome. *PLoS*
Genet **8**, e1003014–e1003014.

Nicholson, P., Gkratsou, A., Josi, C., Colombo, M. and Mühlemann, O. (2018). Dissecting
the functions of SMG5, SMG7, and PNRC2 in nonsense-mediated mRNA decay of human
cells. *RNA* **24**, 557–573.

Oates, A. C. and Ho, R. K. (2002). *Hairy/E(spl)-related*(*Her*) genes are central components of
the segmentation oscillator and display redundancy with the Delta/Notch signaling pathway
in the formation of anterior segmental boundaries in the zebrafish. *Development* **129**, 2929–
2946.

Oates, A. C., Morelli, L. G. and Ares, S. (2012). Patterning embryos with oscillations: structure,
function and dynamics of the vertebrate segmentation clock. *Development* **139**, 625–639.

Ohtsuka, T., Ishibashi, M., Gradwohl, G., Nakanishi, S., Guillemot, F. and Kageyama, R.
(1999). Hes1 and Hes5 as notch effectors in mammalian neuronal differentiation. *EMBO J*
18, 2196–2207.

Ostareck, D. H., Naarmann-de Vries, I. S. and Ostareck-Lederer, A. (2014). DDX6 and its
orthologs as modulators of cellular and viral RNA expression. *WIREs RNA* **5**, 659–678.

Ostareck-lederer, A., Ostareck, D. H. and Hentze, M. W. (1998). Cytoplasmic regulatory

functions of the KH-domain proteins hnRNPs K and E1/E2. *Trends Biochem Sci* **23**, 409–

411.

Palmeirim, I., Henrique, D., Ish-Horowicz, D. and Pourquié, O. (1997). Avian hairy Gene

Expression Identifies a Molecular Clock Linked to Vertebrate Segmentation and

Somitogenesis. *Cell* **91**, 639–648.

Pimentel, H., Bray, N. L., Puente, S., Melsted, P. and Pachter, L. (2017). Differential analysis

of RNA-seq incorporating quantification uncertainty. *Nat Methods* **14**, 687–690.

Pourquié, O. (2011). Vertebrate segmentation: from cyclic gene networks to scoliosis. *Cell* **145**,

650–663.

Presnyak, V. and Coller, J. (2013). The DHH1/RCKp54 family of helicases: an ancient family of

proteins that promote translational silencing. *Biochim Biophys Acta* **1829**, 817–823.

Rabani, M., Pieper, L., Chew, G.-L. and Schier, A. F. (2017). A Massively Parallel Reporter

Assay of 3' UTR Sequences Identifies In Vivo Rules for mRNA Degradation. *Mol Cell* **68**,

1083-1094.e5.

Raghavan, A., Ogilvie, R. L., Reilly, C., Abelson, M. L., Raghavan, S., Vasdewani, J.,

Krathwohl, M. and Bohjanen, P. R. (2002). Genome-wide analysis of mRNA decay in

resting and activated primary human T lymphocytes. *Nucleic Acids Res* **30**, 5529–5538.

Rattenbacher, B., Beisang, D., Wiesner, D. L., Jeschke, J. C., von Hohenberg, M., St

Louis-Vlasova, I. A. and Bohjanen, P. R. (2010). Analysis of CUGBP1 targets identifies

GU-repeat sequences that mediate rapid mRNA decay. *Mol Cell Biol* **30**, 3970–3980.

Salkind, N. (2007). *Encyclopedia of Measurement and Statistics*. Sage Publications, Inc.

Sanduja, S., Blanco, F. F. and Dixon, D. A. (2011). The roles of TTP and BRF proteins in

regulated mRNA decay. *Wiley Interdiscip Rev RNA* **2**, 42–57.

Schneider, T. D. and Stephens, R. M. (1990). Sequence logos: a new way to display

consensus sequences. *Nucleic Acids Res* **18**, 6097–6100.

Schneider, T. D., Stormo, G. D., Gold, L. and Ehrenfeucht, A. (1986). Information content of

binding sites on nucleotide sequences. *J Mol Biol* **188**, 415–431.

Schoenberg, D. R. and Maquat, L. E. (2012). Regulation of cytoplasmic mRNA decay. *Nat Rev*

Genet **13**, 246–259.

Schröter, C. and Oates, A. C. (2010). Segment Number and Axial Identity in a Segmentation

Clock Period Mutant. *Current Biology* **20**, 1254–1258.

Schulte-Merker, S. and Stainier, D. Y. R. (2014). Out with the old, in with the new: reassessing
morpholino knockdowns in light of genome editing technology. *Development* **141**, 3103–
3104.

Schwanhäusser, B., Busse, D., Li, N., Dittmar, G., Schuchhardt, J., Wolf, J., Chen, W. and
Selbach, M. (2011). Global quantification of mammalian gene expression control. *Nature*
473, 337–342.

Sharova, L. V, Sharov, A. A., Nedorezov, T., Piao, Y., Shaik, N. and Ko, M. S. H. (2009).
Database for mRNA half-life of 19 977 genes obtained by DNA microarray analysis of
pluripotent and differentiating mouse embryonic stem cells. *DNA Res* **16**, 45–58.

Sheth, U. and Parker, R. (2003). Decapping and decay of messenger RNA occur in
cytoplasmic processing bodies. *Science* **300**, 805–808.

Sidák, Z. (1967). Rectangular Confidence Regions for the Means of Multivariate Normal
Distributions. *J Am Stat Assoc* **62**, 626–633.

Siegel, D. A., Le Tonqueze, O., Biton, A., Zaitlen, N. and Erle, D. J. (2022). Massively parallel
analysis of human 3' UTRs reveals that AU-rich element length and registration predict
mRNA destabilization. *G3 (Bethesda)* **12**, jkab404.

Simsek, M. F., Chandel, A. S., Saparov, D., Zinani, O. Q. H., Clason, N. and Özbudak, E. M.
(2023). Periodic inhibition of Erk activity drives sequential somite segmentation. *Nature* **613**,
153–159.

Singh, P. P. and Isambert, H. (2020). OHNOLOGS v2: a comprehensive resource for the
genes retained from whole genome duplication in vertebrates. *Nucleic Acids Res* **48**, D724–
D730.

Soto, X., Biga, V., Kursawe, J., Lea, R., Doostdar, P., Thomas, R. and Papalopulu, N.
(2020). Dynamic properties of noise and Her6 levels are optimized by miR-9, allowing the
decoding of the Her6 oscillator. *EMBO J* **39**, e103558–e103558.

Stainier, D. Y. R., Raz, E., Lawson, N. D., Ekker, S. C., Burdine, R. D., Eisen, J. S., Ingham,
P. W., Schulte-Merker, S., Yelon, D., Weinstein, B. M., et al. (2017). Guidelines for
morpholino use in zebrafish. *PLoS Genet* **13**, e1007000–e1007000.

Stoecklin, G., Mayo, T. and Anderson, P. (2006). ARE-mRNA degradation requires the 5'-3'
decay pathway. *EMBO Rep* **7**, 72–77.

Superina, S., Borovina, A. and Ciruna, B. (2014). Analysis of maternal-zygotic ugdh mutants
reveals divergent roles for HSPGs in vertebrate embryogenesis and provides new insight
into the initiation of left–right asymmetry. *Dev Biol* **387**, 154–166.

Tan, S., Ohtsuka, T., González, A. and Kageyama, R. (2012). MicroRNA9 regulates neural stem cell differentiation by controlling Hes1 expression dynamics in the developing brain. *Genes to Cells* **17**, 952–961.

Teixeira, D., Sheth, U., Valencia-Sanchez, M. A., Brengues, M. and Parker, R. (2005). Processing bodies require RNA for assembly and contain nontranslating mRNAs. *RNA* **11**, 371–382.

Thermes, V., Grabher, C., Ristoratore, F., Bourrat, F., Choulika, A., Wittbrodt, J. and Joly, J.-S. (2002). I-SceI meganuclease mediates highly efficient transgenesis in fish. *Mech Dev* **118**, 91–98.

Thisse, B. and Thisse, C. (2005). High Throughput Expression Analysis of ZF-Models Consortium Clones. *ZFIN Direct Data Submission (http://zfin.org)*.

Thisse, B., Pflumio, S., Furthauer, M., Loppin, B., Heyer, V., Degrave, A., Woehl, R., Lux, A., Steffan, T., Charbonnier, X. Q., et al. (2001). Expression of the zebrafish genome during embryogenesis. *ZFIN Direct Data Submission (http://zfin.org)*.

Thomas, P. D., Ebert, D., Muruganujan, A., Mushayahama, T., Albou, L.-P. and Mi, H. (2022). PANTHER: Making genome-scale phylogenetics accessible to all. *Protein Sci* **31**, 8–22.

Tietz, K. T., Gallagher, T. L., Mannings, M. C., Morrow, Z. T., Derr, N. L. and Amacher, S. L. (2020). Pumilio response and AU-rich elements drive rapid decay of Pnrc2-regulated cyclic gene transcripts. *Dev Biol* **462**, 129–140.

Tutucci, E., Vera, M., Biswas, J., Garcia, J., Parker, R. and Singer, R. H. (2018). An improved MS2 system for accurate reporting of the mRNA life cycle. *Nat Methods* **15**, 81–89.

Vejnar, C. E., Abdel Messih, M., Takacs, C. M., Yartseva, V., Oikonomou, P., Christiano, R., Stoeckius, M., Lau, S., Lee, M. T., Beaudoin, J.-D., et al. (2019). Genome wide analysis of 3' UTR sequence elements and proteins regulating mRNA stability during maternal-to-zygotic transition in zebrafish. *Genome Res* **29**, 1100–1114.

Vidya, E. and Duchaine, T. F. (2022). Eukaryotic mRNA Decapping Activation. *Front Genet* **13**, 832547.

Vlasova, I. A., Tahoe, N. M., Fan, D., Larsson, O., Rattenbacher, B., Sternjohn, J. R., Vasdewani, J., Karypis, G., Reilly, C. S., Bitterman, P. B., et al. (2008). Conserved GU-rich elements mediate mRNA decay by binding to CUG-binding protein 1. *Mol Cell* **29**, 263–270.

Waghray, S., Williams, C., Coon, J. J. and Wickens, M. (2015). Xenopus CAF1 requires NOT1-mediated interaction with 4E-T to repress translation in vivo. *RNA* **21**, 1335–1345.

Wang, X., McLachlan, J., Zamore, P. D. and Hall, T. M. T. (2002). Modular Recognition of RNA
by a Human Pumilio-Homology Domain. *Cell* **110**, 501–512.

Wanglar, C., Takahashi, J., Yabe, T. and Takada, S. (2014). Tbx protein level critical for clock-
mediated somite positioning is regulated through interaction between Tbx and Ripply. *PLoS*
One **9**, e107928–e107928.

Watterston, C., Zeng, L., Onabadejo, A. and Childs, S. J. (2019). MicroRNA26 attenuates
vascular smooth muscle maturation via endothelial BMP signalling. *PLoS Genet* **15**,
e1008163–e1008163.

Waxman, J. S. (2005). Regulation of the early expression patterns of the zebrafish Dishevelled-
interacting proteins Dapper1 and Dapper2. *Developmental Dynamics* **233**, 194–200.

Weber, R. and Chang, C.-T. (2024). Human DDX6 regulates translation and decay of
inefficiently translated mRNAs. *Elife* **13**,.

Weidmann, C. A., Raynard, N. A., Blewett, N. H., Van Etten, J. and Goldstrohm, A. C.
(2014). The RNA binding domain of Pumilio antagonizes poly-adenosine binding protein and
accelerates deadenylation. *RNA* **20**, 1298–1319.

Westerfield, M. (2007). The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish (Danio
rerio), 5th Edition. *University of Oregon Press, Eugene (Book)*.

White, E. K., Moore-Jarrett, T. and Ruley, H. E. (2001). PUM2, a novel murine Puf protein,
and its consensus RNA-binding site. *RNA* **7**, 1855-1866.

White, R. J., Collins, J. E., Sealy, I. M., Wali, N., Dooley, C. M., Digby, Z., Stemple, D. L.,
Murphy, D. N., Billis, K., Hourlier, T., et al. (2017). A high-resolution mRNA expression
time course of embryonic development in zebrafish. *Elife* **6**, e30860.

Yabe, T., Hoshijima, K., Yamamoto, T. and Takada, S. (2016). Quadruple zebrafish mutant
reveals different roles of Mesp genes in somite segmentation between mouse and zebrafish.
Development **143**, 2842–2852.

Yang, E., van Nimwegen, E., Zavolan, M., Rajewsky, N., Schroeder, M., Magnasco, M. and
Darnell Jr, J. E. (2003). Decay rates of human mRNAs: correlation with functional
characteristics and sequence attributes. *Genome Res* **13**, 1863–1872.

Yoshioka-Kobayashi, K., Matsumiya, M., Niino, Y., Isomura, A., Kori, H., Miyawaki, A. and
Kageyama, R. (2020). Coupling delay controls synchronized oscillation in the segmentation
clock. *Nature* **580**, 119–123.

Zamore, P. D., Williamson, J. R. and Lehmann, R. (1997). The Pumilio protein binds RNA
through a conserved domain that defines a new class of RNA-binding proteins. *RNA* **3**,
1421-1433.

Zamore, P. D., Bartel, D. P., Lehmann, R. and Williamson, J. R. (1998). The PUMILIO−RNA

Interaction: A Single RNA-Binding Domain Monomer Recognizes a Bipartite Target

Sequence. *Biochemistry* **38**, 596–604.

Zampedri, C., Tinoco-Cuellar, M., Carrillo-Rosas, S., Diaz-Tellez, A., Ramos-Balderas, J.

L., Pelegri, F. and Maldonado, E. (2016). Zebrafish P54 RNA helicases are cytoplasmic

granule residents that are required for development and stress resilience. *Biol Open* **5**,

1473–1484.

Zheng, D., Ezzeddine, N., Chen, C.-Y. A., Zhu, W., He, X. and Shyu, A.-B. (2008).

Deadenylation is prerequisite for P-body formation and mRNA decay in mammalian cells. *J*

Cell Biol **182**, 89–101.

1222

Figure legends

Fig. 1. Overexpressed transcripts in maternal-zygotic *pnrc2*[oz22] (*MZpnrc2*) mutants are

associated with many biological processes and are enriched for destabilizing motifs. (A) Volcano

plot of RNA-Seq results showing transcripts that are overexpressed (q < 0.05, n = 1769) (blue),

under-expressed (i.e. transcripts with reduced expression; q < 0.05, n = 856) (red) or unaffected

(q > 0.05, n = 32,533) (black) in *MZpnrc2* mutant versus wild-type embryos at mid-

segmentation. Segmentation-associated transcripts (see Fig. 2) are indicated on the volcano

plot. (B) Top 30 most-enriched terms within the aspect biological process using GO analysis of

overexpressed genes in *MZpnrc2* mutants. (C-F) Enriched motifs in overexpressed transcript

3'UTRs (E-value < 0.05) identified using the MEME Suite. FC = fold change; FDR = false

discovery rate; ListHit = number of overexpressed genes per GO term.

1234

Fig. 2. Expression patterns of transcripts associated with segmentation or other developmental

processes are disrupted in *MZpnrc2* mutants. (A-U) Wild-type and *MZpnrc2* mutant embryos (n

= 15 per genotype per probe) were raised to mid-segmentation stages (14-18 hpf) and probed

for *rhov* (A, A' vs B, B', RT-qPCR in C), *sp5l* (D, D' vs E, E', RT-qPCR in F), *mespab* (G, G' vs H,

H', RT-qPCR in I), *ripply2* (J, J' vs K, K', RT-qPCR in L), *dact1* (M, M' vs N, N', RT-qPCR in O),

spry1 (P, P' vs Q, Q', RT-qPCR in R), and *spry4* (S, S' vs T, T', RT-qPCR in U). Coloration

reactions for each probe were developed in parallel for identical reaction times for both

genotypes. RT-qPCR was performed separately from in situ experiments using whole embryos

extracts (n = 15 embryos per biological replicate for each genotype). Fold changes are the

mean of biological replicate values derived from the average of two technical replicates. p-

values were calculated using an unpaired t-test. In situs shown are in lateral view (A-T),

magnified in dorsal view to the right of each embryo (A'-T'). Scale bars = 100 um (A-T); dorsal

view = 50 um (A'-T'). hpf = hours post fertilization.

1248

Fig. 3. Overexpressed oscillatory gene transcripts have shortened poly(A) tails, whereas

overexpressed *ddx61* transcripts are substantially polyadenylated in *MZpnrc2* mutants. (A)

Poly(A)-tail (pA-tail) lengths of *her1* transcripts from wild-type and *MZpnrc2* mutant embryos at

mid-segmentation stages (16-17 hpf) were measured by G/I tailing-based RT-PCR (Martin &

Keller, 1998; Kusov et al., 2001), revealing a predominant pA-tail length of ~45 nts (indicated by

the symbol * in red). (B) Lane profile analysis reveals that the *MZpnrc2* mutant embryo pA-tail

length distribution is reduced relative to wild-type embryo pA tail length distribution. (C-Q)

Transcript quantification from wild-type and *MZpnrc2* mutant embryo extracts at mid-

segmentation (16-17 hpf). Total RNA (n = 50 embryos per extract per biological replicate) was

separated into pA-enriched (D, G, J, M, P) and pA-depleted (E, H, K, N, Q) extracts by oligo-dT

bead-based purification (see methods). Fold changes are presented as the mean of 3 biological

replicates with each biological replicate value derived from the average of 2 technical replicates.

Error bars show standard deviation and p-values show results from unpaired t tests. hpf = hours

post fertilization; pA = poly(A) tail PCR; S = gene-specific PCR.

1263

Fig. 4. Accumulated oscillatory gene transcripts are primarily disengaged from ribosomes,

whereas accumulated *ddx61* transcripts are associated with ribosomes and lead to higher

Ddx6/Ddx61 protein levels in *MZpnrc2* mutants. (A) Representative 260 nm absorbance profiles

for wild-type and *MZpnrc2* mutant embryo cytoplasmic extracts fractionated after cycloheximide

treatment and sucrose density gradient centrifugation (n = 300 embryos per genotype at mid-

segmentation stages (16-17 hpf)) (see methods). Regions between dotted lines indicate

fractions containing mRNAs co-sedimenting in the following groups: ribosome/subunit-unbound

mRNAs (mRNP), 40S small ribosomal subunit (40S), 60S large ribosomal subunit (60S),

monosome-bound mRNAs (80S), and polysome-bound mRNAs (polysome). (B-E) Transcript

quantification from wild-type and *MZpnrc2* mutant embryo extracts at mid-segmentation (16-17

hpf) from unbound/subunit, monosome, and polysome fractions. Fold changes are presented as

the mean of 3 biological replicates (n = 300 embryos per genotype per biological replicate) with

each biological replicate value derived from the average of 3 technical replicates. Error bars

show standard deviation and q-values show adjusted p-values from unpaired t tests adjusted

with a two-stage linear step-up Benjamini, Krieger, and Yekutieli false discovery rate (FDR)

procedure (Benjamini et al., 2006). (G) Immunoblot showing Ddx6/Ddx61 protein levels in

extracts from wild-type and *MZpnrc2* mutant embryos with β-actin as a loading control (n = 2

embryo equivalents per lane at mid-segmentation stages (16-17 hpf)). (H) Bar graph showing

relative Ddx6/Ddx61 protein levels in wild-type and *MZpnrc2* mutant embryo extracts calculated

using pixel intensities from lane profile analysis of immunoblot in panel G. Signal intensities

normalized to β-actin. hpf = hours post fertilization; nm = nanometers; mRNP = messenger

ribonucleoprotein particle; S (panel A) = Svedberg unit.

1286

Fig. 5. Induced expression of a dominant-negative deadenylase has no effect in wild-type

embryos but induces segmentation defects in *MZpnrc2* mutant embryos. Wild-type and

MZpnrc2 mutant embryos were injected with 20 pgs *hsp70l:Myc-cnot7DN* plasmid at the 1-cell

stage and raised alongside uninjected clutch-mate controls to ~14 hpf. Injected and uninjected

controls were then heat shocked at 37°C for 30 minutes and raised at standard temperature

(28.5°C) for 6 hours post-heat shock (pHS) with morphological assessment at hourly intervals

(see methods). (A-D) Representative embryos are shown for (A) uninjected wild-type embryos,

n = 36, (B) *hsp70l:Myc-cnot7DN*-injected wild-type embryos, n = 33, (C) uninjected *MZpnrc2*

mutants, n = 62, and (D) *hsp70:Myc-cnot7DN*-injected *MZpnrc2* mutants, n = 58. Arrow in D

indicates an abnormal segment. (E) Bar graph shows the proportion of embryos with normal

and/or abnormal segments, indicating that injected *MZpnrc2* mutants are the only group shown

in A-D with somite defects. (F-G) A subset of injected embryos (n = 15) was reserved as no

heat-shock controls and did not show leaky *Myc-cnot7DN* expression by anti-Myc

immunohistochemistry (F), whereas mosaic expression was observed in injected embryos 6

hours pHS (G) (representative images shown in F-G). P-value was calculated using a Fisher's

Exact test comparing the number of embryos with or without segment defects. hpf = hours post

fertilization; pHS = post-heat shock; DN = dominant negative; scale bar = 300 um.

1304

Fig. 6. *MZpnrc2* mutant larvae are sensitized to depletion of Ddx6 and Ddx61. (A) At 5 dpf,

ddx6;ddx61 double crispants exhibit mild to severe abnormalities that occur at higher frequency

in *MZpnrc2* mutant larvae (representative images shown in A). (B) Bar graph showing proportion

of embryos with phenotypes shown in A, indicating a synergistic effect from loss of *pnrc2*, *ddx6*,

and *ddx61*. Proportions were calculated from the following groups: uninjected WT, n = 112; WT

ddx6;ddx61 crispant, n = 100; uninjected *MZpnrc2*, n = 81; *MZpnrc2 ddx6;ddx61* crispant, n =

47. Adjusted p-values were calculated using a Fisher's Exact test comparing the number of

embryos within each phenotype class (WT, mildly abnormal, strongly abnormal) and corrected

for multiple pairwise comparisons using a Bonferroni adjustment (Bonferroni, 1935; Bonferroni,

1936; Dunn, 1961). (C) Immunoblot showing increased Ddx6/Ddx61 protein levels in *MZpnrc2*

mutants as well as efficient depletion of Ddx61/Ddx6 protein in *ddx6;ddx61* double crispants.

One-half larval equivalents were loaded in each lane with β-actin shown as a loading control.

1317

Fig. 7. Injection of *ddx6;ddx61* CRISPRs does not affect *her1* mRNA levels in wild-type

embryos, but enhances *her1* accumulation in *MZpnrc2* mutant embryos. Wild-type and *MZpnrc2*

mutant embryos were injected with *ddx6;ddx61* CRISPRs and raised to 18 hpf alongside

uninjected clutch-mates as controls. Uninjected and injected embryos were probed for *her1*

expression by in situ hybridization (n = 20 embryos per genotype per condition) and oscillatory

gene transcript levels were quantified by qRT-PCR (n = 10 embryos per genotype per condition;

5 biological replicates for all conditions with biological replicate values derived from the average

of 2 technical replicates each). (A-D') Although *her1* expression in wild-type embryos injected

with *ddx6;ddx61* CRISPRs is indistinguishable from uninjected clutch-mates (A, A' vs B, B'),

her1 expression appears expanded in *MZpnrc2* mutants injected with *ddx6;ddx61* CRISPRs

relative to uninjected *MZpnrc2* mutant clutch-mates (C, C' vs D, D'). Lateral views are shown in

A-D, magnified dorsal views are shown in A'-D'. (E-H) Bar graphs showing RT-qPCR results with

fold changes derived from the mean of 5 biological replicates. Out of four quantified oscillatory

gene transcripts, only *her1* expression is significantly increased in *ddx6;ddx61* crispant *MZpnrc2*

mutants (E vs F-H). In contrast, *her7* expression is modestly, but significantly reduced in

ddx6;ddx61 crispant *MZpnrc2* mutants (F). Oscillatory gene transcript expression is unaffected

in wild-type embryos injected with *ddx6;ddx61* CRISPRs (E-H). Adjusted p-values calculated

using a one-way ANOVA with a Dunn-Šídák correction for multiple comparisons (Šídák,1967;

Abdi, 2007). Scale bars = 100 um (A-D); dorsal view = 50 um (A'-D').

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348   Fig. 1

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360

1361

1362

1363

1364

1365

1366

1367

1368

1369   Fig. 2

1370

1371   Fig. 3

1372

1373

1374

1375

1376

1377

1378

1379

1380

1381

1382

1383

1384

1385

1386

1387

1388

1389 Fig. 4

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408   Fig. 5

1409

1410

1411

1412

1413

1414

1415

1416

1417

1418

1419

1420

1421

1422   Fig. 6

1423

1424

1425

1426

1427

1428

1429

1430

1431

1432

1433

1434

1435

1436

1437

1438

1439

1440

1441

1442

1443

1444

1445

1446

1447

1448

1449   Fig. 7

1450

1451