Thermal cycling-stimulation via nasal inhalation attenuates Aβ$_{25\text{-}35}$-induced cognitive deficits in C57BL/6 mice

Guan-Bo Lin[1,2,†], Hsu-Hsiang Liu[2,3,†], Yu-Yi Kuo[1,2], You-Ming Chen[2,3], Fang-Tzu Hsu[1,2], Yu-Wei Wang[2,3], Yi Kung [2,3], Chien Ching[2,3] and Chih-Yu Chao[1,2,3*]

[1] Department of Physics, Laboratory for Medical Physics & Biomedical Engineering, National Taiwan University, Taipei 106319, Taiwan; gblin@phys.ntu.edu.tw (G.B.L.); hhliu@phys.ntu.edu.tw (H.H.L.); yykuo@phys.ntu.edu.tw (Y.Y.K.); ymchen@phys.ntu.edu.tw (Y.M.C.); tsyr8924503@phys.ntu.edu.tw (F.T.H.); f10245021@phys.ntu.edu.tw (Y.W.W.); a13956@phys.ntu.edu.tw (Y.K.); albertj8808@phys.ntu.edu.tw (C.C.); cychao@phys.ntu.edu.tw (C.Y.C.)
[2] Molecular Imaging Center, National Taiwan University College of Medicine, Taipei 100233, Taiwan
[3] Graduate Institute of Applied Physics, Biophysics Division, National Taiwan University, Taipei 106319, Taiwan

* Correspondence: cychao@phys.ntu.edu.tw
† These authors contributed equally to this work.

ABSTRACT

Alzheimer's disease (AD) continues to pose a significant public health challenge, with current treatments demonstrating limited effectiveness, partly due to the difficulty of delivering therapeutics across the blood-brain barrier (BBB). The nose-to-brain (N-2-B) pathway offers a promising alternative, enabling pharmacological agents to circumvent the BBB. However, to date, no drugs have been successfully implemented in clinical settings for the treatment of AD via this route, underscoring the necessity for additional research in this area. Mild stress is thought to activate intrinsic protective mechanisms against neurodegeneration, but traditional methods of inducing stress often lack both specificity and practicality. To address this limitation, we propose the inhalation of mildly heated air as a form of thermal stimulation, which utilizes the N-2-B pathway to induce mild stress and stimulate cerebral activity. This study employs the method of thermal cycling-hyperthermia (TC-HT) technique [Chao C.Y. et al., U.S. patent 11753634, 2023 & Chao C.Y. et al., U.S. patent Appl. No. 18/864192, 2024] into a new treatment, adapted as thermal cycling-stimulation via nasal inhalation (TCSNI), which provides cyclic stimulation to maintain pathway activity while minimizing thermal injury. In this study, we assessed the health and olfactory effects of TCSNI in C57BL/6 mice and found no adverse effects. In experimental groups administered with β-amyloid (Aβ), TCSNI treatment resulted in significant enhancements in cognitive performance as evidenced by Y-maze and novel object recognition (NOR) assessments, suggesting an improvement in cognitive function. Protein analyses conducted on the hippocampus of the mice indicated a reduction in Aβ accumulation, alongside increased expression of heat shock protein 70 (HSP70) and insulin-degrading enzyme (IDE) expression, as well as elevated levels of phosphorylated Akt (p-Akt). These results suggest that N-2-B-delivered TCSNI effectively modulates protein expression and enhances cognitive function, highlighting its potential for further exploration in AD treatment.

Keywords: Alzheimer's disease; β-amyloid; hyperthermia; nasal inhalation; mild stress

INTRODUCTION

Alzheimer's disease (AD) is a leading cause of dementia and represents an escalating threat to public health [1]. The abnormal accumulation of β-amyloid (Aβ) in the brain is believed to play a significant role in the oxidative stress and damage that contribute to the pathogenesis of AD [2-4]. Furthermore, Aβ aggregation is linked to synaptic dysfunction and neuronal loss, which are critical factors in cognitive decline among AD patients [5,6]. Consequently, Aβ has become a primary target for therapeutic interventions in AD [5-7]. Recently, drugs developed based on the Aβ hypothesis, such as aducanumab and lecanemab, have received FDA approval [8,9]. However, these drug treatments are expensive and carry risks of adverse effects, including brain edema and abnormal bleeding [10,11], which complicate their broader implementation. While pharmacotherapy remains the predominant treatment strategy, the blood-brain barrier (BBB) constrains advancements in new drug development and application [12,13], highlighting the need for further exploration of alternative therapies. The integration of mild stress-based non-pharmacological interventions into healthcare strategies offers a promising opportunity for enhancing health outcomes [14-18]. The design of optimal and well-controlled mild stress stimuli has garnered considerable interest due to their potential to stimulate neuronal cells, improve cognitive function, and possibly extend lifespan, thereby offering beneficial effects in the context of neurodegenerative diseases [19-21]. These benefits are primarily mediated through the principle of hormesis, wherein exposure to low levels of stressors triggers adaptive cellular responses that enhance resilience against neurodegenerative processes [22-25]. Research has indicated that mild stress may upregulate heat shock proteins (HSPs), which are associated with decreased Aβ aggregation [26,27]. Exercise and sauna use are widely recognized as physiological stimuli capable of inducing mild stress in humans [19-21,28,29]. However, despite their general health benefits, these long-term approaches are time-consuming and lack definitive evidence of curative effects [30-33]. Therefore, the exploration of novel and convenient alternative approaches is warranted.

In order to address this issue, it is imperative to explore alternative viable solutions. The nose-to-brain (N-2-B) pathway presents a promising approach. Unlike conventional drug delivery systems that depend on penetration enhancers to facilitate transport across the BBB, the N-2-B pathway allows for drug administration through nasal inhalation, thereby overcoming this limitation [34,35]. Various pharmacological agents, including donepezil and rivastigmine, as well as natural compounds such as quercetin and curcumin, have been investigated in relation to the N-2-B pathway [36,37]. Moreover, the inhalation of insulin has attracted considerable interest in studies pertaining to this pathway [36,38]. Concurrently, aromatherapy is recognized

as a potentially effective complementary or alternative therapy for dementia treatment
[39-41]. Studies indicate that inhaled aromatherapy may alleviate symptoms
associated with dementia by stimulating olfactory pathways and modulating brain
regions involved in emotion and cognition [42-44]. However, despite extensive
investigations into numerous drugs and bioactive compounds, no medication has yet
been successfully administered via this route for the clinical treatment of AD [36],
indicating a need for further exploration of the N-2-B pathway.

While pharmacological treatments and complementary therapies such as
aromatherapy have demonstrated some progress, their overall efficacy remains limited.
Building on the concept of olfactory stimulation and extending beyond traditional
pharmacological approaches, we propose, for the first time, the innovative application
of inhaling mildly heated air as a form of thermal stimulation via the N-2-B pathway
to activate the brain and induce mild stress. Thermal stimulation has long been
recognized for its potential benefits across various medical disciplines [45-47].
Appropriate hyperthermia (HT) has been shown to enhance the expression of heat
shock proteins (HSPs), particularly heat shock protein 70 (HSP70), which is
considered advantageous for improving AD [48-50]. Additionally, studies indicate
that HT can elevate the expression of insulin-degrading enzyme (IDE), which is
thought to mitigate Aβ accumulation [51,52]. Furthermore, thermal stress has been
demonstrated to activate Akt signaling pathways [53,54], and the neuroprotective
effects of phosphorylated Akt (p-Akt) in AD have been extensively discussed [54-57].
Recently, sauna use has gained attention as a thermal stimulation practice for
dementia prevention [30,31,48]; however, while associated with potential cognitive
benefits, its therapeutic efficacy remains inconclusive [30,31]. Moreover, accessibility
issues and safety concerns, particularly for elderly individuals with cardiovascular
risk factors, may limit its practicality [58,59]. In contrast to systemic heating, this
study proposes a novel strategy using the inhalation of slightly heated air to induce
mild stress, with the aim of modulating protein expression and investigating its
potential cognitive benefits. To mitigate the risks associated with continuous heat
exposure, our research team has adapted the previously established method of our
thermal cycling-hyperthermia (TC-HT) technique [Chao C.Y. et al., U.S. patent
11753634, 2023 & Chao C.Y. et al., U.S. patent Appl. No. 18/864192, 2024] into a
new treatment, termed thermal cycling-stimulation via nasal inhalation (TCSNI). By
employing repeated cycles of heated and ambient-temperature air delivery, TCSNI
may alleviate excessive stress and promote sustained activation of HSP-related
signaling pathways.

In the study, we aimed to explore the potential effects of inhaling slightly heated
air as a novel method of mild thermal stimulation via nasal inhalation. We conducted

a comparative analysis of the effects of TCSNI and continuous hyperthermia
stimulation via nasal inhalation (HTSNI), with a particular emphasis on their
respective influences on olfactory function in C57BL/6 mice. Additionally, we
assessed the capacity of TCSNI to enhance cognitive performance, utilizing the
Y-maze and novel object recognition (NOR) tests [60-63] in a mouse model
exhibiting cognitive impairment induced by intracerebroventricular (i.c.v.) injection
of $A\beta_{25-35}$ [64]. Furthermore, we investigated the expression of heat- and
stress-responsive proteins associated with neurodegeneration, including HSP70, IDE,
and phosphorylated Akt (p-Akt). These proteins are known to play a role in
antioxidant defense, promote cell survival, and facilitate the clearance of $A\beta$
accumulation [48-57]. This study aims to contribute new insights into the therapeutic
potential of nasal inhalation-based thermal stimulation and its implications for the
prevention or mitigation of cognitive decline, thereby establishing a foundation for
future applications in neurodegenerative diseases (NDDs).
142
RESULTS
Effects of HTSNI and TCSNI on olfactory function in mice
The objective of this research is to explore a novel hyperthermic stimulation
technique administered through the nasal cavity. Prior to conducting cognitive
assessments in AD mice, we first evaluated the safety of hyperthermic stimulation
approaches, specifically HTSNI and TCSNI, by monitoring potential adverse effects,
as illustrated in Figure 1. Both HTSNI and TCSNI are methods of hyperthermic
stimulation targeting the nasal cavity, necessitating an examination of the olfactory
function in mice after treatments. The buried food test and odor sensitivity test are
common methods for evaluating the olfactory capabilities of mice [65,66]. Our
experimental results indicated that following TCSNI treatment, the latency to find
food (14.0±3.5 sec) did not significantly differ from that of the control group (17.5±
2.8 sec) in the buried food test (Figure 1A). Conversely, the HTSNI group exhibited a
significantly prolonged latency (70.1±4.0 sec) in comparison to both the control and
TCSNI groups. Additionally, in the odor sensitivity assessment, the time spent
sniffing odor stimuli did not differ significantly between the control and TCSNI
groups across the range of odor dilutions. Notably, at a dilution of 10^{-3}, the HTSNI
group exhibited significantly shorter sniffing time (1.34±0.3 sec) compared to the
control (3.5±0.5 sec) and TCSNI (3.0±0.4 sec) groups. Furthermore, at a dilution of
10^{-2}, mice in the HTSNI group spent significantly less time sniffing (1.1±0.3 sec) than
both the control (4.5±0.6 sec) and TCSNI (4.3±0.5 sec) groups, as shown in Figure 1B.
These results suggest that the mice subjected to the HTSNI exhibited diminished
olfactory sensitivity, indicating a reduced capacity to detect and respond to odor cues

at concentrations that remained effective for the control and TCSNI groups. In view
of this, we applied TCSNI in the subsequent evaluations to determine its potential to
enhance recognition in the AD mouse model.

Figure 1. Effects of HTSNI and TCSNI on olfactory detection in mice. Olfactory
performance was assessed using the buried food test and the olfactory habituation test,
as described in the Materials and Methods section. (A) Buried food test: latency to
locate the buried food. (B) Olfactory sensitivity test: response time to odor cues. Data
are presented as mean ± standard error of the mean. Statistical significance was
determined using one-way analysis of variance (ANOVA) followed by Tukey's
post-hoc test. Significance levels between the indicated groups are denoted as $*P <$
0.05, $**P < 0.01$, $***P < 0.001$, while non-significant differences are labeled as ns.

TCSNI attenuates Aβ-induced cognitive impairments in mice
Given that memory impairment is a defining characteristic of AD [67], research
utilizing mouse models commonly incorporates Y-maze and NOR tests to assess
cognitive deficits [60-63]. Spontaneous alternation in the Y-maze serves as a
well-established indicator of short-term spatial working memory [60,61], while the
NOR test is a widely recognized paradigm for evaluating both short-term and
long-term recognition memory [62,63]. In this study, we employed these assessments
to analyze the effects of TCSNI on Aβ-injected mice. As illustrated in Figure 2A,
there was no significant difference in the number of arm entries in the Y-maze test
across all groups, indicating that i.c.v. injection of phosphate-buffered saline (PBS) or
Aβ, along with TCSNI treatment, did not affect the locomotor activity of the mice.
Furthermore, the spontaneous alternation index was significantly diminished in the Aβ
group compared to the control group, decreasing from 64.8±2.1% to 49.4±1.6%.
Importantly, our TCSNI treatment resulted in an increase in the spontaneous
alternation index, suggesting a potential amelioration of Aβ-induced deficits in
short-term spatial working memory, as demonstrated in Figure 2B. In addition, the
results from the short-term and long-term NOR tests provide compelling evidence for
the efficacy of TCSNI in mitigating recognition memory deficits induced by Aβ. As
shown in Figure 2C, the discrimination index in the short-term NOR tests exhibited a
notable reduction in the Aβ group relative to the control group, decreasing from 63.7±
4.5% to 10.5±7.9%. Notably, it was found that our TCSNI treatment resulted in a
significant improvement in the discrimination index, which was recorded at 56.3±
4.9% following the TCSNI treatment. Additionally, the long-term NOR test revealed a
more pronounced memory impairment due to Aβ, with the discrimination index
declining from 48.2±3.1% to 4.3±7.0%. Following TCSNI treatment, the results

revealed that the discrimination index was restored to 43.4±6.3%, as demonstrated in
Figure 2D. Collectively, these findings suggest that TCSNI treatment effectively
alleviates Aβ-induced cognitive deficits, as evidenced by the results from both the
Y-maze and NOR tests.
208
Figure 2. Effect of TCSNI applications on mice's behavioral performances. The
recovery effects of TCSNI treatment on Aβ-induced recognition impairments were
assessed through (A) the total number of total arm entries and (B) the spontaneous
alternation index in the Y-maze test, and (C) the discrimination index with intertrial
intervals of 1 h and (D) 24 h in the NOR maze test. Data are presented as mean ±
standard error of the mean. Statistical significance was determined using one-way
analysis of variance (ANOVA) followed by Tukey's post-hoc test. Significance levels
between the indicated groups are denoted as ***$P < 0.001$, while non-significant
differences are labeled as ns.
218
TCSNI downregulates Aβ accumulation and elevates HSP70, IDE, and p-Akt
levels in the mouse hippocampus
Given that Aβ is a primary target in AD therapeutic strategies [5-7], this study
initially assessed its accumulation in the hippocampus of mice. As shown in Figure
3A, the Aβ-injected group exhibited a 1.50-fold increase in Aβ levels compared to the
control group. Importantly, TCSNI treatment significantly reduced the hippocampal
Aβ levels, decreasing them from 1.50-fold to 0.90-fold of the control group, thereby
demonstrating a substantial inhibitory effect. Furthermore, HSP70, a critical stress
response protein that acts as a molecular chaperone to prevent protein misfolding, has
been implicated in the modulation of the Aβ aggregation process [68-70]. In vivo
studies have indicated that Aβ administration may lead to a slight increase in HSP70
levels, although this did not reach statistical significance [71,72]. In the present study,
HSP70 levels were also evaluated, revealing that the hippocampal HSP70 levels in the
Aβ-injected group were not significantly different from those in the control group.
Notably, it was found that TCSNI treatment resulted in a significant upregulation of
HSP70 levels, reaching 1.57-fold of the control group, as shown in Figure 3B.
Additionally, IDE has long been recognized for its role in Aβ clearance, with
alterations in its activity or expression potentially contributing to Aβ accumulation
and an increased AD risk [51,73,74]. As shown in Figure 3C, Aβ injection in mice led
to a reduction in hippocampal IDE levels to 0.75-fold of the control. This study found
that TCSNI treatment effectively reversed this decline, significantly elevating IDE
expression to 1.45-fold of the control group. Moreover, Akt is known to play a crucial
role in neuronal survival signaling [75,76], and a reduction in Akt activation has been

documented in AD models [55-57,77,78]. Given that Akt is responsive to stress and
thermal stimuli [53,54,79], it warranted further investigation in this study. As
illustrated in Figure 3D, Aβ injection in mice resulted in a significant downregulation
of the p-Akt/t-Akt ratio (0.83-fold), while TCSNI treatment significantly enhanced
this ratio to 1.57-fold compared to the control. Overall, these findings suggest that
TCSNI-induced Aβ removal may be associated with the upregulation of HSP70, IDE,
and the p-Akt/t-Akt ratio in the hippocampi of the examined mice.

Figure 3. Effect of TCSNI applications on the expression levels of Aβ, p-Akt,
HSP70, and IDE in the hippocampus. Representative western blots and the
quantifications of (A) Aβ, (B) HSP70, (C) IDE, and (D) p-Akt are shown to assess the
anti-Aβ effect in the hippocampus. The protein levels of Aβ, HSP70, and IDE were
normalized to β-actin, while p-Akt was normalized to t-Akt. Each relative expression
level was compared to the control and expressed as a fold change relative to the
control. Data are presented as mean ± standard error of the mean. Statistical
significance was determined using one-way ANOVA followed by Tukey's post-hoc
test. Significance levels between the indicated groups are denoted as $*P < 0.05$, $**P <$
0.01, and $***P < 0.001$, while non-significant differences are labeled as ns.

DISCUSSION
AD remains a significant medical challenge globally. The BBB presents a
considerable impediment to the targeted delivery of drugs for treatment of AD [12,13].
In contrast, the N-2-B pathway has emerged as a promising alternative for
circumventing the BBB, with research indicating its potential therapeutic benefits for
brain disorders and brain cancer [34-36]. Additionally, aromatherapy is recognized as
a complementary or alternative therapeutic approach that may offer advantages in the
treatment of dementia [41]. Despite the increasing interest in the N-2-B pathway as a
means to bypass the BBB, no pharmacological agents have yet received clinical
approval for AD treatment via this route [36]. In light of the absence of a definitive
cure for AD and the limited effectiveness of existing pharmacological treatments,
there has been a shift towards exploring alternative strategies [80-82]. The literature
suggests that mild stress stimuli can enhance the functional capacity of organisms and
yield beneficial biological effects on cellular processes [20,83-85]. Moreover,
non-pharmacological interventions, such as exercise and sauna use, which induce
mild stress, are believed to promote human health, potentially extending lifespan and
improving cognitive function [28,29,84,85]. Among these interventions, whole-body
heating through sauna has garnered significant attention, and sauna-like conditions
have been evaluated [28,30,31,86]. However, due to limitations such as time

consumption, practical challenges, and the lack of curative effects, it is imperative to
investigate more effective and safe methods for inducing the desired mild stress.

In the study, we propose a novel methodology using the inhalation of mildly
heated air as a thermal stimulus to induce mild stress responses. This technique
diverges from traditional whole-body heating methods by specifically targeting the
nasal cavity with warm air. However, it is imperative to assess the safety of this
approach prior to conducting further experimental trials. Previous research has
demonstrated that mice can experience thermal collapse when subjected to extreme
heat conditions [87]. In contrast to the aforementioned scenario, which involves the
exposure of the entire body to elevated temperatures, our study focuses on localized
nasal heating through gas delivery. Additionally, the TC-HT method, developed in our
earlier studies, has shown efficacy in both in vitro and in vivo models with minimal
adverse effects, primarily due to its use of intermittent stimulation intervals that help
to mitigate excessive damage [52,54]. Consequently, we assessed the olfactory
function in mice subjected to HTSNI and TCSNI. As illustrated in Figure 1, mice
receiving HTSNI exhibited impaired olfactory function, as evaluated through the
buried food test and the olfactory sensitivity test, whereas the TCSNI-treated group
maintained olfactory function comparable to that of the untreated control group.
These findings suggest that TCSNI is a safer approach compared to HTSNI.

Given that memory impairment is a hallmark of AD [67], research utilizing
mouse models often employs Y-maze and NOR tests to assess memory deficits
[60-63]. Considering that TCSNI preserves olfactory function in mice, it was selected
as the thermal stimulation method for behavioral assessments in mice with cognitive
impairments, thereby minimizing potential sensory disturbances, particularly in light
of the olfactory decline observed with HTSNI. This study explored the ameliorative
effects of TCSNI on Aβ-induced memory impairment and demonstrated that TCSNI
significantly improved the spontaneous alternation index and the ability to recognize
novel objects, as shown in Figure 2. These results indicate that TCSNI has
considerable potential for improving memory deficits and cognitive dysfunction in
vivo in a safe manner.

While traditional continuous heating HT stimulation holds potential to induce
beneficial mild stress, existing literature suggests that inappropriate HT may yield no
advantages [84,85,88] and could contribute to an increased accumulation of reactive
oxygen species (ROS), thereby exacerbating oxidative stress and resulting in cellular
damage [88-90]. In our previous in vitro study, HT was found to be ineffective in
reducing ROS and exhibited less protective effects compared to TC-HT against
cytotoxicity induced by hydrogen peroxide and Aβ [54]. Furthermore, another study
utilizing thermofoil indicated that continuous HT application was less effective than

TC-HT in ameliorating memory impairment in vivo [52]. These findings suggest that
utilizing cyclical thermal stimulation instead of continuous heating may provoke
distinct biological responses [52,54], potentially linked to the activation and saturation
of stimulated protein pathways. For example, in the domain of olfactory research,
prolonged and continuous exposure to specific stimuli can result in olfactory
adaptation or receptor desensitization [91-94], which may influence the efficacy of
aromatherapy interventions. Consequently, the integration of diverse scents or the
implementation of intermittent scent exposure is deemed a critical strategy to
maintain receptor activation and therapeutic effectiveness [95,96]. Similarly, in
thermal stimulation studies, the phenomenon of thermal adaptation has been
examined and identified as a significant factor contributing to the diminished
effectiveness of traditional HT treatments [97,98]. In contrast to HT, TC-HT
technique employs periodic temperature variations, which may help prevent signaling
desensitization and better sustain heat-induced protein responses. Furthermore,
TC-HT methods are characterized by an improved cellular safety profile, potentially
augmenting both the efficacy and practicality of thermal stimulation, thereby
positioning them as a preferable alternative to conventional HT approaches [52,54]. In
this study, while HTSNI was observed to impair olfactory function, the application of
TCSNI did not exhibit such adverse effects and was found to improve memory and
cognition deficits (Figures 1 and 2).

Given that the clearance of toxic Aβ deposits serves as a critical indicator in AD
therapy [99,100], we conducted an investigation into Aβ expression levels within the
hippocampus of mice from a molecular perspective. As illustrated in Figure 3A, the
application of TCSNI resulted in a significant reduction of Aβ expression levels in the
hippocampus. Additionally, we examined several proteins that may play a role in Aβ
clearance. Notably, HSP70, a prominent heat shock protein activated by thermal
stimuli, has garnered considerable attention for its potential to mitigate the
progression of AD [68-70]. HSP70 primarily functions to prevent protein misfolding
and has also been implicated in influencing the aggregation process of Aβ [68-70].
Our findings indicate that TCSNI application significantly elevated HSP70 expression
in the hippocampus of mice, as shown in Figure 3B. Furthermore, HSP70 has been
suggested to upregulate the expression of IDE, a well-known Aβ-degrading enzyme,
thereby facilitating Aβ clearance [68]. Previous studies have demonstrated that IDE is
induced by stress and possesses heat shock-like characteristics [51,52,101]. Moreover,
IDE has been found to regulate the levels of extracellular Aβ through proteolysis
activity, thereby alleviating its neurotoxic effects [102,103]. Research indicates that
diminished IDE activity and expression may correlate with Aβ accumulation and an
elevated risk of AD [51,73,74], suggesting that enhancing IDE levels could represent

a viable therapeutic strategy for AD [74]. Consistent with these findings, our study reveals that TCSNI treatment significantly restores the Aβ-induced reduction in IDE levels within the hippocampus of mice, as shown in Figure 3C.

In addition to IDE and HSP70, Akt emerged as another key protein of interest in our study. Akt is integral to mediating neuronal survival signaling, growth, and synaptic plasticity, all of which are essential for brain development and function [75,76]. A decline in Akt activation has been observed in AD models, whereas an increase in Akt activation has been linked to protective effects in both in vitro and in vivo models [55-57,77,78]. Importantly, Akt is also recognized as a protein responsive to stress and thermal stimulation [53,54,79]. Furthermore, research has indicated that Akt may mediate HSP70 expression, thereby providing neuroprotection [78]. Additionally, it has been suggested that the Akt signaling pathway may ameliorate cognitive impairment in AD by promoting IDE expression and facilitating Aβ degradation [104]. In this study, we detected Akt protein levels in the hippocampus of mice along with HSP70 and IDE proteins. Our results demonstrate that the Aβ-induced decline in p-Akt was restored following TCSNI application, as illustrated in Figure 3D. Collectively, these findings suggest that TCSNI induces mild thermal stress, which may upregulate the expression of HSP70, IDE, and p-Akt in the hippocampus of Aβ-injected mice. This molecular response may contribute to the observed reduction in hippocampal Aβ levels and the cognitive improvements in the treated mice, consistent with the hypothesis of mild stress-induced self-protection.

This study introduces a novel periodic hyperthermic approach, TCSNI, administered through nasal inhalation to induce mild stress as a potential therapeutic strategy for AD. The stress response is anticipated to activate the cellular self-defense mechanisms [105], with the effects being contingent upon the intensity and duration of the stressor, which may lead to cellular adaptation or apoptosis [106]. Prior research has demonstrated that preconditioning cells with non-lethal stress can reduce cellular damage and enhance cell survival [54,107]. The present study proposes an effective method for inducing mild stress in vivo with minimal impact on olfactory function through TCSNI, which has been shown to enhance cognitive performance in Aβ-injected mice, as evidenced by Y-maze and NOR tests. Additionally, TCSNI was found to decrease Aβ expression and significantly upregulate proteins associated with Aβ degradation and cell survival, such as HSP70, IDE, and p-Akt, in the hippocampus of mice subjected to i.c.v. Aβ injection. Further research is necessary to elucidate the role of additional stress-induced proteins, particularly those linked to neuroprotection and Aβ metabolism, and to validate these findings across different species and in humans. Moreover, the potential synergistic effects of TCSNI in conjunction with pharmacological treatments merit further exploration. Overall, this study provides

394   new insights into the development of alternative therapeutic strategies for AD,
395   underscoring the potential of TCSNI as a novel non-invasive intervention.
396

MATERIALS AND METHODS
Experimental animals and housing
C57BL/6 male mice (6–8 weeks old, 22–26 g) were purchased from the National
Laboratory Animal Center (Taipei, Taiwan), and housed at the National Taiwan
University Animal Resources Center. The mice were provided with ad libitum access
to food and water and were maintained at a controlled temperature of $23°C \pm 2°C$,
under a 12:12 h light/dark cycle (lights switched on at 7:00 a.m.). All experimental
procedures adhered to the established guidelines of the Animal Ethical Committee of
National Taiwan University (approval protocol number: NTU-112-EL-00158). The
mice were randomly allocated into two primary groups. The 1st group was subdivided
into control, TCSNI, and HT subgroups, while the 2nd group was categorized into
control, Aβ, and Aβ+TCSNI subgroups.
409

β-amyloid administration
$Aβ_{25-35}$ (Sigma-Aldrich; Merck KGaA, Darmstadt, Germany) was prepared as a
stock solution at a concentration of 1 mg/mL in 0.9% sterile saline solution and was
fibril-aggregated at 37°C for 7 days. On day 0, 10 µL of the aggregated Aβ was
i.c.v.-injected into the brains of the mice in the Aβ and Aβ-TCSNI groups, while the
mice in the control group received an i.c.v. injection of 10 µL PBS.
416

HTSNI and TCSNI applications
This study employed a novel nasal inhalation-based thermal stimulation system.
As illustrated in Figure 4A, the system utilized a water bath method to heat 3%
isoflurane anesthetic gas and regulated the duration of gas heating and cooling
through a timer-controlled solenoid valve. In the HTSNI application, the solenoid
valve was fixed to ensure a consistent release of heated air, maintaining a temperature
of $47.5 \pm 0.5°C$ for 30 min. On the other hand, in the application of TCSNI treatment,
the timer-controlled solenoid valve was programmed to execute a repetitive 10-cycle
process, with each cycle comprising a 3 min heating phase ($47.5 \pm 0.5°C$) followed by
a 1 min cooling phase. The actual air temperature entering the nasal cavity of the mice
was measured and monitored using a K-type thermocouple positioned at the end of
the pipeline, with the measured temperature results presented in Figure 4B.
429

430   **Figure 4. Experimental design and setup of TCSNI.** (A) The TCSNI system is
431   depicted, which provides inhaling mildly heated air for cyclic thermal stimulation. (B)

432 Actual air temperatures entering the mouse's nasal cavity were recorded using a
433 K-type needle thermocouple throughout the exposure durations of TCSNI and HTSNI
434 treatments.
435
Experimental design
For the 1st group, as depicted in Figure 5, the respective thermal application was
employed to the mice of the HTSNI and TCSNI groups on days 7, 10, and 13. Buried
food and odor sensitivity tests were conducted on days 15 and 16, respectively. In the
2nd group, 10 μL Aβ solution was i.c.v.-injected into the brains of mice in the Aβ and
Aβ+TCSNI groups on day 0, while mice in the control group received an i.c.v.
injection of 10 μL PBS. The TCSNI thermal application was applied to the mice in the
various groups on days 7, 10, and 13. The Y-maze test was conducted on day 14,
followed by the NOR tests on days 15 and 16. Following the NOR test, the mice were
weighed and subsequently euthanized for the collection of brain tissues.
446
Figure 5. TCSNI treatment and assessment of animal behavior. (A) In the 1st
group, the thermal applications were administered to the mice in the HTSNI and
TCSNI groups on days 7, 10 and 13. The buried food and odor sensitivity tests were
conducted on days 15 and 16, respectively. (B) In the 2nd group, Aβ solution was
i.c.v.-injected into the mice's brain of the Aβ and Aβ+TCSNI groups at day 0, and
TCSNI application was performed on days 7, 10, and 13. The Y-maze test was
conducted on day 14, followed by the NOR tests on days 15 and 16, respectively.
After the NOR test, the mice were weighed and subsequently euthanized for brain
tissue collection.
456
Buried food test
On day 14, food pellets were removed from the cages, and the mice underwent
an overnight fasting period. The test was performed on day 15, following a 1 h
acclimatization period in the testing room. Each mouse was then individually placed
into a clean cage containing 3 cm deep of clean bedding, where they were allowed to
acclimate for 5 min. This acclimatization aimed to minimize the potential
confounding effects of exploring a novel environment during the test. A small food
pellet, approximately 5mm in diameter, was buried beneath 1 cm of bedding in a
randomly selected corner of the cage, and the mouse was subsequently introduced
into the cage. The site of mice placement and the location where the small pellet was
buried remained consistent throughout the experiment. The latency, defined as the
time taken for the mice to retrieve the small pellet using their forepaws, was recorded
in seconds with a maximum duration of 15 min.

Odor sensitivity test

The odor sensitivity test in mice was conducted on day 16 after 1 h acclimatization in the testing room. Each mouse was individually placed into a clean cage, which was pre-placed with a dry cotton swab without bedding. The mice were allowed to acclimate to the cage for 30 min to reduce the effects of exploring a novel environment during the test. Following this acclimatization, video recording began and lasted for 3 min, during which the dry cotton swab was replaced with a cotton swab soaked in distilled water. After the initial 3 min, the cotton swab was sequentially replaced with diluted solutions of vanilla extract (Flavorganics, Newark, NJ, USA) at concentrations of 1/100,000, 1/10,000, 1/1,000, and 1/100. An increase in sniffing behavior was anticipated as the odor concentration approached the olfactory threshold for the mice. After acquiring all of the olfactory sensitivity and preference videos, a timer was used to time all investigations of the cotton swabs within three min. The investigation time was defined as any period when the mouse's nares are < 1 mm from the cotton swabs. The recorded latency did not include periods when the mouse was not actively investigating, chewing, or licking the swab

Y-maze test

The Y-maze test was employed to evaluate short-term spatial working memory. The apparatus consisted of three identical symmetric arms (40 cm × 13 cm × 6.5 cm) arranged at 120° angles from a central junction. Each mouse, having no prior familiarity with the maze, was individually placed at the center and allowed to freely explore all three arms to assess spontaneous alteration for 8 min. Following each trial, the maze was cleaned with 70% ethanol to eliminate any olfactory cues. Behavioral performance was documented, and the entries into the maze arms were designated as 'A', 'B', and 'C'. An arm entry was defined as the complete entry of the mouse's hind paws into the arm. A correct alternation was characterized by sequential entries into the three distinct arms (i.e., ABC, ACB, BAC, BCA, CAB, and CBA) in overlapping triplet sets. The spontaneous alternation index was calculated using the following formula: (number of correct alternations) / (total number of arm entries − 2) × 100, with the total number of arm entries also serving as an indicator of general locomotor activity.

Novel object recognition test

The NOR test was conducted to evaluate recognition memory in mice during days 14 to 16 of the study. The test was performed in an open field arena (40 cm × 40 cm × 30.5 cm) over three consecutive days. On day 14 (habituation phase), each

mouse was allowed to freely explore the empty arena for 8 min to acclimate to the
testing environment. On day 15 (familiarization phase and short-term recognition
memory test phase), two identical familiar objects (object A) were placed near the
diagonal corners of the arena, allowing the mouse to explore the arena for 8 min.
Following 1 h retention interval in the home cage, the mouse was reintroduced to the
arena for the short-term recognition memory test, during which one of the original
identical familiar objects (object A) was substituted with a novel object (object B).
The mouse's behavioral responses were recorded via an overhead video camera for 8
min. On day 16, to evaluate long-term recognition memory, the mice were again
placed in the arena and allowed to explore the same familiar object from the training
phase (object A) and a new novel object (object C) for 8 min. This experimental
design facilitated the assessment of both short-term and long-term object recognition
memory. The familiar objects were identical in shapes, material compositions, and
colors, while the novel objects exhibited diverse shapes, sizes, and colors in
comparison to the familiar objects. After each assessment in all phases, all olfactory
cues were eliminated using 70% ethanol. Object exploration was quantified as the
time spent by the mouse sniffing or touching the object with the nose and/or forepaws.
The discrimination index was calculated using the formula $(Tn-To) / (Tn+To)$, where
Tn and To represent the exploration times for the novel and familiar objects,
respectively.
528

Collection of brain tissue and preparation of samples for western blot analysis
After NOR tests, the mice were euthanized using carbon dioxide gas, and their
brains were extracted following confirmation of death. The brains were rinsed with
PBS (HyClone; GE Healthcare Life Sciences, Chicago, IL, USA) supplemented with
1% fresh protease inhibitor cocktail (EMD Millipore, Billerica, MA, USA). The
hippocampus tissue was homogenized in 500 µL of ice-cold RIPA lysis buffer
(MilliporeSigma) containing phosphatase and protease inhibitors using a Q125
Sonicator (Qsonica, Newton, CT, USA). The homogenates were then centrifuged at
23,000×g at 4℃ for 30 min. Protein concentration of the samples were determined
using the Bradford protein assay (cat. no. BRA222; BioShop Canada). The protein
samples were subsequently stored at -80℃ for further western blotting analyses.
540

541 **Western blot analysis**
542 20 µg of homogenized hippocampi protein samples were resolved via 10%
543 sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and
544 subsequently transferred to polyvinylidene fluoride (PVDF) membranes
545 (MilliporeSigma, Burlington, MA, USA). Following a blocking step with 5% bovine

serum albumin (BioShop Canada Inc., ON, Canada) in Tris-buffered saline containing 0.1% Tween-20 (TBST) for 1 h at ambient temperature, the membranes were incubated overnight at 4°C with primary antibodies. The specific primary antibodies utilized included those targeting Aβ (cat. no. sc-28365; Santa Cruz Biotechnology, Inc., Dallas, TX, USA), p-Akt (cat. no. 4060; Cell Signaling Technology, Danvers, MA, USA), Akt (cat. no. 9272; Cell Signaling Technology, Inc.), IDE (cat. no. ab133561; Abcam), HSP70 (cat. no. 4872; Cell Signaling Technology, Inc.), and β-actin (cat. no. GTX110564; GeneTex, Irvine, CA, USA), with β-actin serving as the loading control for protein normalization. After washing with TBST, the membranes were incubated with the horseradish peroxidase (HRP)-conjugated secondary antibodies, goat anti-rabbit IgG (HRP) (cat. no. 111-035-003; Jackson ImmunoResearch Laboratories, Inc.) or goat anti-mouse IgG (HRP) (cat. no. GTX213111-01; GeneTex, Inc.). The dilutions of all primary and secondary antibodies were prepared in accordance with the manufacturer's instructions. The membranes were visualized with an enhanced chemiluminescence substrate (Advansta, San Jose, CA, USA) and detected using the Amersham Imager 600 imaging system (AI600; GE Healthcare Life Sciences). The images were analyzed with Image Lab software (Bio-Rad Laboratories, Inc.).

Statistical analysis

For statistical analysis, results were presented as mean ± standard error of the mean (SEM) and were evaluated using one-way analysis of variance (ANOVA) followed by Tukey's post hoc test, conducted with OriginPro 2022 software (version 2022; OriginLab Corporation, Northampton, MA, USA). The P-value of less than 0.05 was considered statistically significant.

AUTHOR CONTRIBUTIONS: Conceptualization, C.Y.C.; methodology, G.B.L., H.H.L. and C.Y.C.; validation, G.B.L., H.H.L., Y.Y.K., Y.M.C., F.T.H., Y.W.W., Y.K., C.C. and C.Y.C.; formal Analysis, G.B.L., H.H.L., Y.Y.K., Y.M.C. and C.Y.C.; investigation, G.B.L., H.H.L. and C.Y.C.; resources, C.Y.C.; data curation, G.B.L., H.H.L., Y.Y.K., Y.M.C., F.T.H., Y.W.W., Y.K., C.C. and C.Y.C.; writing—original draft preparation, G.B.L., H.H.L. and C.Y.C.; writing – review and editing, G.B.L. and C.Y.C.; supervision, C.Y.C.; funding acquisition, C.Y.C. All authors have read and agreed to the published version of the manuscript.

FUNDING: This work was supported by grants from National Science and Technology Council (NSTC 113-2112-M-002-024 and NSTC 112-2112-M-002-033 to C.Y.C.) and Ministry of Science and Technology (MOST 110-2112-M-002-004 to

C.Y.C.) of the Republic of China. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

INSTITUTIONAL REVIEW BOARD STATEMENT: Not applicable.

INFORMED CONSENT STATEMENT: Not applicable.

DATA AVAILABILITY STATEMENT: The data presented in this study are available on request from the corresponding author.

ACKNOWLEDGMENTS: We thank the Animal Resource Center and the Consortium of Integrative Biomedical Science Key Technology at National Taiwan University for their technical support.

ABBREVIATIONS

AD	Alzheimer's disease
Aβ	β-amyloid
N-2-B	nose-to-brain
BBB	blood-brain barrier
HT	hyperthermia
TC-HT	thermal cycling-hyperthermia
TCSNI	thermal cycling-stimulation via nasal inhalation
HTSNI	hyperthermia stimulation via nasal inhalation
i.c.v.	intracerebroventricular
NOR	novel object recognition
HSP70	heat shock protein 70
IDE	insulin-degrading enzyme
p-Akt	phosphorylated Akt
ROS	reactive oxygen species
ANOVA	analysis of variance

REFERENCES

1. Scheltens, P.; De Strooper, B.; Kivipelto, M.; Holstege, H.; Chételat, G.; Teunissen, C.E.; Cummings, J.; van der Flier, W.M. Alzheimer's disease. *Lancet.* **2021**, 397, 1577-1590. doi: 10.1016/S0140-6736(20)32205-4.
2. Sun, X.; Chen, W.D.; Wang, Y.D. β-Amyloid: the key peptide in the pathogenesis of Alzheimer's disease. *Front Pharmacol.* **2015**, 6, 221. doi: 10.3389/fphar.2015.00221.

3. Leuner, K.; Schütt, T.; Kurz, C.; Eckert, S.H.; Schiller, C.; Occhipinti, A.; Mai, S.; Jendrach, M.;
Eckert, G.P.; Kruse, S.E.; et al. Mitochondrion-derived reactive oxygen species lead to enhanced
amyloid beta formation. *Antioxid Redox Signal.* **2012**, 16, 1421-1433. doi: 10.1089/ars.2011.4173.

4. Miranda, S.; Opazo, C.; Larrondo, L.F.; Muñoz, F.J.; Ruiz, F.; Leighton, F.; Inestrosa, N.C. The role
of oxidative stress in the toxicity induced by amyloid beta-peptide in Alzheimer's disease. *Prog*
Neurobiol. **2000**, 62, 633-648. doi: 10.1016/s0301-0082(00)00015-0.

5. Palop, J.J.; Mucke, L. Amyloid-beta-induced neuronal dysfunction in Alzheimer's disease: from
synapses toward neural networks. *Nat Neurosci.* **2010**, 13, 812-818. doi: 10.1038/nn.2583.

6. Reddy, P.H.; Beal, M.F. Amyloid beta, mitochondrial dysfunction and synaptic damage:
implications for cognitive decline in aging and Alzheimer's disease. *Trends Mol Med.* **2008**, 14, 45-53.
doi: 10.1016/j.molmed.2007.12.002.

7. Chen, G.F.; Xu, T.H.; Yan, Y.; Zhou, Y.R.; Jiang, Y.; Melcher, K.; Xu, H.E. Amyloid beta: structure,
biology and structure-based therapeutic development. *Acta Pharmacol Sin.* **2017**, 38, 1205-1235. doi:
10.1038/aps.2017.28.

8. Rahman, A.; Hossen, M.A.; Chowdhury, M.F.I.; Bari, S.; Tamanna, N.; Sultana, S.S.; Haque, S.N.;
Al Masud, A.; Saif-Ur-Rahman, K.M. Aducanumab for the treatment of Alzheimer's disease: a
systematic review. *Psychogeriatrics.* **2023**, 23, 512-522. doi: 10.1111/psyg.12944.

9. Hossain, M.F.; Husna, A.U.; Kharel, M. Use of lecanemab for the treatment of Alzheimer's disease:
A systematic review. *Brain Behav.* **2024**, 14, e3592. doi: 10.1002/brb3.3592.

10. Cummings, J.; Apostolova, L.; Rabinovici, G.D.; Atri, A.; Aisen, P.; Greenberg, S.; Hendrix, S.;
Selkoe, D.; Weiner, M.; Petersen R.C.; et al. Lecanemab: Appropriate Use Recommendations. *J Prev*
Alzheimers Dis. **2023**, 10, 362-377. doi: 10.14283/jpad.2023.30.

11. Fleck, L.M. Alzheimer's and Aducanumab: Unjust Profits and False Hopes. *Hastings Cent Rep.* **2021**,
51, 9-11. doi: 10.1002/hast.1264.

12. Pardridge, W.M. Alzheimer's disease drug development and the problem of the blood-brain
barrier. *Alzheimers Dement.* **2009**, 5, 427-432. doi: 10.1016/j.jalz.2009.06.003.

13. Wu, D.; Chen, Q.; Chen, X.; Han, F.; Chen, Z.; Wang, Y. The blood-brain barrier: structure,
regulation, and drug delivery. *Signal Transduct Target Ther.* **2023**, 8, 217. doi:
10.1038/s41392-023-01481-w.

14. Rattan, S.I. Hormetic modulation of aging and longevity by mild heat stress. *Dose Response.* **2006**, 3,
533-546. doi: 10.2203/dose-response.003.04.008.

15. Schirrmacher, V. Less Can Be More: The Hormesis Theory of Stress Adaptation in the Global
Biosphere and Its Implications. *Biomedicines.* **2021**, 9, 293. doi: 10.3390/biomedicines9030293.

16. Kouda, K.; Iki, M. Beneficial effects of mild stress (hormetic effects): dietary restriction and health. *J*
Physiol Anthropol. **2010**, 29, 127-132. doi: 10.2114/jpa2.29.127.

17. Calabrese, V.; Giordano, J.; Signorile, A.; Laura Ontario, M.; Castorina, S.; De Pasquale, C.; Eckert,
G.; Calabrese, E.J. Major pathogenic mechanisms in vascular dementia: Roles of cellular stress
response and hormesis in neuroprotection. *J Neurosci Res.* **2016**, 94, 1588-1603. doi:
10.1002/jnr.23925.

18. Rattan, S.I.; Demirovic, D. Hormesis can and does work in humans. *Dose Response.* **2009**, 8, 58-63.
doi: 10.2203/dose-response.09-041.Rattan.

19. Adlard, P.A.; Engesser-Cesar, C.; Cotman, C.W. Mild stress facilitates learning and exercise
improves retention in aged mice. *Exp Gerontol.* **2011**, 46, 53-59. doi: 10.1016/j.exger.2010.10.001.

20. Minois, N. Longevity and aging: beneficial effects of exposure to mild stress. *Biogerontology.* **2000**, 1,
15-29. doi: 10.1023/a:1010085823990.

21. Parihar, V.K.; Hattiangady, B.; Kuruba, R.; Shuai, B.; Shetty, A.K. Predictable chronic mild stress
improves mood, hippocampal neurogenesis and memory. *Mol Psychiatry.* **2011**, 16, 171-183. doi:
10.1038/mp.2009.130.

22. Mattson, M.P. Hormesis defined. *Ageing Res Rev.* **2008**, 7, 1-7. doi: 10.1016/j.arr.2007.08.007.

23. Mao, L.; Franke, J. Hormesis in aging and neurodegeneration-a prodigy awaiting dissection. *Int J Mol Sci*. **2013**, 14, 13109-13128. doi: 10.3390/ijms140713109.

24. Wang, G. Hormesis, cell death, and regenerative medicine for neurode-generative diseases. *Dose Response*. **2012**, 11, 238-254. doi: 10.2203/dose-response.12-019.Wang.

25. Mattson, M.P.; Leak, R.K. The hormesis principle of neuroplasticity and neuroprotection. *Cell Metab*. **2024**, 36, 315-337. doi: 10.1016/j.cmet.2023.12.022.

26. Mane, N.R.; Gajare, K.A.; Deshmukh, A.A. Mild heat stress induces hormetic effects in protecting the primary culture of mouse prefrontal cerebrocortical neurons from neuropathological alterations. *IBRO Rep*. **2018**, 5, 110-115. doi: 10.1016/j.ibror.2018.11.002.

27. Smith Sonneborn, J. Alternative strategy for Alzheimer's disease: stress response triggers. *Int J Alzheimers Dis*. **2012**, 2012, 684283. doi: 10.1155/2012/684283.

28. Patrick, R.P.; Johnson, T.L. Sauna use as a lifestyle practice to extend healthspan. *Exp Gerontol*. **2021**, 154, 111509. doi: 10.1016/j.exger.2021.111509.

29. Ji, L.L. Exercise-induced modulation of antioxidant defense. *Ann N Y Acad Sci*. **2002**, 959, 82-92. doi: 10.1111/j.1749-6632.2002.tb02085.x.

30. Knekt, P.; Järvinen, R.; Rissanen, H.; Heliövaara, M.; Aromaa, A. Does sauna bathing protect against dementia? *Prev Med Rep*. **2020**, 20, 101221. doi: 10.1016/j.pmedr.2020.101221.

31. Laukkanen, T.; Kunutsor, S.; Kauhanen, J.; Laukkanen, J.A. Sauna bathing is inversely associated with dementia and Alzheimer's disease in middle-aged Finnish men. *Age Ageing*. **2017**, 46, 245-249. doi: 10.1093/ageing/afw212.

32. Öhman, H.; Savikko, N.; Strandberg, T.E.; Kautiainen, H.; Raivio, M.M.; Laakkonen, M.L.; Tilvis, R.; Pitkälä, K.H. Effects of Exercise on Cognition: The Finnish Alzheimer Disease Exercise Trial: A Randomized, Controlled Trial. *J Am Geriatr Soc*. **2016**, 64, 731-738. doi: 10.1111/jgs.14059.

33. Valenzuela, P.L.; Castillo-García, A.; Morales, J.S.; de la Villa, P.; Hampel, H.; Emanuele, E.; Lista, S.; Lucia, A. Exercise benefits on Alzheimer's disease: State-of-the-science. *Ageing Res Rev*. **2020**, 62, 101108. doi: 10.1016/j.arr.2020.101108.

34. Pardeshi, C.V.; Belgamwar, V.S. Direct nose to brain drug delivery via integrated nerve pathways bypassing the blood-brain barrier: an excellent platform for brain targeting. *Expert Opin Drug Deliv*. **2013**, 10, 957-972. doi: 10.1517/17425247.2013.790887.

35. Patharapankal, E.J.; Ajiboye, A.L.; Mattern, C.; Trivedi, V. Nose-to-Brain (N2B) Delivery: An Alternative Route for the Delivery of Biologics in the Management and Treatment of Central Nervous System Disorders. *Pharmaceutics*. **2023**, 16, 66. doi: 10.3390/pharmaceutics16010066.

36. Agrawal, M.; Saraf, S.; Saraf, S.; Antimisiaris, S.G.; Chougule, M.B.; Shoyele, S.A.; Alexander, A. Nose-to-brain drug delivery: An update on clinical challenges and progress towards approval of anti-Alzheimer drugs. *J Control Release*. **2018**, 281, 139-177. doi: 10.1016/j.jconrel.2018.05.011.

37. Alexander, A.; Saraf, S. Nose-to-brain drug delivery approach: A key to easily accessing the brain for the treatment of Alzheimer's disease. *Neural Regen Res*. **2018**, 13, 2102-2104. doi: 10.4103/1673-5374.241458.

38. Wong, C.Y.J.; Baldelli, A.; Hoyos, C.M.; Tietz, O.; Ong, H.X.; Traini, D. Insulin Delivery to the Brain via the Nasal Route: Unraveling the Potential for Alzheimer's Disease Therapy. *Drug Deliv Transl Res*. **2024**, 14, 1776-1793. doi: 10.1007/s13346-024-01558-1.

39. Li, B.S.Y.; Chan, C.W.H.; Li, M.; Wong, I.K.Y.; Yu, Y.H.U. Effectiveness and Safety of Aromatherapy in Managing Behavioral and Psychological Symptoms of Dementia: A Mixed-Methods Systematic Review. *Dement Geriatr Cogn Dis Extra*. **2021**, 11, 273-297. doi: 10.1159/000519915.

40. Cha, H.; Kim, S.; Seo, M.S.; Kim, H.S. Effects of olfactory stimulation on cognitive function and behavior problems in older adults with dementia: A systematic literature review. *Geriatr Nurs*. **2021**, 42, 1210-1217. doi: 10.1016/j.gerinurse.2021.07.003.

41. Vora, L.K., Gholap, A.D.; Hatvate, N.T.; Naren, P.; Khan, S.; Chavda, V.P.; Balar, P.C.; Gandhi, J.;
Khatri, D.K. Essential oils for clinical aromatherapy: A comprehensive review. *J Ethnopharmacol.*
2024, 330,118180. doi: 10.1016/j.jep.2024.118180.

42. Cioanca, O.; Hritcu, L.; Mihasan, M.; Hancianu, M. Cognitive-enhancing and antioxidant activities
of inhaled coriander volatile oil in amyloid β(1-42) rat model of Alzheimer's disease. *Physiol Behav.*
2013, 120, 193-202. doi: 10.1016/j.physbeh.2013.08.006.

43. Satou, T.; Hanashima, Y.; Mizutani, I.; Koike, K.   The effect of inhalation of essential oil from
Rosmarinus officinalis on scopolamine-induced Alzheimer's type dementia model mice. *Flavour*
and fragrance journal. **2018**,   33, 230-234. doi: 10.1002/ffj.3435

44. Okuda, M.; Fujita, Y.; Takada-Takatori, Y.; Sugimoto, H.; Urakami, K. Aromatherapy improves
cognitive dysfunction in senescence-accelerated mouse prone 8 by reducing the level of amyloid
beta and tau phosphorylation. *PLoS One.* **2020**, 15, e0240378. doi: 10.1371/journal.pone.0240378.

45. Habash, R.W.; Bansal, R.; Krewski, D.; Alhafid, H.T. Thermal therapy, part 1: an introduction to
thermal therapy. *Crit Rev Biomed Eng.* **2006**, 34, 459-489. doi: 10.1615/critrevbiomedeng.v34.i6.20.

46. Li, L.; Zhang, X.; Zhou, J.; Zhang, L.; Xue, J.; Tao, W. Non-Invasive Thermal Therapy for Tissue
Engineering and Regenerative Medicine. *Small.* **2022**, 18, e2107705. doi: 10.1002/smll.202107705.

47. Biro, S.; Masuda, A.; Kihara, T.; Tei, C. Clinical implications of thermal therapy in lifestyle-related
diseases. *Exp Biol Med (Maywood).* **2003**, 228, 1245-1249. doi: 10.1177/153537020322801023.

48. Hunt, A.P.; Minett, G.M.; Gibson, O.R.; Kerr, G.K.; Stewart, I.B. Could Heat Therapy Be an
Effective Treatment for Alzheimer's and Parkinson's Diseases? A Narrative Review. *Front Physiol.*
2020, 10, 1556. doi: 10.3389/fphys.2019.01556.

49. Von Schulze, A.T.; Deng, F.; Morris, J.K.; Geiger, P.C. Heat therapy: possible benefits for cognitive
function and the aging brain. *J Appl Physiol (1985).* **2020**, 129, 1468-1476. doi:
10.1152/japplphysiol.00168.2020.

50. Smadja, D.M.; Abreu, M.M. Hyperthermia and targeting heat shock proteins: innovative
approaches for neurodegenerative disorders and Long COVID. *Front Neurosci.* **2025**, 19, 1475376.
doi: 10.3389/fnins.2025.1475376.

51. Tundo, G.R.; Sbardella, D.; Ciaccio, C.; Grasso, G.; Gioia, M.; Coletta, A.; Polticelli, F.; Di Pierro, D.;
Milardi, D.; Van Endert, P.; et al. Multiple functions of insulin-degrading enzyme: a metabolic
crosslight? *Crit Rev Biochem Mol Biol.* **2017**, 52, 554-582. doi: 10.1080/10409238.2017.1337707.

52. Kuo, Y.Y.; Chen, W.T.; Lin, G.B.; Chen, Y.M.; Liu, H.H.; Chao, C.Y. Thermal cycling-hyperthermia
ameliorates Aβ25-35-induced cognitive impairment in C57BL/6 mice. *Neurosci Lett.* **2023**, 810,
137337. doi: 10.1016/j.neulet.2023.137337.

53. Mearow, K.M.; Dodge, M.E.; Rahimtula, M.; Yegappan, C. Stress-mediated signaling in PC12 cells
- the role of the small heat shock protein, Hsp27, and Akt in protecting cells from heat stress and
nerve growth factor withdrawal. *J Neurochem.* **2002**, 83, 452-462. doi:
10.1046/j.1471-4159.2002.01151.x.

54. Chen, W.T.; Kuo, Y.Y.; Lin, G.B.; Lu, C.H.; Hsu, H.P.; Sun, Y.K.; Chao, C.Y. Thermal cycling
protects SH-SY5Y cells against hydrogen peroxide and β-amyloid-induced cell injury through
stress response mechanisms involving Akt pathway. *PLoS One.* **2020**, 15, e0240022. doi:
10.1371/journal.pone.0240022.

55. Yin, G.; Li, L.Y.; Qu, M.; Luo, H.B.; Wang, J.Z.; Zhou, X.W. Upregulation of AKT attenuates
amyloid-β-induced cell apoptosis. *J Alzheimers Dis.* **2011**, 25, 337-345. doi:
10.3233/JAD-2011-110104.

56. Ghasemi, R.; Moosavi, M.; Zarifkar, A.; Rastegar, K.; Maghsoudi, N. The Interplay of Akt and ERK
in Aβ Toxicity and Insulin-Mediated Protection in Primary Hippocampal Cell Culture. *J Mol*
Neurosci. **2015**, 57, 325-334. doi: 10.1007/s12031-015-0622-6.

57. Kommaddi, R.P.; Gowaikar, R.; Haseena, P.A.; Diwakar, L.; Singh, K.; Mondal, A. Akt activation
ameliorates deficits in hippocampal-dependent memory and activity-dependent synaptic protein

synthesis in an Alzheimer's disease mouse model. *J Biol Chem*. **2024**, 300, 105619. doi:
10.1016/j.jbc.2023.105619.

58. Hannuksela, M.L.; Ellahham, S. Benefits and risks of sauna bathing. *Am J Med*. **2001**, 110, 118-126.
770 doi: 10.1016/s0002-9343(00)00671-9.

59. Bain, A.R.; Nybo, L.; Ainslie, P.N. Cerebral Vascular Control and Metabolism in Heat Stress. *Compr*
772 *Physiol*. **2015**, 5, 1345-1380. doi: 10.1002/cphy.c140066.

60. Prieur, E.A.K.; Jadavji, N.M. Assessing Spatial Working Memory Using the Spontaneous
774 Alternation Y-maze Test in Aged Male Mice. *Bio Protoc.* **2019**, 9, e3162. doi:
775 10.21769/BioProtoc.3162.

61. Kraeuter, A.K.; Guest, P.C.; Sarnyai, Z. The Y-Maze for Assessment of Spatial Working and
777 Reference Memory in Mice. *Methods Mol Biol*. **2019**, 1916, 105-111. doi:
778 10.1007/978-1-4939-8994-2_10.

62. Lueptow, L.M. Novel Object Recognition Test for the Investigation of Learning and Memory in
780 Mice. *J Vis Exp*. **2017**, 126, 55718. doi: 10.3791/55718.

63. Navarro, D.; Gasparyan, A.; Martí Martínez, S.; Díaz Marín, C.; Navarrete, F.; García Gutiérrez,
782 M.S.; Manzanares, J. Methods to Identify Cognitive Alterations from Animals to Humans: A
783 Translational Approach. *Int J Mol Sci*. **2023**, 24, 7653. doi: 10.3390/ijms24087653.

64. Kim, H.Y.; Lee, D.K.; Chung, B.R.; Kim, H.V.; Kim, Y. Intracerebroventricular Injection of
785 Amyloid-β Peptides in Normal Mice to Acutely Induce Alzheimer-like Cognitive Deficits. *J Vis Exp.*
786 **2016**, 109, 53308. doi: 10.3791/53308.

65. Yang, M.; Crawley, J.N. Simple behavioral assessment of mouse olfaction. *Curr Protoc Neurosci*.
788 **2009**, 48, 8.24. doi: 10.1002/0471142301.ns0824s48.

66. Zhang, J.; Zhao, Z.; Sun, S.; Li, J.; Wang, Y.; Dong, J.; Yang, S.; Lou, Y.; Yang, J.; Li, W.; Li, S.
790 Olfactory Evaluation in Alzheimer's Disease Model Mice. *Brain Sci*. **2022**, 12, 607. doi:
791 10.3390/brainsci12050607.

67. Jahn, H. Memory loss in Alzheimer's disease. *Dialogues Clin Neurosci*. **2013**, 15, 445-454. doi:
793 10.31887/DCNS.2013.15.4/hjahn.

68. Lu, R.C.; Tan, M.S.; Wang, H.; Xie, A.M.; Yu, J.T.; Tan, L. Heat shock protein 70 in Alzheimer's
795 disease. *Biomed Res Int*. **2014**, 2014, 435203. doi: 10.1155/2014/435203.

69. Rivera, I.; Capone, R.; Cauvi, D.M.; Arispe, N.; De Maio, A. Modulation of Alzheimer's amyloid β
797 peptide oligomerization and toxicity by extracellular Hsp70. *Cell Stress Chaperones*. **2018**, 23,
798 269-279. doi: 10.1007/s12192-017-0839-0.

70. Valle-Medina, A.; Calzada-Mendoza, C.C.; Ocharan-Hernández, M.E.; Jiménez-Zamarripa, C.A.;
800 Juárez-Cedillo, T. Heat shock protein 70 in Alzheimer's disease and other dementias: A possible
801 alternative therapeutic. *J Alzheimers Dis Rep*. **2025**, 9, 25424823241307021.
802 doi:10.1177/25424823241307021

71. Abdi, A.; Sadraie, H.; Dargahi, L.; Khalaj, L.; Ahmadiani, A. Apoptosis inhibition can be
804 threatening in Aβ-induced neuroinflammation, through promoting cell proliferation. *Neurochem*
805 *Res*. **2011**, 36, 39-48. doi: 10.1007/s11064-010-0259-3.

72. Hoshino, T.; Suzuki, K.; Matsushima, T.; Yamakawa, N.; Suzuki, T.; Mizushima, T. Suppression of
807 Alzheimer's disease-related phenotypes by geranylgeranylacetone in mice. *PLoS One*. **2013**, 8,
808 e76306. doi: 10.1371/journal.pone.0076306.

73. Cook, D.G.; Leverenz, J.B.; McMillan, P.J.; Kulstad, J.J.; Ericksen, S.; Roth, R.A.; Schellenberg, G.D.;
810 Jin, L.W.; Kovacina, K.S. Craft, S. Reduced hippocampal insulin-degrading enzyme in late-onset
811 Alzheimer's disease is associated with the apolipoprotein E-epsilon4 allele. *Am J Pathol*. **2003**, 162,
812 313-319. doi: 10.1016/s0002-9440(10)63822-9.

74. Zhao, L.; Teter, B.; Morihara, T.; Lim, G.P.; Ambegaokar, S.S.; Ubeda, O.J.; Frautschy, S.A.; Cole,
G.M. Insulin-degrading enzyme as a downstream target of insulin receptor signaling cascade:

implications for Alzheimer's disease intervention. *J Neurosci*. **2004**, 24, 11120-11126. doi:
10.1523/JNEUROSCI.2860-04.2004.

75. Ahn, J.Y. Neuroprotection signaling of nuclear akt in neuronal cells. *Exp Neurobiol*. **2014**, 23,
200-206. doi: 10.5607/en.2014.23.3.200.

76. Manning, B.D.; Toker, A. AKT/PKB Signaling: Navigating the Network. *Cell*. **2017**, 169, 381-405.
doi: 10.1016/j.cell.2017.04.001.

77. Cui, W.; Tao, J.; Wang, Z.; Ren, M.; Zhang, Y.; Sun, Y.; Peng, Y.; Li, R. Neuregulin1beta1
antagonizes apoptosis via ErbB4-dependent activation of PI3-kinase/Akt in APP/PS1 transgenic
mice. *Neurochem Res*. **2013**, 38, 2237-2246. doi: 10.1007/s11064-013-1131-z.

78. Magrané, J.; Rosen, K.M.; Smith, R.C.; Walsh, K.; Gouras, G.K.; Querfurth, H.W. Intraneuronal
beta-amyloid expression downregulates the Akt survival pathway and blunts the stress response. *J
Neurosci*. **2005**, 25, 10960-10969. doi: 10.1523/JNEUROSCI.1723-05.2005.

79. Maroni, P.; Bendinelli, P.; Tiberio, L.; Rovetta, F.; Piccoletti, R.; Schiaffonati, L. In vivo heat-shock
response in the brain: signalling pathway and transcription factor activation. *Brain Res Mol Brain
Res*. **2003**, 119, 90-99. doi: 10.1016/j.molbrainres.2003.08.018.

80. Jensen, C.S.; Hasselbalch, S.G.; Waldemar, G.; Simonsen, A.H. Biochemical Markers of Physical
Exercise on Mild Cognitive Impairment and Dementia: Systematic Review and Perspectives. *Front
Neurol*. **2015**, 6, 187. doi: 10.3389/fneur.2015.00187.

81. Wu, W.; Ji, Y.; Wang, Z.; Wu, X.; Li, J.; Gu, F.; Chen, Z.; Wang, Z. The FDA-approved
anti-amyloid-β monoclonal antibodies for the treatment of Alzheimer's disease: a systematic
review and meta-analysis of randomized controlled trials. *Eur J Med Res*. **2023**, 28, 544. doi:
10.1186/s40001-023-01512-w.

82. Ornish, D.; Madison, C.; Kivipelto, M.; Kemp, C.; McCulloch, C.E.; Galasko, D.; Artz, J.; Rentz, D.;
Lin, J.; Norman, K.; et al. Effects of intensive lifestyle changes on the progression of mild cognitive
impairment or early dementia due to Alzheimer's disease: a randomized, controlled clinical trial.
Alzheimers Res Ther. **2024**, 16, 122. doi: 10.1186/s13195-024-01482-z.

83. Matsumoto, K.; Honda, K.; Kobayashi, N. Protective effect of heat preconditioning of rat liver graft
resulting in improved transplant survival. *Transplantation*. **2001**, 71, 862-868. doi:
10.1097/00007890-200104150-00007.

84. Le Bourg, É. Hormesis, aging and longevity. *Biochim Biophys Acta*. **2009**, 1790, 1030-1039. doi:
10.1016/j.bbagen.2009.01.004.

85. Le Bourg, É. Characterisation of the positive effects of mild stress on ageing and resistance to stress.
Biogerontology. **2020**, 21, 485-493. doi: 10.1007/s10522-020-09870-2.

86. Guisle, I.; Canet, G.; Pétry, S.; Fereydouni-Forouzandeh, P.; Morin, F.; Kérauden, R.; Whittington,
R.A.; Calon, F.; Hébert, S.S.; Planel, E. Sauna-like conditions or menthol treatment reduce tau
phosphorylation through mild hyperthermia. *Neurobiol Aging*. **2022**, 113, 118-130. doi:
10.1016/j.neurobiolaging.2022.02.011.

87. Crane, C.R.; Sanders, D.C. Inhalation Toxicology: VIII. Establishing Heat Tolerance Limits for Rats
and Mice Subjected to Acute Exposures at Elevated Air Temperatures. *Civil Aerospace Medical
Institute*. **1986**, DOT/FAA/AM-86/8.

88. Kassis, S.; Grondin, M.; Averill-Bates, D.A. Heat shock increases levels of reactive oxygen species,
autophagy and apoptosis. *Biochim Biophys Acta Mol Cell Res*. **2021**, 1868, 118924. doi:
10.1016/j.bbamcr.2020.118924.

89. Slimen, I.B.; Najar, T.; Ghram, A.; Dabbebi, H.; Ben Mrad, M.; Abdrabbah, M. Reactive oxygen
species, heat stress and oxidative-induced mitochondrial damage. A review. *Int J Hyperthermia*.
2014, 30, 513-523. doi: 10.3109/02656736.2014.971446.

90. Li, L.; Tan, H.; Yang, H.; Li, F.; He, X.; Gu, Z.; Zhao, M.; Su, L. Reactive oxygen species mediate
heat stress-induced apoptosis via ERK dephosphorylation and Bcl-2 ubiquitination in human
umbilical vein endothelial cells. *Oncotarget*. **2017**, 8, 12902-12916. doi: 10.18632/oncotarget.14186.

91.   Dalton, P.; Wysocki, C.J. The nature and duration of adaptation following long-term odor
exposure. *Percept Psychophys*. **1996**, 58, 781-792. doi: 10.3758/bf03213109.
92.   Kato, A.; Touhara, K. Mammalian olfactory receptors: pharmacology, G protein coupling and
desensitization. *Cell Mol Life Sci*. **2009**, 66, 3743-3753. doi: 10.1007/s00018-009-0111-6. Epub 2009
Aug 4
93.   Pellegrino, R.; Sinding, C.; de Wijk, R.A.; Hummel, T. Habituation and adaptation to odors in
humans. *Physiol Behav*. **2017**, 177, 13-19. doi: 10.1016/j.physbeh.2017.04.006.
94.   Kim, K.; Bae, J.; Jin, Y.; Moon, C. Odor habituation can modulate very early olfactory event-related
potential. *Sci Rep*. **2020**, 10, 18117. doi: 10.1038/s41598-020-75263-7.
95.   Bandiera, B.; Natale, F.; Rinaudo, M.; Sollazzo, R.; Spinelli, M.; Fusco, S.; Grassi, C. Olfactory
stimulation with multiple odorants prevents stress-induced cognitive and psychological
alterations. *Brain Commun*. **2024**, 6, fcae390. doi: 10.1093/braincomms/fcae390.
96.   Okada, K.; Shimatani, K. Effect of olfactory stimulation from aromatherapy on the autonomic
nervous activity during aerobic exercises. *Sci Rep*. **2024**, 14, 11198. doi: 10.1038/s41598-024-61732-w.
97.   Scutigliani, E.M.; Liang, Y.; Crezee, H.; Kanaar, R.; Krawczyk, P.M. Modulating the Heat Stress
Response to Improve Hyperthermia-Based Anticancer Treatments. *Cancers (Basel)*. **2021**, 13, 1243.
doi: 10.3390/cancers13061243.
98.   Ahmed, K.; Zaidi, S.F.; Mati-Ur-Rehman.; Rehman, R.; Kondo, T. Hyperthermia and protein
homeostasis: Cytoprotection and cell death. *J Therm Biol*. **2020**, 91, 102615. doi:
10.1016/j.jtherbio.2020.102615.
99.   Selkoe, D.J.; Hardy, J. The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol Med*.
2016, 8, 595-608. doi: 10.15252/emmm.201606210.
100. Karran, E.; De Strooper, B. The amyloid hypothesis in Alzheimer disease: new insights from new
therapeutics. *Nat Rev Drug Discov*. **2022**, 21, 306-318. doi: 10.1038/s41573-022-00391-w.
101. Tundo, G.R.; Sbardella, D.; Ciaccio, C.; Bianculli, A.; Orlandi, A.; Desimio, M.G.; Arcuri, G.; Coletta,
M.; Marini, S. Insulin-degrading enzyme (IDE): a novel heat shock-like protein. *J Biol Chem*. **2013**,
288, 2281-2289. doi: 10.1074/jbc.M112.393108.
102. Vekrellis, K.; Ye, Z.; Qiu, W.Q.; Walsh, D.; Hartley, D.; Chesneau, V.; Rosner, M.R.; Selkoe, D.J.
Neurons regulate extracellular levels of amyloid beta-protein via proteolysis by insulin-degrading
enzyme. *J Neurosci*. **2000**, 20, 1657-1665. doi: 10.1523/JNEUROSCI.20-05-01657.2000.
103. Farris, W.; Mansourian, S.; Chang, Y.; Lindsley, L.; Eckman, E.A.; Frosch, M.P.; Eckman, C.B.;
Tanzi, R.E.; Selkoe, D.J.; Guenette, S. Insulin-degrading enzyme regulates the levels of insulin,
amyloid beta-protein, and the beta-amyloid precursor protein intracellular domain in vivo. *Proc*
Natl Acad Sci U S A. **2003**, 100, 4162-4167. doi: 10.1073/pnas.0230450100.
104. Tian, Y.; Jing, G.; Zhang, M. Insulin-degrading enzyme: Roles and pathways in ameliorating
cognitive impairment associated with Alzheimer's disease and diabetes. *Ageing Res Rev*. **2023**, 90,
101999. doi: 10.1016/j.arr.2023.101999.
105. Chovatiya, R.; Medzhitov, R. Stress, inflammation, and defense of homeostasis. *Mol Cell*. **2014**, 54,
281-288. doi: 10.1016/j.molcel.2014.03.030.
106. Thomas, M.P.; Lieberman J. Live or let die: posttranscriptional gene regulation in cell stress and
cell death. *Immunol Rev*. **2013**, 253, 237-252. doi: 10.1111/imr.12052.
107. Mesihovic, A.; Iannacone, R.; Firon, N.; Fragkostefanakis, S. Heat stress regimes for the
investigation of pollen thermotolerance in crop plants. *Plant Reprod*. **2016**, 29, 93-105. doi:
10.1007/s00497-016-0281-y.

Figure 1

Figure 2

Figure 3

Figure 4

A

B

Figure 5

1st Group

| 1st treatment | 2nd treatment | 3rd treatment | Odor sensitivity test |

0 — 7 — 10 — 13 — 15 — 16 (day)

Buried food test

2nd Group

| Injection of Aβ solution | 1st treatment | 2nd treatment | 3rd treatment | Y-Maze test | 24 h NOR test |

0 — 7 — 10 — 13 — 14 — 15 — 16 (day)

1 h NOR test