Thermal cycling stimulation via nasal inhalation attenuates Aβ$_{25\text{-}35}$-induced cognitive deficits in C57BL/6 mice

Guan-Bo Lin[1,2,†], Hsu-Hsiang Liu[2,3,†], Yu-Yi Kuo[1,2], You-Ming Chen[2,3], Fang-Tzu Hsu[1,2], Yu-Wei Wang[2,3], Yi Kung[2,3], Chien Ching[2,3] and Chih-Yu Chao[1,2,3*]

[1] Department of Physics, Laboratory for Medical Physics & Biomedical Engineering, National Taiwan University, Taipei 106319, Taiwan; gblin@phys.ntu.edu.tw (G.B.L.); hhliu@phys.ntu.edu.tw (H.H.L.); yykuo@phys.ntu.edu.tw (Y.Y.K.); ymchen@phys.ntu.edu.tw (Y.M.C.); tsyr8924503@phys.ntu.edu.tw (F.T.H.); f10245021@phys.ntu.edu.tw (Y.W.W.); a13956@phys.ntu.edu.tw (Y.K.); albertj8808@phys.ntu.edu.tw (C.C.); cychao@phys.ntu.edu.tw (C.Y.C.)
[2] Molecular Imaging Center, National Taiwan University College of Medicine, Taipei 100233, Taiwan
[3] Graduate Institute of Applied Physics, Biophysics Division, National Taiwan University, Taipei 106319, Taiwan

[*] Correspondence: cychao@phys.ntu.edu.tw
[†] These authors contributed equally to this work.

ABSTRACT

Alzheimer's disease (AD) continues to pose a significant public health challenge, with current treatments demonstrating limited effectiveness, partly due to the difficulty of delivering therapeutics across the blood–brain barrier (BBB). The nose-to-brain (N-2-B) pathway offers a promising alternative, enabling pharmacological agents to circumvent the BBB. However, to date, no drugs have been successfully implemented in clinical settings for the treatment of AD via this route, underscoring the necessity for additional research in this area. Mild stress is thought to activate intrinsic protective mechanisms against neurodegeneration, but traditional methods of inducing stress often lack both specificity and practicality. To address this, we propose inhaling mildly heated air as thermal stimulation, leveraging the N-2-B pathway to elicit mild stress and stimulate the brain. Furthermore, this study employs our thermal cycling hyperthermia (TC-HT) concept, adapted as thermal cycling stimulation via nasal inhalation (TCSNI), which delivers cyclic stimulation to sustain pathway activity while minimizing thermal damage. In this study, we evaluated the health and olfactory effects of TCSNI in C57BL/6 mice, observing no adverse impact. In experimental groups administered with β-amyloid (Aβ), TCSNI treatment resulted in significant enhancements in cognitive performance as evidenced by Y-Maze and novel object recognition (NOR) assessments, suggesting an improvement in cognitive function. Protein analyses conducted on the hippocampus of the mice indicated a reduction in Aβ accumulation, alongside increased expression of heat-shock protein 70 (HSP70) and insulin-degrading enzyme (IDE) expression, as well as elevated levels of phosphorylated Akt (p-Akt). These results suggest that N-2-B-delivered TCSNI effectively modulates protein expression and enhances cognitive function, highlighting its potential for further exploration in AD treatment.

Keywords: Alzheimer's disease; β-amyloid; hyperthermia; nasal inhalation; mild stress

INTRODUCTION

Alzheimer's disease (AD) is one of the leading causes of dementia and poses an increasing threat to human health [1]. The abnormal deposition of β-amyloid (Aβ) in the brain is believed to contribute significantly to the oxidative stress and damage that underlie the development of AD [2-4]. Additionally, Aβ aggregation is associated with synaptic dysfunction and neuronal loss, contributing to cognitive decline in AD patients [5,6]. Therefore, Aβ has emerged as a primary target for AD therapy [5-7]. In recent years, drugs developed based on the Aβ hypothesis have gained FDA approval, namely aducanumab and lecanemab [8,9]. However, these drug treatments are costly and associated with the risk of brain edema and abnormal bleeding [10,11], which pose challenges to their widespread adoption. While medication is the main therapeutic approach, the blood-brain barrier (BBB) constrain advancements in new drug development and application [12,13], highlighting the need for further exploration of alternative therapies. Integrating mild stress-based non-pharmacological interventions into healthcare strategies offers a promising avenue for promoting health [14-18]. How to design optimal and well-controlled mild stress stimuli has attracted considerable interest due to their ability to stimulate brain cells, enhance cognitive function, and potentially extend lifespan, thereby providing beneficial effects in the context of neurodegenerative diseases [19-21]. These benefits are largely mediated through the principle of hormesis, wherein exposure to low levels of stressors induces adaptive cellular responses that enhance resilience against neurodegenerative processes [22-25]. Studies indicated that mild stress may upregulate heat shock proteins (HSPs), which are associated with reduced Aβ aggregation [26,27]. Exercise and sauna are widely recognized as physiological stimuli capable of inducing mild stress in humans [19-21,28,29]. However, despite their general health benefits, these long-term approaches are time-intensive and lack conclusive evidence of curative effects [30-33]. In view of this, exploring novel and convenient alternative approaches is valuable.

In order to address this issue, investigating other viable solutions is of utmost importance. The nose-to-brain (N-2-B) pathway offers an alternative approach. Unlike traditional drug delivery methods that rely on penetration enhancers to facilitate transport across the BBB, the N-2-B pathway enables drug delivery via nasal inhalation, thereby circumventing this limitation [34,35]. Drugs such as donepezil, rivastigmine, and natural plant extracts like quercetin and curcumin have also been tested in N-2-B-related research [36,37]. Additionally, insulin inhalation has garnered significant attention in N-2-B-related studies [36,38]. In parallel, aromatherapy is also considered an effective complementary or alternative therapy that may offer benefits in treating dementia [39-41]. Studies indicate that inhaled aromatherapy may mitigate

dementia-related symptoms through olfactory stimulation and modulation of brain
regions governing emotion and cognition [42-44]. Unfortunately, despite extensive
research on various drugs and bioactive compounds, no medication has yet been
successfully applied through this route for the clinical treatment of AD [36],
suggesting that further research into the N-2-B pathway is necessary.

While pharmacological treatments and complementary therapies such as
aromatherapy have shown progress, their impact remains constrained. Building upon
the concept of olfactory stimulation and beyond conventional pharmacological
strategies, we propose, for the first time, the novel application of inhaling mildly
heated air as a form of thermal stimulation via the N-2-B pathway to activate the brain
and induce mild stress. Thermal stimulation has been regarded as having benefits in
various medical fields for a long time [45-47]. Appropriate hyperthermia (HT) can
enhance the expression of heat shock proteins (HSPs), particularly heat shock protein
70 (HSP70), and is considered beneficial for improving AD [48-50]. In addition to the
HSP family, studies have also indicated that HT can increase the expression of
insulin-degrading enzyme (IDE), which is believed to reduce Aβ accumulation
[51,52]. Furthermore, thermal stress has been shown to activate Akt signaling [53,54],
and the neuroprotective effects of phosphorylated Akt (p-Akt) in AD have also been
discussed [54-57]. As a thermal stimulation practice for preventing dementia, sauna
has garnered considerable interest in recent years [30,31,48]. However, while sauna
use has been associated with potential cognitive benefits, its therapeutic effects are
still inconclusive [30,31]. Besides, accessibility issues and safety considerations,
particularly for elderly individuals with cardiovascular risk factors, may limit its
practicality [58,59]. In contrast to systemic heating, this study proposes a novel
strategy using the inhalation of slightly heated air to induce mild stress, with the aim
of modulating protein expression and explore its potential benefits for cognitive
function. Furthermore, to minimize the potential risks associated with continuous heat
treatment, our team adapted the previously developed concept of thermal cycling
hyperthermia (TC-HT) into the novel method, termed thermal cycling stimulation via
nasal inhalation (TCSNI). Through repeated cycles of heated and
ambient-temperature air delivery, TCSNI may mitigate excessive stress and promote
sustained activation of HSP-related signaling pathways.

In the present study, we aimed to investigate the potential effects of inhaling
slightly heated air as a novel form of mild thermal stimulation via nasal inhalation.
We compared the effects of TCSNI with continuous hyperthermia stimulation via
nasal inhalation (HTSNI), focusing on their respective impacts on olfactory function
in C57BL/6 mice. Furthermore, we examined the potential of TCSNI to enhance
cognitive performance, as assessed by Y-Maze and novel object recognition (NOR)

tests [60-63], in a mouse model of cognitive impairment induced by intracerebroventricular (i.c.v.) injection of Aβ$_{25\text{-}35}$ [64]. In addition, we investigated heat- and stress-responsive proteins related to neurodegeneration, including HSP70, IDE, and phosphorylated Akt (p-Akt). These proteins are believed to contribute to antioxidant defense, support cell survival, and facilitate the clearance of Aβ accumulation [48-57]. This study aims to provide new insights into the therapeutic potential of nasal inhalation-based thermal stimulation and its role in preventing or alleviating cognitive decline, thereby laying the groundwork for future applications in neurodegenerative disease management.

RESULTS

Effects of HTSNI and TCSNI on olfactory function in mice

The aim of this study is to investigate a novel hyperthermic stimulation applied through the nasal cavity. Before conducting cognitive tests in the AD mice model, we first examined safety in mice by administering HTSNI and TCSNI and assessing whether these thermal stimulation methods induced any adverse effects, as shown in Figure 1A. Since both HTSNI and TCSNI are forms of hyperthermic stimulation applied to the nasal cavity, the olfactory function of mice was examined after treatments. The buried food test and odor sensitivity test are common methods for assessing the olfactory function of mice [65,66]. In the experiments, we found that after treatment of TCSNI, the latency to find the food (15.2±3.1 s) did not differ significantly from that of the control group (12.2±2.1 s) in the buried food test. However, in the HTSNI group, the latency was significantly increased (60.8±5.1 s) compared with both the control and TCSNI groups. In addition, in the odor sensitivity test, the time spent sniffing odor stimuli did not differ significantly between the control and TCSNI groups across the range of odor dilutions. Notably, at the 10^{-3} dilution, the HTSNI group exhibited significantly shorter sniffing time (1.2±0.6 s) compared to the control (5.8±1.5 s) and TCSNI (3.3±1.2 s) groups. Furthermore, at the 10^{-2} dilution, the HTSNI group spent significantly less time sniffing (0.9±0.7 s) than both the control (12.3±1.8 s) and TCSNI (5.8±0.9 s) groups, as shown in Figure 1B. These findings suggest that mice in the HTSNI group had impaired olfactory sensitivity or reduced ability to detect and respond to odor cues at concentrations that remained effective in the control and TCSNI groups. In view of this, we applied TCSNI in the subsequent examinations to assess whether it could enhance recognition in the AD mouse model.

Figure 1. Effects of HTSNI and TCSNI on olfactory detection in mice. Olfactory performance was assessed using the buried food test and olfactory habituation test, as

described in the Materials and Methods section. (A) Buried food test: latency to locate the buried food. (B) Olfactory habituation test: response time to odor stimuli. Data are shown as mean ± standard error of the mean. Statistical significance was determined using one-way analysis of variance (ANOVA) followed by Tukey's post-hoc test. Significance levels between indicated groups are denoted as ***P < 0.001, while non-significant differences are indicated as ns.

TCSNI attenuates Aβ-Induced cognitive impairments in mice

Since memory loss is a hallmark of AD [67], studies using mouse models commonly employ Y-maze and NOR tests to assess cognitive deficits [60-63]. Spontaneous alternation in the Y-maze is a well-established measure of short-term and spatial working memory [60,61], whereas the NOR test is a widely used paradigm for assessing both short-term and long-term recognition memory [62,63]. In this study, we utilized these tests to evaluate the effects of TCSNI on Aβ-injected mice. As shown in Figure 2A, there was no significant difference in the number of arm entries in the Y-Maze test among all groups, suggesting that i.c.v. injection of phosphate-buffered saline (PBS) or Aβ, and TCSNI treatment did not influence the locomotor activity of mice. Moreover, the spontaneous alternation index was significantly reduced in the Aβ group compared to the control group, which declined from 62.1±3.5% to 49.8±2.3%. Notably, TCSNI treatment elevated the spontaneous alternation index, suggesting an improvement in Aβ-induced deficits in short-term and spatial working memory, as shown in Figure 2B. In addition, findings from the short-term and long-term NOR tests support the effectiveness of TCSNI in ameliorating Aβ-induced recognition memory impairment. As shown in Figure 2C, the discrimination index in the short-term NOR tests significantly decreased in the Aβ group compared to the control group, dropping from 62.2±7.5% to 8±11.5%, whereas TCSNI treatment significantly improved the discrimination index (59.2±6.6%). Furthermore, in the long-term NOR test, Aβ-induced memory impairment was more pronounced, with the discrimination index declining from 49.8±6.5% to 2.2±10.5%, which was restored to 46.2±5.5% following TCSNI treatment. Collectively, these findings indicate that TCSNI treatment effectively alleviates Aβ-induced cognitive deficits, as demonstrated in the Y-Maze and NOR tests.

Figure 2. Effect of TCSNI applications on mice's behavioral performances. The recovery effects of TCSNI treatment on Aβ-induced recognition impairments were assessed via (A) the total number of total arm entries and (B) the spontaneous alternation index in the Y-maze test, and (C) the discrimination index with intertrial period of 1 h and (D) 24 h in the NOR maze test. Data are shown as mean ± standard

202 error of the mean. Statistical significance was determined using one-way analysis of
203 variance (ANOVA) followed by Tukey's post-hoc test. Significance levels between
204 indicated groups are denoted as ***P < 0.001, while non-significant differences are
205 indicated as ns.
206
TCSNI downregulates Aβ accumulation and elevates HSP70, IDE, and p-Akt
207
levels in the mouse hippocampus
208
With Aβ being a major focus in AD treatment strategies [5-7], this study first
evaluated its accumulation in the hippocampus of mice. As shown in Figure 3A, a
1.98-fold increase in Aβ level was observed in the Aβ-injected group compared to the
control group. Notably, TCSNI treatment significantly reduced the hippocampal Aβ
level, decreasing it from 1.98-fold to 1.12-fold that of the control group,
demonstrating a strong inhibitory effect. Furthermore, HSP70, a key stress response
protein, functions as a molecular chaperone to prevent protein misfolding and has also
been reported to influence the Aβ aggregation process [68-70]. In vivo studies showed
that Aβ administration may lead to a slight elevation in HSP70 levels without
reaching statistical significance [71,72]. In this study, HSP70 was also investigated,
and the results showed that hippocampal HSP70 levels in the Aβ-injected group were
not significant compared to those in the control group, whereas TCSNI treatment
significantly upregulated HSP70 levels, reaching 1.22-fold of the control group
(Figure 3B). On the other hand, IDE has long been recognized for its contribution to
Aβ clearance, and its altered activity or expression may lead to Aβ accumulation and
increased AD risk [51,73,74]. As shown in Figure 3C, Aβ injection reduced
hippocampal IDE levels in mice to 0.85-fold of the control. In this study, we found
that TCSNI treatment reversed this decline, significantly increasing IDE expression to
1.52-fold of the control group. Furthermore, it is known that Akt serves a crucial
function in neuronal survival signaling [75,76], and a decline in Akt activation has
been observed in AD models [55-57,77,78]. In addition, Akt responds to stress and
thermal stimuli [53,54,79], drawing our attention. Therefore, in this study, we
examined the phosphorylation ratio of Akt. As illustrated in Figure 3D, injection of
Aβ led to a marked downregulation of the p-Akt/t-Akt ratio (0.65-fold), while TCSNI
significantly upregulated the ratio to 1.53-fold compared to the control. These
findings suggest that Aβ removal might be associated with the upregulation of HSP70,
IDE, and p-Akt/t-Akt in the hippocampi of mice.
236
Figure 3. Effect of TCSNI applications on the expression levels of Aβ, p-Akt,
HSP70, and IDE in hippocampi. Representative western blots and the
quantifications of (A) Aβ, (B) HSP70, (C) IDE, and (D) p-Akt are presented for

evaluating anti-Aβ effect in hippocampi. The protein levels of Aβ, HSP70, and IDE
were normalized to β-actin, and p-Akt was normalized to t-Akt. Each relative
expression level was compared with the control and represented as fold relative to the
control. Data are shown as mean ± standard error of the mean. Statistical significance
was determined using one-way ANOVA followed by Tukey's post-hoc test.
Significance levels between indicated groups are denoted as $*P < 0.05$, $**P < 0.01$,
and $***P < 0.001$, while non-significant differences are indicated as ns.
247
DISCUSSION
To date, AD continues to pose a major medical challenge for humanity. The BBB
presents a significant obstacle to the targeted delivery of drugs to the brain for treating
AD [12,13]. Alternatively, the N-2-B pathway has been recognized as the ideal route
to bypass the BBB, and studies have indicated its potential to offer therapeutic effects
for brain disorders or brain cancer [34-36]. In addition, aromatherapy is also regarded
as a complementary or alternative therapy that may provide benefits for dementia
treatment [41]. Despite the growing interest in the N-2-B pathway as a means to
bypass the BBB, no medication has yet been clinically approved via this route for the
treatment of AD [36]. Given the absence of a cure for AD and the limited efficacy of
current pharmacological treatments, attention has shifted towards exploring
alternative approaches [80-82]. Literature indicates that mild stress stimuli enhance
the functional capacity of organisms and exert positive biological effects on cells
[20,83-85]. Moreover, non-pharmacological interventions, such as exercise and sauna,
which induce mild stress, are believed to enhance human health, promoting lifespan
extension and cognitive improvement [28,29,84,85]. Among them, whole-body
heating sauna has gained great attention and sauna-like conditions were evaluated
[28,30,31,86]. Nevertheless, due to drawbacks such as being time-consuming,
practical challenges, and the lack of a curative effect, it is crucial to explore more
effective and safe ways to induce desired mild stress.
In the current study, we introduced a novel approach of using mildly heated air
inhalation as a thermal stimulus to induce mild stress. Unlike whole-body heating
methods, this approach delivers warm air specifically to the nasal cavity. Nonetheless,
an evaluation of this approach is necessary to determine its safety implications before
further testing. A previous study indicated that mice experience thermal collapse
under excessive heat conditions [87]. In contrast to the aforementioned condition
where the entire body is exposed to a high-temperature environment, this study
employs localized nasal heating via gas application. Furthermore, the TC-HT method
developed in our previous studies, has demonstrated efficacy in both in vitro and in
vivo models with minimal side effects, primarily due to its intermittent stimulation

intervals, which mitigate excessive damage [52,54]. Accordingly, we evaluated the
olfactory function in mice subjected to HTSNI and TCSNI. As shown in Figure 1,
after administration of HTSNI, the mice showed impaired olfactory function, as
assessed by the buried food test and the olfactory discrimination test, whereas the
TCSNI-treated group maintained olfactory function comparable to the untreated
control group. These findings suggest that TCSNI is a safer approach compared to
HTSNI.

As memory loss is a defining feature of AD [67], research involving mouse
models often utilizes Y-maze and NOR tests to examine memory deficits [60-63].
Given that TCSNI preserves olfactory function in mice, it was chosen as the thermal
stimulation method for behavioral tests in cognitive impairment mice to reduce
potential sensory disturbances, particularly considering the olfactory decline observed
with HTSNI. This study investigated the alleviating effect of TCSNI on Aβ-induced
memory impairment and demonstrated that TCSNI significantly enhanced the
spontaneous index and the ability to recognize novel objects, as shown in Figure 2.
These results indicate that TCSNI holds significant potential for improving memory
deficits and cognitive dysfunction in vivo in a safety manner.

Although traditional continuous HT stimulation hold potential in inducing
positive mild stress, literature indicated that unappropriated HT show no benefit
[84,85,88], and may contribute to an increased accumulation of reactive oxygen
species (ROS), which could exacerbate oxidative stress and lead to cellular damage
[88-90]. In our previous in vitro study, HT lacked the ability to reduce ROS and
exhibited less protective effects than TC-HT against hydrogen peroxide- and
Aβ-induced cytotoxicity [54]. In addition, the other previous study using thermofoil
demonstrated that continuous HT application was less effective than TC-HT in
improving memory impairment in vivo [52]. These studies indicate that replacing
continuous heating with cyclical thermal stimulation may elicit unique biological
responses [52,54]. The underlying mechanism may be related to the activation and
saturation of stimulated protein pathways. For example, in olfactory field, long-term
and continuous exposure may lead to olfactory adaptation or receptors desensitization
[91-94], which might affect the outcome of aromatherapy. Therefore, combining
different scents or using intermittent exposure is an important strategy to sustain
receptor stimulation and therapeutic efficacy [95,96]. In thermal stimulation research,
the phenomenon of thermal adaptation has also been investigated and is considered
one of the key factors contributing to the decreased effectiveness of HT [97,98].
Compared to HT, TC-HT applies periodic temperature fluctuations that might help
prevent signaling desensitization and better sustain heat-induced protein responses. In
addition, with an improved cellular safety profile, TC-HT methods potentially

enhance both the efficacy and feasibility of thermal stimulation, making them a
preferable alternative methodology to traditional HT [52,54]. In this study, while
HTSNI was found to impair olfactory function, TCSNI did not cause such impairment
and was shown to ameliorate deficits in memory and cognition (Figures 1 and 2).

Given that the clearance of toxic Aβ deposits is a significant marker in AD
therapy [99,100], we investigated Aβ expression levels in the hippocampus of mice
from a molecular perspective. As depicted in Figure 3A, TCSNI application
significantly reduced Aβ expression levels in the hippocampus. Furthermore, several
proteins with potential roles in Aβ clearance were investigated. Among these, HSP70,
a key heat shock protein activated by thermal stimulation, has gained significant
attention for its role in suppressing AD progression [68-70]. As HSP70 primarily
functions to prevent protein misfolding, it has also been reported to influence the Aβ
aggregation process [68-70]. The study found that TCSNI application significantly
enhanced HSP70 expression in the hippocampus of mice, as shown in Figure 3B. In
addition, HSP70 has been suggested to upregulate the expression of IDE, a
well-known Aβ-degrading enzyme, thereby facilitating Aβ clearance [68]. Studies
have shown that IDE is induced by stress and identified it as a protein with heat
shock-like characteristics [51,52,101]. Additionally, IDE has been found to regulate
extracellular Aβ levels via proteolysis and mitigate its neurotoxic effects [102,103].
Research suggests that reduced IDE activity and expression may be associated with
Aβ accumulation and an increased risk of AD [51,73,74], and thus upregulating IDE
levels has been proposed as a potential therapeutic approach for AD [74]. In line with
these studies, our study finds that TCSNI treatment significantly restores the
Aβ-induced reduction in IDE in the hippocampus of mice, as shown in Figure 3C.
Furthermore, in addition to IDE and HSP70, Akt is another key protein of interest in
our study. Akt plays a pivotal role in mediating neuron survival signaling, growth, and
synaptic plasticity, which are critical for brain development and function [75,76]. It
was found that a decline in Akt activation occurred in AD models, while the
upregulation of Akt activation conferred protective effects in both in vitro and in vivo
models [55-57,77,78]. It is noteworthy that Akt is also recognized as a protein that
responds to stress and thermal stimulation [53,54,79]. In addition, research indicated
that Akt could mediate HSP70 expression, thereby providing neuroprotection [78].
Furthermore, study suggested that Akt signaling pathway may improve cognitive
impairment in AD by promoting the expression of IDE and the degradation of Aβ
[104]. In this work, along with HSP70 and IDE proteins, Akt protein in the
hippocampus of mice was detected. We found that Aβ-induced decline in p-Akt was
restored by TCSNI application, as shown in Figure 3D. Taken together, these findings
suggest that TCSNI induces mild thermal stress, which has the potential to upregulate

the expression of HSP70, IDE, and phosphorylated Akt in the hippocampus of Aβ-injected mice. This molecular response may contribute to the reduction of hippocampal Aβ levels and underlie the cognitive improvements observed in the treated mice, consistent with the mild stress-induced self-protective hypothesis.

This study proposes a novel periodic hyperthermic method, TCSNI, administered via nasal inhalation to induce mild stress as a potential treatment for AD. Stress response would trigger self-defense system [105], and depending on the intensity and duration of stress, cells would adapt or cause their death [106]. Studies have exhibited that preconditioning cells with non-lethal stress minimizes cell damage and improves cell survival [54,107]. The present study suggests an efficient way to induce mild stress in vivo with minimal impact on olfactory function using TCSNI, which has been shown to improve cognitive performance in mice in Y-maze and NOR tests. Moreover, TCSNI could reduce Aβ expression and significantly upregulate proteins related to Aβ degradation and cell survival, such as HSP70, IDE, and p-Akt, in the hippocampus of mice subjected to i.c.v. Aβ injection. Further studies are required to confirm the involvement of other stress-induced proteins, particularly those associated with neuroprotection and Aβ metabolism, and to validate these findings in other species and humans. Additionally, the potential combinatory effects of TCSNI with pharmacological treatments also warrant further investigation. Overall, this study provides new insights into the development of alternative therapeutic strategies for AD, highlighting the potential of TCSNI as a novel non-invasive intervention.

MATERIALS AND METHODS
Experimental animals and housing

C57BL/6 male mice (6–8 weeks old, 22–26 g) were purchased from National Laboratory Animal Center, and housed at National Taiwan University Animal Resources Center. The mice had ad libitum access to food and water and were maintained at a constant temperature of 23℃ ± 2℃, under a 12:12-hour light/dark cycle (lights switched on at 7:00 AM). All experimental techniques were executed based on established guidelines of the Animal Ethical Committee of National Taiwan University (approval protocol number: NTU-112-EL-00158). All mice were randomly assigned to two main groups. The first group was further divided into control, TCSNI, and HT sub groups, while the second group was divided into control, Aβ, and Aβ+TCSNI sub groups.

β-amyloid administration

Aβ$_{25-35}$ (Sigma-Aldrich; Merck KGaA) was prepared as stock solution at a concentration of 1 mg/mL in 0.9% sterile saline solution and fibril-aggregated at 37℃

392   for 7 days. On day 0, 10 µL of the aggregated form of Aβ was injected i.c.v. into the
393   brains of the mice in the Aβ and Aβ-TCSNI groups, while the control group mice
394   received an i.c.v. injection of 10 µL PBS.
395
HTSNI and TCSNI applications
This experiment utilized a novel nasal inhalation-based thermal stimulation
system. As shown in Figure 4A, the system heats 3% isoflurane anesthetic gas by the
water bath method and controls the duration of gas heating and cooling through a
timer-controlled solenoid valve. In the HTSNI application, the timer-controlled
solenoid valve was fixed to ensure a steady release of heated air, and the temperature
was maintained at 47.5 ± 0.5℃ for 30 minutes. In the TCSNI application, the
timer-controlled solenoid valve was programmed to execute a 10-cycle repetitive
process, where each cycle consists of a 3-minute heating phase (47.5 ± 0.5℃),
followed by a 1-minute cooling phase. The actual air temperature entering the
mouse's nasal cavity is measured and monitored by a K-type thermocouple positioned
at the end of the pipeline, with the measured temperature results shown in Figure 4B.
408
409   **Figure 4. Experimental setup and design of TCSNI.** (A) The TCSNI system was
410   illustrated, which provides inhaling mildly heated air to deliver cyclic thermal
411   stimulation. (B) The actual temperatures in the SH-SY5Y cells were measured using a
412   K-type needle thermocouple at 10 sec intervals throughout the exposure durations of
413   TC-HT and HT treatments.
414
Experimental design
For the 1st group, as shown in Figure 5, on day 7, 10, and 13, the respective
thermal application was employed to the mice of the HTSNI and TCSNI groups.
Buried food and odor sensitivity tests were conducted on day 15 and 16, respectively.
For the 2nd group, 10 µL of Aβ solution was i.c.v.-injected into the brains of mice in
the Aβ and Aβ+TCSNI groups on day 0, while control group mice received an i.c.v.
injection of 10 µL PBS. On day 7, 10, and 13, the TCSNI thermal application was
employed to the mice of different groups. The Y-maze test was performed at day 14,
and the NOR tests were carried out at day 15 and 16. After the NOR test, mice were
weighted and then euthanized to collect brain tissues.
425
Figure 5. TCSNI treatment and animal behavioral assessment. (A) For the 1st
group, the thermal applications were employed to the mice of the HTSNI and TCSNI
groups on days 7, 10 and 13. Buried food and odor sensitivity tests were assessed on
day 15 and 16, respectively. (B) For the 2nd group, Aβ solution was injected

intracerebroventricular (i.c.v.) at day 0, and TCSNI application was applied at day 7,
10, and 13. The Y-maze test was performed at day 14, and the NOR tests were carried
out at day 15 and 16, respectively. After the NOR test, mice were weighted and then
euthanized for brain tissues collection.
434
Buried food test
On day 14 at 4 pm, food pellets were removed from the cages and the testing
mice fasted overnight. The test was performed on day 15 at 10 am after 1-hour
acclimatization in the testing room. Mice were then individually introduced into a
clean cage containing 3 cm deep of clean bedding and allowed to acclimate to the
cage for 5 minutes to reduce the interference from novel environment exploration
during the test. A small piece of pellet with a diameter of about 5 mm was buried
beneath 1 cm of bedding in a random corner of the cage, and the mouse was
introduced into the cage. The site of mice placement and the location where the small
pellet was buried remained constant. Time required for the mice to retrieve the small
pellet with its front paws was measured in seconds (latency), with a maximum
duration of 15 minutes.
447
Odor Sensitivity Test
The odor sensitivity test was performed on day 16 after 1-hour acclimatization in
the testing room for the mice. Mice were then individually introduced into a clean
cage, which was pre-placed with a dry cotton swab without bedding. The mice were
allowed to acclimate to the cage for 30 minutes to reduce the interference from novel
environment exploration during the test. After acclimatization, video recording began
and lasted for 3 minutes, with the dry cotton swab was replaced by a cotton swab
soaked in water. After 3 minutes, replace the cotton swab sequentially with diluted
solutions at concentrations of 1/100,000, 1/10,000, 1/1,000, and 1/100. Sniffing
behavior increases when the odor concentration reaches the olfactory threshold in
mice. After acquiring all of the olfactory sensitivity and preference videos, a timer
was used to time all investigations of cotton swabs within three minutes. The time of
investigation is defined as any period when the mouse's nares are < 1 mm from the
cotton swabs. The recorded latency would not include bouts when the mouse is not
actively investigating, chewing, or licking the paper
463
Y-Maze test
The Y-maze test was employed to assess spatial working memory. The apparatus
consists of three identical symmetric arms (40 cm × 13 cm × 6.5 cm) arranged at 120°
angles from a central junction. Each mouse, with no prior familiarity with the maze,

was individually placed at the center and allowed to freely explore all three arms to
test spontaneous alteration for 8 minutes. After each trial, the maze was cleaned with
70% alcohol to eliminate olfactory cues. The behavioral performance was recorded,
and the maze arm entries were labeled 'A', 'B', and 'C'. An arm entry was defined as
the hind paws of mice completely entering the arm. A correct alternation was defined
as sequential entry into the three different arms (i.e., ABC, ACB, BAC, BCA, CAB,
and CBA) in overlapping triple sequences. The spontaneous alternation index was
calculated using the following formula: (number of correct alternations) / (total
number of arm entries – 2) × 100, and the total number of arm entries was also used
as an indicator of general locomotor activity.
478
Novel object recognition test
The NOR test was conducted to evaluate recognition memory in mice across
days 14 to 16 of the study. The test was performed in an open field arena (40 cm × 40
cm × 30.5 cm) over three consecutive days. On day 14 (habituation phase), each
mouse was allowed to freely explore the empty arena for 8 minutes to acclimate to the
testing environment. On day 15 (familiarization and short-term recognition memory
test phase), two identical familiar objects (object A) were placed near the diagonal
corners of the arena, and the assessed mouse was allowed to explore the arena for 8
minutes. After a 1-hour retention interval in the home cage, the mouse was returned to
the arena for the short-term recognition memory test. During this phase, one of the
original identical familiar objects (object A) was replaced with a novel object (object
B). The mouse's behavioral performance was recorded with an overhead video
camera for 8 minutes. On day 16, to assess long-term recognition memory, the mice
were again placed in the arena and allowed to explore the same familiar object from
the training phase (object A) and a new novel object (object C) for 8 minutes. This
design allowed the evaluation of both short-term and long-term object recognition
memory. The familiar objects were possessing similar shapes, material composition,
and coloration. The novel objects were objects of varying shapes, sizes, and colors in
comparison to the familiar objects. After each assessment in all phases, all olfactory
cues were removed by 70% ethanol. Object exploration was defined as the time spent
by the mouse sniffing or touching the object with the nose and/or forepaws. The
discrimination index was calculated as (Tn-To) / (Tn+To), where Tn and To represent
the exploration time for the novel and familiar objects, respectively.
502
503 **Brain tissue collection and western blot sample preparation**
504   After NOR tests, mice were euthanized using carbon dioxide gas, and mice
505 brains were extracted after confirmation of death and rinsed with PBS (HyClone; GE

Healthcare Life Sciences, Chicago, IL, USA) supplemented with 1% fresh protease inhibitor cocktail (EMD Millipore, Billerica, MA, USA). The hippocampus tissue was homogenized in 500 µL ice-cold RIPA lysis buffer (MilliporeSigma) containing phosphatase & protease inhibitors by the Q125 Sonicator (Qsonica, Newton, CT, USA), and the homogenates were centrifuged at 23,000×g at 4℃ for 30 minutes. The protein concentration levels of the samples were measured using the Bradford protein assay (cat. no. BRA222; BioShop Canada). Protein samples were stored at -80℃ for subsequent western blotting assessments.

Western blot analysis

20 µg of protein samples were separated with 10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis and transferred onto polyvinylidene fluoride membranes (MilliporeSigma) at 180 mA for 150 minutes. After blocking with 5% bovine serum albumin (BioShop Canada Inc.) in Tris-buffered saline with 0.1% Tween-20 (TBST) for 1 hour at room temperature, the membranes were incubated overnight with primary antibodies at 4℃ overnight. The specific primary antibodies against β-Amyloid (Aβ; cat. no. sc-28365; Santa Cruz Biotechnology), phospho-Akt (p-AKT; cat. no. 4060; Cell Signaling Technology, Inc.), Akt (AKT; cat. no. 9272; Cell Signaling Technology, Inc.), IDE (cat. no. ab133561; Abcam), HSP70 (cat. no. 4872; Cell Signaling Technology, Inc.), and β-actin (cat. no. GTX110564; GeneTex, Inc.) were used, with β-actin serving as the loading control for protein normalization. Following washing with TBST, the membranes were incubated with the appropriate horseradish peroxidase (HRP)-conjugated secondary antibodies, including goat anti-rabbit IgG (HRP) (cat. no. 111-035-003; Jackson ImmunoResearch Laboratories, Inc.) or goat anti-mouse IgG (HRP) (cat. no. GTX213111-01; GeneTex, Inc.). Dilutions of all primary and secondary antibodies were performed following the manufacturer's instructions. The membranes were visualized with an enhanced chemiluminescence substrate (Advansta, San Jose, CA, USA) and quantified using the Amersham Imager 600 imaging system (AI600; GE Healthcare Life Sciences).

Statistical analysis

All statistical results were presented as mean ± standard error of the mean (SEM) and analyzed using one-way analysis of variance (ANOVA) with Tukey's post hoc test in OriginPro 2022 software (version 2022; OriginLab Corporation, Northampton, MA, USA). A P-value of less than 0.05 was considered statistically significant.

AUTHOR CONTRIBUTIONS: Conceptualization, C.Y.C.; methodology, G.B.L.,

H.H.L. and C.Y.C.; validation, G.B.L., H.H.L., Y.Y.K., Y.M.C., F.T.H., Y.W.W., Y.K., C.C. and C.Y.C.; formal Analysis, G.B.L., H.H.L., Y.Y.K., Y.M.C. and C.Y.C.; investigation, G.B.L., H.H.L. and C.Y.C.; resources, C.Y.C.; data curation, G.B.L., H.H.L., Y.Y.K., Y.M.C., F.T.H., Y.W.W., Y.K., C.C. and C.Y.C.; writing—original draft preparation, G.B.L., H.H.L. and C.Y.C.; writing – review and editing, G.B.L. and C.Y.C.; supervision, C.Y.C.; funding acquisition, C.Y.C. All authors have read and agreed to the published version of the manuscript.

FUNDING: This work was supported by grants from National Science and Technology Council (NSTC 113-2112-M-002-024 and NSTC 112-2112-M-002-033 to C.Y.C.) and Ministry of Science and Technology (MOST 110-2112-M-002-004 to C.Y.C.) of the Republic of China. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

INSTITUTIONAL REVIEW BOARD STATEMENT: Not applicable.

INFORMED CONSENT STATEMENT: Not applicable.

DATA AVAILABILITY STATEMENT: The data presented in this study are available on request from the corresponding author.

ACKNOWLEDGMENTS: We thank the Animal Resource Center and the Consortium of Integrative Biomedical Science Key Technology at National Taiwan University for their technical support.

ABBREVIATIONS

AD	Alzheimer's disease
Aβ	β-amyloid
N-2-B	nose-to-brain
BBB	blood-brain barrier
HT	hyperthermia
TC-HT	thermal cycling hyperthermia
TCSNI	thermal cycling stimulation via nasal inhalation
HTSNI	hyperthermia stimulation via nasal inhalation
i.c.v.	intracerebroventricular
NOR	novel object recognition
HSP70	heat-shock protein 70
IDE	insulin-degrading enzyme

582	p-Akt	phosphorylated Akt
583	ROS	reactive oxygen species
584	ANOVA	analysis of variance
585		

REFERENCES

1. Scheltens, P.; De Strooper, B.; Kivipelto, M.; Holstege, H.; Chételat, G.; Teunissen, C.E.; Cummings, J.; van der Flier, W.M. Alzheimer's disease. *Lancet.* **2021**, 397, 1577-1590. doi: 10.1016/S0140-6736(20)32205-4.

2. Sun, X.; Chen, W.D.; Wang, Y.D. β-Amyloid: the key peptide in the pathogenesis of Alzheimer's disease. *Front Pharmacol.* **2015**, 6, 221. doi: 10.3389/fphar.2015.00221.

3. Leuner, K.; Schütt, T.; Kurz, C.; Eckert, S.H.; Schiller, C.; Occhipinti, A.; Mai, S.; Jendrach, M.; Eckert, G.P.; Kruse, S.E.; et al. Mitochondrion-derived reactive oxygen species lead to enhanced amyloid beta formation. *Antioxid Redox Signal.* **2012**, 16, 1421-1433. doi: 10.1089/ars.2011.4173.

4. Miranda, S.; Opazo, C.; Larrondo, L.F.; Muñoz, F.J.; Ruiz, F.; Leighton, F.; Inestrosa, N.C. The role of oxidative stress in the toxicity induced by amyloid beta-peptide in Alzheimer's disease. *Prog Neurobiol.* **2000**, 62, 633-648. doi: 10.1016/s0301-0082(00)00015-0.

5. Palop, J.J.; Mucke, L. Amyloid-beta-induced neuronal dysfunction in Alzheimer's disease: from synapses toward neural networks. *Nat Neurosci.* **2010**, 13, 812-818. doi: 10.1038/nn.2583.

6. Reddy, P.H.; Beal, M.F. Amyloid beta, mitochondrial dysfunction and synaptic damage: implications for cognitive decline in aging and Alzheimer's disease. *Trends Mol Med.* **2008**, 14, 45-53. doi: 10.1016/j.molmed.2007.12.002.

7. Chen, G.F.; Xu, T.H.; Yan, Y.; Zhou, Y.R.; Jiang, Y.; Melcher, K.; Xu, H.E. Amyloid beta: structure, biology and structure-based therapeutic development. *Acta Pharmacol Sin.* **2017**, 38, 1205-1235. doi: 10.1038/aps.2017.28.

8. Rahman, A.; Hossen, M.A.; Chowdhury, M.F.I.; Bari, S.; Tamanna, N.; Sultana, S.S.; Haque, S.N.; Al Masud, A.; Saif-Ur-Rahman, K.M. Aducanumab for the treatment of Alzheimer's disease: a systematic review. *Psychogeriatrics.* **2023**, 23, 512-522. doi: 10.1111/psyg.12944.

9. Hossain, M.F.; Husna, A.U.; Kharel, M. Use of lecanemab for the treatment of Alzheimer's disease: A systematic review. *Brain Behav.* **2024**, 14, e3592. doi: 10.1002/brb3.3592.

10. Cummings, J.; Apostolova, L.; Rabinovici, G.D.; Atri, A.; Aisen, P.; Greenberg, S.; Hendrix, S.; Selkoe, D.; Weiner, M.; Petersen R.C.; et al. Lecanemab: Appropriate Use Recommendations. *J Prev Alzheimers Dis.* **2023**, 10, 362-377. doi: 10.14283/jpad.2023.30.

11. Fleck, L.M. Alzheimer's and Aducanumab: Unjust Profits and False Hopes. *Hastings Cent Rep.* **2021**, 51, 9-11. doi: 10.1002/hast.1264.

12. Pardridge, W.M. Alzheimer's disease drug development and the problem of the blood-brain barrier. *Alzheimers Dement.* **2009**, 5, 427-432. doi: 10.1016/j.jalz.2009.06.003.

13. Wu, D.; Chen, Q.; Chen, X.; Han, F.; Chen, Z.; Wang, Y. The blood-brain barrier: structure, regulation, and drug delivery. *Signal Transduct Target Ther.* **2023**, 8, 217. doi: 10.1038/s41392-023-01481-w.

14. Rattan, S.I. Hormetic modulation of aging and longevity by mild heat stress. *Dose Response.* **2006**, 3, 533-546. doi: 10.2203/dose-response.003.04.008.

15. Schirrmacher, V. Less Can Be More: The Hormesis Theory of Stress Adaptation in the Global Biosphere and Its Implications. *Biomedicines.* **2021**, 9, 293. doi: 10.3390/biomedicines9030293.

16. Kouda, K.; Iki, M. Beneficial effects of mild stress (hormetic effects): dietary restriction and health. *J Physiol Anthropol.* **2010**, 29, 127-132. doi: 10.2114/jpa2.29.127.

17. Calabrese, V.; Giordano, J.; Signorile, A.; Laura Ontario, M.; Castorina, S.; De Pasquale, C.; Eckert, G.; Calabrese, E.J. Major pathogenic mechanisms in vascular dementia: Roles of cellular stress

response and hormesis in neuroprotection. *J Neurosci Res.* **2016**, 94, 1588-1603. doi: 10.1002/jnr.23925.

18. Rattan, S.I.; Demirovic, D. Hormesis can and does work in humans. *Dose Response.* **2009**, 8, 58-63. doi: 10.2203/dose-response.09-041.Rattan.

19. Adlard, P.A.; Engesser-Cesar, C.; Cotman, C.W. Mild stress facilitates learning and exercise improves retention in aged mice. *Exp Gerontol.* **2011**, 46, 53-59. doi: 10.1016/j.exger.2010.10.001.

20. Minois, N. Longevity and aging: beneficial effects of exposure to mild stress. *Biogerontology.* **2000**, 1, 15-29. doi: 10.1023/a:1010085823990.

21. Parihar, V.K.; Hattiangady, B.; Kuruba, R.; Shuai, B.; Shetty, A.K. Predictable chronic mild stress improves mood, hippocampal neurogenesis and memory. *Mol Psychiatry.* **2011**, 16, 171-183. doi: 10.1038/mp.2009.130.

22. Mattson, M.P. Hormesis defined. *Ageing Res Rev.* **2008**, 7, 1-7. doi: 10.1016/j.arr.2007.08.007.

23. Mao, L.; Franke, J. Hormesis in aging and neurodegeneration-a prodigy awaiting dissection. *Int J Mol Sci.* **2013**, 14, 13109-13128. doi: 10.3390/ijms140713109.

24. Wang, G. Hormesis, cell death, and regenerative medicine for neurode-generative diseases. *Dose Response.* **2012**, 11, 238-254. doi: 10.2203/dose-response.12-019.Wang.

25. Mattson, M.P.; Leak, R.K. The hormesis principle of neuroplasticity and neuroprotection. *Cell Metab.* **2024**, 36, 315-337. doi: 10.1016/j.cmet.2023.12.022.

26. Mane, N.R.; Gajare, K.A.; Deshmukh, A.A. Mild heat stress induces hormetic effects in protecting the primary culture of mouse prefrontal cerebrocortical neurons from neuropathological alterations. *IBRO Rep.* **2018**, 5, 110-115. doi: 10.1016/j.ibror.2018.11.002.

27. Smith Sonneborn, J. Alternative strategy for Alzheimer's disease: stress response triggers. *Int J Alzheimers Dis.* **2012**, 2012, 684283. doi: 10.1155/2012/684283.

28. Patrick, R.P.; Johnson, T.L. Sauna use as a lifestyle practice to extend healthspan. *Exp Gerontol.* **2021**, 154, 111509. doi: 10.1016/j.exger.2021.111509.

29. Ji, L.L. Exercise-induced modulation of antioxidant defense. *Ann N Y Acad Sci.* **2002**, 959, 82-92. doi: 10.1111/j.1749-6632.2002.tb02085.x.

30. Knekt, P.; Järvinen, R.; Rissanen, H.; Heliövaara, M.; Aromaa, A. Does sauna bathing protect against dementia? *Prev Med Rep.* **2020**, 20, 101221. doi: 10.1016/j.pmedr.2020.101221.

31. Laukkanen, T.; Kunutsor, S.; Kauhanen, J.; Laukkanen, J.A. Sauna bathing is inversely associated with dementia and Alzheimer's disease in middle-aged Finnish men. *Age Ageing.* **2017**, 46, 245-249. doi: 10.1093/ageing/afw212.

32. Öhman, H.; Savikko, N.; Strandberg, T.E.; Kautiainen, H.; Raivio, M.M.; Laakkonen, M.L.; Tilvis, R.; Pitkälä, K.H. Effects of Exercise on Cognition: The Finnish Alzheimer Disease Exercise Trial: A Randomized, Controlled Trial. *J Am Geriatr Soc.* **2016**, 64, 731-738. doi: 10.1111/jgs.14059.

33. Valenzuela, P.L.; Castillo-García, A.; Morales, J.S.; de la Villa, P.; Hampel, H.; Emanuele, E.; Lista, S.; Lucia, A. Exercise benefits on Alzheimer's disease: State-of-the-science. *Ageing Res Rev.* **2020**, 62, 101108. doi: 10.1016/j.arr.2020.101108.

34. Pardeshi, C.V.; Belgamwar, V.S. Direct nose to brain drug delivery via integrated nerve pathways bypassing the blood-brain barrier: an excellent platform for brain targeting. *Expert Opin Drug Deliv.* **2013**, 10, 957-972. doi: 10.1517/17425247.2013.790887.

35. Patharapankal, E.J.; Ajiboye, A.L.; Mattern, C.; Trivedi, V. Nose-to-Brain (N2B) Delivery: An Alternative Route for the Delivery of Biologics in the Management and Treatment of Central Nervous System Disorders. *Pharmaceutics.* **2023**, 16, 66. doi: 10.3390/pharmaceutics16010066.

36. Agrawal, M.; Saraf, S.; Saraf, S.; Antimisiaris, S.G.; Chougule, M.B.; Shoyele, S.A.; Alexander, A. Nose-to-brain drug delivery: An update on clinical challenges and progress towards approval of anti-Alzheimer drugs. *J Control Release.* **2018**, 281, 139-177. doi: 10.1016/j.jconrel.2018.05.011.

37. Alexander, A.; Saraf, S. Nose-to-brain drug delivery approach: A key to easily accessing the brain for the treatment of Alzheimer's disease. *Neural Regen Res.* **2018**, 13, 2102-2104. doi: 10.4103/1673-5374.241458.

38. Wong, C.Y.J.; Baldelli, A.; Hoyos, C.M.; Tietz, O.; Ong, H.X.; Traini, D. Insulin Delivery to the Brain via the Nasal Route: Unraveling the Potential for Alzheimer's Disease Therapy. *Drug Deliv Transl Res.* **2024**, 14, 1776-1793. doi: 10.1007/s13346-024-01558-1.

39. Li, B.S.Y.; Chan, C.W.H.; Li, M.; Wong, I.K.Y.; Yu, Y.H.U. Effectiveness and Safety of Aromatherapy in Managing Behavioral and Psychological Symptoms of Dementia: A Mixed-Methods Systematic Review. *Dement Geriatr Cogn Dis Extra*. **2021**, 11, 273-297. doi: 10.1159/000519915.

40. Cha, H.; Kim, S.; Seo, M.S.; Kim, H.S. Effects of olfactory stimulation on cognitive function and behavior problems in older adults with dementia: A systematic literature review. *Geriatr Nurs.* **2021**, 42, 1210-1217. doi: 10.1016/j.gerinurse.2021.07.003.

41. Vora, L.K.; Gholap, A.D.; Hatvate, N.T.; Naren, P.; Khan, S.; Chavda, V.P.; Balar, P.C.; Gandhi, J.; Khatri, D.K. Essential oils for clinical aromatherapy: A comprehensive review. *J Ethnopharmacol.* **2024**, 330,118180. doi: 10.1016/j.jep.2024.118180.

42. Cioanca, O.; Hritcu, L.; Mihasan, M.; Hancianu, M. Cognitive-enhancing and antioxidant activities of inhaled coriander volatile oil in amyloid β(1-42) rat model of Alzheimer's disease. *Physiol Behav.* **2013**, 120, 193-202. doi: 10.1016/j.physbeh.2013.08.006.

43. Satou, T.; Hanashima, Y.; Mizutani, I.; Koike, K. The effect of inhalation of essential oil from Rosmarinus officinalis on scopolamine-induced Alzheimer's type dementia model mice. *Flavour and fragrance journal*. **2018**, 33, 230-234. doi: 10.1002/ffj.3435

44. Okuda, M.; Fujita, Y.; Takada-Takatori, Y.; Sugimoto, H.; Urakami, K. Aromatherapy improves cognitive dysfunction in senescence-accelerated mouse prone 8 by reducing the level of amyloid beta and tau phosphorylation. *PLoS One*. **2020**, 15, e0240378. doi: 10.1371/journal.pone.0240378.

45. Habash, R.W.; Bansal, R.; Krewski, D.; Alhafid, H.T. Thermal therapy, part 1: an introduction to thermal therapy. *Crit Rev Biomed Eng*. **2006**, 34, 459-489. doi: 10.1615/critrevbiomedeng.v34.i6.20.

46. Li, L.; Zhang, X.; Zhou, J.; Zhang, L.; Xue, J.; Tao, W. Non-Invasive Thermal Therapy for Tissue Engineering and Regenerative Medicine. *Small.* **2022**, 18, e2107705. doi: 10.1002/smll.202107705.

47. Biro, S.; Masuda, A.; Kihara, T.; Tei, C. Clinical implications of thermal therapy in lifestyle-related diseases. *Exp Biol Med (Maywood)*. **2003**, 228, 1245-1249. doi: 10.1177/153537020322801023.

48. Hunt, A.P.; Minett, G.M.; Gibson, O.R.; Kerr, G.K.; Stewart, I.B. Could Heat Therapy Be an Effective Treatment for Alzheimer's and Parkinson's Diseases? A Narrative Review. *Front Physiol*. **2020**, 10, 1556. doi: 10.3389/fphys.2019.01556.

49. Von Schulze, A.T.; Deng, F.; Morris, J.K.; Geiger, P.C. Heat therapy: possible benefits for cognitive function and the aging brain. *J Appl Physiol (1985)*. **2020**, 129, 1468-1476. doi: 10.1152/japplphysiol.00168.2020.

50. Smadja, D.M.; Abreu, M.M. Hyperthermia and targeting heat shock proteins: innovative approaches for neurodegenerative disorders and Long COVID. *Front Neurosci*. **2025**, 19, 1475376. doi: 10.3389/fnins.2025.1475376.

51. Tundo, G.R.; Sbardella, D.; Ciaccio, C.; Grasso, G.; Gioia, M.; Coletta, A.; Polticelli, F.; Di Pierro, D.; Milardi, D.; Van Endert, P.; et al. Multiple functions of insulin-degrading enzyme: a metabolic crosslight? *Crit Rev Biochem Mol Biol*. **2017**, 52, 554-582. doi: 10.1080/10409238.2017.1337707.

52. Kuo, Y.Y.; Chen, W.T.; Lin, G.B.; Chen, Y.M.; Liu, H.H.; Chao, C.Y. Thermal cycling-hyperthermia ameliorates Aβ25-35-induced cognitive impairment in C57BL/6 mice. *Neurosci Lett*. **2023**, 810, 137337. doi: 10.1016/j.neulet.2023.137337.

53. Mearow, K.M.; Dodge, M.E.; Rahimtula, M.; Yegappan, C. Stress-mediated signaling in PC12 cells - the role of the small heat shock protein, Hsp27, and Akt in protecting cells from heat stress and

nerve growth factor withdrawal. *J Neurochem*. **2002**, 83, 452-462. doi: 10.1046/j.1471-4159.2002.01151.x.

54. Chen, W.T.; Kuo, Y.Y.; Lin, G.B.; Lu, C.H.; Hsu, H.P.; Sun, Y.K.; Chao, C.Y. Thermal cycling protects SH-SY5Y cells against hydrogen peroxide and β-amyloid-induced cell injury through stress response mechanisms involving Akt pathway. *PLoS One*. **2020**, 15, e0240022. doi: 10.1371/journal.pone.0240022.

55. Yin, G.; Li, L.Y.; Qu, M.; Luo, H.B.; Wang, J.Z.; Zhou, X.W. Upregulation of AKT attenuates amyloid-β-induced cell apoptosis. *J Alzheimers Dis*. **2011**, 25, 337-345. doi: 10.3233/JAD-2011-110104.

56. Ghasemi, R.; Moosavi, M.; Zarifkar, A.; Rastegar, K.; Maghsoudi, N. The Interplay of Akt and ERK in Aβ Toxicity and Insulin-Mediated Protection in Primary Hippocampal Cell Culture. *J Mol Neurosci*. **2015**, 57, 325-334. doi: 10.1007/s12031-015-0622-6.

57. Kommaddi, R.P.; Gowaikar, R.; Haseena, P.A.; Diwakar, L.; Singh, K.; Mondal, A. Akt activation ameliorates deficits in hippocampal-dependent memory and activity-dependent synaptic protein synthesis in an Alzheimer's disease mouse model. *J Biol Chem*. **2024**, 300, 105619. doi: 10.1016/j.jbc.2023.105619.

58. Hannuksela, M.L.; Ellahham, S. Benefits and risks of sauna bathing. *Am J Med*. **2001**, 110, 118-126. doi: 10.1016/s0002-9343(00)00671-9.

59. Bain, A.R.; Nybo, L.; Ainslie, P.N. Cerebral Vascular Control and Metabolism in Heat Stress. *Compr Physiol*. **2015**, 5, 1345-1380. doi: 10.1002/cphy.c140066.

60. Prieur, E.A.K.; Jadavji, N.M. Assessing Spatial Working Memory Using the Spontaneous Alternation Y-maze Test in Aged Male Mice. *Bio Protoc.* **2019**, 9, e3162. doi: 10.21769/BioProtoc.3162.

61. Kraeuter, A.K.; Guest, P.C.; Sarnyai, Z. The Y-Maze for Assessment of Spatial Working and Reference Memory in Mice. *Methods Mol Biol*. **2019**, 1916, 105-111. doi: 10.1007/978-1-4939-8994-2_10.

62. Lueptow, L.M. Novel Object Recognition Test for the Investigation of Learning and Memory in Mice. *J Vis Exp*. **2017**, 126, 55718. doi: 10.3791/55718.

63. Navarro, D.; Gasparyan, A.; Martí Martínez, S.; Díaz Marín, C.; Navarrete, F.; García Gutiérrez, M.S.; Manzanares, J. Methods to Identify Cognitive Alterations from Animals to Humans: A Translational Approach. *Int J Mol Sci*. **2023**, 24, 7653. doi: 10.3390/ijms24087653.

64. Kim, H.Y.; Lee, D.K.; Chung, B.R.; Kim, H.V.; Kim, Y. Intracerebroventricular Injection of Amyloid-β Peptides in Normal Mice to Acutely Induce Alzheimer-like Cognitive Deficits. *J Vis Exp*. **2016**, 109, 53308. doi: 10.3791/53308.

65. Yang, M.; Crawley, J.N. Simple behavioral assessment of mouse olfaction. *Curr Protoc Neurosci*. **2009**, 48, 8.24. doi: 10.1002/0471142301.ns0824s48.

66. Zhang, J.; Zhao, Z.; Sun, S.; Li, J.; Wang, Y.; Dong, J.; Yang, S.; Lou, Y.; Yang, J.; Li, W.; Li, S. Olfactory Evaluation in Alzheimer's Disease Model Mice. *Brain Sci*. **2022**, 12, 607. doi: 10.3390/brainsci12050607.

67. Jahn, H. Memory loss in Alzheimer's disease. *Dialogues Clin Neurosci*. **2013**, 15, 445-454. doi: 10.31887/DCNS.2013.15.4/hjahn.

68. Lu, R.C.; Tan, M.S.; Wang, H.; Xie, A.M.; Yu, J.T.; Tan, L. Heat shock protein 70 in Alzheimer's disease. *Biomed Res Int*. **2014**, 2014, 435203. doi: 10.1155/2014/435203.

69. Rivera, I.; Capone, R.; Cauvi, D.M.; Arispe, N.; De Maio, A. Modulation of Alzheimer's amyloid β peptide oligomerization and toxicity by extracellular Hsp70. *Cell Stress Chaperones*. **2018**, 23, 269-279. doi: 10.1007/s12192-017-0839-0.

70. Valle-Medina, A.; Calzada-Mendoza, C.C.; Ocharan-Hernández, M.E.; Jiménez-Zamarripa, C.A.; Juárez-Cedillo, T. Heat shock protein 70 in Alzheimer's disease and other dementias: A possible

alternative therapeutic. *J Alzheimers Dis Rep*. **2025**, 9, 25424823241307021.
doi:10.1177/25424823241307021

71. Abdi, A.; Sadraie, H.; Dargahi, L.; Khalaj, L.; Ahmadiani, A. Apoptosis inhibition can be
threatening in Aβ-induced neuroinflammation, through promoting cell proliferation. *Neurochem*
Res. **2011**, 36, 39-48. doi: 10.1007/s11064-010-0259-3.

72. Hoshino, T.; Suzuki, K.; Matsushima, T.; Yamakawa, N.; Suzuki, T.; Mizushima, T. Suppression of
Alzheimer's disease-related phenotypes by geranylgeranylacetone in mice. *PLoS One*. **2013**, 8,
e76306. doi: 10.1371/journal.pone.0076306.

73. Cook, D.G.; Leverenz, J.B.; McMillan, P.J.; Kulstad, J.J.; Ericksen, S.; Roth, R.A.; Schellenberg, G.D.;
Jin, L.W.; Kovacina, K.S. Craft, S. Reduced hippocampal insulin-degrading enzyme in late-onset
Alzheimer's disease is associated with the apolipoprotein E-epsilon4 allele. *Am J Pathol*. **2003**, 162,
313-319. doi: 10.1016/s0002-9440(10)63822-9.

74. Zhao, L.; Teter, B.; Morihara, T.; Lim, G.P.; Ambegaokar, S.S.; Ubeda, O.J.; Frautschy, S.A.; Cole,
G.M. Insulin-degrading enzyme as a downstream target of insulin receptor signaling cascade:
implications for Alzheimer's disease intervention. *J Neurosci*. **2004**, 24, 11120-11126. doi:
10.1523/JNEUROSCI.2860-04.2004.

75. Ahn, J.Y. Neuroprotection signaling of nuclear akt in neuronal cells. *Exp Neurobiol*. **2014**, 23,
200-206. doi: 10.5607/en.2014.23.3.200.

76. Manning, B.D.; Toker, A. AKT/PKB Signaling: Navigating the Network. *Cell*. **2017**, 169, 381-405.
doi: 10.1016/j.cell.2017.04.001.

77. Cui, W.; Tao, J.; Wang, Z.; Ren, M.; Zhang, Y.; Sun, Y.; Peng, Y.; Li, R. Neuregulin1beta1
antagonizes apoptosis via ErbB4-dependent activation of PI3-kinase/Akt in APP/PS1 transgenic
mice. *Neurochem Res*. **2013**, 38, 2237-2246. doi: 10.1007/s11064-013-1131-z.

78. Magrané, J.; Rosen, K.M.; Smith, R.C.; Walsh, K.; Gouras, G.K.; Querfurth, H.W. Intraneuronal
beta-amyloid expression downregulates the Akt survival pathway and blunts the stress response. *J*
Neurosci. **2005**, 25, 10960-10969. doi: 10.1523/JNEUROSCI.1723-05.2005.

79. Maroni, P.; Bendinelli, P.; Tiberio, L.; Rovetta, F.; Piccoletti, R.; Schiaffonati, L. In vivo heat-shock
response in the brain: signalling pathway and transcription factor activation. *Brain Res Mol Brain*
Res. **2003**, 119, 90-99. doi: 10.1016/j.molbrainres.2003.08.018.

80. Jensen, C.S.; Hasselbalch, S.G.; Waldemar, G.; Simonsen, A.H. Biochemical Markers of Physical
Exercise on Mild Cognitive Impairment and Dementia: Systematic Review and Perspectives. *Front*
Neurol. **2015**, 6, 187. doi: 10.3389/fneur.2015.00187.

81. Wu, W.; Ji, Y.; Wang, Z.; Wu, X.; Li, J.; Gu, F.; Chen, Z.; Wang, Z. The FDA-approved
anti-amyloid-β monoclonal antibodies for the treatment of Alzheimer's disease: a systematic
review and meta-analysis of randomized controlled trials. *Eur J Med Res*. **2023**, 28, 544. doi:
10.1186/s40001-023-01512-w.

82. Ornish, D.; Madison, C.; Kivipelto, M.; Kemp, C.; McCulloch, C.E.; Galasko, D.; Artz, J.; Rentz, D.;
Lin, J.; Norman, K.; et al. Effects of intensive lifestyle changes on the progression of mild cognitive
impairment or early dementia due to Alzheimer's disease: a randomized, controlled clinical trial.
Alzheimers Res Ther. **2024**, 16, 122. doi: 10.1186/s13195-024-01482-z.

83. Matsumoto, K.; Honda, K.; Kobayashi, N. Protective effect of heat preconditioning of rat liver graft
resulting in improved transplant survival. *Transplantation*. **2001**, 71, 862-868. doi:
10.1097/00007890-200104150-00007.

84. Le Bourg, É. Hormesis, aging and longevity. *Biochim Biophys Acta*. **2009**, 1790, 1030-1039. doi:
10.1016/j.bbagen.2009.01.004.

85. Le Bourg, É. Characterisation of the positive effects of mild stress on ageing and resistance to stress.
Biogerontology. **2020**, 21, 485-493. doi: 10.1007/s10522-020-09870-2.

86. Guisle, I.; Canet, G.; Pétry, S.; Fereydouni-Forouzandeh, P.; Morin, F.; Kérauden, R.; Whittington,
R.A.; Calon, F.; Hébert, S.S.; Planel, E. Sauna-like conditions or menthol treatment reduce tau

phosphorylation through mild hyperthermia. *Neurobiol Aging.* **2022**, 113, 118-130. doi: 10.1016/j.neurobiolaging.2022.02.011.

87. Crane, C.R.; Sanders, D.C. Inhalation Toxicology: VIII. Establishing Heat Tolerance Limits for Rats and Mice Subjected to Acute Exposures at Elevated Air Temperatures. *Civil Aerospace Medical Institute.* **1986**, DOT/FAA/AM-86/8.

88. Kassis, S.; Grondin, M.; Averill-Bates, D.A. Heat shock increases levels of reactive oxygen species, autophagy and apoptosis. *Biochim Biophys Acta Mol Cell Res.* **2021**, 1868, 118924. doi: 10.1016/j.bbamcr.2020.118924.

89. Slimen, I.B.; Najar, T.; Ghram, A.; Dabbebi, H.; Ben Mrad, M.; Abdrabbah, M. Reactive oxygen species, heat stress and oxidative-induced mitochondrial damage. A review. *Int J Hyperthermia.* **2014**, 30, 513-523. doi: 10.3109/02656736.2014.971446.

90. Li, L.; Tan, H.; Yang, H.; Li, F.; He, X.; Gu, Z.; Zhao, M.; Su, L. Reactive oxygen species mediate heat stress-induced apoptosis via ERK dephosphorylation and Bcl-2 ubiquitination in human umbilical vein endothelial cells. *Oncotarget.* **2017**, 8, 12902-12916. doi: 10.18632/oncotarget.14186.

91. Dalton, P.; Wysocki, C.J. The nature and duration of adaptation following long-term odor exposure. *Percept Psychophys.* **1996**, 58, 781-792. doi: 10.3758/bf03213109.

92. Kato, A.; Touhara, K. Mammalian olfactory receptors: pharmacology, G protein coupling and desensitization. *Cell Mol Life Sci.* **2009**, 66, 3743-3753. doi: 10.1007/s00018-009-0111-6. Epub 2009 Aug 4

93. Pellegrino, R.; Sinding, C.; de Wijk, R.A.; Hummel, T. Habituation and adaptation to odors in humans. *Physiol Behav.* **2017**, 177, 13-19. doi: 10.1016/j.physbeh.2017.04.006.

94. Kim, K.; Bae, J.; Jin, Y.; Moon, C. Odor habituation can modulate very early olfactory event-related potential. *Sci Rep.* **2020**, 10, 18117. doi: 10.1038/s41598-020-75263-7.

95. Bandiera, B.; Natale, F.; Rinaudo, M.; Sollazzo, R.; Spinelli, M.; Fusco, S.; Grassi, C. Olfactory stimulation with multiple odorants prevents stress-induced cognitive and psychological alterations. *Brain Commun.* **2024**, 6, fcae390. doi: 10.1093/braincomms/fcae390.

96. Okada, K.; Shimatani, K. Effect of olfactory stimulation from aromatherapy on the autonomic nervous activity during aerobic exercises. *Sci Rep.* **2024**, 14, 11198. doi: 10.1038/s41598-024-61732-w.

97. Scutigliani, E.M.; Liang, Y.; Crezee, H.; Kanaar, R.; Krawczyk, P.M. Modulating the Heat Stress Response to Improve Hyperthermia-Based Anticancer Treatments. *Cancers (Basel).* **2021**, 13, 1243. doi: 10.3390/cancers13061243.

98. Ahmed, K.; Zaidi, S.F.; Mati-Ur-Rehman.; Rehman, R.; Kondo, T. Hyperthermia and protein homeostasis: Cytoprotection and cell death. *J Therm Biol.* **2020**, 91, 102615. doi: 10.1016/j.jtherbio.2020.102615.

99. Selkoe, D.J.; Hardy, J. The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol Med.* **2016**, 8, 595-608. doi: 10.15252/emmm.201606210.

100. Karran, E.; De Strooper, B. The amyloid hypothesis in Alzheimer disease: new insights from new therapeutics. *Nat Rev Drug Discov.* **2022**, 21, 306-318. doi: 10.1038/s41573-022-00391-w.

101. Tundo, G.R.; Sbardella, D.; Ciaccio, C.; Bianculli, A.; Orlandi, A.; Desimio, M.G.; Arcuri, G.; Coletta, M.; Marini, S. Insulin-degrading enzyme (IDE): a novel heat shock-like protein. *J Biol Chem.* **2013**, 288, 2281-2289. doi: 10.1074/jbc.M112.393108.

102. Vekrellis, K.; Ye, Z.; Qiu, W.Q.; Walsh, D.; Hartley, D.; Chesneau, V.; Rosner, M.R.; Selkoe, D.J. Neurons regulate extracellular levels of amyloid beta-protein via proteolysis by insulin-degrading enzyme. *J Neurosci.* **2000**, 20, 1657-1665. doi: 10.1523/JNEUROSCI.20-05-01657.2000.

103. Farris, W.; Mansourian, S.; Chang, Y.; Lindsley, L.; Eckman, E.A.; Frosch, M.P.; Eckman, C.B.; Tanzi, R.E.; Selkoe, D.J.; Guenette, S. Insulin-degrading enzyme regulates the levels of insulin, amyloid beta-protein, and the beta-amyloid precursor protein intracellular domain in vivo. *Proc Natl Acad Sci U S A.* **2003**, 100, 4162-4167. doi: 10.1073/pnas.0230450100.

104. Tian, Y.; Jing, G.; Zhang, M. Insulin-degrading enzyme: Roles and pathways in ameliorating
cognitive impairment associated with Alzheimer's disease and diabetes. *Ageing Res Rev*. **2023**, 90,
101999. doi: 10.1016/j.arr.2023.101999.
105. Chovatiya, R.; Medzhitov, R. Stress, inflammation, and defense of homeostasis. *Mol Cell*. **2014**, 54,
281-288. doi: 10.1016/j.molcel.2014.03.030.
106. Thomas, M.P.; Lieberman J. Live or let die: posttranscriptional gene regulation in cell stress and
cell death. *Immunol Rev*. **2013**, 253, 237-252. doi: 10.1111/imr.12052.
107. Mesihovic, A.; Iannacone, R.; Firon, N.; Fragkostefanakis, S. Heat stress regimes for the
investigation of pollen thermotolerance in crop plants. *Plant Reprod*. **2016**, 29, 93-105. doi:
10.1007/s00497-016-0281-y.

Figure 1

Figure 2

Figure 3

Figure 4

A

Air pump

Flowmeter
Air

Vaporizer

Time switch

Thermometer

water bath

B

Temperature (°C) vs Time (min)

— TCSNI
--- HTSNI

Figure 5