***Mycobacterium tuberculosis* suppresses protective Th17 responses during**

infection through multiple mechanisms

Authors: Alex Zilinskas[1], Amir Balakhmet[1], Douglas Fox[1], Heyuan Michael Ni[1], Carolina

Agudelo[2], Helia Samani[1], Sarah Stanley[1,2]

Affiliations:

1. Division of Immunology and Molecular Medicine, Department of Molecular and Cell

Biology, University of California, Berkeley, Berkeley, CA.

2. Division of Infectious Disease and Vaccinology, School of Public Health, University of

California, Berkeley, Berkeley, CA.

Short title: M. tuberculosis subverts protective Th17s

Abstract

Mycobacterium tuberculosis (Mtb) causes more deaths annually than any other pathogen, yet an effective vaccine remains elusive. IFN-γ–producing CD4+ T cells are necessary but insufficient for immune protection. Data from humans shows that the development of IL-17a producing Th17 T cells correlates with protection against infection, however not all individuals develop a Th17 response. In mouse models, experimental vaccines can elicit protective Th17 cells, but Th17s are rare in primary infection. Here, we identify factors suppressing Th17 responses during primary Mtb infection. First, using Tbet deficient mice, we demonstrate that Mtb drives a Th1 response that is only partially protective and limits the production of protective Th17 cells in an IFN-γ independent manner. Next, we reveal that the ESX-1 type VII alternative secretion system in Mtb suppresses Th17 responses by inhibiting IL-23 production in dendritic cells. These findings define a new function of the ESX-1 secretion system in Mtb virulence, a long-standing question in tuberculosis research.

Introduction

Mycobacterium tuberculosis (Mtb), the causative agent of tuberculosis (TB), remains one of the leading infectious disease killers worldwide, with 10.6 million new cases and 1.6 million deaths reported in 2021 (World Health Organization, 2022). The current vaccine, Mycobacterium bovis bacillus Calmette-Guérin (BCG), protects against childhood TB, but has highly variable efficacy, wanes over time, and provides minimal protection against adult pulmonary TB (Fine, 1995; World Health Organization, 2018; Kumar, 2021). Although there are 17 vaccine candidates currently in clinical trials, achieving robust vaccine elicited immunity remains a major challenge (Zhou and Zhang, 2023).

Interferon-γ (IFN-γ) produced by type 1 T helper (Th1) cells is essential for control of Mtb infection both in animal models and in humans (Flynn et al., 1993; Condos et al., 1998; Casanova and Abel, 2002). However, the MVA85A vaccine, despite inducing robust Th1 responses as intended, failed to provide protection against Mtb infection (Ndiaye et al., 2015). Patients with active TB can have high levels of IFN-γ in the lungs despite being unable to control infection (Ledesma et al., 2011). Thus, Th1 cells are necessary but not sufficient for controlling Mtb infection suggesting that other factors are important for effective immunity. Without a comprehensive understanding of which types and specific subsets of T helper cells essential to limit Mtb infection, developing an effective Mtb vaccine with consistent high efficacy will remain elusive.

Type 17 T helper (Th17) cells are known for their association with mucosal sites, including the lungs, where they have been shown to provide protection against several bacterial and fungal respiratory pathogens (Guglani and Khader, 2010). This protection relies on IL-17a production by Th17 cells (Gaffen, 2009). Many factors determine whether Th17 cells are elicited by

infection. Antigen-presenting cells (APCs) drive the polarization of naïve T cells into Th17 cells

by producing cytokines including IL-1β, IL-6, TGF-β, and IL-23. Activation of the RORγT

transcription factor is essential for Th17 polarization (Lazarevic et al., 2011). There is also

evidence that the route of infection with a bacterial pathogen influences whether a Th17

response is elicited. Infection with *Listeria monocytogenes* via the intranasal, but not

intravenous route is required to observe a Th17 response to infection (Pepper et al., 2010).

Th17 differentiation is opposed by Th1 differentiation via multiple mechanisms (Huber and

Farrar, 2011). Tbet, the transcription factor essential for commitment to the Th1 lineage, has

been shown to prevent Runx1 activation, thereby preventing Runx1-mediated transactivation of

RORγT and therefore Th17 differentiation (Lazarevic et al., 2011). Additionally, IFN-γ, the

signature cytokine of Th1s, inhibits Th17 differentiation and function via activation of STAT1,

independent of Tbet (Yeh et al., 2014). Finally, production of type I IFN also can suppress Th17

differentiation (Zhang et al., 2011). Thus, several hallmarks of Mtb infection, namely strong Th1

responses, IFN-γ production, and type I interferon production tend to weigh against Th17s.

Increasing evidence that Th17 cells correlate with protection against Mtb infection both in the

non-human primate model (Gideon et al., 2015) and in humans (Perreau et al., 2013; Ogongo et

al., 2021; Nathan et al., 2021; Sun et al., 2024) suggest that Th17 cells are an important

component of protective immunity to Mtb. In mice, Th17 cells are rarely observed during primary

Mtb infection, and the conditions under which Th17 cells are induced during infection remain

unclear (Khader et al., 2007; Gopal et al., 2014; Segueni et al., 2016; Aguilo et al., 2016; Jong

et al., 2022). However, experimental TB vaccines, particularly via a mucosal route, elicit Th17

cells that can confer protection against subsequent challenge in mice (Khader et al., 2007; Jong

et al., 2022), demonstrating that Th17 cells can be protective against Mtb infection. Why Mtb

infection does not robustly elicit Th17 cells during primary infection, despite causing infection at

mucosal surfaces via airway infection, is unclear.

The ESX-1 type VII alternative secretion system is a fundamental determinant of Mtb virulence

(Stanley et al., 2003). The ESX-1 secretion system has been demonstrated to subvert and/or

manipulate immune responses, including by perforating phagosomal membranes, inducing

detrimental host type I interferon, and inducing autophagy (Stanley et al., 2007; Simeone et al.,

2012; Romagnoli et al., 2012). As much of the work examining ESX-1 function has been

conducted in vitro, we lack a clear understanding of all the factors by which ESX-1 mutants are

attenuated during in vivo infection. Interestingly, the Mtb lipid phthiocerol dimycocerosate

(PDIM) is also required for virulence and has been linked to ESX-1 function through induction of

type I IFN, which can suppress Th17 responses (Cox et al., 1999; Barczak et al., 2017). In

addition, the ESX-1 secretion system was shown to suppress production of IL-12 p40 (Stanley

et al., 2007), a subunit shared with IL-23, which promotes Th17 differentiation. These data

suggest that Mtb virulence factors, including ESX-1 and PDIM may suppress the emergence of

a protective Th17 response during Mtb infection.

Here we show that the induction of Th17 cells during Mtb infection is suppressed by multiple

mechanisms. First, we show that the absence of a Th17 response during primary infection is not

dependent on duration or route of infection. Instead, we show that mice lacking Tbet (*Tbx21*$^{-/-}$)

produce Th17 cells after Mtb infection, and that these cells provide IL-17a-dependent protection

to early Mtb infection. The suppression of Th17 production by Th1 cells is independent of IFN-γ.

Furthermore, we observe a strong Th17 response during infection with ΔESX-1 or PDIM-lacking

Mtb in naïve mice, suggesting that these virulence factors function to suppress Th17

differentiation. Surprisingly, the ability of ESX-1/PDIM to suppress Th17 differentiation is not

related to their enhanced induction of type I IFN. Instead, we show that during infection with

these strains type 2 conventional dendritic cells (CD11b$^+$ DCs, cDC2s), which are known to

support Th17 differentiation, produce higher levels of IL-23 secretion in draining lymph nodes of

Mtb infected mice (Schlitzer et al., 2013). These results demonstrate that Mtb utilizes the ESX-1

secretion system and PDIM to bias host T helper response towards Th1s and away from Th17s

in a manner that limits but permits Mtb infection.

Results

Th17 induction during *M. tuberculosis* infection does not depend on duration or route of

infection

We first tested whether the duration or route of infection has an impact on the generation of

Th17 cells during Mtb infection. To test for induction of Th17 cells, we infected wild type mice

with ~100 CFU of WT *M. tuberculosis* via the aerosol route, a standard dose for the mouse

model (Soldevilla et al., 2022). As previously published by multiple groups, we do not observe

Th17 cells during the first month after Mtb infection of mice (Khader et al., 2007; Gopal et al.,

2014; Segueni et al., 2016; Aguilo et al., 2016; Jong et al., 2022). At 4 weeks post infection, we

observed the expected plateau of CFU in the lungs of infected mice indicating the onset of T

cell-based control of infection (Figure 1A). With stimulation using peptide pools from model

antigens ESAT-6 and Antigen 85B (Ag85B) (Andersen et al., 1995; Bekmurzayeva et al., 2013;

Babaki et al., 2017), we observed only IFN-γ^+ CD4$^+$ (Th1) T cells (Figure 1B) and no IL-17a$^+$

CD4$^+$ (Th17) T cells at this or earlier timepoints (Figure 1C). Th17 cells have been observed in

humans, at times presumably long after initial infection (Perreau et al., 2013; Ogongo et al.,

2021; Nathan et al., 2021; Sun et al., 2024). We therefore tested whether Th17 cells arise at late

timepoints after infection in mice. Mice continued to only have a Th1 response up to day 170

post infection (Figures 1D-E). It was previously reported that infection with the virulent HN878

"Beijing" strain of the L2 lineage resulted in induction of Th17 cells (Gopal et al., 2014).

However, we did not observe a significant increase in IL-17a producing T cells in murine lungs

when we infected with a Beijing strain at 3 weeks post infection (Supplementary Figure 1).

The route of bacterial infection can impact which T helper cell types are induced. Infection via

the intranasal route has been shown to elicit Th17 cells in response to infection with *Listeria*

Figure 1. Lack of Th17 induction during WT *M. tuberculosis* infection is not due to duration or route of infection. (A-C) WT B6 mice were aerosol infected with WT *M. tuberculosis* Erdman strain and evaluated at 21 days post infection (dpi) for **(A)** bacterial burdens in lungs by CFU from lungs **(B)** IFN-γ⁺ CD4⁺ T cells and **(C)** IL-17a⁺ CD4⁺ T cells from lungs using flow cytometry. Cytokines were measured using ICS after restimulation with Ag85b or ESAT-6 peptide pools. **(D-E)** Mice were infected and analyzed at 170 dpi for **(D)** IFN-γ⁺ CD4⁺ T cells and **(E)** IL-17a⁺ CD4⁺ T cells in lungs using flow cytometry. **(F-H)** WT B6 mice were either aerosol infected or intranasally infected with WT *M. tuberculosis* Erdman strain and evaluated 28 dpi for **(F)** bacterial burdens in lungs by CFU, **(G)** IFN-γ⁺ CD4⁺ T cells and **(H)** IL-17a⁺ CD4⁺ T cells in lungs by flow cytometry. Cytokines were measured using ICS after restimulation with ESAT-6 peptide pool. **(I)** WT B6 mice were aerosol infected with WT *M. tuberculosis* and treated with PBS control, recombinant murine IL-17a (IL-17a) or recombinant murine IL-17a fused to murine serum albumin (MSA-IL-17a) daily either using the intranasal (i.n.) or intraperitoneal (i.p.) routes between days 11 to 20 post infection. CFU in lungs was measured at 21 dpi. Results are representative of at least 2 biological replicates. For all graphs, each dot represents an individual mouse. *p<0.05, **p<0.01 (unpaired nonparametric Mann-Whitney U test).

monocytogenes and Group A *Streptococcus* (Pepper et al., 2010; Dileepan et al., 2011). We

therefore tested whether the intranasal route of infection results in a Th17 response in the lungs

of Mtb infected mice. Mice were infected intranasally such that we inoculated ~100 CFU of WT

Mtb into the lungs as measured on day 1 after infection. After four weeks of infection, mice had

comparable levels of bacteria in the lungs between infection routes (Figure 1F). At this timepoint

we again observed only IFN-γ producing CD4 T cells and very few IL-17a producing CD4 T cells

in the lungs (Figures 1G-H). Thus, the route of infection with Mtb does not seem to impact the

dominant Th1-biased response observed during primary infection of naïve mice with Mtb. We

next investigated whether exogenous IL-17a treatment could provide protection against WT Mtb

infection during primary infection. WT mice aerosol infected with WT Mtb were intranasally

administered with either PBS control or 1 µg recombinant murine IL-17a daily between days 11

to 20 post infection for a total of 10 doses (Figure 1I). WT mice intranasally (i.n.) administered

1 µg recombinant murine IL-17a daily were slightly more protected with ~3-fold reduction in lung

CFU compared to WT mice given vehicle control. However, we observed no effect of IL-17a

when it was administered via the intraperitoneal (i.p.) route. Cytokines are known to have short

half-lives in vivo. Fusing a cytokine to serum albumin can increase its half-life, permitting greater

efficacy (Tan et al., 2022). We therefore fused IL-17a to murine serum albumin (MSA-IL-17a)

and administered this construct either i.p. or i.n. at equimolar quantities compared to free IL-17a

administered dosages. We found that i.n. delivery of this construct was more effective than

delivery of IL-17a, with ~5-fold decrease in CFU observed in the lungs. In addition, fusion to

albumin allowed us to see efficacy with i.p. administration, resulting in a ~2-fold decrease in lung

CFU. These results indicate that IL-17a can induce protective immunity to Mtb in the context of

a primary infection, and that this may have potential as a therapeutic approach.

$Tbet^{-/-}$ mice are protected by a robust Th17 response during early *M. tuberculosis*

infection

Because Th1 responses are known to inhibit Th17 responses (Lazarevic et al., 2011; Yeh et al.,

2014), we next tested whether Th1 differentiation masks a protective Th17 response during Mtb

infection. *Tbx21*, also referred as Tbet, encodes the essential transcription factor that drives the

Th1 lineage (Lazarevic et al., 2011). Interestingly, $Tbet^{-/-}$ mice lack Th1 T cells but are not as

susceptible to infection with Mtb as are $Ifng^{-/-}$ mice (Sullivan et al., 2005; Flynn et al., 1993). As

Figure 2. *Tbet⁻ᐟ⁻* **mice are protected from susceptibility to Mtb infection by the emergence of Th17 cells.** WT B6, *Ifng⁻ᐟ⁻*, *Tbet⁻ᐟ⁻*, and *Tbet⁻ᐟ⁻Il17a⁻ᐟ⁻* mice were aerosol infected with WT *M. tuberculosis* Erdman strain. *Tbet⁻ᐟ⁻* and *Tbet⁻ᐟ⁻Il17a⁻ᐟ⁻* mice were littermate-controlled. Mice were infected with Mtb and evaluated 21 days post infection for **(A)** CFU in the lungs of indicated genotypes or **(B-F)** cellular immune responses during flow cytometry. CD4⁺ T cells were gated on single, live, MHC-II⁻, Ly6G⁻, CD3⁺, γδTCR⁻, CD4⁺, CD8a⁻ cells for analysis of IFN-γ **(B,C)**, IL-17 **(B,D)** and RORγT **(E,F)**. Results in B-E are representative of 3 independent experiments. *, p < 0.05; **, p < 0.01 (unpaired nonparametric Mann-Whitney U test).

expected, we found that *Ifng⁻ᐟ⁻* mice were extremely susceptible to infection with Mtb, however

Tbet⁻ᐟ⁻ had CFU in the lungs equivalent to wild type mice at 21 days after infection (Figure 2A).

Only WT mice had IFN-γ producing CD4 T cells in the lungs at this timepoint (Figures 2B, 2C).

Importantly, *Tbet⁻ᐟ⁻* mice exhibited a robust population of IL-17a producing CD4 T cells at 3

weeks post infection, confirming that Tbet prevents Th17 differentiation during Mtb infection

(Figures 2B, 2D). Importantly, IFN-γ deficient mice did not exhibit elevated levels of IL-17a

producing CD4 T cells demonstrating that IFN-γ production is not the mechanism by which Th1

T cells limit a Th17 response during Mtb infection (Figure 2D).

A previous study demonstrated that Tbet constrains effective T cell responses by promoting the

differentiation of terminally differentiated, non-protective intravascular Th1 T cells (Sallin et al.,

2017). This study also used antibody blockade of IL-17a in Tbet deficient mice to conclude that

IL-17a provides minimal protection in this setting (Sallin et al., 2017). However, given that

antibody blockade is limited by its short duration and possibly incomplete inhibition, we tested

whether genetically ablating IL-17a production in Tbet mice enhanced their susceptibility to Mtb

infection. We therefore infected $Tbet^{-/-}$ and $Tbet^{-/-}Il17a^{-/-}$ ($Il17a^{icre/icre}$ referred to as $Il17a^{-/-}$)

littermate-controlled mice with Mtb and measured CFU and T cell responses in the lungs after 3

weeks of infection. $Tbet^{-/-}Il17a^{-/-}$ mice were highly susceptible to Mtb infection with a 10-fold

increase in lung CFU compared to WT and $Tbet^{-/-}$ mice, similar to the susceptibility observed in

$Ifng^{-/-}$ mice (Figure 2A). $Tbet^{-/-}Il17a^{-/-}$ mice completely lacked both IFN-γ producing (Figure

2C) and IL-17a producing CD4 T cells (Figure 2D). Notably, $Tbet^{-/-}Il17a^{-/-}$ mice still generated

Th17 cells, as evidenced by the presence of the Th17 master transcription factor RORγT

(Lazarevic et al., 2011), despite their inability to produce IL-17a (Figures 2E, 2F). Together,

these results indicate that IL-17a production by Th17 cells confers significant protection

against Mtb infection in $Tbet^{-/-}$ mice, at least during early stages of infection.

***M. tuberculosis* utilizes ESX-1 and PDIM to limit Th17 response in mice**

We next examined whether Mtb utilizes virulence factors to alter the host T helper response.

The ESX-1 secretion system and the complex lipid PDIM are both known to influence

production of cytokines known to impact Th17 differentiation (Stanley et al., 2007; Barczak et

al., 2017; Hinman et al., 2021). Furthermore, these systems have been proposed to operate in

the same virulence promoting pathway (Barczak et al., 2017; Augenstreich et al., 2017). We

therefore tested whether ESX-1 and PDIM might promote virulence in part by suppressing Th17

differentiation (Figure 3A). To test for a role of the ESX-1 secretion system, we infected mice

Figure 3. *M. tuberculosis* ESX-1 and PDIM virulence factors suppress Th17 response in mice. WT B6 mice were aerosol infected with indicated strains and evaluated at 21 days post infection. **(A)** Representative flow cytometry plots (cells gated on single, live, MHC-II⁻, Ly6G⁻, CD3⁺, γδTCR⁻, CD4⁺, CD8a⁻) of IFN-γ⁺ and IL-17a⁺ CD4⁺ T cell populations. 21 days post infection, mice were sacrificed and evaluated ESAT-6 peptide pool stimulated lung homogenates for **(B)** IFN-γ⁺ CD4⁺ T cells and **(C)** IL-17a⁺ CD4⁺ T cells by flow cytometry. **(E)** Littermate-controlled WT B6 and *Il17a⁻/⁻* mice were aerosol infected with indicated strains at 21 days post infection with lung homogenate plated for CFU. Results are representative of 2 independent experiments. *, p < 0.05; **, p < 0.01 (unpaired nonparametric Mann-Whitney U test).

with Δ*eccC1,* which lacks an ATPase required for ESX-1 function (ΔESX-1) (Crosskey et al.,

2020) (Supplementary Figure 2). We also utilized a mutant with an inactivating transposon

insertion in *fadD28*, a gene required for PDIM biosynthesis (*fadD28:tn*, PDIM-lacking) (Cox et

al., 1999) (Supplementary Figure 2). As expected, ΔESX-1 and PDIM-lacking mutants were

highly attenuated during infection at 3 weeks, exhibiting a 100-fold reduction in bacterial burden

while complemented strains had similar CFU as WT Mtb (Supplementary Figure 3).

Interestingly, ESAT-6 peptide pool stimulation of lung cells in ΔESX-1 and PDIM-lacking Mtb

infected mice had an ~5-fold reduction in %IFN-γ$^+$ of CD4$^+$ T cells (Th1s) (Figure 3B) and ~2-

fold increase in %IL-17a$^+$ of CD4$^+$ T cells (Th17s) (Figure 3C) compared to complements of

either mutant or WT Mtb. Similar results were seen with Ag85b peptide pool stimulation

(Supplementary Figure 4). We tested if the induced Th17s are providing protection mediated by

IL-17a by infecting littermate-controlled WT B6 and *Il17a*$^{-/-}$ mice with either WT, ΔESX-1, or

complemented ΔESX-1 Mtb, or either WT, PDIM-lacking, or complemented PDIM-lacking Mtb.

Under the conditions where Th17s are highly induced, mice infected with either ΔESX-1 or

PDIM-lacking Mtb, the *Il17a*$^{-/-}$ mice had ~3-5 fold higher CFU than WT mice (Figures 3D, 3E).

These results indicate that Mtb utilizes ESX-1 and PDIM to prevent the induction of a Th17

response that enhances protection against Mtb infection.

Type I interferons boost Th1 induction and do not impact Th17 induction

As ESX-1 and PDIM are responsible for the host type I interferon response to Mtb infection

(Stanley et al., 2007; Barczak et al., 2017), and type I IFN is known to suppress Th17 responses

(Zhang et al., 2011), we investigated the impact of type I interferon signaling on Th17 induction

during Mtb infection. We examined the effect of type I IFN both in WT B6 mice, and in *Sp140*$^{-/-}$

mice that are highly susceptible to infection because of an excessive type I IFN response to

infection (Ji et al., 2019). To abrogate type I IFN in each background we used mice deficient for

the type I IFN receptor IFNAR. WT, *Ifnar1*$^{-/-}$, *Sp140*$^{-/-}$, *Sp140*$^{-/-}$*Ifnar1*$^{-/-}$, *Ifng*$^{-/-}$, and *Tbet*$^{-/-}$ mice

were aerosol infected with WT Mtb. As expected, both *Sp140*$^{-/-}$ and *Ifng*$^{-/-}$ mice had up to 10-

fold higher CFU than WT mice while *Ifnar1*$^{-/-}$, *Sp140*$^{-/-}$*Ifnar1*$^{-/-}$, and *Tbet*$^{-/-}$ mice were as

resistant as WT mice at 3 weeks post infection (Figure 4A). *Sp140*$^{-/-}$ mice had an almost 2-fold

increase in %IFN-γ$^+$ of CD4$^+$ (Th1) T cells compared to WT mice, *Ifnar1*$^{-/-}$ had a 2-fold reduction

in %IFN-γ$^+$ of CD4$^+$ (Th1) T cells compared to WT mice, *Sp140*$^{-/-}$*Ifnar1*$^{-/-}$ mice had a slightly

stronger 4-fold reduction in %IFN-γ$^+$ of CD4$^+$ (Th1) T cells compared to *Sp140*$^{-/-}$ mice (Figures

4B-D). Importantly, deletion of IFNAR had no impact on Th17 induction in either WT or *Sp140*$^{-/-}$

Figure 4. Type I interferon signaling boosts Th1 induction and does not impact Th17 induction. WT B6, *Ifnar*[−/−], *Sp140*[−/−], *Sp140*[−/−]*Ifnar*[−/−], *Ifng*[−/−], and *Tbet*[−/−] mice were aerosol infected with WT *M. tuberculosis* Erdman strain. **(B, C)** Representative flow cytometry plots (cells gated on single, live, MHC-II[−], Ly6G[−], CD3[+], γδTCR[−], CD4[+], CD8a[−]) of IFN-γ[+] and IL-17a[+] CD4[+] T cells of these populations. 21 days post infection, lung homogenates were measured for **(A)** CFU and analyzed via flow cytometry for **(D)** IFN-γ[+] CD4[+] T cells and **(E)** IL-17a[+] CD4[+] T cells. Results in A, D, and E are representative of 2 independent experiments. *, $p < 0.05$; **, $p < 0.01$ (unpaired nonparametric Mann-Whitney U test).

mice, although we observed the expected increase in Th17 cells in Tbet deficient animals

(Figure 4E). These results indicate that type I interferon signaling only impacts the induction of

Th1s with positive correlation while type I interferon signaling does not influence Th17 induction,

suggesting that induction of type I IFN by ESX-1/PDIM is not the mechanism by which these

virulence factors suppress Th17 differentiation.

ESX-1 and PDIM alter conventional dendritic cell cytokine signaling in the mediastinal

lymph node

Previous work suggested that ESX-1 mutants fail to suppress IL-12 p40, a subunit shared

between IL-12 and IL-23, expression from infected macrophages in vitro (Stanley et al., 2007).

Figure 5. ESX-1 and PDIM alter conventional dendritic cell cytokine signaling in the mediastinal lymph node. WT B6 mice were aerosol infected with indicated strains and analyzed at 21 days post infection **(A,C)** Flow cytometry schematic for gating of **(A)** cDC1 (gated on single, live, CD3⁻, CD19⁻, CD11c⁺, MHC-II⁺, XCR1⁺, CD11b⁻) and **(B)** cDC2 (gated on single, live, CD3⁻, CD19⁻, CD11c⁺, MHC-II⁺, XCR1⁻, CD11b⁺) populations of mediastinal lymph nodes with representative flow cytometry plots of IL-12 p35⁺ and IL-23 p19⁺ gating of cDC1 and cDC2 populations respectively. 21 days post infection, mediastinal lymph nodes from WT B6 mice were aerosol infected with either WT, ΔeccC1 (ΔESX-1), complemented ΔeccC1::eccC1, transposon-inserted fadD28::tn (PDIM-lacking),or complemented transposon-inserted fadD28::tn + pfadD28 *M. tuberculosis* Erdman strain were extracted and analyzed for IL-12 p35⁺ cDC1s **(C)** and IL-23 p19⁺ cDC2s **(D)** by flow cytometry. Results in B and D are representative of 2 independent experiments. *, p < 0.05; **, p < 0.01 (unpaired nonparametric Mann-Whitney U test).

Because IL-23 promotes Th17 responses, and ESX-1 and PDIM are responsible for the lack of

a Th17 response during Mtb infection, we next investigated if ESX-1 and PDIM suppress IL-23

production by dendritic cells during in vivo infection. We infected mice with ~100 CFU via the

aerosol route and harvested mediastinal lymph nodes at 21 days post infection. We examined

IL-12 and IL-23 production by conventional dendritic cells using ICS. To clearly distinguish

between levels of these two cytokines we used ICS for the cytokine specific subunits: IL-12 p35

and IL-23 p19. First, we found that type 1 conventional dendritic cells (cDC1s), known to be

specialized for IL-12 production, produce robust levels of IL-12 p35 during infection with WT Mtb

(Figures 5A, 5B). However, both ESX-1 and PDIM deficient strains elicited significantly lower

levels of IL-12 p35 from these cells (Figures 5A, 5B), consistent with the lower bacterial burdens

in the lungs of mice infected with these mutants (Supplementary Figure 3). During infection with

wild-type bacteria, we observed very little IL-23 p19 production by type 2 conventional dendritic

cells (cDC2s) in the draining mediastinal lymph node (Figures 5C, 5D). Importantly, ESX-1 and

PDIM-lacking Mtb infection resulted in a dramatic increase in the percentage of cDC2s that

produce IL-23 p19 in mediastinal LNs (Figures 5C, 5D). We saw very little to no IL-23 p19

expression in cDC1s and IL-12 p35 expression in cDC2s with no differences between WT or

ESX-1 and PDIM deficient strains (Supplementary Figure 5). These results indicate that the

ESX-1 and PDIM virulence factors impact naïve T cell differentiation at the draining mediastinal

lymph node by promoting IL-12, a Th1-polarizing cytokine, and limiting IL-23, a Th17-polarizing

cytokine, expression in cDC1s and cDC2s respectively.

268　Discussion

Immunity against *M. tuberculosis* (Mtb) requires T helper cell responses and IFN-γ as

demonstrated by susceptibility of *Rag1*$^{-/-}$ (Lai et al., 2024) and *Ifng*$^{-/-}$ (Flynn et al., 1993) mice,

and high co-morbidity between tuberculosis and low CD4 T helper cell counts in humans with

AIDS (Amelio et al., 2019; Geremew et al., 2020). The requirement of a highly efficacious Mtb

vaccine that elicits a T helper response strong enough to provide protection against Mtb

infection remains desired. The MVA85A vaccine candidate elicited a strong Th1 response yet

failed to provide protection against Mtb infection (Ndiaye et al., 2015) illustrating that a strong

Th1 response is not enough to control Mtb infection. The development of a highly efficacious

Mtb vaccine requires a rigorous understanding of which T helper cell types and subsets provide

protection against Mtb infection. The recent reports of Th17s correlating to protection against

Mtb infection in non-human primates (Gideon et al., 2015) and humans (Perreau et al., 2013;

Ogongo et al., 2021; Nathan et al., 2021; Sun et al., 2024) suggest a protective role for Th17s in

controlling Mtb infection. Herein, we showed that Th17s can provide protection that is IL-17a-

dependent in mice, Mtb's ESX-1 secretion system and PDIM enhance infection at least in part

by preventing a Th17 response in mice, and exogenous IL-17a is protective against Mtb

infection.

We have previously reported that infection in unvaccinated mice does not elicit a Th17

response, only a Th1 response, and that an intranasally administered cyclic dinucleotide

adjuvant Mtb protein subunit vaccine elicits enhanced protection to Mtb infection in mice

mediated by Th17s and in-particular IL-17a (Van Dis et al., 2018; Jong et al., 2022). There have

been conflicting reports about Th17s and in-particular IL-17a providing protection against Mtb

infection in the murine model of tuberculosis (Khader et al., 2007; Gopal et al., 2014; Segueni et

al., 2016; Aguilo et al., 2016; Jong et al., 2022). Here, we report that Mtb prevents induction of

Th17s during primary infection via the virulence factors ESX-1 and PDIM. Others have shown

that intranasal vaccination of BCG, which lacks a functional ESX-1, also induced Th17s that

provide protection against Mtb infection (Aguilo et al., 2016).

ESX-1 and PDIM have both been shown to be required for the permeabilization of *M.*

tuberculosis-containing phagosomes and therefore induce the type I interferon response

(Barczak et al., 2017; Stanley et al., 2007; Augenstreich et al., 2017; Watson et al., 2015). Thus,

we investigated how impactful the type I interferon response is on the induction of Th1s and

Th17s during WT *M. tuberculosis* infection. We found that the type I interferon response boosts

the magnitude of the Th1 response but does not impact the induction of Th17s during *M.*

tuberculosis infection. Additionally, we show that ESX-1 and PDIM promote IL-12 and limit IL-23

expression in conventional dendritic cells in the draining mediastinal lymph node during *M.*

tuberculosis infection. Future work will focus on which immune pathways impacted by ESX-1

and PDIM gate-keep the induction of Th1 vs Th17 responses.

One possible answer for why humans induce Th1s and Th17s to Mtb infection while mice only

induce Th1s might be because humans evolved cDC2s capable of expressing Th1-polarizing

cytokine IL-12 and Th17-polarizing cytokines IL-1β, IL-6, and IL-23 (Rojas et al., 2017), and

mice have evolved cDC2s incapable of IL-12 expression but still capable of expressing Th17-

polarizing cytokines (Schlitzer et al., 2013).

A method of reliably inducing Th17s in mice during wild-type Mtb infection needs to be

established to effectively study the role of Th17-mediated protection. Mice lacking Tbet (as

known as *Tbx21*) respond to WT Mtb infection with a Th17 response and provide protection

mediated by IL-17a during early infection. *Tbet*$^{-/-}$ mice can be used as a model to effectively

study how Th17s can provide protection against early Mtb infection in the absence of Th1s. We

show that exogenous IL-17a treatment promotes control of Mtb infection indicating that there

are some cells capable of sensing IL-17a and responding in a beneficial way for the host. The

mechanisms by which IL-17a signaling promotes clearance of Mtb remains unknown. High Th17

plasticity permits Th17s to express a wide variety of T helper subset profiles and can range from

pro-inflammatory Th17.1s that co-express IFN-γ and IL-17a to anti-inflammatory Tr1-like Th17s

that co-express IL-10 and IL-17a (Stadhouders et al., 2018). Moving forward, it will be important

to delineate in depth which Th17 subsets and which mechanisms downstream of IL-17 receptor

signaling are important for controlling Mtb infection.

We have shown that Mtb utilizes the ESX-1 secretion system and lipid virulence factor PDIM to

prevent Th17 induction during infection. Furthermore, ESX-1 and PDIM promotes expression of

Th1-polarizing cytokine IL-12 in cDC1s and limits expression of Th17-polarizing cytokine IL-23

in cDC2s in the draining mediastinal lymph node. The type I interferon response induced by

ESX-1 and PDIM only boosts Th1 induction and does not impact Th17 induction. Additionally,

we have developed a model of inducing Th17s in mice (*Tbet*$^{-/-}$ mice) during early Mtb infection

which provide protection via IL-17a. And lastly, exogenous IL-17a treatment provides protection

against Mtb infection. With this information, our understanding of how Mtb alters the host

immune response to limit control of infection, and our development of models for studying Th17-

mediated protection against Mtb infection provide insights to guide future Mtb vaccine design

strategies.

Materials and Methods

Ethics statement

All procedures involved with the use of mice were approved by the University of California,

Berkeley's Institutional Animal Care and Use Committee (protocol 2015-09-7979). All protocols

conform to federal regulations, the National Research Council Guide for the Care and Use of

Laboratory Animals, and the Public Health Service Policy on Humane Care and Use of

Laboratory Animals.

Bacterial Culture

Mycobacterium tuberculosis strain Erdman was grown in Middlebrook 7H9 liquid medium

supplemented with 10% oleic acid, albumin, dextrose, and catalase (OADC) (BD 212351), 0.4%

glycerol, and 0.05% Tween 80 or on solid 7H10 agar plates supplemented with 10% OADC and

0.4% glycerol. Frozen stocks of *M. tuberculosis* were made from a single culture and used for all

experiments.

Mice

C57BL/6J (no. 000664), *Ifng*$^{-/-}$ (B6.129S7-*Ifng*^{tm1Ts}/J, no. 002287), *Tbet*$^{-/-}$ (*Tbx21*$^{-/-}$, B6.129S6-

Tbx21^{tm1Glm}/J, no. 004648), and *Il17a*$^{icre/icre}$ (*Il17a*$^{-/-}$, B6.129(SJL)-*Il17a*$^{tm1.1(icre)Stck}$/J, no. 035717)

mice were obtained from The Jackson Laboratory and bred in-house. *Ifnar1*$^{-/-}$, *Sp140*$^{-/-}$, and

Sp140$^{-/-}$*Ifnar1*$^{-/-}$ mice were obtained from the Vance laboratory (UC Berkeley) and bred in-

house. Mice were age matched for all experiments.

Aerosol challenge experiments with *M. tuberculosis*

Mice between the age of 8 to 12 weeks old were infected via the aerosol route with *M.*

tuberculosis strain Erdman using a nebulizer and full-body inhalation exposure system (Glas-

Col, Terre Haute, IN). A total of 9 mL of diluted aerosol stock in MiiliQ water was loaded into the

nebulizer to deliver 30-100 bacteria per mouse as measured by whole lung CFU 1 day post

infection.

Intranasal challenge experiments with *M. tuberculosis*

Mice between the age of 8 to 12 weeks old were infected with 40 uL of *M. tuberculosis* strain

Erdman diluted in PBS administration for mice to inhale 30-100 bacteria per mouse as

measured by whole lung CFU 1 day post infection.

Flow Cytometry

Whole lungs were homogenized using GentleMACS C tubes (Miltenyi Biotec 130-093-237) in

complete RPMI-1640 media supplemented with 10% FBS, 1 mM sodium pyruvate (Sigma

Aldrich S8636), 0.01 mM HEPES, 1% Glutamax (Thermo Scientific 35050061), 1% non-

essential amino acids (Gibco 11140-050), and 55 µM 2-mercaptoethanol (Gibco 21985-023)

containing Liberase TM (Sigma Aldrich 05401119001) and DNase I (Sigma Aldrich

11284932001). Cells were stained with live/dead stain (ThermoFisher L23101), Fc block

(Biolegend 101320), and surface stained with antibodies for CD3, Ly6G, CD4 (BD 740268,

551460, and 569182 respectively), MHC-II, pan-γδTCR, and CD8a (Biolegend 107605, 118124,

and 100714 respectively) in Brilliant stain buffer (BD 566349). Cells were fixed and

permeabilized with Foxp3 staining kit (Invitrogen 00-5523-00) before intracellular staining for

RORγT, Foxp3 (BD 564723 and 562996 respectively), Tbet, IFN-γ, and IL-17a (Biolegend

644817, 505814, and 506903 respectively). Data were collected using a BD Symphony A3 flow

cytometer (BD) and analyzed using FlowJo Software (BD).

Mediastinal lymph nodes were homogenized using complete RPMI-1640 containing Liberase

TM (Sigma Aldrich 05401119001) and DNase I (Sigma Aldrich 11284932001) with sterile

plunger of 3 mL syringe to pass dissociated tissue through 100 micron cell strainer. Cells were

stained with live/dead stain (ThermoFisher L34966), Fc block (Biolegend 101320) and surface

stained with antibodies for XCR1, CD11b, CD11c, Ly6G, MHC-II, CD3, CD19 (Biolegend

148220, 101261, 117318, 127627, 107606, 100233, and 115545 respectively) and Ly6C and

CD64 (BD 755198 and 569507 respectively) in Brillant stain buffer (BD 566349). Cells were

fixed and permeabilized with Foxp3 staining kit (Invitrogen 00-5523-00) before intracellular

staining for IL-12 p35 (Invitrogen MA5-23559) and IL-23 p19 (BD 565317). Data were collected

using a BD Symphony A3 flow cytometer (BD) and analyzed using FlowJo Software (BD).

Exogeneous IL-17a administration

Mice were intranasally administered 1 µg endotoxin-free recombinant murine IL-17a (R&D

systems, 7956-ML-100/CF) in 40 µL sterile PBS daily or 40 µL of vehicle control daily, or

intraperitoneally administered 2 µg endotoxin-free recombinant murine IL-17a in 100 µL sterile

PBS daily or 100 uL of vehicle control dailiy from days 11 to 20 post aerosol *M. tuberculosis*

challenge.

MSA-IL-17a was generated by fusing a (GGGGS)3 linker between the C-terminus of mature

murine serum albumin and N-terminus of mature murine IL-17a. MSA-IL-17a was purchased

from Genscript USA, Inc. in endotoxin-free pH 7.2 PBS. Mice were intranasally administered

5.54 µg MSA-IL-17a (equimolar to 1 µg recombinant murine IL-17a) in 40 µL sterile PBS daily or

40 µL vehicle control, or intraperitoneally administered 11.09 µg MSA-IL-17a (equimolar to 2 µg

recombinant murine IL-17a) in 100 µL PBS or 100 µL vehicle control daily between days 11 to

20 post aerosol *M. tuberculosis* challenge.

Statistical Analysis

All data was analyzed using FlowJo v10 (FlowJo, LCC) and/or Prism 10 (GraphPad Software).

All graphs were analyzed using Mann-Whitney U-test. Significant values are symbolized with

asterisks of $* < 0.05$, $** < 0.01$, and ns for non-significant.

Acknowledgements

We thank Russell Vance's lab for the kind gifts of $Sp140^{-/-}$, $Ifnar1^{-/-}$, and $Sp140^{-/-}Ifnar1^{-/-}$ mice. We thank Jeff Cox's lab for the kind gifts of ΔESX-1 (ΔeccC1), complemented ΔESX-1, PDIM-lacking (fadD28::tn), and complemented PDIM-lacking Erdman strains. We thank Helia Samani, Scott Espich, Vicky Wu, Malia Wilson, and Cielo Kwon for assistance in mouse colony maintenance. We thank members of the Stanley, Cox, and Vance laboratories for helpful discussions. SAS is a member of the Scientific Advisory Board for Xbiotix Therapeutics. This work was funded by NIH P01 AI063302 to S.A.S.

References

References

1. Geneva: World Health Organization. 2022. Global Tuberculosis Report 2022.

https://www.who.int/publications/i/item/9789240061729 (accessed October 5, 2024).

2. Fine, P. E. M. 1995. Variation in protection by BCG: implications of and for heterologous

immunity. *The Lancet* 346(8986):1339-1345. doi:10.1016/S0140-6736(95)92348-9.

3. Geneva: World Health Organization. 2018. Global Tuberculosis Report 2018.

https://apps.who.int/iris/bitstream/handle/10665/274453/9789241565646-

end.pdf?sequence=1&isAllowed=y (accessed October 5, 2024).

4. Kumar, P. 2021. A Perspective on the Success and Failure of BCG. *Front Immunol*

12:778028. doi:10.3389/fimmu.2021.778028.

5. Zhou, F., and D. Zhang. 2023. Recent advance in the development of tuberculosis

vaccines in clinical trials and virus-like particle-based vaccine candidates. *Front Immunol*

14:1238649. doi: 10.3389/fimmu.2023.1238649.

6. Flynn, J. L., J. Chan, K. J. Triebold, D. K. Dalton, T. A. Stewart, and B. R. Bloom. 1993.

An essential role for interferon gamma in resistance to *Mycobacterium tuberculosis*

Infection. *J Exp Med* 178(6):2249-2254. doi: 10.1084/jem.178.6.2249.

7. Condos, R., W. N. Rom, Y. M. Liu, and N. W. Schluger. 1998. Local Immune Response

Correlate with Presentation and Outcome in Tuberculosis. *Am J Respir Crit Care Med*

157:729-735. doi: 10.1164/ajrccm.157.3.9705044.

8. Casanova, J. and L. Abel. 2002. Genetic Dissection of Immunity to Mycobacteria: The

Human Model. *Annu Rev Immunol* 20:581-620. doi:

10.1146/annurev.immunol.20.081501.125851.

9. Ndiaye, B. P., D. Thienemann, M. Ota, B. S. Landry, M. Camara, S. Dièye, T. N. Dieye,

H. Esmail, R. Goliath, K. Huygen, V. January, I. Ndiaye, T. Oni, M. Raine, M. Romano, I.

Satti, S. Shutton, A. Thiam, K. A. Wilkinson, S. Mboup, R. J. Wilkinson, and H. McShane.

445 2015. Safety, immunogenicity, and efficacy of the candidate tuberculosis vaccine

446 MVA85A in healthy adults infected with HIV-1: a randomized, placebo-controlled, phase

447 2 trial. *Lancet Respir Med* 3(3):190-200. doi: 10.1016/S2213-2600(15)00037-5.

448 10. Ledesma, J. R., J. Ma, P. Zheng, J. M. Ross, T. Vos, and H. H. Kyu. 2021. Interferon-

449 gamma release assay levels and risk of progression to active tuberculosis: a systematic

450 review and dose-response meta-regression analysis. *BMC Infect Dis* 21(1):467. doi:

451 10.1186/s12879-021-06141-4.

452 11. Guglani, L. and S. A. Khader. 2010. Th17 cytokines in mucosal immunity and

453 inflammation. Curr Opin HIV AIDS 5(2):120-127. doi: 10.1097/COH.0b013r328335c2f6.

454 12. Gaffen, S. L. 2009. Structure and signalling in the IL-17 receptor family. *Nat Rev*

455 *Immunol* 9(8):556-567. doi: 10.1038/nri2586.

456 13. Lazarevic, V., X. Chen, J. Shim, E. Hwang, E. Jang, A. N. Bolm, M. Oukka, V. K.

457 Kuchroo, and L. H. Glimcher. 2011. Transcription factor T-bet represses T_H17

458 differentiation by preventing Runx1-mediated activation of the RORγT gene. *Nat*

459 *Immunol* 12(1):96-104. doi: 10.1038/ni.1969.

460 14. Pepper, M., J. L. Linehan, A. J. Pagán, T. Zell, T. Dileepan, P. P. Cleary, and M. K.

461 Jenkins. 2010. Different routes of bacterial infection induced long-lived T_H1 memory cells

462 and short-lived T_H17 cells. *Nat Immunol* 11:83-89. doi: 10.1038/ni.1826.

463 15. Huber, J. P., and J. D. Farrar. 2011. Regulation of effector and memory T-cell functions

464 by type I interferon. *Immunology* 132(4):466-474. doi: 10.1111/j.1365-

465 2567.2011.03412.x.

466 16. Yeh, W., I. L. McWilliams, and L. E. Harrington. 2014. IFNγ inhibits Th17 differentiation

467 and function via Tbet-dependent and Tbet-independent mechanisms. *J Neuroimmunol*

468 267(0):20-27. doi: 10.1016/j.jneuroim.2013.12.001.

17. Zhang, L., Yuan, S., Cheng, G., and B. Guo. 2011. Type I IFN Promotes IL-10

Production from T cells to Suppress Th17 Cells and Th17-Associated Autoimmune

Inflammation. *PLoS One* 6(12):e28432. doi: 10.1471/journal.pone.0028432.

18. Gideon, H. P., J. Phuah, A. J. Myers, B. D. Bryson, M. A. Rodgers, M. T. Coleman, P.

Maiello, T. Rutledge, S. Marino, S. M. Fortune, D. E. Kirschner, P. L. Lin, and J. L. Flynn.

2015. Variability in Tuberculosis Granuloma T Cell Responses Exists, but a Balance of

Pro- and Anti-inflammatory Cytokines Is Associated with Sterilization. *PLoS Pathog*

11(1):e1004603. doi: 10.1371/journal.ppat.1004603.

19. Perreau, M., V. Rozot, H. C. Welles, F. Belluti-Enders, S. Vigano, M. Maillard, G. Dorta,

J. Mazza-Stalder, P. Bart, T. Roger, T. Calandra, L. Nicod, and A. Harari. 2013. Lack of

Mycobacterium tuberculosis-specific interleukin-17A-producing CD4[+] T cells in active

disease. *Eur J Immunol* 43(3):939-948. doi: 10.1002/eji.201243090.

20. Ogongo, P., L. B. Tezera, A. Ardain, S. Nhamoyebonde, D. Ramsuran, A. Singh, A.

Ng'oepe, F. Karim, T. Naidoo, K. Khan, K. J. Dullabh, M. Fehlings, B. H. Lee, A. Nardin,

C. S. L. Arlehamn, A. Sette, S. M. Behar, A. J. C. Steyn, R. Madansein, H. N. Kløverpris,

P. T. Elkington, and A. Leslie. 2021. Tissue-resident-like CD4[+] T cells secreting IL-17

control *Mycobacterium tuberculosis* in the human lung. *J Clin Invest* 131(10):e142014.

doi: 10.1172/JCI142014.

21. Nathan, A., J. I. Beynor, Y. Baglaenko, S. Suliman, K. Ishigaki, S. Asgari, C. Huang, Y.

Luo, Z. Zhang, K. Lopez, C. S. L. Arlehamn, J. D. Ernst, J. Jimenez, R. I. Calderón, L.

Lecca, I. Van Rhijn, D. B. Moody, M. B. Murray, and S. Raychaudhuri. 2021.

Multimodally profiling memory T cells from a tuberculosis cohort identifies cell state

associations with demographics, environment and disease. *Nat Immunol* 22(6):781-793.

doi: 10.1038/s41590-021-00933-1.

22. Sun, M., J. M. Phan, N. S. Kieswetter, H. Huang, K. K. Q. Yu, M. T. Smith, Y. E. Liu, C.

Wang, S. Gupta, G. Obermoser, H. T. Maecker, A. Krishnan, S. Suresh, N. Gupta, M.

Rieck, P. Acs, M. Ghanizada, S. Chiou, P. Khatri, W. H. Boom, T. R. Hawn, C. M. Stein,

H. Mayanja-Kizza, M. M. Davis, and C. Seshadri. 2024. Specific CD4$^+$ T cell phenotypes

associate with bacterial control in people who 'resist' infection with *Mycobacterium*

tuberculosis. *Nat Immunol* 25(8):1411-1421. doi: 10.1038/s41590-024-01897-8.

23. Khader, S. A., G. K. Bell, J.E. Pearl, J. J. Fountain, J. Rangel-Moreno, G. E. Cilley, F.

Shen, S. M. Eaten, S. L. Gaffen, S. L. Swain, R. M. Locksley, L. Haynes, T. D. Randall,

and A. M. Cooper. 2007. IL-23 and IL-17 in the establishment of protection pulmonary

CD4$^+$ T cell responses after vaccination and during *Mycobacterium tuberculosis*

challenge. Nat Immunol 8(4):369-377. doi: 10.1038/ni1449.

24. Gopal, R., L. Monin, S. Slight, U. Uche, E. Blanchard, B. A. F. Junecko, R. Ramos-

Payan, C. L. Stallings, T. A. Reinhart, J. K. Kolls, D. Kaushal, U. Nagarajan, J. Rangel-

Moreno, and S. A. Khader. 2014. Unexpected Role for IL-17 in Protective Immunity

against Hypervirulent *Mycobacterium tuberculosis* HN878 Infection. *PLoS Pathog*

10(5):e1004099. doi: 10.1371/journal.ppat.1004099.

25. Segueni, N., E. Tritto, M. Bourigault, S. Rose, F. Erard, M. L. Bert, M. Jacobs, F. D.

Padova, D. P. Stiehl, P. Moulin, D. Brees, S. Chibout, B. Ryffel, M. Kammüller, and V. F.

Quesniaux. 2016. Controlled *Mycobacterium tuberculosis* infection in mice under

treatment with anti-IL-17A or IL-17F antibodies, in contrast to TNFα neutralization. *Sci*

Rep 6:36923. doi: 10.1038/srep36923.

26. Aguilo, N., S. Alvarez-Arguedas, S. Uranga, D. Marinova, M. Monzón, J. Badiola, and C.

Martin. 2016. Pulmonary but Not Subcutaneous Delivery of BCG Vaccine Confers

Protection to Tuberculosis-Susceptible Mice by an Interleukin 17-Dependent

Mechanism. *J Infect Dis* 213(5):831-839. doi: 10.1093/infdis/jiv503.

27. Jong, R. M., E. Van Dis, S. B. Berry, X. Nguyenla, A. Baltodano, G. Pastenkos, C. Xu, D.

Fox, N. Yosef, S. M. McWhirter, and S. A. Stanley. 2022. Mucosal Vaccination with Cyclic

Dinucleotide Adjuvants Induces Effective T Cell Homing and IL-17-Dependent Protection

against *Mycobacterium tuberculosis* Infection. *J Immunol* 208(2):407-419. doi:

10.4049/jimmunol.2100029.

28. Stanley, S. A., S. Raghavan, W. W. Hwang, and J. S. Cox. 2003. Acute infection and

macrophage subversion by *Mycobacterium tuberculosis* require a specialized secretion

system. *PNAS* 100(22):13001-13006. doi: 10.1073/pnas.2235593100.

29. Stanley, S. A., J. E. Johndrow, P. Manzanillo, and J. S. Cox. 2007. The Type I IFN

Response to Infection with *Mycobacterium tuberculosis* Requires ESX-1-Mediated

Secretion and Contributes to Pathogenesis. *J Immunol* 178(5):3143-3152. doi:

10.4049/jimmunol.178.5.3143.

30. Simeone, R., A. Bobard, J. Lippmann, W. Bitter, L. Majlessi, R. Brosch, and J. Enninga.

2012. Phagosomal Rupture by *Mycobacterium tuberculosis* Results in Toxicity and Host

Cell Death. *PLoS Pathog* 8(2):e1002507. doi: 10.1371/journal.ppat.1002507.

31. Romagnoli, A., M. P. Etna, E. Giacomini, M. Pardini, M. E. Remoli, M. Corazzari, L.

Falasca, D. Goletti, V. Gafa, R. Simeone, G. Delogu, M. Piacentini, R. Brosch, G. M.

Fimia, and E. M. Coccia. 2012. ESX-1 dependent impairment of autophagic flux by

Mycobacterium tuberculosis in human dendritic cells. *Autophagy* 8(9):1357-1370. doi:

10.4161/auto.20881.

32. Cox, J. S., B. Chen, M. McNeil, and W. R. Jacobs Jr. 1999. Complex lipid determines

tissue-specific replication of *Mycobacterium tuberculosis* in mice. *Nature* 402(6757):79-

83. doi: 10.1038/47042.

33. Barczak, A. K., R. Avraham, S. Singh, S. S. Luo, W. R. Zhang, M. Bray, A. E. Hinman, M.

Thompsan, R. M. Nietupski, A. Golas, P. Montgomery, M. Fitzgerald, R. S. Smith, D. W.

White, A. D. Tischler, A. E. Carpenter, and D. T. Hung. 2017. Systemic, multiparametric

analysis of *Mycobacterium tuberculosis* intracellular infection offers insight into

coordinated virulence. *PLoS Pathog* 13(5):e1006363. doi:

10.1371/journal.ppat.1006363.

34. Schlitzer, A., N. McGovern, P. Teo, T. Zelante, K. Atarashi, D. Low, A. W. S. Ho, P. See,

A. Shin, P. S. Wasan, G. Hoeffel, B. Malleret, A. Heiseke, S. Chew, L. Jardine, H. A.

Purvis, C. M. U. Hilkens, J. Tam, M. Poidinger, E. R. Stanley, A. B. Krug, L. Renia, B.

Sivasankar, L. G. Ng, M. Collin, P. Ricciardi-Castagnoli, K. Honda, M. Haniffa, and F.

Ginhoux. 2013. IRF4 Transcription Factor-Dependent CD11b[+] Dendritic Cells in Human

and Mouse Control Mucosal IL-17 Cytokine Responses. *Immunity* 38(5):970-983. doi:

10.1016/j.immuni.2013.04.011.\

35. Soldevilla, P., C. Vilaplana, and P. Cardona. 2022. Mouse Models for *Mycobacterium*

tuberculosis Pathogenesis: Show and Do Not Tell. *Pathogens* 12(1):49. doi:

10.3390/pathogens12010049.

36. Andersen, P., Å. B. Andersen, A. L. Sørensen, and S. Nagai. 1995. Recall of Long-Lived

Immunity to *Mycobacterium tuberculosis* Infection in Mice. *J Immunol* 154(7):3359-3372.

doi: 10.4049/jimmunol.154.7.3359.

37. Bekmurzayeva, A., M. Sypabekova, and D. Kanayeva. 2013. Tuberculosis diagnosis

using immunodominant, secreted antigens of *Mycobacterium tuberculosis*. *Tuberculosis*

93(4):381-388. doi: 10.1016/j.tube.2013.03.003.

38. Babaki, M. K. Z., S. Soleimanpour, and S. A. Rezaee. 2017. Antigen 85 complex as a

powerful *Mycobacterium tuberculosis* immunogene: Biology, immune-pathogenicity,

applications in diagnosis, and vaccine design. *Microb Pathog* 112:20-29. doi:

10.1016/j.micpath.2017.08.040.

39. Dileepan, T., J. L. Linehan, J. J. Moon, M. Pepper, M. K. Jenkins, and P. P. Cleary. 2011.

Robust antigen specific th17 T cell response to group A Streptococcus is dependent on

IL-6 and intranasal route of infection. *PLoS Pathog* 7(9):e1002252. doi:

10.1371/journal.ppat.1002252.

40. Tan, H., Sattler, M., and P. Zou. 2022. Albumin-binding as a universal strategy for half-life

extension. *Japan J Res* 3(1):1-5. doi: 10.3342512690-8077.1043.

41. Sullivan, B. M., O. Jobe, V. Lazarevic, K. Vasquez, R. Bronson, L. H. Glimcher, and I.

Kramnik. 2005. Increased Susceptibility of Mice Lacking T-bet to Infection with

Mycobacterium tuberculosis Correlates with Increased IL-10 and Decreased IFN-γ

Production. *J Immunol* 175(5):4593-4602. doi: 10.4049/jimmunol.175.7.4593.

42. Sallin, M. A., S. Sakai, K. D. Kauffman, H. A. Young, J. Zhu, and D. L. Barber. 2017. Th1

Differentiation Drives the Accumulation of Intravascular, Non-protective CD4 T Cells

during Tuberculosis. *Cell Rep* 18(13):3091-3104. doi: 10.1016/j.celrep.2017.03.007.

43. Hinman, A. E., C. Jani, S. C. Pringle, W. R. Zhang, N. Jain, A. J. Martinot, and A. K.

Barczak. 2021. *Mycobacterium tuberculosis* canonical virulence factors interfere with a

late component of the TLR2 response. *eLife* 10:e73984. doi: 10.7554/eLife.73984.

44. Augenstreich, J., A. Arbues, R. Simeone, E. Haanappel, A. Wegener, F. Sayes, F. Le

Chevalier, C. Chalut, W. Malaga, C. Guilhot, R. Brosch, and C. Astarie-Dequeker. 2017.

ESX-1 and phthiocerol dimycocerosates of *Mycobacterium tuberculosis* act in concert to

cause phagosomal rupture and host cell apoptosis. *Cell Microbiol* 19(7):e12726. doi:

10.1111/cmi.12726.

45. Crosskey, T. D., K. S. H. Beckham, and M. Wilmanns. 2020. The ATPases of the

mycobacterial type VII secretion system: Structural and mechanistic insights into

secretion. *Prog Biophys Mol Biol* 152:25-34. doi: 10.1016/j.pbiomolbio.2019.11.008.

46. Ji, D. X., L. H. Yamashiro, K. J. Chen, N. Mukaida, I. Kramnik, K. H. Darwin, and R. E.

Vance. 2019. Type I interferon-driven susceptibility to *Mycobacterium tuberculosis* is

mediated by IL-1Ra. *Nat Microbiol* 4(12):2128-2135. doi: 10.1038/s41564-019-0578-3.

47. Lai, R., T. Williams, T. Rakib, J. Lee, and S. M. Behar. 2024. Heterogeneity in lung

macrophage control of *Mycobacterium tuberculosis* is modulated by T cells. *Nat

Commun* 15(1):5710. doi: 10.1038/s41467-024-48515-7.

48. Amelio, P., D. Portevin, J. Hella, K. Reither, L. Kamwela, O. Lweno, A. Tumbo, L.

Geoffrey, K. Ohmiti, S. Ding, G. Pantaleo, C. Daubenberger, and M. Perreau. 2019. HIV

Infection Functionally Impairs *Mycobacterium tuberculosis*-Specific CD4 and CD8 T-Cell

Responses. *J Virol* 93(5):e01728-18. doi: 10.1128/jvi.01728-18.

49. Geremew, D., M. Melku, A. Endalamaw, B. Woldu, A. Fasil, M. Negash, H. W. Baynes,

H. Geremew, T. Teklu, T. Deressa, B. Tessema, and U. Sack. 2020. Tuberculosis and its

association with CD4+ T cell count among adult HIV positive patients in Ethiopian

settings: a systemic review and meta-analysis. *BMC Infect Dis* 20(1):325. doi:

10.1186/s12879-020-05040-4.

50. Van Dis, E., K. M. Sogi, C. S. Rae, K. E. Sivick, N. H. Surh, M. L. Leong, D. B. Kanne, K.

Metchette, J. J. Leong, J. R. Bruml, V. Chen, K. Heydari, N. Cadieux, T. Evans, S. M.

McWhirter, T. W. Dubensky Jr., D. A. Portnoy, and S. A. Stanley. 2018. STING-Activating

Adjuvants Elicit a Th17 Immune Response and Protect against *Mycobacterium*

tuberculosis Infection. *Cell Rep* 23(5):1435-1447. doi: 10.1016/j.celrep.2018.04.003.

51. Watson, R. O., S. L. Bell, D. A. MacDuff, J. M. Kimmey, E. J. Diner, J. Olivas, R. E.

Vance, C. L. Stallings, H. W. Virgin, and J. S. Cox. 2015. The cytosolic sensor cGAS

detects *Mycobacterium tuberculosis* DNA to induce type I interferons and activate

autophagy. *Cell Host Microbe* 17(6):811-819. doi: 10.1016/j.chom.2015.05.004.

52. Rojas, I. M. L., W. Mok, F. E. Pearson, Y. Minoda, T. J. Kenna, R. T. Barnard, and K. J.

Radford. 2017. Human Blood CD1c[+] Dendritic Cells Promote Th1 and Th17 Effector

Function in Memory CD4[+] T cells. Front Immunol 8:971. doi: 10.3389/fimmu.2017.00971.

53. Stadhouders, R., E. Lubberts, and R. W. Hendriks. 2018. A cellular and molecular view

of T helper 17 cells plasticity in autoimmunity. *J Autoimmun* 87:1-15. doi:

10.1016/j.jaut.2017.12.007.

624 **Figure S1.** Mice were infected with either the Erdman or Beijing strain of Mtb via the aerosol

625 route and evaluated at 21dpi for **(A)** CFU in the lungs or **(B-G)** IL-17a$^+$ CD4$^+$ T cells in the lungs.

626 **(B,C)** unstimulated, **(D,E)** restimulated with Antigen 85B peptide, **(F,G)** restimulated with ESAT-6

627 peptide. Representative experiment of 2.

Figure S2: Mtb Erdman PDIM mutants secrete ESAT-6 and ESX-1 mutants produce PDIM

(A) WT, ΔeccC1 (ESX), complemented ΔeccC1::eccC1 (ESX(C), *fadD28::tn* (PDIM), or

fadD28::tn + p*fadD28* (PDIM(C)) *M. tuberculosis* Erdman strain were cultured in Sauton's

complete media without Tween-80 for 5 days. Cell lysate and culture supernatant were

processed through SDS-PAGE and Western blot analysis. Anti-ESAT-6 antibody was used for

detecting ESAT-6. Anti-GroEL antibody was used as a loading control for the supernatant

fraction. Results shown are from 1 experiment representative of 2 independent experiments. **(B)**

WT, ΔeccC1 (ESX), complemented ΔeccC1::eccC1 (ESX(C), *fadD28::tn* (PDIM), or *fadD28::tn* +

p*fadD28* (PDIM(C)) *M. tuberculosis* Erdman strain were cultured in 7H9 media. Outer leaflet of

mycomembrane was removed via hexanes wash, concentrated and separated on TLC plate

with 98:2 petroleum ether:acetone mobile phase. TLC plate was stained with 0.2% Coomassie

blue in 20% methanol. Results shown are from 1 experiment representative of 2 independent

experiments.

649 **Figure S3. Attenuation of ESX-1 and PDIM mutants at 21 days post infection.** WT B6 mice

650 were aerosol infected with either WT, Δ*eccC1,* Δ*eccC1::eccC1*, *fadD28::tn*, or *fadD28::tn* +

651 p*fadD28 M. tuberculosis* Erdman strain. 21 days post infection, mice were sacrificed, and CFU

652 were enumerated from lung lysates. *, $p < 0.05$; **, $p < 0.01$ (unpaired nonparametric Mann-

653 Whitney U test).

Figure S4: Antigen 85b stim resembles ESAT-6 stimulation for IFN-γ and IL-17a staining

of CD4⁺ T cells. As in Figure 3, WT B6 mice were aerosol infected with either WT, Δ*eccC1,*

Δ*eccC1::eccC1*, *fadD28::tn*, or *fadD28::tn* + p*fadD28 M. tuberculosis* Erdman strain. 21 days

post infection, mice were sacrificed, lung single cell homogenates were stimulated with Antigen

85b peptide pool and measured for lung **(A)** IFN-γ⁺ CD4⁺ T cells and **(B)** IL-17a⁺ CD4⁺ T cells by

flow cytometry. Results in A and B are representative of 2 independent experiments. *, p < 0.05;

**, p < 0.01 (unpaired nonparametric Mann-Whitney U test

Figure S5: There is little to no IL-23 p19 expression in cDC1s and IL-12 p35 expression in

cDC2s during Mtb infection. As in Figure 5, WT B6 mice were aerosol infected with WT,

Δ*eccC1,* Δ*eccC1::eccC1*, *fadD28::tn*, or *fadD28::tn* + p*fadD28 M. tuberculosis* Erdman strain.

21 days post infection, mice were sacrificed, mediastinal lymph nodes were extracted, and

processed for ICS. Analysis of **(A)** IL-23 p19$^+$ cDC1s and **(B)** IL-12 p35$^+$ cDC2s by flow

cytometry. Results in A and B are representative of 2 independent experiments. (unpaired

nonparametric Mann-Whitney U test).