# Cell cycle checkpoint activity in the malaria parasite

# *Plasmodium falciparum*

Monique K. Johnson[1], Juliana Naldoni[1], William H. Lewis[2], Ross F. Waller[2], Catherine J. Merrick[1*]

[1] Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge, CB2 1QP, UK.

[2] Department of Biochemistry, University of Cambridge, Tennis Court Road, Cambridge, CB2 1QW, UK.

* Address correspondence to:

Department of Pathology, Cambridge University, Tennis Court Road, Cambridge, CB2 1QP, UK

Tel: (+44) 1223 333330

Email: cjm48@cam.ac.uk

# Abstract

*Plasmodium spp.* have different modes of cell division from most eukaryotes. Little is known about how these are controlled and cell-cycle checkpoints are particularly poorly characterised. However, parasites can arrest their cell cycle when treated with the frontline antimalarial drug artemisinin, and artemisinin-resistant parasites can modulate their cell-cycle progression, so it is very important to understand these aspects of *Plasmodium* biology. Here, we show that *P. falciparum* displays hallmarks of an intra-S-phase checkpoint when exposed to DNA damage, including acute reduction of DNA replication and phosphorylation of a putative damage-marker histone. Compounds that inhibit human checkpoint kinases can inhibit this arrest of DNA replication, and synergise with DNA damage in parasite killing. This suggests the existence of checkpoint kinase activity in *P. falciparum*, yet these kinases have no clear orthologues in *Plasmodium* genomes. We hypothesise that

23 phosphatidylinositol 3-kinase – an essential lipid kinase – may moonlight in this role: it is the closest

24 homologue to a checkpoint kinase and is reportedly up-regulated in artemisinin-resistant parasites.

25 Finally, we show that the cryptic checkpoint-kinase activity may also regulate the ring-stage survival

26 phenotype after artemisinin damage, which resembles a G1/S checkpoint.  Hence we suggest that

27 checkpoint kinase inhibitors are candidates for synergy with artemisinin.

## 28 Author summary

Malaria parasites cause illness by infecting human red blood cells, wherein they replicate to produce

many new parasites within ~1-3 days.  This is unusual because most cells (such as human cells)

replicate simply by copying their genome and splitting in half – which is called binary fission – but

malaria parasites make ~20 genome copies and then partition them simultaneously into 20 new cells

– which is called schizogony.  Here we studied how these parasites control schizogony.  In particular,

do they have 'checkpoints': the ability to pause the cell cycle for repair if the genome is damaged?

We found that DNA damage during active replication did indeed cause hallmarks of a checkpoint.

Key proteins that enforce this in other cells are surprisingly absent in malaria parasites, although

DNA repair pathways and other regulatory pathways remain present.  Furthermore, this checkpoint

activity may be involved in the response to the antimalarial drug artemisinin: a response that

parasites make by pausing their cycle before active replication begins.  This could be important

because it implies that inhibiting the checkpoint could exacerbate parasite killing by artemisinin-

based drugs, similar to the way that some cancer therapies work by damaging DNA and also

preventing the cancer cells from repairing it.

## 43 Keywords

44 Malaria, *Plasmodium,* cell cycle, checkpoint, PI3K

## 45 Introduction

*Plasmodium falciparum* is the most important cause of human malaria, giving rise to over half a

million deaths and hundreds of millions of clinical cases per year [1]. Besides being a major human

pathogen, it is also a highly unusual protozoan organism, distantly related to humans and other

model organisms, with distinctive features of cell and molecular biology. One of these is a non-

canonical cell cycle. Instead of dividing by binary fission, as in most model organisms, *Plasmodium*

parasites divide primarily by schizogony. After invading a host erythrocyte, they execute multiple,

asynchronous rounds of genome replication and nuclear division, generating a syncytium with

multiple nuclei within the same cytoplasm. This is followed by a relatively synchronous mass

cytokinesis event, partitioning as many as 20-30 nuclei and organelles into individual merozoites.

The complex events of schizogony are now being dissected in considerable molecular and

cellular detail [2-5] but fundamental questions remain about how the process is regulated.

Schizogony is not easily mapped onto the well-characterised cell cycle phases of G1, S, G2 and M.

After invading a new host cell, the parasite undergoes a pre-replicative growth phase broadly

analogous to G1. The haploid genome is then repeatedly replicated in a continuous S-phase without

once-and-only-once regulation, generating many new haploid nuclei. Different nuclei cease to

replicate at different times, and divide at different times, so there is no well-defined G2 and no

single M-phase [3].

*Plasmodium* replication is not controlled by conventional waves of cyclin and cyclin-

dependent kinase activity [6]. Nor is it apparently governed by conventional cell cycle checkpoints,

which usually operate at the G1/S transition, within S-phase, and at G2/M. Respectively, these

checkpoints delay the start of S-phase, arrest replication if DNA damage or replication stress is

detected, and prevent mitosis if a genome is incompletely replicated [7]. The *Plasmodium* genus,

however, encodes no clear homologues of the checkpoint kinases conserved in most eukaryotes

(animals, fungi and plants): Mec1 and Rad53 in *S. cerevisiae* and their homologues ATM/ATR and

Chk1/Chk2 in mammalian cells [6]. (By contrast, some parasites in the apicomplexan phylum,

including *Toxoplasma*, do retain homologues of the 'master regulators' Mec1/ATM/ATR, which are

phosphoinositide-3-kinase-like kinases (PIKKs), and checkpoints have been better characterised in

*Toxoplasma* [8-10]).

Nevertheless, eukaryotic cells generally do pause their cell cycles to ensure effective DNA

repair and faithful DNA replication – not only in metazoans, which must guard against cancer, but

also in protozoans. *Plasmodium* encodes most of the known DNA repair pathways [6, 11] and

generates relatively low rates of SNPs, indels and translocations during normal replication [12], so it

seems unlikely that it is entirely without checkpoints.

There is some evidence for a G1/S-like checkpoint: pre-replicative rings can undergo

'temporary growth arrest (TGA)' [13] in response to nutritional [14], temperature [15] or

antimalarial-drug [16, 17] stresses. TGA is particularly important under stress caused by the

antimalarial drug artemisinin, because it can contribute to treatment failure. However, there is no

molecular evidence as-yet that conventional checkpoint effectors are involved. A G1/S checkpoint

usually involves the cyclin/CDK complexes that drive S-phase entry, influenced positively by growth

factors or negatively by damage responsive kinases like PIKKs. TGA, by contrast, is defined only as a

'non-specific stress-response survival mechanism' [13]: induced by multiple stresses, possibly via

multiple molecular pathways.

An intra-S-phase checkpoint, meanwhile, should arrest S-phase specifically after DNA

damage or replication stress. After DNA damage, the transcriptional cascade that drives the

*Plasmodium* cell cycle can slow down, with DNA repair proteins and chromatin modifications being

upregulated [18], but the signals and transducers for this potential checkpoint are unknown.

(*Plasmodium* generally lacks validated markers for DNA damage and for direct responses to it.)

Indeed, the multi-nucleate, asynchronous nature of schizogony raises questions about how an intra-

S-phase checkpoint should or could be enforced. Replication itself is apparently regulated per-

nucleus, probably via individual accumulations of PCNA, together with non-canonical, chromatin-

bound CDK-related kinases like CRK4 [19]. Nevertheless, diffusible checkpoint kinases, potentially

affecting the whole cell, may not be incompatible with schizogony – at least in response to

exogenous DNA damage.

Finally, a G2/M checkpoint may be entirely absent because some *P. falciparum* mutants

perform grossly aberrant genome segregation but still undergo cytokinesis [20]. In fact, even in non-

mutants, certain phases of the lifecycle produce a high rate of zoids – particularly the rapid process

of gametogenesis inside the mosquito, when male cells produce and segregate eight new genomes

into mature, motile gametes in less than 15 minutes. In gametogenesis, cytokinesis proceeds even if

genome replication fails entirely [21, 22].

Here, we set out to investigate checkpoint responses to DNA damage in *Plasmodium,* using

newly developed molecular and cellular tools. We show that the hallmarks of an S-phase checkpoint

do occur in *P. falciparum*, despite the apparent absence of canonical PIKKs.

## Results

## Evidence for DNA-damage-responsive checkpoint activity in *P.*

## *falciparum*

We used two complementary tools to investigate the response to DNA damage during S-phase in *P.*

*falciparum*, focussing at the levels of DNA and protein. Firstly, we labelled nascent DNA in

replicating parasites with a quantifiable modified nucleotide. This tested whether DNA replication

could be acutely arrested after damage: a hallmark of the intra-S-phase checkpoint [23, 24].

Secondly, we measured phosphorylation of histone H2A, which was recently identified as being

analogous to histone H2AX phosphorylation in mammalian cells [25]. Canonically, H2AX is

phosphorylated at chromatin sites of DNA damage [26, 27]. H2AX is absent in *Plasmodium*, but the

non-variant histone H2A can nevertheless be phosphorylated [25].  Importantly, this indicates an *in*

*trans*, kinase-driven, genuinely 'checkpoint-like' response.

To induce DNA damage, we used the well-characterised alkylating agent methyl methane

sulphonate (MMS), which alkylates DNA bases, particularly guanine and adenine [28].  In mammalian

cells, it induces an intra-S-phase checkpoint [23].  In *P. falciparum* parasites, it leads to DNA

breakage, as measured by comet assays [29] and concomitantly delays the cell cycle [18], but its

effect as a checkpoint inducer has not been explicitly addressed.  In the *P. falciparum* 3D7 strain,

MMS had a 48h $IC_{50}$ of ~96µM (Fig S1A).  Having established this, we performed highly acute assays

over just 1-2h, using 10x or 20x $IC_{50}$, to minimise non-specific or secondary effects.  We confirmed

that parasites could substantially recover from this level of MMS exposure within one cell cycle,

although their maturation was delayed, as reported previously [18], and there was also a reduction

in merozoite numbers in mature schizonts (Fig S2).

DNA replication was acutely reduced in the 30 minutes following a 30-minute exposure to

MMS at 10x or 20x $IC_{50}$.  Some reduction occurred throughout S-phase (~28-36 hours post invasion,

hpi) but it was most pronounced when DNA was damaged in early trophozoites (~32hpi) (Fig 1A).

Within a similar timeframe, histone H2A phosphorylation was moderately elevated (Fig 1B).  We

recently reported that several DNA damaging agents, including MMS, could cause H2A

phosphorylation [30], but that the upregulation was transient and modest – usually less than 2-fold

over background levels – making it difficult to measure consistently.  This was equally true when

MMS was used to induce H2A phosphorylation here (Fig 1B).  (Previously published work measured

H2A phosphorylation in pre-replicative rings exposed to ionising radiation [25], whereas we

measured it in actively replicating trophozoites.)  Therefore, we chose the replication-reduction

assay as a more robust readout for checkpoint activity in subsequent experiments.

141                    -----------------------------------------

**Fig 1:** Evidence for DNA-damage-responsive checkpoint activity in *P. falciparum*

A.  BrdU incorporation into populations of trophozoites at three stages, after treatment with

MMS, as measured by ELISA. Parasites were treated with MMS for 1 h and BrdU was added

for the latter 30 mins.  Means of two biological replicates, conducted in technical triplicate,

are shown; error bars show range. MMS treatments: 10x $IC_{50}$, 960 µM; 20x $IC_{50}$, 1.92 mM.

For each parasite stage, statistical testing was via ordinary one-way ANOVA and posthoc

Tukey's multiple comparisons test. *, P-value <0.05; comparisons not shown, ns. Pictures

show examples of trophozoites at each timepoint. Scale bar, 2µm.

B.  Representative western blot of phosphorylated histone in trophozoite parasite lysate

treated with MMS (2 h, 10x $IC_{50}$). Phosphorylated histone signal was quantified versus

control histone H4: relative quantification is shown below the blot.  Graph shows

quantifications from three biological replicate blots, relative to either H4 or Hsp70 as a

control protein. Unpaired two-tailed t-test: P-value = 0.0354.

155                       -----------------------------------------

## Possible mediators of checkpoint activity in *P. falciparum*

When the intra-S-phase checkpoint is triggered in other eukaryotes, both histone phosphorylation

and arrest of DNA replication are transduced through PIKKs [24, 26].  Since Figure 1 shows that these

phenotypes also appear in *P. falciparum,* a transducing kinase activity should be present.  However,

orthologues of ATM/ATR are not obvious by similarity searches in the *P. falciparum* genome [6].  To

test for the presence or loss of such orthologues more thoroughly, we performed an extensive

phylogenetic analysis of PIKK homologues throughout apicomplexans and other major eukaryotic

lineages (Fig 2).  In this analysis, PI3K and PI4K kinases formed an outgroup relative to all ATM/ATR-

like proteins.  ATR homologues were found in most apicomplexans, and some – like *T. gondii* –

clearly retained homologues of both ATM and ATR.  Other apicomplexans, including

*Cryptosporidium*, *Babesia* and *Theileria*, had apparently independently lost the ATM homologue,

indicating a propensity for loss of PIKKs in Apicomplexa.  *Plasmodium* spp. were unique in having lost

both ATM and ATR, retaining no apparent PIKKs at all.  The closest PIKK homologues in any

*Plasmodium* genome were the lipid kinases PI3K and PI4K (Fig 2, Fig S3), which are also present

throughout most apicomplexans [31].  (*Theileria* was seemingly unique in having lost its PI3K

homologue, possibly due to a lifecycle conducted free in the cytoplasm of leukocytes, which may

supply this activity.)  Notably, the closest *Plasmodium* homologue, PI3K, has been reported to be

overexpressed in artemisinin resistant parasites [32], which can alter their cell cycle dynamics.

Accordingly, we hypothesised that PI3K might act as a checkpoint kinase in *Plasmodium.*

----------------------------------------

**Fig 2:** *Plasmodium encodes no homologues of PIKK checkpoint kinases*

A phylogeny inferred for amino acid protein sequences of ATR, ATM, and TOR homologues sampled

from broad eukaryote groups. Apicomplexa homologues of PI3K and PI4K (the most closely related

proteins to ATR and ATM present in *Plasmodium*) and characterised homologues of PI3K and PI4K

from *Saccharomyces cerevisiae* and *Homo sapiens* were included as an outgroup. Support values,

displayed as percentages, were generated from 1000 ultrafast bootstrap replicates [33] and only

support values ≥ 90, indicating strongly supported clades, are shown. The scale bar represents the

number of amino acid substitutions per site. A fully-expanded version of this tree with no collapsed

clades and displaying sequence accessions is provided in the source data.

*Pf*PI3K is the only phosphoinositide 3-kinase of the six phosphoinositide kinases in

*Plasmodium* [31]: it is a well-validated, essential lipid kinase [34] with roles in nutrient uptake and

vesicle trafficking [35, 36].  We hypothesised that it might also 'moonlight' as a protein kinase, which

would not be unprecedented, as other PI3Ks can to act on proteins as well as lipids.  For example,

humans encode a large family of PI3Ks, and the type-1 PI3Ks can phosphorylate protein substrates

such as cytokine receptors, as well as auto-phosphorylating themselves [37, 38].  The sole

*Plasmodium* enzyme is a type-3 PI3K, and thus far these are proven only to phosphorylate lipids [34].

193   If our hypothesis was correct, then *Pf*PI3K should have some activity within the nucleus,

194 phosphorylating substrates such as histone H2A.  To track the protein *in cellulo,* we raised peptide

195 antibodies (modelling for peptide selection is shown in Fig S4A), which detected a protein of the

196 correct 256kDa size (Fig 3A).  In immunofluorescence, *Pf*PI3K appeared dispersed and cytoplasmic

197 (Fig 3B, Fig S4B), consistent with a previous publication that suggested a location in transport

198 vesicles [36].  We did not detect any substantial re-location to the nucleus upon DNA damage (Fig

199 3C).  However, we recently showed that the ATM and ATR homologues in *Toxoplasma* are likewise

200 dispersed throughout the cell and do not detectably relocate to the nucleus after DNA damage, but

201 they are nevertheless able to mediate checkpoint phenotypes such as histone phosphorylation [39].

202        ----------------------------------------

203 **Fig 3:** *Pf*PI3K does not visibly relocate to the nucleus upon DNA damage

204  A. Western blots showing detection of *Pf*PI3K by two antisera raised to *Pf*PI3K peptides.

205    Replicate lanes from the same blot of protein lysate from late-stage *P. falciparum* were

206    exposed to either pre-immune or immune serum. Histone H4 was used as a loading control.

207  B. Immunofluorescence assay (IFA) showing location of *Pf*PI3K (detected by antiserum 1) in *P.*

208    *falciparum.* 2-3 representative cells are shown per stage. Scale bar 1 μm.

209  C. IFA showing *Pf*PI3K in *P. falciparum* after DNA damage: 1 h treatment with 5x or 10x $IC_{50}$ of

210    DHA or MMS on late-stage parasites.  One representative cell per condition is shown. Scale

211    bar 1μm.

212        ----------------------------------------

213   Further molecular-genetic experiments were precluded because the *Pf*PI3K enzyme is

214 essential (probably due to its unique lipid kinase activity).  Several attempts were made to 3' tag the

215 gene for a partial, non-lethal conditional knockdown instead, but these failed, probably because the

216 enzyme's active site is encoded at the extreme 3' end and may become mis-folded after 3' tagging.

217 Similar attempts by other groups to manipulate this gene were either unsuccessful or gave no

phenotype [40, 41].  The large size of the *PfPI3K* gene (6.6kb) precluded overexpression, dominant-

negative overexpression or recombinant protein production for enzyme-activity assays.  Attempts

were also made to immunoprecipitate the native protein with its potential substrates, but only a

single *Pf*PI3K peptide was detected across several immunoprecipitation attempts – probably because

the kinase is lowly expressed, very large and unstable *in vitro*, or lacking suitable tryptic digest sites

(notably, it was likewise undetected, despite its large size, in two dedicated proteomic studies that

examined the nuclear versus cytoplasmic proteomes in *P. falciparum* [42, 43]).  Thus, the putative

PIKK-like activity of *Pf*PI3K remained neither evidenced nor disproved.  Future work using state-of-

the-art proteomic methods like hyperLOPIT for protein localisation (recently achieved in *T. gondii*

[44]) or proximity-labelling of protein partners via BioID [45] could shed light on a putative intra-

nuclear role for *Pf*PI3K.

# The mediator of DNA-damage checkpoint responses in *Plasmodium*

# is sensitive to PIKK inhibitors

Despite the lack of conclusive evidence for *Pf*PI3K as a checkpoint kinase, a divergent PIKK-like

kinase may nevertheless exist in *P. falciparum*, and hence we attempted to inhibit it with PIKK

inhibitors.  We chose two compounds that had validated specificity for human ATM and ATR

respectively (but not for human PI3Ks), KU-55933 and VE-821 [46, 47].  If an ATM-like and/or ATR-

like activity existed in *P. falciparum*, either in *Pf*PI3K or in another protein, these compounds might

specifically inhibit it.  We first determined their 48h $IC_{50}$ values on *P. falciparum* (3D7 strain): KU-

55933 had an $IC_{50}$ in the micromolar range and VE-821, in the nanomolar range (Fig 4A, Fig S1B).

238                                 ----------------------------------------

239   **Fig 4:** Inhibitors of human PIKKs synergise with the DNA-damaging agent MMS in *P. falciparum*

240       A.   Inhibition of parasite growth by VE-821 and KU-55933 (molecular structures of each

241            inhibitor are shown).  Growth curves from five independent malaria SYBR green I-based

fluorescence (MSF) assays: percentage parasite growth is plotted as a function of drug

concentration, error bars are standard errors of the means. Mean $IC_{50}$s from all experiments

are shown (individual $IC_{50}$ values from replicate experiments are in Fig S1B).

B.  Isobolograms plotted as fractional inhibitory concentrations (FIC) of combinations of MMS

and VE-821 or KU-55933, as determined by MSF assay. FIC calculations were completed for

each combination by dividing the measured "apparent" $IC_{50}$ values for individual drugs in the

different combinations by $IC_{50}$ values obtained when the drugs were used alone. Each curve

represents a different biological replicate experiment.  The mean FIC ($\Sigma FIC(FIC_A+FIC_B)$) is

shown, with range.  The type of relationship was defined as moderately synergistic based on

well-established criteria: $\Sigma FIC$ <0.5, substantial synergism; $\Sigma FIC$ <1, moderate synergism;

$\Sigma FIC$ ≥1 and <2, additive interaction; $\Sigma FIC$ ≥2 and <4, slight antagonism and $\Sigma FIC$ >4, marked

antagonism [48].

C.  BrdU incorporation into 30-32 hpi trophozoites after treatment with MMS, VE-821, KU-

55933, geneticin and combinations of these, measured by ELISA. Parasites were treated with

10x $IC_{50}$ of drug for 1 h and BrdU was added for the latter 30 mins. Six biological replicates

are plotted, each completed in technical triplicate. Mean and standard deviation error bars

are plotted, values normalised to control (set to 100%) in each replicate. Concentrations of

drugs used: MMS 960 $\mu$M, VE-821 1.62 $\mu$M, KU-55933 400 $\mu$M, geneticin 1500 $\mu$g/ml.

Ordinary one-way ANOVA and posthoc Tukey's multiple comparisons test of control vs all

and MMS vs all was completed; P-values: *, <0.05; **, <0.01; ***, <0.001; ****, <0.0001;

comparisons not shown, ns.

----------------------------------------

Both inhibitors clearly synergised with MMS in a 48h isobologram analysis, particularly

strongly in the case of VE-821, suggesting activity within the same pathway (Fig 4B).  Control

compounds that kill parasites but are not known to cause DNA damage – the antibiotics blasticidin

and geneticin – did not synergise with MMS (Fig S5).  Furthermore, when parasites were exposed to

high-dose MMS (10x IC50, as in Figure 1) simultaneously with KU-55933 or VE-821, the inhibitors

ablated the acute reduction in DNA replication caused by MMS, whereas co-exposure to the control

antibiotic geneticin did not (Fig 4C). In this short 1h assay, we used the PIKK inhibitors at only their

48h IC50 levels, to minimise any negative effects from the inhibitors alone. Indeed, within the

timeframe of the assay the inhibitors alone did not significantly affect DNA replication (Fig 4C).

Overall, these data strongly supported the existence of a cryptic PIKK activity, responsible for a

genuine intra-S-phase checkpoint via *in trans* kinase activity.

Having established this, we sought to link the inhibitors to their putative target of *Pf*PI3K, on

the basis that parasites with more or less *Pf*PI3K would be expected to have, respectively, lower or

higher inhibitor sensitivity. Although no successful inducible knockdown of *Pf*PI3K has yet been

created (as noted above), the first inducible knockout was very recently published [49]. We

obtained this and tested knockout parasites for any change in sensitivity to PIKK inhibitors.

Unfortunately, knockout parasites did not prove amenable to standard 48h (or 72h) drug-sensitivity

assays (Fig S6). This was because they displayed a delayed-death phenotype, well described in [49],

that rendered them relatively healthy through cycle-2 after induction of the knockout, but dead

before cycle-3. Accordingly, they were not differently sensitive to VE-821 or KU-55933 in MSF assays

conducted over 48h in cycles 1-2, but all parasites died before a longer 72h MSF assay could be read.

The converse approach of overexpressing PI3K for a reduction in inhibitor sensitivity was also

attempted. Due to the prohibitively large size of *Pf*PI3K, we used a PbAC artificial chromosome to

express an extra copy of the *P. berghei* PI3K homologue in *P. knowlesi*: a system recently

published as a novel chemogenomic screening platform [50]. Unfortunately, *Pb*PI3K transgene

overexpression proved in this instance to be weak and resultant changes in drug sensitivity were

marginal and not statistically-significant (similar to the results shown in figure S6).

## The antimalarial drug dihydroartemisinin induces DNA replication arrest in *P. falciparum*

MMS is a useful, well-characterised DNA-damaging agent, but it is not an antimalarial drug. We were particularly interested in the potential DNA-damaging activity of antimalarial drugs, because if parasites do make checkpoint responses to such drugs, then their antimalarial activity might be potentiated by inhibiting checkpoint kinases. Artemisinin is such a drug: it has non-specific alkylating activity that may affect both proteins and DNA [51, 52]. There are similarities between the parasite's response to dihydroartemisinin (DHA, the active metabolite of artemisinin) and MMS in terms of DNA breakage [51, 53], delay in the parasite's transcriptional cascade, and upregulation of DNA repair genes [18]. Indeed, we found that DHA induced H2A phosphorylation, at all stages of the cycle but particularly in pre-replicative ring stages (~3-fold upregulation after 4h exposure to $IC_{50}$ DHA, (Fig 5A)). This confirmed that DHA does indeed cause damage that activates a putative histone kinase, and furthermore that it can cause this damage prior to S-phase.

Using the replication-reduction assay, we saw that in replicating parasites, DHA caused an acute reduction in nascent DNA replication, similar to MMS (Fig 5B). However, unlike MMS, this was not ablated by the PIKK inhibitors KU-55933 and VE-821. Consistent with this difference, the inhibitors did not synergise with DHA in isobologram analysis (Fig 5C). This suggested that although both DHA and MMS may damage DNA, they differ in either the type of damage or the cellular response thereto in terms of inducing an intra-S-phase checkpoint.

----------------------------------------

**Fig 5:** The antimalarial drug dihydroartemisinin induces a DNA replication arrest in *P. falciparum*

  A. Western blot of phosphorylated histones in ring, trophozoite and schizont-stage parasites treated with DHA (4h, $IC_{50}$ (3nM)). Phosphorylated histone signal was quantified versus control histone H4: relative quantification is shown below the blot.

315        B.  BrdU incorporation into 30-32 hpi trophozoites after treatment with DHA, VE-821, KU-55933

316           and combination treatments. Parasites were treated with 10X $IC_{50}$ of drug for 1 h and BrdU

317           was added for the latter 30 mins. Five biological replicates were plotted, each completed in

318           technical triplicate. Mean and standard deviation error bars are plotted; values were

319           normalised to control (set to 100%) in each replicate. Concentrations of drugs used: DHA 45

320           nM, VE-821 1.62 $\mu$M, KU-55933 400 $\mu$M. Statistical testing by ordinary one-way ANOVA and

321           posthoc Tukey's multiple comparisons test of control vs all and DHA vs all; P-values: *, <0.05;

322           comparisons not shown, ns.

323        C.  Isobolograms plotted as Figure 4B. Each curve is a biological replicate and the mean FIC

324           ($\Sigma$FIC($FIC_A$+$FIC_B$)) is shown, with range. The relationship between DHA and either VE-821 or

325           KU55933 was defined as additive.

326                                ----------------------------------------

## Ring-stage survival after artemisinin damage is affected by PIKK inhibitors

Cell-cycle checkpoint activity in *P. falciparum* may not be limited to an intra-S-phase checkpoint:

most eukaryotic cells also operate a G1/S checkpoint, delaying S-phase entry if cells are damaged,

starved or stressed. This may be analogous to the 'TGA' arrest or 'dormancy' that occurs in ring-

stage parasites after various stressors, including artemisinin treatment [13]. We therefore set out to

establish whether ring-stage dormancy induced by DHA actually represents a classical, PIKK-

mediated G1/S checkpoint, responsive to the DNA damage that can be exerted upon ring stages by

DHA (as shown in figure 5A).

We conducted the ring-stage survival assay (RSA) on parasites treated with DHA in the

presence or absence of PIKK inhibitors. In this assay, pre-replicative early-ring-stage parasites are

damaged with a pulse of DHA: most of them die, but some successfully arrest and thence survive.

The more parasites are able to do this, the higher the subsequent recovery of viable parasites.   We

reasoned that if successful arrest is due to PIKK-mediated checkpoint activity, then PIKK inhibitors

should reduce ring-stage survival.  These experiments were conducted in the 3D7 strain and also in

the Cambodian strain MRA-1252.  MRA-1252 was derived from a naturally-arising DHA-resistant

strain, which was genetically reverted back to a sensitive phenotype by reverting the resistance

mutation in the gene encoding Kelch13 [54].  Its genetic background is otherwise that of a S.E. Asian

DHA-resistant line.  Both 3D7 and MRA-1252 are highly sensitive to artemisinin, so we used the

'extended recovery ring-stage survival assay' (eRRSA), which detects recovered parasites very

sensitively after 72h via quantitative PCR [55].

Figure 6A shows that PIKK inhibitors, particularly KU-55933, did indeed exacerbate the killing

of ring-stage parasites by DHA.  Differences were only subtle and parallel assays using the more

common but less sensitive flow cytometric assessment of live/dead cells to read the RSA likewise

showed only marginal differences (Fig S7).  However, in both 3D7 and MRA-1252, triplicate eRRSAs

showed that survival trended downwards when DHA was combined with either inhibitor.  Neither

condition (DHA + VE-821 or DHA + KU-55933) reached statistical significance versus survival after

DHA alone, but the p-values versus control were $< 0.0001$ rather than 0.001.  As above, we

minimised any negative effects of the inhibitors themselves by using them at a low level – only half

of the 48h $IC_{50}$.  We confirmed that this exposure alone for 6h did not affect parasite survival after

72h (Fig 6B).

358                                        -----------------------------------------

**Fig 6**: Ring-stage survival is affected by PIKK inhibitors

Data in A-D are from eRRSA assays on the strains 3D7, MRA-1252 and MRA-1240. Samples were

treated with DMSO (the control condition), VE-821, KU-55933, DHA and combinations of these. Drug

concentrations: VE-821, 81 nM (0.5X $IC_{50}$); KU-55933, 20 μM (0.5X $IC_{50}$); DMSO, 0.02%; DHA, 700 nM.

Values plotted are the means of -log(foldchange = 2^(DHA-DMSO)) for each treatment. Midline

represents the mean and boxes show min to max from three biological replicates, each conducted in

technical triplicate.  Statistical testing by ordinary one-way ANOVA and Dunnett's multiple

comparisons test, comparing all means to the mean value from DMSO- or DHA-treated samples.  P-

values: *, <0.05; **, <0.01; ***, <0.001; ****, <0.0001. Differences between DHA-treated samples

and treatments other than the control did not reach statistical significance.

A.  Ring-stage DHA-sensitive parasite strains, 3D7 or MRA-1252, were treated with DMSO (control

condition), DHA and inhibitors VE-821 or KU-55933 in combination with DHA for a 6 h pulse, then

left to recover for 72 h.

B.  Control experiments without DHA: ring-stage 3D7 or MRA-1252 parasites were treated with

DMSO or inhibitors VE-821 or KU-55933 for a 6 h pulse, then left to recover for 72 h.

C.  Ring-stages of the DHA-resistant parasite strain MRA-1240 were treated with DMSO (control

condition), DHA and inhibitors VE-821 or KU-55933 in combination with DHA for a 6 h pulse, then

left to recover for 72 h.

D.  Control experiment without DHA: ring-stage MRA-1240 parasites were treated with DMSO or

inhibitors VE-821 or KU-55933 for a 6 h pulse, then left to recover for 72 h.

E.  Inhibition of growth over a full 48h growth cycle by MMS, VE-821 and KU55933 was measured in

the strains 3D7, MRA-1252 and MRA-1240 via MSF assay.  $IC_{50}$s were calculated from at least

three independent experiments and all values are plotted, with the midline showing the mean

and error bars showing standard deviation. MSF growth curves are in Fig S1C.

383                                    ----------------------------------------

These RSAs did not show conclusively that 'TGA' after artemisinin damage represents a true

G1/S checkpoint.  However, they did suggest that such a checkpoint, mediated by a cryptic PIKK, may

contribute to ring-stage survival.  Hence, the cryptic PIKK must be active throughout the cell cycle,

not only in actively replicating trophozoites.

When the same assay was conducted on DHA-resistant parasites (the parent strain of MRA-

1252, called MRA-1240) the result differed. As expected, DHA survival was much better in MRA-

1240, and PIKK inhibitors had no negative effect on this survival (Fig 6C). If anything, there was a

trend towards better growth and better DHA survival when MRA-1240 was exposed to PIKK

inhibitors (Fig 6D). Hence, when a Kelch13 mutation was present, conferring DHA resistance, any

contribution of checkpoint activity to parasite survival was apparently ablated.

## 394 Artemisinin-resistant parasites are cross-resistant to MMS and VE-

## 395 821, but not to KU-55933

We hypothesised that artemisinin-resistant (Kelch13 mutant) parasites might not require 'G1/S'

checkpoint activity to survive DHA damage because they are more efficient at DNA repair. Several

mutations in DNA repair genes have been identified in resistant strains [52], although the pathway(s)

leading to improved DNA repair remain to be fully characterised. We checked whether the DHA-

resistant line MRA-1240 was cross-resistant to MMS, and its $IC_{50}$ was indeed significantly higher than

that of the matched sensitive line MRA-1252 (and also higher than that of 3D7) (Fig 6E, left panel; Fig

S1C). Since the point mutation in Kelch13 is the only difference between the two Cambodian MRA

lines, a genetic background of enhanced DNA repair cannot fully account for this.

One other explanation for an improved DNA repair capacity in resistant parasites could be

that they have hyper-active checkpoint activity (possibly due to Kelch13-associated over-expression

of PI3K [32]) and that this stimulates hyperactive DNA repair. This explanation *would* be consistent

with MRA-1240 (Kelch13-mutant) and MRA-1252 (revertant) having different sensitivities to MMS.

If so, the Kelch13 mutant state would also be expected to correlate with reduced sensitivity to PIKK

inhibitors. Indeed, MRA-1240 was much less sensitive to the VE-821 inhibitor (Fig 6E, middle panel;

Fig S1C) – so much so that it was difficult to measure a consistent $IC_{50}$ via MSF assay. Curiously,

however, the same line was not less sensitive to KU-55933 (Fig 6E, right panel; Fig S1C). This was the

first instance in which VE-821 and KU-55933 had shown a clear difference in their effects on *P.*

*falciparum.* It may suggest that more than one checkpoint protein is required, at least in the MRA-

1240/1252 genetic background, and that one of the inhibitors has a broader inhibitory spectrum

than the other. For example, *Pf*PI3K may indeed moonlight as a checkpoint kinase, uniquely

unregulated after Kelch13 mutation and uniquely inhibited by VE-821, whereas *Pf*PI4K and/or other

'off-target' kinases may also act in certain checkpoint pathways, and KU-55933 may inhibit this

broader kinase spectrum, which is unaffected by the Kelch13 mutation in MRA-1240.

These speculations urgently require the conditional genetic manipulation of at least two

large, essential kinase-encoding genes, *PfPI3K* and *PfPI4K* – in parallel, in a molecular-genetic system

that can achieve meaningful levels of knockdown without cell death.

## 422 Discussion

This work provides the first evidence that *Plasmodium* parasites during blood-stage schizogony can

enforce an intra-S-phase checkpoint. The checkpoint is evidenced by an acute reduction in the rate

of DNA replication and increased phosphorylation of histone H2A in a position analogous to the

canonical DNA damage marker H2AX. *P. falciparum* was previously shown to slow its cell-cycle

progression after DNA damage [18] but it was not clear if this occurred through cell-cycle checkpoint

activity (partly because the *Plasmodium* system lacked validated DNA damage markers, signals or

transducers thereof). Markers still remain limited – there are, for example, no cyclin markers for

cell-cycle phases, and the H2A phosphorylation that has been reported as a DNA damage marker in

ring stages [25] is not, in our hands, a very strong marker in trophozoite stages (possibly because

background levels are higher in actively-replicating parasites). However, the ability to measure

nascent DNA replication in *Plasmodium* [56] does now allow intra-S-phase checkpoint activity to be

measured. Thus we showed that DNA replication was acutely arrested by two distinct alkylating

agents, MMS and the antimalarial drug DHA.

The transducer(s) of checkpoint activity in *Plasmodium* remain cryptic. *Plasmodium* spp.

lack identifiable orthologues of PIKK kinases, in contrast to many apicomplexans including *T. gondii*

[39]. Nevertheless, such activity is apparently present, because the reduction in replication caused

by MMS was ablated by well-validated inhibitors of human PIKKs. Hence it must be induced *in trans*

by a genuine kinase-mediated signalling cascade (also consistent with H2A phosphorylation), rather

than occurring simply because alkylated bases physically block the movement of replisomes.

Interestingly, DHA differed from MMS in this respect: it also affected DNA replication, but PIKK

inhibitors did not ameliorate this. Possibly the acute effect of DHA – which is a larger molecule than

MMS – *is* primarily to block replisomes via bulky DNA adducts (a documented effect of alkylating

agents in human cells [23]), and these cannot be bypassed whether a checkpoint kinase is active or

not, while a lesser role is played by *in trans* checkpoint signalling after DHA damage.

Other, more complex, explanations for the difference in response to MMS versus DHA are

also possible. For example, slow DNA replication can, in general, stimulate compensatory activation

of extra replication origins [57], but when DNA damage is sensed, checkpoint kinases act *in trans* to

suppress such origin firing [23]. If checkpoint kinases are inhibited, then extra origins may fire

regardless, hence MMS-damaged cells may keep replicating and encountering more damage, so

genome replication ultimately fails to complete. This would explain how MMS and PIKK inhibitors

ultimately synergise in cell killing. Perhaps the broader-spectrum damage that DHA causes to

proteins as well as DNA [51], and/or the oxidative stress that it generates [53] means that

compensatory origin firing does not happen, whether a checkpoint is activated or not. Hence PIKK

inhibitors would not synergise with DHA in cell killing. To evidence these rather complex

hypotheses, precision measurement of DNA replication dynamics at single-molecule resolution after

different types of damage will be needed. This has been achieved in human cells [23], and is now in

development for *Plasmodium*.

Turning to the possibility of a pre-replicative G1/S checkpoint, the evidence for this being

mediated by a PIKK-like kinase is not conclusive, but nevertheless tantalising.  In the RSA, which

measures DHA killing of pre-replicative rings, PIKK inhibitors exacerbated parasite killing.  The effect

was modest, so if a classical (kinase-enforced) G1/S checkpoint does exist, it is probably not the sole

defining factor in survival of artemininin-induced dormancy.   Nevertheless, the trend was consistent

(albeit not statistically significant) in triplicate assays across two genetically-distinct DHA-sensitive

strains, 3D7 and MRA-1252.  Checkpoint kinase activity could help to protect ring-stage cells,

perhaps by detecting DNA damage, stimulating repair, and preventing S-phase entry until repair is

complete.  By contrast, a DHA-resistant strain showed no effect of PIKK inhibitors.  This may be

because it can already suppress DHA activation in early rings and thus protect itself from DNA

damage; or alternatively, it may still sustain some damage, but be rescued by hyperactive DNA

repair [52].  Consistent with this latter hypothesis, it was recently shown that when *Pf*PI3K is

inducibly knocked out in a DHA-resistant Kelch-13 mutant background, these parasites become less

DHA resistant – perhaps because they can no longer resist or repair DHA-related DNA damage when

the mediating kinase, PI3K, is lost [49].

If there is indeed hyperactive DNA repair in the MRA-1240 line, it cannot be conferred by a

DHA-resistant genetic background because the matched strain MRA-1252 was not cross-resistant to

MMS, whereas MRA-1240 was.  However, it could be because the Kelch13 mutation, uniquely

present in MRA-1240, confers hyper-checkpoint activity, thus stimulating hyperactive DNA repair.  In

any case, these data raise the intriguing possibility that artemisinin drugs could synergise with

checkpoint inhibitors in killing ring-stage parasites; the caveat being that this might only work well in

sensitive, not resistant, strains.

It is an attractive idea that *Pf*PI3K is the cryptic checkpoint kinase, particularly if Kelch13

mutant strains contain more *Pf*PI3K [32].  However, we were not yet able to prove or disprove this,

owing to the difficulty of manipulating the large, essential *PfPI3K* gene.  Broad-spectrum inhibitors

such as wortmannin were used in previous work [36], but these are slow-killing and unable to

separate lipid-kinase from putative protein-kinase activity.  We used PIKK inhibitors instead, hoping

to isolate a putative protein kinase activity.  The fact that our PIKK inhibitors did not synergise with

DHA in MSF assays, whereas specific inhibitors of distinct human PI3Ks (i.e. lipid kinases) *have* been

reported to do so [58], may suggest that this succeeded, and that VE-821 and KU-55933 do target a

distinct activity in *Plasmodium*.  They may, of course, merely affect irrelevant off-targets – multiple

or unexpected kinases, or perhaps even non-kinases.  As an example, a repurposed human-kinase

inhibitor was recently deployed to target *Plasmodium* PK6, but it actually targeted haem

detoxification pathways as well [59].  More likely, however, is that these inhibitors genuinely inhibit

a *Plasmodium* checkpoint kinase – which may reside in *Pf*PI3K, *Pf*PI4K, or another radically divergent

protein. *Plasmodium* contains many kinases whose roles are yet-unknown [60], and the *Plasmodium*

cell cycle is driven by markedly divergent kinases such as CRK4 [61], so it is entirely possible that

*Plasmodium* checkpoints have evolved to be enforced by novel kinase(s) as well.  In conclusion, we

present new evidence for measurable cell-cycle checkpoint activity in *Plasmodium*.  We also raise

the considerable challenge of unambiguously identifying its mediators in this divergent eukaryotic

parasite.

# Methods

## Parasite cultures

503    *Plasmodium falciparum* cultures were grown in 2% or 4% haematocrit human erythrocytes

504    (Research Red Cells obtained from the NHS Blood & Transplant service), in RPMI 1640 (Sigma,

505    R4130) supplemented with 2.3 g/L sodium bicarbonate, 50 mg/L hypoxanthine (Sigma, H9377), 25

506    μg/L gentamicin (Melford Laboratories, G38000-1), 2.5 g/L Albumax II and 5% human serum.

507    Parasites were cultured under a 1% $O_2$/ 3% $CO_2$/ 96% $N_2$ gaseous atmosphere at 37 °C.

## Parasite strains

Parasite strains used were: 3D7, 3D7 pTK, iPI3K-ko (also in the 3D7 background) [49], MRA-1240 and

MRA-1252. 3D7 pTK carries a thymidine kinase transgene to allow incorporation of

bromodeoxyuridine (BrdU) for monitoring DNA replication [56]. The iPI3K-ko line carries a D-Cre-

inducible deletion module for the C-terminal part of the *PI3K* gene [49].  MRA-1240 and MRA-1252

are matched artemisinin-resistant and sensitive Cambodian field lines, respectively [54], obtained

through BEI Resources (NIAID, NIH). Parasites were synchronised as in [62].  The PI3K inducible

knockout was enforced by treating ring-stage parasites with 100nM rapamycin (or DMSO vehicle for

the control), as described by the original authors [49], before conducting MSF assays 24-72h or 24-

96h later.

## Database sampling and phylogenetic analysis

The phylogeny (Fig 2) for ATM, ATR, TOR, PI3K and PI4K homologues was inferred by following a

computational workflow (https://github.com/camwallerlab/Methods-for-phylogenetic-analysis-of-

plastid-translocons) developed as part of a previous study [63]. In summary, this workflow involved

sampling a custom protein sequence database that included data for major eukaryotic groups from

UniProt [64] and the Marine Microbial Eukaryote Transcriptome Sequencing Project [65], as well as

data for apicomplexans and close relatives from VEuPathDB [66] and two previous studies [67, 68].

This custom database was then sampled for ATM, ATR and TOR homologues using the program

blastp (BLAST+ version 2.11.0 [69]), for which characterised homologues of these proteins from

selected model organisms were used as search queries. The dataset of possible ATM, ATR and TOR

homologues obtained was then clustered with CD-HIT [70] to remove highly similar sequences,

thereby reducing overall redundancy of the dataset.  PI3K and PI4K homologues from *Plasmodium*

and other apicomplexans, as well as characterised homologues from *Saccharomyces cerevisiae* and

*Homo sapiens* were incorporated into the dataset as a phylogenetic outgroup for the ATM, ATR and

TOR proteins. Iterative rounds of alignment were then performed using mafft (MAFFT version 7.475

[71]), conserved site selection using trimal with "gappyout" mode (trimAl version 1.4 [72]), and tree

inference with FastTreeMP (FastTree version 2.1.11 [73]) using the default settings.  For each

iterlation, sequences that aligned poorly or were so dissimilar from the rest of the sequences in the

dataset that they were unlikely to be ATM, ATR or TOR homologues, but rather false positives of the

described sampling strategy, were manually identified and removed from the dataset. The final

curated dataset that these iterations obtained was then aligned using mafft-linsi (MAFFT version

7.475 [71]) and conserved sites were selected using trimal with "gappyout" mode (trimAl version 1.4

[72]). A phylogeny was then inferred from this dataset using the program iqtree2 (IQ-TREE version

2.1.2 [74]) with 1000 ultrafast bootstrap replicates (UFBoot2 [33]) and using the best-fitting model,

LG+F+I+G4 ([75]) that was chosen according to the Bayesian Information Criterion by ModelFinder

[76], all implemented within iqtree2.  The protein sequences, alignment, and tree inference output

files used to generate the phylogeny are provided as source data.

## BrdU Enzyme-linked immunosorbent assay (ELISA)

ELISA was carried out as per [56]. In brief, tightly synchronised *P. falciparum* 3D7 pTK parasites were

collected at 4% parasitaemia and 4% haematocrit at 28 hpi, 32 hpi and 36 hpi. They were labelled

with BrdU (100 μM, Sigma) for 30 min and then collected for ELISA, or treated with MMS at 10 x $IC_{50}$

(960 μM) or 20 x $IC_{50}$ (1,920 μM) for 30 min prior to BrdU labelling. In further experiments, parasites

at 30-32 hpi were exposed to either MMS alone (10x $IC_{50}$), MMS+VE-821 (1x $IC_{50}$), MMS+KU-55933

(1x $IC_{50}$), VE-821 alone, KU-55933 alone, DHA alone (10x $IC_{50}$), DHA+VE-821, DHA+KU-55933 or no

drugs, then processed as above.

**Immunofluorescence**

Air-dried thick blood smears were fixed with 4% paraformaldehyde in PBS for 10 min and

permeabilised for 10 min with 0.05% triton in PBS. Slides were then blocked for 1 h in 1% BSA/PBS.

The primary antibody was added in blocking buffer for 1 h under a coverslip. After 3x 5 min washes

in PBS, the secondary antibody was added to slides for 1 h. The slides were washed with PBS for 5

min, 120 µl of 2 µg/ml DAPI solution was added for 10 min and then the slides washed again with

PBS for 5 min. Coverslips were mounted on slides in Prolong® Diamond (Invitrogen) and allowed to

dry overnight at 4 °C. Images were acquired using a Nikon Eclipse Ti widefield microscope with a

Nikon objective lens (Plan APO, 100x/1.45 oil), and a Hamamatsu C11440, ORCA Flash 4.0 camera.

Images were processed using the NIS-Elements software and ImageJ (v.1.51w).

## Malaria SYBR Green-1 Fluorescence (MSF) assay

566    MSF assays were conducted as in [77].   100 µM chloroquine (Merck) was used as a control for

567    parasite death. Data were analysed and graphs plotted in GraphPad Prism.

## Isobolograms

Parasites were prepared as for MSF assays. Isobologram design followed the procedure in [78].

Serial dilutions were planned so that the $IC_{50}$s of each drug would be in the second two-fold dilution

out of 6 total dilutions.  $IC_{50}$s for each combination solution were calculated using sigmoidal best-fit

curves assigned in Graphpad Prism.  Fractional inhibitory concentrations (FIC values) were calculated

in Excel [79] and isobolograms plotted in Graphpad Prism. Source data for the isobolograms are

provided in figure S8.

## Ring-stage Survival Assay (RSA)

Parasite cultures at 8-10% schizonts were synchronised using percoll and Compound 2 as in [62],

then allowed to reinvade. Early ring-stage (0-3 hpi) parasites at 0.5% parasitaemia, 2% haematocrit

were exposed for 6 h to 700 nM DHA or 0.02% DMSO (vehicle control) under normal conditions,

before washing three times with incomplete media, resuspending in complete media and incubating

under the same conditions for a further 66 h. In addition to the DHA 700 nM pulse, VE-821 and KU-

55933 inhibitors were used at 0.5x IC$_{50}$ with DHA, or by themselves, throughout the 6 h pulse period.

72 h after initial drug exposure, 20 µl was frozen for RT-PCR as per eRRSA protocol [55].

## qPCR-based RSA analysis

qPCR amplification was used to quantify live parasites in the eRRSA [55].  The Phusion Blood Direct

PCR kit was used (ThermoFisher, cat # F547L), supplemented with 1x SYBR Green I (ThermoFisher)

and 50 nM Low ROX (ThermoFisher), according to the manufacturer's instructions. qPCR

amplification was measured using the fast mode of the ABI 7900HT, with a 20 s denaturation at 95

°C, followed by 35 cycles of 95 °C for 1 s, 62.3 °C for 30 s, and 65 °C for 15 s. Cycle threshold (Ct)

values were calculated; then the fold change was calculated by determining the mean difference in

Ct (ΔCt) for the three technical replicates between the untreated and DHA-treated samples by

applying the following equation:  Fold change =2^(ΔCt of treated − ΔCt of untreated). This was -log

transformed and data were plotted in GraphPad Prism.

## Flow cytometry-based RSA analysis

Flow cytometry wash buffer 'FC wash' (2% FBS (Gibco) in HBSS (calcium, magnesium, no phenol red,

Gibco) was prepared.  The remaining 180 µl of each RSA assay, not harvested for eRRSA, was

centrifuged and packed cells were washed three times in 200 µL of FC wash, then resuspended in FC

wash to make a 10% haematocrit suspension. 25 µL of suspension from each assay (conducted in

technical triplicate) were mixed with 25 µL of 0.6X SYBR Green I (SYBR) with 1.5 nM MitoTracker

Deep Red FM (MT, Invitrogen) in FC wash. Compensation controls were prepared with suspension

from one DMSO-treated triplicate, labelled with either 0.6X SYBR, 1.5 nM MT, or FC wash alone.

Samples were incubated in the dark at 37°C for 30 min, then washed three times in 200 µL and

resuspended in 180 µL of FC wash.

10,000 events of SYBR positive cells were captured at a flow rate of 25 µL/min on an Attune NxT

Acoustic Focusing Cytometer (Invitrogen). SYBR and MT fluorescence were detected on the BL1 (488

nm excitation, 530/30 nm emission, 340 V) and RL1 (638 nm excitation, 670/14 nm emission, 510 V)

channels respectively. Debris and doublets were excluded by gating on a forward scatter area (FSC-

A) versus side scatter height (SSC-H) plot and forward scatter area (FSC-A) versus forward scatter

height (FSC-H). SYBR/MT quadrant plots were generated using the Attune NxT software v4.2.0

(Invitrogen).

## Antibody production

Peptides were designed from *P. falciparum* PI3K (PF3D7_0515300) with help from Eurogentec using

parameters such as size, amino acid composition, hydrophobicity and secondary structure. Two

peptides were chosen - one predicted to be in the catalytic domain and one upstream of this

domain, both predicted to be surface-displayed. N-terminal cysteines were added to the peptide

sequences to target the coupling site at the carrier protein.

Antibodies were raised by Eurogentec. Two rabbits were bled for pre-immune sera, inoculated with

both peptides, then further immunised with 3 booster inoculations at 12 days, ~1 month and

~2months. Immune serum was collected after ~9 weeks.

## Western blotting

Parasites were extracted from infected erythrocytes by mixing with 0.5 volumes of 0.2% saponin in

PBS and incubating for 10 min on ice. Saponin lysates were centrifuged at 15,000 *xg* at 4 °C for 10

min, washed 3 x in ice-cold PBS, centrifuged at 15,000 *xg* for 5 min and resuspended in PBS before

freezing at -20°C. Parasites obtained by saponin lysis were resuspended in 1x RIPA buffer (150 mM

sodium chloride, 1.0% NP-40 or Triton X-100, 0.5% sodium deoxycholate, 0.1% SDS and 50 mM Tris

pH 8) with protease inhibitors (ROCHE, Complete Mini) added directly before use, then freeze-

thawed 3 times. Cell debris was removed by centrifugation at 4 °C, 13,000 *xg*, and supernatant was

mixed with 1 x NuPage LDS Sample Buffer (ThermoFisher: NP0007), supplemented with 0.1% 2-

mercaptoethanol, then boiled at 90 °C for 5 min. Protein lysates were resolved on 4-15% TRIS-

glycine gels (BIO-RAD Mini -PROTEAN TGX) for 1–2 h at 100 V in 1X TGS buffer. Wet transfer was

completed onto nitrocellulose membrane 0.2-µM-pore-size (Amersham Protran®, GE Healthcare)

at 60 V for 45 min.

Membranes were blocked with 3% BSA TBST for 1 h, washed once in TBST and incubated overnight

in primary antibody diluted in blocking buffer at 4 °C. Blots were washed 3x 5 min in TBST and

incubated in a secondary antibody diluted 1:5000 in blocking buffer for 1 h. Blots were washed again

(3x 5 min, TBST), developed using SuperSignal West Pico Plus Chemiluminescent Substrate and

imaged on a gel imager (Azure, Q500). Some membranes were stripped prior to re-blocking (10 ml

20% SDS and 12.5 ml 0.5M Tris HCL, pH 6.8, added to 77.5 ml distilled water, with 0.8 ml β-

mercaptoethanol added just before use).  5 ml stripping buffer (warmed to 50 °C) was added to a

blot and incubated at 50 °C for 45 min with agitation. The stripping buffer was removed, and the

membrane was washed under a running tap for 1-2 min, then washed 3x 5 min with TBST. The

membrane was then ready for re-blocking.

Antibodies used were pre-immune or immune sera for *Pf*PI3K peptides, diluted 1:200, with

secondary goat anti-rabbit HRP 1:5000 (Abcam, ab97080); or rabbit $\alpha$-$\gamma$-H2aX (Cell signalling, 9718S)

with goat $\alpha$-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 594 (Invitrogen, A-

11012).

# Acknowledgements

We would like to thank Dr Holly Matthews for early work on this project, Dr Ibtissam Jabre for help

with H2A-P blots, Prof Kai Wengelnik for sharing the PI3K inducible knockout line swiftly during

revision of this manuscript, and Profs Marcus Lee and Paul Roepe for helpful early discussions.

This work was funded by the European Research Council (ERC) under the European Union's Horizon

2020 research and innovation programme (ERC-2016-COG 725126 to CJM); and a PhD studentship

from the University of Cambridge Department of Pathology to MKJ. The funders had no role in study

design, data collection, interpretation or the decision to submit the work for publication.

## References

1. WHO. World Malaria Report 2023. 2023. Epub 2023.

2. Klaus S, Binder P, Kim J, Machado M, Funaya C, Schaaf V, et al. Asynchronous nuclear cycles
in multinucleated Plasmodium falciparum facilitate rapid proliferation. Sci Adv. 2022;8(13):eabj5362.
doi: 10.1126/sciadv.abj5362. PubMed PMID: 35353560.

3. McDonald J, Merrick CJ. DNA replication dynamics during erythrocytic schizogony in the
malaria parasites Plasmodium falciparum and Plasmodium knowlesi. PLoS pathogens. 2022;18(6):
e1010595.

4. Simon CS, Sturmer VS, Guizetti J. How Many Is Enough? - Challenges of Multinucleated Cell
Division in Malaria Parasites. Front Cell Infect Microbiol. 2021;11:658616. doi:
10.3389/fcimb.2021.658616. PubMed PMID: 34026661; PubMed Central PMCID: PMCPMC8137892.

5. Liffner B, Cepeda Diaz AK, Blauwkamp J, Anaguano D, Frolich S, Muralidharan V, et al. Atlas
of Plasmodium falciparum intraerythrocytic development using expansion microscopy. Elife.
2023;12. doi: 10.7554/eLife.88088. PubMed PMID: 38108809; PubMed Central PMCID:
PMCPMC10727503.

6. Matthews H, Duffy CW, Merrick CJ. Checks and balances? DNA replication and the cell cycle
in Plasmodium. Parasit Vectors. 2018;11(1):216. doi: 10.1186/s13071-018-2800-1. PubMed PMID:
29587837; PubMed Central PMCID: PMCPMC5872521.

7. Zhou BB, Elledge SJ. The DNA damage response: putting checkpoints in perspective. Nature.
2000;408(6811):433-9. doi: 10.1038/35044005. PubMed PMID: 11100718.

8. Hawkins LM, Naumov AV, Batra M, Wang C, Chaput D, Suvorova ES. Novel CRK-Cyclin
Complex Controls Spindle Assembly Checkpoint in Toxoplasma Endodyogeny. mBio.
2021;13(1):e0356121. doi: 10.1128/mbio.03561-21. PubMed PMID: 35130726; PubMed Central
PMCID: PMCPMC8822342.

9. Naumov AV, Wang C, Chaput D, Ting LM, Alvarez CA, Keller T, et al. Restriction Checkpoint
Controls Bradyzoite Development in Toxoplasma gondii. Microbiol Spectr. 2022;10(3):e0070222. doi:
10.1128/spectrum.00702-22. PubMed PMID: 35652638; PubMed Central PMCID: PMCPMC9241953.

10. Alvarez CA, Suvorova ES. Checkpoints of apicomplexan cell division identified in Toxoplasma
gondii. PLoS pathogens. 2017;13(7):e1006483. doi: 10.1371/journal.ppat.1006483. PubMed PMID:
28671988; PubMed Central PMCID: PMCPMC5510908.

11. Lee AH, Symington LS, Fidock DA. DNA repair mechanisms and their biological roles in the
malaria parasite Plasmodium falciparum. Microbiol Mol Biol Rev. 2014;78(3):469-86. doi:
10.1128/MMBR.00059-13. PubMed PMID: 25184562; PubMed Central PMCID: PMCPMC4187680.

12. Claessens A, Hamilton WL, Kekre M, Otto TD, Faizullabhoy A, Rayner JC, et al. Generation of
antigenic diversity in Plasmodium falciparum by structured rearrangement of Var genes during
mitosis. PLoS genetics. 2014;10(12):e1004812. doi: 10.1371/journal.pgen.1004812. PubMed PMID:
25521112; PubMed Central PMCID: PMCPMC4270465.

13.    Platon L, Menard D. Plasmodium falciparum ring-stage plasticity and drug resistance. Trends
Parasitol. 2024;40(2):118-30. doi: 10.1016/j.pt.2023.11.007. PubMed PMID: 38104024.
14.    Babbitt SE, Altenhofen L, Cobbold SA, Istvan ES, Fennell C, Doerig C, et al. Plasmodium
falciparum responds to amino acid starvation by entering into a hibernatory state. Proceedings of
the National Academy of Sciences of the United States of America. 2012;109(47):E3278-87. doi:
10.1073/pnas.1209823109. PubMed PMID: 23112171; PubMed Central PMCID: PMCPMC3511138.
15.    Yuan L, Hao M, Wu L, Zhao Z, Rosenthal BM, Li X, et al. Refrigeration provides a simple
means to synchronize in vitro cultures of Plasmodium falciparum. Exp Parasitol. 2014;140:18-23. doi:
10.1016/j.exppara.2014.03.010. PubMed PMID: 24632190; PubMed Central PMCID:
PMCPMC4018460.
16.    Bohorquez EB, Juliano JJ, Kim HS, Meshnick SR. Mefloquine exposure induces cell cycle delay
and reveals stage-specific expression of the pfmdr1 gene. Antimicrobial agents and chemotherapy.
2013;57(2):833-9. doi: 10.1128/AAC.01006-12. PubMed PMID: 23208721; PubMed Central PMCID:
PMCPMC3553685.
17.    Witkowski B, Lelievre J, Barragan MJ, Laurent V, Su XZ, Berry A, et al. Increased tolerance to
artemisinin in Plasmodium falciparum is mediated by a quiescence mechanism. Antimicrobial agents
and chemotherapy. 2010;54(5):1872-7. doi: 10.1128/AAC.01636-09. PubMed PMID: 20160056;
PubMed Central PMCID: PMCPMC2863624.
18.    Gupta DK, Patra AT, Zhu L, Gupta AP, Bozdech Z. DNA damage regulation and its role in drug-
related phenotypes in the malaria parasites. Sci Rep. 2016;6:23603. doi: 10.1038/srep23603.
PubMed PMID: 27033103; PubMed Central PMCID: PMCPMC4817041.
19.    Voss Y, Klaus S, Guizetti J, Ganter M. Plasmodium schizogony, a chronology of the parasite's
cell cycle in the blood stage. PLoS pathogens. 2023;19(3):e1011157. doi:
10.1371/journal.ppat.1011157. PubMed PMID: 36862652; PubMed Central PMCID:
PMCPMC9980825 commercial or financial relationships that could be construed as a potential
conflict of interest.
20.    Absalon S, Dvorin JD. Depletion of the mini-chromosome maintenance complex binding
protein allows the progression of cytokinesis despite abnormal karyokinesis during the asexual
development of Plasmodium falciparum. Cellular microbiology. 2021;23(3):e13284. doi:
10.1111/cmi.13284. PubMed PMID: 33124706; PubMed Central PMCID: PMCPMC8058698.
21.    Janse CJ, van der Klooster PF, van der Kaay HJ, van der Ploeg M, Overdulve JP. DNA synthesis
in Plasmodium berghei during asexual and sexual development. Molecular and biochemical
parasitology. 1986;20(2):173-82. doi: 10.1016/0166-6851(86)90029-0. PubMed PMID: 3092048.
22.    Matthews H, McDonald J, Totanes FIG, Merrick CJ. Dynamics of DNA replication during male
gametogenesis in the malaria parasite Plasmodium falciparum. Cellular microbiology. 2022. doi:
doi.org/10.1101/2021.12.18.473304
23.    Merrick CJ, Jackson D, Diffley JF. Visualization of altered replication dynamics after DNA
damage in human cells. The Journal of biological chemistry. 2004;279(19):20067-75. PubMed PMID:
14982920.
24.    Santocanale C, Diffley JF. A Mec1- and Rad53-dependent checkpoint controls late-firing
origins of DNA replication. Nature. 1998;395(6702):615-8. doi: 10.1038/27001. PubMed PMID:
9783589.
25.    Goyal M, Heinberg A, Mitesser V, Kandelis-Shalev S, Singh BK, Dzikowski R. Phosphorylation
of the Canonical Histone H2A Marks Foci of Damaged DNA in Malaria Parasites. mSphere. 2021;6(1).
doi: 10.1128/mSphere.01131-20. PubMed PMID: 33441412; PubMed Central PMCID:
PMCPMC7845613.
26.    Helt CE, Cliby WA, Keng PC, Bambara RA, O'Reilly MA. Ataxia telangiectasia mutated (ATM)
and ATM and Rad3-related protein exhibit selective target specificities in response to different forms
of DNA damage. The Journal of biological chemistry. 2005;280(2):1186-92. doi:
10.1074/jbc.M410873200. PubMed PMID: 15533933.

27.     Rogakou EP, Pilch DR, Orr AH, Ivanova VS, Bonner WM. DNA double-stranded breaks induce
histone H2AX phosphorylation on serine 139. The Journal of biological chemistry.
1998;273(10):5858-68. doi: 10.1074/jbc.273.10.5858. PubMed PMID: 9488723.
28.     Beranek DT. Distribution of methyl and ethyl adducts following alkylation with
monofunctional alkylating agents. Mutat Res. 1990;231(1):11-30. doi: 10.1016/0027-5107(90)90173-
2. PubMed PMID: 2195323.
29.     Gopalakrishnan AM, Kumar N. Opposing roles for two molecular forms of replication protein
A in Rad51-Rad54-mediated DNA recombination in Plasmodium falciparum. mBio.
2013;4(3):e00252-13. doi: 10.1128/mBio.00252-13. PubMed PMID: 23631919; PubMed Central
PMCID: PMCPMC3648904.
30.     Craven HM, Nettesheim G, Cicuta P, Blagborough AM, Merrick CJ. Effects of the G-
quadruplex-binding drugs quarfloxin and CX-5461 on the malaria parasite Plasmodium falciparum.
Int J Parasitol Drugs Drug Resist. 2023;23:106-19. doi: 10.1016/j.ijpddr.2023.11.007. PubMed PMID:
38041930; PubMed Central PMCID: PMCPMC10711401.
31.     Hassett MR, Roepe PD. PIK-ing New Malaria Chemotherapy. Trends Parasitol.
2018;34(11):925-7. doi: 10.1016/j.pt.2018.06.003. PubMed PMID: 29934102.
32.     Mbengue A, Bhattacharjee S, Pandharkar T, Liu H, Estiu G, Stahelin RV, et al. A molecular
mechanism of artemisinin resistance in Plasmodium falciparum malaria. Nature.
2015;520(7549):683-7. doi: 10.1038/nature14412. PubMed PMID: 25874676; PubMed Central
PMCID: PMCPMC4417027.
33.     Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. UFBoot2: Improving the
Ultrafast Bootstrap Approximation. Mol Biol Evol. 2018;35(2):518-22. doi: 10.1093/molbev/msx281.
PubMed PMID: 29077904; PubMed Central PMCID: PMCPMC5850222.
34.     Hassett MR, Sternberg AR, Riegel BE, Thomas CJ, Roepe PD. Heterologous Expression,
Purification, and Functional Analysis of Plasmodium falciparum Phosphatidylinositol 3'-Kinase.
Biochemistry. 2017;56(33):4335-45. doi: 10.1021/acs.biochem.7b00416. PubMed PMID: 28719180.
35.     Tawk L, Chicanne G, Dubremetz JF, Richard V, Payrastre B, Vial HJ, et al. Phosphatidylinositol
3-phosphate, an essential lipid in Plasmodium, localizes to the food vacuole membrane and the
apicoplast. Eukaryotic cell. 2010;9(10):1519-30. doi: 10.1128/EC.00124-10. PubMed PMID:
20709789; PubMed Central PMCID: PMCPMC2950420.
36.     Vaid A, Ranjan R, Smythe WA, Hoppe HC, Sharma P. PfPI3K, a phosphatidylinositol-3 kinase
from Plasmodium falciparum, is exported to the host erythrocyte and is involved in hemoglobin
trafficking. Blood. 2010;115(12):2500-7. doi: 10.1182/blood-2009-08-238972. PubMed PMID:
20093402; PubMed Central PMCID: PMCPMC2918364.
37.     Thomas D, Powell JA, Green BD, Barry EF, Ma Y, Woodcock J, et al. Protein kinase activity of
phosphoinositide 3-kinase regulates cytokine-dependent cell survival. PLoS biology.
2013;11(3):e1001515. doi: 10.1371/journal.pbio.1001515. PubMed PMID: 23526884; PubMed
Central PMCID: PMCPMC3601961.
38.     Foukas LC, Shepherd PR. Phosphoinositide 3-kinase: the protein kinase that time forgot.
Biochem Soc Trans. 2004;32(Pt 2):330-1. doi: 10.1042/bst0320330. PubMed PMID: 15046601.
39.     Johnson MK, Chelaghma S, Koreny L, Waller RF, Merrick CJ. Characterisation of cell cycle
checkpoint kinases in Toxoplasma gondii. BioRxiv. 2024:doi.org/10.1101/2024.09.10.612042. doi:
doi.org/10.1101/2024.09.10.612042;.
40.     Arendse LB, Murithi JM, Qahash T, Pasaje CFA, Godoy LC, Dey S, et al. The anticancer human
mTOR inhibitor sapanisertib potently inhibits multiple Plasmodium kinases and life cycle stages. Sci
Transl Med. 2022;14(667):eabo7219. doi: 10.1126/scitranslmed.abo7219. PubMed PMID: 36260689;
PubMed Central PMCID: PMCPMC9951552.
41.     Mukherjee A, Crochetiere ME, Sergerie A, Amiar S, Thompson LA, Ebrahimzadeh Z, et al. A
Phosphoinositide-Binding Protein Acts in the Trafficking Pathway of Hemoglobin in the Malaria
Parasite Plasmodium falciparum. mBio. 2022;13(1):e0323921. doi: 10.1128/mbio.03239-21. PubMed
PMID: 35038916; PubMed Central PMCID: PMCPMC8764524.

42.       Oehring SC, Woodcroft BJ, Moes S, Wetzel J, Dietz O, Pulfer A, et al. Organellar proteomics
reveals hundreds of novel nuclear proteins in the malaria parasite Plasmodium falciparum. Genome
biology. 2012;13(11):R108. doi: 10.1186/gb-2012-13-11-r108. PubMed PMID: 23181666; PubMed
Central PMCID: PMCPMC4053738.
43.       Briquet S, Ourimi A, Pionneau C, Bernardes J, Carbone A, Chardonnet S, et al. Identification
of Plasmodium falciparum nuclear proteins by mass spectrometry and proposed protein annotation.
PloS one. 2018;13(10):e0205596. doi: 10.1371/journal.pone.0205596. PubMed PMID: 30379851;
PubMed Central PMCID: PMCPMC6209197.
44.       Barylyuk K, Koreny L, Ke H, Butterworth S, Crook OM, Lassadi I, et al. A Comprehensive
Subcellular Atlas of the Toxoplasma Proteome via hyperLOPIT Provides Spatial Context for Protein
Functions. Cell host & microbe. 2020;28(5):752-66 e9. doi: 10.1016/j.chom.2020.09.011. PubMed
PMID: 33053376; PubMed Central PMCID: PMCPMC7670262.
45.       Birnbaum J, Scharf S, Schmidt S, Jonscher E, Hoeijmakers WAM, Flemming S, et al. A Kelch13-
defined endocytosis pathway mediates artemisinin resistance in malaria parasites. Science (New
York, NY. 2020;367(6473):51-9. doi: 10.1126/science.aax4735. PubMed PMID: 31896710.
46.       Reaper PM, Griffiths MR, Long JM, Charrier JD, Maccormick S, Charlton PA, et al. Selective
killing of ATM- or p53-deficient cancer cells through inhibition of ATR. Nat Chem Biol. 2011;7(7):428-
30. doi: 10.1038/nchembio.573. PubMed PMID: 21490603.
47.       Hickson I, Zhao Y, Richardson CJ, Green SJ, Martin NM, Orr AI, et al. Identification and
characterization of a novel and specific inhibitor of the ataxia-telangiectasia mutated kinase ATM.
Cancer research. 2004;64(24):9152-9. doi: 10.1158/0008-5472.CAN-04-2727. PubMed PMID:
15604286.
48.       Safeukui I, Ware RE, Mohandas N, Haldar K. Simultaneous adjunctive treatment of malaria
and its coevolved genetic disorder sickle cell anemia. Blood Adv. 2023;7(19):5970-81. doi:
10.1182/bloodadvances.2022009124. PubMed PMID: 37093647; PubMed Central PMCID:
PMCPMC10580175 interests.
49.       Al Monla R, Penzo M, Vallentin A, Lohia R, Vincent J, Berry L, et al. PI3-kinase has multiple
functions in asexual blood stages of Plasmodium falciparum. Sci Rep. 2025;15(1):16762. doi:
10.1038/s41598-025-01397-1. PubMed PMID: 40369090; PubMed Central PMCID:
PMCPMC12078608.
50.       Bower-Lepts C, Rawat M, Keith WC, Girling G, Luth MR, Carolino K, et al. A novel
chemogenomic screening platform for scalable antimalarial drug target identification. bioRxiv.
2024:2024.08.27.609875. doi: 10.1101/2024.08.27.609875.
51.       Ismail HM, Barton V, Phanchana M, Charoensutthivarakul S, Wong MH, Hemingway J, et al.
Artemisinin activity-based probes identify multiple molecular targets within the asexual stage of the
malaria parasites Plasmodium falciparum 3D7. Proceedings of the National Academy of Sciences of
the United States of America. 2016;113(8):2080-5. doi: 10.1073/pnas.1600459113. PubMed PMID:
26858419; PubMed Central PMCID: PMCPMC4776496.
52.       Xiong A, Prakash P, Gao X, Chew M, Tay IJJ, Woodrow CJ, et al. K13-Mediated Reduced
Susceptibility to Artemisinin in Plasmodium falciparum Is Overlaid on a Trait of Enhanced DNA
Damage Repair. Cell Rep. 2020;32(5):107996. doi: 10.1016/j.celrep.2020.107996. PubMed PMID:
32755588; PubMed Central PMCID: PMCPMC7408483.
53.       Gopalakrishnan AM, Kumar N. Antimalarial action of artesunate involves DNA damage
mediated by reactive oxygen species. Antimicrobial agents and chemotherapy. 2015;59(1):317-25.
doi: 10.1128/AAC.03663-14. PubMed PMID: 25348537; PubMed Central PMCID: PMCPMC4291367.
54.       Straimer J, Gnadig NF, Witkowski B, Amaratunga C, Duru V, Ramadani AP, et al. Drug
resistance. K13-propeller mutations confer artemisinin resistance in Plasmodium falciparum clinical
isolates. Science (New York, NY. 2015;347(6220):428-31. doi: 10.1126/science.1260867. PubMed
PMID: 25502314; PubMed Central PMCID: PMCPMC4349400.
55.       Davis SZ, Singh PP, Vendrely KM, Shoue DA, Checkley LA, McDew-White M, et al. The
extended recovery ring-stage survival assay provides a superior association with patient clearance

half-life and increases throughput. Malaria journal. 2020;19(1):54. doi: 10.1186/s12936-020-3139-6.
PubMed PMID: 32005233; PubMed Central PMCID: PMCPMC6995136.
56. Merrick CJ. Transfection with thymidine kinase permits bromodeoxyuridine labelling of DNA
replication in the human malaria parasite Plasmodium falciparum. Malar J. 2015;14(490). doi:
10.1186/s12936-015-1014-7.
57. Courtot L, Hoffmann JS, Bergoglio V. The Protective Role of Dormant Origins in Response to
Replicative Stress. Int J Mol Sci. 2018;19(11). doi: 10.3390/ijms19113569. PubMed PMID: 30424570;
PubMed Central PMCID: PMCPMC6274952.
58. Mott BT, Eastman RT, Guha R, Sherlach KS, Siriwardana A, Shinn P, et al. High-throughput
matrix screening identifies synergistic and antagonistic antimalarial drug combinations. Sci Rep.
2015;5:13891. doi: 10.1038/srep13891. PubMed PMID: 26403635; PubMed Central PMCID:
PMCPMC4585899.
59. Nardella F, Jiang T, Wang L, Bohmer MJ, Chakraborty S, Okombo J, et al.
Plasmodiumfalciparum protein kinase 6 and hemozoin formation are inhibited by a type II human
kinase inhibitor exhibiting antimalarial activity. Cell Chem Biol. 2025;32(7):926-41 e23. doi:
10.1016/j.chembiol.2025.06.003. PubMed PMID: 40628257.
60. Ward P, Equinet L, Packer J, Doerig C. Protein kinases of the human malaria parasite
Plasmodium falciparum: the kinome of a divergent eukaryote. BMC genomics. 2004;5:79. doi:
10.1186/1471-2164-5-79. PubMed PMID: 15479470; PubMed Central PMCID: PMCPMC526369.
61. Ganter M, Goldberg JM, Dvorin JD, Paulo JA, King JG, Tripathi AK, et al. Plasmodium
falciparum CRK4 directs continuous rounds of DNA replication during schizogony. Nat Microbiol.
2017;2:17017. doi: 10.1038/nmicrobiol.2017.17. PubMed PMID: 28211852; PubMed Central PMCID:
PMCPMC5328244.
62. Totanes FIG, Gockel J, Chapman SE, Bartfai R, Boemo MA, Merrick CJ. A genome-wide map
of DNA replication at single-molecule resolution in the malaria parasite Plasmodium falciparum.
Nucleic acids research. 2023. doi: 10.1093/nar/gkad093. PubMed PMID: 36808528.
63. Lewis WH, Paris G, Beedessee G, Koreny L, Flores V, Dendooven T, et al. Plastid translocon
recycling in dinoflagellates demonstrates the portability of complex plastids between hosts. Curr
Biol. 2024;34(23):5494-506 e3. doi: 10.1016/j.cub.2024.10.034. PubMed PMID: 39571577; PubMed
Central PMCID: PMCPMC7617431.
64. UniProt C. UniProt: the Universal Protein Knowledgebase in 2023. Nucleic acids research.
2023;51(D1):D523-D31. doi: 10.1093/nar/gkac1052. PubMed PMID: 36408920; PubMed Central
PMCID: PMCPMC9825514.
65. Keeling PJ, Burki F, Wilcox HM, Allam B, Allen EE, Amaral-Zettler LA, et al. The Marine
Microbial Eukaryote Transcriptome Sequencing Project (MMETSP): illuminating the functional
diversity of eukaryotic life in the oceans through transcriptome sequencing. PLoS biology.
2014;12(6):e1001889. doi: 10.1371/journal.pbio.1001889. PubMed PMID: 24959919; PubMed
Central PMCID: PMCPMC4068987.
66. Alvarez-Jarreta J, Amos B, Aurrecoechea C, Bah S, Barba M, Barreto A, et al. VEuPathDB: the
eukaryotic pathogen, vector and host bioinformatics resource center in 2023. Nucleic acids research.
2024;52(D1):D808-D16. doi: 10.1093/nar/gkad1003. PubMed PMID: 37953350; PubMed Central
PMCID: PMCPMC10767879.
67. Mathur V, Kolisko M, Hehenberger E, Irwin NAT, Leander BS, Kristmundsson A, et al.
Multiple Independent Origins of Apicomplexan-Like Parasites. Curr Biol. 2019;29(17):2936-41 e5.
doi: 10.1016/j.cub.2019.07.019. PubMed PMID: 31422883.
68. Janouskovec J, Paskerova GG, Miroliubova TS, Mikhailov KV, Birley T, Aleoshin VV, et al.
Apicomplexan-like parasites are polyphyletic and widely but selectively dependent on cryptic plastid
organelles. Elife. 2019;8. doi: 10.7554/eLife.49662. PubMed PMID: 31418692; PubMed Central
PMCID: PMCPMC6733595.

69.      Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
architecture and applications. BMC bioinformatics. 2009;10:421. doi: 10.1186/1471-2105-10-421.
PubMed PMID: 20003500; PubMed Central PMCID: PMCPMC2803857.
70.      Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation
sequencing data. Bioinformatics. 2012;28(23):3150-2. doi: 10.1093/bioinformatics/bts565. PubMed
PMID: 23060610; PubMed Central PMCID: PMCPMC3516142.
71.      Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol. 2013;30(4):772-80. doi:
10.1093/molbev/mst010. PubMed PMID: 23329690; PubMed Central PMCID: PMCPMC3603318.
72.      Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. trimAl: a tool for automated alignment
trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25(15):1972-3. doi:
10.1093/bioinformatics/btp348. PubMed PMID: 19505945; PubMed Central PMCID:
PMCPMC2712344.
73.      Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood trees for large
alignments. PloS one. 2010;5(3):e9490. doi: 10.1371/journal.pone.0009490. PubMed PMID:
20224823; PubMed Central PMCID: PMCPMC2835736.
74.      Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-
TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol
Evol. 2020;37(5):1530-4. doi: 10.1093/molbev/msaa015. PubMed PMID: 32011700; PubMed Central
PMCID: PMCPMC7182206.
75.      Gu X, Fu YX, Li WH. Maximum likelihood estimation of the heterogeneity of substitution rate
among nucleotide sites. Mol Biol Evol. 1995;12(4):546-57. doi:
10.1093/oxfordjournals.molbev.a040235. PubMed PMID: 7659011.
76.      Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: fast
model selection for accurate phylogenetic estimates. Nature methods. 2017;14(6):587-9. doi:
10.1038/nmeth.4285. PubMed PMID: 28481363; PubMed Central PMCID: PMCPMC5453245.
77.      Smilkstein M, Sriwilaijaroen N, Kelly JX, Wilairat P, Riscoe M. Simple and inexpensive
fluorescence-based technique for high-throughput antimalarial drug screening. Antimicrobial agents
and chemotherapy. 2004;48(5):1803-6. PubMed PMID: 15105138; PubMed Central PMCID:
PMCPMC400546.
78.      Huang RY, Pei L, Liu Q, Chen S, Dou H, Shu G, et al. Isobologram Analysis: A Comprehensive
Review of Methodology and Current Research. Front Pharmacol. 2019;10:1222. doi:
10.3389/fphar.2019.01222. PubMed PMID: 31736746; PubMed Central PMCID: PMCPMC6830115.
79.      Fivelman QL, Adagu IS, Warhurst DC. Modified fixed-ratio isobologram method for studying
in vitro interactions between atovaquone and proguanil or dihydroartemisinin against drug-resistant
strains of Plasmodium falciparum. Antimicrobial agents and chemotherapy. 2004;48(11):4097-102.
doi: 10.1128/AAC.48.11.4097-4102.2004. PubMed PMID: 15504827; PubMed Central PMCID:
PMCPMC525430.
80.      Yang J, Yan R, Roy A, Xu D, Poisson J, Zhang Y. The I-TASSER Suite: protein structure and
function prediction. Nature methods. 2015;12(1):7-8. doi: 10.1038/nmeth.3213. PubMed PMID:
25549265; PubMed Central PMCID: PMCPMC4428668.

# Supporting information

**Fig S1: Growth curves determining IC$_{50}$ values for MMS, VE-821 and KU-55933 in the parasite**

**strains 3D7, MRA-1252 and MRC-1240.**

Each panel shows growth curves from 3-5 independent malaria SYBR green I-based fluorescence

(MSF) assays: percentage parasite growth is plotted as a function of drug concentration; error bars

are standard errors of the means.

A.  Inhibition of growth of 3D7 strain by MMS.

B.  $IC_{50}$ values from the growth curves shown in main figure 3A.

C.  Inhibition of strains MRA-1252 and MRA-1240 by the drugs VE-821, KU55933 and MMS.

**Fig S2: 3D7 parasites survive a short pulse of MMS or DHA at the trophozoite stage, but their**

**maturation is delayed and progeny number per schizont, reduced.**

A.  Images were taken 12h after trophozoites at 30-32hpi had been exposed to a 1 h pulse of

10X $IC_{50}$ MMS or DHA (2 independent experiments, '1' and '2'). Images show that at this

timepoint (i.e. 42-44hpi), cultures were dominated by a mixture of immature and mature

schizonts, without clear evidence of parasite death in the damaged cultures. Parasites were

then treated with 75nM ML10 for 10 h, to arrest mature schizonts before egress, and re-

imaged. Mature schizonts then dominated in all cultures.

B.  Parasite stages imaged in (A) were categorised as either: early rings, small rings, large rings,

early trophozoites, mid trophozoites, late trophozoites, immature schizonts or mature

schizonts (2 independent experiments, n ≥ 100 from each experiment). Differences between

the proportions of stages in control vs DHA-treated vs MMS-treated cultures were tested via

two-way ANOVA and posthoc Tukey's multiple comparisons test. P-values are shown in a

table below the graph: *, <0.05; **, <0.01; ***, <0.001. Stars of matched colours across

rows denote a significant difference between those conditions (e.g. at 44hpi, there were still

more immature schizonts after MMS treatment than in the control (pink stars); 10h later,

this difference disappeared, with more mature and fewer immature schizonts (denoted by

963            pairs of yellow and blue stars).  The only other difference was in the small numbers of rings

964            that escaped and reinvaded after prolonged arrest.

965     C.  Merozoites per schizont were counted in each of the cultures shown in (A) (n = 26-30 per

966            culture).  Numbers of merozoites substantially increased in all cultures after the long ML10-

967            arrest period, which allowed mature schizonts to accumulate even after the damage-

968            induced delay seen in (B).  A reduction in merozoite number was, however, still seen after

969            MMS damage, although this was not significant after DHA damage.

971   **Fig S3: Protein sequence alignment of *P. falciparum* PI3K, PI4K and PIKKs from *S. cerevisiae* and**

972   **human.**

973     A.  MUSCLE (3.8) alignment of catalytic domains of *P. falciparum* PI3K and PI4K against human

974            ATM, ATR, PI3K, and yeast Mec1 and Tel1. Colour key is ClustalW standard. Highlighted in

975            the yellow boxes are the catalytic loops in ATR and ATM. Percentage identity matrix is

976            shown below the alignment.

978   **Fig S4: Anti-PfPI3K antibody production**

979     A.  Model made in i-TASSER [80], threading 1500 C-terminal residues of the 2133-amino acid

980            *Pf*PI3K against human ATR. Protein coloured blue to red, N-terminal to C-terminal. Peptides

981            used as antigens are highlighted in white and circled: amino acids 1359-1374 and 2011-2025

982            of full length *Pf*PI3K.

983     B.  Controls for specificity of immune versus pre-immune antisera against *Pf*PI3K.

984            Representative images of single cells are shown. Scale bar 1 μm. Primary antibodies used

985            were rabbit antisera, pre-immune or immune, diluted 1:200 in 3% BSA.

**Fig S5: Isobolograms showing that two antibiotic drugs do not synergise with MMS**

Isobolograms plotted as fractional inhibitory concentrations (FIC) of combinations of MMS and

geneticin (G-418), or blasticidin (BSD), as determined by MSF assay. Data were processed as in Figure

4B. The type of relationship was defined as additive for blasticidin and borderline antagonistic for

geneticin, based on well-established criteria ($\sum$FIC <0.5, substantial synergism; $\sum$FIC <1, moderate

synergism; $\sum$FIC $\geq$1 and <2, additive interaction; $\sum$FIC $\geq$2 and <4, slight antagonism and $\sum$FIC >4,

marked antagonism).

**Fig S6: Growth curves determining IC$_{50}$ values for VE-821 and KU-55933 when PI3K is knocked out**

A.  Example growth curves from malaria SYBR green I-based fluorescence (MSF) assays

conducted on the iPI3K-ko line. (As in figure S1, percentage parasite growth is plotted as a

function of drug concentration; error bars are standard errors of the means.) Ring-stage

parasites of the iPI3K-ko line were treated with 100nM rapamycin to induce gene deletion,

or with DMSO vehicle for the control. MSF assays with VE-821 and KU-55933 were set up

24h later, with drug exposure over 48h, i.e. from trophozoites in cycle-1 to trophozoites in

cycle-2. At 72h, when assays were read, morphology of knockout parasites did not appear

overtly unhealthy. However, if assays were extended to 96h (after reinvasion to cycle-3), all

rapamycin-treated parasites appeared dead, whereas control parasites still appeared

healthy. This delayed-death phenotype precluded any meaningful results from 3-day (72h)

MSF assays, so exclusively 48h assays are shown here.

B.  IC$_{50}$ values from the growth curves shown A, and from 3 further replicate assays conducted

on knockout parasites over the same 2-day window (24-72h post-rapamycin). Mean IC$_{50}$s

did not differ markedly from those of the control, nor from those of the 3D7 parent line

(shown in figure S1).

1011

1012   **Fig S7: Ring stage survival assessed by flow cytometry**

1013    A ring-stage survival assay was assessed by flow cytometry as well as eRRSA as used in Figure 5.

1014    Density plot quadrant graphs show one biological replicate of an RSA conducted on 3D7. Cell counts

1015    for each quadrant are shown. Percentage survival was calculated as the proportion of live versus

1016    dead cells (SYBR+ MITO+/SYBR+ MITO-) for each condition.

1017

1018    **Fig S8: Raw growth-curve data for all isobolograms plotted in figures 4B, 5C and S5**

1019        A.   Data for isobolgrams in figure 4B, VE-821 with MMS

1020        B.   Data for isobolgrams in figure 4B, KU-55933 with MMS

1021        C.   Data for isobolgrams in figure 5C, VE-821 and KU55933 with DHA

1022        D.   Data for isobolgrams in figure S5, Blasticidin and G-418 with MMS

1023

1024    **Source data files and spreadsheets 1-7:**

1025    Files 1-6: Source data for phylogenetic tree shown in figure 2

1026    Spreadsheet 7: Source data for all other figures and supplementary figures

1027