Title:

Cell cycle checkpoint activity in the malaria parasite

Plasmodium falciparum

Running title: Cell cycle checkpoints in *Plasmodium*

Monique K. Johnson[a], Juliana Naldoni[a,b], William H. Lewis[b], Ross F. Waller[b], Catherine J. Merrick[a#]

[a] Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge, CB2 1QP, UK.

[b] Department of Biochemistry, University of Cambridge, Tennis Court Road, Cambridge, CB2 1QW,

UK.

Address correspondence to:

Department of Pathology, Cambridge University, Tennis Court Road, Cambridge, CB2 1QP, UK

Tel: (+44) 1223 333330

Email: cjm48@cam.ac.uk

15 Abstract

Plasmodium spp. have different modes of cell division from most eukaryotes. Little is known about

how these are controlled and cell-cycle checkpoints are particularly poorly characterised. However,

parasites can arrest their cell cycle when treated with the frontline antimalarial drug artemisinin,

and artemisinin-resistant parasites can modulate their cell-cycle progression, so it is important to

understand these aspects of *Plasmodium* biology. Here, we show that *P. falciparum* displays

hallmarks of an intra-S-phase checkpoint when exposed to DNA damage, including acute reduction

of DNA replication and phosphorylation of a putative damage-marker histone. Compounds that

inhibit human checkpoint kinases can inhibit this arrest of DNA replication, and synergise with DNA

damage in parasite killing. This suggests the existence of checkpoint kinase activity in *P. falciparum*,

yet these kinases have no clear homologues in *Plasmodium* genomes. Their closest homologues are

the phosphatidylinositol lipid kinases. We hypothesise that phosphatidylinositol 3-kinase – which is

reportedly up-regulated in artemisinin-resistant parasites – may moonlight in this role, and we

characterise this essential kinase for the first time via expansion microscopy. Finally, we show that

the cryptic checkpoint-kinase activity may also regulate the ring-stage survival phenotype after

artemisinin damage, which resembles a G1/S checkpoint. Hence we suggest that checkpoint kinase

inhibitors are candidates for synergy with artemisinin.

Importance

Malaria parasites infect red blood cells, wherein they replicate to produce many new parasites. This

is unusual because most cells replicate simply by copying their genome and splitting in half (called

binary fission) but malaria parasites make ~20 genome copies, then partition them simultaneously

into 20 new cells (called schizogony). Here we studied how schizogony is controlled: in particular,

are there 'checkpoints'? Checkpoints pause the cycle for repair if the genome is damaged. We found

that DNA damage *did* cause checkpoint hallmarks, yet key proteins that enforce this in other cells

are absent in malaria parasites. Furthermore, this checkpoint activity may be involved in the

response to an antimalarial drug, when parasites pause their cycle before active replication begins.

This implies that inhibiting the checkpoint could exacerbate parasite killing by such drugs. Cancer

therapies often work like this – by damaging DNA and also preventing the cancer cells from repairing

it.

Keywords

45   Malaria, *Plasmodium,* cell cycle, checkpoint, PI3K

Introduction

Plasmodium falciparum is the most important cause of human malaria, giving rise to over half a

million deaths and hundreds of millions of clinical cases per year (1). Besides being a major human

pathogen, it is also a highly unusual protozoan organism, distantly related to humans and other

model organisms, with distinctive features of cell and molecular biology. One of these is a non-

canonical cell cycle. Instead of dividing by binary fission, as in most model organisms, *Plasmodium*

parasites divide primarily by schizogony. After invading a host erythrocyte, they execute multiple,

asynchronous rounds of genome replication and nuclear division, generating multiple nuclei within

the same cytoplasm. This is followed by a relatively synchronous mass cytokinesis event,

partitioning ~20 nuclei and organelles into individual merozoites.

The complex events of schizogony are now being dissected in molecular and cellular detail

(2-5) but fundamental questions remain about how the process is regulated. Schizogony is not easily

mapped onto the cell cycle phases of G1, S, G2 and M. After invading a host cell, the parasite

undergoes a pre-replicative growth phase broadly analogous to G1. The haploid genome is then

repeatedly replicated in a continuous S-phase without once-and-only-once regulation, generating

many new haploid nuclei. Different nuclei cease to replicate at different times, and divide at

different times, so there is no well-defined G2 and no single M-phase (3).

Plasmodium replication is not controlled by conventional cyclin-dependent kinase activity

(6), nor is it apparently governed by conventional cell cycle checkpoints, which usually operate at the

G1/S transition, within S-phase, and at G2/M. Respectively, these checkpoints delay the start of S-

phase, arrest replication if DNA damage or replication stress is detected, and prevent mitosis if a

genome is incompletely replicated (7). The *Plasmodium* genus, however, encodes no clear

homologues of the checkpoint kinases conserved in most eukaryotes (animals, fungi and plants) e.g.

S. cerevisiae Mec1 and Rad53 or human ATM/ATR and Chk1/Chk2 (6). (By contrast, some parasites

in the apicomplexan phylum, like *Toxoplasma*, do retain homologues of the 'master regulators'

Mec1/ATM/ATR, which are phosphoinositide-3-kinase-like kinases (PIKKs), and checkpoints are

better characterised in *Toxoplasma* (8-10)).

Nevertheless, it seems unlikely that *Plasmodium* is entirely without checkpoints. Eukaryotic

cells (not only in metazoans, which must guard against cancer, but also protozoans) generally do

pause their cell cycles to ensure effective DNA repair. *Plasmodium* encodes most of the known DNA

repair pathways (6, 11) and generates relatively low rates of SNPs, indels and translocations during

normal replication (12).

There is some evidence for a G1/S-like checkpoint: pre-replicative rings can undergo

'temporary growth arrest (TGA)' (13) in response to nutritional (14), temperature (15) or

antimalarial-drug (16, 17) stresses. TGA is particularly important under stress caused by the

antimalarial drug artemisinin, because it can contribute to treatment failure. However, there is no

molecular evidence as-yet that conventional checkpoint effectors are involved. A G1/S checkpoint

usually involves the cyclin/CDK complexes that drive S-phase entry, influenced positively by growth

factors or negatively by damage responsive kinases like PIKKs. TGA in *Plasmodium*, by contrast, is

defined only as a 'non-specific stress-response survival mechanism' (13): induced by multiple

stresses, possibly via multiple molecular pathways.

An intra-S-phase checkpoint should arrest S-phase specifically after DNA damage or

replication stress. In *Plasmodium*, the transcriptional cascade that drives the cell cycle can slow

down after DNA damage, with DNA repair proteins and chromatin modifications being upregulated

(18), but the signals and transducers for any potential checkpoint are unknown. *Plasmodium*

generally lacks validated markers for DNA damage and for direct responses to it. Indeed, the multi-

nucleate, asynchronous nature of schizogony raises questions about how an intra-S-phase

checkpoint should or could be enforced. Replication itself is apparently regulated per-nucleus,

probably via individual accumulations of PCNA, together with non-canonical, chromatin-bound CDK-

related kinases like CRK4 (19). Nevertheless, diffusible checkpoint kinases, potentially affecting the

whole cell, may not be incompatible with schizogony – at least in response to exogenous DNA

damage.

Finally, a G2/M checkpoint may be entirely absent because some *P. falciparum* mutants

perform grossly aberrant genome segregation but still undergo cytokinesis (20). In fact, even in non-

mutants, certain phases of the lifecycle produce a high rate of zoids – this occurs particularly

gametogenesis, wherein cytokinesis proceeds even if genome replication fails entirely (21, 22).

Here, we set out to investigate checkpoint responses to DNA damage in *Plasmodium,* using

newly developed molecular and cellular tools. We show that the hallmarks of an S-phase checkpoint

do occur in *P. falciparum*, despite the apparent absence of canonical PIKKs.

Results

Evidence for DNA-damage-responsive checkpoint activity in *P. falciparum*

We used two complementary tools to investigate the response to DNA damage during S-phase in *P.*

falciparum, focussing at the levels of DNA and protein. Firstly, we labelled nascent DNA in

replicating parasites with a quantifiable modified nucleotide. This tested whether DNA replication

would acutely arrest after damage: a hallmark of the intra-S-phase checkpoint (23, 24). Secondly,

we measured phosphorylation of histone H2A, which was recently identified as being analogous to

histone H2AX phosphorylation in mammalian cells (25). Canonically, H2AX is phosphorylated at

chromatin sites of DNA damage (26, 27). H2AX is absent in *Plasmodium*, but the non-variant histone

H2A can nevertheless be phosphorylated (25). Importantly, this indicates an *in trans*, kinase-driven,

response that is genuinely 'checkpoint-like'.

To induce DNA damage, we used the well-characterised alkylating agent methyl methane

sulphonate (MMS), which alkylates DNA bases, particularly guanine and adenine (28). In mammalian

cells, it induces an intra-S-phase checkpoint (23). In *P. falciparum* parasites, it leads to DNA

breakage, as measured by comet assays (29) and concomitantly delays the cell cycle (18), but its

effect as a checkpoint inducer has not been explicitly addressed. In the *P. falciparum* 3D7 strain,

MMS had a 48h IC_{50} of ~96µM (Fig S1A). Having established this, we performed highly acute assays

over just 1-2h, using 10x or 20x IC_{50}, to minimise non-specific or secondary effects. We confirmed

that parasites could substantially recover from this level of MMS exposure within one cell cycle,

although their maturation was delayed, as reported previously (18), and there was also a reduction

in merozoite numbers in mature schizonts (Fig S2).

DNA replication was acutely reduced in the 30 minutes following a 30-minute exposure to

MMS at 10x or 20x IC_{50}. Some reduction occurred throughout S-phase (~28-36 hours post invasion,

hpi) but it was most pronounced when DNA was damaged in early trophozoites at ~32hpi (Fig 1A).

Within a similar timeframe, histone H2A phosphorylation was moderately elevated (Fig 1B). We

recently reported that several DNA damaging agents, including MMS, could cause H2A

phosphorylation (30), but that the upregulation in trophozoite stages was modest – usually less than

2-fold over background levels – making it difficult to measure consistently. This was equally true

when MMS was used to induce H2A phosphorylation here (Fig 1B). (Previously published work

measured H2A phosphorylation in pre-replicative rings exposed to ionising radiation (25), whereas

we measured it in actively replicating trophozoites, wherein the background signal may be higher.)

Therefore, we chose the replication-reduction assay as a more robust readout for checkpoint activity

in subsequent experiments.

Absence of clear checkpoint mediators in *P. falciparum*

When the intra-S-phase checkpoint is triggered in other eukaryotes, both histone phosphorylation

and arrested DNA replication are transduced through PIKKs (24, 26). Since Figure 1 shows that these

phenotypes also appear in *P. falciparum,* a transducing kinase should be present. However,

orthologues of ATM/ATR are not found in the *P. falciparum* genome (6). To test for the presence or

loss of such orthologues more thoroughly, we performed an extensive phylogenetic analysis of PIKK

homologues throughout apicomplexans and other major eukaryotic lineages (Fig 2). In this analysis,

the lipid kinases PI3K and PI4K formed an outgroup relative to all ATM/ATR-like proteins. In

eukaryotes, PI3/4Ks are generally the closest relatives of PIKKs (31) and we did not seek to re-

establish this (e.g. by surveying apicomplexan genomes for any other kinases resembling PIKKs), we

simply sought to establish patterns of PIKK presence/loss in apicomplexans.

ATR homologues were found in most apicomplexans, and some – like *T. gondii* – clearly

retained homologues of both ATM and ATR. Other apicomplexans, including *Cryptosporidium*,

Babesia and *Theileria*, had independently lost the ATM homologue, indicating a propensity for loss

of PIKKs in Apicomplexa. *Plasmodium* spp., however, had lost both ATM and ATR, retaining no

apparent PIKKs at all. The closest PIKK homologues in any *Plasmodium* genome were the lipid

kinases PI3K and PI4K (Fig 2, Fig S3), which were also present throughout apicomplexans (32).

(*Theileria*, unusually, was found to have lost its PI3K homologue, possibly due to a lifecycle

conducted free in the cytoplasm of leukocytes, which may supply this activity.

Notably, the closest *Plasmodium* PIKK homologue, PI3K, is reportedly overexpressed in

artemisinin resistant parasites (33), which can alter their cell cycle dynamics. Furthermore, a recent

analysis of the steady state locations of mature schizont-stage *P. falciparum* proteins by spatial

proteomics assigned PI3K as a nucleus-located protein, whereas PI4K was assigned to cytoplasmic

transport vesicles (Fig S4A) (34). This is consistent with PI3K having access to nuclear substrates such

as histone H2A for phosphorylation. Accordingly we hypothesised that PI3K might moonlight as a

checkpoint kinase in *Plasmodium.*

*Pf*PI3K is the only phosphoinositide 3-kinase of the six phosphoinositide kinases in

Plasmodium (32): it is a well-validated, essential lipid kinase (35), with roles in nutrient uptake and

vesicle trafficking (36, 37). However, its moonlighting as a protein kinase would not be

unprecedented because other PI3Ks can to act on proteins as well as lipids. For example, humans

encode a large family of PI3Ks, and their type-1 PI3Ks can phosphorylate protein substrates like

cytokine receptors, as well as auto-phosphorylating themselves (38, 39). The sole *Plasmodium*

enzyme is a type-3 PI3K, and thus far these are proven only to phosphorylate lipids (35).

We sought orthogonal validation of the spatial proteomics via immunofluorescence after

raising peptide antibodies to *Pf*PI3K (Fig 3A, Fig S4B, C). The protein appeared in puncta throughout

the cytoplasm, consistent with previous work (37) (Fig 3B). This highly dispersed location could

obscure any minor nuclear signal, and we did not detect substantial re-location to nuclei upon DNA

damage (Fig 3C). We then used expansion microscopy to better resolve the location of *Pf*PI3K,

observing that puncta grew more abundant as cells matured, and were densest in an area free of

nuclei that is probably the food vacuole – a known hub for vesicle trafficking (Fig 3D). Some *Pf*PI3K

was also detected inside nuclei in all cells imaged (Fig 3E, Fig S5). Of note, we recently showed that

the ATM and ATR homologues in *Toxoplasma* are likewise dispersed throughout the cell and do not

substantially relocate to the nucleus after DNA damage, but they are nevertheless able to mediate

checkpoint phenotypes such as histone phosphorylation (40).

We attempted to use the same antibodies to immunoprecipitate native *Pf*PI3K with any

potential protein substrates, but only a single *Pf*PI3K peptide was detected across several attempts.

It is probably highly unstable *in vitro*, or lacking suitable tryptic digest sites, because it was likewise

completely undetected, despite its very large size, in two dedicated proteomic studies of the nuclear

versus cytoplasmic proteomes in *P. falciparum* (41, 42); and it was only sparsely detected in spatial

proteomics (34). Further validation of an intra-nuclear role for *Pf*PI3K would probably require

methods focussed less on the kinase itself and more on its targets – such as BioID for proximity-

labelling of protein partners (43).

The mediator of DNA-damage checkpoint responses in *Plasmodium* is sensitive to PIKK inhibitors

Assuming that a divergent PIKK-like kinase does exist in *P. falciparum*, we attempted to inhibit it with PIKK inhibitors. We chose two compounds with validated specificity for human ATM and ATR respectively (but not for human PI3Ks), KU-55933 and VE-821 (44, 45). If an ATM-like or ATR-like activity existed in *P. falciparum* – either in *Pf*PI3K or elsewhere – these compounds might specifically inhibit it.

We first determined their 48h IC_{50} values on *P. falciparum* (3D7 strain): KU-55933 had an IC_{50} in the micromolar range and VE-821, in the nanomolar range (Fig 4A, Fig S1B). Both inhibitors clearly synergised with MMS in a 48h isobologram analysis, particularly strongly in the case of VE-821, suggesting activity in the same pathway (Fig 4B). Control compounds that kill parasites but are not known to cause DNA damage – the antibiotics blasticidin and geneticin – did not synergise with MMS (Fig S6). Furthermore, when parasites were exposed to high-dose MMS (10x IC_{50}, as in Figure 1) simultaneously with KU-55933 or VE-821, the inhibitors ablated the acute reduction in DNA replication caused by MMS, whereas co-exposure to the control antibiotic geneticin did not (Fig 4C). In this short 1h assay, we used the PIKK inhibitors at only their 48h IC_{50} levels, to minimise any negative effects from the inhibitors alone. Indeed, within the timeframe of the assay the inhibitors alone did not significantly affect DNA replication (Fig 4C). Overall, these data strongly supported the existence of a cryptic PIKK activity, responsible for a genuine intra-S-phase checkpoint via *in trans* kinase activity.

The antimalarial drug dihydroartemisinin induces DNA replication arrest in *P. falciparum*

MMS is a useful, well-characterised DNA-damaging agent, but it is not an antimalarial drug. We were particularly interested in the potential DNA-damaging activity of antimalarial drugs, because if

parasites do make checkpoint responses to such drugs, then their antimalarial activity might be

potentiated by inhibiting checkpoint kinases. Artemisinin is such a drug: it has non-specific alkylating

activity that may affect both proteins and DNA (46, 47). There are similarities between the

parasite's response to dihydroartemisinin (DHA, the active metabolite of artemisinin) and MMS in

terms of DNA breakage (46, 48), delay in the parasite's transcriptional cascade, and upregulation of

DNA repair genes (18). Indeed, we found that DHA induced H2A phosphorylation, at all stages of the

cycle but particularly in pre-replicative ring stages (~3-fold upregulation after 4h exposure to IC_{50}

DHA, (Fig 5A)). This confirmed that DHA does indeed cause damage that activates a putative histone

kinase, and furthermore that it can cause this damage prior to S-phase.

Using the replication-reduction assay, we saw that in replicating parasites, DHA caused an

acute reduction in nascent DNA replication, similar to MMS (Fig 5B). However, unlike MMS, this was

not ablated by the PIKK inhibitors KU-55933 and VE-821. Consistent with this difference, the

inhibitors did not synergise with DHA in isobologram analysis (Fig 5C). This suggested that although

both DHA and MMS may damage DNA, they differ in either the type of damage or the cellular

response thereto in terms of inducing an intra-S-phase checkpoint.

Ring-stage survival after artemisinin damage is affected by PIKK

inhibitors

Cell-cycle checkpoint activity in *P. falciparum* may not be limited to an intra-S-phase checkpoint:

most eukaryotic cells also operate a G1/S checkpoint, delaying S-phase entry if cells are damaged,

starved or stressed. This may be analogous to the 'TGA' arrest or 'dormancy' that occurs in ring-

stage parasites after artemisinin treatment (13). We therefore set out to establish whether ring-

stage dormancy induced by DHA actually represents a classical, G1/S checkpoint, responsive to the

DNA damage caused in ring stages by DHA (as shown in figure 5A), and mediated by a potential PIKK-

like kinase.

We conducted the ring-stage survival assay (RSA) on parasites treated with DHA in the

presence or absence of PIKK inhibitors. In this assay, pre-replicative early-ring-stage parasites are

damaged with a pulse of DHA: most of them die, but some successfully arrest and thence survive.

The more parasites are able to do this, the higher the subsequent recovery of viable parasites.   We

reasoned that if successful arrest is due to PIKK-mediated checkpoint activity, then PIKK inhibitors

should reduce ring-stage survival. These experiments were conducted in the 3D7 strain and also in

the Cambodian strain MRA-1252, derived from a naturally-arising DHA-resistant strain reverted back

to a sensitive phenotype by specific correction of the resistance mutation in the Kelch13 gene (49):

its genetic background is otherwise that of a S.E. Asian DHA-resistant line.

Both 3D7 and MRA-1252 are highly sensitive to artemisinin, so we used the 'extended

recovery ring-stage survival assay' (eRRSA), which detects recovered parasites very sensitively after

72h via quantitative PCR (50). Figure 6A shows that PIKK inhibitors, particularly KU-55933, did

exacerbate the killing of ring-stage parasites by DHA. Differences were subtle and parallel assays

reading the RSA with the less sensitive flow cytometric assessment of live/dead cells likewise

showed marginal differences (Fig S7). However, in triplicate eRRSAs on two independent lines, 3D7

and MRA-1252, survival trended consistently downwards when DHA was combined with either

inhibitor – so the effect, although not strong, was highly reproducible. As above, we minimised any

negative effects of the inhibitors themselves by using them at a low level: only half of the 48h IC_{50}.

We confirmed that this exposure alone for 6h did not affect parasite survival after 72h (Fig 6B).

These RSAs did not show conclusively that 'TGA' after artemisinin damage represents a true

G1/S checkpoint but they did suggest that such a checkpoint, mediated by a cryptic PIKK, may

contribute to ring-stage survival. Hence, the cryptic PIKK must be active throughout the cell cycle,

not only in actively replicating trophozoites.

When the same assay was conducted on DHA-resistant parasites (the parent strain of MRA-

1252, called MRA-1240) the result differed. As expected, DHA survival was much better in MRA-

1240, and PIKK inhibitors had no negative effect on this survival (Fig 6C). There was even a trend

towards better growth and DHA survival when MRA-1240 was exposed to PIKK inhibitors (Fig 6D).

Hence, when a Kelch13 mutation was present, conferring DHA resistance, any contribution of

checkpoint activity to parasite survival was apparently ablated.

270 Artemisinin-resistant parasites are cross-resistant to MMS and VE-

271 821, but not to KU-55933

We hypothesised that artemisinin-resistant parasites like MRA-1240 might not require 'G1/S'

checkpoint activity to survive DHA damage because they are more efficient at DNA repair. Several

mutations in DNA repair genes have been identified in resistant strains (47), although the pathway(s)

leading to improved DNA repair are not fully characterised. We checked whether the DHA-resistant

line MRA-1240 was cross-resistant to MMS, and its IC$_{50}$ was indeed significantly higher than that of

the matched sensitive line MRA-1252 (and also higher than that of 3D7) (Fig 6E, left panel; Fig S1C).

Since the point mutation in Kelch13 is the only difference between the two Cambodian MRA lines, a

genetic background of enhanced DNA repair cannot fully account for this.

Another explanation could be that MRA-1240 has hyper-active checkpoint activity due to

Kelch13-associated over-expression of PI3K (33), and that this stimulates hyperactive DNA repair.

This explanation *would* be consistent with the Kelch13 mutant and revertant having different

sensitivities to MMS. If so, the Kelch13 mutant state would also be expected to correlate with

reduced sensitivity to PIKK inhibitors. Indeed, MRA-1240 was much less sensitive to the VE-821

inhibitor (Fig 6E, middle panel; Fig S1C) – so much so that it was difficult to measure a consistent IC$_{50}$

via MSF assay. Curiously, however, the same line was not less sensitive to KU-55933 (Fig 6E, right

panel; Fig S1C). This was the first instance in which VE-821 and KU-55933 had shown a clear

difference in their effects on *P. falciparum.* It may suggest that in the MRA-1240/1252 genetic

background (perhaps unlike 3D7), more than one checkpoint protein is required, and that one of the

inhibitors has a broader inhibitory spectrum than the other. For example, *Pf*PI3K may indeed

moonlight as a checkpoint kinase, uniquely unregulated after Kelch13 mutation and uniquely

inhibited by VE-821, whereas *Pf*PI4K and/or other 'off-target' kinases may also be needed when

broader-spectrum damage induces more complex checkpoint responses, and KU-55933 may inhibit

this broader kinase spectrum.

295   Validating these speculations would require challenging experiments, such as biochemical

assays on recombinant PI3/4K proteins (which are ~250kDa in size), or robust conditional

manipulation of the two large, essential genes encoding *PfPI3K* and *PfPI4K* in parallel. We did make

several attempts to tag the *PfPI3K* gene for conditional knockdown, but the enzyme's active site is

encoded at the extreme 3' end and may become mis-folded after 3' tagging, because all our

attempts failed. Similar efforts by other groups to tag or manipulate this gene were likewise

unsuccessful or gave no phenotype (51, 52), while a recently-published mutant that irreversibly

deletes the 3' end of the *PfPI3K* gene was lethal (53). The very large size of the gene (6.6kb) also

precludes its overexpression or dominant-negative overexpression, as well as recombinant protein

production for enzyme-activity assays. Overall, these large and essential genes proved near-

intractable to molecular genetics. Thus, their putative PIKK-like activity was not conclusively

evidenced or disproved, but the dual nuclear/cytoplasmic location of *Pf*PI3K does suggest that it

could moonlight on nuclear substrates.

308 Discussion

This work provides the first evidence that *Plasmodium* parasites during blood-stage schizogony can

enforce an intra-S-phase checkpoint. The checkpoint is evidenced by an acute reduction in the rate

of DNA replication and increased phosphorylation of histone H2A in a position analogous to the

canonical DNA damage marker H2AX. *P. falciparum* was previously shown to slow its cell-cycle

progression after DNA damage (18) but it was not clear if this occurred through cell-cycle checkpoint

activity (partly because the *Plasmodium* system lacked validated DNA damage markers, signals or

transducers thereof). Markers still remain limited – there are, for example, no cyclin markers for

cell-cycle phases, and the H2A phosphorylation that was reported as a DNA damage marker in ring

stages (25) is, in our hands, a consistent but low-dynamic-range marker in trophozoite stages,

probably because background levels of H2A-P are higher in actively-replicating parasites. However,

the ability to measure nascent DNA replication in *Plasmodium* (54) does now allow intra-S-phase

checkpoint activity to be measured. Thus, we showed that DNA replication was acutely arrested by

two distinct alkylating agents, MMS and the antimalarial drug DHA.

The transducer(s) of checkpoint activity in *Plasmodium* remain cryptic. *Plasmodium* spp.

lack identifiable orthologues of PIKK kinases, in contrast to many apicomplexans including *T. gondii*

(40). Nevertheless, such activity is apparently present, because the reduction in replication caused

by MMS was ablated by well-validated inhibitors of human PIKKs. Hence it must be induced *in trans*

by a kinase-mediated signalling cascade (also consistent with H2A phosphorylation), rather than

occurring simply because alkylated DNA bases physically block the replisome. Interestingly, DHA

differed from MMS in this respect: it also induced H2A phosphorylation, and it also affected DNA

replication, but PIKK inhibitors did not significantly ameliorate this. Possibly the acute effect of DHA

– which is a much larger molecule than MMS – *is* primarily to block replisomes via bulky DNA

adducts (a documented effect of alkylating agents in human cells (23)). If such adducts cannot be

quickly bypassed whether a checkpoint kinase is active or not, then PIKK inhibitors might not

ameliorate the acute reduction in DNA replication.

Other, more complex, explanations are also possible for the difference in response to MMS

versus DHA. For example, slow DNA replication can, in general, stimulate compensatory activation

of extra replication origins (55), but when damage to DNA is sensed, checkpoint kinases act *in trans*

to suppress such origin firing (23). If checkpoint kinases are inhibited, then extra origins may fire

regardless, hence MMS-damaged cells may keep replicating and encountering more damage, so

genome replication ultimately fails to complete. This would explain how MMS and PIKK inhibitors

ultimately synergise in cell killing. Perhaps the broader-spectrum damage that DHA causes to

proteins as well as DNA (46), and/or the oxidative stress that it generates (48) means that

compensatory origin firing does not happen, whether a checkpoint is activated or not. Hence PIKK

inhibitors would not synergise with DHA in cell killing. To evidence these rather complex

hypotheses, precision measurement of DNA replication dynamics at single-molecule resolution after

different types of damage will be needed. This has been achieved in human cells (23), and is now in

development for *Plasmodium*.

Turning to the possibility of a pre-replicative G1/S checkpoint, the evidence for this being

mediated by a PIKK-like kinase is tantalising albeit not conclusive. In the RSA, which measures DHA

killing of pre-replicative rings, PIKK inhibitors exacerbated parasite killing. The effect was modest, so

if a classical (kinase-enforced) G1/S checkpoint does exist, it is probably not the sole defining factor

in survival of artemisinin-induced dormancy.   Nevertheless, the trend was consistent in triplicate

assays across two genetically-distinct DHA-sensitive strains, 3D7 and MRA-1252. Checkpoint kinase

activity could therefore help to protect ring-stage cells, perhaps by detecting DNA damage,

stimulating repair, and preventing S-phase entry until repair is complete. By contrast, a DHA-

resistant strain showed no effect of PIKK inhibitors. This may be because it can already suppress

DHA activation in early rings and thus protect itself from DNA damage; or alternatively, it may still

sustain some damage, but be rescued by hyperactive DNA repair (47). Consistent with this latter

hypothesis, it was recently shown that when *Pf*PI3K is inducibly knocked out in a DHA-resistant

Kelch-13 mutant background, the parasites become less DHA resistant – perhaps because they can

no longer resist or repair DHA-related DNA damage when the mediating kinase (*Pf*PI3K) is lost (53).

If there is indeed hyperactive DNA repair in the MRA-1240 line, it cannot be conferred per se

by a DHA-resistant genetic background because the matched strain MRA-1252 was not cross-

resistant to MMS, whereas MRA-1240 was. However, it could be because the Kelch13 mutation,

uniquely present in MRA-1240, confers hyper-checkpoint activity, thus stimulating hyperactive DNA

repair. In any case, these data raise the intriguing possibility that artemisinin drugs could synergise

with checkpoint inhibitors in killing ring-stage parasites; the caveat being that this might only work

well in sensitive, not resistant, strains.

It is an attractive possibility that *Pf*PI3K is the cryptic checkpoint kinase, particularly if

Kelch13 mutant strains do contain more *Pf*PI3K (33). However, we were not yet able to prove or

disprove this, owing to the extreme difficulty of manipulating the large, essential *PfPI3K* gene.

Broad-spectrum inhibitors such as wortmannin were used in previous work (37), but these are slow-

killing and unable to separate lipid-kinase from putative protein-kinase activity. We used PIKK

inhibitors instead, hoping to isolate a putative protein kinase activity. The fact that our PIKK

inhibitors did not synergise with DHA in MSF assays, whereas specific inhibitors of distinct human

PI3Ks (i.e. lipid kinases) *have* been reported to do so (56), may suggest that this succeeded, and that

VE-821 and KU-55933 do target a distinct activity in *Plasmodium*. Of course, they may affect

irrelevant off-targets, i.e. multiple or unexpected kinases, or even non-kinases. As an example, a

repurposed human-kinase inhibitor was recently deployed to target the PK6 kinase in *Plasmodium*,

but it actually targeted haem detoxification as well (57). More likely, however, is that these

inhibitors genuinely inhibit a *Plasmodium* checkpoint kinase – which may reside in *Pf*PI3K, *Pf*PI4K, or

another radically divergent protein. *Plasmodium* contains many kinases whose roles are yet-

unknown (58), and the *Plasmodium* cell cycle is driven by markedly divergent kinases such as CRK4

(59), so it is entirely possible that *Plasmodium* checkpoints are enforced by novel kinase(s) as well.

In conclusion, we present new evidence for measurable cell-cycle checkpoint activity in

Plasmodium. We also raise the considerable challenge of unambiguously identifying its mediators in

this divergent eukaryotic parasite.

Methods

Full Methods as follows are provided as Supplementary material:

• Parasite culture & parasite strains

• Database sampling and phylogenetic analysis

• BrdU Enzyme-linked immunosorbent assay (ELISA)

• Immunofluorescence

• Immunofluorescence with expansion microscopy

• Malaria SYBR Green-1 Fluorescence (MSF) assay

• Isobolograms

• Ring-stage Survival Assay (RSA)

• RSA analysis: qPCR based & flow cytometry-based

• Antibody production

• Western blotting

Acknowledgements

We thank Dr Holly Matthews for early work on this project, Dr Ibtissam Jabre for help with H2A-P

blots, Prof Kai Wengelnik for sharing a PI3K inducible-knockout line, and Profs Marcus Lee and Paul

Roepe for helpful early discussions.

This work was funded by the European Research Council (ERC) under the European Union's Horizon

2020 research and innovation programme (ERC-2016-COG 725126 to CJM); a PhD studentship from

the University of Cambridge Department of Pathology to MKJ; a Biotechnology and Biological

Sciences Research Council (BBSRC) grant APP54272 to RFW; and a Gordon and Betty Moore

Foundation Investigator Award to RFW (https://doi.org/10.37807/GBMF9194). The funders had no

role in study design, data collection, interpretation or the decision to submit the work for

publication.

References

1.   WHO. 2023. World Malaria Report 2023.

2.   Klaus S, Binder P, Kim J, Machado M, Funaya C, Schaaf V, Klaschka D, Kudulyte A, Cyrklaff M,
Laketa V, Hofer T, Guizetti J, Becker NB, Frischknecht F, Schwarz US, Ganter M. 2022.
Asynchronous nuclear cycles in multinucleated Plasmodium falciparum facilitate rapid
proliferation. Sci Adv 8:eabj5362.

3.   McDonald J, Merrick CJ. 2022. DNA replication dynamics during erythrocytic schizogony in
the malaria parasites Plasmodium falciparum and Plasmodium knowlesi. Plos Pathogens 18:
e1010595.

4.   Simon CS, Sturmer VS, Guizetti J. 2021. How Many Is Enough? - Challenges of Multinucleated
Cell Division in Malaria Parasites. Front Cell Infect Microbiol 11:658616.

5.   Liffner B, Cepeda Diaz AK, Blauwkamp J, Anaguano D, Frolich S, Muralidharan V, Wilson DW,
Dvorin JD, Absalon S. 2023. Atlas of Plasmodium falciparum intraerythrocytic development
using expansion microscopy. Elife 12.

6.   Matthews H, Duffy CW, Merrick CJ. 2018. Checks and balances? DNA replication and the cell
cycle in Plasmodium. Parasit Vectors 11:216.

7.   Zhou BB, Elledge SJ. 2000. The DNA damage response: putting checkpoints in perspective.
Nature 408:433-9.

8.   Hawkins LM, Naumov AV, Batra M, Wang C, Chaput D, Suvorova ES. 2021. Novel CRK-Cyclin
Complex Controls Spindle Assembly Checkpoint in Toxoplasma Endodyogeny. mBio
13:e0356121.

9.   Naumov AV, Wang C, Chaput D, Ting LM, Alvarez CA, Keller T, Ramadan A, White MW, Kim K,
Suvorova ES. 2022. Restriction Checkpoint Controls Bradyzoite Development in Toxoplasma
gondii. Microbiol Spectr 10:e0070222.

10.   Alvarez CA, Suvorova ES. 2017. Checkpoints of apicomplexan cell division identified in
Toxoplasma gondii. PLoS Pathog 13:e1006483.

11.   Lee AH, Symington LS, Fidock DA. 2014. DNA repair mechanisms and their biological roles in
the malaria parasite Plasmodium falciparum. Microbiol Mol Biol Rev 78:469-86.

12.   Claessens A, Hamilton WL, Kekre M, Otto TD, Faizullabhoy A, Rayner JC, Kwiatkowski D.
2014. Generation of antigenic diversity in Plasmodium falciparum by structured
rearrangement of Var genes during mitosis. PLoS Genet 10:e1004812.

13.   Platon L, Menard D. 2024. Plasmodium falciparum ring-stage plasticity and drug resistance.
Trends Parasitol 40:118-130.

14.   Babbitt SE, Altenhofen L, Cobbold SA, Istvan ES, Fennell C, Doerig C, Llinas M, Goldberg DE.
2012. Plasmodium falciparum responds to amino acid starvation by entering into a
hibernatory state. Proc Natl Acad Sci U S A 109:E3278-87.

15.   Yuan L, Hao M, Wu L, Zhao Z, Rosenthal BM, Li X, He Y, Sun L, Feng G, Xiang Z, Cui L, Yang Z.
2014. Refrigeration provides a simple means to synchronize in vitro cultures of Plasmodium
falciparum. Exp Parasitol 140:18-23.
16.   Bohorquez EB, Juliano JJ, Kim HS, Meshnick SR. 2013. Mefloquine exposure induces cell cycle
delay and reveals stage-specific expression of the pfmdr1 gene. Antimicrob Agents
Chemother 57:833-9.
17.   Witkowski B, Lelievre J, Barragan MJ, Laurent V, Su XZ, Berry A, Benoit-Vical F. 2010.
Increased tolerance to artemisinin in Plasmodium falciparum is mediated by a quiescence
mechanism. Antimicrob Agents Chemother 54:1872-7.
18.   Gupta DK, Patra AT, Zhu L, Gupta AP, Bozdech Z. 2016. DNA damage regulation and its role
in drug-related phenotypes in the malaria parasites. Sci Rep 6:23603.
19.   Voss Y, Klaus S, Guizetti J, Ganter M. 2023. Plasmodium schizogony, a chronology of the
parasite's cell cycle in the blood stage. PLoS Pathog 19:e1011157.
20.   Absalon S, Dvorin JD. 2021. Depletion of the mini-chromosome maintenance complex
binding protein allows the progression of cytokinesis despite abnormal karyokinesis during
the asexual development of Plasmodium falciparum. Cell Microbiol 23:e13284.
21.   Janse CJ, van der Klooster PF, van der Kaay HJ, van der Ploeg M, Overdulve JP. 1986. DNA
synthesis in Plasmodium berghei during asexual and sexual development. Mol Biochem
Parasitol 20:173-82.
22.   Matthews H, McDonald J, Totanes FIG, Merrick CJ. 2022. Dynamics of DNA replication during
male gametogenesis in the malaria parasite Plasmodium falciparum. Cellular Microbiology
doi:doi.org/10.1101/2021.12.18.473304
23.   Merrick CJ, Jackson D, Diffley JF. 2004. Visualization of altered replication dynamics after
DNA damage in human cells. J Biol Chem 279:20067-75.
24.   Santocanale C, Diffley JF. 1998. A Mec1- and Rad53-dependent checkpoint controls late-
firing origins of DNA replication. Nature 395:615-8.
25.   Goyal M, Heinberg A, Mitesser V, Kandelis-Shalev S, Singh BK, Dzikowski R. 2021.
Phosphorylation of the Canonical Histone H2A Marks Foci of Damaged DNA in Malaria
Parasites. mSphere 6.
26.   Helt CE, Cliby WA, Keng PC, Bambara RA, O'Reilly MA. 2005. Ataxia telangiectasia mutated
(ATM) and ATM and Rad3-related protein exhibit selective target specificities in response to
different forms of DNA damage. J Biol Chem 280:1186-92.
27.   Rogakou EP, Pilch DR, Orr AH, Ivanova VS, Bonner WM. 1998. DNA double-stranded breaks
induce histone H2AX phosphorylation on serine 139. J Biol Chem 273:5858-68.
28.   Beranek DT. 1990. Distribution of methyl and ethyl adducts following alkylation with
monofunctional alkylating agents. Mutat Res 231:11-30.
29.   Gopalakrishnan AM, Kumar N. 2013. Opposing roles for two molecular forms of replication
protein A in Rad51-Rad54-mediated DNA recombination in Plasmodium falciparum. mBio
4:e00252-13.
30.   Craven HM, Nettesheim G, Cicuta P, Blagborough AM, Merrick CJ. 2023. Effects of the G-
quadruplex-binding drugs quarfloxin and CX-5461 on the malaria parasite Plasmodium
falciparum. Int J Parasitol Drugs Drug Resist 23:106-119.
31.   Smith GCM, Jackson SP. 2010. Chapter 77 - The PIKK Family of Protein Kinases. Handbook of
Cell Signaling (Second Edition), doi:10.1016/B978-0-12-374145-5.00077-2:575-580.
32.   Hassett MR, Roepe PD. 2018. PIK-ing New Malaria Chemotherapy. Trends Parasitol 34:925-
927.
33.   Mbengue A, Bhattacharjee S, Pandharkar T, Liu H, Estiu G, Stahelin RV, Rizk SS, Njimoh DL,
Ryan Y, Chotivanich K, Nguon C, Ghorbal M, Lopez-Rubio JJ, Pfrender M, Emrich S, Mohandas
N, Dondorp AM, Wiest O, Haldar K. 2015. A molecular mechanism of artemisinin resistance
in Plasmodium falciparum malaria. Nature 520:683-7.

34. Chisholm SA, Flores V, Kemp A, Breckels MA, Koreny L, Dos Santos Pacheco N, Barylyuk K,
Kuroshchenkova A, Lilley KS, Rayner JC, Waller RF. 2025. The spatial proteome of the
Plasmodium falciparum schizont illuminates the composition and evolutionary trajectories
of its organelles. BioRxiv doi:10.1101/2025.11.27.690946

35. Hassett MR, Sternberg AR, Riegel BE, Thomas CJ, Roepe PD. 2017. Heterologous Expression,
Purification, and Functional Analysis of Plasmodium falciparum Phosphatidylinositol 3'-
Kinase. Biochemistry 56:4335-4345.

36. Tawk L, Chicanne G, Dubremetz JF, Richard V, Payrastre B, Vial HJ, Roy C, Wengelnik K. 2010.
Phosphatidylinositol 3-phosphate, an essential lipid in Plasmodium, localizes to the food
vacuole membrane and the apicoplast. Eukaryot Cell 9:1519-30.

37. Vaid A, Ranjan R, Smythe WA, Hoppe HC, Sharma P. 2010. PfPI3K, a phosphatidylinositol-3
kinase from Plasmodium falciparum, is exported to the host erythrocyte and is involved in
hemoglobin trafficking. Blood 115:2500-7.

38. Thomas D, Powell JA, Green BD, Barry EF, Ma Y, Woodcock J, Fitter S, Zannettino AC, Pitson
SM, Hughes TP, Lopez AF, Shepherd PR, Wei AH, Ekert PG, Guthridge MA. 2013. Protein
kinase activity of phosphoinositide 3-kinase regulates cytokine-dependent cell survival. PLoS
Biol 11:e1001515.

39. Foukas LC, Shepherd PR. 2004. Phosphoinositide 3-kinase: the protein kinase that time
forgot. Biochem Soc Trans 32:330-1.

40. Johnson MK, Chelaghma S, Koreny L, Waller RF, Merrick CJ. 2024. Characterisation of cell
cycle checkpoint kinases in Toxoplasma gondii. BioRxiv
doi:doi.org/10.1101/2024.09.10.612042;:doi.org/10.1101/2024.09.10.612042.

41. Oehring SC, Woodcroft BJ, Moes S, Wetzel J, Dietz O, Pulfer A, Dekiwadia C, Maeser P, Flueck
C, Witmer K, Brancucci NM, Niederwieser I, Jenoe P, Ralph SA, Voss TS. 2012. Organellar
proteomics reveals hundreds of novel nuclear proteins in the malaria parasite Plasmodium
falciparum. Genome Biol 13:R108.

42. Briquet S, Ourimi A, Pionneau C, Bernardes J, Carbone A, Chardonnet S, Vaquero C. 2018.
Identification of Plasmodium falciparum nuclear proteins by mass spectrometry and
proposed protein annotation. PLoS One 13:e0205596.

43. Birnbaum J, Scharf S, Schmidt S, Jonscher E, Hoeijmakers WAM, Flemming S, Toenhake CG,
Schmitt M, Sabitzki R, Bergmann B, Frohlke U, Mesen-Ramirez P, Blancke Soares A,
Herrmann H, Bartfai R, Spielmann T. 2020. A Kelch13-defined endocytosis pathway mediates
artemisinin resistance in malaria parasites. Science 367:51-59.

44. Reaper PM, Griffiths MR, Long JM, Charrier JD, Maccormick S, Charlton PA, Golec JM, Pollard
JR. 2011. Selective killing of ATM- or p53-deficient cancer cells through inhibition of ATR. Nat
Chem Biol 7:428-30.

45. Hickson I, Zhao Y, Richardson CJ, Green SJ, Martin NM, Orr AI, Reaper PM, Jackson SP, Curtin
NJ, Smith GC. 2004. Identification and characterization of a novel and specific inhibitor of the
ataxia-telangiectasia mutated kinase ATM. Cancer Res 64:9152-9.

46. Ismail HM, Barton V, Phanchana M, Charoensutthivarakul S, Wong MH, Hemingway J, Biagini
GA, O'Neill PM, Ward SA. 2016. Artemisinin activity-based probes identify multiple
molecular targets within the asexual stage of the malaria parasites Plasmodium falciparum
3D7. Proc Natl Acad Sci U S A 113:2080-5.

47. Xiong A, Prakash P, Gao X, Chew M, Tay IJJ, Woodrow CJ, Engelward BP, Han J, Preiser PR.
2020. K13-Mediated Reduced Susceptibility to Artemisinin in Plasmodium falciparum Is
Overlaid on a Trait of Enhanced DNA Damage Repair. Cell Rep 32:107996.

48. Gopalakrishnan AM, Kumar N. 2015. Antimalarial action of artesunate involves DNA damage
mediated by reactive oxygen species. Antimicrob Agents Chemother 59:317-25.

49. Straimer J, Gnadig NF, Witkowski B, Amaratunga C, Duru V, Ramadani AP, Dacheux M, Khim
N, Zhang L, Lam S, Gregory PD, Urnov FD, Mercereau-Puijalon O, Benoit-Vical F, Fairhurst

RM, Menard D, Fidock DA. 2015. Drug resistance. K13-propeller mutations confer
artemisinin resistance in Plasmodium falciparum clinical isolates. Science 347:428-31.

50. Davis SZ, Singh PP, Vendrely KM, Shoue DA, Checkley LA, McDew-White M, Button-Simons
KA, Cassady Z, Sievert MAC, Foster GJ, Nosten FH, Anderson TJC, Ferdig MT. 2020. The
extended recovery ring-stage survival assay provides a superior association with patient
clearance half-life and increases throughput. Malar J 19:54.

51. Arendse LB, Murithi JM, Qahash T, Pasaje CFA, Godoy LC, Dey S, Gibhard L, Ghidelli-Disse S,
Drewes G, Bantscheff M, Lafuente-Monasterio MJ, Fienberg S, Wambua L, Gachuhi S,
Coertzen D, van der Watt M, Reader J, Aswat AS, Erlank E, Venter N, Mittal N, Luth MR,
Ottilie S, Winzeler EA, Koekemoer LL, Birkholtz LM, Niles JC, Llinas M, Fidock DA, Chibale K.
2022. The anticancer human mTOR inhibitor sapanisertib potently inhibits multiple
Plasmodium kinases and life cycle stages. Sci Transl Med 14:eabo7219.

52. Mukherjee A, Crochetiere ME, Sergerie A, Amiar S, Thompson LA, Ebrahimzadeh Z, Gagnon
D, Lauruol F, Bourgeois A, Galaup T, Roucheray S, Hallee S, Padmanabhan PK, Stahelin RV,
Dacks JB, Richard D. 2022. A Phosphoinositide-Binding Protein Acts in the Trafficking
Pathway of Hemoglobin in the Malaria Parasite Plasmodium falciparum. mBio 13:e0323921.

53. Al Monla R, Penzo M, Vallentin A, Lohia R, Vincent J, Berry L, Gomes AR, Cerdan R,
Wengelnik K. 2025. PI3-kinase has multiple functions in asexual blood stages of Plasmodium
falciparum. Sci Rep 15:16762.

54. Merrick CJ. 2015. Transfection with thymidine kinase permits bromodeoxyuridine labelling
of DNA replication in the human malaria parasite Plasmodium falciparum. Malar J 14.

55. Courtot L, Hoffmann JS, Bergoglio V. 2018. The Protective Role of Dormant Origins in
Response to Replicative Stress. Int J Mol Sci 19.

56. Mott BT, Eastman RT, Guha R, Sherlach KS, Siriwardana A, Shinn P, McKnight C, Michael S,
Lacerda-Queiroz N, Patel PR, Khine P, Sun H, Kasbekar M, Aghdam N, Fontaine SD, Liu D,
Mierzwa T, Mathews-Griner LA, Ferrer M, Renslo AR, Inglese J, Yuan J, Roepe PD, Su XZ,
Thomas CJ. 2015. High-throughput matrix screening identifies synergistic and antagonistic
antimalarial drug combinations. Sci Rep 5:13891.

57. Nardella F, Jiang T, Wang L, Bohmer MJ, Chakraborty S, Okombo J, Calla J, Silva TM, Pazicky
S, Che J, Jeon J, Vincent E, Boonyalai N, Coyle R, Buchanan MJ, Schaefer S, Chen D, Khan A,
Mayville E, De Souza ML, Treat M, Charlton J, Tumwebaze PK, Tjia S, Montejo L, Cover K,
Rosenthal PJ, Cooper RA, Bozdech Z, Lee MCS, Chakrabarti R, Desai SA, Fidock DA, Wang J,
Gray NS, Winzeler EA, Chakrabarti D. 2025. Plasmodiumfalciparum protein kinase 6 and
hemozoin formation are inhibited by a type II human kinase inhibitor exhibiting antimalarial
activity. Cell Chem Biol 32:926-941 e23.

58. Ward P, Equinet L, Packer J, Doerig C. 2004. Protein kinases of the human malaria parasite
Plasmodium falciparum: the kinome of a divergent eukaryote. BMC Genomics 5:79.

59. Ganter M, Goldberg JM, Dvorin JD, Paulo JA, King JG, Tripathi AK, Paul AS, Yang J, Coppens I,
Jiang RH, Elsworth B, Baker DA, Dinglasan RR, Gygi SP, Duraisingh MT. 2017. Plasmodium
falciparum CRK4 directs continuous rounds of DNA replication during schizogony. Nat
Microbiol 2:17017.

60. Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the
Ultrafast Bootstrap Approximation. Mol Biol Evol 35:518-522.

61. Safeukui I, Ware RE, Mohandas N, Haldar K. 2023. Simultaneous adjunctive treatment of
malaria and its coevolved genetic disorder sickle cell anemia. Blood Adv 7:5970-5981.

62. Yang J, Yan R, Roy A, Xu D, Poisson J, Zhang Y. 2015. The I-TASSER Suite: protein structure
and function prediction. Nat Methods 12:7-8.

Figure legends

Fig 1: <u>Evidence for DNA-damage-responsive checkpoint activity in *P. falciparum*</u>

A. BrdU incorporation into populations of trophozoites at three stages, after treatment with

MMS, as measured by ELISA. Parasites were treated with MMS for 1 h and BrdU was added

for the latter 30 mins. Means of two biological replicates, conducted in technical triplicate,

are shown; error bars show range. MMS treatments: 10x IC_{50}, 960 µM; 20x IC_{50}, 1.92 mM.

For each parasite stage, statistical testing was via ordinary one-way ANOVA and posthoc

Tukey's multiple comparisons test. *, P-value <0.05; comparisons not shown, ns. Pictures

show examples of trophozoites at each timepoint. Scale bar, 2µm.

B. Representative western blot of phosphorylated histone in trophozoite parasite lysate

treated with MMS (2 h, 10x IC_{50}). Phosphorylated histone signal was quantified versus

control histone H4: relative quantification is shown below the blot. Graph shows

quantifications from three biological replicate blots, relative to either H4 or Hsp70 as a

control protein. Unpaired two-tailed t-test: P-value = 0.0354.

Fig 2: <u>*Plasmodium* encodes no homologues of PIKK checkpoint kinases</u>

C. A phylogeny inferred for amino acid protein sequences of ATR, ATM, and TOR homologues

sampled from broad eukaryote groups. Apicomplexa homologues of PI3K and PI4K (the most

closely related proteins to ATR and ATM present in *Plasmodium*) and characterised

homologues of PI3K and PI4K from *Saccharomyces cerevisiae* and *Homo sapiens* were

included as an outgroup. Support values, displayed as percentages, were generated from

1000 ultrafast bootstrap replicates (60) and only support values ≥ 90, indicating strongly

supported clades, are shown. The scale bar represents the number of amino acid

substitutions per site. A fully-expanded version of this tree with no collapsed clades and

displaying sequence accessions is provided in the source data.

Fig 3: _Pf_PI3K is dispersed throughout the cell and does not visibly relocate to the nucleus upon DNA

damage

A. Western blots showing detection of _Pf_PI3K by two antisera raised to _Pf_PI3K peptides.

Replicate lanes from the same blot of protein lysate from late-stage _P. falciparum_ were

exposed to either pre-immune or immune serum. Histone H4 was used as a loading control.

B. Immunofluorescence assay (IFA) showing location of _Pf_PI3K (detected by antiserum 1) in _P._

falciparum. 2 representative cells are shown per stage. Scale bar 1 µm.

C. IFA showing _Pf_PI3K in _P. falciparum_ after DNA damage: 1 h treatment with 5x or 10x IC_{50} of

DHA or MMS on late-stage parasites. One representative cell per condition is shown. Scale

bar 1µm.

D. IFA showing _Pf_PI3K after expansion: maximum projections of 2-3 representative cells per

condition, as in (C). DAPI, blue; _Pf_PI3K, green; scale bar 10µm.

E. Single sections from the IFA in (D): nuclear _Pf_PI3K signals are highlighted with arrows. Full Z

stacks are shown in supplementary data.

Fig 4: Inhibitors of human PIKKs synergise with the DNA-damaging agent MMS in _P. falciparum_

A. Inhibition of parasite growth by VE-821 and KU-55933 (molecular structures of each

inhibitor are shown). Growth curves from five independent malaria SYBR green I-based

fluorescence (MSF) assays: percentage parasite growth is plotted as a function of drug

concentration, error bars are standard errors of the means. Mean IC_{50}s from all experiments

are shown (individual IC_{50} values from replicate experiments are in Fig S1B).

B. Isobolograms plotted as fractional inhibitory concentrations (FIC) of combinations of MMS

and VE-821 or KU-55933, as determined by MSF assay. FIC calculations were completed for

each combination by dividing the measured "apparent" IC_{50} values for individual drugs in the

different combinations by IC_{50} values obtained when the drugs were used alone. Each curve

represents a different biological replicate experiment. The mean FIC (ΣFIC(FIC$_A$+FIC$_B$)) is

shown, with range. The type of relationship was defined as moderately synergistic based on

well-established criteria: $\sum$FIC <0.5, substantial synergism; $\sum$FIC <1, moderate synergism;

$\sum$FIC ≥1 and <2, additive interaction; $\sum$FIC ≥2 and <4, slight antagonism and $\sum$FIC >4, marked

antagonism (61).

C.   BrdU incorporation into 30-32 hpi trophozoites after treatment with MMS, VE-821, KU-

55933, geneticin and combinations of these, measured by ELISA. Parasites were treated with

10x IC$_{50}$ of drug for 1 h and BrdU was added for the latter 30 mins. Six biological replicates

are plotted, each completed in technical triplicate. Mean and standard deviation error bars

are plotted, values normalised to control (set to 100%) in each replicate. Concentrations of

drugs used: MMS 960 μM, VE-821 1.62 μM, KU-55933 400 μM, geneticin 1500 μg/ml.

Ordinary one-way ANOVA and posthoc Tukey's multiple comparisons test of control vs all

and MMS vs all was completed; P-values: *, <0.05; **, <0.01; ***, <0.001; ****, <0.0001;

comparisons not shown, ns.

Fig 5: <u>The antimalarial drug dihydroartemisinin induces a DNA replication arrest in *P. falciparum*</u>

A.   Western blot of phosphorylated histones in ring, trophozoite and schizont-stage parasites

treated with DHA (4h, IC$_{50}$ (3nM)). Phosphorylated histone signal was quantified versus

control histone H4: relative quantification is shown below the blot.

B.   BrdU incorporation into 30-32 hpi trophozoites after treatment with DHA, VE-821, KU-55933

and combination treatments. Parasites were treated with 10X IC$_{50}$ of drug for 1 h and BrdU

was added for the latter 30 mins. Five biological replicates were plotted, each completed in

technical triplicate. Mean and standard deviation error bars are plotted; values were

normalised to control (set to 100%) in each replicate. Concentrations of drugs used: DHA 45

nM, VE-821 1.62 μM, KU-55933 400 μM. Statistical testing by ordinary one-way ANOVA and

posthoc Tukey's multiple comparisons test of control vs all and DHA vs all; P-values: *, <0.05;

comparisons not shown, ns.

C. Isobolograms plotted as Figure 4B. Each curve is a biological replicate and the mean FIC

(ΣFIC(FIC_A+FIC_B)) is shown, with range. The relationship between DHA and either VE-821 or

KU55933was defined as additive.

Fig 6: Ring-stage survival is affected by PIKK inhibitors

Data in A-D are from eRRSA assays on the strains 3D7, MRA-1252 and MRA-1240. Samples were

treated with DMSO (the control condition), VE-821, KU-55933, DHA and combinations of these. Drug

concentrations: VE-821, 81 nM (0.5X IC_{50}); KU-55933, 20 μM (0.5X IC_{50}); DMSO, 0.02%; DHA, 700 nM.

Values plotted are the means of -log(foldchange = 2^(DHA-DMSO)) for each treatment. Midline

represents the mean and boxes show min to max from three biological replicates, each conducted in

technical triplicate. Statistical testing by ordinary one-way ANOVA and Dunnett's multiple

comparisons test, comparing all means to the mean value from DMSO- or DHA-treated samples. P-

values: *, <0.05; **, <0.01; ***, <0.001; ****, <0.0001. Differences between DHA-treated samples

+/- PIKK inhibitors did not reach statistical significance, but the p-values for DHA + VE-821 or DHA +

KU-55933 versus control were < 0.0001, rather than 0.001 for DHA-alone versus control.

A. Ring-stage DHA-sensitive parasite strains, 3D7 or MRA-1252, were treated with DMSO (control

condition), DHA and inhibitors VE-821 or KU-55933 in combination with DHA for a 6 h pulse, then

left to recover for 72 h.

B. Control experiments without DHA: ring-stage 3D7 or MRA-1252 parasites were treated with

DMSO or inhibitors VE-821 or KU-55933 for a 6 h pulse, then left to recover for 72 h.

C. Ring-stages of the DHA-resistant parasite strain MRA-1240 were treated with DMSO (control

condition), DHA and inhibitors VE-821 or KU-55933 in combination with DHA for a 6 h pulse, then

left to recover for 72 h.

D. Control experiment without DHA: ring-stage MRA-1240 parasites were treated with DMSO or

inhibitors VE-821 or KU-55933 for a 6 h pulse, then left to recover for 72 h.

E. Inhibition of growth over a full 48h growth cycle by MMS, VE-821 and KU55933 was measured in

the strains 3D7, MRA-1252 and MRA-1240 via MSF assay. IC50s were calculated from at least

three independent experiments and all values are plotted, with the midline showing the mean

and error bars showing standard deviation. MSF growth curves are in Fig S1C.

Supporting information

Fig S1: Growth curves determining IC50 values for MMS, VE-821 and KU-55933 in the parasite

strains 3D7, MRA-1252 and MRC-1240.

Each panel shows growth curves from 3-5 independent malaria SYBR green I-based fluorescence

(MSF) assays: percentage parasite growth is plotted as a function of drug concentration; error bars

are standard errors of the means.

A.   Inhibition of growth of 3D7 strain by MMS.

B.   IC50 values from the growth curves shown in main figure 3A.

C.   Inhibition of strains MRA-1252 and MRA-1240 by the drugs VE-821, KU55933 and MMS.

Fig S2: 3D7 parasites survive a short pulse of MMS or DHA at the trophozoite stage, but their

maturation is delayed and progeny number per schizont, reduced.

A.   Images were taken 12h after trophozoites at 30-32hpi had been exposed to a 1 h pulse of

718   10X IC_{50} MMS or DHA (2 independent experiments, '1' and '2'). Images show that at this

719   timepoint (i.e. 42-44hpi), cultures were dominated by a mixture of immature and mature

720   schizonts, without clear evidence of parasite death in the damaged cultures. Parasites were

then treated with 75nM ML10 for 10 h, to arrest mature schizonts before egress, and re-

imaged. Mature schizonts then dominated in all cultures.

B.   Parasite stages imaged in (A) were categorised as either: early rings, small rings, large rings,

early trophozoites, mid trophozoites, late trophozoites, immature schizonts or mature

schizonts (2 independent experiments, n ≥ 100 from each experiment). Differences between

the proportions of stages in control vs DHA-treated vs MMS-treated cultures were tested via

two-way ANOVA and posthoc Tukey's multiple comparisons test. P-values are shown in a

table below the graph: *, <0.05; **, <0.01; ***, <0.001. Stars of matched colours across

rows denote a significant difference between those conditions (e.g. at 44hpi, there were still

more immature schizonts after MMS treatment than in the control (pink stars); 10h later,

this difference disappeared, with more mature and fewer immature schizonts (denoted by

pairs of yellow and blue stars). The only other difference was in the small numbers of rings

that escaped and reinvaded after prolonged arrest.

C.   Merozoites per schizont were counted in each of the cultures shown in (A) (n = 26-30 per

culture). Numbers of merozoites substantially increased in all cultures after the long ML10-

arrest period, which allowed mature schizonts to accumulate even after the damage-

induced delay seen in (B). A reduction in merozoite number was, however, still seen after

MMS damage, although this was not significant after DHA damage.

Fig S3: Protein sequence alignment of *P. falciparum* PI3K, PI4K and PIKKs from *S. cerevisiae* and

human.

MUSCLE (3.8) alignment of catalytic domains of *P. falciparum* PI3K and PI4K against human ATM,

ATR, PI3K, and yeast Mec1 and Tel1. Colour key is ClustalW standard. Highlighted in the yellow boxes

are the catalytic loops in ATR and ATM. Percentage identity matrix is shown below the alignment.

Fig S4: Predicted cellular locations of *Pf*PI3K and *Pf*PI4K, and anti-PfPI3K antibody production

A. Data plotted from (34), highlighting the clusters of *P. falciparum* proteins classified via

748   spatial proteomics as nuclear (yellow) and vesicular (brown). *Pf*PI3K (encoded by gene

749   PF3D7_0515300, red arrow)) is classified as nuclear, but falls at the boundary of the two

750   clusters, whereas *Pf*PI4K (encoded by gene PF3D7_0509800, green arrow) is classified as

751   vesicular and falls in the centre of the vesicular cluster.

752   B. Model made in i-TASSER (62), threading 1500 C-terminal residues of the 2133-amino acid

753   *Pf*PI3K against human ATR. Protein coloured blue to red, N-terminal to C-terminal. Peptides

754   used as antigens are highlighted in white and circled: amino acids 1359-1374 and 2011-2025

755   of full length *Pf*PI3K.

756   C. Controls for specificity of immune versus pre-immune antisera against *Pf*PI3K.

757   Representative images of single cells are shown. Scale bar 1 µm. Primary antibodies used

758   were rabbit antisera, pre-immune or immune, diluted 1:200 in 3% BSA.

Fig S5: Immunofluorescence after expansion, showing the location of PfPI3K

Examples of full Z-stacks taken during imaging of expanded cells: these stacks generated the

maximum projections shown in Figure 3D for control cell (iii), DHA-treated cell (i) and MMS-treated

cell (ii). They include the single sections extracted for Figure 3E. DAPI, blue; *Pf*PI3K, green; scale bar

10µm.

Fig S6: Isobolograms showing that two antibiotic drugs do not synergise with MMS

Isobolograms plotted as fractional inhibitory concentrations (FIC) of combinations of MMS and

geneticin (G-418), or blasticidin (BSD), as determined by MSF assay. Data were processed as in Figure

4B. The type of relationship was defined as additive for blasticidin and borderline antagonistic for

geneticin, based on well-established criteria ($\sum$FIC <0.5, substantial synergism; $\sum$FIC <1, moderate

synergism; $\sum$FIC $\geq$1 and <2, additive interaction; $\sum$FIC $\geq$2 and <4, slight antagonism and $\sum$FIC >4,

marked antagonism).

Fig S7: Ring stage survival assessed by flow cytometry

A ring-stage survival assay was assessed by flow cytometry as well as eRRSA as used in Figure 5.

Density plot quadrant graphs show one biological replicate of an RSA conducted on 3D7. Cell counts

for each quadrant are shown. Percentage survival was calculated as the proportion of live versus

dead cells (SYBR+ MITO+/SYBR+ MITO-) for each condition.

Fig S8: Raw growth-curve data for all isobolograms plotted in figures 4B, 5C and S5

A. Data for isobolgrams in figure 4B, VE-821 with MMS

B. Data for isobolgrams in figure 4B, KU-55933 with MMS

C. Data for isobolgrams in figure 5C, VE-821 and KU55933 with DHA

D. Data for isobolgrams in figure S5, Blasticidin and G-418 with MMS

Source data files 1-6

Files 1-6: Source data for phylogenetic tree shown in figure 2

A

Rabbit 1

| Pre-immune | Immune |

250 kDa

15 kDa
10 kDa
α-H4

Rabbit 2

| Pre-immune | Immune |

250 kDa

15 kDa
10 kDa
α-H4

B

| | Phase | PI3K | DAPI | merge |

Rings

Trophozoites

Schizonts

C

Control

10x IC50 DHA

10x IC50 MMS

D

Maximum Intensity Projection

| DAPI | PI3K | merge |

Control (i)

(ii)

(iii)

10x IC50 DHA (i)

(ii)

10x IC50 MMS (i)

(ii)

E

Single sections

Control (iii)   DHA (i)   MMS (ii)

A

	Ring		Trophozoite		Schizont		
	−	+	−	+	−	+	DHA

15kDa
10kDa — a-H4

15kDa — a-γ-H2aX

1 : 3.1   1 : 1.6   1 : 2.3

B

normalised percentage BrdU

Control, DHA, VE-821, KU-55933, DHA+VE-821, DHA+KU-55933

C

FIC$_{50}$ (DHA) vs FIC$_{50}$ (VE-821)

FIC$_{50}$= 1.32(1.18-1.41)

FIC$_{50}$ (DHA) vs FIC$_{50}$ (KU-55933)

FIC$_{50}$= 1.15(1.09-1.20)