Regulation of somatic stem cell precursor fates and proliferation during *Drosophila*

***melanogaster* pupal ovary development resembles the signaling framework for adult**

stem cell behavior

Short title: Signals regulating Drosophila ovarian FSC development

Rachel Misner, Amy Reilein, Helen V. Kogan and Daniel Kalderon*

Department of Biological Sciences, Columbia University, 1212 Amsterdam Avenue, New York,

NY10027 USA

* corresponding author: ddk1@columbia.edu

13
14
15

Abstract

Follicle Stem Cells (FSCs) in the Drosophila melanogaster ovary are maintained through independent regulation of division and differentiation to become proliferative Follicle Cells (FCs) to their posterior or quiescent Escort Cells (ECs) to their anterior. These behaviors are guided by graded extracellular Hedgehog (Hh) and Wnt signals emanating from cells anterior to FSCs and an inverse gradient of JAK-STAT pathway activity. Here, we used lineage analyses to investigate the development of ECs, FSCs and FCs from a common set of precursors during pupation. Previous studies found that the most anterior precursors divide slowest, with quiescence spreading from the anterior over time to include all ECs, that FSCs are specified simply by their location at eclosion, and that the first FCs derive from cells that accumulate posterior to the developing germline over the first 48h of pupation. We now provide evidence that those accumulating cells derive from migration out of the developing germarium. We found that Wnt pathway activity favored conversion of precursors to more anterior adult derivatives (ECs rather than FCs), while JAK-STAT pathway activity favored posterior outcomes. Posterior bias and faster division, explored by altering Cyclin E activity, both favored a precursor becoming an FSC. Both JAK-STAT and Hh signaling could increase precursor division rate. All of these characteristics resemble regulation of adult FSC behavior. We suggest that similar signaling networks and division rate dependence during maintenance and development may be general features for stem cells that are specified in parallel with tissue development and that exhibit division-independent differentiation.

Article Summary

Drosophila ovarian Follicle Stem Cell (FSC) behavior and regulation by graded Hedgehog (Hh), Wnt and JAK-STAT signals is well understood, motivating us to study mechanisms guiding development of FSCs and the neighboring cell types they maintain. The marked progeny of single cells labeled at pupariation were compared in adults for lineages with genetic alterations affecting signal transduction. Wnt and JAK-STAT had strong opposing influences on the anterior-posterior location, and hence identity of derivatives, while JAK-STAT and Hh (via Yorkie) stimulated cell division. Conversion of precursors to FSCs was favored by higher division rates.

Introduction

Many adult tissues are maintained by adult stem cells (CLEVERS AND WATT 2018; POST AND CLEVERS 2019; KALDERON 2022; GAO *et al.* 2023; BEUMER AND CLEVERS 2024). The two universal stem cell behaviors of cell division and initiating differentiation at appropriate, balanced frequencies are inevitably regulated by their physical and signaling environment, often known as the stem cell niche (VOOG AND JONES 2010; BITEAU *et al.* 2011; LOSICK *et al.* 2011). Moreover, the further differentiation of immediate stem cell derivatives is often guided by additional cues from the mature adult tissue. Thus, stem cell function depends on the nature and spatial organization of the stem cells themselves, neighboring niche cells and more distant cells guiding full differentiation. The behavior and regulation of different types of adult stem cell is being studied productively for several paradigms. By contrast, the developmental mechanisms that place an appropriate number of stem cells in a suitable niche and larger tissue context are largely unknown. Understanding that process would have the potential

benefit of facilitating the creation of self-sustaining organoids and exploring developmental
abnormalities leading to impaired adult stem cell function or cancer susceptibility.
The Drosophila ovary is an excellent paradigm for studying the development of adult germline
and somatic stem cells (GIEDT AND TOOTLE 2023; ST JOHNSTON 2023). Each adult ovary consists of 15-20
ovarioles, with an anterior germarium and a series of developing egg chambers of increasing
maturity, with each ovariole producing a mature egg as often as every 12h (Fig. 1A) (SPRADLING *et al.*
1997; DUHART *et al.* 2017; GIEDT AND TOOTLE 2023). This formidable cell production is fueled by 2-3
Germline Stem Cells (GSCs) (ELIAZER AND BUSZCZAK 2011) and about sixteen somatic Follicle Stem Cells
(FSCs) (REILEIN *et al.* 2017; HAYASHI *et al.* 2020; KALDERON *et al.* 2021; KALDERON 2022). The known
biology of these two types of stem cell includes numerous interesting contrasts. GSCs generally
remain in place at the anterior of the germarium, are individually long-lived, maintained by single-cell
asymmetry and depend on anti-differentiation signals from immediately adjacent niche cells (Cap
cells) (KAHNEY *et al.* 2019; ZHANG AND CAI 2020; ST JOHNSTON 2023). The differentiation of the immediate
"cystoblast" derivative of a GSC depends on interactions with non-dividing somatic Escort Cells (ECs)
(KIRILLY *et al.* 2011) during posterior migration and is manifest as a set of four synchronized divisions
with incomplete cytokinesis through region 1 to form a sixteen-cell region 2 cyst.
FSC cell bodies reside principally in two adjacent rings along the anterior-posterior (A/P) axis,
surrounding a central germline stage 2b cyst and bounded on the other side by the basement
membrane lining the germarium (Fig. 1A) (REILEIN *et al.* 2017; KALDERON 2022). FSCs can exchange A/P
locations and move circumferentially. They are maintained by population asymmetry with
independent cell division and differentiation, characterized by frequent and variable, apparently
stochastic, loss and expansion of individual cell lineages (REILEIN *et al.* 2017; REILEIN *et al.* 2018;
KALDERON 2022). This picture of FSC organization and behavior differs markedly from an earlier model
as a result of more thorough investigation and discarding earlier expectations of single-cell
asymmetry, which led to highlighting only a minority of all FSC lineages (MARGOLIS AND SPRADLING 1995;
KALDERON 2022). FSCs are responsible for producing about six founder Follicle Cells (FCs) for each
future egg chamber. A few of these FCs arrest division soon afterwards and form specialized polar
cells at the anterior and posterior termini of each egg chamber, as well as adjacent stalk cells, which
connect adjacent egg chambers and allow budding of an egg chamber from the germarium (Fig. 1A)
(GRAMMONT AND IRVINE 2001; ASSA-KUNIK *et al.* 2007; DAI *et al.* 2017). Other FCs continue division in
growing egg chambers to form an expanding single-layer epithelium of about 650 cells by mid-
oogenesis (stage 6). These FCs acquire different identities according to their location, guided by JAK-
STAT signals produced in polar cells and EGFR-stimulating signals from the oocyte, initially at the
posterior of the germline cyst, and then from the oocyte nucleus after migration to an anterior
cortical location that defines the future dorsal aspect of a developing embryo (MCGREGOR *et al.* 2002;
XI *et al.* 2003; MERKLE *et al.* 2020; ST JOHNSTON 2023). FSCs also produce new ECs in adults, albeit at a
much lower rate than FCs (REILEIN *et al.* 2017; MELAMED AND KALDERON 2020). The A/P location and
differentiation of FSCs are guided principally by long-range signals, emanating from the anterior
(multiple Wnt ligands from Cap cells and ECs) and posterior (JAK-STAT ligand Unpaired from newly-
differentiating polar cells) to produce inverse gradients of those pathway activities across the stem
cell domain, effectively defining the A/P extent of the FSC domain (REILEIN *et al.* 2017; MELAMED AND
KALDERON 2020). JAK-STAT pathway activity also stimulates FSC division in a graded fashion (MELAMED
AND KALDERON 2020; MELAMED *et al.* 2023). Hedgehog pathway activity also stimulates FSC division but
may not contribute to the spatial pattern of FSC division (VIED AND KALDERON 2009; HUANG AND KALDERON
2014). Other factors supporting the much faster division of posterior FSCs remain to be identified.
The division rate of posterior FSCs approximately matches the rate of their conversion to FCs, so the

exchange of FSCs between anterior and posterior layers is roughly at equilibrium (MELAMED *et al.*
2023).
110   During development, germline and somatic gonadal cells are specified separately in the
embryo and coalesce after germ cell migration to form an embryonic gonad (BOYLE AND DINARDO 1995;
MURRAY *et al.* 2010; JEMC 2011). During larval stages, the ovary develops from anterior to posterior
(initially termed apical to basal), starting with the specification of somatic Terminal Filament (TF)
cells, which resolve into separate intercalated stacks that seed each individual ovariole (GODT AND LASKI
1995; GILBOA 2015). A Notch signal from TF cells recruits adjacent cells from a pool of somatic cells
(Intermingled Cells: ICs) intermingled with germ cells, to become quiescent Cap cells (WARD *et al.*
2006; SONG *et al.* 2007; PANCHAL *et al.* 2017; ZAMFIRESCU *et al.* 2022). Germ cells that contact Cap Cells
become the future GSCs, with other germ cells, in more posterior locations, initiating differentiation
soon afterwards (ZHU AND XIE 2003; ASAOKA AND LIN 2004; GANCZ *et al.* 2011; GANCZ AND GILBOA 2013). As
Cap cells and GSCs form in third instar larvae, somatic Swarm Cells migrate basally (towards the
future posterior) around the TF cells to separate sets of developing ICs and germ cells into individual
units (BANISCH *et al.* 2021). Then, during pupation, germ cells sequentially mature (towards the
posterior) into sixteen-cell cysts, while somatic ICs amplify. These IC precursors eventually become
ECs, FSCs and FCs, such that the newly-eclosed adult has a mature germarium structure, followed by
four egg chambers of increasing size, allowing egg-laying to commence within about a day (REILEIN *et*
al. 2021).
127   The specification of different ovarian somatic cell types during pupation has been studied by
lineage analyses and direct morphological observation, including live imaging, yielding three major
conclusions. First, FSCs are not specified at a particular time during development; instead, single-cell
lineages generally include ECs and FCs together with FSCs, with the latter defined simply by occupying
the appropriate A/P locations at eclosion (REILEIN *et al.* 2021). Thus, in this paradigm, the stem cells
are not set aside early in development but derive from the same process that forms the whole adult
tissue. Another study had suggested separate precursors for ECs and FSCs at the start of pupation,
based on lineage analysis initiated by two GAL4 drivers (*Con-GAL4* and *bond-GAL4*) together with
temperature sensitive GAL80 (SLAIDINA *et al.* 2020). That suggestion was, however, based on assuming
that the additional presence of marked ECs in most adult ovarioles with marked FSCs was due to the
presence of two or more lineages. Sparse induction of cell lineages showed this explanation to be
incorrect; FSC-containing single-cell lineages almost always included ECs, revealing a common origin
(REILEIN *et al.* 2021). Moreover, using the same reagents, it was apparent that neither the frequency,
nor the time of lineage initiation could be controlled satisfactorily using the temperature-sensitive
GAL80 technique, contrasting with lineage initiation using a heat-shock to activate recombinase
encoded by a *hs-flp* transgene (REILEIN *et al.* 2021). Second, anterior somatic cell precursors divide
more slowly than posterior precursors and terminate division during pupation. This feature appears
to limit the number of the most anterior derivatives, Escort Cells (REILEIN *et al.* 2021). Third, the
formation of the first FCs and the first egg chamber is morphologically very different from the process
in adults, where newly-formed FCs are adjacent to a germline cyst and that germline cyst is
subsequently surrounded by a monolayer of FCs before budding from the posterior of the
germarium. Instead, many somatic cells accumulate posterior to the germarium (harboring all germ
cells) during the first half of pupation and are then invaded by a germline cyst released from the
germarium and migrating a considerable distance towards the posterior (REILEIN *et al.* 2021).
151   We would like to understand what orchestrates the movements and proliferation of somatic
cells during pupation required to adopt appropriate adult identities and assemble a morphologically
normal adult ovariole. As a first step, we have investigated cell-autonomous responses of somatic

precursors to genetic alterations by using lineage analyses during pupation to measure final
outcomes as ECs, FSCs and FCs. We began by investigating signals and cellular properties known to
influence the behavior of adult FSCs. This includes examining signals instrumental to FSC maintenance
and differentiation (Wnt and JAK-STAT), signals and effectors important for FSC division (JAK-STAT,
Hedgehog (Hh) and Yorkie (Yki)), and the effect of cell division rate on cell fate (ZHANG AND KALDERON
2001; SONG AND XIE 2003; VIED AND KALDERON 2009; WANG AND KALDERON 2009; VIED *et al.* 2012; WANG *et*
al. 2012; HUANG AND KALDERON 2014; WANG AND PAGE-MCCAW 2014; REILEIN *et al.* 2017; HAYASHI *et al.*
2020; MELAMED AND KALDERON 2020; MELAMED *et al.* 2023). Among adult FSCs, there is competition for
residence in the FSC domain, with conversion of FSCs to FCs or ECs balanced by FSC amplification
through cell division. Hence, maintenance and amplification of an individual FSC lineage is cell-
autonomously favored by genetic alterations that either increase the division rate or reduce the
frequency of differentiation (REILEIN *et al.* 2017; REILEIN *et al.* 2018). We found that this principle and
the roles of Wnt and JAK-STAT signaling apply also to the pupal development of FSCs, while responses
to Hh and Yki are at least partly shared for stem cell development and adult maintenance. A key
source of anterior signals (Wnt, Hh) is present at the start of pupal development. The original source
of the ligand activating the JAK-STAT pathway, which guides FC formation from the start of pupation,
is not defined but likely has a posterior bias; the major source, which instructs a robust gradient of
pathway activity, emerges after the first egg chamber forms midway through pupation. These
experimental findings and their rationalization suggest principles that may be shared generally among
adult stem cells with division-independent differentiation that is guided by location: early
establishment of niche signaling centers, continuous competition between division and
differentiation for an individual cell to become or remain a stem cell, and regulation of division and
cell location by the same signaling networks during adult stem cell development and maintenance.
177
178
179
180   **Materials and Methods**
181
182   **Clone induction via MARCM**
183
GFP-marked MARCM clones (LEE AND LUO 2001) were induced on different chromosome arms (2L, 2R,
and 3R) using flies of the genotypes listed below ("X" and "Y" identities are listed subsequently; ">"
indicates an FRT element, "*tub*" refers to the α*tub84B* promoter (LEE AND LUO 2001) and "*act*" to the
act5C promoter (PIGNONI AND ZIPURSKY 1997)). In all cases, recombination is initiated at *FRT* sites by
heat-shock induction of FLP recombinase, leading to marked clones lacking a *Gal80* transgene and
expressing *UAS-GFP* from *tub-GAL4* and *act>GAL4* drivers. Simultaneously, a recessive allele ("X")
becomes homozygous or a transgene ("*UAS-(Y)*") initiates expression, or both, or neither (controls).
191
192   (2L) *hs-FLP UAS-nGFP tub-Gal4*; *(X) FRT40A/tub-Gal80 FRT 40A*; *UAS-*
193   *(Y)/act > CD2 > GAL4*
194   (2R) *hs-FLP UAS-nGFP tub-Gal4*; *FRT42D (X) / FRT 42D tub-Gal80*; *UAS-*
195   *(Y)/act > CD2 > GAL4*
196   (3R) *hs-FLP UAS-nGFP tub-Gal4*; *+ / act > CD2 > GAL4, UAS-GFP*; *FRT 82B (X) / FRT 82B tub-Gal80*.
197
198   Studies conducted at 29C utilized flies of the genotype below (where an additional *act-GAL80*
199   element on 2R is necessary to silence GAL4):

200   (2R) *hs-FLP UAS-nGFP tub-Gal4*; *FRT42D (X) / FRT 42D act-Gal80 tub-Gal80*; *UAS-(Y)/act > CD2 > GAL4.*
201

For 2L chromosome studies, (X) included *NM* (control), *cycE^{wx}*, *smo^2*, *pka-C1^{H2}*, *smo^2 pka-C1^{H2}*,
cutlet^{4.5.43}, and (Y) included *DIAP1*, *CycE*, *Yki^{S168A}*; also, a *Su(fu)^{LP}/ Su(fu)^{LP}* background was used for
smo pka clones.
For 2R chromosome studies, (X) included *sha* (control), *ptc^{S2}*, *arr^2*, *yki^{B5}*, *ptc^{S2} yki^{B5}*, *hpo^{42-47}* and (Y)
included *Hop* together with *UAS-Dap*, as well as a *tub-yki* transgene that does not include the *UAS*
promoter.
For 3R chromosome studies, (X) includes *NM* (control), *apc1^1 apc2^2*, *axn^{E77}*, *kibra^{del}*, *wts^{x1}*, and
stat^{85C9}.
210

Lineage analysis by MARCM in newly-eclosed flies
211
212

From the time that pre-mated adults were put into a fresh vial (for 24h or less) to eclosion of the first
several adult progeny, sufficient for analysis, was most commonly 11 days (REILEIN *et al.* 2021). Timing
of heat shocks to initiate lineages was determined with this 11-day course in mind. The larval-to-
pupal transition, where third-instar larvae are crawling up the sides of the vial and beginning
puparium formation, was generally 5 days before eclosion. Hence, adults collected 5d after heat-
shock were considered to have lineages initiated at pupariation ("0h APF"). At the time of heat-shock
(generally 6d after parents were first added to the vial), some animals were at pupariation; others
were slightly more advanced or delayed. Vials were cleared of any eclosed adults the evening before
collecting newly-eclosed adults over a period of no more than 6h centered on 120h after heat-shock.
Similar procedures were used for collection of adults with lineages designated as initiated 2 days
prior to pupariation ("-2d APF": heat-shock 4d after introducing parents; collect adult progeny over
6h, centered on 168h after heat-shock) and 36h after pupariation ("36h APF": heat-shock 7.5d after
introducing parents; collect adult progeny over 6h, centered on 84h after heat-shock). Vials were
heat shocked at 33C for 12 min (0h APF, 36h APF) or 20 min (-48h APF). These conditions were
chosen as the mildest conditions consistently inducing a similar yield of marked lineages at 0h APF, so
that many ovarioles would harbor a single marked lineage. To achieve a similar result, a longer period
of heat-shock was used at -48h APF because fewer precursors are present. Afterwards, vials were
incubated at 25C or 29C. Higher temperature increases GAL4 activity and was used for expression
of *UAS-Hop* and controls for that test (analogous tests in adult ovaries showed that 29C is necessary
to achieve strong activation of the JAK-STAT pathway (MELAMED AND KALDERON 2020)); all other flies
were incubated at 25C or room temperature (23C-25C). Flies were dissected on the day of eclosion
(0d).
235

Adult ovary fixation and staining
236
237

Adults were dissected in PBS using forceps by separating the entire posterior region of the abdomen,
revealing the ovaries within. Non-ovary material, such as fat bodies and other organs, was gently
shaken or teased out with forceps, and the ovaries (still attached to the cuticle) were transferred
using forceps to tubes containing 400 μl fixative. Ovaries were fixed in 4% paraformaldehyde in PBS
for 10 min at room temperature, rinsed 3x in PBS with 0.1% Triton and 0.05% Tween-20 (PBST), and
blocked in 10% normal goat serum (Jackson ImmunoResearch Laboratories) in PBST. Ovarioles were
incubated in primary antibody for 1 hr. Ovarioles were rinsed in PBST 3x and incubated for 1 hr in
secondary antibodies diluted 1:1000. Ovaries were rinsed 2x in PBST and 1x in PBS for 5 min each.
Ovaries were transferred using a p1000 with the narrow portion of the tip cut off to a well containing

PBS. The cuticle was then separated from the ovaries, which were transferred using forceps to a slide
containing a drop of PBS. Each slide accommodates two drops, and each drop contained the ovaries
from one individual animal. Ovaries were gently teased apart with extra sharp forceps, separating
into clusters of 1-4 ovarioles, taking care to ensure that entire ovarioles remained intact and did not
separate, as tracing the clones throughout the entire ovariole is crucial. After separation was
complete, 25 µl of Flouromount-DAPI was applied to a coverslip and carefully placed over the portion
of the slide with each sample.
254
255 **Immunohistochemistry**
256
Monoclonal antibodies against Fasciclin III (Fas3) and Vasa were obtained from the Developmental
Studies Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of Iowa,
Department of Biology, Iowa City, Iowa 52242. 7G10 anti-Fasciclin III was deposited to the DSHB by C.
Goodman, and was used at 1:250. Anti-Vasa (used at 1:10) was deposited by A. C. Spradling/D.
Williams and used for staining adult ovaries. For pupal ovaries, the monoclonal rat anti-Vasa was not
effective and instead rabbit anti-Vasa (gift from R. Lehmann, MIT) was used. Other primary antibodies
used were anti-GFP (A6455, Molecular Probes) at 1:1,000; and goat FITC-anti-GFP (Abcam ab6662) at
1:400. Secondary antibodies were Alexa-488, Alexa-546, Alexa-594 or Alexa-647 from Molecular
Probes. DAPI-Fluoromount-G (Southern Biotech) was used as mounting medium for all experiments.
Images were collected with a Zeiss LSM700 or LSM800 laser scanning confocal microscope (Zeiss)
using a 63x 1.4 N.A. lens.
268
269 **Scoring and Analysis of marked lineages in adult ovaries**
270
Ovaries were stained using antibodies to Vasa, Fas3 and GFP. Vasa staining assists in classifying cysts
of regions 1 (fewer than 16 cells) versus regions 2a and 2b (16-cell cysts). The anterior border of
strong Fas3 expression corresponds to the border between FSCs and FCs, with cells immediately
adjacent and anterior to this boundary, together with their immediate anterior neighbors (three rings
of cells anterior to the boundary of strong Fas3 staining), classified as FSCs. Escort cells were scored
as r1 or r2a based on measurements that were performed with wild-type ovaries to determine the
lengths of these regions. We previously examined EdU incorporation in ovaries of newly -eclosed flies
and measured the distance from cap cells to the end of region 1 as determined by the presence of
cysts that incorporated EdU. Region 1 cysts are going through mitosis and Region 2a cysts have the
complete complement of 16 germ cells. The length ratio of regions 1:2a is about 3:1 in newly-eclosed
adults. To measure the percentage of each terminal egg chamber covered by marked FCs, the ratio of
GFP-marked cells to unmarked cells (stained with DAPI) was estimated throughout all z-sections.
The total number of ovarioles containing any GFP-positive cells ("clones" or "lineages") was recorded,
along with the total number of ovarioles, for each coverslip, which has a pair of ovaries from one
animal. Each ovariole was scored for GFP-positive cells by counting the number of r1 ECs, r2a ECs, and
FSCs. FCs were scored for presence (yes or no) in the following regions: "Immediate Daughter"
(region 2b), "Germarium" (region 3), "1", "2", "3", or "4" (corresponding to the four egg chambers
typically present in ovaries of newly-eclosed flies; sometimes "5" also), and "BS" (Basal Stalk; a cluster
of about 10 cells around the polar cells at the posterior of the terminal, most mature, egg chamber).
Notes were also added if labeled cells of a specific genotype were present at unusual frequencies in
certain locations, including polar cells and stalk cells. Each scored ovariole was then labeled with its
broad clone type (EC, EC/FSC, EC/FSC/FC, EC/FC, FSC, FSC/FC, or FC). This classification, together with
the exact number of GFP-labeled r1 and r2a ECs, FSCs, and the presence or absence of FCs in the

regions listed above, was recorded in each row of a spreadsheet for a given genotype in an
experiment. For each genotype in an experiment, we aimed to score at least 50 marked ovarioles
across at least three coverslips. The data from each such spreadsheet were aggregated in order to
measure several parameters, as described in the next section.
298
299   **Estimating single-cell lineage outcomes underlying raw observations**
300
Although we used a mild and brief heat-shock in order to induce only low activity of a *hs-flp*
transgene, it is inevitable that some adult ovarioles will include lineages from more than one marked
precursor. Previously, two methods were considered for estimating single-cell lineage outcomes from
raw observations (REILEIN *et al.* 2021). One assumed that every marking event is independent (zero-
clone frequency method, below). Another used information about the frequency of ovarioles with
marked ECs and FCs (but no marked FSCs) as a likely indicator of the presence of more than one
lineage. For wild-type lineages, the latter method (of the two) indicated a higher frequency of
ovarioles with more than one lineage for a given sample, and was the preferred method (because we
believed it to be more accurate). Here, that method cannot be used because different genotypes may
greatly alter EC and FC frequencies, as well as producing single lineages with only marked ECs and FCs
(from selective loss of prospective FSCs). Past experience suggested that the "zero-clone" frequency
method used here will underestimate the number of ovarioles with more than one lineage. An
indicator of that potential error is the proportion of deduced single-cell lineages with only EC and FCs;
the average frequency of that lineage category was 5% for controls. However, during the course of
the studies reported here we learned that lineages with just ECs and FCs can arise from loss of FSCs
from a single EC/FSC/FC lineage rather than representing simultaneous EC-only and FC-only lineages.
Thus, the error in estimating single-cell lineage frequency and composition by the method described
below is less than initially anticipated and overestimated by the EC/FC frequency.
319
320   **Zero-clone frequency method**
321
We assume all marking events are independent and use binomial calculations to derive the expected
frequency of ovarioles with 1 and 2 lineages by deriving the probability of a recombination event in
each precursor cell.
Although we need to use an estimate of the total number of cells present at the time of
recombination, the numerical outcomes are not very sensitive to this exact number.
From prior published work (REILEIN *et al.* 2021), lineage data estimated 24 or 17 precursor cells at
pupariation (different experiments and labeling techniques). The number of precursor cells directly
observed was 38 at 12h APF and 51 at 24h APF. This leads to an estimated number at 0h APF
(assuming similar expansion from 0-12h and 12-24h APF) of 38 x 38/51= 28 cells
Since MARCM (the same technique used here) was used to deduce 24 precursors, we use that
number.
333
If the frequency of ovarioles with no marked cells = f and p is the probability of recombination in a
single cell,
$f = (1-p)^{24}$
The proportion of ovarioles with a single clone will be $24p(1-p)^{23} = f(24p)/(1-p)$.
Hence, the frequency of single-cell clones among labeled ovarioles = $f(24p)/((1-p)(1-f))$.
The frequency of ovarioles with 3 or more clones (estimated by this method) will likely be <20% of
remaining ovarioles. Hence, we approximate, for simplicity, that all of the remaining ovarioles (non-

zero and non-single cell clones) contain two clones. We also round up to a whole number for the
estimated percentages of ovarioles with zero, 1 or 2 lineages (adding to 100).
343
344 For lineages induced 48h before pupariation, we previously estimated there might be 10 precursor
345 cells.
346 So, $f = (1-p)^{10}$
347 and the frequency of single-cell clones among labeled ovarioles = $f(10p)/((1-p)(1-f))$.
348

349 **Estimating single-cell lineage type frequencies**
350
351 After establishing the estimated number of ovarioles with one and two lineages for a given
352 experimental sample, we deduce the total number of single-cell lineages present and then estimate
353 the fraction of those of specific types. The major types are outlined first and then sub-divisions are
354 brought in.
355
356 We start with raw data of major phenotypes: EC, EC/FSC, FSC, FC, EC+FC
357 Single-cell clone frequencies are named: EC(q), EC/FSC (r), FSC(s), FC (t), EC+FC(m)
358 Single lineage frequency = p1, two-lineage frequency = p2
359
360 Observed EC only = $q.p1 + q^2.p2 = q^2.p2 + q.p1$
361 Hence, we can calculate q directly
362
363 Observed FC only = $t.p1 + t^2.p2 = t^2.p2 + t.p1$
364 Hence, we can calculate t directly
365
366 Observed FSC only = $s.p1 + s^2.p2 + 2s.t.p2 = s^2.p2 + s(p1+2.t.p2)$
367
368 Observed EC+FC = $m.p1 + m^2.p2 + 2.p2.m(q+t) + 2.p2.q.t$
369 $= m^2.p2 + m (p1+2.p2. (q+t)) + 2.p2.q.t$
370
371 Observed EC/FSC = $1- (q+s+t+m)$
372
373 Then, for r1 (a) and terminal FCs (z)
374 Observed r1 only = $a.p1 + a^2.p2 = a^2.p2 + a.p1$
375
376 Observed "Terminal FC only" = $z.p1 + z^2.p2 = z^2.p2 + z.p1$
377 Note- here, "Terminal FC only" means there are marked cells in the terminal egg chamber and there
378 are no marked cells other than FCs.
379
380 r1 ECs + FC (b) and EC + terminal FC only (y) can also be inferred.
381
382 Observed r1+FC = $b.p1 + b^2.p2 + 2.p2.b(q+t) + 2.p2.a.t$
383 $= b^2.p2 + b (p1 + 2.p2 (q+t)) + 2.p2.a.t$
384
Observed EC + terminal FC only = $y.p1 + y^2.p2 + 2.p2.y(q+t) + 2.p2.q.z$
$= y^2.p2 + y (p1 + 2.p2 (q+t)) + 2.p2.q.z$

The above two equations exclude EC + FC with no r1 only or terminal FCs but that is a very small
error.

For "only Term FC", x, meaning no marked cells (even FCs) anterior to the penultimate egg chamber,
and a similar category of EC + only Terminal FC, w, there are analogous equations.

Observed "only Terminal FC" = $x.p1 + x^2.p2 = x^2.p2 + x.p1$

Observed EC+ only Terminal FC = $w.p1 + w^2.p2 + 2.p2.w(q+x) + 2.p2.q.x$
$= w^2.p2 + w (p1 + 2.p2 (q+x)) + 2.p2.q.x$

Added categories
400

(a) "Normalized" numbers to acknowledge that EC+FC category sometimes includes an EC-only and
an FC-only lineage.
We acknowledge that the zero-clone frequency method might underestimate multiple clone
frequency. Hence, the EC+FC category may include some EC-only and FC-only clones.
However, it is (i) reasonable that some prospective FSCs might be lost for WT from EC/FSC/FC
lineages to give EC+FC lineages, (ii) genotypes with reduced division rate may cause selective loss of
FSCs (this work), converting prospective EC/FSC/FC to EC+FC lineages. This may also occur to a degree
for all genotypes. So, taking EC+FC as possibly composite (2 lineages) should be avoided if there is a
division defect and is likely an overestimate of composite clones in all cases.

Method: assume all EC+FC (fraction m) are really single EC and single FC clones. Hence, m is added to
EC-only & FC-only proportions (same for EC+FC specifically with term FCs or r1 ECs) and the total
clone number is multiplied by (1+m) to give "normalized" EC and FC values.
These values appear in supplementary spreadsheets but are not discussed in the text.

(b) Clones with any terminal FC or r1 EC representation.
This is useful for considering cell allocation to the most extreme A/P locations (independent of other
issues).
After deriving single-cell lineage frequencies,
Observed raw # of ovarioles with marked terminal FCs = "ObsTerm"
Raw ovarioles with clones = x
Deduced fraction of ovarioles with two lineages = f
Fraction of single-cell lineages with marked terminal FCs = T

ObsTerm = $x(1-f)T + xf(T^2+2T(1-T))$
$= xT-xfT+xfT^2+2xfT-2xfT^2$
$= T(x+xf)-T^2xf$

$(xf)T^2 - x(1+f)T + ObsTerm = 0$

Quadratic solution = $((x(1+f)-SQRT[(x+xf)^2-4xf.ObsTerm]))/(2xf)$

Same for r1, using Obsr1 to calculate R

Criticisms/shortcomings
436

If clone frequency varies more from fly to fly than ovariole to ovariole (within the same fly), then
these corrections ought theoretically to be applied to each fly/coverslip. However, this is impractical
(the methods cannot work well when n is low). To minimize this deficiency, it is best if all scored
coverslips have roughly similar clone frequencies. We therefore instituted the criterion that coverslips
(containing ovaries from a single fly) scored for clone phenotypes should have ovarioles with labeled
cells at a frequency of 75% or lower (there were one or two experiments where we included
coverslips with higher frequencies because there was no choice). We also instituted a minimum clone
frequency of 15%. This was for two reasons. First, so that all clone frequencies were in a narrower
range, so that errors due to single-cell lineage estimations would not vary widely among samples.
Second, we had observed previously that there is occasional cell marking in the absence of deliberate
heat-shock. Such lineages arise at unknown times (if originating from recombination; most appear to
be single EC marks of unknown origin). Such background is less than 5% and has minimal impact if
deliberately-induced lineages are prevalent. For controls, the average frequency of marked ovarioles
scored was 42% at pupariation and 63% 48h before pupariation.
Because any error in single-cell lineage estimations likely depends on clone frequency, we
experimented with splitting each control sample into two of similar number, with the lowest lineage
frequency flies segregated from the highest lineage frequency flies. We processed the data separately
and then aggregated. The final outcomes were almost identical, so we treated all fly ovaries from one
experiment as a single batch for single-cell lineage estimations.

Terminal FC labeling with or without other FCs
458

"Only terminal FC" means labeled cells in the basal stalk and/or terminal egg chamber (which may
extend to the penultimate egg chamber) but nowhere else in the ovariole.
"FC-only with terminal FCs" includes the category above, but also includes lineages with terminal FCs
and additional FCs anterior to the penultimate egg chamber, but no ECs or FSCs.
Subtracting the category above from "FC-only" gives the FC-only lineages that do not include any
labeled terminal FCs.
Thus, within the FC-only category, frequencies can be derived for only Terminal FCs, no terminal FCs,
and a mix of terminal and non-terminal FCs.
These calculations were performed only for control lineages in order to determine if there were
common precursors of FCs of terminal and non-terminal egg chambers.

Cells per clone: EC-only lineages
471

We considered only EC-only, and not EC+FC, raw data because some of the latter may derive from
EC/FSC/FC clones that have lost FSCs.
We calculated raw ECs per EC-only clone (and used the same principle for r1 ECs only) and multiplied
by the ratio of raw EC-only clone frequency to calculate single-clone EC-only frequency. Because this
is a derived number, we use raw observations of EC number in EC-only clones to estimate standard
deviation (and then, standard error).

Cells per clone: EC/FSC/(FC) lineages

480
Single-cell EC/FSC clones will all be within this raw data category. This is an inclusive category- most
clones have FCs but those without FCs are also included. However, (a) such clones in raw data will
sometimes also include an EC-only or an FSC-only clone and (b) the correct number of EC/FSC clones
deduced by single-cell lineage estimations should be used.
Single-clone EC/FSC frequency = r
Observed EC/FSC = $r.p1 + r^2.p2 + 2.p2.r.(q+m) + 2.p2.r.s + 2.p2.(q+m).s$
487
488   (a) Subtractions from total counted cells
489   Additional ECs come from EC only clones at (average cells per EC-only clone)(total number of single
490   clones= actual clone number plus deduced number of double-clones)(2p2.(q+m).r)
491   Additional ECs come from EC only clones at (average cells per EC-only clone)(total number of single
492   clones= actual clone number plus deduced number of double-clones)(2p2.s.r)
493
494   (b) Total number of single-cell EC/FSC clones in observed EC/FSC samples
495   = r. total number of single-cell clones in sample
496   = r. total clones in sample (1+p2)
497
498   Because the final estimates are derived numbers, we use raw observations of EC and FSC number in
499   EC/FSC clones to estimate standard deviation (and then, standard error)
500
501   All of the above calculations are in the supplementary spreadsheet that includes all samples for
502   lineages induced at different times: "Single-cell 0hAPF calculations" and "Single-cell -2dAPF
503   calculations".
504
505   **Evaluation of Controls**
506
Expected variability
If we assume that we are examining single-cell clones and that the chance of hitting each type
(location) of cell in a given developing ovariole is independent (and given by the average frequency of
that clone type (p) in a sample with n single clones), we expect to see a binomial distribution of the
frequency of outcome p.
We therefore calculated the "expected" SEM for this process for each average frequency of lineage
category over all 12 controls for lineages induced at pupariation. For binomial distributions the
expected standard deviation= SQRT[np(1-p)]
We also calculated the observed SEM among the 12 individual values for each category.
The ratio of the two values (observed/expected) was mostly in the range of 1-2, consistent with the
random sampling (by recombination events) of the starting A/P location of precursors being a major
contributor to variation.
These data are presented in the supplementary spreadsheet "0h APF Controls Variation".
520
Significant Differences
The significance of differences for the frequency of a specific type of lineage for an experimental
sample versus controls was estimated using Fisher's exact two-tailed test. The data on the
spreadsheets described above was copied (as values) on to separate spreadsheets to add some extra
summary categories. Calculated p values (from an online calculator) were then added when these

were either very low or of importance to acknowledge as not being low (i.e., values above 0.05 were
generally not entered). Altogether, we tested the following types of condition as both reduced or
increased by genetic manipulation- cell division rate, Wnt signaling, JAK-STAT signaling, Yki activity,
and Hedgehog signaling, translating to 10 broad conditions. It may therefore be of special interest to
know when p values are below 0.05 and below 0.005, so we designated those values by "*" and "**"
in graphs. Supplementary spreadsheets with these p values are named "0h APF Significance Fisher"
and "-2d APF Significance Fisher".
533
534
Inferring clone types, FSC and EC numbers in newly-eclosed adults from scoring 2d-old adults
In order to make quantitative conclusions about the developmental process up to eclosion from
scoring clonal outcomes in 2d-old adults we made the following assumptions and adjustments. We
assumed that EC numbers and locations did not change during this 2d interval because ECs generally
appear to be quite long-lived. ECs are, however, produced from FSCs during adulthood, so it is likely
that a small number of marked ECs scored at +2d were produced from FSCs after eclosion. The
proportion of marked ECs arising from FSCs is very likely small based on known rates of FSC
conversion to ECs and the observed high frequency of FSC clones with no ECs (44%) from control -2d
samples. Individual marked FSCs can be lost, mainly by becoming FCs, or amplified at high rates
during adulthood. If one or more marked FSCs were present at 0d but became FCs by +2d those
marked FCs would reside around the two germarial cysts or in the two youngest egg chambers
because egg chambers bud roughly every 12h, allowing four cycles of FC recruitment over 2d. We
therefore scored an ovariole as containing an FSC clone (FSC-only or FSC/EC, according to EC content)
if there were any marked FCs up to and including the second egg chamber, even if there were no
marked FSCs at +2d. In those cases, we scored the number of marked FSCs as zero. On average, the
number of marked FSCs should remain constant from 0d to +2d, so we should obtain a very good
estimation of FSC numbers at 0d by scoring the numbers of marked FSCs at +2d, provided we include
all examples where marked FSCs were lost altogether as containing zero FSCs. By using these
guidelines, we could surmise the exact number of different types of lineage at 0d and the exact
number of constituent ECs and FSCs from data collected by scoring clones in 2d-old adults.
555
556
Dissection and staining of pupal ovaries
Pupae were removed from the vial wall by adding a drop of water and transferred with forceps into a
well of a Corning PYREX glass spot plate (Corning, cat. no. 7220–85) containing either PBS or 4%
paraformaldehyde solution with care taken not to damage or pierce the posterior end of the pupa. If
the dissection was done in fixative solution, it could take no longer than 10 min (ovaries were rocked
in clean fixative for a minimum of 15 min, but for no longer than 30 min). Ovaries from 72h APF and
older had fewer tears in them when dissected directly into fixative. Using a dissecting
microscope, pupae were held against the bottom of the well using forceps and the posterior tip was
cut off using Vannas Spring Scissors (2.5mm straight edge, Fine Science Tools, Foster City, CA, item
number 15000-08) about a third of the way in from the posterior end. The anterior part of the pupa
was removed to a discard well. The posterior end was searched for ovaries (clear, spherical, striated
structures) by gently dislodging fat tissue surrounding the ovaries so as not to cloud the well with
debris. This was done by gently tearing apart the contents of the pupal case or swirling them in the
well using two pairs of forceps. As soon as they were spotted, ovaries were transferred to a new well
of fixative by the following process. A 20µl pipette tip with a pipettor set to 20µl was coated in 10%
Normal Goat Serum (NGS) by pipetting up and down several times. The pipettor was set to 5µl to

pipette up the ovary from the dissection well to transfer into the fixative well. Ovaries were fixed by
rocking in the well for 15 min at room temperature. Fixative (and all liquid in subsequent steps) was
removed in either of two ways: with a 1000µl pipette set to 270µl to slowly and carefully pipette up
the liquid; or, using the same method to transfer ovaries above, with a 20µl pipette tip with a
pipettor set to 20µl, which was coated in 10% Normal Goat Serum (NGS) by pipetting up and down
several times. The pipettor was set to 5µl to pipette up the ovary from the fixative well to transfer
into a fresh well with PBST. Ovaries were rinsed 3x in 2% PBST 0.5% Tween solution for 5, 10, and 45
min at 4C by rocking very slowly in a horizontal plane. The glass dissection dish was covered with a
large pipette tip box. Ovaries were rinsed for 30-60 min in 10% NGS solution at 4C, and then rocked
gently in primary antibody overnight. Ovaries were rinsed 3x with 0.5% PBST and then incubated for
1 hour (covered, cold room, rocker set on 2) in secondary antibody diluted 1:1000 with 0.5% Triton.
Ovaries were rinsed 2x with 0.5% PBST and 1x with PBS. To mount ovaries, a 20µl pipette tip coated
in 10% NGS solution, as described above, was set to 5µl and used to capture the ovary and transfer to
a glass slide. 20-25µl of DAPI Fluoromount was added to a coverslip, and then placed on top of the
slide with the ovaries. Care was taken to avoid pressing the coverslip and potentially disfiguring the
mounted ovaries. In some studies of older pupal ovaries, specifically 72h APF and onwards, forceps
were used to gently tear apart the ovarioles to separate germaria for clearer imaging. A sharpie was
used to draw arrows to mark the location of the ovaries on the coverslip.
591
592   **Staging of pupae (Fz3-RFP experiments)**
593
Third-instar larvae were sorted to select females, transferred to a vial with food and checked every
hour to mark the time each individual developed into a puparium (white, but immobile, with small
anterior spiracles). This time is 0 hours APF (after puparium formation). We found that keeping
animals in the light at night and in the dark during the day on a 12h/12h cycle, allowed more of them
to pupate during the daytime. We used a programmable outlet timer (Nearpow) together with a
manual LED soft white nightlight (Energizer cat. 37099) installed in a dark incubator at 25C.
600
601   **Imaging and quantitation of Fz3-RFP profiles**
602
Fz3-RFP flies (made by Ramanuj DasGupta, Genome Institute of Singapore and a gift from Erika Bach,
NYU, USA) were dissected and stained for Fas3 and DAPI. Ovaries were imaged with a 63 × 1.4 N.A
lens on a Zeiss LSM 800 confocal microscope (Carl Zeiss). Zeiss Zen Blue was used to acquire
microscope images of 1024x1024 pixels at 16-bit depth with line averaging 2. Zeiss Zen was used to
measure the intensity of RFP in cells along an edge of a germarium in a line starting adjacent to Cap
cells and ending at the border of Fas3 expression. Measurements were taken in 5 µm distance
increments within a z-stack range that did not exceed 8 µm. If there was more than one cell within
the same 5 µm increment, both cells would be measured. If there was no suitable cell within the 8
µm z-stack range at a specific 5 µm increment, no measurement was collected. Fz3-RFP intensities
were measured by circling the nucleus of a cell in the DAPI channel using the spline contour tool and
obtaining the average intensity value in the 561nm channel for the enclosed area from the Z-stack
with the highest RFP intensity. The 561nm laser intensity was set at 1.1% for 21-36h APF germaria
and reduced to 0.3% beginning at 48h-APF through adult in order to keep RFP within the linear range.
Because overall RFP levels varied from germarium to germarium, individual RFP profiles were
constructed for each germarium; RFP intensities for each germarium were normalized to the
brightest cell on the edge of that gemarium. Normalized intensities were then averaged for each time
point. In order to reflect the relative RFP brightness from 27h to 36h-APF, the averaged, normalized

values were divided by the laser intensity at which they were taken. The averaged normalized values
for every distance along the 27h-APF and 36-APF curves were divided by 1.1. The averaged
normalized values for every distance along the 48h-APF, 60h-APF, and adult-stage curves were
divided by 0.3. The 27h-APF, 36h-APF, 48h-APF, 60h-APF, and adult were then normalized to the
highest value in all of the graphs to reflect relative intensities and display the gradient profiles.
Because the 21h-36h germaria were necessarily acquired with a higher laser intensity, output levels
in Photoshop were reduced from 255 to 100 to indicate the greatly reduced RFP intensity of those
samples. The distances from Cap cells to the border of strong Fas3 expression on the edge of the
germarium and to the posterior end of the germline in the center of the germarium were also
measured for every germarium and averaged for each time-point.
630
631
632 **Results**
633
634 **Cell lineage methods and analysis**
635
The cell-autonomous effects of genetic alterations on cell fate can generally be studied by
comparing outcomes for marked cells, with or without a genetic alteration, over the relevant
developmental period. Here, we are interested in examining all somatic cells of an ovariole from
pupariation (when mature third instar larvae cease movements) to adulthood. We therefore induced
FRT-catalyzed recombination events to mark and modify any ovarian somatic cell at a chosen
developmental stage by using a heat-shock inducible Flp recombinase transgene. Prior studies using
this approach for wild-type cells suggested that the number and identity of adult somatic cells in a
marked lineage depends principally on the A/P location of the marked precursor at pupariation
(REILEIN *et al.* 2021). Anterior precursors become ECs, posterior precursors become FCs, and
precursors in intermediate locations produce mixed lineages that include ECs, FSCs and FCs (Fig. 1A).
Moreover, posterior precursors divide faster and, unlike EC-only precursors, they do not cease
division during pupation (REILEIN *et al.* 2021). Here, we use equivalent lineage studies with specific
genetic alterations to identify factors that guide patterned cell division, differentiation of FCs during
pupation, and anterior-posterior (A/P) migration of somatic ovarian precursors, which dictates EC and
FSC fates in the adult germarium. There are, however, some significant technical challenges and
limitations.
First, the marked cells in a single adult ovariole ideally derive from a single precursor within a
pool of about two dozen precursors at pupariation. We therefore used a very mild heat-shock
protocol, carefully administered for uniformity, to induce mitotic recombination in a MARCM (Mosaic
Analysis with a Repressible Marker) strategy (LEE AND LUO 2001). In practice, this generally leads to
about 30-50% of all ovarioles with at least one labeled cell (Fig. 1B). Some of these ovarioles will,
however, include derivatives of two or more initially marked cells. We therefore estimated this
fraction from the percentage of unlabeled ovarioles by assuming that all recombination events are
independent, and we then inferred the frequency and composition of single-cell lineages (see
Methods). An example of the results of this conversion of raw data to deduced single-cell lineage
data is shown for a control sample (Fig. 1C). In all other presented data, only the deduced single-cell
lineage data are shown (with sample numbers in figure legends referring to raw data; all raw data
and conversion details are in supplementary data spreadsheets).
664

Figure 1: Estimation of single-cell lineage outcomes and cell numbers in control samples.
(A) Illustration of somatic cells of an ovariole in a newly-eclosed fly. The germarium (anterior, left) produces budded egg chambers (posterior, right), four of which (numbered 1-4) are typically present in each ovariole in a newly-eclosed fly. Escort Cells (ECs) are divided according to their location, as region 1 (r1: magenta) and region 2a (r2a: yellow). FSCs (cyan) reside principally in two A/P rings, just anterior to the anterior border of strong Fas3 cell surface expression ("Fas3 border": red line). FCs (mostly without a filled color), posterior to the Fas3 border, are classified by their location: Immediate (Region 2b, I), Germarium (Region 3, G), egg chamber 1, 2, 3, and 4. Specialized FC sub-types colored are polar cells (dark green), stalk cells (light green) and basal stalk cells (light green). **(B)** MARCM controls from 12 experiments, analyzed in newly-eclosed flies after 0h APF clone induction. Colors indicate the recombinant FRT site: 40A (yellow), 42D (blue), 82B (red). Black is the non-weighted average of all control values. The first cluster reports the percentage of all ovarioles with any marked cells. The number of marked ovarioles ranged from 36 to 121 (36, 41, 51, 54, 58, 56, 48, 121, 62, 98, 49, 65, respectively) and the total number of ovarioles from 102 to 277 (128, 112, 135, 135, 140, 134, 111, 277, 136, 207, 102, 119, respectively). Other clusters report the deduced frequency of the named lineage categories among marked lineages after estimating single-cell lineage frequencies (see methods). The number of FSC-containing lineages without ECs is generally very low and is therefore combined with lineages containing ECs and FSCs (written as (EC)/FSC); almost all of these lineages also contain marked FCs. Ovarioles with ECs and FCs but no FSCs are thought mostly to contain two lineages. The EC+FC category therefore gives an indication of the limitations of the method used to convert raw data to estimates of single-cell lineages. Here, the (EC+FC) and FC-only frequencies have also been added to give the

(EC)/FC category. The final two clusters show the percentage of all lineages that includes at least one marked r1 EC or
terminal FC. **(C)** An individual control from (**B**) (control C, third control) to show raw values (blue) compared to deduced
single-cell lineage values (orange). **(D, D')** Deduced average number of **(D)** cells per EC-only clone (total cell number: 54,
68, 103, 103, 116, 80, 85, 195, 110, 107, 106, 87, respectively) and **(D')** EC plus FSC cells in EC/FSC clones (total EC + FSC
cell number: 40, 24, 79, 56, 71, 116, 12, 266, 58, 194, 85, 281, respectively; the number of EC/FSC lineages for samples B
and G are notably low) for all MARCM controls analyzed in newly-eclosed flies after 0h APF clone induction. (**B-D**) SEMs
shown. For raw data and calculations please see supplementary spreadsheets titled 0h APF Graphs Fig1,3; 0h APF Controls
Variation; Compilation of Numbers Figure 1 Controls.
693
Second, although the chosen approach can simultaneously report on the behavior of all
precursors in a variety of AP locations, it necessarily includes a sampling error. For example, if we are
examining the fate of the roughly 20% of precursors that would normally become FCs, the sample of
such cells will not always be exactly 20%. We have performed many sets of experiments, each
including a control ("wild-type") genotype. Those multiple controls allow experimental evaluation of
sampling variability. They also provide a cumulatively large control sample to produce reliable
measures for each parameter deduced from lineage studies. For MARCM clones induced at
pupariation, the standard deviation among controls for the frequency of a given lineage type (Fig. 1B)
was generally similar to that expected for random sampling of individual cells from the estimated
number of somatic precursors present at pupariation (see Methods and cited supplementary
spreadsheet).
Third, clone sizes provide only limited insight into the spatial regulation of division rates. First,
the deduced average number of marked ECs per EC-only clone was consistently between 2 and 3 for
controls, consistent with limited division of the most anterior precursors and providing little
opportunity to measure significant quantitative deficits (Fig. 1D). Second, almost all lineages with
FSCs also include ECs and FCs. The number of ECs and FSCs in such lineages (average of about 8 for
controls) (Fig. 1D') provides some indication of the amplification of a subset of the precursors, but
this excludes the large and variable numbers of FCs generally present in such lineages. It is not
possible to measure the number of marked FCs that first associate with a germline cyst (which would
be the key number to measure divisions in and around the prospective FSC domain) because each
such cell generally divides many further times as the egg chamber forms and grows. Thus, the sum of
ECs and FSCs in an FSC-containing lineage only provides limited information about division rates in
regions that give rise to ECs, FSCs and FCs. The fraction of cysts with marked FCs in FSC-containing
lineages provides a second measure. The number of marked cells in the most posterior egg chamber
can report on proliferation of the most posterior precursors. Additional information comes from
measuring the frequency with which such lineages extend into the penultimate egg chamber.
Fourth, the MARCM method has the key virtue of clear positive marking, so that every lineage
(even a single cell) and every marked cell within a lineage can be identified readily. A well-known
disadvantage is, however, that GAL80 gene products must first be diluted and degraded before GAL4-
driven *UAS-X* transgene products can accumulate. Strong accumulation of GFP from *UAS-GFP* in
somatic ovarian cells takes at least 3d using our genetic reagents, and a similar delay is therefore
expected before any *UAS*-driven transgene begins to exert maximal effect. Thus, both the onset and
plateau of phenotypic responses to activation of a *UAS*-driven transgene in a MARCM clone can only
be estimated and will generally be later than for the generation of a loss-of-function homozygous
mutation. These and other relevant methodological considerations are discussed at greater length in
the context of specific results obtained for variant genotypes.
730
731

Terminal FC source is not distinct from other EC, FSC and FC precursors

We are interested in understanding how somatic cells are instructed to occupy various positions in the adult germarium to function subsequently as FSCs and ECs. We are also especially interested in how the first FCs are specified during pupation. This is an intriguing question because FC formation in adults relies on signals from previously produced FCs. We therefore first sought to clarify the origins of the FCs on the first-formed egg chamber.

At the start of pupation, about 20-30 somatic cells are dispersed among germline cells to form a primitive germarium (Fig. 2A). These intermingled cells (ICs) express the transcription factor, Traffic Jam (TJ) (Li *et al.* 2003). Numerous additional TJ-negative somatic cells, referred to as basal cells (Swarm cells after migration; not shown in Fig. 2A and not the same as basal stalk cells), are present in a more posterior location, as a seemingly contiguous group, rather than associated with individual developing ovarioles. Over the next 48h, TJ-positive somatic cells accumulate posterior to the developing germarium (beyond the most posterior germline cyst) to form a structure termed the Extra-Germarial Crown (EGC) (Reilein *et al.* 2021). In the most posterior portion of the EGC, somatic cells are intercalated to form a "basal stalk", which also includes some TJ-negative cells at the posterior (King *et al.* 1968; Godt and Laski 1995; Vlachos *et al.* 2015; Reilein *et al.* 2021). Thereafter (at about 56h APF on average), the most posterior germline cyst moves away from the germarium, into the EGC and basal stalk, as cells in that extra-germarial territory, including some originally TJ-negative cells, move anteriorly along the cyst to form a monolayer epithelium, leaving a short residue of basal stalk at the posterior end that persists until eclosion (Reilein *et al.* 2021). Most of the mid-pupal (around 48h APF) EGC and basal stalk cells become FCs (or adult basal stalk cells) on the first budded (terminal) egg chamber but some contribute to the next (penultimate) egg chamber, alongside FCs derived from IC precursors present within the germarium around 48h APF (Fig. 2A). These deductions derive from fixed images at successive stages, live imaging cell tracing and lineage analyses initiated by a heat-shock at 36h APF (Reilein *et al.* 2021). Many such 36h APF lineages contained only marked FCs or basal stalk cells on the terminal egg chamber (purple lineage in Fig. 2A), while other lineages had marked FCs only in more anterior locations (cyan and yellow lineages in Fig. 2A). Each type of FC-only lineage sometimes included marked FCs in the penultimate egg chamber. Thus, at 36-48h APF, only precursors in the EGC or basal stalk contribute terminal FCs and only IC precursors yield FCs anterior to the two terminal egg chambers.

The remaining question is whether the precursors of the first FCs (EGC and basal stalk cells) emerge from germarial ICs over the first 48h of pupation. This hypothetical migration out of the developing germarium has not been visualized directly. There is extensive division outside the germarium, so there may be only a modest ongoing exodus from the germarium that escaped detection during live imaging. An alternative possibility is that the EGC and basal stalk cells seen at 48h APF have a separate origin from IC cells, conceivably deriving from extra-germarial basal cells present at pupariation (King *et al.* 1968) and acquiring TJ expression over the following days. These possibilities can be distinguished by lineage analysis. An FC-only lineage (induced prior to 36h APF) that includes FCs in both the terminal egg chamber ("terminal FCs") and at least one location anterior to the penultimate egg chamber ("non-terminal" FCs) would provide evidence of a common precursor of ICs and EGC/basal stalk cells present before 36h APF.

Figure 2: Terminal and non-terminal Follicle Cells derive from germarial ICs at pupariation.
(A) Illustration of selected individual precursors at pupariation (0h APF) and their progression over time. At 0h APF, the somatic cells are interspersed with single germline cells and 2-cell cysts. By 48h APF, the most mature germline cysts have 16 cells and a single such cyst occupies the most posterior position in the germarium. At this time, more posterior somatic cells express Fas3 (anterior border of expression indicated by the red line). The relationships between precursors at 48h APF and adult derivatives have previously been established. Specifically, precursors within the Extra-Germarial Crown (EGC) and Basal Stalk, which contain no germline cells and are therefore considered extra-germarial, give rise to FCs in the terminal egg chamber or adult basal stalk (with some derivatives extending into the penultimate egg chamber but no

further anterior). Precursors within the germarium at 48h APF give rise to adult ECs only (orange), ECs, FSCs and non-
terminal FCs (yellow) or non-terminal FCs only (blue). It was not known whether cells in the EGC and basal stalk migrated
out of the germarium between 0 and 48h APF (as indicated by the colors in the diagram) or if they derived from
precursors in another location. The existence of precursors at pupariation producing adult FCs in terminal and non-
terminal locations (as indicated for green cells) would support a germarial origin of terminal FCs. It would still be expected
that some precursors at 0h APF would give rise only to terminal FCs (as shown for purple cells). The outcomes shown for
green and purple cells were found, as illustrated in (b) and (D). **(B-G)** Representative images of lineages induced at 0h APF
in ovarioles of newly-eclosed flies. **(B)** Terminal FC-only; **(C)** non-Terminal FCs only (egg chambers 2 and 3); **(D)** non-
Terminal and Terminal FCs, showing that all FCs derive from germarial cells at pupariation; **(E)** EC/FSC/FC (designated as
EC/FSC in graphs), **(F)** EC-only, **(G)** EC plus Terminal FC (presumed to be two lineages in the same ovariole because
lineages derived from a single cell generally occupy largely contiguous territory). Scale Bar 50 μm for all images. White
arrowheads indicate marked Follicle Cells; magenta arrows, r1 ECs; yellow arrows, r2a ECs; cyan arrows, FSCs. DAPI, blue;
GFP, green; Fas3, red.
800
801
From 12 separate control experiments for lineages initiated at pupariation, the average
fraction of FC-only lineages with terminal FCs (n=101) that also included marked FCs anterior to the
penultimate egg chamber was (0.253/0.685) 0.37. This argues strongly for a single cell often acting as
a precursor to both terminal and non-terminal FCs. Such precursors, like other non-terminal FC
precursors, are therefore within or contiguous with the IC population, meaning that EGC and basal
stalk cells derive from posterior migration of ICs out of a primitive germarium. Thus, it appears that
some precursors in the germarium at pupariation produce only terminal FCs (Fig. 2B; purple cell in
Fig. 2A), others produce only non-terminal FCs (Fig. 2C; blue cell in Fig. 2A), while several produce
both terminal and non-terminal FCs (Fig. 2D; green cell in Fig. 2A). More anterior precursors produce
EC/FSC/FC lineages (Fig. 2E; yellow cell in Fig. 2A) or only ECs (Fig. 2F; orange cell in Fig. 2A).
It is possible that our method for estimating single-cell lineage composition from raw
observations underestimates the frequency of "double-clones" in a single ovariole (see Methods). To
estimate the potential impact of an underestimation of this type we looked at the average frequency
of lineages with terminal FCs that also include ECs (Fig. 2G). This incidence (0.039/(0.102 + 0.039)=
0.26) was previously considered mostly due to the presence of two lineages. To estimate the
frequency of a second lineage including non-terminal FCs rather than ECs (which are much more
frequently marked) due to "double-clones", we multiply by the relative frequency of the latter two
lineages ((0.154-0.064)/0.644) = 0.14). This gives an expected incidence of (0.26 x 0.14) = 0.036,
which is far below the observed incidence of 0.37. In fact, further studies reported here suggest that
some lineages containing only marked ECs and FCs are potential EC/FSC/FC lineages that have lost
FSCs rather than indicating two superimposed lineages. Hence, the incidence of "double-clones" in
our deduced single-cell lineages is likely even lower than estimated above. We can therefore be
confident that ICs and extra-germarial somatic cells at 36-48h APF have a common origin at
pupariation, with the terminal FC precursors migrating out of the nascent germarium. This means
that we can use altered genotypes in lineage analyses to look for differences in the production of
terminal FCs in order to identify factors that regulate posterior migration out of the developing
germarium over the first 48h of pupation. The lineage studies provide evidence that the migration
must occur prior to 48h APF but do not define when the migration occurs. There are some Traffic-
Jam-positive cells posterior to all germ cells even by 12h APF (Reilein *et al.* 2021), so it is likely that
some migration is early.
832
833
834

Consequences of altered precursor division rate

In an adult ovariole, the division rate of a marked FSC over a period of several days cannot simply be assessed by counting the number of marked derivatives, principally because FC derivatives are initially highly proliferative, so that the number of FCs directly produced from marked FSCs cannot be counted. The analogous limitation applies to any pupal precursors that give rise to a mixture of FCs and other cell types during pupation. The number of divisions for EC-only lineages can be measured, but the average size of control EC-only clones is only 2.7 (Figs. 1D and 3I'), providing only a small range for detecting reduced division rates. The most posterior somatic precursors at pupariation contribute principally to FCs of the first-formed egg chamber (purple cell in Fig. 2A). The division frequency of a posterior precursor can therefore theoretically be estimated as the proportion of FCs in the terminal egg chamber that are marked (ignoring lineages that only mark polar cell and stalk cells, which cease division soon after specification).

We wanted first to establish whether changes in division rate can be measured in pupal ovary precursors in various locations and then to determine whether altering the rate of cell division has a systematic effect on developmental outcomes. We have previously shown that a specific hypomorphic *cycE* allele (*cycE^WX*) reduces the division rate of adult FSCs without affecting differentiation significantly, and that this leads to loss of marked FSCs over time (WANG AND KALDERON 2009; REILEIN *et al.* 2018). The reduced competitiveness of slowly dividing FSCs is an important characteristic that results from division and differentiation being independent processes for these stem cells (as in some other population asymmetry paradigms) (VERMEULEN AND SNIPPERT 2014; REILEIN *et al.* 2018). Essentially, this is because the marked *cycE* mutant stem cells drain from the stem cell domain at the normal rate through differentiation but are replenished by division at a lower rate.

We found that the average number of marked ECs per EC-only clone for *cycE* induced at 0h APF was 1.9, compared to 2.7 for controls (Fig. 3I'). This is consistent with reduced division of the most anterior somatic ovarian precursors but likely does not report the magnitude of the effect well because wild-type *cycE* products likely perdure for the short duration of pre-EC divisions. The most posterior somatic precursors at pupariation contribute principally to FCs of the first-formed egg chamber (purple cell in Fig. 2A). The division frequency of a posterior precursor can therefore theoretically be estimated as the proportion of FCs in the terminal egg chamber that are marked. The average fraction of marked FCs among terminal FCs with main-body FC labeling was 3.0% for *cycE*, compared to nearly 50% for the direct control (Fig. 3A, B). This measure might be distorted if the altered genotype affects the likelihood of precursors becoming stalk or polar cells, which proliferate less than other FCs. However, marked stalk or polar cells were no more prevalent for *cycE* (2/14 samples) than for controls (5/10 samples). Thus, it appears that reduced CycE activity associated with the *cycE^WX* allele greatly impaired cell division rates for posterior ovarian precursors.

Reduced precursor division frequency might be anticipated to have different consequences on lineage outcomes depending on precursor location. First, we might hypothesize that slowly-dividing *cycE* cells in regions giving rise to future FSCs will be outcompeted by normal precursors, depleting marked FSCs, exactly as seen for adult-induced FSC lineages. The logic here is that since there is a certain A/P domain where precursors reside to become FSCs, the survival of a lineage in this domain requires cell division to balance posterior loss. When induced at pupariation and analyzed in newly-eclosed adult ovaries, there was a large reduction in the frequency of *cycE* clones that included one or more FSCs (2% vs 15% for controls; p<0.005) (Fig. 3I). Out of all marked ECs and FSCs, only 1% of *cycE* cells were FSCs vs 23% of control cells, confirming the substantial deficiency in FSC production. The frequency of ovarioles harboring marked cells of any type was not significantly

altered (47% *cycE*, 42% in the direct, side-by-side control), suggesting that the *cycE* mutation does not impact cell survival and is therefore unlikely to be depleting FSCs through selective cell death. The results therefore support the hypothesis that severely reduced cell division substantially reduces the probability of a cell emerging from pupal development as an FSC. Selective loss of FSCs from prospective EC/FSC/FC lineages would be expected to result in an unusually high frequency of lineages containing only marked ECs and FCs. This was observed for *cycE* lineages (Fig. 3G, I; p<0.005).

Figure 3: Effects of altered division frequency on lineage outcomes.
(A-F) Fractional occupancy of terminal egg chamber by marked FCs, when present, as an approximate measure of division frequency for posterior precursors. **(A-C)** show lineages induced at pupariation and **(D-F)** are lineages induced 48h prior to

puapariation for the indicated genotypes. **(F)** Percentage of marked cells out of total cells in terminal egg chamber for lineages of the indicated genotypes induced 48h before puapariation. Direct controls precede other genotypes for each experiment. **(G, H)** Ovarioles with **(G)** a *cycE* EC/FC lineage induced at 0h APF, illustrating the increased frequency of such lineages, potentially due to selective loss of FSCs from EC/FSC/FC lineages and **(H)** a *UAS-CycE* EC/FSC/FC lineage, present at elevated frequency, induced 48h before puapariation. White arrowheads indicate Follicle Cells; magenta arrows, r1 ECs; yellow arrows, r2a ECs; cyan arrows, FSCs. DAPI, blue; GFP, green; Fas3, red. Scale Bar **(A-E)** 50 µm and **(G-H)** 20 µm). **(I, J)** Deduced frequency of the indicated types of single-cell lineage as a percentage of all lineages and **(I', J')** average number of cells per EC-only or EC/FSC lineage (ECs and FSCs only) induced at **(I)** 0h APF and **(J)** -48h APF. Error bars show SEM. One asterisk (*) indicates p<0.05; two asterisks (**) indicates p<0.005 (Fisher's two-tailed exact test). **(I)** Number of marked ovarioles = 739 (control), 50 (*cycE*), 60 (*cutlet*), 78 (*cutlet; UAS-CycE*). **(J)** Number of marked ovarioles = 319 (control), 93 (*cycE*), 119 (*cutlet*), 85 (*UAS-CycE*), 75 (*cutlet; UAS-CycE*). **(I,' J')** Number of lineages indicated above bars. For raw data and calculations please see supplementary spreadsheets titled 0h APF Graphs Fig1,3; -2d APF Graphs; FC occupancy aggregate**;** Compilation of Numbers Figure 3 Division -5d; Compilation of Numbers Figure 3 Division -7d.

A second hypothesis is that reduced division rate may result in reduced posterior movement of precursors. The somatic cell population grows and expands posteriorly over the first 48h of pupation to occupy a wider A/P territory prior to the budding of the first egg chamber. While it is more likely that posterior expansion is fueled by pressure from growth and division of germline cysts and somatic cells that lie anterior, it is possible that the division of a cell has some cell-autonomous contribution to its posterior movement. If such an effect were significant, we might expect that *cycE* mutant cells would less frequently contribute to FCs of the terminal egg chamber and would more frequently become ECs, especially region 1 ECs. Contrary to that hypothesis, the fraction of lineages including terminal FCs was higher (32% vs 22%; p= 0.09) and the fraction with r1 ECs was lower (53% vs 67%; p<0.05) for *cycE* lineages compared to controls (Fig. 3I). The frequency of EC-only lineages was also reduced for *cycE* (43% vs 63%; p<0.005), while the frequency of FC-only lineages was increased (32% vs 17%; p<0.05) (Fig. 3I). These results therefore suggest an unexpected cell-autonomous posterior movement of *cycE* mutant cells relative to surrounding precursors over the whole A/P range.

To test whether the posterior movement of *cycE* mutant precursors was due to reduced division or another action of CycE we tested an allele of the replication factor gene *cutlet*, which reduced division rate in adult FSCs (WANG *et al.* 2012). The mean number of marked ECs per EC-only was 2.4 compared to 2.7 for controls (Fig. 3I'), while the average fraction of marked FCs among terminal FCs was 7.4%, compared to 24.3% for the direct control (Fig. 3C). These results are consistent with reduced cell division rates for *cutlet* mutant cells, albeit to a lesser degree than observed for the *cycE^WX* allele. The fraction of lineages with marked FSCs was 7%, compared to 15% for controls (Fig. 3I). EC-only clone frequency increased slightly relative to controls (70% vs 63%) but the frequencies of FC-only lineages (17% vs 17%), labeling of r1 ECs (66% vs 67%) and of terminal FCs (19% vs 22%) were largely unchanged (Fig. 3I). Thus, there was selective loss of *cutlet* mutant FSCs without any significant A/P bias of precursor fates. Loss of FSCs for both *cycE* (15% to 2%) and *cutlet* (15% to 7%) is consistent with the conclusion that reduced cell division rates lower FSC production. By contrast, a large deficit of r1 EC-only lineages, an increase in FC-only lineages far exceeding that expected just from FSC loss, and an increase in terminal FCs were observed only for *cycE,* suggesting that these measures of net posterior movement of precursors do not result from a reduced division rate.

We then assessed the effects of overexpression of CycE, using *UAS-CycE* expression with *tub-GAL4* plus *act-GAL4* as drivers of expression (our standard MARCM condition) in marked lineages induced 48h before puapariation. The timing was chosen so that increased CycE levels might start to accrue roughly coincident with puapariation. In adult FSC lineage studies, *UAS-CycE* increased the

division rate of FSCs but not ECs, which are quiescent in adulthood (MELAMED AND KALDERON 2020;
MELAMED *et al.* 2023). EC precursors are still dividing early in pupal development, so they might be
stimulated to divide more frequently if excess CycE is provided at the start of pupation. Excess CycE
would certainly be expected to increase the division rate of more posterior somatic cell precursors.
Hence, by the converse of the logic explained earlier for reduced CycE, faster division might increase
the representation of marked FSCs. The number of marked cells per EC-only clone was higher for
UAS-CycE (5.3) than for controls (3.2) (Fig. 3J'). The average fraction of marked FCs among terminal
FCs was also increased, from 45% to 55% (Fig. 3C). Among EC/FSC/FC lineages (where FC production
cannot be counted, limiting the accuracy of estimating division rates), the average number of marked
germarial cells (ECs and FSCs) rose from 14.5 (including 9.5 ECs) for controls to 24.8 (including 17.4
ECs) for *UAS-CycE* (Fig. 3J'). Thus, *UAS-CycE* increased the division rate of precursors in all locations.
UAS-CycE produced a slightly higher clone rate (frequency of ovarioles with marked cells) than the
side-by-side control (65% vs. 56%), demonstrating that excess CycE does not reduce cell survival.
Importantly, *UAS-CycE* lineages contained FSCs (40%) more frequently than controls (23% for direct
control; 22% for average controls; p<0.005) (Fig. 3H, J). These results support the hypothesis that
increased division rate during pupal stages can increase FSC allocation, just as faster division
increases FSC representation in adult FSC lineages.
The observed increased representation of EC/FSC/FC lineages is expected to be because
potential FSCs are less frequently lost to yield lineages with just ECs and FCs. Indeed, the frequency of
(EC+FC) lineages was lower for UAS-CycE (4% vs 11% controls for lineages induced 48h prior to
pupariation) and higher for reduced *cycE* activity induced 48h prior to pupation (17% vs 11%) or at
pupariation (24% vs 6%). These results suggest that a wild-type precursor in a location that can yield
ECs, FSCs and FCs because of modest A/P spread during pupal development will occasionally contain
only adult ECs and FCs because cells in prospective FSC locations were not replenished fast enough to
counter continuous drain to more posterior FC locations. That, in turn, suggests that a significant
fraction of (EC+FC) lineages result from single precursors, rather than the super-position of an EC-only
and an FC-only lineage, as previously thought (REILEIN *et al.* 2021) and that our estimation of single-
cell lineages numbers derived from raw data is better than we had believed on the basis of the
deduced (EC+FC) lineage frequency.
The frequencies of both EC-only (43% vs 55%; p=0.1) and r1-only lineages (24% vs 43%;
p<0.005) were decreased for *UAS-CycE*. This could simply be because the increased number of cells
per lineage inevitably results in several clones crossing posterior boundaries, into r2a EC and FSC
territory, respectively. The frequency of FC-only lineages was unaltered (14% vs 13%), suggesting
there is no large systematic displacement of all marked cells towards the posterior. Rather, the
principal response was that the representation of a central population, which become FSCs, was
increased.
Since *UAS-CycE* was found to restore survival of adult FSC lineages with *cutlet* mutations
(albeit assessed prior to more definitive measures of FSC competition introduced in 2017) (WANG *et*
al. 2012), we tested whether excess CycE might ameliorate the key phenotypes of *cutlet* mutant
lineages induced 48h prior to pupariation (deficiency of FSCs and indications of reduced division rate).
UAS-CycE increased the average number of number of cells in EC-only lineages from 2.5 to the control
value of 3.2 (Fig. 3J'). The average fraction of marked FCs among terminal FCs was greatly increased,
from 2% to 17%, indicating substantial rescue of division rate towards control values (Fig. 3C). The
frequency of lineages with FSCs was also partially restored (from 2% to 12%) towards the control
value (22%) (Fig. 3D, E, J). In the EC/FSC lineages there were now similar numbers of ECs (8 vs 9.5) and
FSCs (3.5 vs 5.0) to controls (Fig. 3J', sum shown). The FC productivity of FSC-containing lineages can

be roughly estimated by counting the proportion of egg chambers, region 2b and region 3a germline
cysts with associated marked FCs; this measure would be increased by recruitment of more future
FSCs or a higher division rate of those cells. The low proportion of cysts with marked FCs for *cutle*t
(1/11; 9%) was increased by *UAS-CycE* (33/54; 61%) towards the control value for this experiment
(128/156; 82%). These results are all consistent with substantial correlated restoration of cell division
rate and FSC production. A smaller restoration of FSC-containing lineages (from 7% to 10%) towards
control values (15%) was seen for the addition of *UAS-CycE* to *cutlet* loss of function in lineages
initiated at pupariation (Fig. 3I). The proportion of cysts with surrounding marked FCs in FSC-
containing lineages was also increased from 14% (3/21) for *cutlet* to 46% (25/54) by the addition of
UAS-CycE, towards the control value of 65% (26/40). The delayed expression of excess CycE in
lineages initiated at pupariation likely accounts for the lesser rescue.
Reduced CycE activity was also tested in lineages induced 2d prior to pupariation (Fig. 3J). The
major phenotype was a large loss of lineages with FSCs, similar to *cutlet*. There was also an increase
in FC-only and EC/FC lineages combined ("(EC)/FC") for *cycE* (38%) relative to *cutlet* (22%) and
controls (23%), similar to results for lineages induced at pupariation. However, there was neither an
increase for terminal FC representation (27% vs control 35%), nor a large reduction in r1 EC
representation (80% vs 75%) seen for lineages initiated at pupariation. Thus, the strong phenotype of
FSC loss was seen for both stages of lineage induction, while the unexpected posterior movement of
precursors due to loss of *cycE* was prominent only for lineages induced at pupariation.
This set of studies suggests that the primary consequence of cell-autonomously reducing the
rate of precursor division within a competitive environment of normal cells during pupation is a large
reduction in the frequency of lineages containing FSCs in newly-eclosed adults. This was observed for
both loss of *cycE* and loss of *cutlet*, initiating lineages at pupariation and 2d earlier. The converse was
seen in response to increasing the division rate. No other consequence was consistently seen for both
cycE and *cutlet* loss of function mutations, while excess CycE also appeared only to affect FSC
representation.
1012
1013 **Wnt signaling magnitude strongly influences precursor fates across the whole A/P domain**
1014
We wished to investigate the roles of Wnt and JAK-STAT signaling on somatic ovarian
precursor development because these are the two major pathways regulating adult FSC behavior
(VIED *et al.* 2012; REILEIN *et al.* 2017; MELAMED AND KALDERON 2020). Canonical Wnt signaling is initiated
by binding of ligand to Frizzled and LRP5/6 (Arrow in Drosophila) family co-receptors, leading to
release of β-catenin from a destruction complex scaffolded by Axin (Axn) and its subsequent
association with nuclear TCF/LEF (TCF in Drosophila) to convert TCF from a repressor to an activator
of transcription (WIESE *et al.* 2018).
We used the strong *axn^{E77}* allele (which truncates the protein after Q406 to eliminate DIX,
GSK3, β-catenin and protein phosphatase 2A binding domains to behave like a null in several assays)
to create marked homozygous *axn* mutant precursors at pupariation and measure the spectrum of
lineages obtained consequent to increasing Wnt pathway activity cell autonomously during pupal
development. Almost all lineages consisted of only marked ECs (91%, compared to 63% in controls)
(Fig. 4A, D, F), with an elevated yield of lineages confined to r1 ECs (61% vs 49% control) and of
lineages containing at least one marked r1 EC (81% vs 67% control). Only 2% of lineages contained
marked FCs without FSCs (22% for controls) and only 6% contained any marked FSCs (15% for
controls) (Fig. 4F). Thus, it appears that all marked precursors have preferentially moved anteriorly
during pupal development in a cell-autonomous response to increased Wnt pathway activity, leading

to a near-absence of marked posterior derivatives, FSCs and FCs. In adult FSCs, loss of *axn* leads to
strong net anterior movement, producing a high frequency of marked ECs over time but it also
severely reduces cell division (REILEIN *et al.* 2017; MELAMED AND KALDERON 2020). Based on *cycE* and
cutlet mutant phenotypes over pupal development, reduced division would be expected to reduce
the yield of FSCs but some of the "lost" FSCs would become FCs. The drastic loss of both FSCs and FCs
for *axn* mutant precursors argues for a strongly altered behavior (anterior migration) other than
reduced division. The average number of cells per EC clone was slightly lower for *axn* (2.5) than the
control value (2.7). However, this is not a sensitive test and the *axn* mutant ECs, on average, also
occupy a slightly more anterior set of EC positions than control derivatives, which may lead to earlier
cessation of division. Since almost all *axn* precursors become ECs, there were too few FSC and FC *axn*
derivatives to assess cell division rates of more posterior precursors. Thus, it is not clear whether
increased Wnt signaling directly affects cell division during pupation but there is a strong phenotype
of anterior bias for adult derivatives of precursors in all locations.
1045   To investigate the effects of loss of Wnt pathway activity we used a strong or null *arrow* allele
(*arr²*) that introduces a premature stop codon. We also added *UAS-DIAP1*, which was found in adult
FSC derivatives to increase the yield of labeled *arr* mutant ECs (MELAMED AND KALDERON 2020). Other
studies have found that adult ECs, especially those at the far anterior, are subject to apoptosis when
Wnt pathway activity is severely reduced (WANG *et al.* 2015; WANG AND PAGE-MCCAW 2018). Relative to
controls, *arr* mutant lineages induced at pupariation (with *UAS-DIAP1*) had a strong deficit of lineages
with only r1 ECs (23% vs 49% control), only ECs (34% vs 63% control) or any r1 EC (36% vs 67%
control) (Fig. 4B, C, F). By contrast, there was an increased fraction of lineages with marked FSCs (27%
vs 15% control), only FCs (33% vs 17%) or containing terminal FCs (41% vs 22% control). The average
number of marked germarial cells in each EC-only (2.1 vs 2.7) or EC/FSC/FC lineage (7.4 vs 8.6) was
largely unchanged. Thus, as in adult FSC lineages, loss of Wnt pathway activity appears to have no
major effect on the rate of cell division but causes a strong posterior shift in location, leading to a
deficit of ECs and an excess of FCs. The increase in lineages with FSCs can be rationalized as due to a
greater numerical shift of would-be ECs (which are the most numerous) to FSCs than of would-be
FSCs to FCs. These conclusions explain the unexpected observation long ago that loss of *arr* increased
FSC clonal yield if induced at larval stages and assayed 10 days later, despite reducing FSC longevity in
adult FSC lineages because of a posterior shift increasing conversion of FSCs to FCs (VIED *et al.* 2012).
In other words, the same primary consequence of preferential posterior movement due to
autonomous loss of Wnt signaling reduces FSC yield in adults because marked FSCs cannot be
replenished from marked ECs, but increases FSC yield during pupation because marked EC precursors
can move into FSC locations; in both situations there is faster depletion of marked FSCs posteriorly to
become FCs.
1067   The effects of altered Wnt pathway activity on the conversion of precursors to adult ECs and
FSCs can also be evaluated by counting the total number of these derivatives over all examined
lineages (FC numbers cannot be evaluated in this way). This analysis has the virtue that it does not
rely on whether marked cells in an ovariole derive from one or more precursors. In line with the
analysis of lineage types above, the proportion of derivatives that became FSCs (control 20%) was
greatly reduced for *axn* clones (6%) and markedly increased for *arr* clones (35%), while the proportion
that became r1 ECs (control 54%) was similar for *axn* clones (50%) and markedly decreased for *arr*
clones (31%). The unchanged proportion of r1 ECs for *axn* is somewhat deceptive because FCs are not
included in this analysis. Control samples include many FCs whereas *axn* lineages do not. Hence, r1 EC
frequency as a proportion of all marked cells (including FCs) is actually lower for controls than for *axn*.

Figure 4: Higher Wnt pathway activity promotes anterior precursor movement during pupation.

(A-E) Representative lineages induced at **(A-D)** 0h APF or **(E)** 36h APF for the indicated genotypes. White arrowheads indicate Follicle Cells; magenta arrows, r1 ECs; yellow arrows, r2a ECs; cyan arrows, FSCs. DAPI, blue; GFP, green; Fas3, red. **(A-C)** The r1 EC, r2a EC, FSC and FC domains are shown in corresponding colors in the horizontal bars above. Characteristic phenotypes with elevated frequency included **(A)** an *axn* r1 EC-only lineage and **(B, C)** *arr; UAS-DIAP1* lineages containing FSCs and FCs, **(B)** with or **(C)** without ECs (the latter is rarely observed for control lineages). **(D, E)** Ovarioles with an *axn* lineage induced at **(D)** 0h APF, with only ECs, and at **(E)** 36h APF, with marked terminal FCs in the basal stalk (arrowhead). Scale Bar 20 μm for **(A-C)** and 50 μm for **(D, E)**. **(F)** Deduced frequency of the indicated types of single-cell lineage as a percentage of all lineages induced at 0h APF. Error bars show SEM. One asterisk (*) indicates p<0.05; two asterisks (**) indicates p<0.005 (Fisher's two-tailed exact test). Number of marked ovarioles = 739 (control), 44 (*axn*), 95 (*arr; UAS-DIAP1*). For raw data and calculations please see supplementary spreadsheets titled 0h APF Graphs Fig4-10; Compilation of Numbers Figure 4 Wnt.

Wnt pathway activity limits cell migration out of the germarium during the first half of pupation

We have shown that EGC and basal stalk cells, the progenitors of terminal egg chamber FCs, derive from precursors that migrate posteriorly out of the germarium over the first 48h of pupation, and that increased Wnt signaling throughout pupation strongly reduces the acquisition of terminal FC identity cell-autonomously (Fig. 4F). To refine the key period of action, we induced control and *axn* lineages at 36h APF, estimating that genetically increased Wnt signaling would be delayed by a further few hours to coincide roughly with the completion of EGC/basal stalk formation and the migration of the most posterior germline cyst out of the germarium. At 36h APF there are many more somatic cell precursors per ovariole than at pupariation, so even the mildest heat-shock conditions produce MARCM clones in almost all ovarioles. Consequently, most ovarioles will contain more than one lineage and it is not possible to derive single-cell lineage frequencies accurately; raw

observations are therefore presented for that time-point. The clone frequency for *axn* (97%) was
higher than for the direct control (58%), so that multiple lineages per ovariole are likely more
frequent. Nevertheless, there was clearly no deficit in the fraction of marked ovarioles with labeled
terminal FCs (46% vs 31% control) (Fig. 4E; Supplementary spreadsheet "-3.5d axn August 2023
Numbers"). This is markedly different to the result for lineages initiated at pupariation, indicating that
the near absence of terminal FCs for those samples is caused by failure of precursors to migrate
posteriorly out of the germarium over the first 48h of pupation, with no subsequent deficit in
surviving as terminal FCs.
The frequency of lineages that included terminal FCs was much higher for *arr* (41%) than for
control lineages (22%) when induced at pupariation, indicating that Wnt signaling normally limits
terminal FC formation (Fig. 4F). When lineages were induced 24h APF there was a similar clone
frequency for control (84%) and *arr* plus *UAS-DIAP1* (73%) and a similar proportion of ovarioles that
included marked terminal FCs (29% control vs 30%; Supplementary spreadsheet "-4d arr DIAP
Numbers"). Thus, the increased representation of terminal FCs seen for lineages induced at
pupariation is absent 24h later. We deduce that normal Wnt signaling restrains posterior movement
of precursors out of the germarium at the start of pupation. These consequences of elevated and
reduced Wnt signaling on generation of the first FCs parallel the effects seen in adults for conversion
of FSCs to FCs, in which a precise level of Wnt signal is required for normal FSC production and
optimal FSC maintenance, even though FC formation in adults and for the first egg chamber in pupae
are morphologically very different.
1123
1124 **Wnt activity is influential throughout pupation**
1125
The above results mostly showed a strong effect of Wnt signaling favoring more anterior
precursor locations when changes in Wnt pathway activity were applied throughout pupal
development. Additionally, strong effects on terminal FC formation were seen for genetic changes
initiated at pupariation but not if initiated at 24h APF (*arr*) or 36h APF (*axn*), showing that Wnt
signaling levels are critical very early in pupariation for migration of precursors out of the germarium
to form the first FCs. In theory, further insight into critical times of action can be gained by initiating
lineage tracing and a change in Wnt signaling at a full range of developmental times. However, a
limitation of using MARCM clones is that a strong GFP signal only develops after 3d because of
perdurance of GAL80 products. We therefore induced clones at 6d, 5d, 4d, 3d and 2d before eclosion
and examined lineages in adults two days after eclosion. A drawback of this approach is that the first
egg chambers produced during pupal development have matured into deposited eggs and are no
longer present in 2d-old adults, so marked FCs cannot be scored comprehensively. Hence, our
analysis focused only on ECs and FSCs. Also, FSCs present at eclosion are not scored directly but must
be inferred (see Methods). The experiment was performed for *axn* and also for loss of a different β-
catenin destruction complex member (using a double mutation, *apc1 apc2*) with similar results, for
arr, and for three control genotypes. Results, expressed as the percentage of all marked ECs/FSCs
(Supplementary Fig. 1), and were summed for all controls, two *arr* tests and the two genotypes (*axn*
and *apc1 apc2*) producing excess Wnt pathway activity. A small deficit of FSCs was apparent for
axn/apc in lineages initiated 2d before eclosion (48% vs 55% control) and a larger deficit was seen at
5d and 6d prior to eclosion (14% vs 33% control in each case), suggesting that increased Wnt pathway
activity provokes movement of precursors away from FSC-producing AP locations throughout pupal
development (Supplementary Fig. 1). Loss of *arr* activity produced a strong r1 EC deficit (5% vs 16%
control) and a substantial FSC surfeit (84% vs 55%) even from 2d prior to eclosion, showing that

normal Wnt signaling influences A/P movement of precursors in the last two days of pupal
development. The excess frequency of FSCs among ECs and FSCs for *arr* relative to controls averaged
over -2d and -3d clone initiation times was 32%; it was 64% averaged over -6d and -5d initiation
points. (Supplementary Fig. 1 and supporting information spreadsheet). These data suggest that loss
of Wnt activity, and hence normal Wnt signaling, has an effect on cell location within EC/FSC
precursor territory, as well as on terminal FC precursors, early in pupal development. The above
deductions have the limitation that they are based on FSC or r1 EC representation among non-FC
precursors rather than all precursors. Both *axn* and *arr* alter the proportion of precursors that
become FCs and the magnitude of this effect may vary during pupation, contributing to changes in
the measured representation of FSCs and r1 ECs over time.
1159
1160 **Pattern of Wnt pathway activity during pupal ovary development; an A/P gradient**
1161
Wnt pathway activity in adult germaria, measured with a Fz3-RFP reporter that responds to
genetic pathway manipulations in that tissue (WANG AND PAGE-MCCAW 2014; DAI *et al.* 2017; REILEIN *et*
al. 2017), is uniformly high in ECs and declines over the FSC domain to zero in the earliest FCs (Fig.
5L). The sources of Wnt in the adult germarium include high levels of Wg and Wnt6 production in Cap
cells and lower levels of Wnt 2 and Wnt4 in ECs (SAHAI-HERNANDEZ AND NYSTUL 2013; WANG AND PAGE-
MCCAW 2014; WANG AND PAGE-MCCAW 2018).
In pupal germaria, Fz3-RFP was first detected 27h APF and showed a marked increase in
intensity between 36h and 48h (Fig. 5A-G, J). At all stages, Fz3-RFP was higher in anterior than
posterior regions, consistent with the possibility that Cap Cells, which are specified at the anterior of
the germarium very close to the start of pupation, and even developing EC precursors, may be major
sources of Wnt ligand, as in the adult. The delay in reaching high levels of pathway activity might be
due to maturation of Cap Cell properties but is perhaps more likely due to functional differentiation
of ECs from precursors, most of which cease division, starting from the anterior, by 48h of pupation
(REILEIN *et al.* 2021). At 48h APF the spatial decline in Fz3-RFP was spread over the posterior two-
thirds of the germarium (Fig. 5G, J). By 60h APF, the domain of consistently high Fz3-RFP expression
occupied a larger anterior territory, extending roughly halfway to the region of strong Fas3
expression, followed by a steeper decline to background levels (Fig. 5H, K). At all stages, the anterior
border of strong Fas3 expression coincided with very low levels of Wnt pathway reporter activity. In
adult germaria and in pupal ovarioles prior to release of the first germline cyst, cells at, and posterior
to, the strong Fas3 border have committed to becoming FCs. The persistent coincidence of this Fas3
border with a decline of Wnt pathway activity to near zero suggests that low Wnt signaling may be a
pre-requisite for FC formation at all stages. This hypothesis is supported by the near-complete block
of FC formation during pupal development in *axn* mutant precursors and the increased propensity of
arr mutant precursors to form FCs throughout pupal development. These responses to altered
signaling appear to be mediated by net A/P migration throughout the precursor domain. Thus, *arr*
mutant cells are not depleted immediately anterior to potential FC-producing zones because there is
also enhanced migration to those positions from the EC precursor domain; in fact, adult FSC
production from *arr* precursor cells was increased overall. Conversely, *axn* mutant cells that do not
become FCs do not accumulate extensively in the neighborhood of FSCs; there was, instead, a net
loss of FSC fates in favor of EC production.
1192

1193
1194
Figure 5: Development of a Wnt signaling gradient in pupal germaria.
(A-L) Results from pupae expressing Fz3-RFP as a reporter for Wnt pathway activity. **(A)** Diagram of a 30h APF ovary. An
individual germarium is highlighted with anterior at the top. **(D-F, J-L)**. Individual germaria at the noted stages, oriented
anterior to the left, with the germarial region (containing germline cells) outlined in yellow and the anterior border of
strong Fas3 staining indicated by a blue arrowhead. Samples were stained for Fas3 (white) and pupal samples were
imaged intact, without separating developing ovarioles. **(B, C, G-I)** RFP profiles were constructed by measuring RFP
intensities (y-axes) at 5 µm intervals along the edge of the germarium where most somatic cell bodies are located, starting
adjacent to Cap Cells (0 on x-axes) and moving posteriorly. Individual germaria varied in RFP brightness and therefore the
profile for each germarium was normalized to the brightest cell in that germarium. Normalized curves were averaged and
SEM calculated to show the average Fz3-RFP profiles derived from germaria at **(B)** 27h APF (n=50 cells in 7 germaria), **(C)**
36h APF (n=108 cells in 14 germaria), **(G)** 48h APF (n=138 cells in 15 germaria), **(H)** 60h APF (n=178 cells in 14 germaria)
and **(I)** adults (n=178 cells in 14 germaria). The average location of the border of strong Fas3 expression (blue vertical
line), Fz3-RFP levels at that location (horizontal dashed line) and the location of the average posterior end of the germline
cells in the germarium (green vertical line; measured from the Cap Cells to the central point at the posterior of the
germline) are indicated. **(D-F, J-L)** Multiple z-sections were combined in a maximum intensity projection to show Fz3-RFP
in cells along the length of germaria. **(D)** Fz3-RFP was not detected at 21h APF, **(E)** was present at low levels with an
anterior bias at 27h (4 µm-thick z projection) and **(F)** at 36h APF (5 µm-thick z projection) before **(J)** accumulating to high
anterior levels with a steep decline ending close to the strong Fas3 border at 48h APF (4 µm-thick z projection) and **(K)**
60h-APF (5 µm-thick z projection), similar to **(L)** the profile in newly-eclosed adults (10 µm-thick z projection). Scale bar of
20 µm applies to all images. For raw data please see supplementary spreadsheet titled Fig5 Fz3RFP Graph.
1215
1216

JAK-STAT activity promotes FC formation and favors more posterior ovarian precursor fates

In Drosophila, Unpaired ligands activate a receptor to promote cross-phosphorylation and activation of the receptor-associated Janus kinase (JAK), Hopscotch (Hop), followed by phosphorylation and activation of the STAT transcription factor (ZEIDLER AND BAUSEK 2013). The effect of altered JAK-STAT signaling was explored using a strong loss of function *stat* mutation and overexpression of Hop, as used in studies of adult FSC behavior (VIED *et al.* 2012; MELAMED AND KALDERON 2020). The spectrum of adult derivatives of *stat* mutant precursors induced at pupariation was very similar to that of *axn* mutant cells. There was a drastic loss of lineages with any marked terminal FCs (2% vs 22%), of FC-only lineages (4% vs 17%) and of lineages with FSCs (5% vs 15%), while EC-only lineages (90% vs 63%) increased greatly in frequency (Fig. 6A, B, G). However, in contrast to the behavior of *axn* mutant cells, the frequency of lineages containing only r1 ECs (46% vs 49% control) or at least one r1 EC (64% vs 67% control) was not increased for *stat* mutant cells (Fig. 6G). The frequency of labeled ovarioles was similar for *stat* mutant cells (43%) and the direct control (47%) in the same experiment, suggesting that marked cells had not been lost extensively. Thus, we deduce that all precursors posterior to EC-producing territory have preferentially moved anteriorly during pupal development in response to loss of JAK-STAT pathway activity, leading to a near-absence of posterior derivatives, FSCs and FCs.

We also examined *stat* mutant lineages induced at 36h APF. As for *axn*, marked ovarioles with labeled terminal FCs (38%) were now as frequent as for controls (31%) (Fig. 6F; Supplementary spreadsheet "-3.5d stat November 2021 Numbers"), consistent with a conclusion that JAK-STAT signaling is critical in the first 48h of pupation for promoting posterior migration of precursors out of the germarium into the EGC and basal stalk. Marked ovarioles with labeled FSCs were very rare (3% vs control 13%) and the fraction of all labeled ECs and FSCs that were FSCs was much reduced (3% vs control 21%), as for lineages induced at pupariation. This is consistent with the possibility that JAK-STAT signaling is required for FSC formation during the second half of pupation, perhaps by promoting cell division.

Expression of a *UAS-Hop* gene at 29C (or two copies at 25C) in adult FSCs leads to increased JAK-STAT activity and a variety of strong phenotypes, including increased FSC division and differentiation to FCs (VIED *et al.* 2012; MELAMED AND KALDERON 2020). In adults, the effects of increased JAK-STAT activity on cell location and differentiation could be studied in the absence of greatly altered cell division by co-expressing the CycE-Cdk2 inhibitor Dacapo (Dap) (LANE *et al.* 1996; MELAMED AND KALDERON 2020). Adult ovaries included very large numbers of labeled cells in *UAS-Hop* lineages induced at pupariation (Fig. 6D), indicating greatly increased cell division. The highly populated lineages included anterior, EC regions, which normally only include 2-4 marked cells. In adult ovaries, excess JAK-STAT can stimulate significant division in otherwise quiescent cells in the EC domain (VIED *et al.* 2012; MELAMED AND KALDERON 2020). The abundance of marked cells in pupal lineages precluded useful classification into clone types, other than noting that almost all lineages included ECs, FSCs and FCs.

We therefore added *UAS-Dap*. MARCM clones induced at pupariation to initiate *UAS-Hop* (and *UAS-Dap*) expression might be expected to increase JAK-STAT activity only after a delay. However, we found that the fraction of lineages that included terminal FCs was increased significantly (50% vs 22%, Fig. 6G), suggesting that increased JAK-STAT promotes terminal FC production and that the transgene markedly alters JAK-STAT activity within the first 48h after clone induction. Precursors with increased JAK-STAT activity also produced an increased proportion of lineages containing FSCs (29% vs 15%) (Fig. 6C), while greatly reducing the frequency of EC-only lineages (40% vs 63%), r1 only lineages (33%

vs 49%) and lineages containing any r1 ECs (48% vs 68%) (Fig. 6G). The average number of cells in EC-only *UAS-Hop UAS-Dap* lineages was lower than for controls (1.1 vs 2.7). This is likely partly because many of the more posterior precursors that would normally produce only ECs shifted posteriorly to also produce FSCs (and FCs), so that EC-only lineages derived from only the most anterior precursors, which are the first to terminate division. The average number of ECs and FSCs in EC/FSC/FC lineages was similar to controls for *UAS-Hop UAS-Dap* lineages (8.0 vs 7.9 cells). Altogether, precursor division rates were likely similar to controls for this genotype, suggesting that the increase in FSC representation was due to direct effects on cell migration and not an increase in cell division rate, as observed for excess CycE expression.

Figure 6: JAK-STAT pathway promotes posterior migration of precursors during pupation.
(A-F) Representative lineages induced at (**A-E**) 0h APF or (**F**) 36h APF for the indicated genotypes. White arrowheads indicate Follicle Cells; magenta arrows, r1 ECs; yellow arrows, r2a ECs; cyan arrows, FSCs. DAPI, blue; GFP, green; Fas3, red. (**A-C**) The r1 EC, r2a EC, FSC and FC domains are shown in corresponding colors in the horizontal bars above. **(A)** control EC/FSC lineage that includes r1 ECs, **(B)** *stat* EC-only lineage, **(C)** *UAS-Dap UAS-Hop* EC/FSC/FC lineage, **(D)** *UAS-Hop* lineage, illustrating an exceptionally high number of marked cells, likely due to increased cell division, and **(E)** *arr; UAS-Dap UAS-Hop* lineage, illustrating the common FC-only lineage outcome. **(F)** A *stat* lineage induced at 36h APF illustrating terminal FC marking, which is virtually absent for *stat* lineages induced at 0h APF. Scale Bar 20 μm for **(A-C)** and 50 μm for **(D-F)**. **(G)** Deduced frequency of the indicated types of single-cell lineage as a percentage of all lineages induced at 0h APF. Error bars show SEM. One asterisk (*) indicates p<0.05; two asterisks (**) indicates p<0.005 (Fisher's two-tailed exact test). Number of marked ovarioles = 739 (control), 142 (*stat*), 50 (*UAS-Dap UAS-Hop*), 57 (*arr; UAS-Dap UAS-Hop*), 95 (*arr; UAS-DIAP1*). For raw data and calculations please see supplementary spreadsheets titled 0h APF Graphs Fig4-10; Compilation of Numbers Figure 6 JAK-STAT.

Thus, increased JAK-STAT activity has the potential to promote posterior movement of all precursors, even from the most anterior r1 EC-producing locations. The restriction of the opposite phenotype for *stat* loss of function to cells posterior to the EC-producing domain (i.e., no increase in r1 EC-containing lineages) suggests that anterior regions of the developing ovariole may not normally experience significant levels of JAK-STAT signaling.

We examined JAK-STAT pathway activity using reporters containing either ten tandem STAT
binding sites upstream of GFP (Bach *et al.* 2007) or STAT-responsive promoter sequences from the
Socs36E gene upstream of RFP (He *et al.* 2019). Both reporters showed similar patterns in adult
ovaries (Melamed and Kalderon 2020) and in pupal ovaries. There was very strong fluorescence
surrounding each developing ovariole in both cases but this did not prevent signal detection in
individual confocal sections of the developing ovarioles themselves (Fig. 7 and Supplementary Fig. 2).
Activity was first detected clearly at about 30h APF, strongest around and anterior to the border
between Fas3-positive and Fas3-negative somatic cells (Fig. 7A). Thereafter, reporter activity
increased and spread posteriorly into Fas3-positive cells. The signal was lower in the anterior half of
the developing germarium than in more posterior Fas3-negative somatic cells (Fig. 7B-D). These
results are not sufficient to point to a specific cellular source of ligand. However, Fas3-positive cells
are among the candidates. They may themselves initially be less responsive to ligand than Fas3-
negative cells but reporter activity was as strong in Fas3-positive cells as in their anterior neighbors
by 36h APF and thereafter (Fig. 7B-D; Supplementary Fig. 2). The higher pathway activity observed in
somatic cells around, and posterior to, the three or four most mature germline cysts is consistent
with the possibility that pathway activity in this region is necessary for a cell to maintain its position
or to promote more posterior migration in order to seed FCs of the first budded egg chamber. We did
not observe a clear gradient within this region (Fig. 7A-D).
Shortly after the most posterior germline cyst had left the germarium to form the first egg
chamber (around 60h APF), somatic cells anterior to that cyst had particularly strong reporter activity,
similar to nascent stalk cells during adult oogenesis (Supplementary Fig. 2). By that time, both Castor
staining and higher Fas3 staining than in surrounding cells indicate the presence of polar cells in the
first egg chamber (Reilein *et al.* 2021). Polar cells were similarly evident in the second egg chamber by
about 84h APF, with strong STAT-GFP reporter activity in nearby somatic cells (Supplementary Fig. 2).
The STAT-GFP signal extending into the germarium at 84h APF was significantly stronger than at
earlier times and strongly resembled the adult pattern, with very little activity anterior to region 2a
cysts (Supplementary Fig. 2). It is possible that the higher STAT-GFP signal is due to polar cells
contributing JAK-STAT ligand. However, the overall spatial pattern of STAT-GFP was similar earlier and
it is possible that the technical necessity of maintaining an intact ovary during dissection and staining
of early pupal ovaries (rather than releasing individual ovarioles) limited GFP detection by antibody
prior to 84h APF. Thus, there clearly is detectable JAK-STAT pathway activity prior to any polar cell
formation in a pattern that largely excludes the anterior half of the developing germarium. It also
appears that this pattern is significantly enhanced once polar cells have formed in the second budded
egg chamber. The minimal levels of JAK-STAT pathway activity in anterior regions throughout the
pupal period are consistent with the observation that marked r1 EC frequency is not altered by the
absence of STAT activity, even though ectopic JAK-STAT can convert their precursors to adult cells in
more posterior locations.
In adults, altering JAK-STAT activity in either direction greatly affects the division rate of FSCs,
while excess JAK-STAT can promote significant division of otherwise quiescent ECs (Melamed and
Kalderon 2020; Melamed *et al.* 2023). During pupation, loss of STAT activity barely altered the average
number of cells in EC-only lineages (2.5 vs 2.7). This is likely partly because some of the precursors
were originally in more posterior locations and set to divide faster. It is also consistent with the lack
of net migration of r2a EC precursors to become r1 ECs and the observation of only low JAK-STAT
activity in the normal EC precursor domain. There were insufficient *stat* mutant lineages in terminal
egg chambers to measure average FC occupancy and insufficient FSC-containing lineages to measure
the average number of marked ECs and FSCs. Thus, the strong anterior movement of *stat* mutant

derivatives precluded direct measurement of the normal role of JAK-STAT signaling in supporting the
division of posterior precursors through pupation.
1341

1342
1343
Figure 7: JAK/STAT signaling in pupal ovaries.
(A-D) Pupal ovaries expressing STAT-RFP were stained for RFP (red), Fasciclin 3 (green), and with DAPI (blue). STAT-RFP
was strongly expressed in epithelial sheath cells surrounding each developing germarium and the entire ovary. STAT-RFP
was first detectable in developing germaria at 30h APF, and increased in intensity over time. **(A)** At 30h APF, STAT-RFP was
detected at low levels around the most mature cysts in the germarium and in the most anterior Fasciclin III-expressing
cells posterior to the germarium (in the Extra-Germarial Crown (EGC)). **(B-D)** From 36h APF to 60h APF, STAT-RFP
expression was seen around the most mature egg chambers and throughout the EGCs and basal stalks. **(D)** At 60h APF,
STAT-RFP was also seen in follicle cells of the budding egg chamber. The most mature cyst in each germarium is marked
with an asterisk. Scale Bars, 20 μm.
1353
Since loss of Wnt and increased JAK-STAT signaling have similar outcomes of promoting
posterior precursor movement, it might be expected that combining these changes would produce
still stronger phenotypes, as seen for adult FSCs. Adding *UAS-Hop UAS-Dap* to loss of *arr* produced
very little change in the various measured parameters, other than an increase in FC-only lineages
(46% vs 33%, Fig. 6E, G). Comparison of the phenotype of *arr UAS-Hop UAS-Dap* with *UAS-Hop UAS-*
Dap is less straightforward for technical reasons: the clone frequency for the latter genotype was

significantly higher, likely leading to incomplete resolution of double clones (suggested also by the
comparatively high EC+FC frequency of 12% vs the exceptionally low value of 1% for the former
genotype). Nevertheless, there are some clear changes imposed by loss of Wnt signaling. The
frequency of r1 EC-only clones (25% vs 33%) and lineages containing any r1 ECs (32% vs 48%) was
reduced, while the frequency of FC-only lineages was increased (46% vs 19%; or 47% vs 31% if EC+FC
lineages are included), as was the fraction of lineages containing FCs but no FSC (46% vs 28%). These
indicate greater posterior shifts for precursors in the normal EC and FSC producing domains. The
effect on terminal FC production is less clear, likely because of the difference in clone frequencies and
consequent limitations in deducing single lineages. Overall, the formation of terminal FCs, already
increased by raising JAK-STAT activity, does not appear to be enhanced by additional elimination of
Wnt pathway activity, contrasting with the effects on precursors in more anterior locations.
Altogether, the loss of Wnt signaling was more influential in anterior regions, while excess JAK-STAT
had a greater influence in more posterior regions. A similar conclusion was made regarding the much
narrower spatial domain of FSCs in adults (MELAMED AND KALDERON 2020).
1374
1375 **Yki activity primarily affects ovarian precursor survival and cell division**
1376
We were interested in examining the roles of Hedgehog (Hh) signaling and the Hippo/Yorkie
pathway for two reasons. First, both have major inputs into adult FSC division rates (HARTMAN *et al.*
2013; HUANG AND KALDERON 2014; HSU *et al.* 2017) and we hoped to learn more about the regulation of
somatic ovarian precursor proliferation. In adults, Hh stimulates FSC cell division through Yorkie (Yki)
(HUANG AND KALDERON 2014). Second, Hh ligand is known to emanate from anterior locations in pupal
ovaries (LAI *et al.* 2017), so that it might influence precursor cell migration, even though neither Hh
nor Yki are known to have a significant effect on the migration or differentiation of adult FSCs.
Yki is a transcriptional co-activator frequently implicated in the regulation of cell and tissue
growth. Yki activity is generally regulated, principally through inactivating phosphorylation, by
upstream members of the Hippo (Hpo) pathway, including Hpo and Warts (Wts) protein kinases and a
variable assortment of more upstream proteins (including Kibra) capable of sensing mechanical
stimuli, including tension (MISRA AND IRVINE 2018). However, in adult ovaries, Yki is also
transcriptionally regulated by the Hh signaling pathway (HUANG AND KALDERON 2014). In adult FSC
lineages, loss of *yki* resulted in reduced division, increased apoptosis and rapid loss of FSCs. FSC
maintenance was substantially improved by expression of excess CycE and excess DIAP1, both known
transcriptional targets of Yki, to counter those deficiencies (HUANG AND KALDERON 2014).
When *yki* mutant lineages were induced at pupariation, the proportion of ovarioles with
labeled cells was lower (19%) than for the direct side-by-side control (37%) but this deficit was largely
restored by addition of *UAS-DIAP1* (32%) to inhibit apoptosis. Both the reduction relative to controls
for *yki* (13% vs 59%) and restoration by *UAS-DIAP1* (to 60%) were observed in a second experiment.
Although variation in the frequency of clone induction is potentially quite high because of the brief
and moderate heat-shock protocol, these results suggest that some *yki* mutant lineages are lost to
apoptosis during pupal development.
For each experiment, FSCs were rarely present in either *yki* (0% and 6%) or *yki; UAS-DIAP1*
lineages (4% and 3%) relative to control averages (15%) (Fig. 8A, H). For *yki; UAS-DIAP1*, there was no
significant difference in the fraction of lineages with either r1 ECs or terminal FCs relative to controls.
This suggests that there is no strong overall A/P shift in precursor outcomes, contrasting with
manipulations of Wnt or JAK-STAT pathways, and that FSC loss is therefore most likely to result from
reduced division rates, as observed for *cycE* and *cutlet* mutations.

1407
Figure 8: Yki activity counters posterior derivative cell apoptosis, and promotes cell division rate to increase FSC
representation.
(A-G) Representative lineages induced at (**A-C, E, F**) 0h APF or (**D, G**) 48h before pupariation for the indicated genotypes.
White arrowheads indicate Follicle Cells; magenta arrows, r1 ECs; yellow arrows, r2a ECs; cyan arrows, FSCs. DAPI, blue;
GFP, green; Fas3, red. The r1 EC, r2a EC, FSC and FC domains are shown in corresponding colors in the horizontal bars
above images other than **(B)**. **(A)** *yki* commonly produced EC-only lineages, whereas **(B)** the addition of *UAS-DIAP1* greatly
increased the representation of FCs, including terminal FCs, as shown here. **(C-D)** The further addition of *UAS-CycE*
restored FSC-containing lineages for clones induced at **(D)** 48h before pupariation, as shown here, but not at **(C)** 0h APF;
here showing an EC+FC lineage, characteristic of FSC loss. When *yki* mutant cells became FCs (generally requiring the
presence of *UAS-DIAP1*)), they were frequently polar cells (arrowheads in **(C)**) but sometimes produced small patches of
main-body FCs, as in **(B)**. **(E-G)** show lineages with increased Yki activity. **(E)** a *wts* lineage, illustrating the common r2a EC-
only lineage, owing to a reduction in r1 EC fates, and **(F)** a *hpo* lineage, illustrating the increased frequency of EC/FSC/FC
lineages, again in the absence of r1 ECs. **(G)** A UAS-Yki* (S168A) EC/FSC/FC lineage induced 48h before pupariation,
illustrating the increase in such lineages and a large number of labeled cells, indicative of increased division rates. Scale
Bar 20 μm for **(A, C-G)** and 50 μm for **(B)**. **(H, I)** Deduced frequency of the indicated types of single-cell lineage as a

percentage of all lineages induced at **(H)** 0h APF and **(I)** -48h APF. Error bars show SEM. One asterisk (*) indicates p<0.05; two asterisks (**) indicates p<0.005 (Fisher's two-tailed exact test). (**H**) Number of marked ovarioles = 739 (control), 84 (*yki*), 127 (*yki; UAS-DIAP1*), 67 (*yki; UAS-DIAP1 UAS-CycE*), 44 (*kibra*), 42 (*wts*), 82 (*hpo)*, 49 (*UAS-Yki**). (**I**) Number of marked ovarioles = 319 (control), 57 (*yki*), 101 (*yki; UAS-DIAP1*), 94 (*yki; UAS-CycE UAS-DIAP1*), 85 (*UAS-yki**). For raw data and calculations please see supplementary spreadsheets titled 0h APF Graphs Fig4-10; -2d APF Graphs**;** Compilation of Numbers Figure 8 yki hpo -5d; Compilation of Numbers Figure 8 yki hpo -5d.

Other changes in lineage outcomes suggest that loss of Yki does have some effect on final A/P cell locations, either by affecting precursor migration or spatially selective apoptosis. FC-only lineages were seen at lower frequency for *yki* (6% and 10%) relative to controls (average 17%), with no increase of EC/FC lineages (1% and 0% vs 5% for controls); correspondingly, there was a large increase in EC-only clones (95% and 86% vs 63% for controls). For *yki; UAS-DIAP1*, however, the average frequency of FC-only (20% vs 17%) and EC/FC lineages (3% vs 5%) was similar to controls. Addition of *UAS-CycE* together with *UAS-DIAP1* also resulted in frequencies of FC-only (20%) and EC/FC lineages (5%) similar to controls. Thus, the addition of *UAS-DIAP1* restored FC representation, including terminal FCs to control levels in two separate genetic tests, suggesting that posterior (would-be FCs) derivatives preferentially die in the absence of Yki. A caveat to this conclusion is that the restoration of FC lineages by *UAS-DIAP1* was seen only in one of two experiments for *yki; UAS-DIAP1* (5% and 35% for FC-only; 4% and 2% for EC/FC) (Fig. 8B, H). Moreover, the FC deficit seen for *yki* (with no *UAS-DIAP1*) in both experiments was not reproduced for lineages initiated 2d before pupariation (see below). A plausible explanation is that both a posterior bias of apoptosis induced by loss of Yki and elimination of that bias by addition of DIAP1 are variable (perhaps poised close to a threshold). In that case, the modest increase of EC-only lineage proportion and failure of lost FSC lineages to be converted to FC lineages for *yki; UAS-DIAP1*, as expected for reduced division, may be due to some residual preferentially posterior apoptosis, with no underlying direct effect of loss of Yki on A/P cell migration.

To examine further whether loss of lineages with FSCs was due to lower division rates, we tested the effect of adding *UAS-CycE* to *yki; UAS-DIAP1* lineages. This did not substantially restore FSCs (present in only 4% of lineages, Fig. 8C) for lineages induced at pupariation. However, the MARCM system introduces a delay in the accumulation of *UAS-CycE* product because perduring GAL80 gene products must first be diluted or degraded. When lineages were induced 2d prior to pupariation the frequency of FSC-containing lineages was increased from 1% (*yki*) and 3% (*yki; UAS-DIAP1*) to 18% (Fig. 8D), closer to the control average of 22% (Fig. 8I), suggesting that reduced division is indeed the primary reason for the deficit of FSCs in *yki* mutant lineages.

Other results from lineages induced 2d before pupariation showed some similarities and some differences to results from lineages induced at pupariation. The frequency of marked ovarioles for *yki; UAS-DIAP1* (74%) and *yki; UAS-CycE UAS-DIAP1* (73%) were once again higher than for *yki* (43%), suggesting some apoptotic loss of lineages for loss of Yki activity. The frequency of lineages with FCs but no FSCs (FC-only plus EC/FC) was similar to controls (23%) for the two types of *yki* lineage expressing DIAP1 (25% and 27%), as noted for clones induced at pupariation. However, *yki* lineages produced a similar result (28%) (Fig. 8I), rather than a reduction of these FC lineages, as observed for clones induced at pupariation (Fig. 8H). The fraction of lineages with labeled terminal FCs was also similar among all three *yki* genotypes (25%, 23% and 22%). Interestingly, these three values were all somewhat lower than the control average of 35%. This difference is particularly notable for the *yki; UAS-CycE UAS-DIAP1* genotype because all other lineage characteristics are similar to controls. Thus, loss of Yki results in reduced division, leading to a severe loss of FSCs, apoptosis with variable posterior cell (FC) selectivity, and a small decline in lineages with terminal FCs.

In all cases, FCs that lacked *yki* function shared some characteristics. Labeled cells were
preferentially found only in polar cells (most commonly), stalk cells or the basal stalk (Fig. 8C, data
not shown). Labeled FCs in other locations were approximately equally split between two categories:
either a pair of cells with strongly elevated Fas3, resembling ectopic polar cells, or a relatively normal
FC patch (Fig. 8B). Adult-induced lineages were previously described as frequently including polar
cells or ectopic polar cells (CHEN *et al.* 2011). Our results suggest that loss of *yki* greatly increase the
chances that a FC derivative acquires the position or characteristics of a precursor of stalk and polar
cells; cells that nevertheless become main-body FCs can develop further in a relatively normal fashion
despite the absence of *yki* activity.
The consequences of increasing Yki activity were examined using strong loss of function alleles
for upstream regulators that can limit Yki activity. *kibra*, wts and *hpo* lineages, induced at pupariation,
shared the phenotype of a reduced fraction of both r1 EC-only lineages (31% *kibra*, 34% *wts* and 26%
hpo vs 49% control) and lineages containing any r1 ECs (37%, 35% and 42% vs 67%, respectively) (Fig.
8H). These manipulations did not reproducibly change the proportion of lineages with marked
terminal FCs (28%, 18% and 21% vs 22% control), while only specific mutations increased the
proportion of lineages with FSCs (*hpo*; 27% vs 15%; others were 11%) (Fig. 8E, F, H). Neither the
number of cells in EC-only lineages (*kibra* 1.8, *wts* 1.8 and *hpo* 2.2 vs control 2.7), nor the average
sum of ECs and FSCs in EC/FSC/FC lineages were increased by these mutations (7.3, 9.8, 6.8 vs control
8.6). Thus, it appears that cell division rates were not significantly increased.
We also tested the effects of expressing a Yki derivative (Yki-S168A [*UAS-Yki**]) partially
resistant to down-regulation by upstream pathway components (CHEN *et al.* 2011; HUANG AND
KALDERON 2014). This resulted in an increase in lineages with FSCs (25% vs 15% control), as for *hpo*
(but not the other Hpo pathway mutations), consistent with an expected consequence of an
increased division rate. UAS-Yki* also produced a slightly reduced fraction of r1 EC-only lineages
induced at pupariation (38% vs 49% control) but a normal fraction of lineages with any r1 ECs (67% vs
67%) (Fig. 8H). The former decrease might simply be a consequence of increased division expanding
some of the most anterior clones into region 2a.
In *UAS-Yki** lineages initiated 2d before pupariation, the frequency of r1 EC-only (15% vs 43%)
and EC-only lineages (35% vs 55%) was greatly reduced while lineages with marked FSCs were greatly
increased (46% vs 22%) (Fig. 8G, I). Lineages with any marked r1 ECs (75% vs 75%) or terminal FCs
(42% vs 35% control) were not greatly changed in frequency. The average size of EC-only clones (3.5
vs 3.2) and the average number of ECs and FSCs in EC/FSC/FC clones (17.9 vs 14.5) were slightly
increased. These results suggest that *UAS-Yki** is more effective when induced prior to pupariation,
allowing an increase in cell division rate, that greatly increases FSC production; the increase was
slightly greater than achieved by excess CycE expression overv the same time period. The depletion
of r1 only and EC-only lineages without affecting the frequency of lineages containing an r1 EC or an
EC may result from increased clone sizes, with some spreading into more posterior domains, rather
than indicating an anterior to posterior shift of all derivatives. By contrast, there was a depletion of r1
ECs amongst all clones for *kibra*, *wts* and *hpo* lineages induced at pupariation and the former two
genotypes did not appear to affect cell division rate. Thus, Hippo pathway components primarily
affected A/P movement of the most anterior precursors, whereas excess Yki primarily affected
division rate and, hence, FSC representation.
1512
1513
1514

Hedgehog signaling promotes cell division, counters posterior derivative cell apoptosis and can influence A/P migration

The normal effect of Hh signaling on somatic ovarian precursors is most simply investigated by generating lineages homozygous for a strong *smoothened* (*smo^2*) loss of function mutation, preventing all responses to Hh (INGHAM 2022). When induced at pupariation, the proportion of ovarioles with labeled cells was consistently lower than for controls over three experiments (10% vs 28%, 12% vs 40%, 17% vs 34%). The yield of labeled ovarioles remained lower for clones of genotype *smo; UAS-DIAP1* (47% vs 73%, 19% and 17% vs 34%). Given the strong restoration of *yki* mutant clone frequency by *UAS-DIAP1* under analogous conditions, the results suggest that some *smo* mutant lineages are lost for reasons other than apoptosis or that the apoptotic signal is stronger than induced by *yki* and is not efficiently countered by *UAS-DIAP1*. The nature of recovered lineages was nevertheless influenced by the presence of *UAS-DIAP1*, as described below, suggesting some selective loss of cells by apoptosis. Expression of *UAS-DIAP1* alone did not alter any control phenotype significantly (Fig. 9D, E).

The phenotype shared by *smo* and *smo; UAS-DIAP1* (each cited as the average of three tests) was a reduction of lineages that include FSCs (3% each vs 15% control) (Fig. 9A, B, D). The average number of cells per EC-only clone (1.4 and 1.4 vs 2.7) and per EC/FSC clone (1.2 and 3.1 (both from fewer than five clones) vs 8.6) were reduced (Fig. 10H), as was the average size of a terminal FC patch (*smo* 15% vs 26% control, *smo; UAS-DIAP1* 15% vs 20% control; Fig. 10J). All of these data are consistent with a reduced division rate leading to loss of FSCs. That deficit alone might be expected to produce an increase in FC-only or EC/FC lineages. The frequency of FC-only plus EC/FC lineages was indeed significantly increased for *smo; UAS-DIAP1* lineages (37% + 7% = 44% vs 17% + 6% = 23% for control) but not for *smo* lineages (19% + 1% = 20%). There was also a reduction in lineages that included terminal FCs for *smo* (11% vs 22% control) but not for *smo; UAS-DIAP1* (24%) (Fig. 9A, B, D). These results suggest that some *smo* mutant FCs, including some of those contributing to the terminal egg chamber, are lost to apoptosis, reminiscent of the *yki* mutant genotype. For *smo; UAS-DIAP1* cells, there was a small reduction in the fraction of lineages including r1 ECs (56% vs 67%). The increase in FC-only lineages (37% vs 17%) noted above was also greater than expected to result only from loss of FSCs. Hence, loss of Hh signaling in the presence of *UAS-DIAP1* appears also to result in a posterior shift in precursor derivatives. This does not appear to extend to the most posterior precursors, since the proportion of lineages with labeled terminal FCs remained almost unchanged (24% vs 22%).

The phenotypes of *smo* and *smo; UAS-DIAP1* lineages were reproduced when induced 2d prior to pupariation. Clone frequency was reduced (17% and 21%, respectively versus 53% for the control, which expressed *UAS-DIAP1*) and lineages with FSCs were greatly reduced in both cases (1% and 1% vs 22% control average, Fig. 9E). The fraction of lineages with marked FCs in the absence of FSCs (FC-only plus EC/FC) was significantly increased only in the presence of DIAP1 (*smo* 26% and *smo; UAS-DIAP1* 44% vs 23% control), with a marked decrease in representation of terminal FCs amongst all lineages only in the absence of *UAS-DIAP1* (*smo* 20% and *smo; UAS-DIAP1* 31% vs 35% controls, Fig. 9E). Again, there was a decrease of lineages containing r1 ECs when DIAP1 was present (47% vs 75% control). These results are consistent with a posterior shift of all but the most posterior FC precursors in the absence of Hh signaling, together with some loss of FCs through apoptosis. This posterior shift would by itself increase FSC representation, based on the *arr* mutant phenotype, so the observed loss of FSCs is almost certainly due to reduced division rates. The addition of *UAS-CycE* to the *smo; UAS-DIAP1* genotype did not, however, increase FSC representation (Fig. 9E), contrasting with the

1561 analogous rescue observed for *yki* mutant lineages. One possibility is that the reduction in cell
1562 division due to loss of *smo*, which may be more profound than from loss of *yki* (consistent with a
1563 greater FSC deficit), is not rescued effectively or in a timely fashion by *UAS-CycE*.
1564

1565
Figure 9. Hedgehog signaling promotes cell division and FSC representation, counters posterior derivative cell apoptosis
and can influence A/P migration
(A-C) Representative lineages induced at 0h APF for the indicated genotypes. White arrowheads indicate Follicle Cells;
magenta arrows, r1 ECs; yellow arrows, r2a ECs; cyan arrows, FSCs. DAPI, blue; GFP, green; Fas3, red. **(A)** A *smo* EC-only
lineage and **(B)** a *smo; UAS-DIAP1* lineage with terminal FC marking, rarely seen in the absence of *UAS-DIAP1*. **(C)** A *smo*
pka Su(fu) lineage, illustrating the common EC-only outcome. FSCs were rarely observed for all three illustrated
genotypes. Scale Bar 20 μm for **(C)** and 50 μm for **(A, B)**. **(D, E)** Deduced frequency of the indicated types of single-cell
lineage as a percentage of all lineages induced at **(D)** 0h APF and **(E)** -48h APF. Error bars show SEM. One asterisk (*)
indicates p<0.05; two asterisks (**) indicates p<0.005 (Fisher's two-tailed exact test). **(D)** Number of marked ovarioles =
739 (control), 59 (*UAS-DIAP1*), 98 (*smo*), 142 (*smo; UAS-DIAP1*), 38 (*pka*), 51 (*smo pka Su(fu)*), 85 (*ptc*). **(E)** Number of
marked ovarioles = 319 (control), 58 (*UAS-DIAP1*), 37 (*smo*), 88 (*smo; UAS-DIAP1*), 55 (*smo; UAS-DIAP1 UAS-CycE*). For raw
data and calculations please see supplementary spreadsheets titled 0h APF Graphs Fig4-10; -2d APF Graphs**;** Compilation
of Numbers Figure 9 Hh -5d; Compilation of Numbers Figure 9 Hh -5d.
1579

We also investigated the effects of partially compromising Hh signaling. In *Drosophila* wing
discs, the loss of Protein Kinase A (PKA) compromises Smo activation by Hh but also blocks proteolytic
processing of the full-length latent transcriptional activator Ci to a repressor form (Jɪᴀ *et al.* 2003;
Aᴘɪᴏɴɪsʜᴇᴠ *et al.* 2005; Sᴍᴇʟᴋɪɴsᴏɴ *et al.* 2007; Lɪᴛᴛʟᴇ *et al.* 2020). It therefore produces an intermediate
level of Hh pathway activity (high levels of full-length Ci but limited activation of Ci-155 by Smo-

activated Fused kinase). Similarly, *smo pka: Su(fu)* mutant cells allow no activation of Fused but
partially alleviate the requirement for Fu kinase activity because of the absence of Su(fu) (Suppressor
of fused) restraint on full-length Ci activity (OHLMEYER AND KALDERON 1998). In wing discs, both *pka* and
smo pka; Su(fu) genotypes produce relatively high levels of pathway activity but lower than maximal
Hh signaling (manifest as strong transcriptional *patched* (*ptc*) induction but no induction of anterior
Engrailed). In developing ovarioles, there was no significant reduction in clone frequency for these
lineages induced at pupariation. Both genotypes led to a reduced fraction of lineages containing FSCs
(12% and 3% vs 15% control, Fig. 9D). The effect was strongest for *smo pka Su(fu)* (Fig. 9C), suggesting
that precursors in locations destined to produce FSCs normally receive high levels of Hh during some
or all of pupal development. The FSC deficit was accompanied by an increase in lineages with FCs but
no FSCs for *pka* and *smo pka; Su(fu)* (25% and 41% vs 22% for controls), without large changes in the
fraction of lineages with terminal FCs (18% and 24% vs 22% control) or r1 ECs (54% and 66% vs 67%)
(Fig. 9D). Thus, even intermediate reductions in Hh pathway activity reduce FSC production, likely
through reduced cell division rates, while complete loss of pathway activity also results in a posterior
shift of all but terminal FC precursors, accompanied by some loss of posterior precursor lineages
through apoptosis.
The maximal physiological level of Hh pathway activity is generally reproduced by loss of *ptc*
function (INGHAM 2022). The phenotype of *ptc* mutant lineages induced at pupariation was consistent
with an elevated division rate. The fraction of lineages that included an FSC was increased (24% vs
15% control, Fig. 9D, Fig. 10A), as was the average size of an EC/FSC lineage (13.4 vs 8.6 control EC
plus FSC cells; Fig. 10H). The fraction of all marked ECs and FSCs that were FSCs was also higher (28%
vs 19% control average). The average size of a terminal FC patch was slightly higher (29% vs 24%; this
measure excludes clones exclusively composed of polar cells, stalk cells or basal stalk cells; Fig. 10K).
Since *ptc* marked FCs were generally very abundant (Fig. 10D) we wondered if these measures fully
represented the division rate of *ptc* mutant cells. We therefore also measured indicators of the
spread of clones as a further indicator of division rate. A precursor at pupariation that is destined to
contribute to the terminal egg chamber also sometimes contributes to the penultimate egg chamber.
As noted in the first section of Results, the lineage sometimes also contributes FCs occupying more
anterior positions. A lineage that produces more cells is more likely to spread into each of these
locations. Control lineages contributing to the terminal FC also populated the penultimate egg
chamber with a frequency of 38% and more anterior egg chambers with a frequency of 25%. Those
values were increased to 55% and 66%, respectively for *ptc* lineages (Fig. 10L). This measure does not
correct for single-cell lineages and in other experiments the control values for penultimate egg
chamber occupancy were higher (to give an average of 49%) (Fig. 10L). Thus, our cumulative evidence
suggests that loss of *ptc* function may not increase the division rate of the most posterior precursors.
The spread of FSC-containing lineages was estimated by counting the fraction of cysts with adjacent
marked FCs. This value was much higher for *ptc* (80%) than for control FSC-containing lineages (Fig.
10I), supporting the deduction from marked EC and FSC numbers in such lineages (Fig. 10H) that *ptc*
increases division rate for precursors in the regions that can produce FSCs. The average size of an EC-
only clone was not increased for *ptc* lineages (2.0 vs 2.7 control cells; Fig. 10H), suggesting that
division is not increased in the most anterior regions, closest to the source of Hh.
There also appears to be a posterior bias in the distribution of *ptc* mutant lineages, with fewer
lineages including r1 ECs (57% vs 67% control) and more lineages including terminal FCs (34% vs
22%), reflected also in an increased fraction of lineages containing FCs but no FSC (38% vs 22%) (Fig.
9D). A posterior shift can also contribute to an increase of lineages with FSCs, as observed for *arr*
lineages. Nevertheless, the *ptc* phenotype shows a lesser posterior shift than *arr*, a greater increase in

EC/FSC lineages and some direct evidence of increased division rates, affirming a significant
contribution of division rate to FSC representation. The major effect of Hh signaling on cell division
rate and FSC production was manifest by opposite *smo* and *ptc* phenotypes. Effects on cell location
appear to be more complex: increased Hh signaling caused posterior displacement throughout the
precursor domain, while loss of Hh signaling produced the same result (not the converse, as might be
expected) for all but the most posterior precursors.
1637
1638 **Relationship between Hh signaling and Yki activity**
1639
In adult FSC lineages, *ptc* increased *yki* RNA levels, *yki* was fully epistatic to *ptc* and
substituting *yki* with a *tub-yki* transgene, preventing normal transcriptional regulation of *yki*,
suppressed all *ptc* phenotypes (to produce roughly wild-type FSC properties) (HUANG AND KALDERON
2014). It was therefore concluded that *ptc* increased FSC representation (through enhanced division)
via transcriptional induction of *yki*. Another report suggested a post-transcriptional regulation of Yki
activity through the Hh pathway (LI *et al.* 2015), while the study of adult ECs deficient for *yki* or *smo*
activity led to a conclusion of independent (albeit similar) Hh and Yki effects, each leading to EC
dysfunction and consequently delayed or deficient germ cell differentiation (HUANG *et al.* 2017).
We explored the relationship between Hh signaling and Yki activity in ovarian precursors by
manipulating both simultaneously. *ptc yki* lineages initiated at pupariation had phenotypes similar to
yki phenotypes (Fig. 8A, Fig. 10B). Lineages with FSCs were less frequent than for controls (6% vs 15%;
yki 2%), as were FC-only lineages (11% vs 17%; *yki* 8%) and the subset with only terminal FCs (5% vs
10%; *yki* 3%), while EC-only lineages were more frequent (71% vs 63%; *yki* 90%), as were r1 EC-only
lineages (61% vs 49%; *yki* 82%) (Fig. 10G). Correspondingly, lineages that included labeled terminal
FCs were less frequent (10% vs 22%; *yki* 3%) and those with r1 ECs were more abundant (82% vs 67%;
yki 89%). All of these deviations from wild-type behavior are similar to those of *yki* lineages, though
smaller in magnitude, and opposite to changes characteristic of *ptc* alone (Figs. 8 and 10). Thus, loss
of *yki* is largely epistatic to loss of *ptc*, regarding the distribution of lineage types. All terminal FC
samples for *ptc yki*, like *yki*, were mostly composed of polar cells or basal stalk cells, so a reliable
measure of division rate for terminal epithelial FC precursors is not available. Similarly, there were
too few FSC-containing lineages (just three) to provide a reliable estimate of division rate of EC/FSC
precursors. Thus, the loss of FSC-containing *ptc yki* lineages is plausibly due to a reduced rate of
precursor division, as deduced for *yki* lineages through rescue with *UAS-CycE*, but there is no direct
supporting evidence.
The phenotype of marked *ptc yki* FCs commonly included the characteristic *yki* phenotype of
occupying polar cell locations or producing ectopic polar-like cells with high Fas3, but also included
multi-layering (Fig. 10E), with the marked cells in an outer layer, or the presence of multiple excess
labeled cells between egg chambers. Those phenotypes were previously noted for *ptc* mutant FCs
induced in adults (ZHANG AND KALDERON 2000) and were evident also in *ptc* mutant pupal lineages (Fig.
10D), where the FC patches were generally much larger than for *ptc yki* (supporting the hypothesis of
a significantly lower division rate). We deduce that the multi-layering phenotype, indicative of a
different affinity of *ptc* mutant FCs for germline cells and wild-type FCs, is not dependent on Yki
activity.
Substitution of *yki* function with *tub-yki* resulted in lineages with phenotypes similar to
controls (Fig. 10G). This result suggests that the levels of *yki* provided by the transgene are roughly
normal (consistent with maintenance of healthy stocks with *tub-yki* replacing endogenous *yki* activity)

and that *yki* need not be regulated transcriptionally for roughly normal ovarian somatic cell precursor
behavior.
1678

1679
1680
Figure 10: Dependence of *ptc* mutant phenotypes on transcriptional modulation of *yki*.
(A-F) Representative lineages induced at 0h APF for the indicated genotypes. White arrowheads indicate Follicle Cells;
magenta arrows, r1 ECs; yellow arrows, r2a ECs; cyan arrows, FSCs. DAPI, blue; GFP, green; Fas3, red. The r1 EC, r2a EC,
FSC and FC domains are shown in corresponding colors in the horizontal bars for (A-C). (A) A *ptc* EC/FSC/FC lineage
showing a large number of marked cells, likely due to increased division. (B) A *ptc yki* EC-only lineage resembling a

common *yki* phenotype (Fig. 6A), and **(C)** a *ptc yki; tub-yki* EC/FSC/FC lineage with characteristically fewer cells than for *ptc*. **(D-F)** Ovarioles with *ptc* mutations induced at 0h APF produced ovarioles with multilayered FCs (paired white arrowheads) for **(D)** *ptc*, **(E)** *ptc yki* (which has relatively few marked FCs) and **(F)** *ptc yki; tub-yki*. Scale Bar 20 μm for **(A-C)** and 50 μm for **(D-F)**. **(G)** Deduced frequency of the indicated types of single-cell lineage as a percentage of all lineages induced at 0h APF. Error bars show SEM. One asterisk (*) indicates p<0.05; two asterisks (**) indicates p<0.005 (Fisher's two-tailed exact test). Animals occasionally lose the *tub-yki* transgene. Hence, data are reported for all scored ovaries for the *ptc yki; tub-yki* genotype and also as "revised" after removing samples from animals thought likely to have lost the *tub-yki* transgene based on phenotypes (low clone frequency, low frequency of FSCs and FCs) of all examined ovarioles. Number of marked ovarioles = 739 (control), 178 (*yki; tub-yki*), 59 (*ptc yki*), 76 (*ptc yki; tub-yki)*, 65 (revised *ptc yki; tub-yki*), 85 (*ptc*). **(H-K)** Measures of division rate and composition of FSC-containing lineages. **(H)** Average number of ECs in EC-only lineages, and the average sum of EC and FSC number in FSC-containing lineages. SEMs are based on raw data (prior to correction for single-cell lineages). n values (number of lineages scored) are above columns. **(I)** Percentage of cysts (includes region 2b, region 3 and all egg chambers) with adjacent marked FCs among FSC-containing lineages. n values (number of cysts scored) are above columns. **(J)** Percentage of all lineages with marked FSC but no EC, marked EC and FSC, marked FSC with or without EC. Number of marked ovarioles as in (G) for first four genotypes; 95 for *arr; UAS-DIAP1* and 50 for *UAS-Hop UAS-DIAP1*. **(K)** Left: average percentage of all FCs around the terminal egg chamber that are marked among lineages that include terminal FCs (but excluding those with only polar cells, stalk cells or basal stalk cells). Right: percentage of lineages with marked terminal FCs (no exclusions) with marked FCs on the penultimate egg chamber. n values (number of terminal egg chambers scored) are above columns. For raw data and calculations please see supplementary spreadsheets titled 0h APF Graphs Fig4-10; FC occupancy aggregate; Division Measures Hh/Yki**;** Compilation of Numbers Figure 10 Hh Hpo.

We noticed that the *tub-yki* transgene, marked by w^+, is sometimes spontaneously excised in stocks. This is, unfortunately not directly visible in stocks carrying the *yki^{B3}* allele, which itself has strong w^+ expression. We noted, however, that in our initial tests, many lineages expected to be of genotype *yki; tub-yki* showed characteristic *yki* mutant FC phenotypes (a high proportion of polar and stalk cells or elevated Fas3 expression). We therefore obtained the lineage results cited above by constructing a stock that is homozygous for *yki* together with one copy of *tub-yki*, so that the active transgene must be maintained for viability. The unstable *tub-yki* transgene also was lost over time from *ptc yki; tub-yki* stocks, as revealed by crosses to *yki* stocks to test viability, but here there was no solution to enforce *tub-yki* retention. We therefore measured *ptc yki; tub-yki* lineage phenotypes for two recently-derived stocks. In two flies derived from each stock, lineages resembled those seen for *ptc yki* (we score ovaries from individual flies separately). Results are therefore presented without any adjustment, likely representing a mix of *ptc yki; tub-yki* and *ptc yki* genotypes, and also after discarding the four sets of ovarioles very likely lacking *tub-yki*. Results were not greatly altered by the latter step, but we discuss here the second set of results ("revised *ptc yki; tub-yki*"), which likely represents the correct genotype.

The addition of *ptc* (to *yki; tub-yki*) increased the fraction of lineages with FSCs (Fig. 10C, F) from 18% to 23%, both higher than controls (15%) but not significantly (p>0.05) (Fig. 10D, J). The average number of cells per EC/FSC clone (7.4) was similar to the *yki; tub-yki* genotype (6.6) and to controls (8.6) but much lower than for *ptc* (13.4) (Fig. 10H). To probe division rate further in the neighborhood of precursors that can yield FSCs we also counted the frequency of marked FC patches in FSC-containing lineages. The very high proportion of germline cysts contacting marked FCs for *ptc* lineages (99/124; 80%), was not matched by *ptc yki; tub-yki* (38/56; 68%), yielding a value similar to the direct control for that experiment (35/51; 69%), slightly higher than the average of six such controls (121/209; 58%), and higher than for *yki;tub-yki* (98/209; 47%) (Fig. 10I). Thus, marked cells in FSC-containing lineages were less abundant in *ptc yki; tub-yki* ovarioles than for *pt*c when assessed both by EC/FSC numbers and a measure of FC production, suggesting that the faster division promoted by *ptc* is largely suppressed by *yki* substitution with *tub-yki*. That is consistent with the

possibility that *ptc* increases division of precursors in intermediate FSC-producing locations largely
through transcriptional induction of *yki*, as in adult FSC lineages. The average terminal FC clone size
(35%) was very similar for *ptc yki; tub-yki* and the direct control (36%) but higher than for *yki; tub-yki*
(29%) in the same experiment (Fig. 10K). The percentage of terminal FC clones that spread to the
penultimate egg chamber was higher for *ptc yki; tub-yki* (63%) than both the control (55%) and *yki;*
tub-yki (48%) (Fig. 10K). Thus, there may be a small loss of division of posterior precursors for *yki; tub-*
yki and a small increase due to loss of *ptc* that does not rely on *yki* transcriptional induction. The
phenotype of marked *ptc yki; tub-yki* FCs included multilayering and the accumulation of cells
between egg chambers (Fig. 10F), confirming that these characteristics are caused by Hh signaling
without using *yki* as an intermediate.
The fractions of lineages including r1 ECs (47% vs 67%) and of EC-only lineages (48% vs 63%)
were lower than controls, while the frequency of FC-only lineages was increased (22% vs 17%) for *ptc*
yki; tub-yki, as observed for *ptc* alone (Fig. 10G). However, terminal FC labeling among all lineages
was similar to controls (20% vs 22%), unlike *ptc* (34%). These results suggest that the posterior bias
produced by excessive Hh signaling is not dependent on *yki* induction for precursors other than the
most posterior, potential terminal FC precursors. It is interesting that loss of *yki* can lead to reduced
terminal FC representation, while excess Yki increased representation (Fig. 8I). Those results are
consistent with the possibility of *ptc* increasing terminal FC representation by inducing *yki*.
Conversely, it is possible that reduced *yki* induction in *smo* mutant cells reduces terminal FC
representation. This might balance a *yki*-independent consequence of *smo* that produces net
posterior movement to result in the observed *smo* phenotype of posterior displacement of all but the
most posterior clones. If correct, *smo* and *ptc* would have opposite *yki*-dependent changes in
terminal FC formation, with Yki acting positively to promote terminal FC identity. For more anterior
precursors, *smo* and *ptc* would both promote posterior movement in a Yki-independent manner.
Increased posterior movement in the EC/FSC precursor domain, together with a roughly
normal rate of cell division might account for the unusually high proportion of *ptc yki; tub-yki* lineages
with labeled FSCs but no labeled ECs (15% vs 2% for controls, Fig. 10G, J). The other genotypes with a
similar phenotype were *arr* (loss of Wnt signaling) and *UAS-DAP UAS-Hop* (increased JAK-STAT with
an inhibitor of increased division), which causes posterior movement of all precursors with no
apparent effect on cell division (Fig. 10J). The phenotype is not shared by *ptc*, presumably because
increased division additionally replenishes EC precursors in EC/FSC lineages to balance posterior loss.
Thus, although FSC-containing lineages are not much more abundant for *ptc* than for *ptc yki; tub-yki*,
the contribution of increased division rate, which appears to rely on *yki* induction, is manifest by the
markedly different frequencies of EC/FSC lineages (Fig. 10G and 10J).
1769
1770
1771

Discussion

The replenishment of lost differentiated cells by adult stem cells is a vital physiological function for tissues with high cellular turnover, including human epidermal and intestinal cells. Adult stem cells must execute this function over a lifetime, overcoming natural variations and occasional extreme challenges, while their necessary continued replication and potential longevity make stem cells vulnerable to the accumulation of cancer-causing mutations. Adult stem cell studies are slowly revealing the cellular organizations and behaviors selected by these evolutionary pressures. In one large class of paradigms, which includes Drosophila FSCs and mammalian intestinal stem cells, stem cell division and differentiation are independent (JONES 2010; REILEIN *et al.* 2017; CLEVERS AND WATT 2018; REILEIN *et al.* 2018; BEUMER AND CLEVERS 2021; GREULICH *et al.* 2021; KALDERON 2022). Hence, maintenance of a specific stem cell requires that division competes successfully with differentiation; this balance can be altered by genetic mutation, to produce a functional stem cell deficit or a hyper-competitive lineage with oncogenic potential. Moreover, differentiation in these paradigms is governed by the spatial distribution of extracellular signals and is initiated by changes in cell location (along the AP axis for FSCs and the crypt bottom/top axis for intestinal stem cells).

Equally relevant, but less studied, are the developmental processes that create the requisite organization of stem cells and niche cells that produce key extracellular signals. Drosophila FSCs belong to a class of paradigm where stem cells are not set aside during development (GUIU *et al.* 2019; OBERNIER AND ALVAREZ-BUYLLA 2019; BOND *et al.* 2021; REILEIN *et al.* 2021). Instead, a set of precursors simultaneously produces stem cells and the adult differentiated cells they will support. If differentiated cells are continuously irreversibly lost from the precursor pool through migration away from the future stem cell domain, as is true for FCs in the pupal Drosophila ovary and likely holds also for mammalian intestine, the survival of a specific precursor to produce a stem cell may depend on sufficient division activity during development to counter differentiation losses (Fig. 11 and Fig. 12). Also, if stem cell division and migration (leading to differentiation) must be balanced during development and in adults, the use of a similar set of extracellular factors might be expected. Our studies of the responses of single cells to specific genetic manipulations during Drosophila ovary development support both of these hypotheses (Fig. 11). These conclusions, based on experimental evidence for FSCs, are likely to apply generally to tissues supported by stem cells that develop alongside the mature tissue and act as a community with independent regulation of division and differentiation (Fig. 12).

Origin of the first FCs

It was previously shown that most FC-producing lineages induced at 36h APF contributed either no terminal FCs or only terminal FCs, suggesting segregation of terminal FC precursors by that stage (REILEIN *et al.* 2021). This deduction was supported by live imaging, showing that only cells posterior to germline cells (in the Extra-Germarial Crown and Basal Stalk) from about 48h APF populated the terminal egg chamber (REILEIN *et al.* 2021). Some of these cells accumulated anterior to the terminal egg chamber, destined to become FCs on the penultimate egg chamber, consistent with lineage studies. By contrast, FCs in more anterior egg chambers, together with ECs and FSCs, derived from germarial ICs. In this work, we found that lineages induced at pupariation often included marked FCs in the terminal egg chamber and also anterior to the penultimate egg chamber. The high frequency of those occurrences could not be explained by chance induction of two separate lineages; instead, most such occurrences report the behavior of single-cell lineages. Hence, we deduce that

somatic cells intermingled with germline cells (ICs) in the developing germarium at pupariation can
give rise to adult somatic ovary cells in every location, from the most anterior ECs to the most
posterior FCs and the basal stalk protruding from the posterior of the terminal egg chamber (Fig.
11A). Accordingly, the terminal FC precursors seen accumulating over the next 48h must largely
derive from migration of ICs out of the germarium, distancing themselves from germline cells, even
though such movements have not been directly observed by live imaging. Our lineage studies cannot
exclude the possibility that some of the terminal FC population derive from a separate source but
there is no specific evidence supporting that possibility. It seems most likely that all terminal FCs
derive from ICs at pupariation.
The alternative potential source of terminal FCs previously considered were "basal cells",
already present posterior to developing germaria at pupariation but without the characteristic Fas3
expression of cells in the subsequently enlarging Extra-Germarial Crown. Recent studies identified a
population of cells, named Swarm cells, in larval ovaries, according to selective expression of Single-
minded and Crossveinless-2 (Banisch *et al.* 2021). By using corresponding antibodies and GAL4 lines
for lineage tracing, it was shown that these apical cells migrate basally from 72h to 120h after egg-
laying to form the "basal cell" population at the larval/pupal transition. Swarm cell migration is
initiated by an ecdysone pulse, while a second ecdysone pulse triggers the now-posterior Swarm cells
to send a differentiation signal to posterior PGCs. Thereafter, Swarm cell lineages are lost during
pupal development, leaving little residue in adults, consistent with an earlier surmise of basal cell
death during pupation from electron micrographs (King *et al.* 1968). These deductions are consistent
with our conclusion that ICs, and not basal cells, are the precursors of terminal FCs.
1839
1840 **Cell division rate strongly affects the likelihood of a precursor becoming a FSC**
1841
Division rate is a key parameter for adult stem cells. For FSCs and all adult stem cells
maintained by population asymmetry with division-independent differentiation, a genetic change
that alters the division rate of a single stem cell necessarily has a strong effect on the likelihood that
the stem cell lineage will amplify, survive or be extinguished (Reilein *et al.* 2018). A stem cell with
increased division rate will likely be maintained, with an elevated chance of amplifying to occupy an
entire niche, immortalizing the genetic change and possibly seeding a cancer. Conversely, a stem cell
with reduced division that cannot contribute efficiently to the physiological function of the stem cell
community tends to be eliminated quite rapidly. We found experimentally that there is an analogous
effect of division rate on the likelihood of becoming an FSC during pupal development. Thus, very few
precursors with reduced division due to *cycE* or *cutlet* mutations became FSCs (Fig. 11B), while a
greater proportion than normal became FSCs as a consequence of faster division due to excess CycE
or increased JAK-STAT or Hh signaling (Fig. 11C). The likely rationale for this outcome is similar to the
situation for adult FSCs. Precursors in the A/P territory of a developing germarium that may become
FSCs are continually depleted during pupation by irreversibly becoming FCs (Fig. 11A). This loss is
potentially balanced by influx from more anterior positions and by precursor division. In the adult
germarium there is no net flux of ECs to become FSCs (Kalderon 2022). It appears that the major
means of FSC precursor survival during pupal development is also through replication, even though
there is net posterior expansion of the germarium early in pupation and hence, the potential for
substantial replenishment of the FSC precursor domain from the anterior. Thus, the competitive
advantage of a faster-dividing potential FSC actually initiates even prior to FSC formation, likely
beginning at pupariation. The genetic factors that regulate division rate may differ between pupal
and adult stages but it appears from our studies that several are in common (*cycE*, *cutlet*, JAK-STAT

and Hh signaling) (Fig. 11B, C). These findings can explain the stronger effect on FSC lineage
prevalence of mutations that increase or reduce division rates when induced in larvae rather than
adults, as noted long ago for alterations to Hh signaling (Vɪᴇᴅ *et al.* 2012). An analogous principle may
hold for some human stem cells with division-dependent differentiation, allowing the amplification of
potential cancer-initiating mutations even before adulthood or full tissue maturation.
1869
1870 **Regulation of cell division rate during pupation**
1871
There is prominent spatial regulation of somatic ovary precursor cell division rate during pupal
development, with EdU incorporation and FUCCI cell cycle reporters providing evidence of lower
rates and early cessation of division anteriorly (Rᴇɪʟᴇɪɴ *et al.* 2021). Indeed, this spatial regulation of
division is potentially instrumental in setting the appropriate number of adult ECs. Unfortunately, we
can only measure division rate in selected regions and with some limitations during pupation,
counting ECs in EC-only lineages, the ECs and FSCs of EC/FSC/FC lineages, and FCs in terminal FC
lineages. These measures sufficed to provide reliable evidence of reduced or increased division in
response to genetic manipulations but cannot provide precise details of changes in the spatial
distribution of division rates. Thus, we have evidence of decreased division for loss of function
alterations to *cycE*, *cutlet*, *smo* and *yki*, together with increased division for excess CycE, activated Yki,
ptc and increased JAK-STAT signaling, measured over the whole duration of pupation. Very little
information was revealed for alterations, like loss of *stat* and excess Wnt pathway activity, which
resulted in the production of almost exclusively ECs.
Since Hh pathway activity had a positive effect on precursor division and Hh pathway activity
is known to decline from anterior to posterior in pupal ovaries (Lᴀɪ *et al.* 2017), this pathway cannot
contribute to the greater division rate of posterior cells. Excess JAK-STAT activity produced lineages
with very large numbers of marked cells, partly obscuring the likely A/P position of the original
precursor. Nevertheless, there were very few examples of the highly prevalent (>50%) typical wild-
type EC-only clone with just 2-3 labeled cells. It seems clear, therefore, that excess JAK-STAT signaling
increased division of even the most anterior precursors. Thus, it is surely important to limit JAK-STAT
pathway activity in anterior precursors. Once an egg chamber has budded to produce polar cells,
which are a key source of JAK-STAT pathway ligand, we observed strong pathway reporter signal in all
egg chambers and the posterior half of the germarium, as in adults (Vɪᴇᴅ *et al.* 2012; Mᴇʟᴀᴍᴇᴅ ᴀɴᴅ
Kᴀʟᴅᴇʀᴏɴ 2020). It therefore seems inevitable that, as in the adult, JAK-STAT pathway activity
promotes division of posterior FSC/FC precursors but not anterior EC precursor germarial cells during
the second half of pupation.
A key question is whether JAK-STAT is an important stimulator of FSC or FC precursor division
over the first 48h. There were few or no FSC- or FC-containing *stat* mutant lineages to measure cell
numbers directly. Moreover, the severe loss of *stat* FSC lineages, which might be caused by reduced
division rates, by anterior precursor movement or both, could have occurred during the second half
of pupation. We did not see a strong JAK-STAT pathway reporter signal prior to 30h APF. Thereafter,
however, reporter signal was stronger in posterior germarial cells and extra-germarial cells than in
the anterior half of the germarium (Fig. 7), so it is plausible that graded JAK-STAT pathway activity
may also contribute to faster division of more posterior cells over the first half of pupation (Fig. 12).
Inferred division rates of germarial pupal precursors are roughly inversely correlated to Wnt
pathway activity reported by Fz3-RFP. Thus, EC precursors have low or undetectable pathway activity
only early in pupation while anterior precursors are still dividing; more posterior precursors
consistently have only low Wnt pathway activity and do not arrest. Loss of Wnt pathway activity (*arr*)

did not, however, increase the average size of EC-only lineages or EC/FSC lineages. Almost all *axn*
mutant precursors become ECs, so it is not known whether abnormally high (and potentially supra-
physiological) Wnt pathway activity would impede posterior precursor cell division, as found for adult
FSCs (MELAMED AND KALDERON 2020). Thus, despite the spatial correlation between higher Wnt
signaling levels and lower division rates, we have no direct evidence of Wnt signaling reducing
precursor division rates during pupation.

Figure 11: Summary of key influences on FSC formation and formation of the first FCs
(A) Ovariole diagram with ECs in yellow, FSCs in cyan, specialized stalk and polar cell FCs in green and other FCs in white.
Representative precursors of EC-only (yellow), EC/FSC/FC (red) and terminal FC lineage (green) precursors are shown for
oh APF. EC-only precursors are the most abundant, so two yellow cells are shown. An intermediate developmental time
(such as 48h APF) is shown for EC/FSC/FC precursors to highlight the increased A/P range of derivates destined to become
ECs (yellow) and FSCs (cyan), as well as the constant loss of EC/FSC precursors to become FCs to the posterior (horizontal
arrow). The more posterior EC/FSC precursors divide faster than anterior EC/FSC precursors, especially later in pupation
(larger renewal arrow). Our studies show that this renewal is important for FSC precursors to persist, compensating for

losses from continued FC formation. **(B-D)** Differences from wild-type behavior that affect precursors becoming FSCs. Experimental results are summarized and the underlying rationalization is illustrated (showing only relevant precursors). (**B**) Cell-autonomous reduction of division rate of an EC/FSC/FC (red) precursor among other wild-type precursors leads to progressive loss of FSC precursors throughout pupation because FSC precursor division no longer matches losses from differentiation to FCs. Consequently, many potential EC/FSC/FC lineages become EC/FC lineages. This behavior was seen most clearly for reduced activity of *cycE* and *cutlet*, but also for *smo* and *yki* when apoptosis was inhibited. **(C)** Conversely, a genetic cell-autonomous increase in division rate relative to other precursors leads to a progressive increase of FSC precursors during pupation, ensuring survival of FSC precursors and giving rise to EC/FSC/FC lineages with more FSCs than usual. Experimentally, this condition increased the frequency of EC/FSC/FC lineages, implying that some potential FSC lineages are normally lost in wild-type ovarioles because of insufficient division. This behavior was seen most clearly for increased activity of *cycE*. Similar results were seen for excess activated Yki (but not for regulators of Hippo pathway activity expected to increase Yki activity) and lesser, similar changes were observed for *ptc*. Excess JAK-STAT (*UAS-Hop*) produced many FSC-containing lineages but there were too many marked cells altogether to quantify lineage results. (**D**) A genetic change deduced to cause cell-autonomous posterior movement of all precursors (as seen for *arr*) resulted in fewer EC-only lineages, more FSC-containing lineages and more FC-only lineages. The increase in FSC-producing is because there are initially more potential EC-only (yellow) precursors than EC/FSC/FC precursors. Hence, a posterior shift relative causes more precursors originally in EC-only locations to move into FSC-producing territory than the number of precursors moving out of EC/FSC/FC territory to become FCs. Many of the resulting FSC-containing lineages lacked ECs. That is presumably because precursors entering the EC/FSC/FC-producing domain can continue to move posteriorly through that domain (red arrows) and are sustained, in part, through normal rates of cell division. Division continues in the FSC-producing (cyan) region but not in the EC-producing (yellow) region in the second half of pupation, so that many lineages maintain FSCs but not ECs. A high proportion of FSC lineages without ECs was seen also for *UAS-Hop; UAS-Dap* and *ptc yki; tub-yki* genotypes, supporting the deduction that they cause posterior movement around the FSC precursor region without affecting cell division rate. **(E)** The most posterior ICs at pupariation (green in (**A**)) were deduced from lineage studies here to migrate out of the germarium over the first 48h of pupation and then become the first FCs, occupying the terminal egg chamber (and sometimes also the penultimate egg chamber). The frequency of this outcome (producing Terminal FCs) was greatly decreased by loss of JAK-STAT or increased Wnt signaling; it was increased by reciprocal changes in those pathways. Reduced terminal FC contributions in the absence of *yki* and *smo* activity appeared to be due to spatially selective apoptosis. Increased Hh signaling (*ptc*) increased terminal FC lineage frequency, apparently dependent on normal *yki* gene regulation.

Wnt signaling promotes anterior migration

We found strong and consistent alterations in precursor outcomes for genetic alterations of Wnt pathway activity. Loss of activity caused an extensive posterior shift in outcomes, with a large increase in lineages containing terminal FCs and only FCs, together with a large reduction in lineages containing only ECs or containing any r1 ECs. The frequency of lineages containing FSCs was substantially increased and an unusually high fraction of FSC lineages did not include any ECs. This is an interesting outcome: loss of Wnt signaling causes increased FSC loss in adults (because of greater conversion to FCs and apparently no replenishment from the anterior) but increased FSC production during pupal development. This can be rationalized on the basis that there are normally many more EC-only than FC-only precursors (63% vs 17%), so more cells have the potential to enter FSC-producing regions from the anterior than from the posterior (Fig. 11D). Moreover, continued posterior movement appears to convert many potential EC/FSC/FC lineages to FSC/FC lineages. The resulting high frequency of FSC/FC lineages plausibly constitutes a distinguishing characteristic of genotypes that promote posterior migration without altering division rates for a given location (Fig. 10J and 11D).

Increased Wnt pathway activity produced strong phenotypes, which were broadly the converse of those observed for loss of Wnt pathway activity (more EC-only lineages; fewer FC-only and terminal FC-containing lineages). Amongst EC outcomes, there was only a small shift in favor of r1 ECs (contrasting with the large converse effect of loss of pathway activity), suggesting that

precursor migration in the most anterior regions of the developing germarium, where Wnt signaling
appears to be roughly uniform, may not normally be driven by differential Wnt pathway activity.
There was also a reduction in *axn* FSC-containing lineages, all of which included ECs, from 15% to 6%.
If anterior displacement caused all prospective FSC precursors to become ECs and all FC-precursors to
form FSC-containing lineages, there would be a net gain in FSC lineages because the initial FC-only
precursor pool is larger than the FSC precursor pool. Since FSC lineages are depleted we might infer
that FC-only precursors migrated beyond the FSC domain to form EC-only lineages. An additional
contribution to FSC loss could be a reduction in cell division rate. There were too few FC- and FSC-
containing *axn* lineages to assess division rate in these posterior regions. However, adult FSC division
is severely reduced by loss of *axn* (REILEIN *et al.* 2017; MELAMED AND KALDERON 2020) and the properties
of FSC precursors during the last 2 days of pupation are likely highly similar to adult FSCs because
they appear to exist in a very similar environment. Hence, some contribution of reduced division rate
to the observed paucity of FSC-containing lineages is likely.
We attempted to measure the temporal effects of Wnt pathway activity by initiating genetic
alterations at different times during pupation. These results cannot easily be interpreted
quantitatively but indicated directly that Wnt signaling influenced outcomes in the latter stages and
indirectly (inferred from greater severity when initiated early) that Wnt signaling levels were also
influential early in pupation. Also, since terminal FC fates were greatly reduced for *axn* lineages
induced at pupariation but not when induced 36h later, we infer that the primary deficit is likely
migration of precursors posteriorly out of the germarium into the EGC and basal stalk during the first
48h of pupation. Thus, terminal FC formation relies on limiting Wnt pathway activity (Fig. 11E), as
found for FC production from adult FSCs (REILEIN *et al.* 2017; MELAMED AND KALDERON 2020). Throughout
pupal development, Wnt pathway activity, reported by Fz3-RFP, declined to undetectable levels
roughly coincident with the anterior extent of strong Fas3 expression, which appears to mark FC
precursors. Thus, very low or absent Wnt pathway activity appears to be a significant factor in
allowing FC formation throughout pupation, as in adults.
2004
2005 **JAK-STAT signaling promotes posterior cell migration**
2006
The situation for JAK-STAT pathway signaling is roughly the mirror-image of that described for
Wnt signaling, with a couple of significant differences (Fig. 11). First, increased JAK-STAT activity
greatly stimulated precursor division. Second, although increased JAK-STAT activity (together with
UAS-Dap to block changes in cell division rate) reduced the incidence of lineages with r1 ECs, loss of
stat activity did not have the converse effect. Thus, even the most anterior precursors can be shifted
more posterior in response to JAK-STAT but slightly more posterior EC precursors do not rely on JAK-
STAT pathway activity to prevent anterior movement (unlike FC precursors and FSC precursors). The
most likely explanation is that JAK-STAT activity is low, as observed directly, and functionally
insignificant in the anterior half of the germarium throughout pupation. Interestingly, as for *arr*
mutations, there was an increase in FSC-containing lineages lacking marked ECs for the *UAS-Hop UAS-*
Dap genotype, consistent with the phenotype originating from posterior migration from EC-producing
regions, rather than an increase in division rate (there was no indication of increased division for this
genotype).
The large deficit of *stat* mutant lineages with marked terminal FCs (Fig. 11E) was a particularly
important finding. The deficit was not observed for lineages induced 36h after pupariation and
therefore appears to originate from a failure of precursors to migrate posteriorly out of the
germarium over the first 48h of pupation. This is consistent with JAK-STAT pathway activity

universally favoring FC formation in adults and pupae. The *stat* mutant phenotype suggests that low
Wnt pathway activity, as seen in the most posterior pupal precursors, is not sufficient for normal
terminal FC production, although the condition of eliminating both Wnt and JAK-STAT pathway
simultaneously has not been directly tested.

In adults there is always a nascent or just-formed egg chamber budding from the germarium
with anterior polar cells producing a JAK-STAT ligand that produces a posteriorly-biased gradient in
the vicinity of FSCs to promote FC formation (VIED *et al.* 2012). Prior to formation of the first egg
chamber there are no polar cells as a source of ligand. The genetic results show that STAT must have
significant activity during this period because very few precursors lacking *stat* activity become
terminal FCs. The small number of mutant terminal FCs formed were found in different locations in
different ovarioles. Had all marked FCs been found, for example, in the basal stalk, that might have
suggested that the most posterior precursors do not need to respond to a JAK-STAT signal to
maintain their position and might therefore be the source of JAK-STAT pathway ligand.

We were able to detect pathway activity from 30h APF onwards. Activity was strongest in
regions surrounding and posterior to the most posterior germline cyst, consistent with the possibility
that this heightened activity promotes posterior migration of precursors out of the germarium prior
to production of the first egg chamber. It will be important to determine which cells might be a
source of ligand for the JAK-STAT pathway in the period prior to polar cell production. The pattern of
pathway activity at 48h APF is consistent with the possibility of production of ligand by Fas3-positive
cells (and/or efficient passage through Fas3-positive cells) and is similar, albeit weaker, to the pattern
in adult germaria. That observation raises the question of whether the graded JAK-STAT pathway
activity in adult germaria results entirely from ligand produced in polar cells or might also be
contributed by early Fas3-positive FCs. Selective inhibition of Upd ligand production in adult polar
cells using *neur-GAL4* strongly reduced STAT-GFP with partial penetrance (around 70%), while
inhibition with *109-30-GAL4*, which is expressed in all early FCs and slightly higher in polar cells, was
also effective (VIED *et al.* 2012). Those results argue strongly for a primary role of polar cells but also
allow the possibility that early FCs may provide an additional source of JAK-STAT pathway ligand in
adult germaria.

2052

2053 **Hedgehog signaling and Yki as an intermediate; division rate and some location effects**
2054

The contributions of Hh signaling and Yki activity to the behavior of adult germarial cells are
relatively clear. Both Hh signaling and Yki promote FSC division and are required to limit apoptosis,
while both are necessary in ECs to limit BMP production and hence allow normal germ cell
maturation. In FSCs, Hh pathway activity can increase Yki activity through transcriptional regulation of
yki (HUANG AND KALDERON 2014), while in ECs different reports suggested independent actions of Hh
and Yki (HUANG *et al.* 2017) or post-transcriptional regulation of Yki activity by Hh (LI *et al.* 2015).

Manipulation of Hh signaling and Yki activity during pupal development also produced some
related phenotypes, reminiscent of effects on adult FSCs. Both increased Hh signaling (*ptc* mutation)
and increased activated Yki (*UAS-YkiS168A*) expression increased measures of precursor division and
increased FSC representation, as would be predicted as a consequence (Fig. 11C). However, Hpo
pathway inactivation (via *hpo*, *wts* and *kibra*), which is expected to activate Yki post-transcriptionally,
did not appear to increase division or FSC representation consistently. The increase in semi-
quantitative measures of precursor division and of FSC representation observed for *ptc* were mostly
reduced when *tub-yki* substituted for *yki*, suggesting that the increased division rate may, as in adult
FSCs, depend on transcriptional induction of *yki*. However, there were exceptions, limiting the

evidence that responses to *ptc* require normal transcriptional regulation of *yki*. Moreover, the
inferred division rates and overall distribution of cell types for the *yki; tub-yki* genotype were quite
similar to controls, suggesting that transcriptional regulation of *yki* activity can be substituted quite
well by a constitutive promoter, as in adult FSCs.
There was a clear loss of *yki* mutant lineages that was suppressed by expressing DIAP1 to
inhibit apoptosis but this appeared only to affect posterior precursors giving rise to FC lineages. The
paucity of *yki; UAS-DIAP1* FSC-containing lineages was likely due primarily to reduced precursor
division because it was suppressed by adding excess CycE together with DIAP1 (in lineages initiated
2d before pupariation to allow timely initiation of expression of the *UAS*-driven transgenes in MARCM
lineages), strongly suggesting that Yki normally contributes to requisite precursor division rates.
A variety of other observations suggest some additional contributions of Hh signaling and Yki
activity but these deductions are tempered by the limited magnitude or significant variability in some
of the noted effects. First, there did appear to be some consequences for A/P movement of
precursors. Loss of *yki* in the presence of *UAS-DIAP1* did not result in any notable changes of
representation of r1 ECs but did reduce the representation of terminal FCs in lineages initiated 2d
before pupariation. Increased Yki activity consistently (*kibra*, *wts*, *hpo*, *UAS-YkiS168A*) reduced r1 EC
representation but only *UAS-YkiS168A* modestly (and not significantly) increased terminal FC
production. Increased Hh signaling modestly decreased r1 EC representation, while increasing
terminal FC representation. The effect on r1 ECs was retained when *yki* was substituted by *tub-yki*,
while the effect on terminal FC representation was lost. This suggests that increased Hh signaling
reduces r1 EC production independent of Yki, even though increased Yki activity has the same effect,
while terminal FC representation, which is modestly increased by *UAS-YkiS168A* and decreased by
loss of *yki*, is only increased by excess Hh signaling when transcriptional regulation of *yki* is allowed.
These deductions align with earlier conclusions about adult ovaries: Hh signaling acts through Yki to
influence more posterior cells (FSCs), while Hh and Yki act independently, albeit with similar
consequences, for anterior cells (ECs) (HUANG AND KALDERON 2014; HUANG *et al.* 2017).
The interpretation of *smo* phenotypes, representing loss of Hh pathway activity, has an
additional caveat. Clone frequency was lower than for controls and was not restored by *UAS-DIAP1*,
even though lineage phenotypes were altered by DIAP1. Thus, there is significant precursor loss or
death (either non-apoptotic or apoptosis is not efficiently rescued by *UAS-DIAP1*) with unknown
spatial selectivity. In *smo; UAS-DIAP1* lineages there was a loss of FSCs, plausibly due to limited cell
division (Fig. 11B). This would be expected to increase the frequency of FC-only or EC/FC lineages. FC-
only lineage frequency was increased but beyond that expected from FSC loss alone, and r1 EC
representation was modestly reduced, with no change in terminal FC representation. Thus, loss of Hh
signaling, at least among surviving cells, appears to cause a modest posterior shift over most of the
precursor domain, excluding the most posterior regions. This deduction stands alongside a deduction
of (stronger) posterior shifts also for increased Hh signaling. The latter deduction relates to potential
consequences of Hh signaling, whereas the former reports the normal influence of Hh signaling. Thus,
with significant caveats, it appears that Hh signaling might normally promote retention of anterior
positions for most of the precursor domain. This is easily reconciled with the anterior bias of Hh
pathway activity and the likelihood that posterior precursors soon fall out of the range of Hh
signaling.
Another study examined the consequences of manipulating Hh pathway activity in all somatic
cell ovary precursors (rather than clonally) and concentrated mainly on genetic changes initiating
prior to pupariation, to conclude that Hh signaling promotes retention of ICs in the developing gonad
(LAI *et al.* 2017). It is conceivable that the shortfall of *smo* lineages we observed is due to migration of

some precursors away from the developing ovary. However, such losses might plausibly affect the
most posterior precursors most strongly, and we found no selective shortage of terminal FC
representation in *smo; UAS-DIAP1* lineages. Hence, we favor limited cell death as a more likely
explanation. The effects of Hpo pathway inhibition (to increase Yki activity) have also been reported
for early ovary development using manipulation of gene activity in all precursors, leading to a
conclusion that the IC pool was increased in response, plausibly due to increased division (SARIKAYA
AND EXTAVOUR 2015). This result is commensurate with our deductions from clonal analyses with UAS-
YkiS168A and loss of *hpo*, though neither *kibra* nor *wts* mutations elicited a discernible indicator of
increased division.
2125
2126   **Summary conclusions and open questions**
2127
In the adult ovary, FSCs are maintained within a narrow A/P domain. At the posterior edge of
that domain, some FSCs move posteriorly into territory with lower Wnt signaling and higher JAK-STAT
signaling to become FCs (Fig. 12). Genetic reduction of Wnt or increase of JAK-STAT signaling greatly
increase the chance, cell autonomously, of an individual FSC becoming an FC, suggesting that the
magnitudes of these pathways are major drivers of FSC movement (VIED *et al.* 2012; REILEIN *et al.*
2017; MELAMED AND KALDERON 2020). The converse genetic argument holds for FSC conversion to ECs
but the frequency of EC production is much lower than for FCs, perhaps because FSCs less frequently
stray into more anterior territory, which is stably occupied by quiescent ECs, while space to the
posterior of FSCs is constantly vacated by the process of egg chamber formation and budding. We
previously found that the nature or behavior of somatic ovarian precursor cells, like FSCs, appears to
be guided solely by location (REILEIN *et al.* 2021). Thus, the precursors that become FSCs are simply
those that end up in the location of FSCs at the end of pupation.
Here, we found that the external signals guiding the location of precursors during pupal
development overlap extensively with those used in adults (Fig. 12). This is relatively easily
rationalized for Wnt signaling because Cap cells, located at the anterior of the germarium, are already
differentiated at pupariation and appear to be a source of Wnt signals throughout pupation and
adulthood. It is likely that an additional source of Wnt arises progressively during pupation as EC
precursors stop dividing and adopt mature cell characteristics of producing Wnt ligands (SAHAI-
HERNANDEZ AND NYSTUL 2013; WANG AND PAGE-MCCAW 2018) while the distance from Cap cells to FSC and
FC precursors increases (Fig. 12). The net consequence is that from early pupation onwards there is
graded Wnt signaling that declines to undetectable levels at the point of FC specification, which
coincides with the onset of strong Fas3 expression (Fig. 5).
The involvement of JAK-STAT signaling was not predictable because a major source of ligand
in adults are the anterior polar cells associated with an egg chamber budding from the germarium
(Fig. 12). The first egg chamber produced during pupation has no antecedent and there is therefore
no polar cell source of JAK-STAT ligand as a subset of somatic cells move posteriorly out of the
developing germarium during the first 48h to acquire strong Fas3 expression and become the first
FCs. We nevertheless detected JAK-STAT pathway activity around this precursor to FC transition zone
and found that formation of the first FCs is strongly dependent on JAK-STAT activity, just as it
depends on low Wnt activity. We do not know the source of the JAK-STAT ligand or whether pathway
activity is graded over this transition zone, though it is clearly lower in more anterior regions. Hh
pathway activity declines from anterior to posterior in both pupal and adult ovaries (FORBES *et al.*
1996; VIED *et al.* 2012; LAI *et al.* 2017). Although there were some effects of manipulating Hh signaling
on A/P precursor outcomes (contrasting with adult studies, where only cell division rate appears to

be affected), these were much weaker than observed for Wnt or JAK-STAT pathways, suggesting that
the latter two pathways are the major determinants of developmental formation of ECs, FSCs and
FCs, just as they have the greatest influence on FSCs becoming ECs or FCs or remaining as FSCs in
adults.
2166

2167
Figure 12. General principles illustrated by FSC development.
The cartoon depicts a pool of precursors on top and the adult tissue they form at the bottom. If a set of precursors (in the
pale green space) produce both stem cells and differentiated tissue extending away from the source during development,
then the probability that a given precursor will remain in the stem cell zone and become an adult stem cell depends on its
division rate competing successfully against movement out of the stem cell zone. In paradigms with independent division
and differentiation of adult stem cells, the same competition applies to maintaining a given stem cell. Both division rate
and cell movements are likely to be regulated by external signals. It would generally be economical to use largely similar
signals to regulate those processes during both development and adulthood, as we have found for Drosophila ovarian
FSCs. Most simply, this is achieved by signals from niche cells established early in development. In the Drosophila ovary,
two such signals come from Cap cells, which form prior to pupation: Hh promotes division and Wnt promotes anterior
movement. There are two other features of ovary development, which might also be used in other paradigms. First, the
primitive Cap cell niche is elaborated or expanded during pupal development by the production of additional niche cells
(Escort Cells; pale yellow region) that supplement signaling from the anterior by producing additional Wnt ligands.
Second, there is an important signal in adults (activating the JAK-STAT pathway) that originates from differentiated cells
posterior to the stem cells. This signal promotes both stem cell division and posterior exit from the stem cell zone. During
development, the same pathway promotes exit from the stem cell zone and has the potential to stimulate division; our
data do not show whether division is stimulated at the earliest times during pupation or what is the exact cellular source
of JAK-STAT pathway ligand prior to production of the first differentiated cells. Some stem cell paradigms, like mouse gut
crypts, may not have the feature of FSCs that differentiated cells are produced in two directions (the yellow territory may
be absent altogether or not replenished by the stem cells). The general principles still remain that (i) division rate is
important for both development and maintenance of stem cells because it competes with cells moving out of the stem
cell zone, and (ii) niche cells may produce a set of signals that act in a graded fashion over distance to regulate division
rate and movement of both stem cell precursors and their adult stem cell derivatives.
2191

The factors regulating proliferation of pupal precursors also were found to be very similar to
those regulating adult FSC division rate, although the developmental data are much more limited and
semi-quantitative at best. JAK-STAT and Hh signaling, via *yki* induction, are major stimulators of cell
division, while Wnt signal normally has no significant influence (Fig. 12). The role of cell division rate
in producing FSCs during pupal development was also found to be analogous to its role in maintaining
FSCs. In both situations, it appears that the major loss of cells in a location with potential to act as an
FSC is from posterior movement to become FCs. Although some cells may move from an EC-centered

precursor domain to final FSC positions during pupal development, the maintenance of precursors
that become FSCs and of adult FSCs themselves, despite posterior loss, both depend principally on a
sufficient cell division rate. Hence, both the formation and the maintenance or expansion of FSC
lineages is favored by genetic cell-autonomous changes that increase division rate (Fig. 11).
We suggest that two major findings of our study are likely to apply to other stem cell
paradigms that share major characteristics with FSCs (Fig. 12). First, if stem cells of a specific tissue
are specified during the entire period that the rest of the tissue is specified, rather than being set
aside early during development, it is likely that the same signals contribute to stem cell development
and stem cell maintenance in adults. Second, in such scenarios where, additionally, the division and
differentiation of adult stem cells are separate processes, the probability of an appropriately located
precursor becoming a stem cell and of an adult stem cell being maintained both depend on the
division rate of that cell relative to competitors vying to become stem cells or remain as adult stem
cells.
2212
2213

2214 **Data Availability**
2215
2216 Drosophila stocks used in this study are available upon request. The authors affirm that all data
2217 necessary for confirming the conclusions of the article are present within the article, figures, and
2218 supplementary spreadsheets deposited with the article.
2219
2220

2221 **Acknowledgments**
2222
We thank the Bloomington Drosophila stock center, the Developmental Studies Hybridoma Bank
(DSHB), and Laura Johnston (Columbia University) for reagents, FlyBase as an information resource,
and the confocal microscope resource provided by the Department of Biological Sciences, Columbia
University. We thank Aron Choi, Brian Heubel, Pegah Khosravi-Kamrani and Karen Sophia Park for
experimental assistance, David Melamed, Hoyon Kim, Laura Johnston and Iva Greenwald for
discussions.
2229

2230 **Study funding**
2231
2232 This study was funded by NIH grant RO1 GM079351 to DK.

References

Apionishev, S., N. M. Katanayeva, S. A. Marks, D. Kalderon and A. Tomlinson, 2005 Drosophila
Smoothened phosphorylation sites essential for Hedgehog signal transduction. Nature Cell
Biology 7**:** 86-92.
Asaoka, M., and H. Lin, 2004 Germline stem cells in the Drosophila ovary descend from pole cells in
the anterior region of the embryonic gonad. Development 131**:** 5079-5089.
Assa-Kunik, E., I. L. Torres, E. D. Schejter, D. S. Johnston and B. Z. Shilo, 2007 Drosophila follicle cells
are patterned by multiple levels of Notch signaling and antagonism between the Notch and
JAK/STAT pathways. Development 134: 1161-1169.
Bach, E. A., L. A. Ekas, A. Ayala-Camargo, M. S. Flaherty, H. Lee *et al.*, 2007 GFP reporters detect the
activation of the Drosophila JAK/STAT pathway in vivo. Gene Expr Patterns 7**:** 323-331.
Banisch, T. U., M. Slaidina, S. Gupta, M. Ho, L. Gilboa *et al.*, 2021 A transitory signaling center controls
timing of primordial germ cell differentiation. Dev Cell 56**:** 1742-1755 e1744.
Beumer, J., and H. Clevers, 2021 Cell fate specification and differentiation in the adult mammalian
intestine. Nat Rev Mol Cell Biol 22**:** 39-53.
Beumer, J., and H. Clevers, 2024 Hallmarks of stemness in mammalian tissues. Cell Stem Cell 31**:** 7-24.
Biteau, B., C. E. Hochmuth and H. Jasper, 2011 Maintaining tissue homeostasis: dynamic control of
somatic stem cell activity. Cell Stem Cell 9**:** 402-411.
Bond, A. M., G. L. Ming and H. Song, 2021 Ontogeny of adult neural stem cells in the mammalian
brain. Curr Top Dev Biol 142**:** 67-98.
Boyle, M., and S. DiNardo, 1995 Specification, migration and assembly of the somatic cells of the
Drosophila gonad. Development 121**:** 1815-1825.
Chen, H. J., C. M. Wang, T. W. Wang, G. J. Liaw, T. H. Hsu *et al.*, 2011 The Hippo pathway controls
polar cell fate through Notch signaling during Drosophila oogenesis. Dev Biol 357**:** 370-379.
Clevers, H., and F. M. Watt, 2018 Defining Adult Stem Cells by Function, not by Phenotype. Annu Rev
Biochem 87**:** 1015-1027.
Dai, W., A. Peterson, T. Kenney, H. Burrous and D. J. Montell, 2017 Quantitative microscopy of the
Drosophila ovary shows multiple niche signals specify progenitor cell fate. Nat Commun 8**:**
1244.
Duhart, J. C., T. T. Parsons and L. A. Raftery, 2017 The repertoire of epithelial morphogenesis on
display: Progressive elaboration of Drosophila egg structure. Mech Dev 148**:** 18-39.
Eliazer, S., and M. Buszczak, 2011 Finding a niche: studies from the Drosophila ovary. Stem Cell Res
Ther 2**:** 45.
Forbes, A. J., A. C. Spradling, P. W. Ingham and H. Lin, 1996 The role of segment polarity genes during
early oogenesis in Drosophila. Development 122**:** 3283-3294.
Gancz, D., and L. Gilboa, 2013 Insulin and Target of rapamycin signaling orchestrate the development
of ovarian niche-stem cell units in Drosophila. Development 140**:** 4145-4154.
Gancz, D., T. Lengil and L. Gilboa, 2011 Coordinated regulation of niche and stem cell precursors by
hormonal signaling. PLoS Biol 9**:** e1001202.
Gao, S., Y. Zhang and F. Liu, 2023 Revisiting the lineage contribution of hematopoietic stem and
progenitor cells. Development 150.
Giedt, M. S., and T. L. Tootle, 2023 The Vast Utility of Drosophila Oogenesis. Methods Mol Biol 2626**:**
1-36.
Gilboa, L., 2015 Organizing stem cell units in the Drosophila ovary. Curr Opin Genet Dev 32**:** 31-36.

Godt, D., and F. A. Laski, 1995 Mechanisms of cell rearrangement and cell recruitment in Drosophila
ovary morphogenesis and the requirement of bric a brac. Development 121**:** 173-187.
Grammont, M., and K. D. Irvine, 2001 fringe and Notch specify polar cell fate during Drosophila
oogenesis. Development 128**:** 2243-2253.
Greulich, P., B. D. MacArthur, C. Parigini and R. J. Sanchez-Garcia, 2021 Universal principles of lineage
architecture and stem cell identity in renewing tissues. Development 148.
Guiu, J., E. Hannezo, S. Yui, S. Demharter, S. Ulyanchenko *et al.*, 2019 Tracing the origin of adult
intestinal stem cells. Nature 570**:** 107-111.
Hartman, T. R., T. I. Strochlic, Y. Ji, D. Zinshteyn and A. M. O'Reilly, 2013 Diet controls Drosophila
follicle stem cell proliferation via Hedgehog sequestration and release. J Cell Biol 201**:** 741-
757.
Hayashi, Y., Y. Yoshinari, S. Kobayashi and R. Niwa, 2020 The regulation of Drosophila ovarian stem
cell niches by signaling crosstalk. Curr Opin Insect Sci 37**:** 23-29.
He, L., R. Binari, J. Huang, J. Falo-Sanjuan and N. Perrimon, 2019 In vivo study of gene expression with
an enhanced dual-color fluorescent transcriptional timer. Elife 8.
Hsu, T. H., C. Y. Yang, T. H. Yeh, Y. C. Huang, T. W. Wang *et al.*, 2017 The Hippo pathway acts
downstream of the Hedgehog signaling to regulate follicle stem cell maintenance in the
Drosophila ovary. Sci Rep 7**:** 4480.
Huang, J., and D. Kalderon, 2014 Coupling of Hedgehog and Hippo pathways promotes stem cell
maintenance by stimulating proliferation. J Cell Biol 205**:** 325-338.
Huang, J., A. Reilein and D. Kalderon, 2017 Yorkie and Hedgehog independently restrict BMP
production in escort cells to permit germline differentiation in the Drosophila ovary.
Development 144**:** 2584-2594.
Ingham, P. W., 2022 Hedgehog signaling. Curr Top Dev Biol 149**:** 1-58.
Jemc, J. C., 2011 Somatic gonadal cells: the supporting cast for the germline. Genesis 49**:** 753-775.
Jia, J., C. Tong and J. Jiang, 2003 Smoothened transduces Hedgehog signal by physically interacting
with Costal2/Fused complex through its C-terminal tail. Genes & Development 17**:** 2709-2720.
Jones, P. H., 2010 Stem cell fate in proliferating tissues: equal odds in a game of chance. Dev Cell 19**:**
489-490.
Kahney, E. W., J. C. Snedeker and X. Chen, 2019 Regulation of Drosophila germline stem cells. Curr
Opin Cell Biol 60**:** 27-35.
Kalderon, D., 2022 Investigating Adult Stem Cells Through Lineage analyses. Stem Cell Rev Rep 18**:** 2-
22.
Kalderon, D., D. Melamed and A. Reilein, 2021 The number of Follicle Stem Cells in a Drosophila
ovariole. bioRxiv**:** 2021.2010.2025.465475
King, R. C., S. K. Aggarwal and U. Aggarwal, 1968 The development of the female Drosophila
reproductive system. J Morphol 124**:** 143-166.
Kirilly, D., S. Wang and T. Xie, 2011 Self-maintained escort cells form a germline stem cell
differentiation niche. Development 138**:** 5087-5097.
Lai, C. M., K. Y. Lin, S. H. Kao, Y. N. Chen, F. Huang *et al.*, 2017 Hedgehog signaling establishes
precursors for germline stem cell niches by regulating cell adhesion. J Cell Biol 216**:** 1439-
1453.
Lane, M. E., K. Sauer, K. Wallace, Y. N. Jan, C. F. Lehner *et al.*, 1996 Dacapo, a cyclin-dependent kinase
inhibitor, stops cell proliferation during Drosophila development. Cell 87**:** 1225-1235.

Lee, T., and L. Luo, 2001 Mosaic analysis with a repressible cell marker (MARCM) for Drosophila
neural development.[erratum appears in Trends Neurosci 2001 Jul;24(7):385]. Trends in
Neurosciences 24**:** 251-254.
Li, C., L. Kan, Y. Chen, X. Zheng, W. Li *et al.*, 2015 Ci antagonizes Hippo signaling in the somatic cells of
the ovary to drive germline stem cell differentiation. Cell Res 25**:** 1152-1170.
Li, M. A., J. D. Alls, R. M. Avancini, K. Koo and D. Godt, 2003 The large Maf factor Traffic Jam controls
gonad morphogenesis in Drosophila. Nat Cell Biol 5**:** 994-1000.
Little, J. C., E. Garcia-Garcia, A. Sul and D. Kalderon, 2020 Drosophila hedgehog can act as a
morphogen in the absence of regulated Ci processing. Elife 9.
Losick, V. P., L. X. Morris, D. T. Fox and A. Spradling, 2011 Drosophila stem cell niches: a decade of
discovery suggests a unified view of stem cell regulation. Dev Cell 21**:** 159-171.
Margolis, J., and A. Spradling, 1995 Identification and behavior of epithelial stem cells in the
Drosophila ovary. Development 121**:** 3797-3807.
McGregor, J. R., R. Xi and D. A. Harrison, 2002 JAK signaling is somatically required for follicle cell
differentiation in Drosophila. Development 129**:** 705-717.
Melamed, D., A. Choi, A. Reilein, S. Tavare and D. Kalderon, 2023 Spatial regulation of Drosophila
ovarian Follicle Stem Cell division rates and cell cycle transitions. PLoS Genet 19: e1010965.
Melamed, D., and D. Kalderon, 2020 Opposing JAK-STAT and Wnt signaling gradients define a stem
cell domain by regulating differentiation at two borders. Elife 9.
Merkle, J. A., J. Wittes and T. Schupbach, 2020 Signaling between somatic follicle cells and the
germline patterns the egg and embryo of Drosophila. Curr Top Dev Biol 140**:** 55-86.
Misra, J. R., and K. D. Irvine, 2018 The Hippo Signaling Network and Its Biological Functions. Annu Rev
Genet 52**:** 65-87.
Murray, S. M., S. Y. Yang and M. Van Doren, 2010 Germ cell sex determination: a collaboration
between soma and germline. Curr Opin Cell Biol 22**:** 722-729.
Obernier, K., and A. Alvarez-Buylla, 2019 Neural stem cells: origin, heterogeneity and regulation in the
adult mammalian brain. Development 146.
Ohlmeyer, J. T., and D. Kalderon, 1998 Hedgehog stimulates maturation of Cubitus interruptus into a
labile transcriptional activator. Nature 396**:** 749-753.
Panchal, T., X. Chen, E. Alchits, Y. Oh, J. Poon *et al.*, 2017 Specification and spatial arrangement of
cells in the germline stem cell niche of the Drosophila ovary depend on the Maf transcription
factor Traffic jam. PLoS Genet 13**:** e1006790.
Pignoni, F., and S. L. Zipursky, 1997 Induction of Drosophila eye development by decapentaplegic.
Development 124**:** 271-278.
Post, Y., and H. Clevers, 2019 Defining Adult Stem Cell Function at Its Simplest: The Ability to Replace
Lost Cells through Mitosis. Cell Stem Cell 25**:** 174-183.
Reilein, A., H. V. Kogan, R. Misner, K. S. Park and D. Kalderon, 2021 Adult stem cells and niche cells
segregate gradually from common precursors that build the adult Drosophila ovary during
pupal development. Elife 10.
Reilein, A., D. Melamed, K. S. Park, A. Berg, E. Cimetta *et al.*, 2017 Alternative direct stem cell
derivatives defined by stem cell location and graded Wnt signalling. Nat Cell Biol 19**:** 433-444.
Reilein, A., D. Melamed, S. Tavare and D. Kalderon, 2018 Division-independent differentiation
mandates proliferative competition among stem cells. Proc Natl Acad Sci U S A 115**:** E3182-
E3191.
Sahai-Hernandez, P., and T. G. Nystul, 2013 A dynamic population of stromal cells contributes to the
follicle stem cell niche in the Drosophila ovary. Development 140**:** 4490-4498.

Sarikaya, D. P., and C. G. Extavour, 2015 The Hippo pathway regulates homeostatic growth of stem
cell niche precursors in the Drosophila ovary. PLoS Genet 11**:** e1004962.
Slaidina, M., T. U. Banisch, S. Gupta and R. Lehmann, 2020 A single-cell atlas of the developing
Drosophila ovary identifies follicle stem cell progenitors. Genes Dev 34**:** 239-249.
Smelkinson, M. G., Q. Zhou and D. Kalderon, 2007 Regulation of Ci-SCFSlimb binding, Ci proteolysis,
and hedgehog pathway activity by Ci phosphorylation. Dev Cell 13**:** 481-495.
Song, X., G. B. Call, D. Kirilly and T. Xie, 2007 Notch signaling controls germline stem cell niche
formation in the Drosophila ovary. Development 134**:** 1071-1080.
Song, X., and T. Xie, 2003 Wingless signaling regulates the maintenance of ovarian somatic stem cells
in Drosophila. Development 130**:** 3259-3268.
Spradling, A. C., M. de Cuevas, D. Drummond-Barbosa, L. Keyes, M. Lilly *et al.*, 1997 The Drosophila
germarium: stem cells, germ line cysts, and oocytes. Cold Spring Harbor Symposia on
Quantitative Biology 62**:** 25-34.
St Johnston, D., 2023 Polarity and axis formation in the Drosophila female germ line. Curr Top Dev
Biol 154**:** 73-97.
Vermeulen, L., and H. J. Snippert, 2014 Stem cell dynamics in homeostasis and cancer of the intestine.
Nat Rev Cancer 14**:** 468-480.
Vied, C., and D. Kalderon, 2009 Hedgehog-stimulated stem cells depend on non-canonical activity of
the Notch co-activator Mastermind. Development 136**:** 2177-2186.
Vied, C., A. Reilein, N. S. Field and D. Kalderon, 2012 Regulation of stem cells by intersecting gradients
of long-range niche signals. Dev Cell 23**:** 836-848.
Vlachos, S., S. Jangam, R. Conder, M. Chou, T. Nystul *et al.*, 2015 A Pak-regulated cell intercalation
event leading to a novel radial cell polarity is involved in positioning of the follicle stem cell
niche in the Drosophila ovary. Development 142**:** 82-91.
Voog, J., and D. L. Jones, 2010 Stem cells and the niche: a dynamic duo. Cell Stem Cell 6**:** 103-115.
Wang, S., Y. Gao, X. Song, X. Ma, X. Zhu *et al.*, 2015 Wnt signaling-mediated redox regulation
maintains the germ line stem cell differentiation niche. Elife 4**:** e08174.
Wang, X., and A. Page-McCaw, 2014 A matrix metalloproteinase mediates long-distance attenuation
of stem cell proliferation. J Cell Biol 206**:** 923-936.
Wang, X., and A. Page-McCaw, 2018 Wnt6 maintains anterior escort cells as an integral component of
the germline stem cell niche. Development 145.
Wang, Z. A., J. Huang and D. Kalderon, 2012 Drosophila follicle stem cells are regulated by
proliferation and niche adhesion as well as mitochondria and ROS. Nat Commun 3**:** 769.
Wang, Z. A., and D. Kalderon, 2009 Cyclin E-dependent protein kinase activity regulates niche
retention of Drosophila ovarian follicle stem cells. Proc Natl Acad Sci U S A 106**:** 21701-21706.
Ward, E. J., H. R. Shcherbata, S. H. Reynolds, K. A. Fischer, S. D. Hatfield *et al.*, 2006 Stem cells signal
to the niche through the Notch pathway in the Drosophila ovary. Curr Biol 16**:** 2352-2358.
Wiese, K. E., R. Nusse and R. van Amerongen, 2018 Wnt signalling: conquering complexity.
Development 145.
Xi, R., J. R. McGregor and D. A. Harrison, 2003 A gradient of JAK pathway activity patterns the
anterior-posterior axis of the follicular epithelium. Developmental Cell 4**:** 167-177.
Zamfirescu, A. M., A. S. Yatsenko and H. R. Shcherbata, 2022 Notch signaling sculpts the stem cell
niche. Front Cell Dev Biol 10**:** 1027222.
Zeidler, M. P., and N. Bausek, 2013 The Drosophila JAK-STAT pathway. JAKSTAT 2**:** e25353.
Zhang, H., and Y. Cai, 2020 Signal transduction pathways regulating Drosophila ovarian germline stem
cells. Curr Opin Insect Sci 37**:** 1-7.

Zhang, Y., and D. Kalderon, 2000 Regulation of cell proliferation and patterning in Drosophila
oogenesis by Hedgehog signaling. Development 127**:** 2165-2176.
Zhang, Y., and D. Kalderon, 2001 Hedgehog acts as a somatic stem cell factor in the Drosophila ovary.
Nature 410**:** 599-604.
Zhu, C. H., and T. Xie, 2003 Clonal expansion of ovarian germline stem cells during niche formation in
Drosophila. Development 130**:** 2579-2588.

Supplementary Materials

Supplementary Figure 1: Wnt signaling influences over different periods of pupation.

Supplementary Figure 2: JAK/STAT signaling in pupal germaria after stalk and polar cell formation.

Supplementary spreadsheets with raw data and calculations underlying all presented data

Single-cell 0hAPF calculations
All summary clone data for lineages induced at 0h APF together with the calculations converting data to estimations of single-cell lineage data and averaging of all control values and other genotypes tested in more than one experiment.

Single-cell -2dAPF calculations
All summary clone data for lineages induced at -2d APF together with the calculations converting data to estimations of single-cell lineage data and averaging of all control values and other genotypes tested in more than one experiment.

0h APF Significance Fisher and **-2d APF Significance Fisher**
Copies of the two previous spreadsheets (numbers only, no formulae exposing the calculation methods) with additional columns to show p values of significant differences to controls if p<0.05 using Fisher's exact two-tailed test.

0h APF Controls Variation
Calculation of SD and SEM for single-cell lineage parameters among control experiments and comparison to expectations from probabilistic labeling of precursors in different A/P locations.

0h APF Graphs Fig1,3
First sheet is 0h APF master sheet (from the single-cell calculation sheet). Others show data for Figs. 1B-D and Fig. 3I.

0h APF Graphs Fig4-10
First sheet is 0h APF master sheet (from the single-cell calculation sheet). Others show data for Figs. 4, 6, 8H, 9D, 10G

-2d APF Graphs
First sheet is -2d APF master sheet (from the single-cell calculation sheet). Others show data for Figs. 3J, 8I, 9E.

-3.5d axn August 2023 Numbers
Control and *axn* lineages initiated 3.5d before eclosion (36h APF); raw data and deduced frequencies of clone types.

-4d arr DIAP Numbers
Control and *arr; UAS-DIAP1* lineages initiated 4d before eclosion (24h APF); raw data and deduced frequencies of clone types.

Fig. S1 Graph Data
Data for Supplementary Fig. 1.

Fig5 Fz3RFP Graph
Data for Fig. 5 graph of Fz3-RFP Wnt reporter intensities over the germarium at different times.

-3.5d stat November 2021 Numbers
stat lineages initiated 3.5d before eclosion (36h APF); raw data and deduced frequencies of clone types.

FC occupancy aggregate
Raw data and aggregation for fraction of terminal egg chamber occupied by marked FCs and frequency of additional FC labeling in penultimate or more anterior egg chambers for cycE and cutlet genotype lineages (initiated at 0h APF and -2d APF, leading to graph in Fig. 3F), and for selected Hedgehog and Yorkie genotypes (0h APF Hh Yki TermFC). Raw cell number data for EC-only and EC/FSC lineages of select Hh Yki genotypes to estimate SEM (Cell#). Summary data for fraction of available cysts with marked FCs in FSC-containing lineages (%cystsMarkedFCs 0h APF).

Division Measures Hh/Yki
Summary data for graphs in Fig. 10H-K.

Raw data lineage spreadsheets: "**Compilation of Numbers**" named by Figure and time of lineage initiation ("-5d" = 0h APF, "-7d" = -2d APF).
First sheet in a pair is the raw tabulation of marked cells for each sample of a given genotype. The second sheet compiles the raw data to give a variety of measures, including those transferred to the master spreadsheets used to estimate single-cell lineage parameters ("calculations" spreadsheets).