Contrasting roles for IKK regulated inflammatory signalling pathways for development and maintenance of type 1 and adaptive γδ T cells

Farjana Islam[1,2], Cayman Williams[1], Thea Hogan, Ines Boal-Carvalho* and Benedict Seddon[1]*

[1]Institute of Immunity and Transplantation, Division of Infection and Immunity, University College London, The Pears Building, Hampstead, London, UK

[2]Current address : Department of Biochemistry and Molecular Biology, Shahjalal University of Science and Technology, Sylhet 3114, Bangladesh

*Corresponding authors

Running title : IKK regulates γδ T cell homeostasis

Address correspondence to :

Benedict Seddon, Division of Infection and Immunity, UCL Institute of Immunity and Transplantation, The Pears Building, Rowland Hill Street, London NW3 2PP, U.K.

Non-standard abbreviation list

CBM - CARD11 BCL10 MALT1

CIAP - Cellular inhibitor of apoptosis protein

IKK - inhibitor of kappa B kinase

IκB - Inhibitors of kappa B

Nec1 - Necrostatin-1

RIPK1 - Receptor-interacting protein kinase 1

TAK1 - Transforming growth factor-beta activated kinase 1

TNFR1 - Tumour necrosis factor receptor 1

TRADD - Tumour necrosis factor receptor 1-associated death domain protein

TRAF - Tumour necrosis factor receptor-associated factor

Abstract

The inhibitor of kappa B kinase complex (IKK) is a critical regulator of cell death and inflammatory signaling in multiple cell types. Phosphorylation of IκB proteins by IKK results in their degradation and consequent activation of NF-κB transcription factors. RIPK1, a critical cell death regulator, is also a direct target of IKK kinase activity, thereby repressing its cell death activity. In αβ T cells, the RIPK1 kinase activity of IKK is critical for normal thymic development while mature αβ T cells require IKK for both activation of NF-κB dependent survival programmes, and repression of RIPK1. γδ T cells play a unique and versatile role in host immunity with specific effector functions that enables them to act as early responders in immune defense. The role of IKK regulated pathways in their development and survival is not known. Here, we use mouse genetics to dissect the function of IKK and downstream pathways in the normal homeostasis of γδ T cells. We find that IKK expression is critical to establish a replete γδ T cell compartment, but that requires vary between different subsets. Type 1 γδ T cells require IKK dependent NF-κB activation for their generation, while IKK is redundant for development of adaptive γδ T cells. Instead, IKK dependent NF-κB activation is required for their longterm survival. We also find evidence that IKK repression of RIPK1 is required for survival of peripheral but not thymic γδ T cells. Ablation of CASPASE8 did not rescue γδ T cells in the absence of IKK but rather revealed a potent sensitivity of all γδ subsets to necroptosis, that was rescued by kinase dead RIPK1. Overall, we reveal critical requirements for IKK regulated inflammatory pathways by γδ T cells that contrast with those of αβ T cells, and between different subsets, highlighting the complexity of the regulation of these pathways in the adaptive immune system.

Introduction

The NF-κB family of transcription factors play critical roles in controlling development and function of many cell types (Bonizzi and Karin, 2004). Canonical NF-κB signalling is mediated by hetero or homodimers of p50, RELA and cREL family members that are sequestered in the cytoplasm by inhibitory proteins, the Inhibitors of kappa B (IκB) family and the related protein NFκB1. The key regulator of NF-κB dimer release is the inhibitor of kappa-B kinase (IKK) complex, a trimeric complex of two kinases, IKK1 (IKKα) and IKK2 (IKKβ), and a third regulatory component, NEMO (IKKγ). IKK phosphorylates IκB proteins, targeting them for degradation by the proteasome and releasing NF-κB dimers to enter the nucleus.

In αβ T cells, activation of NF-κB is a well-recognised critical early event during T cell receptor antigen recognition (Gerondakis and Siebenlist, 2010). In the absence of REL subunits, or upstream NF-κB activators, such as TAK1 or IKK complex, T cells fail to blast transform or enter cell cycle (Webb et al., 2019; Xing et al., 2016). Consequently, mice with T cell-specific ablation of RELA and/or cREL have peripheral naive T cells, but lack effector or memory phenotype T cells (Webb et al., 2019; Zheng et al., 2003b). As well as regulating activation, the NF-κB signaling pathway also has important functions in the development and survival of αβ T cells. While activation of NF-κB by TCR appears redundant for normal selection of thymocytes during thymic development (Schmidt-Supprian et al., 2004; Webb et al., 2019), signals from TNF and other TNF receptor superfamily members (TNFRSF) are important for survival and differentiation of post selection single positive thymocytes. NF-κB is required for re-expression of *Il7r* following thymic development, that is necessary for long term IL-7 dependent survival of peripheral naive T cells (Miller et al., 2014; Silva et al., 2014). Additionally, long-term survival of fully mature naive CD4[+] T cells is dependent upon tonic NF-κB since CD4[CreERT] induced deletion of REL subunits results in a substantial loss of T cells (Carty et al., 2023).

In addition to activating NF-κB dependent transcriptional activity to regulate T cell survival, TNF-induced NF-κB signalling pathways also control cell death through the activity of IKK. In addition to its function as an inhibitor of kappa-B kinase, the IKK complex also directly controls cell survival independent of NF-κB activation. While thymic development is largely normal in the absence of REL subunits (Webb et al., 2019), IKK deficiency in αβ T cells results in a profound block in thymic development that is arrested at the SP stage (Schmidt-Supprian et al., 2003) due to TNF-induced apoptosis (Webb et al., 2016). Ligation of TNFR1 causes recruitment of TRADD, TRAF2, and the serine/threonine kinase RIPK1. The ubiquitin ligases TRAF2, cellular inhibitor of apoptosis proteins (cIAPs) and the linear ubiquitin chain assembly complex (LUBAC), add ubiquitin chain modifications to themselves and RIPK1, creating a scaffold that allows recruitment and activation of the TAB/TAK and IKK complexes that in turn activate NF-κB. This is termed complex I (Annibaldi and Meier, 2018; Vandenabeele et al., 2010). A failure to maintain the stability of this

complex results in the formation of cell death inducing complexes composed of TRADD, FADD, CASPASE8 and RIPK1 that induce apoptosis, a function dependent upon RIPK1 kinase activity (Annibaldi and Meier, 2018; Dondelinger et al., 2016; Ting and Bertrand, 2016). Phosphorylation of RIPK1 by IKK blocks RIPK1 kinase activity and therefore its capacity to induce apoptosis (Dondelinger et al., 2015). In thymocytes, it is this function of IKK, and not NF-κB activation, that is critical for their survival and onward development and accounts for the phenotype observed in IKK deficiency (Blanchett et al., 2022; Webb et al., 2019). Survival of mature T cells depends upon both the IKK capacity to repress RIPK1 and its function to activate NF-κB (Carty et al., 2023). RIPK1-dependent cell death of thymocytes and mature naive T cells is mediated by apoptosis, rather than necroptosis, since cell death is CASPASE8 dependent (Carty et al., 2023). Thymocytes are not susceptible to necroptosis as they lack expression of MLKL (Webb et al., 2019), a key effector molecule for mediating necroptotic cell death. Similarly, resting peripheral αβ T cells are resistant to necroptotic cell death, since *Casp8* deletion has little impact on peripheral αβ T cell compartments (Ch'en et al., 2008; Ch'en et al., 2011). In contrast, activated T cells are acutely susceptible to necroptosis and readily undergo necroptotic cell death in the absence of *Casp8* expression during LCMV infection or following activation in vitro (Ch'en et al., 2008; Ch'en et al., 2011).

γδ T cells are a distinct subset of T lymphocytes that play a unique and versatile role in host immunity. While conventional αβ T cells rely on MHC-restricted antigen presentation, γδ T cells recognise stress-induced ligands, phosphoantigens, and non-peptidic molecules directly (Deseke and Prinz, 2020). Distinct subsets of innate-like cells also develop with specific effector functions, such as type 1 and type 17 γδ T cells (Munoz-Ruiz et al., 2017; Vantourout and Hayday, 2013). This enables them to act as early responders in immune defense. As such, while it seems likely that inflammatory NF-κB and cell death signalling pathways would be important regulators of γδ T cell development and homeostasis, as is the case for αβ T cells, very little is currently known. Evidence that NF-κB signalling could be important comes from the observation that the TNFRSF member CD27 is an important regulator of type 1 γδ T cell development (Ribot et al., 2009) and is also an activator of NF-κB signalling required for survival of αβ T cells (Silva et al., 2014). However, whether and how inflammatory signalling pathways mediated by IKK and NF-κB for the normal homeostasis of γδ T cells remains unknown.

In the present study, we used mouse genetics to investigate the role of IKK and NF-κB survival and cell death pathways for homeostasis of γδ T cells. We found evidence that thymic development of type I γδ T cells was dependent on these pathways, but redundant for adaptive γδ T cells. However, long-term survival of adaptive γδ T cells was dependent upon both NF-κB and, in part, RIPK1-dependent survival pathways and both populations were highly susceptible to necroptotic cell death.

Results

IKK signalling is required for generation of type 1 γδ T cells and maintenance of adaptive γδ T cells

We first asked whether IKK signalling was required for γδ T cell specification and development of adaptive and type 1 γδ T cells in the thymus of adult mice lacking expression of IKK proteins in the T cell compartment. Type 17 γδ T cells develop as a finite wave in a very specific window of embryogenesis (Munoz-Ruiz et al., 2017). Since our analysis was focused on adult genetic mutants we only assessed type 17 γδ T cells in secondary lymphoid organs. We generated mice with conditional genes encoding IKK1 and IKK2 protein (*Chuk^flox* and *Ikbkb^flox* respectively) and an iCre transgenic construct controlled by huCD2 expression elements (IKKΔT^CD2 mice hereon). huCD2iCre is expressed in CLP in bone marrow, and Cre-mediated gene deletion is evident almost all of the earliest thymic progenitors that enter the thymus (Siegemund et al., 2015), so it is ideal to target gene deletion prior to γδ T cell specification that occurs during DN2 (Fiala et al., 2020). In confirmation, we analysed huCD2^iCre mediated Rosa26RmTom *Cre* reporter expression in adaptive, type 1 and type 17 γδ T cell subsets and found ubiquitous expression of reporter in all subsets (Fig. S1). Analysing thymi from IKKΔT^CD2 mice revealed normal representation and numbers of both CD25+ CD27+ TCRδ+ progenitor cells and total CD25– CD27+ TCRδ+ T cells, suggesting that IKK signalling was not required for either specification of the γδ T cell lineage from uncommitted DN precursors or subsequent thymic development of γδ T cells. HSA expression was high in both populations in IKKΔT^CD2 mice, confirming their immature status (Fig. 1A). Type 1 γδ T cells can be identified by their expression of CD122. Analysing subsets of CD25– CD27+ TCRδ+ T cells revealed that CD122– adaptive γδ T cells were present in normal numbers while numbers of CD122+ type 1 γδ T cells were significantly and substantially reduced in the absence of IKK activity (Fig. 1B).

The thymic phenotype of IKKΔT^CD2 mice suggested that IKK signalling is required for the development of type 1 γδ T cells. Analysing numbers of type 1 γδ T cells recovered from lymph node and spleen combined confirmed this view since IKKΔT^CD2 mice were almost completely devoid of this subset (Fig. 1C-D). Strikingly, adaptive γδ T cells were also largely absent from secondary lymphoid organs. Only a small population of CD27– type 17 γδ T cells were detectable in periphery of IKKΔT^CD2 mice, but these were also significantly reduced in number compared to Cre –ve littermates (FIg. 1D). Together, these data suggest that IKK signalling is required for development of type 1 γδ T cells and for the long term survival of adaptive γδ T cells.

Casp8 deficiency rescues thymic generation of type 1 γδ development but not maintenance of peripheral γδ T cells

In αβ T cells, IKK is required to repress acute induction of *Casp8*-dependent cell death in thymocytes and peripheral T cells, independent of NF-κB, but in mature peripheral T cells, does also trigger NF-κB transcriptional activity that also promotes their survival (Carty et al., 2023). To test which functions of IKK are required by different γδ T cell populations, were first analysed IKKΔT^{CD2} mice with additional deletion of *Casp8* (Casp8.IKKΔT^{CD2} mice). In the thymus, numbers of progenitor γδ T cells in Casp8.IKKΔT^{CD2} mice were similar to those of Cre –ve littermates (Fig. 2A), as observed in IKKΔT^{CD2} mice. In contrast, numbers of adaptive γδ T cells in Casp8.IKKΔT^{CD2} mice exhibited a modest but statistically significant reduction, while numbers of type 1 γδ T cells appeared to be rescued to levels similar to Cre –ve controls, suggesting that development of this subset was *Casp8* dependent in the absence of IKK expression. However, this rescue appeared limited in nature, since analysing the periphery of Casp8.IKKΔT^{CD2} mice failed to reveal any detectable rescue of cell numbers of any γδ T cell subsets in the lymph node and spleen (Fig. 2B).

Partial rescue of IKK-deficient peripheral γδ T cells by kinase dead RIPK1

In αβ T cells, IKK kinase activity blocks CASPASE8-dependent apoptosis by directly phosphorylating RIPK1, which acts to repress its kinase activity that is required to induce CASPASE8-dependent cell death (Blanchett et al., 2022; Webb et al., 2019). The failure of CASPASE8 ablation to rescue peripheral γδ T cells potentially implied that IKK was not required to repress extrinsic cell death pathways in γδ T cells. However, while *Casp8* deletion would serve to protect cells from CASPASE8-dependent apoptosis, it could also result in triggering death instead by necroptosis. In T cells, extrinsic death pathways triggering CASPASE8-dependent apoptosis or necroptosis both depend on the kinase activity of RIPK1. Therefore, to test whether *Casp8* deletion was instead triggering necroptosis in Casp8.IKKΔT^{CD2} mice, we analysed peripheral γδ T cell numbers in IKKΔT^{CD2} mice expressing kinase dead RIPK1^{D138N}. As described earlier, IKKΔT^{CD2} mice are almost completely devoid of γδ T cells, except for type 17 cells that were present, albeit in reduced numbers (Fig. 3A-B). Analysing IKKΔT^{CD2} RIPK1^{D138N} mice revealed a small but significant population of CD27$^+$ γδ T cells in the periphery of mice, that included both adaptive and type 1 subsets (Fig. 3A). In contrast, type 17 γδ T cells were not additionally affected in IKKΔT^{CD2} RIPK1^{D138N} mice. These results suggest that IKK is required by γδ T cells to repress cell death pathways. They also provided evidence that γδ T cells are susceptible to necroptosis, and that the failure of CASPASE8 ablation to rescue cells from cell death was due to a switch from apoptotic to necroptotic death in the absence of IKK expression.

Adaptive and type 1 γδ T cells are highly susceptible to necroptosis

In order to directly test whether γδ T cells are susceptible to necroptosis, we analysed mice in which *Casp8* alone was deleted in T cells, in Casp8ΔT[CD2] mice. Analysing γδ T cell development in the thymus revealed normal representation (Fig. 4A) and numbers (Fig. 4B) of progenitor and newly generated adaptive and type 1 γδ T cells. In the periphery, however, numbers of both adaptive and type 1 γδ T cells were profoundly reduced with only a small population of CD27+ γδ T cells remaining (Fig. 4C-D). In contrast, no significant difference in the total number of type 17 γδ T cells was apparent in the periphery, suggesting that CASPASE8 is not required for either development or maintenance of this subset (Fig. 4D).

To confirm that the loss of peripheral γδ T cells was due to necroptosis in the absence of *Casp8* expression, we generated Casp8ΔT[CD2] RIPK1[D138N] mice expressing kinase dead RIPK1, to see if this could rescue cells from death. Analysing the periphery of these mice revealed normal representation (Fig. 5A) and numbers (Fig. 5B) of adaptive, type 1 and type 17 γδ T cells, confirming that loss of adaptive and type 1 subsets was the result of necroptosis in the absence of *Casp8* expression. We further confirmed that rescued cells in Casp8ΔT[CD2] RIPK1[D138N] mice were the functional counterparts of the same subsets in Cre –ve controls by assessing cytokine production in vitro. IL-17A and IFN-gamma production was similar in extent and restricted to the corresponding type 17 and type 1 subsets (Fig. 5C).

Alternative NF-κB activation is redundant for normal γδ T cell homeostasis

Our data demonstrated that generation of type 1 and long-term maintenance of adaptive γδ T cells was highly dependent upon IKK signalling. IKK kinase activity is required by T cells to both trigger NF-κB activation but also to directly block extrinsic cell death pathways by inhibiting RIPK1 activity (Blanchett et al., 2021). Kinase dead RIPK1 only achieved a modest rescue of peripheral γδ T cell numbers, suggesting that these cells also require the IκB kinase activity of IKK for their persistence, and that they also require NF-κB. We therefore dissected the specific role of NF-κB for γδ T cell homeostasis. IKK proteins regulate activation of both canonical NF-κB, mediated by Rela/p50 and cRel/p50 heterodimers, and non-canonical or alternative NF-κB, mediated by RelB/p52 dimers. A trimeric complex of IKK1, IKK2 and regulatory subunit NEMO is responsible for triggering canonical activation, while alternative NF-κB is triggered exclusively by homodimers of IKK1. In IKKΔT[CD2] mice, γδ T cells are unable to activate either of these NF-κB pathways.

To determine if alternative NF-κB pathways contribute to the phenotype of IKKΔT[CD2] mice, we analysed IKK1ΔT[CD2] mice that lack IKK1 expression while retaining IKK2 expression. Canonical NF-κB activation in T cells is largely normal in the absence of IKK1, while alternative NF-κB is completely blocked (Lawrence, 2009). Analysing thymus from these mice showed that development of both adaptive and type 1 γδ T cells was normal (Fig. 6A-B). Similarly, numbers and representation of adaptive, type 1 and type 17 γδ T cells in the periphery were also normal in these

mice as compared with Cre –ve littermates. Therefore, this suggests that alternative NF-κB pathways are redundant for both the development and persistence of γδ T cells.

Canonical NF-κB is essential for maintenance of peripheral γδ T cells

The phenotype of IKK1ΔT^{CD2} mice appeared to exclude a role for alternative NF-κB pathways in regulating γδ T cell homeostasis. Therefore, we next sought to dissect the role of canonical NF-κB pathways. The p50 subunit that derives from processing of NFKB1, lacks a transactivation domain, so while it can bind DNA, it cannot activate transcription alone. Therefore, combined ablation of cREL and RELA is sufficient to completely block canonical NF-κB pathways. Amongst conventional αβ T cells, cREL expression is redundant for naive T cells that can be supported by RELA alone, while memory phenotype T cells depend upon both cREL and RELA expression for their generation and/or persistence (Webb et al., 2019; Zheng et al., 2003a). Therefore, we analysed mice whose T cells lacked expression of only RELA (RelaΔT^{CD2}), RELA and Nfkb1 (RelaΔT^{CD2} Nfkb1$^{-/-}$) or RELA and cREL (Rela.RelΔT^{CD2}). We first analysed the thymus of these strains to assess the development of adaptive and type 1 γδ T cells. Representation of progenitor and adaptive γδ T cells appeared normal (Fig. 7A), while there was some evidence of a modest reduction in absolute numbers of both these subsets in some of the REL deficient strains (Fig. 7B). This was contrary to the lack of phenotype observed in IKKΔT^{CD2} mice (Fig. 1). Since representation of these populations appeared normal, we also compared representation of progenitors γδ T cells with DN3 cell numbers, that reflect upstream progenitor pool. This ratio was similar across strains (Fig. 7B), suggesting that the observed modest reductions were not cell intrinsic but may rather reflect differences in overall thymic cellularity from these strains. In contrast, analysing representation and numbers of type 1 γδ T cells revealed similar reductions as observed in IKK1ΔT^{CD2} mice, suggesting that development of this subset is indeed dependent upon canonical NF-κB. Development of this subset appeared highly dependent upon NF-κB, since even ablation of RELA alone was sufficient to reduce the numbers of newly generated cells in the thymus (Fig. 7B).

Finally, we analysed the peripheral compartments of REL deficient strains to determine the specific requirements for NF-κB by different subsets. In mice lacking RELA and/or cREL, type 1 and type 17 γδ T cells were substantially reduced (Fig. 7C). The reduction in numbers of type 1 γδ T cells in the periphery of these mice mirrored reductions observed in the thymus, reinforcing the view that NF-κB signalling was essential for their normal generation. The reduction of type 17 γδ T cells suggests that NF-κB is also necessary for the development and/or maintenance of this population. Type 1 and type 17 γδ T cells were similarly reduced in all three REL deficient strains, suggesting a strong dependence upon NF-κB. Peripheral numbers of adaptive γδ T cells were also profoundly reduced in RELA/cREL deficient mice, but there was evidence for redundancy between subunits, since specific ablation of RELA alone was well tolerated (FIg. 7C). Overall, these data

show that maintenance of replete peripheral γδ T cell compartments is highly dependent on tonic NF-κB signals either for development or maintenance of different subsets.

Discussion

The IKK complex is a critical regulator of cell death, differentiation and inflammation in multiple cell types, including αβ T cells. In the present study, we sought to understand whether IKK signalling was also important for the development and maintenance of γδ T cells, which are implicated as early responders in a host of different inflammatory settings. We found that IKK was critical for establishing a mature γδ T cell compartment, but found evidence for contrasting roles and mechanisms of downstream pathways for the development and maintenance of different subsets.

The inhibitor of κB kinase activity of the IKK complex is critical for activating NF-κB signalling, so we could demonstrate the essential role of this transcription factor by two independent means - either ablating the upstream IKK complex or direct ablation of canonical REL subunits. Numbers of type 1 γδ T cells in the thymus were substantially reduced in both settings, suggesting that generation of this pre-differentiated subset requires NF-κB signalling to support their ontogeny. In contrast, NF-κB signalling was redundant for both specification of progenitors to a γδ T cell lineage and generation of adaptive γδ T cells, as numbers of progenitors and adaptive γδ T cells were largely normal in both REL and IKK deficient strains. Strikingly, however, mature adaptive γδ T cells were largely absent in both REL and IKK deficient mice, revealing that NF-κB is critical for maintenance of mature γδ T cells. Assessing the role of different REL subunits also revealed distinct requirements for development vs maintenance that mirrored preferences observed in αβ T cells. Generation of effector αβ T cells is highly dependent upon NF-κB, since loss of cREL alone is sufficient to block development of αβ T cell memory (Webb et al., 2019; Zheng et al., 2003a). Similarly, we found that loss of RELA alone was sufficient to impair development of type 1 γδ T cells, suggesting the cREL and p50 alone are insufficient for their development. Furthermore, in naive αβ T cell, there is evidence of redundancy between REL subunits. Expression of cREL and p50 in the absence of RELA is sufficient to maintain naive αβ T cells, while expression of cREL alone, in RELA/p50 double knockouts is not (Webb et al., 2019). We also found this to be the exact case for adaptive γδ T cells. These subtly distinct requirements for NF-κB probably reflect both distinct biological processes involved, ontogeny vs maintenance, but also potentially the receptors involved in triggering signalling. TNFRSF members appear to be implicated in triggering survival signalling in naive αβ T cells (Silva et al., 2014) while generation of differentiated αβ T effectors is dependent on strong agonist triggers from TCR and also TNFRSF members (Layzell et al., 2024). This is also true of Foxp3+ regulatory T cells, that require both RELA and cREL for their development (Isomura et al., 2009; Messina et al., 2016), and are induced by a combination of

both agonist TCR and TNFRSF signalling, amongst other receptors. Intrathymic development of type 1 γδ T cells also depend on agonist TCR signalling.

In αβ T cells, IKK also plays a crucial role in controlling cell survival directly by its repressive activity upon the cell death regulator, RIPK1. Developing thymocytes do not require NF-κB signalling for development but do become exquisitely sensitive to TNF-induced cell death in the absence of IKK (Webb et al., 2019), while mature naive T cells require IKK expression to both repress RIPK1 and activate an NF-κB dependent survival signal. The balance between control of NF-κB and blocking cell death by IKK appears different in γδ T cells, as does the control of cell death processes themselves. Neither αβ nor γδ thymocytes undergo necroptosis in the absence of CASPASE8, while, in stark contrast to resting αβ T cells, adaptive and type 1 γδ T cells in the periphery are exquisitely sensitive to the induction of necroptosis in the absence of CASPASE8. This appears to be determined in part by expression of MLKL that is present in γδ T cells but absent from resting αβ T cells (ImmGen browser enquiry (Heng and Painter, 2008)). Using kinase dead RIPK1, it was possible to test the importance of regulation of RIPK1-dependent cell death pathways by IKK, since this RIPK1 mutant blocks both necroptosis and apoptosis, while CASPASE8 ablation simply redirected cell death processes. While there was some detectable restoration of the peripheral γδ T cell compartment in IKKΔT[CD2] mice expressing kinase dead RIPK1, it was far less than achieved amongst naive αβ T cells following CASPASE8 ablation. Together with the phenotype of REL-ablated mice, this suggests that the loss of NF-κB activity in IKK deficiency has the biggest impact on the maintenance of peripheral γδ T cells.

Type 17 γδ T cells are primarily generated during embryonic stages of development (Munoz-Ruiz et al., 2017), and so we could not directly observe their intrathymic generation in adult mice. As such, the presence or absence of cells in adult mice could reflect either developmental processes and/or their defective survival. Nevertheless, there were still some useful conclusions we could draw. The maintenance of the mature type 17 γδ T cells does depend on NF-κB signalling. Peripheral numbers of type 17 γδ T cells were reduced by around a two-thirds in both IKK and REL deficient strains. However, in contrast to other subsets, there was no evidence that kinase dead RIPK1 mediated any rescue of type 17 subset, and they did not appear to be sensitive to necroptosis in the absence of CASPASE8. Their numbers were also not restored in Casp8.IKKΔT[CD2] mice, strongly suggesting that the sole defect in IKK deficient strains was the absence of NF-κB activation. The stark contrast in sensitivity of type 17 and other subsets to necroptosis in the absence of CASPASE8 was surprising, as they all appear to express MLKL (ImmGen browser enquiry (Heng and Painter, 2008)). It is possible this reflects the distinct fetal liver derived progenitors from which type 17 cells are derived. Whichever the case, if confirmed, it would be of interest to identify the mechanism by which these cells are resistant to necroptosis, since they appear to express the basic elements required to form the necrosome.

In conclusion, our study reveals the complexity of regulation of inflammatory and cell death pathways in T cells of the adaptive immune system. In other cell types and tissues, IKK mediates well defined pro-inflammatory and pro-suvival functions. In contrast, the functions of IKK and regulation of cell death pathways varies with differentiation state and lineage. Repression of RIPK1 by IKK is critical for thymic development of αβ T cells, while this function is largely redundant in γδ T cells, that instead require the NF-κB activating functions of IKK for either development or survival. Resting αβ T cells are resistant to necroptosis induction in the absence of CASPASE8 while γδ T cells become exquisitely sensitive as soon as they enter the periphery. This difference may reflect the role of γδ T cells as first line early responders in immune responses, that may expose them to microbial interference of cell death pathways, that necroptosis has evolved to resist. Whichever is the case, a detailed understanding of these pathways will be critical to understand the full impact of therapies that target these potent inflammatory pathways, and that will have diverse and complex impact on adaptive immunity.

Materials and methods

Mice

Mice with the following mutations were used in this study; B6.129-Casp8[tm1Hed]/J (*Casp8[fx]*), Ikbkbtm2Mka (Li et al., 2003) *(Ikbkb[fx])*, Chukrm1Mpa (Gareus et al., 2007) *(Chuk[fx])*, Relatm1Asba (Steinbrecher et al., 2008) *(Rela[fx])*, B6.129S1-Reltm1Ukl/J(Heise et al., 2014) *(Rel[fx])*, Nfkb1tm1Bal *(Nfkb1[null])*, *Cre* transgenes expressed under the control of the human CD2, B6.Cg-Tg(CD2-icre)4Kio/J (*huCD2[iCre]*) (de Boer et al., 2003) mice with a D138N mutation in *Ripk1, B6.129-Ripk1[tm1Geno]/J* (RIPK1[D138N]*)* (Newton et al., 2014). The following strains were bred using these alleles for this study; *Chuk[fx/fx] Ikbkb[fx/fx] huCD2[iCre]* (IKKΔT[CD2]), *Chuk[fx/fx] huCD2[iCre]* (IKK1ΔT[CD2]), *Chuk[fx/fx] Ikbkb[fx/fx] Casp8[fx/fx] huCD2[iCre]* (Casp8.IKKΔT[CD2]), *Chuk[fx/fx] Ikbkb[fx/fx] huCD2[iCre] Ripk1[D138N]* (IKKΔT[CD2] RIPK1[D138N]), *Casp8[fx/fx] huCD2[iCre]* (Casp8ΔT[CD2]), *Casp8[fx/fx] huCD2[iCre] Ripk1[D138N]* (Casp8ΔT[CD2]RIPK1[D138N]), *Rela[fx] huCD2[iCre]* (RelaΔT[CD2]), *Rela[fx/fx]* Rel[fx/fx]*huCD2[iCre]* (Rela.RelΔT[CD2]), *Rela[fx/fx] huCD2[iCre] Nfkb1[−/−]*(RelaΔT[CD2] Nfkb1[−/−]). All mice were bred in the Comparative Biology Unit of the Royal Free UCL campus and at Charles River laboratories, Manston, UK. Animal experiments were performed according to institutional guidelines and Home Office regulations.

Flow cytometry and electronic gating strategies

Flow cytometric analysis was performed with 10×10^6 thymocytes, 5×10^6 lymph node (LN) or spleen cells. Cell concentrations of thymocytes, lymph node and spleen cells were determined with a Scharf Instruments Casy Counter. Cells were incubated with saturating concentrations of antibodies in 100 µl of Dulbecco's phosphate-buffered saline (PBS) containing bovine serum albumin (BSA, 0.1%) for 1 hour at 4°C followed by two washes in PBS-BSA. Panels used the following mAb: BV421-conjugated antibody against CD27 (Biolegend), PE-conjugated antibody against CD127 (ThermoFisher Scientific), BV785-conjugated CD44 antibody (Biolegend), BUV395-conjugated antibody against CD25 (BD horizon), PE-Cy7-conjugated antibody against CD122 (Biolegend), PerCP-cy5.5-conjugated antibody against CD3 (eBioscience), APC-conjugated antibody against TCRɣδ (eBioscience), FITC-conjugated antibody against IFN-ɣ (BD Biosciences), PE-Cy7-conjugated antibody against IL-17A (eBioscience). Cell viability was determined using LIVE/DEAD cell stain kit (Invitrogen Molecular Probes), following the manufacturer's protocol. multi-color flow cytometric staining was analyzed on a LSRFortessa (Becton Dickinson) instrument, and data analysis and color compensations were performed with FlowJo V10 software (TreeStar). The following gating strategies were used : ɣδ progenitor T cells - CD3[+] TCRɣδ[hi] CD25[+] CD27[+] , adaptive ɣδ T cells - CD3[+]TCRɣδ[+]CD25[-]CD27[+]CD44[lo]CD122[lo], type 1 ɣδ T cells - CD3[+]TCRɣδ[+]CD25[-]CD27[+]CD44[+]CD122[+], type 17 ɣδ T cells - CD3[+]TCRɣδ[+]CD25[-]CD27[-] respectively.

Intracellular cytokine staining

LN T cells (cervical, auxiliary, brachial, inguinal and mesenteric) were cultured at 37°C with 5% CO_2 in RPMI-1640 (Gibco, Invitrogen Corporation, CA) supplemented with 10% (v/v) fetal bovine serum (FBS) (Gibco Invitrogen), 0.1% (v/v) 2-mercaptoethanol βME (Sigma Aldrich) and 1% (v/v) penicillin-streptomycin (Gibco Invitrogen) (RPMI-10). Approximately 5×10^6 LN cells were stimulated with a protein transport inhibitor cocktail (Brefeldin and Monensin) at 12.6 μM concentration and cell stimulation cocktail (PMA and Ionomycin) at 1.42 μM concentration resuspended in RPMI. The stimulated cells were incubated for 4 hours, followed by the cell surface and intracellular cytokine staining.

Statistics

Statistical analysis and bar charts were performed using Graphpad Prism 9.2. Column data compared by two-tailed T-test (non-parametric) Mann-Witney t-test. Results were denoted as ns = nonsignificant, * $p<0.05$, ** $p<0.01$, *** $p<0.001$ and **** $p<0.0001$.

Acknowledgements

We thank UCL Comparative Biology Unit staff for assistance with mouse breeding and maintenance. We thank the following for generously sharing of their mouse strains: Prof Manolis Pasparakis for *Chuk* conditional strain, Prof Michael Karin for *Ikbkb* conditional strain, Prof Vishva Dixit for the RIPK1[D138N] strain, Prof Albert Baldwin for *Rela* conditional strain. The authors declare no competing financial interests. The work in the Seddon lab is supported by the Medical Research Council UK under programme codes MR/P011225/1. FIS was supported by a scholarship from the Commonwealth Scholarship Commission of the United Kingdom.

Bibliography

Annibaldi, A., and P. Meier. 2018. Checkpoints in TNF-Induced Cell Death: Implications in Inflammation and Cancer. *Trends Mol Med* 24:49-65.

Blanchett, S., I. Boal-Carvalho, S. Layzell, and B. Seddon. 2021. NF-kappaB and Extrinsic Cell Death Pathways - Entwined Do-or-Die Decisions for T cells. *Trends Immunol* 42:76-88.

Blanchett, S., Y. Dondelinger, A. Barbarulo, M.J.M. Bertrand, and B. Seddon. 2022. Phosphorylation of RIPK1 serine 25 mediates IKK dependent control of extrinsic cell death in T cells. *Front Immunol* 13:1067164.

Bonizzi, G., and M. Karin. 2004. The two NF-kappaB activation pathways and their role in innate and adaptive immunity. *Trends Immunol* 25:280-288.

Carty, F., S. Layzell, A. Barbarulo, F. Islam, L.V. Webb, and B. Seddon. 2023. IKK promotes naive T cell survival by repressing RIPK1-dependent apoptosis and activating NF-kappaB. *Sci Signal* 16:eabo4094.

Ch'en, I.L., D.R. Beisner, A. Degterev, C. Lynch, J. Yuan, A. Hoffmann, and S.M. Hedrick. 2008. Antigen-mediated T cell expansion regulated by parallel pathways of death. *Proc Natl Acad Sci U S A* 105:17463-17468.

Ch'en, I.L., J.S. Tsau, J.D. Molkentin, M. Komatsu, and S.M. Hedrick. 2011. Mechanisms of necroptosis in T cells. *J Exp Med* 208:633-641.

de Boer, J., A. Williams, G. Skavdis, N. Harker, M. Coles, M. Tolaini, T. Norton, K. Williams, K. Roderick, A.J. Potocnik, and D. Kioussis. 2003. Transgenic mice with hematopoietic and lymphoid specific expression of Cre. *Eur J Immunol* 33:314-325.

Deseke, M., and I. Prinz. 2020. Ligand recognition by the gammadelta TCR and discrimination between homeostasis and stress conditions. *Cell Mol Immunol* 17:914-924.

Dondelinger, Y., S. Jouan-Lanhouet, T. Divert, E. Theatre, J. Bertin, P.J. Gough, P. Giansanti, A.J. Heck, E. Dejardin, P. Vandenabeele, and M.J. Bertrand. 2015. NF-kappaB-Independent Role of IKKalpha/IKKbeta in Preventing RIPK1 Kinase-Dependent Apoptotic and Necroptotic Cell Death during TNF Signaling. *Mol Cell* 60:63-76.

Dondelinger, Y., P. Vandenabeele, and M.J. Bertrand. 2016. Regulation of RIPK1's cell death function by phosphorylation. *Cell Cycle* 15:5-6.

Fiala, G.J., A.Q. Gomes, and B. Silva-Santos. 2020. From thymus to periphery: Molecular basis of effector gammadelta-T cell differentiation. *Immunol Rev* 298:47-60.

Gareus, R., M. Huth, B. Breiden, A. Nenci, N. Rosch, I. Haase, W. Bloch, K. Sandhoff, and M. Pasparakis. 2007. Normal epidermal differentiation but impaired skin-barrier formation upon keratinocyte-restricted IKK1 ablation. *Nat Cell Biol* 9:461-469.

Gerondakis, S., and U. Siebenlist. 2010. Roles of the NF-kappaB pathway in lymphocyte development and function. *Cold Spring Harb Perspect Biol* 2:a000182.

Heise, N., N.S. De Silva, K. Silva, A. Carette, G. Simonetti, M. Pasparakis, and U. Klein. 2014. Germinal center B cell maintenance and differentiation are controlled by distinct NF-kappaB transcription factor subunits. *J Exp Med* 211:2103-2118.

Heng, T.S., and M.W. Painter. 2008. The Immunological Genome Project: networks of gene expression in immune cells. *Nature immunology* 9:1091-1094.

Isomura, I., S. Palmer, R.J. Grumont, K. Bunting, G. Hoyne, N. Wilkinson, A. Banerjee, A. Proietto, R. Gugasyan, L. Wu, A. McNally, R.J. Steptoe, R. Thomas, M.F. Shannon, and S. Gerondakis. 2009. c-Rel is required for the development of thymic Foxp3+ CD4 regulatory T cells. *J Exp Med* 206:3001-3014.

Lawrence, T. 2009. The nuclear factor NF-kappaB pathway in inflammation. *Cold Spring Harb Perspect Biol* 1:a001651.

Layzell, S., A. Barbarulo, G. van Loo, R. Beyaert, and B. Seddon. 2024. NF-kappaB regulated expression of A20 controls IKK dependent repression of RIPK1 induced cell death in activated T cells. *Cell Death Differ*

Li, Z.W., S.A. Omori, T. Labuda, M. Karin, and R.C. Rickert. 2003. IKK beta is required for peripheral B cell survival and proliferation. *J Immunol* 170:4630-4637.

Messina, N., T. Fulford, L. O'Reilly, W.X. Loh, J.M. Motyer, D. Ellis, C. McLean, H. Naeem, A. Lin, R. Gugasyan, R.M. Slattery, R.J. Grumont, and S. Gerondakis. 2016. The NF-kappaB transcription factor RelA is required for the tolerogenic function of Foxp3(+) regulatory T cells. *J Autoimmun* 70:52-62.

Miller, M.L., M. Mashayekhi, L. Chen, P. Zhou, X. Liu, M. Michelotti, N. Tramontini Gunn, S. Powers, X. Zhu, C. Evaristo, M.L. Alegre, and L.L. Molinero. 2014. Basal NF-kappaB controls IL-7 responsiveness of quiescent naive T cells. *Proc Natl Acad Sci U S A* 111:7397-7402.

Munoz-Ruiz, M., N. Sumaria, D.J. Pennington, and B. Silva-Santos. 2017. Thymic Determinants of gammadelta T Cell Differentiation. *Trends Immunol* 38:336-344.

Newton, K., D.L. Dugger, K.E. Wickliffe, N. Kapoor, M.C. de Almagro, D. Vucic, L. Komuves, R.E. Ferrando, D.M. French, J. Webster, M. Roose-Girma, S. Warming, and V.M. Dixit. 2014. Activity of protein kinase RIPK3 determines whether cells die by necroptosis or apoptosis. *Science* 343:1357-1360.

Ribot, J.C., A. deBarros, D.J. Pang, J.F. Neves, V. Peperzak, S.J. Roberts, M. Girardi, J. Borst, A.C. Hayday, D.J. Pennington, and B. Silva-Santos. 2009. CD27 is a thymic determinant of the balance between interferon-gamma- and interleukin 17-producing gammadelta T cell subsets. *Nat Immunol* 10:427-436.

Schmidt-Supprian, M., G. Courtois, J. Tian, A.J. Coyle, A. Israel, K. Rajewsky, and M. Pasparakis. 2003. Mature T cells depend on signaling through the IKK complex. *Immunity* 19:377-389.

Schmidt-Supprian, M., J. Tian, E.P. Grant, M. Pasparakis, R. Maehr, H. Ovaa, H.L. Ploegh, A.J. Coyle, and K. Rajewsky. 2004. Differential dependence of CD4+CD25+ regulatory and natural killer-like T cells on signals leading to NF-kappaB activation. *Proc Natl Acad Sci U S A* 101:4566-4571.

Siegemund, S., J. Shepherd, C. Xiao, and K. Sauer. 2015. hCD2-iCre and Vav-iCre mediated gene recombination patterns in murine hematopoietic cells. *PLoS One* 10:e0124661.

Silva, A., G. Cornish, S.C. Ley, and B. Seddon. 2014. NF-kappaB signaling mediates homeostatic maturation of new T cells. *Proc Natl Acad Sci U S A* 111:E846-855.

Steinbrecher, K.A., E. Harmel-Laws, R. Sitcheran, and A.S. Baldwin. 2008. Loss of epithelial RelA results in deregulated intestinal proliferative/apoptotic homeostasis and susceptibility to inflammation. *J Immunol* 180:2588-2599.

Ting, A.T., and M.J.M. Bertrand. 2016. More to Life than NF-kappaB in TNFR1 Signaling. *Trends Immunol* 37:535-545.

Vandenabeele, P., W. Declercq, F. Van Herreweghe, and T. Vanden Berghe. 2010. The role of the kinases RIP1 and RIP3 in TNF-induced necrosis. *Sci Signal* 3:re4.

Vantourout, P., and A. Hayday. 2013. Six-of-the-best: unique contributions of gammadelta T cells to immunology. *Nat Rev Immunol* 13:88-100.

Webb, L.V., A. Barbarulo, J. Huysentruyt, T. Vanden Berghe, N. Takahashi, S. Ley, P. Vandenabeele, and B. Seddon. 2019. Survival of Single Positive Thymocytes Depends upon Developmental Control of RIPK1 Kinase Signaling by the IKK Complex Independent of NF-κB. *Immunity* 50:348-361.e344.

Webb, L.V., S.C. Ley, and B. Seddon. 2016. TNF activation of NF-kappaB is essential for development of single-positive thymocytes. *J Exp Med* 213:1399-1407.

Xing, Y., X. Wang, S.C. Jameson, and K.A. Hogquist. 2016. Late stages of T cell maturation in the thymus involve NF-κB and tonic type I interferon signaling. *Nature Immunology* 17:565-573.

Zheng, Y., M. Vig, J. Lyons, L. Van Parijs, and A.A. Beg. 2003a. Combined deficiency of p50 and cRel in CD4+ T cells reveals an essential requirement for nuclear factor kappaB in regulating mature T cell survival and in vivo function. *The Journal of experimental medicine* 197:861-874.

Zheng, Y., M. Vig, J. Lyons, L. Van Parijs, and A.A. Beg. 2003b. Combined deficiency of p50 and cRel in CD4+ T cells reveals an essential requirement for nuclear factor kappaB in regulating mature T cell survival and in vivo function. *The Journal of Experimental Medicine* 197:861-874.

Figure 1 - Development of type 1 and persistence of adaptive and type 17 γδ T cells depends on IKK expression. Thymi, lymph nodes and spleen from IKKΔT^{CD2} mice (n=13) and Cre -ve littermates (n=7) were enumerated and analysed by flow. (A) Representative flow plots are of thymocytes with the indicated gates, showing gates used to identify progenitor, adaptive and type 1 subsets intrathymically. (B) Bar charts are of total cell numbers of the indicated subset from the thymus. (C) Representative flow plots illustrate the gating strategy to identify adaptive, type 1 and type 17 subsets in lymph nodes (shown) and spleen. (D) Bar charts are of total cell numbers recovered from both spleen and lymph nodes of the indicated subset. Data are pooled from multiple batches of mice analysed. Error bars show standard deviations.

A

Thymus

B   Periphery

Figure 2 - CASPASE8 ablation does not rescue peripheral γδ T cell compartment of IKKΔT^{CD2} mice.
Thymi, lymph nodes and spleen from Casp8.IKKΔT^{CD2} mice (n=14) and Cre –ve littermates (n=9) were enumerated and analysed by flow. (A) Bar charts are of total cell numbers of the indicated subset recovered from thymi of the indicated strain. (B) Representative flow plots from lymph nodes of the indicated mouse strains (rows), with the indicated gates (columns). Bar charts are of total cell numbers recovered from both spleen and lymph nodes of the indicated subset. Data are pooled from multiple batches of mice analysed. Error bars show standard deviations.

Figure 3 - Kinase dead RIPK1 mediates partial rescue of peripheral γδ T cell compartments of IKKΔT^CD2 mice. Lymph nodes and spleen from IKKΔT^CD2 (n=4) IKKΔT^CD2RIPK1^D138N mice (n=4) and Cre –ve littermates (n=4 each) were enumerated and analysed by flow. (A) Representative flow plots are of lymph node cells from different strains (rows) with the indicated gates (columns). (B) Bar charts are of total cell numbers of the indicated subset recovered from lymph nodes and spleen combined of the indicated IKKΔT^CD2 strain. Data are pooled from two independent experiments. Error bars show standard deviations.

Figure 4 - CASPASE8 expression is critical for long-term survival of peripheral γδ T cells. Thymi, lymph nodes and spleen from Casp8ΔT^CD2 mice (n=10) and Cre –ve littermates (n=13) were enumerated and analysed by flow. (A) Representative flow plots are of thymocytes with the indicated gates, showing gates used to identify progenitor, adaptive and type 1 subsets. (B) Bar charts are of total cell numbers of the indicated subset from thymus. (C) Representative flow plots illustrate gating strategy to identify adaptive, type 1 and type 17 subsets in lymph nodes (shown) and spleen. (D) Bar charts are of total cell numbers recovered from both spleen and lymph nodes combined, of the indicated subset. Data are pooled from five independent experiments. Error bars show standard deviations.

Figure 5 - Kinase dead RIPK1 fully restores the peripheral γδ T cell compartments of Casp8ΔTCD2 mice. Lymph nodes and spleen from Casp8ΔTCD2 RIPK1D138N mice (n=16) and Cre –ve littermates (n=7) were enumerated and analysed by flow. (A) Representative flow plots are of lymph nodes from the indicated strains (rows) with the indicated gates (columns), showing gates used to identify type 17, adaptive and type 1 subsets. (B) Bar charts are of total cell numbers of the indicated subsets from both lymph nodes and spleen combined. (C) Lymph node cells were stimulated in vitro with calcium ionophore and phorbyl esters for 4 hours with brefeldin A, and then analysed for the indicated intracellular cytokines. Representative flow plots illustrate gating strategy to identify adaptive, type 1 and type 17 subsets on the basis of CD44 and CD122 expression. Quad gates for cytokine detection were set against negative controls of matched unstimulated cells. Bar charts are of total % of cells stained for IFN-gamma or IL-17A. Data are pooled from multiple batches of mice analysed (A-B) or are the pooled from 4 independent experiments (C). Error bars show standard deviations.

Figure 6 - Alternative NF-κB signalling is redundant for generation and maintenance of γδ T cell compartments. Thymi, lymph nodes and spleen from IKKΔT^CD2 mice (n=13) and Cre –ve littermates (n=7) were enumerated and analysed by flow. (A) Representative flow plots are of thymocytes with the indicated gates, showing gates used to identify progenitor, adaptive and type 1 subsets intrathymically. (B) Bar charts are of total cell numbers of the indicated subset from thymus. (C) Representative flow plots illustrate gating strategy to identify adaptive, type 1 and type 17 subsets in lymph nodes (shown) and spleen. (D) Bar charts are of total cell numbers recovered from both spleen and lymph nodes of the indicated subset. Data are pooled from multiple batches of mice analysed. Error bars show standard deviations.

Figure 7 - Canonical NF-κB signalling is essential for development of type 1 and maintenance of adaptive γδ T cell compartments. Thymi, lymph nodes and spleen from RelaΔT^{CD2} (n=9), RelaΔT^{CD2} *Nfkb1$^{-/-}$* mice (n=5), Rela.RelΔT^{CD2} (n=17) and Cre –ve littermates (n=24) were enumerated and analysed by flow. (A) Representative flow plots are of thymocytes with the indicated gates, showing gates used to identify progenitor, adaptive and type 1 subsets intrathymically. (B) Bar charts are of total cell numbers of the indicated subset from thymus, and the ratio of progenitor:DN3 subsets for the indicated strains. + indicates WT allele, – indicates gene deletion. (C) Bar charts are of total cell numbers recovered from both spleen and lymph nodes of the indicated subset. Data are pooled from multiple batches of mice analysed. Error bars show standard deviations.

Chapter 4

Figure S1 - huCD2iCre targets Cre activity to γδ T cells in the thymus

Thymi and lymph nodes (LNs) from 9-15 weeks old: wild-type mouse was analysed by flow. The density plot number represents the percentage of cells in each gate. (A) Representative flow plots are of thymocytes profiles: Histograms display RFP expression by γδ progenitor, effector γδ and type 17 γδ T cells. (B) Representative flow plots display LNs profiles: Histograms display RFP expression by effector γδ and type 17 γδ T cells.