Micron-scale, liquid-liquid phase separation in ternary lipid membranes containing DPPE

Gunnar J. Goetz[1], Sasha Naomi[1], Angelique M. Madrigal[1], Catherine L. A. Chang[1†], Caitlin E. Cornell[1††], and Sarah L. Keller[1]

[1] Department of Chemistry, University of Washington, Seattle, WA, 98195

Keywords: lipid, vesicle, liposome, phase separation, three-phase coexistence
Running title: Coexisting liquid phases with PE-lipids

*   Corresponding author: slkeller@uw.edu
†   Current address:
        Department of Psychiatry, University of Nevada, Reno, NV, 89557.
†† Current address:
        Department of Bioengineering, University of California, Berkeley, CA, 94720

ORCID:
    G.J. Goetz   0000-0002-5316-7950
    S. Naomi   0000-0003-3941-6946
    A.M. Madrigal   0009-0006-3632-7314
    C.L.A. Chang   0000-0003-4103-3303
    C.E. Cornell   0000-0001-7675-1415
    S.L. Keller   0000-0001-5919-0185

ABSTRACT

Micron-scale, liquid-liquid phase separation occurs in membranes of living cells, with physiological consequences. To discover which lipids might support phase separation in cell membranes and how lipids might partition between phases, miscibility phase diagrams have been mapped for model membranes. Typically, model membranes are composed of ternary mixtures of a lipid with a high melting temperature, a lipid with a low melting temperature, and cholesterol. Phospholipids in ternary mixtures are chosen primarily to favor stable membranes (phosphatidylcholines and sphingomyelins) or add charge (phosphatidylglycerols and phosphatidylserines). A major class of phospholipids missing from experimental ternary diagrams has been the phosphatidylethanolamines (PEs). PE-lipids constitute up to 20 mol% of common biological membranes, where they influence protein function and facilitate membrane fusion. These biological effects are often attributed to PE's smaller headgroup, which leads to higher monolayer spontaneous curvatures and higher melting temperatures. Taken alone, the higher melting points of saturated PE-lipids imply that liquid-liquid phase separation should persist to higher temperatures in membranes containing PE-lipids. Here, we tested that hypothesis by substituting a saturated PE-lipid (DPPE) for its corresponding PC-lipid (DPPC) in two well-studied ternary membranes (DOPC/DPPC/cholesterol and DiphyPC/DPPC/cholesterol). We used fluorescence microscopy to map full ternary phase diagrams for giant vesicles over a range of temperatures. Surprisingly, we found no micron-scale, liquid-liquid phase separation in vesicles of the first mixture (DOPC/DPPE/cholesterol), and only a small region of liquid-liquid phase separation in the second mixture (DiphyPC/DPPE/cholesterol). Instead, coexisting solid and liquid phases were widespread, with the solid phase enriched in DPPE. An unusual feature of these ternary membranes is that solid and liquid-ordered phases can be distinguished by fluorescence microscopy, so tie-line directions can be estimated throughout the phase diagram, and transition temperatures to the 3-phase region (containing a liquid-disordered phase, a liquid-ordered phase, and a solid phase) can be accurately measured.

SIGNIFICANCE STATEMENT

Under physiological conditions, yeast vacuole membranes phase separate into liquid phases. The resulting domains are microns in size, and they are important for the cell's function. Yeast membranes contain a significant fraction of lipids with phosphatidylethanolamine headgroups. It was unknown whether these lipids enhanced liquid-liquid phase separation or hindered it. Here, we produced model membranes containing a saturated phosphatidylethanolamine lipid, we mapped miscibility phase diagrams over broad temperature ranges, and we compared our results to existing diagrams for membranes of other lipid types. We were surprised to find that the new lipid suppressed liquid-liquid phase separation compared to a lipid with a larger, phosphatidylcholine headgroup. Previous simulations incorporated phosphatidylethanolamine lipids, and our results provide an experimental basis for future simulations.

INTRODUCTION

Liquid-liquid phase separation of membranes is key to the function of some cells, which adjust their membrane compositions with respect to the phase transition (1, 2). Independent of whether membranes are derived from a cell with hundreds of lipids (2–5), or from minimal model systems with only two lipids (6), all membranes that demix into micron-scale liquid phases contain at least three moieties: lipid chains with high order, lipid chains with low order, and sterols (7–9).

Figure 1. Structures of lipids. Phospholipids are ordered from left to right in terms of increasing melting temperature (T_{melt}). Vesicles were made from ternary mixtures of lipids with a high T_{melt}, lipids with a low T_{melt}, and a sterol. Lipids have noncylindrical shapes when they have highly branched or unsaturated chains (as for DiPhyPC at the left) and/or smaller headgroups (as for PE lipids on the right). Headgroups of PC- and PE-lipids differ only by the number of methyl groups on the amine.

To discover which types of lipids support liquid-liquid phase separation and how the lipids partition between the phases, researchers have mapped phase diagrams, typically of ternary mixtures of a lipid with saturated chains, a lipid with either unsaturated or methylated chains, and cholesterol (Fig. 1). When researchers are faced with the question of exactly which lipids to incorporate into ternary membranes, they often choose PC-lipids, which are widely commercially available and produce stable membranes over broad experimental temperatures (10). Although researchers have made forays into producing ternary membranes containing other types of lipids (e.g., sphingomyelins (11, 12), ceramides (13, 14), and charged PS- or PG-lipids (15, 16)), mapping phase diagrams can be expensive, in both materials and time. As a result, there have been major gaps in the literature, for example in the lack of ternary membranes containing phosphatidylethanolamine (PE) lipids. PE-lipids are distinguished by

smaller headgroups than PC-lipids and by the potential to hydrogen bond with neighboring phosphates, both of which contribute to the cone shape of PE-lipids (17, 18).

There are several strong reasons to expand the known set of miscibility phase diagrams to include membranes with PE-lipids:

- First, some channels and proteins function and fold differently in membranes containing PE-lipids (e.g., (19–22), and other results reviewed in (23)). Phase separation of membranes would likely concentrate PE-lipids in domains, locally enhancing the effect.

- Second, PE-lipids have higher chain order than corresponding PC-lipids, and they impart higher chain order in neighboring lipids (24). Higher chain order parameters hold whether they are assessed at equivalent absolute temperatures or at equivalent reduced temperatures (25). Higher chain order results in higher T_{melt} values. For example, for di(16:0)PE, also known as DPPE, T_{melt} = 62.3 ± 5.0°C (26), whereas for di(16:0)PC, also known as DPPC, T_{melt} = 41.3 ± 1.8°C (10).

- Third, PE-lipids are common in biological membranes. They constitute ~10% of red blood cell membranes (3), ~8% of giant plasma membrane vesicles from zebrafish ZF4 cells (2), ~15% of total yeast membranes (27), and 10-20% of yeast vacuole membranes (4, 5).

- Fourth, the spontaneous curvature that cone-shaped lipids impart to each leaflet of a membrane (or to the whole of an asymmetric membrane) has biological implications beyond protein function. Curvature can influence membrane fusion (28) as well as the size and shape of phase-separated domains ((29), and reviewed in (30)). Likewise, high fractions of PE-lipids are associated with transitions from flat, lamellar sheets to curved, inverted hexagonal (H_{II}) and cubic phases (26, 31). The pressure dependence of the lamellar-H_{II} transition (32) explains why deep-sea creatures such as ctenophores undergo "tissue melting" at atmospheric pressure (33).

- Fifth, as scientists incorporate more diverse lipids into their simulations (reviewed in (34)), data for complex membranes containing PE-lipids are needed as ground truth, particularly in the context of curvature stress of asymmetric membranes (35).

Here, we produce giant unilamellar vesicles (GUVs) composed of ternary mixtures of cholesterol, a PE-lipid with a high melting temperature (DPPE, or di(16:0)PE), and a PC-lipid with a low melting temperature (either DOPC/di(18:1)PC or DiphyPC/di(16:0-4Me)PC). We chose DPPE as our PE-lipid because it has been reported to form a lamellar phase at common experimental temperatures (26). Even so, membranes with DPPE likely experience some curvature frustration because DPPE transitions into an inverted hexagonal phase at 120°C (26), and a cubic phase may lie nearby (32, 36). In our experiments, we use fluorescence microscopy to identify coexisting phases in vesicle membranes, and we map their miscibility phase diagrams. We compare our data with existing phase diagrams for membranes without PE-lipids,

and we adjust the methylation of lipid headgroups to understand intermediate chemical structures between PE-lipids and PC-lipids.

MATERIALS AND METHODS

Lipids. 1,2-dipalmitoyl-sn-glycero-3-phosphoethanolamine (DPPE or di(16:0)PE, T_{melt} = 62.3 ± 5.0°C, T_{hex} = 120.6 ± 3.6°C (26)), 1,2-dipalmitoyl-sn-glycero-3-phosphoethanolamine-N-methyl (DPPE-1Me or di(16:0)PE-1Me, T_{melt} = 57.6 ± 1.5°C (26)), 1,2-dipalmitoyl-sn-glycero-3-phosphoethanolamine-N,N-dimethyl (DPPE-2Me or di(16:0)PE-2Me, T_{melt} = 43.2 ± 2.0°C (26)), 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC or di(16:0)PC, T_{melt} = 41.3 ± 1.8°C (10)), 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC or di(16:1)PC, T_{melt} = -18.3 ± 3.6°C (10)), 1,2-diphytanoyl-sn-glycero-3-phosphocholine (DiPhyPC or di(16:0-4Me)PC, T_{melt} = <-120°C (37)), and a fluorescent dye, rhodamine-DPPE, were obtained from Avanti Polar Lipids (Alabaster, AL). Cholesterol was obtained from Sigma Aldrich (St. Louis, MO). All lipids were used without further purification. Cholesterol was mixed from powder in chloroform to a concentration of 10 mg/ml. DPPE, DPPE-1Me, and DPPE-2Me were mixed from powder in 9:1 chloroform/methanol to a concentration of 10 mg/ml. All other unlabeled lipids were supplied from Avanti in chloroform, and their nominal concentrations of 10 mg/ml were not verified. Lipid stocks were sealed with Teflon tape and parafilm and stored at -20°C.

Electroformation. Vesicles were made by electroformation (38). Two halves of a glass slide coated with indium-tin-oxide (ITO) (Delta Technologies, Loveland, CO) were heated on a heat block at 60°C, which is just below the boiling point of chloroform. A comparison of data from ITO-coated slides on a heat block at 75°C appears in Fig. S1. Stocks of phospholipids and cholesterol (with 0.8 mol% rhodamine-DPPE) in chloroform or chloroform/methanol were mixed to achieve a total of 0.25 mg total lipid and heated to 60°C for 15 min on the heat block. The lipid solutions were spread evenly across the slides with the side of a glass Pasteur pipette. As the solvent evaporated from the lipid layer, thin-film interference was visible by eye. The slide-halves were exposed to vacuum for ≥30 min to allow any remaining chloroform to evaporate. The two halves were assembled face-to-face, separated by Teflon spacers of 0.3 mm thickness held in place with vacuum grease. The chamber was filled with 18 MΩ–cm water, sealed with vacuum grease, and attached to metal electrodes using binder clips. An AC voltage of 1.0 V at 10 Hz was applied across the electrodes for 1-2 hours at 75°C while vesicles formed (60°C for DiphyPC/DPPC/cholesterol). This temperature was chosen to be well above the highest T_{melt} of any lipid used, to adequately incorporate all lipids into vesicles and to minimize uncertainty in transition temperatures (39). For vesicles containing DPPE, 60°C is not a sufficiently high electroformation temperature (Fig. S2). After electroformation, vesicles were transferred into an Eppendorf tube and typically imaged immediately (or within two hours). From limited data on how lipid ratios in vesicles differ from ratios in stock solutions, we expect at least three of the lipids in our study (saturated PE-lipid, DOPC, and cholesterol) to faithfully incorporate into electroformed vesicles (40).

Vesicle Imaging. Vesicle solutions were diluted to achieve roughly 100-200 vesicles per field of view (usually a 10:1 dilution) and placed between #1.5 glass coverslips. Coverslip edges were coated with vacuum grease, a second coverslip was placed on top to seal the vesicle solution inside, and the coverslip assembly was placed on a layer of Omegatherm thermal paste (Omega Engineering, Norwalk, CT) coating the top plate of a Peltier temperature controller on a home-built microscope stage. Vesicles were imaged on a Nikon (Melville, NY) Eclipse ME600L upright epifluorescence microscope using a Hamamatsu (Bridgewater, NJ) C13440 camera. Contrast between phases in the membrane arises from preferential partitioning of the lipid dye into the phase with less lipid order. For every sample containing membranes with coexisting phases, a representative image was selected for a field of view of vesicles at ~30°C (Fig. S4-6 and S9-11). Fiji (41) was used to crop the image to 300 x 300 pixels, to adjust brightness and contrast so that vesicle features were visible (gamma = 1), and to add scalebars.

Transition Temperatures and Plotting. We use T_{mix} to denote three possible transitions that result in an increase in the number of phases in a membrane as the temperature decreases: L→ L_o/L_d, L→ S/L, and S/L→ S/L_o/L_d. The corresponding phase transition temperatures were first estimated, then measured. To make estimates, an initial coverslip assembly was heated at ~1°C/min, and the approximate range of temperatures over which phase-separated domains disappeared was recorded. Temperature was then set 5-10°C above the estimated transition, and the initial coverslip assembly was discarded and replaced with a new one. Fields of vesicles (usually >100 vesicles total) were imaged as temperature was decreased stepwise over a total span of ≥12°C. At each step, temperature was allowed to equilibrate for ~1-2 min before images were collected. For each field of view, the number of vesicles with phase-separated membranes and the number with uniform membranes were tabulated by hand. Circular domains that immediately coalesced upon collision were categorized as liquid phase. Solid phases were categorized as having domains with irregular shapes and/or domains that did not coalesce upon collision. In vesicles with very small area fractions of ordered phase, domains are so small that L_o and solid phases are difficult to distinguish. To determine the transition temperature, T_{mix}, a non-linear least squares algorithm from the open-source Python library, SciPy (42), was used to fit a sigmoidal curve of the form

$$\% \text{ Phase separated} = Max / [1 + e^{-k (T - T_{mix})}]$$

where *Max* is the maximum percent of vesicles that phase separate, *k* is the steepness of the curve, T_{mix} is the phase transition temperature located at the inflection point, and *T* is the temperature input variable. Plots of sigmoidal curves and fits were generated using the open-source Python library, Matplotlib (43), and are shown in Fig. S13-18. Plots of ternary phase diagrams were generated using the open-source Python library, mpltern (44). Interpolation between data points was achived with the griddata function in SciPy (42), which uses a Clough-Tocher scheme. All phase transition temperatures and errors are recorded in Tables S1-S8.

Measurement Errors. Error was quantified in four ways: **1)** For each sigmoidal curve, a 95% confidence interval (shown in Fig. S13-S18) was calculated, which provides errors of the fit

parameters. For T_{mix}, these fit errors are very small (<0.5°C) and are recorded in Tables S1-S8. **2)** For each sigmoidal curve, ½($|T_{90\%} - T_{10\%}|$) was calculated; this is half the range between the temperatures at which 90% of vesicles and 10% of vesicles are phase separated (Fig. S13-S18). These errors are recorded in Tables S1-S8. **3)** Three researchers tabulated numbers of phase-separated vesicles. To estimate the error that each individual introduced, all three researchers tabulated the same four data sets in Fig. S19. The resulting values of T_{mix} were typically within ±0.5°C. **4)** Sample-to-sample variation was measured by standard deviations across three duplicate sample sets and nineteen triplicate sample sets, shown in bolded data in Tables S1-8. The average standard deviation across all duplicate and triplicate sample sets was 0.7 ± 0.6°C and ranged from 0.1°C to 2.8°C.

RESULTS AND DISCUSSION

Vesicles are stable over wide ranges of DPPE mole fractions.

Ternary lipid mixtures containing DPPE produce stable, giant vesicles over broad spans of temperature and mole fraction (Fig. 2). The solubility of cholesterol in PE-membranes is lower than in equivalent PC-membranes – we maintained cholesterol fractions at ≤50 mol% to avoid exceeding cholesterol's solubility limit of 51 ± 3 mol% in DPPE membranes (45, 46). Although DPPE's saturated chains make it less conical than unsaturated PE-lipids, challenges persist in forming ternary vesicles with DPPE. Nearly every vesicle has high-curvature tubules (Fig. 2), and nearly every field of view contains a bright aggregate of lipids, unlike in corresponding samples with PC-lipids (47, 48). Moreover, when the combined mole fraction of DPPE and cholesterol exceeds 80-90 mol%, vesicle yields are too small for data collection. Although tubulation is not a necessary outcome of producing vesicles with PE-lipids (49), tubulation is enhanced by temperature fluctuations near solid-liquid and liquid-liquid transitions (49, 50) and by inclusion of cholesterol, which can reduce the temperature of the lamellar-H_{II} transition by 20°C (18).

Figure 2. Lipid mixtures containing DPPE can form stable vesicles that phase separate, often with tubes. A) At high temperature (50°C), vesicle membranes of 20:50:30 DOPC/DPPE/cholesterol are in one liquid phase. **B)** At lower temperature (46°C), vesicles in the same sample phase separate into coexisting solid (dark) and liquid (bright) phases. Solid phases are characterized by noncircular domains that do not coalesce over time. **C)** Coexisting L_o and L_d liquid phases appear for some ratios of DiphyPC/DPPE/cholesterol, shown here for 30:40:30 at 45°C. Liquid phases are characterized by circular domains that merge in taut vesicles. **D)** Long tubules often emanate from vesicles containing DPPE, shown here for a ratio of 10:60:30 of DOPC/DPPE/cholesterol at 52°C. Contrast between the two phases arises from 0.8 mol % fluorescently labeled rhodamine-DPPE, which preferentially partitions to the L_d phase.

Replacing DPPC with DPPE eradicates (or severely limits) liquid-liquid coexistence in ternary membranes.

Ternary membranes containing DPPE differ starkly from membranes containing the corresponding PC-lipid. In Fig. 3, liquid-liquid coexistence is obliterated across the entire phase diagram of DOPC/DPPE/cholesterol (Fig. 3B), whereas it persists across broad ranges of temperature and lipid composition in vesicles of corresponding PC-lipids (Fig. 3C) (47, 51). In place of liquid-liquid coexistence, solid-liquid coexistence is observed (Fig. 3B). Solid domains are often identified by non-circular shapes, similar to the hexagonal and flower-shaped domains in binary DOPC/DPPC vesicles (52, 53). The area fraction of dark, noncircular, solid domains in vesicles monotonically increases as the mole fraction of DPPE increases (toward the bottom right vertex in the small images in Fig. 3A and in the large fields of view in Fig. S4-S6). This

observation is in line with the expectation that solid phases are enriched in lipids with high melting temperatures. Transition temperatures along the binary DOPC/DPPE axis follow the same trend but are consistently ≥2°C higher than values reported for vesicles made by a related technique (gentle hydration at 70.0°C (54)), even at low fractions of DPPE. Even though this shift is small, it is not expected because DOPC and a similar PE-lipid (DLPE; di(12:0)PE) incorporate equally well in vesicles made by electroformation and by gentle hydration (40).

The lack of liquid-liquid phase separation in Fig. 3A is surprising because DPPE's melting temperature (T_{melt} = 62.3 ± 5.0°C (26)) is close to the melting temperature of DAPC (di(20:0)PC, T_{melt} = 65.3 ± 1.5°C (10)), and yet coexisting liquid phases persist in equivalent ternary membranes with DAPC (47). Increasing the T_{melt} of PC-lipids or sphingomyelins in ternary mixtures tends to simply increase liquid-liquid transition temperatures (11, 47, 55) rather than suppress liquid-liquid coexistence. Instead, Fig. 3A is reminiscent of ternary phase diagrams in which palmitoyl-ceramide (PCer) is the high melting temperature lipid (14). This is surprising because PCer is structurally very different from DPPE: PCer has no polar headgroup at all, which gives it an extraordinarily high melting temperature of 90.0°C (56).

Figure 3. Vesicles of DOPC/DPPE/cholesterol exhibit coexisting solid and liquid phases (rather than coexisting liquid phases). A) Representative images at 30°C of vesicles of DOPC/DPPE/cholesterol. Solid phases are dark, and liquid phases are bright. Scale bars: 20 µm. Black X's mark compositions for which vesicle yields were too small to measure transition temperatures. Cholesterol mole fractions above 50% were avoided (stipple) because they likely exceed maximum cholesterol levels that incorporate in membranes with PE-lipids (45). Wider fields of view with larger images are in Fig. S4-S6. **B)** Transition temperatures in vesicles of DOPC/DPPE/cholesterol, from one liquid phase at high temperature to coexisting solid and liquid phases at lower temperatures (filled squares). Unfilled circles indicate that only one liquid phase was observed. Errors in T_{mix} are shown in Fig. S3. Sigmoidal curves used to calculate T_{mix} are in Fig. S13-S15. All values are recorded in Table S1. **C)** Transition temperatures in vesicles of DOPC/DPPC/cholesterol, from one liquid phase at high temperature to either coexisting L_o and L_d phases (filled circles) or coexisting solid and liquid phases (squares), reproduced from (47).

We hypothesized that replacing the low-T_{melt} lipid of DOPC with DiphyPC (also known as di(16:0-4Me)PC) might produce liquid-liquid phase separation in membranes containing DPPE (Fig. 4). The methylated chains of DiphyPC give it a very low T_{melt} of <-120°C (37). Even though DiphyPC is typically associated with robust liquid-liquid phase separation over broad composition ranges of PC-lipids (6, 48), in vesicles containing DPPE, coexisting liquid phases appear over only a small island of mole ratios near the center of the phase diagram (Fig. 4B). Liquid phases are characterized by round domains (Fig. 2C). In taut vesicles, these domains diffuse, collide, and coalesce until only one domain of each phase remains. The liquid ordered (L_o) phase is identified by being enriched in high-T_{melt} lipids and by excluding the dye relative to the liquid disordered (L_d) phase, as in Fig. S8. When coexisting solid and liquid phases are observed, the area fraction of solid phase increases with DPPE mole fraction, indicating the solid is enriched in DPPE (Fig. 4A, and large fields of view in Fig. S9-S11).

Figure 4. Vesicles of DiPhyPC/DPPE/cholesterol exhibit coexisting liquid phases or coexisting solid and liquid phases. A) Representative images at 30°C of vesicles of DiphyPC/DPPE/cholesterol. Dark, noncircular domains are solid. L_d domains are bright with respect to both L_o and solid domains. Scale bars: 20 μm. Black X's mark compositions for which vesicle yields were too small to measure transition temperatures. Cholesterol mole fractions above 50% were avoided (stipple) because they likely exceed maximum cholesterol levels that incorporate in membranes with PE-lipids (45). Wider fields of view are in Fig. S9-11. **B)** Transition temperatures in vesicles of DiphyPC/DPPE/cholesterol, from one liquid phase at high temperature to either coexisting solid and liquid phases (grey squares) or coexisting L_o and L_d phases (red circles). Unfilled circles indicate that only one liquid phase was observed. At three lipid ratios (10:60:30, 10:70:20, and 20:60:20 DiphyPC/DPPE/cholesterol), vesicle membranes were in one liquid phase at high temperature, coexisting solid and liquid phases at an intermediate temperature, and three coexisting phases (solid, L_o, and L_d) at a lower temperature. Errors in T_{mix} are shown in Fig. S7. Sigmoidal curves used to calculate T_{mix} values are in Fig. S16-S18. All values are recorded in Tables S2-S3. **C)** Transition temperatures in vesicles of DiphyPC/DPPC/cholesterol, from one liquid phase at high temperature to either coexisting L_o and L_d phases (filled circles) or coexisting solid and liquid phases (squares), reproduced from (48). Areas between data points are interpolated and do not reflect uncertainties in data points, which are quoted as midpoints in temperature ranges that span ±2°C (48).

Replacing DPPC with DPPE makes Lo and solid phases distinguishable by microscopy.

An unexpected advantage of incorporating DPPE into ternary membranes is that L_o and solid phases become distinguishable by fluorescence microscopy, and subsequent transitions into the 3-phase region (L_o, L_d, and solid) are clearly visible (Fig. 5). Previously, 3-phase coexistence has been observed by fluorescence microscopy for only a few combinations of dyes, lipids, and solutions (e.g., (57, 58)). As a result, identification of 3-phase regions by fluorescence microscopy has required observation of changes in domain shapes (55) or in the inability of domains to diffuse and coarsen quickly (48). When L_o and solid phases are not distinguishable, alternative spectroscopic techniques (51, 55, 59) or atomic force microscopy (58, 60) must be deployed to conclusively identify the 3-phase region.

Figure 5 shows an example of a liquid phase at a high temperature (57°C) demixing into coexisting liquid and solid phases at a lower temperature (53°C). As temperature decreases still further (48°C), three fluorescence levels are visible, a bright L_d phase, a gray L_o phase, and a dark solid phase. In many cases, as in Fig. 5, solid domains constrain the motion of the liquid domains. Additional images of vesicles with clear 3-phase coexistence are shown in Fig. S12.

Viewed in the 3-dimensional space of a ternary phase diagram, the volume in which all three phases (L_o, L_d, and solid) coexist is most easily imagined as an irregular triangular pyramid, with its tip pointing up toward high temperatures. In traditional ternary membranes with PC-lipids (e.g., DOPC/DPPC/cholesterol), the pyramid is shaped like a shard lying on its side, with two long edges defined by long tie-lines between the L_d phase and the other phases (59). In contrast, in the DiphyPC/DPPE/cholesterol phase diagram in Fig. 4B, the pyramid must be more compact because the L_o/L_d coexistence region is smaller. The face of the pyramid that touches the L_o/solid coexistence region must have a shallow slope because transitions as in Fig. 5 (from a 2-phase, L_o/solid region to a 3-phase, L_o/L_d/solid region) occur over lipid compositions that span at least 10 mol%.

Because L_o and solid phases are distinguishable by fluorescence microscopy, we can learn more about how the substitution of DPPE for DPPC results in a phase diagram with primarily solid-liquid coexistence (Fig. 4), as opposed to a diagram with only liquid-liquid coexistence (DiphyPC/DPPC/cholesterol, (48)). The change from DPPE to DPPC involves the addition of only three methyl groups to the lipid headgroup. In Fig. 6, we add these methyl groups one-by-one for three lipid ratios. Addition of each methyl group is expected to decrease the monolayer spontaneous curvature due to the lipid, based on trends for DOPE, DOPE-1Me, and DOPC (61). We find that largest change in the phase behavior occurs upon addition of the first methyl group, from DPPE to singly-methylated DPPE ("1Me"), whereupon temperatures of the 2-phase (solid/liquid) and 3-phase (L_o/L_d/solid) transitions become nearly indistinguishable (Fig. 6).

Figure 5. Liquid-liquid phase separation occurs at lower temperature than solid-liquid separation in 10:70:20 DiPhyPC/DPPE/cholesterol membranes. A) Vesicle membranes are in one liquid phase at high temperature. **B)** Solid domains nucleate at ~56°C. **C)** Domains of dark, noncircular, solid phase coexist with a bright, liquid phase. **D)** At ~52°C, the liquid phase begins demixing into L_o and L_d phases. The L_o phase is identified by an intermediate fluorescence level between the dark solid phase and bright L_d phase. **E)** At 48°C, micron-scale regions of three fluorescence levels are observed, corresponding to solid, L_o, and L_d phases. Solid phases are wet by L_o domains. **E-F)** With time, domains diffuse, and liquid domains can merge. **F-J)** Phase transitions are reversible with temperature. Images correspond to Video S1; timepoints for images A–J are 0, 16, 31, 40, 48, 54, 70, 76, 96 and 119 sec, respectively. These transitions occur in the same order in vesicles of 10:60:30 and 20:60:20 DiphyPC/DPPE/cholesterol.

Figure 6. Methylation of DPPE headgroup eradicates solid-liquid coexistence at high temperatures. As temperature is decreased in vesicles of 10:70:20, 10:60:30, and 20:60:20 DiphyPC/DPPE/cholesterol (denoted "PE" on each x-axis) transitions are observed from one liquid phase at high temperature, to coexisting solid and liquid phases, and then to three phases (L_o, L_d, and solid). In each of the three mixtures, DPPE was replaced by singly- ("1Me"), doubly- ("2Me"), or triply-methylated ("PC") versions of the molecule. In all three panels, gray squares show temperatures at which coexisting solid and liquid phases appear, and red circles show temperatures at which both L_o and L_d phases appear, for three independent measurements. Error bars are standard deviations of the triplicate measurements. Lines through triplicate averages are to guide the eye and are not fits to a model. Corresponding data are tabulated in Tables S2-8.

Of course, many PE-lipids beyond DPPE are relevant and interesting. One strong outcome of our research would be to facilitate future investigations of liquid-liquid phase separation in membranes containing other PE-lipids. Of high interest would be PE-lipids with unsaturated chains and with vinyl linkages to the chains (plasmalogens), both of which are prevalent in biological membranes and have been the subject of molecular dynamics simulations (3, 4, 33, 62–65).

CONCLUSIONS

In this manuscript, we substituted a PE-lipid (DPPE) for a PC-lipid (DPPC) in ternary lipid membranes. Changes to the molecular structure are small: these two lipids differ by only three methyl groups. However, subsequent changes to miscibility phase diagrams of ternary membranes containing these lipids are large. Whereas liquid-liquid phase separation is present in vesicles across a wide band of compositions composed of DOPC/DPPC/cholesterol, it is entirely absent from the phase diagram of vesicles of DOPC/DPPE/cholesterol, and only a small island of liquid-liquid phase separation appears in the phase diagram of vesicles of DiphyPC/DPPE/cholesterol. Inside that island, membranes of a small set of compositions begin

first in a liquid phase at high temperature, then separate into coexisting solid and liquid phases as temperature decreases, and then undergo a transition into a three-phase region after temperature is decreases further. This three-phase region contains coexisting liquid-ordered, liquid-disordered, and solid phases, all of which are distinguishable by fluorescence microscopy. In most other membranes, transitions into the three-phase region are difficult to observe by microscopy, due to solid phases nucleating within liquid-ordered phases and/or due to poor contrast of fluorophores between liquid-ordered and solid phases. Ideally, our investigation of ternary membranes containing DPPE will serve as a starting point for other researchers to map the phase behavior of membranes containing other PE-lipids, especially the unsaturated PEs or plasmalogens prevalent in biological membranes.

AUTHOR CONTRIBUTIONS

GJG, CEC, and SLK designed research and protocols. GJG, CLAC, and SN performed research. GJG, SN, and AMM analyzed data. GJG and SLK wrote the manuscript. GJG and SLK edited the manuscript.

ACKNOWLEDGEMENTS

This research was supported by National Science Foundation grant MCB-1925731 and MCB-2325819 to SLK. The authors thank Bob Weng for preliminary experiments.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

1. Leveille, C.L., C.E. Cornell, A.J. Merz, and S.L. Keller. 2022. Yeast cells actively tune their membranes to phase separate at temperatures that scale with growth temperatures. *Proc. Natl. Acad. Sci. USA*. 119:e2116007119.

2. Burns, M., K. Wisser, J. Wu, I. Levental, and S.L. Veatch. 2017. Miscibility Transition Temperature Scales with Growth Temperature in a Zebrafish Cell Line. *Biophys. J.* 113:1212–1222.

3. Lorent, J.H., K.R. Levental, L. Ganesan, G. Rivera-Longsworth, E. Sezgin, M. Doktorova, E. Lyman, and I. Levental. 2020. Plasma membranes are asymmetric in lipid unsaturation, packing and protein shape. *Nat. Chem. Biol.* 16:644–652.

4. Reinhard, J., C.L. Leveille, C.E. Cornell, A.J. Merz, C. Klose, R. Ernst, and S.L. Keller. 2023. Remodeling of yeast vacuole membrane lipidomes from the log (one phase) to stationary stage (two phases). *Biophys. J.* 122:1043–1057.

5. Kim, H., and I. Budin. 2024. Intracellular sphingolipid sorting drives membrane phase separation in the yeast vacuole. *J. Biol. Chem.* 300:105496.

6. Wilson, K.J., H.Q. Nguyen, J. Gervay-Hague, and S.L. Keller. 2024. Sterol–lipids enable large-scale, liquid–liquid phase separation in bilayer membranes of only two components. *Proc. Natl. Acad. Sci. USA*. 121:e2401241121.

7. Dietrich, C., L.A. Bagatolli, Z.N. Volovyk, N.L. Thompson, M. Levi, K. Jacobson, and E. Gratton. 2001. Lipid Rafts Reconstituted in Model Membranes. *Biophys. J.* 80:1417–1428.

8. Samsonov, A.V., I. Mihalyov, and F.S. Cohen. 2001. Characterization of Cholesterol-Sphingomyelin Domains and Their Dynamics in Bilayer Membranes. *Biophys. J.* 81:1486–1500.

9. Veatch, S.L., and S.L. Keller. 2002. Organization in Lipid Membranes Containing Cholesterol. *Phys. Rev. Lett.* 89:268101.

10. Koynova, R., and M. Caffrey. 1998. Phases and phase transitions of the phosphatidylcholines. *BBA - Rev. Biomembr.* 1376:91–145.

11. Veatch, S.L., and S.L. Keller. 2005. Miscibility Phase Diagrams of Giant Vesicles Containing Sphingomyelin. *Phys. Rev. Lett.* 94:148101.

12. Ionova, I.V., V.A. Livshits, and D. Marsh. 2012. Phase Diagram of Ternary Cholesterol/Palmitoylsphingomyelin/Palmitoyloleoyl-Phosphatidylcholine Mixtures: Spin-Label EPR Study of Lipid-Raft Formation. *Biophys. J.* 102:1856–1865.

13. Lin, W.-C., C.D. Blanchette, and M.L. Longo. 2007. Fluid-Phase Chain Unsaturation Controlling Domain Microstructure and Phase in Ternary Lipid Bilayers Containing GalCer and Cholesterol. *Biophys. J.* 92:2831–2841.

14. Castro, B.M., L.C. Silva, A. Fedorov, R.F.M. de Almeida, and M. Prieto. 2009. Cholesterol-rich Fluid Membranes Solubilize Ceramide Domains: Implications for the Structure and

Dynamics of Mammalian Intracellular and Plasma Membranes. *J. Biol. Chem.* 284:22978–22987.

15. Blosser, M.C., J.B. Starr, C.W. Turtle, J. Ashcraft, and S.L. Keller. 2013. Minimal Effect of Lipid Charge on Membrane Miscibility Phase Behavior in Three Ternary Systems. *Biophys. J.* 104:2629–2638.

16. Vequi-Suplicy, C.C., K.A. Riske, R.L. Knorr, and R. Dimova. 2010. Vesicles with charged domains. *BBA - Biomembr.* 1798:1338–1347.

17. Pitman, M.C., F. Suits, K. Gawrisch, and S.E. Feller. 2005. Molecular dynamics investigation of dynamical properties of phosphatidylethanolamine lipid bilayers. *J. Chem. Phys.* 122:244715.

18. Suits, F., M.C. Pitman, and S.E. Feller. 2005. Molecular dynamics investigation of the structural properties of phosphatidylethanolamine lipid bilayers. *J. Chem. Phys.* 122:244714.

19. Keller, S.L., S.M. Bezrukov, S.M. Gruner, M.W. Tate, I. Vodyanoy, and V.A. Parsegian. 1993. Probability of alamethicin conductance states varies with nonlamellar tendency of bilayer phospholipids. *Biophys. J.* 65:23–27.

20. Brown, M.F. 1994. Modulation of rhodopsin function by properties of the membrane bilayer. *Chem. Phys. Lipids*. 73:159–180.

21. Bogdanov, M., and W. Dowhan. 1998. Phospholipid-assisted protein folding: phosphatidylethanolamine is required at a late step of the conformational maturation of the polytopic membrane protein lactose permease. *EMBO J.* 17:5255–5264.

22. Wikström, M., A.A. Kelly, A. Georgiev, H.M. Eriksson, M.R. Klement, M. Bogdanov, W. Dowhan, and Å. Wieslander. 2009. Lipid-engineered *Escherichia coli* Membranes Reveal Critical Lipid Headgroup Size for Protein Function. *J. Biol. Chem.* 284:954–965.

23. Andersen, O.S., and R.E. Koeppe 2nd. 2007. Bilayer Thickness and Membrane Protein Function: An Energetic Perspective. *Annu. Rev. Biophys.* 36:107–130.

24. Cullis, P.R., M.J. Hope, and C.P.S. Tilcock. 1986. Lipid polymorphism and the roles of lipids in membranes. *Chem. Phys. Lipids*. 40:127–144.

25. Davies, M.A., W. Hubner, A. Blume, and R. Mendelsohn. 1992. Acyl chain conformational ordering in 1,2 dipalmitoylphosphatidylethanolamine. Integration of FT-IR and 2H NMR results. *Biophys. J.* 63:1059–1062.

26. Koynova, R., and M. Caffrey. 1994. Phases and phase transitions of the hydrated phosphatidylethanolamines. *Chem. Phys. Lipids*. 69:1–34.

27. Klose, C., M.A. Surma, M.J. Gerl, F. Meyenhofer, A. Shevchenko, and K. Simons. 2012. Flexibility of a Eukaryotic Lipidome – Insights from Yeast Lipidomics. *PLOS ONE*. 7:e35063.

28. Siegel, D.P. 1993. Energetics of intermediates in membrane fusion: comparison of stalk and inverted micellar intermediate mechanisms. *Biophys. J.* 65:2124–2140.

29. Allender, D.W., and M. Schick. 2023. On the force between "rafts." *Eur. Phys. J. E*. 46:85.

30. Cornell, C.E., A.D. Skinkle, S. He, I. Leventaľ, K.R. Leventaľ, and S.L. Keller. 2018. Tuning Length Scales of Small Domains in Cell-Derived Membranes and Synthetic Model Membranes. *Biophys. J.* 115:690–701.

31. Shyamsunder, E., S.M. Gruner, M.W. Tate, D.C. Turner, P.T.C. So, and C.P.S. Tilcock. 1988. Observation of inverted cubic phase in hydrated dioleoylphosphatidylethanolamine membranes. *Biochem.* 27:2332–2336.

32. So, P.T.C., S.M. Gruner, and S. Erramilli. 1993. Pressure-induced topological phase transitions in membranes. *Phys. Rev. Lett.* 70:3455–3458.

33. Winnikoff, J.R., D. Milshteyn, S.J. Vargas-Urbano, M.A. Pedraza-Joya, A.M. Armando, O. Quehenberger, A. Sodt, R.E. Gillilan, E.A. Dennis, E. Lyman, S.H.D. Haddock, and I. Budin. 2024. Homeocurvature adaptation of phospholipids to pressure in deep-sea invertebrates. *Science*. 384:1482–1488.

34. Marrink, S.J., V. Corradi, P.C.T. Souza, H.I. Ingólfsson, D.P. Tieleman, and M.S.P. Sansom. 2019. Computational Modeling of Realistic Cell Membranes. *Chem. Rev.* 119:6184–6226.

35. Hossein, A., and M. Deserno. 2020. Spontaneous Curvature, Differential Stress, and Bending Modulus of Asymmetric Lipid Membranes. *Biophys. J.* 118:624–642.

36. Keller, S.L., S.M. Gruner, and K. Gawrisch. 1996. Small concentrations of alamethicin induce a cubic phase in bulk phosphatidylethanolamine mixtures. *BBA - Biomembr.* 1278:241–246.

37. Lindsey, H., N.O. Petersen, and S.I. Chan. 1979. Physicochemical characterization of 1,2-diphytanoyl-*sn*-glycero-3-phosphocholine in model membrane systems. *BBA - Biomembr.* 555:147–167.

38. Veatch, S.L. 2007. Electro-formation and fluorescence microscopy of giant vesicles with coexisting liquid phases. *Methods Mol. Biol.* 398:59–72.

39. Veatch, S.L., and S.L. Keller. 2005. Seeing spots: Complex phase behavior in simple membranes. *BBA - Mol. Cell Res.* 1746:172–185.

40. Weakly, H.M.J., K.J. Wilson, G.J. Goetz, E.L. Pruitt, A. Li, L. Xu, and S.L. Keller. 2024. Several common methods of making vesicles (except an emulsion method) capture intended lipid ratios. *Biophys. J.* 123:3452–3462.

41. Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for biological-image analysis. *Nat. Methods*. 9:676–682.

42. Virtanen, P., R. Gommers, T.E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E. Burovski, P. Peterson, W. Weckesser, J. Bright, S.J. van der Walt, M. Brett, J. Wilson, K.J. Millman, N. Mayorov, A.R.J. Nelson, E. Jones, R. Kern, E. Larson, C.J. Carey, İ. Polat, Y. Feng, E.W. Moore, J. VanderPlas, D. Laxalde, J. Perktold, R. Cimrman, I. Henriksen, E.A. Quintero, C.R. Harris, A.M. Archibald, A.H. Ribeiro, F. Pedregosa, and P. van Mulbregt.

2020. SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods*. 17:261–272.

43. Hunter, J.D. 2007. Matplotlib: A 2D Graphics Environment. *Comput. Sci. Eng.* 9:90–95.

44. Ikeda, Y. 2023. yuzie007/mpltern: 1.0.2. *Zenodo*. https://doi.org/10.5281/zenodo.8289090.

45. Huang, J., J.T. Buboltz, and G.W. Feigenson. 1999. Maximum solubility of cholesterol in phosphatidylcholine and phosphatidylethanolamine bilayers. *BBA - Biomembr.* 1417:89–100.

46. Shaikh, S.R., V. Cherezov, M. Caffrey, S.P. Soni, D. LoCascio, W. Stillwell, and S.R. Wassall. 2006. Molecular Organization of Cholesterol in Unsaturated Phosphatidylethanolamines: X-ray Diffraction and Solid State 2H NMR Reveal Differences with Phosphatidylcholines. *J. Am. Chem. Soc.* 128:5375–5383.

47. Veatch, S.L., and S.L. Keller. 2003. Separation of Liquid Phases in Giant Vesicles of Ternary Mixtures of Phospholipids and Cholesterol. *Biophys. J.* 85:3074–3083.

48. Veatch, S.L., K. Gawrisch, and S.L. Keller. 2006. Closed-Loop Miscibility Gap and Quantitative Tie-Lines in Ternary Membranes Containing Diphytanoyl PC. *Biophys. J.* 90:4428–4436.

49. Sakuma, Y., T. Taniguchi, T. Kawakatsu, and M. Imai. 2013. Tubular Membrane Formation of Binary Giant Unilamellar Vesicles Composed of Cylinder and Inverse-Cone-Shaped Lipids. *Biophys. J.* 105:2074–2081.

50. Talbot, E.L., J. Kotar, L.D. Michele, and P. Cicuta. 2019. Directed tubule growth from giant unilamellar vesicles in a thermal gradient. *Soft Matter*. 15:1676–1683.

51. Davis, J.H., J.J. Clair, and J. Juhasz. 2009. Phase Equilibria in DOPC/DPPC-d62/Cholesterol Mixtures. *Biophys. J.* 96:521–539.

52. Chen, D., and M.M. Santore. 2014. 1,2-Dipalmitoyl-sn-glycero-3-phosphocholine (DPPC)-Rich Domain Formation in Binary Phospholipid Vesicle Membranes: Two-Dimensional Nucleation and Growth. *Langmuir*. 30:9484–9493.

53. Wan, H., G. Jeon, W. Xin, G.M. Grason, and M.M. Santore. 2024. Flower-shaped 2D crystals grown in curved fluid vesicle membranes. *Nat. Commun.* 15:3442.

54. Sakuma, Y., M. Imai, M. Yanagisawa, and S. Komura. 2008. Adhesion of binary giant vesicles containing negative spontaneous curvature lipids induced by phase separation. *Eur. Phys. J. E*. 25:403–413.

55. Zhao, J., J. Wu, F.A. Heberle, T.T. Mills, P. Klawitter, G. Huang, G. Costanza, and G.W. Feigenson. 2007. Phase studies of model biomembranes: Complex behavior of DSPC/DOPC/Cholesterol. *BBA - Biomembr.* 1768:2764–2776.

56. Shah, J., J.M. Atienza, R.I. Duclos, A.V. Rawlings, Z. Dong, and G.G. Shipley. 1995. Structural and thermotropic properties of synthetic C16:0 (palmitoyl) ceramide: effect of hydration. *J. Lipid Res.* 36:1936–1944.

57. Pataraia, S., Y. Liu, R. Lipowsky, and R. Dimova. 2014. Effect of cytochrome c on the phase behavior of charged multicomponent lipid membranes. *BBA - Biomembr.* 1838:2036–2045.

58. Balleza, D., A. Mescola, N. Marín–Medina, G. Ragazzini, M. Pieruccini, P. Facci, and A. Alessandrini. 2019. Complex Phase Behavior of GUVs Containing Different Sphingomyelins. *Biophys. J.* 116:503–517.

59. Veatch, S.L., O. Soubias, S.L. Keller, and K. Gawrisch. 2007. Critical fluctuations in domain-forming lipid mixtures. *Proc. Natl. Acad. Sci. USA.* 104:17650–17655.

60. Aufderhorst-Roberts, A., U. Chandra, and S.D. Connell. 2017. Three-Phase Coexistence in Lipid Membranes. *Biophys. J.* 112:313–324.

61. Dymond, M.K. 2021. Lipid monolayer spontaneous curvatures: A collection of published values. *Chem. Phys. Lipids.* 239:105117.

62. Rog, T., and A. Koivuniemi. 2016. The biophysical properties of ethanolamine plasmalogens revealed by atomistic molecular dynamics simulations. *BBA - Biomembr.* 1858:97–103.

63. Ferraro, M., M. Masetti, M. Recanatini, A. Cavalli, and G. Bottegoni. 2015. Modeling lipid raft domains containing a mono-unsaturated phosphatidylethanolamine species. *RSC Adv.* 5:37102–37111.

64. West, A., V. Zoni, W.E.Jr. Teague, A.N. Leonard, S. Vanni, K. Gawrisch, S. Tristram-Nagle, J.N. Sachs, and J.B. Klauda. 2020. How Do Ethanolamine Plasmalogens Contribute to Order and Structure of Neurological Membranes? *J. Phys. Chem. B.* 124:828–839.

65. Girard, M., and T. Bereau. 2023. Induced asymmetries in membranes. *Biophys. J.* 122:2092–2098.

Supporting Material for:

**Micron-scale, liquid-liquid phase separation
in ternary lipid membranes containing DPPE**

Gunnar J. Goetz, Sasha Naomi, Angelique M. Madrigal, Catherine L. A. Chang,
Caitlin E. Cornell, and Sarah L. Keller

This file includes:

Captions for Videos S1-S3
Figures S1-S19
Data Tables S1-S8

Video S1

Timelapse video of membrane phase transitions for vesicles composed of 10:70:20 DiphyPC/DPPE/cholesterol. Temperature is first ramped from high to low. Solid domains nucleate at ~56°C. Domains of dark, noncircular, solid phase coexist with a bright, liquid phase. At ~52°C, the liquid phase demixes into L_o and L_d phases. The L_o phase is identified by an intermediate fluorescence level between the dark solid and bright L_d phases. At ~48°C, micron-scale regions of three fluorescence levels are observed, corresponding to solid, L_o, and L_d phases. Liquid domains can diffuse and merge over time but may be constrained by the solid phase. Solid phases are usually wet by L_o domains. Temperature is ramped back up to 52°C, and then to 56°C, showing that the phase transitions are reversible. Video corresponds to images shown in Fig. 5 of the main text.

Video S2

Timelapse video of membrane phase transitions for vesicles composed of 10:60:30 DiphyPC/DPPC/cholesterol. Temperature is first ramped from high to low. Solid domains nucleate at ~53°C. Domains of dark, noncircular, solid phase coexist with a bright, liquid phase. At ~51°C, the liquid phase demixes into L_o and L_d phases. The L_o phase is identified by an intermediate fluorescence level between the dark solid and bright L_d phases. At ~50°C, micron-scale regions of three fluorescence levels are observed, corresponding to solid, L_o, and L_d phases. Liquid domains can diffuse and merge over time but may be constrained by the solid phase. Solid phases are usually wet by L_o domains. Temperature is ramped back up to 52°C, and then to 56°C, showing that the phase transitions are reversible.

Video S3

Timelapse video of membrane phase transitions for vesicles composed of 20:60:20 DiphyPC/DPPC/cholesterol. Temperature is first ramped from high to low. Solid domains nucleate at ~54°C. Domains of dark, noncircular, solid phase coexist with a bright, liquid phase. At ~51°C, the liquid phase demixes into L_o and L_d phases. The L_o phase is identified by an intermediate fluorescence level between the dark solid and bright L_d phases. At ~50°C, micron-scale regions of three fluorescence levels are observed, corresponding to solid, L_o, and L_d phases. Liquid domains can diffuse and merge over time but may be constrained by the solid phase. Solid phases are usually wet by L_o domains. Temperature is ramped back up to 54°C, and then to 58°C, showing that the phase transitions are reversible.

Figure S1. Lipid films spread on slides at 60°C or 75°C (and then electroformed at 75°C) produce vesicles with similar T_{mix} values. In all experiments in the main text, vesicles were electroformed at 75°C. This temperature was chosen to be well above the T_{melt} of DPPE, which is 62.3 ± 5.0°C (26). Ideally, to prepare lipid films for electroformation, they should have been spread on indium-tin-oxide slides at the same high temperature. However, it is technically difficult to do so, given that chloroform boils at 61°C. Therefore, we assessed if spreading the films at 60°C, a temperature just below chloroform's boiling point, affected the miscibility phase transition temperature of subsequently electroformed vesicles. We would expect bigger effects for mixtures with higher mole fractions of DPPE, which are the first three mixtures in the plot. For these mixtures, differences between T_{mix} of vesicles made from films spread at 60°C and 75°C are inconsequential, and justify our decision to spread all films at 60°C. We were perplexed but not concerned by the larger difference in T_{mix} for the mixture with 20% DPPE (the rightmost mixture in the plot). For all four samples, the average difference in T_{melt} between samples spread at 60°C and 75°C is ±1.2°C (range 0.4°C to 2.4°C). For comparison, sample-to-sample variations are ±0.1 to ±1.3°C (range 0.1°C to 2.8°C).

Figure S2. DOPC/DPPE/cholesterol vesicles electroformed at 60°C (below the T_{melt} for DPPE) exhibit lower T_{mix} values than vesicles electroformed at 75°C, especially at higher DPPE mole fractions. In all experiments in the main text, vesicles were electroformed at 75°C, well above T_{melt} for DPPE (62.3 ± 5.0°C (26)). To illustrate the importance electroforming above T_{melt}, several ternary lipid mixtures composed of DOPC/DPPE/cholesterol were electroformed at 60°C, a temperature commonly used for electroforming vesicles containing PC-lipids such as POPC, DOPC, and DPPC. At higher DPPE compositions, the measured Tmix value is as much as ~10°C lower for samples electroformed at 60°C than at 75°C. These results imply that insufficient DPPE is incorporated into vesicles at 60°C and are consistent with (39).

Figure S3. Two methods of assessing error from single measurements of T_{mix} for vesicles of DOPC/DPPE/Cholesterol. A) Temperatures of phase transitions from one liquid at high temperature to coexisting solid and liquid phases at lower temperatures, from Fig. 3B. **B)** Error in the T_{mix} fit parameter calculated from 95% confidence intervals of the sigmoidal fits (Fig. S13-S15), centered at $T = T_{mix}$. This method yields very small errors in T_{mix} (<0.5°C), indicating the sigmoidal fits are good. However, the errors do not give information about the breadth of the transition. **C)** Error in T_{mix} calculated as $\frac{1}{2}(|T_{90\%} - T_{10\%}|)$, which is half the difference between the temperature at which 90% of vesicles are phase separated and that at which 10% are phase separated. Errors calculated by this method give more information about sample preparation and the shape of the phase boundary. Smaller errors reflect more uniformity in vesicle-to-vesicle composition and/or more constant transition temperatures from composition to composition. In the absence of independent measurements to measure standard deviations, this method can provide a useful way to assess experimental error. Corresponding data are tabulated in Table S1.

Figure S4. Representative fields of view for vesicles of DOPC/DPPE/cholesterol, covering the left third of the phase diagram in Figure 3b of the main text. The vertex at the left represents 100% DOPC, and high cholesterol is toward the top. Scalebars are 50 µm.

Figure S5. Representative fields of view for vesicles of DOPC/DPPE/cholesterol, covering the middle third of the phase diagram in Figure 3b of the main text. High DOPC fractions are on the left, high DPPE fractions on the right, and high cholesterol toward the top. Scalebars are 50 μm.

Figure S6. Representative fields of view for vesicles of DOPC/DPPE/cholesterol, covering the right third of the phase diagram in Figure 3b of the main text. The vertex at the right represents 100% DPPE, and high cholesterol is toward the top. Scalebars are 50 μm.

Figure S7. Two methods of assessing error from single measurements of T_{mix} for vesicles of DiphyPC/DPPE/Cholesterol. A) Temperatures of phase transitions from one liquid to coexisting solid and liquid phases (squares) or to coexisting L_o and L_d phases (circles), from Fig. 4B. **B)** Error in the T_{mix} fit parameter calculated from 95% confidence intervals of the sigmoidal fits (Fig. S16-S18), centered at $T = T_{mix}$. This method yields very small errors in T_{mix} (typically <0.5°C), indicating good sigmoidal fits. However, the errors do not give information about the breadth of the transition. Compositions for which the transition is from one liquid phase to coexisting L_o and L_d phases are bounded by a dotted black line. Compositions for which we could clearly observe a transition into a 3-phase region are bounded by a solid black line. **C)** Error in T_{mix} calculated as $\frac{1}{2}(|T_{90\%} - T_{10\%}|)$, which is half the difference between the temperature at which 90% of vesicles are phase separated and that at which 10% are phase separated. Errors calculated by this method give more information about sample preparation and the shape of the phase boundary. Smaller errors reflect more uniformity in vesicle-to-vesicle composition and/or more constant transition temperatures from composition to composition. In the absence of independent measurements to measure standard deviations, this method can provide a useful way to assess experimental error. Corresponding data are tabulated in Tables S2 and S3.

Figure S8. Representative fields of view for vesicles of DiphyPC/DPPE/cholesterol in the liquid-liquid coexistence region in Figure 4b in the main text. Vesicle images were collected between 46-47°C. The region with a red background approximates the L_o/L_d region at this temperature, and the region with a yellow background approximates the 3-phase triangle. Because data were taken in increments of 10 mol%, the fade from red to yellow represents uncertainty in the location of the left edge of the 3-phase triangle. Scalebars are 50 µm.

Figure S9. Representative fields of view for vesicles of DiphyPC/DPPE/cholesterol, covering the left third of the phase diagram in Figure 4b of the main text. The vertex at the left represents 100% DiphyPC. High cholesterol is toward the top. Scalebars are 50 μm.

Figure S10. Representative fields of view for vesicles of DiphyPC/DPPE/cholesterol, covering the middle third of the phase diagram in Figure 4b of the main text. High DiphyPC fractions are on the left, high DPPE fractions on the right, and high cholesterol toward the top. Scalebars are 50 μm.

Figure S11. Representative fields of view for vesicles of DiphyPC/DPPE/cholesterol, covering the right third of the phase diagram in Figure 4b of the main text. The vertex at the right represents 100% DPPE, and high cholesterol is toward the top. Scalebars are 50 μm.

Figure S12. Images of vesicles composed of 10:70:20 DiphyPC/DPPE/cholesterol with three fluorescence intensities due to 3-phase coexistence of L_d-phase domains (bright), a L_o-phase domains (middle gray), and solid-phase domains (dark). In all cases, vesicles are close to a transition temperature in which vesicles with S/L coexistence were cooled into a region of S/L_d/L_o coexistence. Shapes of the L_d domains are not static; they are typically noncircular due to low line tension, but they can also be noncircular due to constraints imposed by neighboring solid domains. **A-B)** Images of two different vesicles, collected with a 10x objective. **C-D)** Images of the top and bottom hemispheres of a single vesicle exhibiting three distinct fluorescence levels for coexisting L_o, L_d, and solid phases. As would be expected, images collected with a 60x objective are at higher resolution than those collected with a 10x objective, and it is easier to distinguish the three phases.

Figure S13. Sigmoidal fits with 95% confidence intervals (gray bands) for T_{mix} for vesicles of DOPC/DPPE/cholesterol, covering the left third of the phase diagram in Figure 3b of the main text. The vertex at the left represents 100% DOPC, and high cholesterol is toward the top. X-axes show temperatures (°C), y-axes show percent of phase-separated vesicles. Transition temperatures shown in gray font are transitions from one liquid phase at a high temperature to coexisting solid and liquid phases at a lower temperature.

Figure S14. Sigmoidal fits with 95% confidence intervals (gray bands) for vesicles of DOPC/DPPE/cholesterol, covering the middle third of the phase diagram in Figure 3b of the main text. High DOPC fractions are on the left, high DPPE fractions on the right, and high cholesterol toward the top. X-axes show temperatures (°C), y-axes show percent of phase-separated vesicles. Transition temperatures shown in gray font are transitions from one liquid phase at a high temperature to coexisting solid and liquid phases at a lower temperature.

Figure S15. Sigmoidal fits with 95% confidence intervals (gray bands) for vesicles of DOPC/DPPE/cholesterol, covering the right third of the phase diagram in Figure 3b of the main text. The vertex at the right represents 100% DPPE, and high cholesterol is toward the top. X-axes show temperatures (°C), y-axes show percent of phase-separated vesicles. Transition temperatures shown in gray font are transitions from one liquid phase at a high temperature to coexisting solid and liquid phases at a lower temperature.

Figure S16. Sigmoidal fits with 95% confidence intervals (gray bands) for vesicles of DiphyPC/DPPE/cholesterol, covering the left third of the phase diagram in Figure 4b of the main text. The vertex at the left represents 100% DiphyPC, and high cholesterol is toward the top. X-axes show temperatures (°C), y-axes show percent of phase-separated vesicles. Transition temperatures shown in gray font are transitions from one liquid phase at a high temperature to coexisting solid and liquid phases at a lower temperature.

Figure S17. Sigmoidal fits with 95% confidence intervals (red and gray bands) for vesicles of DiphyPC/DPPE/cholesterol, covering the middle third of the phase diagram in Figure 4b of the main text. High DiphyPC fractions are on the left, high DPPE fractions on the right, and high cholesterol toward the top. X-axes show temperatures (°C), y-axes show percent of phase-separated vesicles. Transition temperatures shown in gray font are transitions from one liquid phase at a high temperature to coexisting solid and liquid phases at a lower temperature. Transition temperatures shown in red font are transitions in which coexisting L_o and L_d phases appear as temperature decreases.

Figure S18. Sigmoidal fits with 95% confidence intervals (red and gray bands) for vesicles of DiphyPC/DPPE/cholesterol, covering the right third of the phase diagram as in Figure 4b of the main text. The vertex at the right represents 100% DPPE, and high cholesterol is toward the top. X-axes show temperatures (°C), y-axes show percent of phase-separated vesicles. Transition temperatures shown in gray font are transitions from one liquid phase at a high temperature to coexisting solid and liquid phases at a lower temperature. Transition temperatures shown in red font are transitions in which coexisting L_o and L_d phases appear as temperature decreases.

Figure S19. Values of T_{mix} are independent of which researcher analyzed the data. In this manuscript, three different researchers tabulated numbers of phase-separated vesicles from fields of view as in Fig. S4-6. To estimate the error that this introduced, all three researchers (represented by the square, triangle, and circle for Counters 1, 2, and 3) tabulated data from the same fields of view for vesicles made from the four compositions shown on the x-axis. In each case, the analyzed values of T_{mix} typically differed by ±0.5°C or less. For comparison, sample-to-sample variations are on the order of ±0.1 to ±1.3°C (range 0.1°C to 2.8°C).

DOPC (mol%)	DPPE (mol%)	Chol (mol%)	L→S/L T_{mix} (°C)	Error in T_{mix} fit parameter (°C)	½($|T_{90\%} - T_{10\%}|$) (°C)
90	10	00	27.8	0.1	2.5
80	20	00	38.9	0.3	2.7
80	10	10	30.2	0.4	2.9
70	30	00	44.4	0.2	2.0
70	20	10	37.8	0.2	2.2
70	10	20	26.7	0.4	2.6
60	40	00	47.4	0.2	1.1
60	30	10	43.6	0.1	1.3
60	20	20	39.2	0.1	2.0
60	10	30	24.5	0.3	2.4
50	50	00	53.1	0.0	1.3
50	40	10	48.3	0.1	1.3
50	30	20	43.3	0.2	1.6
50	20	30	38.0	0.1	2.1
50	10	40	21.9	0.3	2.1
40	60	00	54.3	0.0	0.7
40	50	10	52.1	0.1	1.1
40	40	20	47.4	0.1	1.3
40	30	30	41.3	0.4	1.7
40	20	40	30.9	0.4	1.9
40	10	50	17.6	0.3	2.0
30	60	10	55.6	0.0	0.8
30	50	20	50.7	0.3	2.1
30	40	30	44.1	0.1	1.1
30	30	40	35.7	0.1	1.0
30	20	50	23.3	0.5	2.9
20	70	10	55.3	0.2	1.7
20	60	20	55.1	0.1	1.3
20	50	30	49.2	0.5	1.5
20	40	40	39.8	0.4	2.9
20	30	50	26.0	0.5	3.6

Table S1. Data for transition temperatures (T_{mix}) from one liquid phase at high temperature to coexisting solid and liquid phases at low temperature (L→S/L) for vesicles composed of varying ratios of DOPC/DPPE/cholesterol.

DiphyPC (mol%)	DPPE (mol%)	Chol (mol%)	L→S/L T_{mix} (°C)	Error in Tmix fit parameter (°C)	½($\|T_{90\%} - T_{10\%}\|$) (°C)
90	10	00	28.4	0.7	2.5
80	**20**	**00**	**39.7**	**0.5**	**2.7**
80	**20**	**00**	**38.6**	**0.1**	**2.9**
80	10	10	29.8	1.4	2.0
70	**30**	**00**	**43.4**	**0.0**	**2.2**
70	**30**	**00**	**39.4**	**1.7**	**2.6**
70	20	10	40.2	0.2	1.1
70	10	20	30.5	1.3	1.3
60	40	00	46.6	0.4	2.0
60	30	10	45.2	1.2	2.4
60	20	20	41.1	0.3	1.3
60	10	30	33.0	0.4	1.3
50	50	00	53.7	0.1	1.6
50	40	10	49.3	0.2	2.1
50	**30**	**20**	**44.5**	**0.3**	**2.1**
50	**30**	**20**	**44.4**	**0.2**	**0.7**
50	20	30	40.7	0.1	1.1
40	60	00	54.7	0.2	1.3
40	50	10	54.6	0.3	1.7
40	20	40	43.0	0.2	1.9
30	60	10	55.6	0.2	2.0
20	70	10	57.0	0.0	0.8
20	**60**	**20**	**54.5**	**0.1**	**0.9**
20	**60**	**20**	**53.3**	**0.1**	**1.3**
20	**60**	**20**	**54.2**	**0.3**	**1.8**
10	**70**	**20**	**55.6**	**0.0**	**0.7**
10	**70**	**20**	**55.9**	**0.1**	**1.0**
10	**70**	**20**	**56.0**	**0.2**	**0.8**
10	**60**	**30**	**52.4**	**0.4**	**1.9**
10	**60**	**30**	**52.2**	**1.2**	**1.5**
10	**60**	**30**	**53.3**	**0.6**	**1.6**

Table S2. Data for transition temperatures (T_{mix}) from one liquid phase at high temperature to coexisting solid and liquid phases at low temperature (L→S/L) for vesicles composed of varying ratios of DiphyPC/DPPE/cholesterol. Rows in bold font indicate repeated samples from duplicate or triplicate independent experiments.

DiphyPC (mol%)	DPPE (mol%)	Chol (mol%)	$L \rightarrow L_o/L_d$ T_{mix} (°C)	Error in Tmix fit parameter (°C)	½($|T_{90\%} - T_{10\%}|$) (°C)
40	40	20	49.6	0.2	2.5
40	30	30	46.8	0.3	2.7
30	50	20	50.8	0.1	2.9
30	40	30	47.9	0.2	2.0
30	30	40	47.1	0.2	2.2
20	**60**	**20**	**49.9**	**0.1**	**2.6**
20	**60**	**20**	**50.4**	**0.1**	**1.1**
20	**60**	**20**	**51.5**	**0.2**	**1.3**
20	50	30	50.2	0.0	2.0
20	40	40	50.5	0.1	2.4
10	**70**	**20**	**51.3**	**0.0**	**1.3**
10	**70**	**20**	**51.0**	**0.0**	**1.3**
10	**70**	**20**	**51.4**	**0.1**	**1.6**
10	**60**	**30**	**51.1**	**0.2**	**2.1**
10	**60**	**30**	**51.2**	**0.0**	**2.1**
10	**60**	**30**	**51.3**	**0.0**	**0.7**
10	50	40	50.3	0.1	1.1

Table S3. Data for transition temperatures (T_{mix}) for vesicles with a liquid phase (and perhaps a solid phase) at high temperatures, and coexisting L_o and L_d phases at lower temperatures (L→ L_o/L_d), for vesicles composed of varying ratios of DiphyPC/DPPE/cholesterol. Rows in bold font indicate repeated samples from duplicate or triplicate independent experiments.

DiphyPC (mol%)	DPPE-1Me (mol%)	Chol (mol%)	L→S/L T_{mix} (°C)	Error in Tmix fit parameter (°C)	½($\|T_{90\%} - T_{10\%}\|$) (°C)
20	60	20	48.3	0.2	1.0
20	60	20	49.1	0.2	0.6
20	60	20	48.6	0.4	1.5
10	70	20	50.3	0.5	1.4
10	70	20	50.6	0.0	0.7
10	70	20	50.5	0.1	0.9
10	60	30	49.5	*	0.1
10	60	30	49.6	0.1	0.4
10	60	30	47.8	*	2.1

Table S4. Data for transition temperatures (T_{mix}) from one liquid phase at high temperature to coexisting solid and liquid phases at low temperature (L→S/L) for vesicles composed of varying ratios of DiphyPC/DPPE-1Me/cholesterol. Rows in bold font indicate repeated samples from duplicate or triplicate independent experiments. Star symbols (*) indicate T_{mix} fit parameter values that were non-physical due to the proximity of the transition to the 3-phase region at temperatures just below the L→S/L T_{mix}. The temperature range to observe S/L coexistence narrows with increasing headgroup methylation and increasing DOPC composition.

DiphyPC (mol%)	DPPE-1Me (mol%)	Chol (mol%)	L→ L_o/L_d T_{mix} (°C)	Error in Tmix fit parameter (°C)	½($\|T_{90\%} - T_{10\%}\|$) (°C)
20	60	20	48.8	0.1	0.4
20	60	20	48.8	0.1	0.5
20	60	20	48.6	0.0	0.4
10	70	20	49.4	0.0	0.2
10	70	20	49.2	0.0	0.4
10	70	20	49.9	0.1	0.6
10	60	30	49.5	0.0	0.3
10	60	30	49.3	0.0	0.3
10	60	30	49.7	0.0	0.3

Table S5. Data for transition temperatures (T_{mix}) for vesicles with a liquid phase (and perhaps a solid phase) at high temperatures, and coexisting L_o and L_d phases at lower temperatures (L→ L_o/L_d), for vesicles composed of varying ratios of DiphyPC/DPPE-1Me/cholesterol. Rows in bold font indicate repeated samples from duplicate or triplicate independent experiments.

DiphyPC (mol%)	DPPE-2Me (mol%)	Chol (mol%)	T_{mix} (°C)	Error in Tmix fit parameter (°C)	½($\|T_{90\%} - T_{10\%}\|$) (°C)
10	70	20	46.7	0.1	0.5
10	70	20	44.7	0.1	0.5
10	70	20	45.5	0.1	0.6

Table S6. Data for transition temperatures (T_{mix}) from one liquid phase at high temperature to coexisting solid and liquid phases at low temperature (L→S/L) for vesicles composed of varying ratios of DiphyPC/DPPE-2Me/cholesterol. Rows in bold font indicate repeated samples from duplicate or triplicate independent experiments.

DiphyPC (mol%)	DPPE-2Me (mol%)	Chol (mol%)	L→ L_o/L_d T_{mix} (°C)	Error in Tmix fit parameter (°C)	½($\|T_{90\%} - T_{10\%}\|$) (°C)
20	60	20	45.8	0.0	0.6
20	60	20	45.2	0.1	0.8
20	60	20	46.3	0.0	0.4
10	70	20	44.4	0.1	0.7
10	70	20	46.3	0.1	0.3
10	70	20	45.3	0.0	0.6
10	60	30	47.2	0.0	1.1
10	60	30	47.3	0.1	0.3
10	60	30	46.3	0.1	0.9

Table S7. Data for transition temperatures (T_{mix}) for vesicles with a liquid phase (and perhaps a solid phase) at high temperatures, and coexisting L_o and L_d phases at lower temperatures (L→ L_o/L_d), for vesicles composed of varying ratios of DiphyPC/DPPE-2Me/cholesterol. Rows in bold font indicate repeated samples from duplicate or triplicate independent experiments.

DiphyPC (mol%)	DPPC (mol%)	Chol (mol%)	L→ L$_o$/L$_d$ T$_{mix}$ (°C)	Error in Tmix fit parameter (°C)	½(|T$_{90\%}$ − T$_{10\%}$|) (°C)
20	60	20	41.4	0.4	2.0
20	60	20	42.9	0.3	2.3
20	60	20	40.8	0.3	2.4
10	70	20	40.4	0.2	1.1
10	70	20	38.5	0.4	2.5
10	70	20	39.1	0.2	2.1
10	60	30	43.0	0.2	2.2
10	60	30	43.8	0.3	1.9
10	60	30	44.8	0.1	1.8

Table S8. Data for transition temperatures (T_{mix}) for vesicles with a liquid phase (and perhaps a solid phase) at high temperatures, and coexisting L$_o$ and L$_d$ phases at lower temperatures (L→ L$_o$/L$_d$), for vesicles composed of varying ratios of DiphyPC/DPPC/cholesterol. Rows in bold font indicate repeated samples from duplicate or triplicate independent experiments.