Molecular mechanism of exchange coupling in CLC chloride/proton antiporters

Deniz Aydin[1,2,3,4,*], Chih-Ta Chien[5,6,*], Jürgen Kreiter[1,*], Amy Nava[1,*], Jasmina M. Portasikova[7,8], Lukas Fojtik[7,8], Catalina Mosquera Salcedo[1], Petr Man[8], Ron O. Dror[1,2,3,4,†], Wah Chiu[5,9,†], Merritt Maduke[1,†]

[1]Stanford University, Department of Molecular and Cellular Physiology

[2]Department of Computer Science, Stanford University, Stanford, CA 94305

[3]Department of Structural Biology, Stanford University, Stanford, CA 94305

[4]Institute for Computational and Mathematical Engineering, Stanford University, Stanford, CA 94305

[5]Department of Bioengineering and Department of Microbiology and Immunology, Stanford University, Stanford, 94305

[6]National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), National Institutes of Health, Bethesda, MD, 20892

[7]Department of Biochemistry, Faculty of Science, Charles University, Hlavova 2030/8, 128 43, Prague 2, Czech Republic

[8]Institute of Microbiology of the Czech Academy of Sciences, Division BioCeV, Prumyslova 595, 252 50 Vestec, Czech Republic

[9]Division of CryoEM and Bioimaging, SSRL, SLAC National Accelerator Laboratory, Stanford University, Menlo Park 94025

[*]co-first authors

[†]co-corresponding authors

Correspondence:

Dr. Merritt Maduke	Dr. Wah Chiu	Dr. Ron O. Dror
279 Campus Drive West	318 Campus Drive West	353 Jane Stanford Way
Stanford, CA 94035	Stanford, CA 94305	Stanford, CA 94305
Email: maduke@stanford.edu	wahc@stanford.edu	ron.dror@stanford.edu

Abstract

The ubiquitous CLC membrane transporters are unique in their ability to exchange anions for cations. Despite extensive study, there is no mechanistic model that fully explains their 2:1 Cl^-/H^+ stoichiometric exchange mechanism. Here, we provide such a model. Using differential hydrogen-deuterium exchange mass spectrometry, cryo-EM structure determination, and molecular dynamics simulations, we uncovered new conformational dynamics in CLC-ec1, a bacterial CLC homolog that has served as a paradigm for this family of transporters. Simulations based on a cryo-EM structure at pH 3 revealed critical steps in the transport mechanism, including release of Cl^- ions to the extracellular side, opening of the inner gate, and novel water wires that facilitate H^+ transport. Surprisingly, these water wires occurred independently of Cl^- binding, prompting us to reassess the relationship between Cl^- binding and Cl^-/H^+ coupling. Using isothermal titration calorimetry and quantitative flux assays on mutants with reduced Cl^- binding affinity, we conclude that, while Cl^- binding is necessary for coupling, even weak binding can support Cl^-/H^+ coupling. By integrating our findings with existing literature, we establish a complete and efficient CLC 2:1 Cl^-/H^+ exchange mechanism.

The CLC "chloride channels" are a biophysically fascinating gene family consisting of both

passive chloride (Cl^-) channels and active transporters that exchange chloride for protons (H^+) [1-5].

CLC transporters are ubiquitously expressed in intracellular membranes, and their dysfunction

disrupts ion homeostasis and causes Dent's disease, osteopetrosis, and neurodegenerative

disorders [4,6-12]. CLC transporters are also found widely in plants and microorganisms [4,13,14]. In bacteria,

CLC transporters support pathogenesis by facilitating extreme acid tolerance [15,16].

CLC transporters catalyze Cl^-/H^+ exchange transport with a stoichiometry of 2 Cl^- ions per H^+.

This 2:1 stoichiometry is observed in all CLC transporter homologs under all experimental conditions

measured – including Cl^- and H^+ concentrations that vary by orders of magnitude. CLC-ec1 is an *E.*

coli homolog that was initially examined due to its functional similarity to mammalian homologs,

together with its superior suitability for structural analysis [17-19]. CLC-ec1 was the first CLC

demonstrated to be an active transporter instead of a channel[1], and has proven valuable for guiding

studies of mammalian CLCs, both transporters and channels [2,3,20-22]. CLC-ec1 remains a useful

paradigm, as it is currently the only transporter homolog for which multiple conformations have been

determined at high resolution [19,23-25] and the only homolog whose ion-binding properties have been

quantified by isothermal titration calorimetry [26,27]. In addition, the availability of a functional cysteine-

less variant makes it amenable to spectroscopic approaches [28-30].

Studies of CLC-ec1's structure, function, and dynamics have provided essential clues to the

CLC Cl^-/H^+ transport mechanism. Structurally, CLCs are homodimers in which each subunit can act

independently [4]. Each subunit contains permeation pathways for Cl^- and H^+, which are shared along

the extracellular section and then diverge toward the intracellular side (**Fig. 1a**). The Cl^- pathway is

gated by a conserved glutamate ("E_{gate}" – E148 in CLC-ec1), which also functions as a H^+-transfer

site (**Fig. 1a**). This pathway is defined by two anion-binding sites, S_{ext} and S_{cen}, in the middle of each

subunit. These sites can be occupied by the negatively charged E_{gate} side chain ("middle" and "down"

conformations in **Fig.1b**) or by Cl^- ions. The H^+ permeation pathway diverges from the Cl^- pathway

toward the intracellular side, as illustrated in **Fig.1a**. This positioning of the H^+ pathway is supported

by several lines of evidence. First, mutations along the intracellular segment of the pathway,

particularly at "Glu_{in}" (E203), severely repress H^+ transport [31,32]. Second, E_{gate} can occupy this

proposed H^+ pathway in its "out" conformation (see **Fig.1b)**. Third, molecular dynamics simulations

on CLC-ec1 detect water wires (chains of hydrogen-bonded water molecules) that provide a pathway

for H^+ conduction in this region [24,25,32-39]. Within this framework, the key to achieving 2:1 Cl^-/H^+

coupling must lie in coupling of the E_{gate} rotameric conformational changes to Cl^- and H^+ movement

in opposite directions through the pathways.

Although some important aspects of this coupling are understood—such as the correlation

between coupling and Cl^- binding to S_{cen} [26,32,37,40,41] and the synergistic binding of Cl^- and H^+ [26,27] —a

satisfactory model to fully explain the 2:1 exchange mechanism has remained elusive. Existing

models explain some of CLC-ec1's known properties but fall short of capturing the full picture. For

instance, proposed models tend to be coherent in only one direction of the transport cycle [24,38,42],

whereas CLC-ec1 facilitates transmembrane Cl^-/H^+ exchange with similar efficiency in both directions

[43]. Furthermore, all available structures show the inner gate of the Cl^- pathway as closed, leaving the

mechanism of Cl^- movement to and from the intracellular side unresolved.

A major challenge in the study of CLC antiporters is the apparently subtle nature of the protein

conformational changes, which is in stark contrast to the large conformational changes that are seen

in other secondary active transporters [44-46]. For instance, the crystal structure of WT CLC-ec1 shows

E_{gate} in the "middle" conformation (**Fig.1b**). When E_{gate} is mutated to glutamine (to simulate the

protonation that occurs as part of the transport cycle), the resulting structure is nearly

indistinguishable from that of WT, differing only in the rotation of the side chain (from "middle" to "up"),

and with the extracellular pathway still narrower than a Cl^- ion. This observation raised an important

question: are conformational changes in CLC transporters inherently subtle compared to those in

other transporters, or do significant conformational shifts remain undiscovered? This question drove

us, and others, to explore these changes outside the constraints of crystallization [30,47-51]. To do so, we

utilized low-resolution spectroscopic techniques to analyze conformational dynamics in E_{gate} mutant

proteins, as well as in wild-type proteins under conditions of lowered pH to facilitate the protonation

of E_{gate}. Through these studies, we identified a mutant ("QQQ") that was stabilized in the protonated

state, allowing us to determine its high-resolution structure and reveal molecular details of

conformational change, including widening of the extracellular pathway and the positioning of E_{gate} in

the "out" conformation [24]. Subsequently, structures of WT CLC-ec1 at pH 4.5 recapitulated many of

the conformational features observed in the QQQ structure as well as new conformational changes

in the N-terminal region and at the dimer interface[25]. However, none of these structures illuminate the

mechanism of inner-gate opening. This void leaves lingering questions and suggests we may still be

missing important conformational changes, particularly given conformational dynamics observed

spectroscopically in structures lining the inner Cl^- pathway [47,50].

Here, we explored CLC-ec1's pH-dependent conformational change at even lower pH. CLC-

ec1 chloride transport rates increase as the pH is lowered from 4.5 to 3.0 [52], demonstrating the

protein's ability to undergo conformational cycling under these conditions. Additionally, low pH aligns

with CLC-ec1's physiological role in extreme acid tolerance within the mammalian stomach [15]. When

we lowered the pH from 4.5 to 3.0, we observed a sharp increase in protein conformational dynamics

detected by hydrogen deuterium exchange spectrometry. This observation prompted us to explore

the molecular details of this conformational change using single particle cryo-electron microscopy

(cryo-EM) structure determination. Our cryo-EM structures revealed subtle but consequential

conformational changes between pH 4.0 and 3.0, with molecular dynamics (MD) simulations of the

pH 3.0 structure providing unprecedented insights, including new water wires and opening of the inner

gate. Integrating results from structure, function, and simulation studies, we present a model to

explain the bidirectional 2:1 Cl^-/H^+ stoichiometric exchange mechanism of CLC exchange

transporters.

RESULTS

Hydrogen-deuterium exchange mass spectrometry reveals two pH-dependent conformational

transitions in CLC-ec1

To investigate pH-dependent changes in CLC-ec1 conformational dynamics, we first

employed hydrogen-deuterium exchange mass spectrometry (HDX-MS). This technique offers an

unbiased (label-free) spectrometric approach to revealing structural properties and conformational

dynamics in proteins [53,54] with virtually no limitation in experimental conditions (pH, temperature, buffer

composition). It is especially advantageous for studying CLC-ec1 because it can identify subtle

conformational changes in confined areas, such as the inner gate, without requiring a bulky label.

HDX is based on the principle that structurally rigid, H-bonded protein regions incorporate deuterium

into backbone amides slowly, whereas water-accessible dynamic protein regions do so rapidly.

CLC-ec1 was overexpressed in *E. coli* and purified using established methods [19,24,52]. The

concentrated protein sample was incubated in deuterated buffer for varying amounts of time to allow

hydrogen atoms in the protein backbone to exchange with deuterium from the solvent. After the

incubation period, the exchange reaction was quenched to stop further deuterium uptake, and the

sample digested with protease. The resulting peptides were analyzed using mass spectrometry (MS),

which measures the mass-to-charge ratio of the peptides, allowing for the determination of the extent

of deuterium incorporation and rates of hydrogen deuterium exchange (HDX). MS analysis of CLC-

ec1 identified 237 peptides, providing nearly complete (89%) coverage of the protein sequence

(**Extended Data Fig.1**).

Fig. 2a shows a differential heat map for deuterium incorporation observed in CLC-ec1 at pH

4.5 compared to pH 6.5. Each row shows comparisons at specific time points where the samples had

equivalent opportunity for deuterium exchange, accounting for the intrinsic pH-dependence of the

hydrogen deuterium exchange reaction, which is 100-fold faster at pH 6.5 than at pH 4.5. Several

regions of the CLC-ec1 protein exhibited increased HDX at pH 4.5, particularly around helices A, G,

H, I, P, and R, which are located on the intracellular side of the protein and along the subunit interface

(as mapped onto the protein structure at right in **Fig.2a**). This result closely aligns with previous cryo-

EM structural analyses at pH 7.5 and pH 4.5, which showed conformational changes consistent with

enhanced water accessibility in these regions at the lower pH[25]. **Fig.2b** shows a differential heat map

for deuterium incorporation observed in CLC-ec1 at pH 3.0 compared to pH 4.0. In this case, many

peptides displayed increased HDX at the lower pH. This result indicates a second conformational

transition, encompassing a broad swath of the protein. Importantly, the CLC-ec1 protein remained

stable under the pH 3.0 condition, as evaluated by size exclusion chromatography (**Extended Data**

Fig.2), indicating that the increase in HDX is not merely a result of protein unfolding.

The deuterium uptake plots used to create the HDX heat maps are presented in **Extended**

Data Fig.3. Fig.2c illustrates examples of these uptake plots for peptides selected to represent the

HDX properties and conformational transitions observed in this study. The location of each peptide

(residue numbers indicated above each plot) was mapped in color onto the protein structure shown

at center. For reference, a diagram of the protein with all helices labeled is shown in **Fig. 2d.** Peptides

336-344 (helix M, dark green) and 41-50 (helix B, pink) (plots shown at top) represent regions of the

protein that display little to no HDX at any pH. These peptides are in helices within the hydrophobic

core of the protein/membrane region. Peptides 17-25 (helix A, yellow) and 200-208 (helix H, blue)

(plots at left) represent the first pH-dependent conformational transition, as detected by the major shift

in the deuterium incorporation curve that occurs at pH 4.5 compared to pH 6.5. Peptides 146-163

(helix F, orange) and 110-116 (helix D, teal green) (plots at right) represent the second pH-dependent

conformational transition, detected by the shift in the deuterium update plot occurring at pH 3.0

compared to 3.5. Notably, these peptides encompass helices that constitute the outer and inner gates

of the Cl^- pathway, where enhanced water accessibility may indicate increased mobility of Cl^- and H^+

in these regions at pH 3.0. In contrast, water accessibility of other membrane helices (B, I, M, and N)

remained low at pH 3.0 (**Extended Data Fig.3**). Accessibility of helices C and J was also low, although

HDX coverage of these helices was not complete. Taken together, these data demonstrate two

distinct pH-dependent conformational changes in CLC-ec1, one of which had not been previously

detected.

Cryo-EM structures of CLC-ec1

To visualize these pH-dependent changes at high resolution, we determined cryo-EM

structures of CLC-ec1 at pH 7.5, 4.0, and 3.0, as summarized in **Fig. 3** and **Table 1**. Notably, the E_{gate}

was well-resolved in all three structures, as evidenced in the density maps and by the Q-scores [55]

(**Extended Data Figs.4-6**). In the pH 7.5 structure (3.18 Å resolution), E_{gate} occupies the S_{ext} anion-

binding site in the "middle" orientation (**Fig. 3a,d, Extended Data Fig. 4**). This structure resembles

the original CLC-ec1 crystal structure (1OTS, 2.51 Å resolution [23]), with a C-α r.m.s.d of 1.00 Å. In the

pH 4.0 structure (3.21 Å resolution), we observed a mixture of E_{gate} in both the "middle" and "out"

orientations, suggesting a transitional phase (**Fig. 3c,e, Extended Data Fig. 5**). In contrast, at pH 3.0

(3.14 Å resolution), E_{gate} is observed only in the "out" orientation (**Fig.3c,f, Extended Data Fig. 6**).

These results confirm the pH-dependent modulation of E_{gate} conformations.

In the pH 4.0 cryo-EM structure, the primary conformational change compared to pH 7.5 is an

outward shift of helices A and G-H-I, which opens access to the H^+ permeation pathway (**Fig. 3g**).

These changes at pH 4.0 compared to pH 7.5 are consistent with the increased water accessibility

observed by HDX at pH 4.0 (**Extended Data Fig. 2**) and were also seen in the published pH 4.5 cryo-

EM structure of CLC-ec1[25].

In the pH 3.0 cryo-EM structure, there is a subtle conformational change compared to pH 4.0,

with an overall C-α r.m.s.d of 0.92 Å. Notably, the pH 3.0 conformation resembles the "QQQ" CLC-

ec1 crystal structure (6V2J) (overall C-α r.m.s.d 0.84 Å) more closely than the pH 4.0 structure (overall

C-α r.m.s.d 1.27 Å) (**Fig. 3h**). Additionally, while the pore radius profiles of the pH 3.0 and 4.0

structures are similar to one another (**Fig. 3i**), at pH 3.0 E_{gate} is shifted towards the pore, and residue

F357, which in simulations lowers the energy barrier for Cl^- movement [38], is observed in the energy-

lowering "up" position (**Fig. 3j**). Together, these subtle conformational changes at pH 3.0 vs 4.0

motivated us to perform molecular dynamics simulations on the pH 3.0 structure.

Molecular dynamics simulations reveal water wires, Cl^- leaving, and inner-gate opening

Connected water wires can facilitate proton conduction via a Grotthuss mechanism[56,57]. Early

simulations on CLC-ec1 showed water wires transiently connecting E_{gate} to Glu_{in}[32-39], which is a

residue near the intracellular side of the transmembrane domain (**Fig. 1a**) whose neutralization has

a major impact on H^+ transport[24,31]. More recently, water wires connecting E_{gate} directly to the

intracellular solution were observed in MD simulations of the QQQ mutant crystal structure[24] and in

simulations of WT CLC-ec1 pH 4.5 cryo-EM structure [25]. These water wires provide a plausible

explanation for the transfer of H^+ to and from E_{gate} when it is in the "out" position. However, they do

not account for all H^+ transport steps needed to model a complete, reversible Cl^-/H^+ transport cycle,

including how E_{gate} becomes protonated when it is in the "down" position within the anion pathway.

Simulations of the new pH 3.0 cryo-EM structure could provide further insights into this question or

shed light on other aspects of the transport cycle.

We performed extended molecular dynamics simulations on the pH 3.0 cryo-EM structure in

order to evaluate protein dynamics and water-wire formation. We observed three classes of water

wires. Two classes (wires 2 and 3) resembled water wires seen in previous simulations, connecting

the intracellular solution to E_{gate} in the "out" position, whereas the third class (wire 1) was a novel

water wire, connecting the intracellular solution to the anion permeation pathway (**Fig. 4a**).

Water wires 2 and 3 offer a plausible explanation for how a proton moving to the inside is

coupled to two chloride ions moving out, as illustrated in **Fig. 4b.** Following the delivery of a proton

from the "out" E_{gate} to the intracellular solution, the resulting negatively charged E_{gate} will be

energetically favored to move from the hydrophobic core of the protein to the anion-friendly pore. To

achieve this, E_{gate} must displace any Cl^- ions occupying the pore. Since CLCs are antiporters with a

strict 2:1 Cl^-/H^+ exchange stoichiometry, the transport of a proton to the intracellular solution would

require the deprotonated E_{gate} to displace 2 Cl^- ions to the outside (**Fig. 4b**).

Capturing the displacement of 2 Cl^- ions by E_{gate} in simulations would provide strong support

for this proposed water-wire mechanism. We therefore performed simulations on the CLC-ec1 pH 3.0

structure with E_{gate} deprotonated and evaluated the presence of Cl^- in the permeation pathway by

plotting the distance from pore-lining inner-gate residue Y445 to the nearest Cl^- ion. Strikingly, in 8 of

10 independent 1.5-us trajectories, both Cl^- ions left the permeation pathway (**Fig. 4c**). In stark

contrast, Cl^- remained bound throughout all 10 simulations with E_{gate} protonated (**Extended data Fig.**

7). Thus, the deprotonation of E_{gate} is a crucial event that triggers expulsion of Cl^- from the permeation

pathway. Inspection of the trajectories revealed that the Cl^- ions always left to the extracellular side,

in harmony with the experimentally observed 2:1 Cl^-/H^+ exchange. Surprisingly, movement of E_{gate}

into the anion pathway was accompanied by reorientation of inner-gate residues S107 and Y445

(**Movie 1, Fig. 4d**).

Water wires do not depend on chloride

The binding of Cl^- to the S_{cen} site (**Fig. 1b**) is recognized as a critical feature in the CLC

coupling mechanism and is supported by multiple lines of evidence. First, mutations that weaken Cl^-

binding to S_{cen} also reduce coupling efficiency, resulting in fewer H^+ ions being transported per Cl^- ion

[26,40]. Second, polyatomic anions that bind less strongly to S_{cen} than Cl^- also exhibit weaker coupling

to H^+ [41]. Finally, Cl^- binding at S_{cen} is critical for water wires that connect Glu_{in} to E_{gate}, and mutations

that disrupt these water wires reduce coupling efficiency [32,37]. Given these previous findings, we

anticipated that the water wires connecting E_{gate} to the intracellular solution would depend on Cl^-

binding at S_{cen}. To test this hypothesis, we quantified water-wire formation in our two sets of

simulations: with E_{gate} protonated (Cl^- always present in the permeation pathway) and with E_{gate}

deprotonated (Cl^- absent during >70% of the simulation time). Surprisingly, we found that water wires

were similarly abundant in both simulations (**Fig. 4e**), indicating that water wires do not depend on Cl^-

binding.

Weak Cl^- binding at S_{cen} does not necessarily lead to uncoupling

This finding inspired us to investigate the relationship between Cl^- binding and the coupling

mechanism. Prior studies showed that any disruption to anion binding at S_{cen} results in uncoupling

[26,40,41]. As a result, it has been reasonably suggested that a key distinction between CLC transporters

and channel homologs lies in their Cl^- binding affinity at S_{cen}, with CLC channels functioning as

uncoupled channels rather than coupled transporters due to their reduced Cl^- binding affinity [27]. But

does weak Cl^--binding inevitably result in uncoupling? The mutagenesis studies that established the

link between weakened anion binding and uncoupling in CLC transporters involved mutation of

residues S107 and Y445. These residues not only directly interact with Cl^- at S_{cen}, but they also

directly form the inner gate that controls Cl^- passage to and from S_{cen} (**Fig.4d**). Consequently,

mutations at these positions physically alter the inner gate, and it is therefore possible that the

uncoupling caused by these mutations is more a result of gate disruption than of weakened Cl^-

binding.

To test the hypothesis that weak Cl^- binding alone does not cause uncoupling in the CLC

transporter, we needed to disrupt Cl^- binding while leaving the inner gate intact. To do so, we focused

on K131, as free energy calculations identified it as the most influential side chain in stabilizing Cl^-

binding at S_{cen} [58]. K131 does not directly interact with Cl^-; rather, the positively charged sidechain is

buried within the transmembrane domain, with the positively charged sidechain pointing towards S_{cen}

and stabilizing Cl⁻ binding from a distance of 7-9 Å from S_{cen} [58]. We first experimentally confirmed

weak Cl⁻ binding by K131A CLC-ec1, as well as the overall structural integrity of this mutant. The

cryo-EM structure of K131A CLC-ec1 exhibited minimal differences in overall structure compared with

WT CLC-ec1 (overall C-α r.m.s.d 0.52 Å), except for a clear reduction of Cl⁻ density at S_{cen} (**Fig. 5a,**

Extended Data Fig. 9). Isothermal titration calorimetry (ITC) confirmed weak Cl⁻ binding by CLC-ec1

K131A (**Fig. 5b**), with the lack of heat detection indicating Kd > 20 mM [26].

To evaluate the transport function of CLC-ec1 K131 mutants, we used quantitative ion flux

assays on purified transporters reconstituted into phospholipid vesicles [59] (**Fig. 5c**). We examined

both K131A (for which we have structural and ITC data) and K131M, which was previously studied

using planar lipid bilayer recordings[31]. We found that the unitary Cl⁻ turnover rate (Cl⁻ ions/s per

transporter) was greatly reduced in K131 mutant transporters compared to WT (**Fig. 5d,e** and **Table**

2). Their unitary H⁺ turnover rates were also greatly reduced, to levels just measurable above

background (**Fig. 5d,e** and **Table 2**). Importantly, we structured this flux-assay experiment to ensure

that any H⁺ movement into the vesicles was a result of Cl⁻-dependent H⁺ pumping rather than leakage

(**Fig. 5c**). Therefore, the low H⁺ transport rates measured for K131 mutant transporters represent

bona fide H⁺ pumping rates. By analyzing the ratio of the Cl⁻ and H⁺ transport rates, we determined

the Cl⁻/H⁺ transport stoichiometry (and thus the strength of coupling) for WT and mutant CLC-ec1

transporters. Strikingly, coupling in the K131 mutant transporters was largely intact, with a just a slight

increase in Cl⁻/H⁺ transport stoichiometry from ~2 for WT to ~3 for both K131 mutants (**Fig. 5f, Table**

2). This modest reduction in Cl⁻/H⁺ coupling efficiency is consistent with findings from K131M studied

in planar lipid bilayers, where shifts in reversal potentials suggested slight uncoupling [31]. Together,

these results illustrate that weak binding at S_{cen} does not on its own cause uncoupling.

The dramatic slowdown in transport rates by K131A and K131M mutant transporters (**Fig.**

5d,e) could be due directly to their weak Cl⁻ affinity, with Cl⁻ binding being rate limiting for transport.

Alternatively, it could be due to an indirect effect, with decreased Cl⁻ binding at S_{cen} causing the K131

transporters to stall at one or more of the H⁺-transport steps in the cycle. To distinguish these

possibilities, we evaluated the effect of K131 mutations in the background of the E_{gate} mutation,

E148A. CLC-ec1 E148A, lacking a titratable residue at the critical E_{gate} position, is completely unable

to transport H^+ and operates strictly as a passive Cl^- transporter [1]. If K131 mutations slow transport

because they weaken Cl^- binding, then they should affect Cl^- transport rates in both WT and E148A

backgrounds. On the other hand, if K131A mutations slow transport because they cause the

transporter to stall at one of the H^+-transport steps, then they should have no effect in the E148A

background, where there are no H^+-transport steps. We therefore tested CLC-ec1 K131A/E148A and

CLC-ec1 K131M/E148A mutant transporters using the quantitative Cl^- flux assay (**Fig. 5c**) (measuring

only Cl^- since E148 transporters do not transport H^+). K131 mutations had no discernable effect on

E148A Cl^- transport rates (**Fig. 5g, Table 2**). These results align with the "stalled transport cycle"

model for K131 mutants: when these low Cl^--binding mutant transporters are untethered from obligate

Cl^-/H^+ coupled transport, Cl^- movement becomes faster, unimpeded by the need to wait for a stalled

proton-transport step.

DISCUSSION

A comprehensive model to explain the CLC transporters' 2:1 Cl^-/H^+ exchange mechanism has

been elusive. This study was conducted with the hypothesis that investigating CLC conformational

dynamics across a broader expanded pH range – encompassing the physiological pH associated with

maximal activity – would yield new insights into the exchange mechanism.

Conformational Dynamics at pH 3.0

Previous spectroscopic and cryo-EM studies revealed that the CLC-ec1 transporter

undergoes conformational change at pH 4.5 in comparison to pH 7.5 and that this conformational

change is essential for activity [25,30]. Our HDX-MS results align with those previous results and reveal

additional conformational dynamics that occur at lower pH values, down to pH 3.0. Our cryo-EM

structure at pH 4.0 reveals a conformational change that opens access to the H^+ permeation pathway

(**Fig. 3g**), similar to that reported in cryo-EM studies of CLC-ec1 at pH 4.5 [25]. At pH 3.0, we observe

further conformational changes, particularly involving residues near the ion permeation pathways

(**Fig. 3h,j**).

In a recent cryo-EM study on CLC-ec1 by Fortea et al., a "Twist" conformation featuring a

notable 28.4-degree rotation between subunits was resolved at pH 4.5 [25]. Interestingly, we did not

observe this conformation in our pH 4.0 dataset. Our initial hypothesis was that the difference might

stem from the membrane mimetics used: we employed Salipro lipid nanodiscs, whereas Fortea et al.

used the detergent decyl maltoside (DM). To test this hypothesis, we determined a cryo-EM structure

under our original conditions (pH 4.0, 150 mM NaCl) but using DM instead of Salirpo nanodiscs.

However, we still did not detect the "Twist" conformation; thus, the difference in membrane

environment does not account for our lack of detection of the Twist conformation. Other differences

between our experimental preparation and that of Fortea et al. include pH (4.0 vs 4.5), NaCl

concentration (150 mM vs 100 mM), and buffer (citrate vs acetate). Given that the Twist conformation

is not essential for function (preventing its formation with cross-links has no effect on activity [25,28]), it

is perhaps not surprising that it is not detected under all conditions that support activity.

Conformational dynamics revealed by molecular dynamics simulations

Our molecular dynamics simulations on the pH 3.0 cryo-EM structure provided unprecedented

visualization of key steps in the Cl^-/H^+ exchange mechanism. With E_{gate} deprotonated, we dynamically

observed the conformational change of E_{gate} from the "out" to the "down" position. Remarkably, this

conformational change occurred in concert with the displacement of two Cl^- ions into the extracellular

solution, directly illustrating CLC-ec1's 2:1 exchange stoichiometry. Additionally, the accompanying

reorientation of inner-gate residues S107 and Y445 (**Movie 1, Fig. 4d**) was an unexpected

observation that provides further clarity on the exchange mechanism. This reorientation unveils a

long-awaited snapshot of the "inward-open" conformational state of CLC transporters.

Our simulations revealed water wires that connect the intracellular solution to the CLC-ec1

protein core. These wires occur both through the canonical H^+ pathway, as have been reported

previously[24,25,32-39] and through the Cl^- pathway (**Fig. 4a**). Surprisingly, these water wires occur

regardless of whether Cl^- is bound (**Fig. 4e**), contradicting prior simulations that indicated water wires

depend on Cl^- [32,37]. However, the prior simulations were conducted on a pH-7 structure now thought

to represent an inactive or poorly active conformation [25]. Additionally, the water wires observed in

prior simulations connected E_{gate} only to Glu_{in}, which is beneficial but non-essential for H^+ transport

[24], and not to the intracellular solution. Nonetheless, the simulation result is surprising because the

water wires that form in the absence of Cl^- appear to create a pathway for uncoupled transport. In

this scenario, H^+ transfer from the intracellular solution to E_{gate} could be followed by movement of

protonated E_{gate} to the "out" position, allowing a proton to be transferred to the extracellular solution

without concomitant Cl^- movement. However, the risk of uncoupled proton occurring in the Cl^--free

transporter may be alleviated if the protonation energy for E_{gate} is excessively high in the absence of

Cl^-. Indeed, free energy calculations conducted on CLC-ec1 in its low-activity state have predicted

this phenomenon, with experimental ion-binding studies supporting the computational prediction [27].

Therefore, the observed abundance of water wires, regardless of the presence of Cl^-, does not

inherently pose a risk to coupling.

The mechanism of reversible Cl^-/H^+ exchange

Our findings, combined with previous research, provide an unprecedented opportunity to propose a

complete and efficient CLC 2:1 Cl^-/H^+ exchange mechanism, free from steps that are energetically

unviable (**Fig. 6**). The transport mechanism involves conformational states identified in structural

studies— "up," "out," "down," and "middle"—alongside transient states observed in simulations. The

cycle, which can proceed in either direction, facilitates the transmembrane exchange of two Cl^- ions

for one H^+, with each step characterized by energetically plausible mechanisms. Beginning at the top

left with E_{gate} in the "up" conformation (state **A**) and moving through the cycle clockwise, protonation

from the extracellular solution enables the transition to the "out" conformation, with E_{gate} in the

hydrophobic core of the protein (state **B**). From there, water wires facilitate transfer of a proton to the

intracellular solution. (For simplicity of illustration, we show only one water wire, though multiple water

wires may form.) The resulting negatively charged E_{gate} moves from the hydrophobic core of the

protein into the anion-friendly pore (transient state **C**), expelling Cl^- to the outside, either two ions at

once, as observed in our simulations, or one at a time, as modeled here (states **D** and **E**). From the

single-site occupancy states (**D** and **E**), which have been observed crystallographically ("down" and

"middle"), the transient opening of the inner gate, as observed in our simulations, provides a pathway

for Cl^- to enter from the intracellular solution. From state **E**, Cl^- entry expels E_{gate} to the "up"

conformation, allowing it to be protonated from the extracellular side, thus initiating another transport

cycle. All steps in this mechanism are inherently reversible.

Notably, the newly identified water wire (Wire 1 in **Figure 4**, illustrated here in states **C** and

D), which connects the intracellular solution to E_{gate} positioned in the "down" conformation, offers a

physically plausible pathway for the transport cycle to proceed in the counterclockwise direction, with

Cl^- ions moving inward and H^+ ions moving outward. Starting at the top left in state **A** and moving

counterclockwise, the negatively charged E_{gate} moves into the Cl^- pathway, displacing a Cl^- ion to the

intracellular side and generating state **E**. Subsequent entry of a Cl^- ion from the extracellular side

knocks E_{gate} to the "down" position and displaces a second Cl^- ion, generating state **D.** Water wires

can connect to E_{gate} in this conformation, but protonation of E_{gate} is not energetically favorable at this

stage, with only 1 Cl^- ion in the pathway. However, when a second Cl^- enters (state **C**), protonation

of E_{gate} becomes energetically favorable. Following protonation, the neutralized E_{gate} is favored

towards the "out" conformation (state **B**). A subsequent shift to the "up" conformation (state **A**) enables

release of a proton to the extracellular solution, thereby completing the transport cycle in that direction.

K131 is universally conserved across the CLC family, including both transporter and channel

homologs. This conservation, combined with K131's location - entirely buried within the

transmembrane domain, which is rare for lysine residues [60] – suggests that it plays a crucial functional

role. Indeed, mutations at this position in CLC channels shift voltage dependence and lead to

myotonia [61-64]. In CLC-ec1, we found that K131 mutations weaken Cl^- binding affinity and reduce

Cl^-/H^+ transport rates (**Fig. 5, Table 2**). The exchange mechanism illustrated in **Fig. 6a** provides a

framework for understanding how this weakened Cl^- binding results in reduced Cl^-/H^+ transport rates.

In this context, the impact of reduced Cl^- binding at S_{ext} and S_{cen} must be considered relative to E_{gate}

binding at these sites. Although E_{gate} binding affinity may also be weakened in the mutants, its physical

tethering gives it a competitive advantage over free Cl^- ions. Supporting this idea, the K131A structure

shows that the cryo-EM density for E_{gate} is less affected compared to that of Cl^-. Consequently, Cl^-

will struggle to compete with E_{gate} for the S_{ext} and S_{cen} sites, thereby slowing the $D \rightleftarrows E$, $D \rightarrow C$, and

$E \rightarrow A$ transitions and stalling the transport cycle.

In addition to reducing Cl^- binding and transport rates by more than tenfold, K131 mutations

cause mild uncoupling, increasing the Cl^-/H^+ stoichiometry from 2 to ~3. A potential mechanism for

this experimentally observed uncoupling may involve off-cycle transitions through states with

unoccupied anion-binding sites, as illustrated in **Fig. 6b**. During these off-cycle transitions, weakened

Cl^- binding allows Cl^- to exit the permeation pathway independently, without being displaced by

incoming Cl^- from the opposite side. As a result, these off-cycle transitions facilitate the transport of

a Cl^- ion independently of H^+ transport. Within this context, Cl^- movement to and from the interior is

likely facilitated by a slight weakening of the inner gate, due to the loss of hydrogen bonds between

K131 and the conserved inner-gate loop residues A104 and G106. Nevertheless, since the

uncoupling is mild, these off-cycle events do not predominate but rather occur approximately once

per transport cycle.

The proposed transport mechanism (**Fig. 6a**) is relevant to CLC transporters across all

kingdoms, which share fundamental features including 2:1 anion/proton coupling and E_{gate} as an

essential coupling element [4]. The mechanism elucidates how transport occurs in the direction of Cl^-

ions moving inward across the membrane and H^+ ions moving outward, which is the direction of

physiological relevance for CLC-ec1 in facilitating extreme acid tolerance [15] and for mammalian CLCs

in acidifying intracellular compartments [22,65]. While this mechanism addresses the fundamental Cl^-/H^+

exchange function, questions remain concerning how this core function is regulated in the various

CLC transporters. For example, mammalian CLCs contain large cytoplasmic domains that mediate

regulation by signaling lipids [65-67] and by nucleotides [68-71], but these regulatory mechanisms are only

partially understood. Many prokaryotic homologs also feature such domains, although their functional

roles have yet to be investigated. Gaining insight into how these regulatory mechanisms interact with

the core Cl^-/H^+ exchange process will be crucial for fully understanding the diverse functions of CLC

transporters in cellular physiology.

METHODS

Mutagenesis

Mutations were introduced into the WT CLC-ec1 gene in pASK [17] using New England BioLabs

Q5 Site Directed Mutagenesis kit (New England Biolabs, Ipswich, MA, USA). Each mutation was

subsequently verified with whole plasmid sequencing and analyzed using SnapGene® software 2023

(from Dotmatics; available at snapgene.com).

Protein Production and Purification

CLC ec-1 with an N-terminal polyhistidine tag, in the pASK90 expression vector [72], was

transformed into BL21 *E. coli* cells and plated onto LB plates containing 100 µg/mL ampicillin. The

following morning, the colonies from 1-2 plates were scraped into 10 mL LB, added to 1 L of Terrific

Broth [73] in a 2.8-L baffled flask, and cultured at 37°C with vigorous shaking (~220 RPM). When the

culture reached 1.0 OD_{600}, it was induced with 0.2 mg/L anhydrotetracycline (added from a 0.2 mg/mL

solution in dimethylformamide). The cultures were grown for an additional 3 hours then pelleted at

2246*xg* for 20 minutes and flash-frozen in liquid nitrogen for future purification.

For purification, CLC-ec1 pellets were extracted with 50 mM n-Decyl-α-D-Maltopyranoside

(DM**)** (Anatrace, sol grade) for two hours then centrifuged at 25,759*xg* for 45 minutes. The protein

was then isolated using a cobalt affinity chromatography step (3 mL of 50% cobalt metal affinity resin

slurry, Takara Bio USA, Inc. San Jose, CAI for each 1L prep), eluting with cobalt wash buffer (200

mM NaCl, 20 mM DM (sol grade), 400mM Imidazole, 40 mM TrisCl, pH 7). The eluate was incubated

with Endoproteinase Lys-C (Santa Cruz Biotechnology, Inc.), 0.17 U per L culture. The final

purification by size exclusion chromatography (SEC) (Enrich SEC 650 10 X 300 column, Bio Rad

Laboratories Pleasanton, CA) was carried out in SEC Buffer **(**10 mM NaHepes, 150 mM NaCl, 5 mM

DM (anagrade), pH 7.5). Purified protein was concentrated using Amicon® Ultra-15 centrifugal filters

with a 50-kD cut-off.

For single-particle cryo-EM experiments at pH 7.5 and 4.0, CLC-ec1 purified by SEC as above,

was reconstituted into Saposin-lipoprotein nanoparticle (Salipro

(https://doi.org/10.1038/nmeth.3801)) at either pH 7.5 and pH 4. We followed the reconstitution

protocol described in Chien et al. [74] with some modifications. For the pH 7.5 Salipro sample

preparation, 0.5 mg of CLC-ec1 in DM at pH 7.5 was mixed with 1.35 mM of POPE, 450 µM of POPG,

0.2 % (w/v) of DDM, 120 µM of Saposin-A, and Gel-filtration buffer (GFB) (10 mM Hepes, 150 mM

NaCl, pH 7.5) to a total volume of 2.5 mL. The mixture was incubated at 4 °C for 45 min. Then 50 %

(w/v) of Amberlite XAD-2 (Millipore Sigma) was added to the mixture followed by 15 min incubation

at 4 °C. The Amberlite XAD-2 beads were removed by centrifugation. The mixture was concentrated

to 200 µL using a 10 kDa cutoff Amicon (Millipore Sigma). A final SEC step was carried out using

Superdex 200 Increase (10/300) (Cytiva) equilibrated in GFB. The fractions corresponding to CLC-

ec1 in Salipro were collected and concentrated using a 3-kDa cutoff Amicon concentrator (Millipore

Sigma) to a final concentration of 2.2 mg/mL. For the pH 4.0 Salipro sample, 1.35 mM of POPE, 450

µM of POPG, 0.2 % (w/v) of DDM, 120 µM of Saposin-A and the pH 4 buffer (1.8 mM Citric Acid, 8.2

mM Sodium Citrate, 150 mM NaCl, pH 4) were mixed first and incubated at 37 °C for 10 min. Then

1.5 mg of CLC-ec1 in DM at pH 7.5 was added to the mixture making a total volume of 5 mL. The

mixture was incubated at 37 °C for 5 min. The rest of the steps including Amberlite XAD-2 beads and

gel-filtration are the same as for the Salipro pH 7.5 sample.

For cryo-EM of WT CLC-ec1 at pH 3, protein was generated using the method as described

above but with modification to generate a final sample in the detergent lauryl maltose neopentyl glycol

(LMNG). During the cobalt-affinity chromatography step, DM gradually replaced with LMNG

(Anatrace), to a final concentration of 20 µM LMNG. The exchange was as follows 50:50, 25:75,

10:90, and finally 5:95 percent by volume DM to LMNG. The final SEC step was performed in pH 3

buffer, 8.2mM Citric Acid, 1.8mM sodium Citrate, 150mM NaCl and 20 µM LMNG. For cryo-EM of

K131A, the LMNG protocol was used as described above, except the pH in the final SEC buffer was

7.5 (10 mM NaHepes, 150 mM NaCl, 20 µM LMNG, pH 7.5).

Hydrogen/deuterium exchange mass spectrometry

For HDX experiments, CLC-ec1 purification was done using the standard protocol with a final

SEC step in 7.5 mM HEPES, 100 mM NaCl, 1.8 mM DM (anagrade), pH 7.5. This stock was then

buffer exchanged to H$_2$O buffer of different pHs, incubated afterwards at least 1hr and then used to

run HDX.

For HDX, McIlavaine buffers of the desired pH were prepared. Solution A (0.2 M Na$_2$HPO$_4$)

and solution B (0.1 M citric acid) were mixed in the recommended proportions for each pH (pH 3.0 –

A 206 µl, B 795 µl; pH 3.5 – A 304 µl, B 697 µl; pH 4.0 – A 386 µl, B 615 µl; pH 4.5 – A 454 µl, B 546

µl; pH 6.5 – A 710 µl, B 290 µl). Buffers were diluted 5-fold into water, and then 100 mM NaCl and

1.8 mM DM were added. Buffers were prepared in both H$_2$O and D$_2$O. The protein was transferred to

McIlvaine buffers of pH 6.5, 4.5, 4.0, 3.5, and 3.0 using Zeba Spin 7k MWCO desalting columns

(Thermo Scientific).

The pH dependent changes in CLC-ec1 were followed on a detergent solubilized protein at

pH 6.5, 4.5, 4.0, 3.5, 3.0. CLC-ec1 in McIlvaine buffer was diluted 5-fold into a D$_2$O-based buffer of

identical composition and pD. Protein concentration during exchange was 3.2µM and temperature 21

°C. Aliquots were taken at 20s, 63s, 200s, 633s, 2000s, 6325s, 20000s, 63000s, where 20s, 633s,

20000s were triplicated. The exchange was stopped by adding phosphoric acid (mixing ratio 1:1) and

freezing in liquid nitrogen. Different phosphoric acid concentrations matching the exchange buffer's

pH and containing 1.8 mM DM were used (pH 6.5 – 60mM, pH 4.5 – 40mM, pH 4.0 – 35mM, pH 3.5

– 30mM, 3.0 – 25mM). Fully deuterated control was prepared to correct for deuterium loss during the

analysis.

Each sample was quickly thawed and injected onto a protease column (bed volume 70µl)

containing immobilized pepsin and nepenthesin-2 [75]. Digestion was done at 19 °C and was driven by

a flow of 0.4% formic acid in water delivered by an Agilent 1260 Infinity II Quaternary pump (Agilent

Technologies, Waldbronn, Germany) at 200 µL.min-1. Peptides were trapped on a guard column

(SecurityGuard™ ULTRA Cartridge UHPLC Fully Porous Polar C18, 2.1mm ID, Phenomenex,

Torrance, CA) and desalted. Next, separation on an analytical column (Luna Omega Polar C18, 1.6

µm, 100 Å, 1.0x100 mm, Phenomenex, Torrance, CA) was done by an acetonitrile gradient (10%–

45%; solvent A: 0.1% FA in water, solvent B: 0.1%FA, 2% water in ACN). Solvents were delivered by

the Agilent 1290 Infinity II LC system pumping at 50 µL.min-1. To minimize carry-over and remove

the detergent, the protease column was washed by two consecutive injections of 4M urea, 500mM

glycine HCl pH 2.3 and 5% acetonitrile, 5% isopropanol, 20% acetic acid in water [76]. The analytical

column, together with the guard column, was washed with solvent B for 4 min and then with 70%

methanol, 20% isopropanol, and 1% formic acid in water for 4 min. The LC setup was cooled to 2°C

to minimize the deuterium loss. The analytical column outlet was directly connected to an ESI source

of a timsTOF Pro (Bruker Daltonics, Bremen, Germany) operating in MS mode with 1Hz data

acquisition. The data processing started with peak picking in Data Analysis 5.3 and exported to simple

text files, which were then further handled in DeutEx. [77]. Data visualization was done using MSTools

[78] and PyMol. Identification of peptides arising from pepsin/nepenthesin-2 digestion was done using

the same LCMS system, but the mass spectrometer operated in data-dependent MSMS mode with

PASEF enabled. The data were searched using MASCOT (v. 2.7, Matrix Science, London, UK)

against a database containing sequences of CLC-ec1, acid proteases, and cRAP.fasta

(https://www.thegpm.org/crap/). The search parameters were: precursor tolerance 10 ppm, fragment

ion tolerance 0.05 Da, decoy search enabled with FDR <1%, IonScore > 20, and peptide length >5.

The mass spectrometry data have been deposited to the ProteomeXchange Consortium via the

PRIDE partner repository with the dataset identifier PXD058693.

For the stability control experiment, purified CLC-ec1 in 20 mM HEPES (pH 7.5), 100 mM

NaCl, and 1.8 mM DM was transferred to McIlvaine buffer (pH 3.0) containing 100 mM NaCl and 1.8

mM DM using Zeba Spin 7K MWCO desalting columns (Thermo Scientific). After buffer exchange,

the protein was incubated for 0, 1, 4, and 8 hours before separation on a size-exclusion

chromatography (SEC) column (Enrich™ SEC 650, 10 × 300 mm, 24 mL; Bio-Rad). ClC-ec1

maintained at pH 7.5 served as a control. SEC separation was performed using a 1 mL sample

containing 1 mg of protein at a flow rate of 1 mL/min. The elution buffer consisted of 20 mM HEPES

(pH 7.5), 100 mM NaCl, and 1.8 mM DM. All samples were analyzed under identical conditions.

Cryo-EM sample preparation, data collection, and image processing

Purified samples with a concentration of 2-4 mg/mL were used for cryo-EM grid preparation.

1 mM fluorinated FOS-choline-8 was added prior to freezing grids to improve ice quality and avoid

preferred orientation due to air-water interface interactions. 3 μL of the sample was applied to glow

discharged Quantifoil R1.2/1.3 Cu200 grids, then blotted with a Whatman filter paper for 3 s before

plunge-frozen in liquid ethane using a Vitrobot Mark IV (Thermo Fisher Scientific) at 4 °C and 100%

humidity.

All cryo-EM data were collected with Thermo Fisher Scientific Titan-Krios cryo-electron

microscopes operating at 300 keV. The WT pH 7.5 dataset was collected with a Falcon 4 camera

without an energy filter. The WT pH 4 dataset and K131A dataset were collected with a Gatan K3

camera and Bio-quantum energy filter set to 20 eV. The WT pH 3 dataset was collected with a Falcon

4 and Selectris energy filter set to 10 eV. Data collection parameters are reported in Table 1.

Automated data collection was done using EPU software (Thermo Fisher Scientific).

The complete data processing workflows with specific details for each sample/dataset are

reported in **Extended Data Figs. 4-6**. The overall data processing strategy is similar for all four

datasets (WT at pH 7.5, 4, and 3, and K131 at pH 7.5). The raw movies were pre-processed (motion

correction and CTF correction) in CryoSPARC live. Micrographs were curated with defined criteria

including CTF fit and relative ice thickness. A blob picker or a template picker was used to pick a

small subset of the dataset. The picked particles were pruned by 2D classifications, ab-initio

reconstructions, and heterogeneous refinements in cryoSPARC. The good particles that could be

reconstructed to a map with recognizable protein features were used to train a Topaz particle picker.

This Topaz picker was used to pick the full dataset. Several rounds of ab-initio reconstructions and

heterogeneous refinements with 3 classes were used to clean the particle stacks. For WT pH 7.5 and

pH 3 datasets, 3D classification without alignment in Relion with a mask that covers only the proteins

was also used for particle cleanup. The final non-uniform refinement with C2 symmetry imposed was

done in cryoSPARC.

Model fitting and refinement

A CLC-ec1 crystal structure (PDB: 1OTS) was used as an initial atomic model for WT CLC-ec

at pH 7.5 and pH 4 and for K131A. The QQQ crystal structure (PDB: 6V2J) was used for pH 3 model

building. The initial models were first modified based on our construct (i.e. K131A or WT sequences).

These models were rigid-body docked into the cryo-EM map and refined using ISOLDE

(10.1107/S2059798318002425) and PHENIX (10.1107/S2059798318006551). All atomic models

were validated in PHENIX validation job and Q-score [55]. All structure figures were prepared with UCSF

ChimeraX [79].

System setup for molecular dynamics simulations

We performed simulations of CLC-ec1 under two conditions: (1) simulations with E148 (Glu_{ex}),

E203 (Glu_{in}), and E113 protonated; (2) simulations with E148 (Glu_{ex}) deprotonated, and E203 (Glu_{in})

and E113 protonated. We initiated all simulations from a CLC-ec1 structure based on the cryo-EM

data reported in this manuscript (specifically, from a model very similar to that presented here but

based on an earlier refinement). The 3 Cl^- ions bound at S_{cen}, S_{ext}, and S_{int} anion-binding sites in each

of the two subunits were preserved. For each simulation condition, we performed ten independent

simulations, each 1.5 μs in length. For each simulation, initial atom velocities were assigned randomly

and independently.

For all simulation conditions, the protein structure was aligned to the Orientations of Proteins

in Membranes [80] entry for 6V2J (CLC-ec1 triple mutant (E113Q, E148Q, E203Q) [24] using PyMOL,

and crystal waters from 6V2J were incorporated. Prime (Schrödinger) [81] was used to add capping

groups to protein chain termini. Protonation states of all titratable residues except for E148 (Glu_{ex}),

E203 (Glu_{in}) and E113 were assigned at pH 4.5, which is the pH at which activity is most often

measured (eg. **Fig. 5**). Using Dabble [82], the prepared protein structures were inserted into a pre-

equilibrated palmitoyl-oleoyl-phosphatidylcholine (POPC) bilayer, the system was solvated, and

sodium and chloride ions were added to neutralize the system and to obtain a final concentration of

150 mM. The final systems comprised approximately 123,000 atoms, and system dimensions were

approximately 160x120x90 Å.

Molecular dynamics simulation and analysis protocols

We used the CHARMM36m force field for proteins, the CHARMM36 force field for lipids and

ions, and the TIP3P model for waters [83-85]. All simulations were performed using the Compute Unified

Device Architecture (CUDA) version of particle-mesh Ewald molecular dynamics (PMEMD) in

AMBER20 on graphics processing units (GPUs).

Systems were first minimized using three rounds of minimization, each consisting of 500 cycles of

steepest descent followed by 500 cycles of conjugate gradient optimization. 10.0 and 5.0

$kcal \cdot mol^{-1} \cdot Å^{-2}$ harmonic restraints were applied to the protein and lipids for the first and second rounds

of minimization, respectively. 1 $kcal \cdot mol^{-1} \cdot Å^{-2}$ harmonic restraints were applied to the protein for the

third round of minimization. Systems were then heated from 0 K to 100 K in the NVT ensemble over

12.5 ps and then from 100 K to 310 K in the NPT ensemble over 125 ps, using 10.0 $kcal \cdot mol^{-1} \cdot Å^{-2}$

harmonic restraints applied to protein heavy atoms. Subsequently, systems were equilibrated at 310

K and 1 bar in the NPT ensemble, with harmonic restraints on the protein non-hydrogen atoms

tapered off by 1.0 $kcal \cdot mol^{-1} \cdot Å^{-2}$ starting at 5.0 $kcal \cdot mol^{-1} \cdot Å^{-2}$ in a stepwise fashion every 2 ns for 10

ns, and then by 0.1 $kcal \cdot mol^{-1} \cdot Å^{-2}$ every 2 ns for 20 ns. Production simulations were performed without

restraints at 310 K and 1 bar in the NPT ensemble using the Langevin thermostat and the Monte

Carlo barostat, and using a timestep of 4.0 fs with hydrogen mass repartitioning [86]. Bond lengths were

constrained using the SHAKE algorithm [87]. Non-bonded interactions were cut off at 9.0 Å, and long-

range electrostatic interactions were calculated using the particle-mesh Ewald (PME) method with an

Ewald coefficient of approximately 0.31 $Å^{-1}$, and 4th order B-splines. The PME grid size was chosen

such that the width of a grid cell was approximately 1 Å. Trajectory frames were saved every 200 ps

during the production simulations. The AmberTools17 CPPTRAJ package was used to reimage

trajectories [88]. Simulations were visualized and analyzed using Visual Molecular Dynamics (VMD) [89]

and PyMOL [90].

Isothermal Titration Calorimetry

CLC-ec1 WT and K131A protein purification was carried out as described above for WT CLC-

ec1 at pH 7.5 in DM, with the following differences. First, to obtain quantities CLC ec-1 needed for

ITC experiments (2 mL at ~15 µM), we used cells from 2-3 L of pellets. Second, during the final SEC

step, we eluted the protein with Cl$^-$-free buffer (buffer A: 10 mM HEPES, 150 mM Na-isethionate, 5

mM anagrade DM, pH 7.5). The protein was then dialyzed at 4℃ against the same buffer (250 mL)

with buffer exchange every three hours for a total of four exchanges to remove trace Cl⁻. Following

dialysis, 2 mL of the protein sample was added to the sample chamber of the VP-ITC microcalorimeter

(Malverne Panalytical, Malvern, United Kingdom). Titrant (30 mM KCl in buffer A) was injected into

the sample chamber 10 µL at a time, once per minute for a total 180 minutes. Reference data were

obtained by titrating buffer A into the protein-containing solution. The data were analyzed using

MicroCal ITC-ORIGIN Analysis Software, with fitting using the 'one set of sites' model (keeping n =

1).

Reconstitution and transport assays

WT and mutant CLC-ec1 were reconstituted into liposomes as described elsewhere [24,59,91]. In

brief, 25 mg/ml E.coli polar lipid extract (Avanti Polar Lipids, Alabaster, Al, USA) were evaporated

under vacuum, washed with pentane, and then resuspended in reconstitution buffer (300 mM KCl, 40

mM citric acid, pH = 4.3 with NaOH and 21.5 mg/mL CHAPS (3-[(3-

Cholamidopropyl)dimethylammonio]-1-propanesulfonate, Biotium, Freemont, CA, USA)) to a final

concentration of 20 mg/ml. CLC-ec1 protein (1-2 mg/mL in DM) was then added to achieve a

protein:lipid ratio of 0.4 µg/mg and a lipid concentration between 19.5 and 19.8 mg/ml.

Proteoliposomes were then aliquoted into 200-µL reconstitution samples and dialyzed over two days

with 4 x 1 L reconstitution buffer. Proteoliposomes were subject to four freeze-thaw cycles and then

used directly for transport assays or stored at -80°C until measured.

Flux assays were performed as described [24]. Briefly, 60 µL of reconstituted sample was buffer-

exchanged using a Sephadex spin column, then diluted into 600 µL pH-4.5 flux assay buffer (300 mM

K-isethionate, 2 mM citric acid and 50 µM KCl, pH 4.5). Each 200-µL reconstitution sample was used

for a maximum of three flux-assay measurements (60 µL each) and the results averaged to yield a

single data point (see Source Data). For each flux-assay experimental condition, reconstitutions from

at least two independent protein preparations were used. Chloride and proton transport were

measured in parallel using Ag/AgCl electrodes and a micro pH electrode. Data were acquired using

AxosScope 9.0 (Axon Instruments, San Jose, CA, USA) and processed using Clampfit 9.0 (Axon

Instruments) and Python routines. Data and transport rates shown in the manuscript were visualized

in SigmaPlot 14 (Systat Software Inc., San Jose, CA, USA).

DATA AVAILABILILTY

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium

via the PRIDE [1] partner repository with the dataset identifier PXD058693" where the reference is

PubMed ID: 34723319. The DNA constructs used in this study are available from M.M. upon request.

Models and associated cryo-EM maps have been deposited into the Protein Data Bank (PDB) and

the Electron Microscopy Data Bank (EMDB) under the following accession codes: CLC-ec1 pH 7.5

(PDB ID 9095; EMD-70242); CLC-ec1 pH 4.0 (PDB ID 9096; EMD-70243); CLC-ec1 pH 3.0 (PDB ID

9097; EMD-70244); CLC-ec1 K131A pH 7.5 (PDB ID 9098; EMD-70245). The MD simulation data

that support this study are available from R.O.D. upon request.

ACKNOWLEDGMENTS

We thank Anindita Das and Shwetha Srinivasan for comments on the manuscript. This research was

funded by NIH R01GM113195 (MM, ROD, and WC). For the mass spectrometry aspects of this

research, financial support from MEYS/EU project OP JAK – Photomachines

(CZ.02.01.01/00/22_008/0004624) is gratefully acknowledged. Access to the Centre of Molecular

Structure (CMS), BioCeV, was supported by CIISB LM2023042 and ERDF "UP CIISB"

(CZ.02.1.01/0.0/0.0/18_046/0015974). Cryo-EM was performed at the Stanford-SLAC Cryo-EM

Center (S2C2), which is supported by the National Institutes of Health Common Fund Transformative

High-Resolution Cryo-Electron Microscopy program (U24 GM129541). A.N was supported by an

administrative supplement to NIH R01GM113195 and by a Stanford Propel Postdoctoral Fellowship.

AUTHOR CONTRIBUTIONS

D.A. designed, performed, and analyzed the MD simulations, wrote methods; C.C. collected and

analyzed cryo-EM data, built and refined experimental models, modeled transport cycle, wrote

methods; J.K designed, performed, and analyzed flux-assay experiments, modeled transport cycle,

wrote methods; A.N. prepared and purified protein samples, designed, performed, and analyzed

molecular biology and ITC experiments, wrote methods; J.M.P. prepared and purified protein

samples, performed and analyzed mass spectrometry experiments, wrote methods; L.F. prepared

and purified protein samples, performed and analyzed mass spectrometry experiments, wrote

methods; C.M.S. prepared and purified protein samples; P.M., R.O.D., W.C., and M.M. oversaw the

research. M.M. wrote the initial draft of the paper. All authors edited and approved the manuscript.

REFERENCES:

1. Accardi, A. & Miller, C. Secondary active transport mediated by a prokaryotic homologue of ClC Cl-channels. *Nature* **427**, 803-7 (2004).

2. Miller, C. ClC chloride channels viewed through a transporter lens. *Nature* **440**, 484-9 (2006).

3. Lisal, J. & Maduke, M. The ClC-0 chloride channel is a 'broken' Cl-/H+ antiporter. *Nat Struct Mol Biol* **15**, 805-10 (2008).

4. Jentsch, T.J. & Pusch, M. CLC Chloride Channels and Transporters: Structure, Function, Physiology, and Disease. *Physiol Rev* **98**, 1493-1590 (2018).

5. Koster, A.K. & Maduke, M. CLC Chloride Channels and Transporters. in *Textbook of Ion Channels* Vol. II Properties, Function, and Pharmacology of the Superfamilies (eds. Zheng, J. & Trudeau, M.C.) 193-208 (CRC Press, Boca Raton, FL, 2023).

6. Bose, S., He, H. & Stauber, T. Neurodegeneration Upon Dysfunction of Endosomal/Lysosomal CLC Chloride Transporters. *Front Cell Dev Biol* **9**, 639231 (2021).

7. Duncan, A.R. et al. Unique variants in CLCN3, encoding an endosomal anion/proton exchanger, underlie a spectrum of neurodevelopmental disorders. *Am J Hum Genet* **108**, 1450-1465 (2021).

8. Zifarelli, G. The Role of the Lysosomal Cl(-)/H(+) Antiporter ClC-7 in Osteopetrosis and Neurodegeneration. *Cells* **11**(2022).

9. Coppola, M.A. et al. Biophysical Aspects of Neurodegenerative and Neurodevelopmental Disorders Involving Endo-/Lysosomal CLC Cl(-)/H(+) Antiporters. *Life (Basel)* **13**(2023).

10. Palmer, E.E. et al. Functional and clinical studies reveal pathophysiological complexity of CLCN4-related neurodevelopmental condition. *Mol Psychiatry* **28**, 668-697 (2023).

11. Dieguez, L. et al. Dent's Disease: A Cause of Monogenic Kidney Stones and Nephrocalcinosis. *J Pers Med* **14**(2024).

12. Iyer, H. & Talbot, W.S. The Cl- transporter ClC-7 is essential for phagocytic clearance by microglia. *J Cell Sci* **137**(2024).

13. McIlwain, B.C., Ruprecht, M.T. & Stockbridge, R.B. Membrane Exporters of Fluoride Ion. *Annu Rev Biochem* **90**, 559-579 (2021).

14. Liu, L., Li, X., Wang, C., Ni, Y. & Liu, X. The Role of Chloride Channels in Plant Responses to NaCl. *Int J Mol Sci* **25**(2023).

15. Iyer, R., Iverson, T.M., Accardi, A. & Miller, C. A biological role for prokaryotic ClC chloride channels. *Nature* **419**, 715-8 (2002).

16. Kim, M., Choi, N., Choi, E. & Lee, E.J. ClC Chloride Channels in Gram-Negative Bacteria and Its Role in the Acid Resistance Systems. *J Microbiol Biotechnol* **33**, 857-863 (2023).

17. Maduke, M., Pheasant, D.J. & Miller, C. High-level expression, functional reconstitution, and quaternary structure of a prokaryotic ClC-type chloride channel. *J Gen Physiol* **114**, 713-22 (1999).

18. Mindell, J.A., Maduke, M., Miller, C. & Grigorieff, N. Projection structure of a ClC-type chloride channel at 6.5 A resolution. *Nature* **409**, 219-23 (2001).

19. Dutzler, R., Campbell, E.B., Cadene, M., Chait, B.T. & MacKinnon, R. X-ray structure of a ClC chloride channel at 3.0 A reveals the molecular basis of anion selectivity. *Nature* **415**, 287-94 (2002).

20. Dutzler, R. A structural perspective on ClC channel and transporter function. *FEBS Lett* **581**, 2839-44 (2007).

21. Accardi, A. Structure and gating of CLC channels and exchangers. *J Physiol* **593**, 4129-38 (2015).

22. Picollo, A. Vesicular CLC chloride/proton exchangers in health and diseases. *Front Pharmacol* **14**, 1295068 (2023).

23. Dutzler, R., Campbell, E.B. & MacKinnon, R. Gating the selectivity filter in ClC chloride channels. *Science* **300**, 108-12 (2003).

24. Chavan, T.S. et al. A CLC-ec1 mutant reveals global conformational change and suggests a unifying mechanism for the CLC Cl(-)/H(+) transport cycle. *Elife* **9**(2020).

25. Fortea, E. et al. Structural basis of pH-dependent activation in a CLC transporter. *Nat Struct Mol Biol* **31**, 644-656 (2024).

26. Picollo, A., Malvezzi, M., Houtman, J.C. & Accardi, A. Basis of substrate binding and conservation of selectivity in the CLC family of channels and transporters. *Nat Struct Mol Biol* **16**, 1294-301 (2009).

27. Picollo, A., Xu, Y., Johner, N., Berneche, S. & Accardi, A. Synergistic substrate binding determines the stoichiometry of transport of a prokaryotic H(+)/Cl(-) exchanger. *Nat Struct Mol Biol* **19**, 525-31, S1 (2012).

28. Nguitragool, W. & Miller, C. CLC Cl /H+ transporters constrained by covalent cross-linking. *Proc Natl Acad Sci U S A* **104**, 20659-65 (2007).

29. Chadda, R. & Robertson, J.L. Measuring Membrane Protein Dimerization Equilibrium in Lipid Bilayers by Single-Molecule Fluorescence Microscopy. *Methods Enzymol* **581**, 53-82 (2016).

30. Khantwal, C.M. et al. Revealing an outward-facing open conformational state in a CLC Cl(-)/H(+) exchange transporter. *Elife* **5**(2016).

31. Accardi, A. et al. Separate ion pathways in a Cl-/H+ exchanger. *J Gen Physiol* **126**, 563-70 (2005).

32. Han, W., Cheng, R.C., Maduke, M.C. & Tajkhorshid, E. Water access points and hydration pathways in CLC H+/Cl- transporters. *Proc Natl Acad Sci U S A* **111**, 1819-24 (2014).

33. Kuang, Z., Mahankali, U. & Beck, T.L. Proton pathways and H+/Cl- stoichiometry in bacterial chloride transporters. *Proteins* **68**, 26-33 (2007).

34. Wang, D. & Voth, G.A. Proton transport pathway in the ClC Cl-/H+ antiporter. *Biophys J* **97**, 121-31 (2009).

35. Ko, Y.J. & Jo, W.H. Secondary water pore formation for proton transport in a ClC exchanger revealed by an atomistic molecular-dynamics simulation. *Biophysical Journal* **98**, 2163-9 (2010).

36. Kieseritzky, G. & Knapp, E.W. Charge transport in the ClC-type chloride-proton anti-porter from Escherichia coli. *Journal of Biological Chemistry* **286**, 2976-86 (2011).

37. Jiang, T., Han, W., Maduke, M. & Tajkhorshid, E. Molecular Basis for Differential Anion Binding and Proton Coupling in the Cl(-)/H(+) Exchanger ClC-ec1. *J Am Chem Soc* **138**, 3066-75 (2016).

38. Leisle, L. et al. Divergent Cl(-) and H(+) pathways underlie transport coupling and gating in CLC exchangers and channels. *Elife* **9**(2020).

39. Wang, Z., Swanson, J.M.J. & Voth, G.A. Local conformational dynamics regulating transport properties of a Cl(-) /H(+) antiporter. *J Comput Chem* **41**, 513-519 (2020).

40. Accardi, A., Lobet, S., Williams, C., Miller, C. & Dutzler, R. Synergism between halide binding and proton transport in a CLC-type exchanger. *J Mol Biol* **362**, 691-9 (2006).

41. Nguitragool, W. & Miller, C. Uncoupling of a CLC Cl-/H+ exchange transporter by polyatomic anions. *J Mol Biol* **362**, 682-90 (2006).

42. Miller, C. Q-cubed mutant cues clues to CLC antiport mechanism. *J Gen Physiol* **153**(2021).

43. Matulef, K. & Maduke, M. Side-dependent inhibition of a prokaryotic ClC by DIDS. *Biophys J* **89**, 1721-30 (2005).

44. Bosshart, P.D. & Fotiadis, D. Secondary Active Transporters. *Subcell Biochem* **92**, 275-299 (2019).

45. Garaeva, A.A. & Slotboom, D.J. Elevator-type mechanisms of membrane transport. *Biochem Soc Trans* **48**, 1227-1241 (2020).

46. Licht, J.A., Berry, S.P., Gutierrez, M.A. & Gaudet, R. They all rock: A systematic comparison of conformational movements in LeuT-fold transporters. *Structure* **32**, 1528-1543 e3 (2024).

47. Bell, S.P., Curran, P.K., Choi, S. & Mindell, J.A. Site-directed fluorescence studies of a prokaryotic ClC antiporter. *Biochemistry* **45**, 6773-82 (2006).

48. Elvington, S.M. & Maduke, M. Thinking outside the crystal: complementary approaches for examining transporter conformational change. *Channels (Austin)* **2**, 373-9 (2008).

49. Elvington, S.M., Liu, C.W. & Maduke, M.C. Substrate-driven conformational changes in ClC-ec1 observed by fluorine NMR. *EMBO J* **28**, 3090-102 (2009).

50. Abraham, S.J. et al. 13C NMR detects conformational change in the 100-kD membrane transporter ClC-ec1. *J Biomol NMR* **61**, 209-26 (2015).

51. Heath, G.R. et al. Localization atomic force microscopy. *Nature* **594**, 385-390 (2021).

52. Accardi, A., Kolmakova-Partensky, L., Williams, C. & Miller, C. Ionic currents mediated by a prokaryotic homologue of CLC Cl- channels. *J Gen Physiol* **123**, 109-19 (2004).

53. Coppieters 't Wallant, K. & Martens, C. Hydrogen-deuterium exchange coupled to mass spectrometry: A multifaceted tool to decipher the molecular mechanism of transporters. *Biochimie* **205**, 95-101 (2023).

54. Ozohanics, O. & Ambrus, A. Hydrogen-Deuterium Exchange Mass Spectrometry: A Novel Structural Biology Approach to Structure, Dynamics and Interactions of Proteins and Their Complexes. *Life (Basel)* **10**(2020).

55. Pintilie, G. et al. Measurement of atom resolvability in cryo-EM maps with Q-scores. *Nat Methods* **17**, 328-334 (2020).

56. Wraight, C.A. Chance and design--proton transfer in water, channels and bioenergetic proteins. *Biochim Biophys Acta* **1757**, 886-912 (2006).

57. Kratochvil, H.T. et al. Transient water wires mediate selective proton transport in designed channel proteins. *Nat Chem* **15**, 1012-1021 (2023).

58. Faraldo-Gomez, J.D. & Roux, B. Electrostatics of ion stabilization in a ClC chloride channel homologue from Escherichia coli. *J Mol Biol* **339**, 981-1000 (2004).

59. Walden, M. et al. Uncoupling and turnover in a Cl-/H+ exchange transporter. *J Gen Physiol* **129**, 317-29 (2007).

60. Baker, J.A., Wong, W.C., Eisenhaber, B., Warwicker, J. & Eisenhaber, F. Charged residues next to transmembrane regions revisited: "Positive-inside rule" is complemented by the "negative inside depletion/outside enrichment rule". *BMC Biol* **15**, 66 (2017).

61. Zhang, X.D., Li, Y., Yu, W.P. & Chen, T.Y. Roles of K149, G352, and H401 in the channel functions of ClC-0: testing the predictions from theoretical calculations. *J Gen Physiol* **127**, 435-47 (2006).

62. Engh, A.M., Faraldo-Gomez, J.D. & Maduke, M. The role of a conserved lysine in chloride- and voltage-dependent ClC-0 fast gating. *J Gen Physiol* **130**, 351-63 (2007).

63. Yuan, J.H. et al. Genetic spectrum and founder effect of non-dystrophic myotonia: a Japanese case series study. *J Neurol* **269**, 6406-6415 (2022).

64. Brenes, O., Pusch, M. & Morales, F. ClC-1 Chloride Channel: Inputs on the Structure-Function Relationship of Myotonia Congenita-Causing Mutations. *Biomedicines* **11**(2023).

65. Leray, X. et al. Tonic inhibition of the chloride/proton antiporter ClC-7 by PI(3,5)P2 is crucial for lysosomal pH maintenance. *Elife* **11**(2022).

66. Carpaneto, A., Boccaccio, A., Lagostena, L., Di Zanni, E. & Scholz-Starke, J. The signaling lipid phosphatidylinositol-3,5-bisphosphate targets plant CLC-a anion/H(+) exchange activity. *EMBO Rep* **18**, 1100-1107 (2017).

67. He, H. et al. Mutations in CLCN6 as a Novel Genetic Cause of Neuronal Ceroid Lipofuscinosis in Patients and a Murine Model. *Ann Neurol* **96**, 608-624 (2024).

68. Grieschat, M., Guzman, R.E., Langschwager, K., Fahlke, C. & Alekov, A.K. Metabolic energy sensing by mammalian CLC anion/proton exchangers. *EMBO Rep* **21**, e47872 (2020).

69. He, J. et al. Cryo-EM structure of the plant nitrate transporter AtCLCa reveals characteristics of the anion-binding site and the ATP-binding pocket. *J Biol Chem* **299**, 102833 (2023).

70. Zhang, B. et al. Molecular basis of ClC-6 function and its impairment in human disease. *Sci Adv* **9**, eadg4479 (2023).

71. Wan, Y. et al. Structural basis of adenine nucleotides regulation and neurodegenerative pathology in ClC-3 exchanger. *Nat Commun* **15**, 6654 (2024).

72. Skerra, A. Use of the tetracycline promoter for the tightly regulated production of a murine antibody fragment in E.coli. *Gene* **151**, 131-135 (1994).

73. Sambrook, J., Fritsch, E.F. & Maniatis, T. *Molecular Cloning, A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

74. Chien, C.H. et al. An Adaptable Phospholipid Membrane Mimetic System for Solution NMR Studies of Membrane Proteins. *J Am Chem Soc* **139**, 14829-14832 (2017).

75. Yang, M. et al. Recombinant Nepenthesin II for Hydrogen/Deuterium Exchange Mass Spectrometry. *Anal Chem* **87**, 6681-7 (2015).

76. Majumdar, R. et al. Minimizing carry-over in an online pepsin digestion system used for the H/D exchange mass spectrometric analysis of an IgG1 monoclonal antibody. *J Am Soc Mass Spectrom* **23**, 2140-8 (2012).

77. Trcka, F. et al. Human Stress-inducible Hsp70 Has a High Propensity to Form ATP-dependent Antiparallel Dimers That Are Differentially Regulated by Cochaperone Binding. *Mol Cell Proteomics* **18**, 320-337 (2019).

78. Kavan, D.a.M., P. MSTools—Web based application for visualization and presentation of HXMS data. *International Journal of Mass Spectrometry* **302**, 53-58 (2011).

79. Pettersen, E.F. et al. UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci* **30**, 70-82 (2021).

80. Lomize, M.A., Lomize, A.L., Pogozheva, I.D. & Mosberg, H.I. OPM: orientations of proteins in membranes database. *Bioinformatics* **22**, 623-5 (2006).

81. Jacobson, M.P., Friesner, R.A., Xiang, Z. & Honig, B. On the role of the crystal environment in determining protein side-chain conformations. *J Mol Biol* **320**, 597-608 (2002).

82. Betz, R. Dabble. (2017).

83. Huang, J. et al. CHARMM36m: an improved force field for folded and intrinsically disordered proteins. *Nat Methods* **14**, 71-73 (2017).

84. Klauda, J.B. et al. Update of the CHARMM all-atom additive force field for lipids: validation on six lipid types. *J Phys Chem B* **114**, 7830-43 (2010).

85. Beglov, D. & Roux, B. Finite representation of an infinite bulk system: Solvent boundary potential for computer simulations. *J. Chem. Phys.* **100**, 9050-9063 (1994).

86. Hopkins, C.W., Le Grand, S., Walker, R.C. & Roitberg, A.E. Long-Time-Step Molecular Dynamics through Hydrogen Mass Repartitioning. *J Chem Theory Comput* **11**, 1864-74 (2015).

87. Ryckaert, J.-P., Ciccotti, G., & Berendsen, H. J. C. . Numerical Integration of the Cartesian Equations of Motion of a System with Constraints: Molecular Dynamics of n-Alkanes. *Journal of Computational Physics* **23**, 327-341 (1977).

88. Roe, D.R. & Cheatham, T.E., 3rd. PTRAJ and CPPTRAJ: Software for Processing and Analysis of Molecular Dynamics Trajectory Data. *J Chem Theory Comput* **9**, 3084-95 (2013).

89. Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J Mol Graph* **14**, 33-8, 27-8 (1996).

90. The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC.

91. Lim, H.H., Shane, T. & Miller, C. Intracellular proton access in a Cl(-)/H(+) antiporter. *PLoS Biol* **10**, e1001441 (2012).

92. Schrecker, M., Korobenko, J. & Hite, R.K. Cryo-EM structure of the lysosomal chloride-proton exchanger CLC-7 in complex with OSTM1. *Elife* **9**(2020).

93. Park, E. & MacKinnon, R. Structure of the CLC-1 chloride channel from Homo sapiens. *Elife* **7**(2018).

94. Feng, L., Campbell, E.B., Hsiung, Y. & MacKinnon, R. Structure of a eukaryotic CLC transporter defines an intermediate state in the transport cycle. *Science* **330**, 635-41 (2010).

95. Xu, M. et al. CryoEM structures of the human CLC-2 voltage-gated chloride channel reveal a ball-and-chain gating mechanism. *Elife* **12**(2024).

96. Pintilie, G. & Chiu, W. Assessment of structural features in Cryo-EM density maps using SSE and side chain Z-scores. *J Struct Biol* **204**, 564-571 (2018).

Figure 1: CLC structure overview. (**a**) Side view of CLC-ec1 (pdb 1OTS). The Cl^- and H^+ permeation pathways are indicated by green and blue arrows, respectively. Bound Cl^- is shown as a green sphere. Key glutamate residues E_{gate} and Glu_{in} are shown space filled. E_{gate} physically gates the Cl^- permeation pathway and serves as a conduit for H^+; Glu_{in} lines the intracellular portion of the H^+ permeation pathway. (**b**) Cartoon depictions of the Cl^- pathway, showing the conformations of E_{gate} in high-resolution structures: "middle" (CLC-ec1[19]), "up" (CLC-ec1[23], CLC-7 [92]), "out" (CLC-ec1[24,25], CLC-1[93]), and "down" (cmCLC[94], CLC-2[95]).

Fig. 2: HDX-MS analysis of CLC-ec1 reveals dynamics at low pH. The pH-driven structural transitions as captured by HDX-MS. (**a**) Deuteration levels of CLC-ec1 at pH 4.5 were subtracted from those obtained at pH 6.5 and the results were visualized via a differential heat map. Each colored line represents a time point of the HDX kinetics. Due to differences in the chemical exchange rate at pH 4.5 and 6.5 (100-fold), time points with equal exchange were compared (e.g. 20s at pH 6.5 vs. 2000s at pH 4.5).The blue-white-red color gradient indicates deuteration levels. Differences at the final time point (denoted by a black square dot on right of the heat map) were plotted onto the CLC-ec1 structure at right. (**b**) Differential heat map capturing the time resolved HDX kinetics of the second transition, shown for time points of matching exchange (10-fold difference due to 1 pH unit shift). The last timepoint was used for CLC-ec1 structure coloring. (**c**) Selected deuterium uptake plots. The peptides monitored are indicated at the top of each plot and are mapped by color onto the CLC-ec1 structure at center (views from the perspective of the membrane and from the intracellular side). Exchange times are normalized to pH 3.0 to compensate for the differences in exchange rates. For replicated time points (20s, 633s and 20000s), error bars indicate standard deviations. Data were corrected for back-exchange. Graphs at left and right depict representative peptides that capture major pH transitions. Regions of no or small pH-induced changes are represented by the graphs at the top. The full set of uptake plots are available in **Extended Data Figure 3**. (**d**) Membrane and intracellular views of CLC-ec1 with helices labeled.

Figure 3: Cryo-EM structures of CLC-ec1. (**a, b, c**) Cryo-EM density maps for CLC-ec1, with each panel showing one of the two subunits in color: pH 7.5, orange; pH 4.0, purple; pH 3.0, cyan. (**d, e, f**) Molecular models of CLC-ec1 for structure determination at pH 7.5, 4.0, and 3.0, with colors as in panels a-c. The zoomed-in views of the models showing key residues and resolved Cl⁻ ions. Cartoons depictions to the right of the zoomed-in views are as in Fig. 1, highlighting the E$_{gate}$ conformations observed at pH 7.5 ("middle" conformation), pH 4 ("middle" and "out" conformations), and pH 3 ("out" conformation). (**g**) View from the intracellular side, highlighting major conformational changes between pH 7.5 and 4.0. Residues lining the H⁺ pathway are colored in green (E113, E148, A182, L186, A189, F190, F199, E202, E203, M204, I402, T407 and Y445), to highlight the opening of this pathway in the pH 4.0 conformation. (**h**) Comparison of pH 4.0 and pH 3.0 cryo-EM structures to the "QQQ" crystal structure. The structural alignment was done in ChimeraX using the matchmaker command aligning the dimer. (**i**) Pore radius profiles calculated using HOLE. (**j**) Zoomed-in view comparing E148 and F357 between pH 4 (out) and pH 3. The inner gate residue Y445 is shown for reference.

Figure 4: Molecular Dynamics simulations reveal water wires, Cl⁻ leaving, and inner-gate opening.
(**a**) Simulation snapshot depicting water wires that connect to the intracellular solution. (**b**) Cartoon model of proton transfer from E_{gate} to the intracellular solution along water wire 2. The resulting negatively charged E_{gate} is predicted to return to the anion pathway, expelling two Cl⁻ ions. (**c**) Cl⁻ leaves during 8 out of 10 1.5-microsecond simulations with E_{gate} deprotonated. Each plot shows the distance from inner-gate residue Y445 to the nearest Cl⁻ ion; >12 Å indicates that Cl⁻ no longer occupies anion binding sites with the pathway. Unsmoothed traces (light purple lines) and traces smoothed with a moving average (dark purple lines) are shown for all simulations. The time traces were smoothed using a moving average with a window size of 20 ns. (**d**) Simulation snapshots showing a representative Cl⁻-leaving event. Concomitant with Cl⁻ leaving to the extracellular side (i.e., away from the viewer), inner-gate residues S107 and Y445 move away from one another, opening the inner gate. Cartoon depictions are shown beneath each snapshot. (**e**) Water wires occurred with similar abundance in E_{gate} protonated and deprotonated simulations, indicating that Cl⁻ binding does not substantially influence water-wire formation. Each dot corresponds to one simulation and shows the fraction of frames in which any of the three classes of water wires is formed. The middle line of each box in the plot is the median across simulations, with the box extending from the 1st to the 3rd quartile and defining the interquartile range. Whiskers extend to last data points that are within 150% of the interquartile range. Note that Cl⁻ is always bound in simulations with E_{gate} protonated, whereas no Cl⁻ is bound in over 70% of simulation frames with E_{gate} deprotonated. A simulation snapshot showing water wires in the absence of Cl⁻ is shown in **Extended Data Fig. 8.**

Video 1: Deprotonated E$_{gate}$ expels Cl$^-$ to the outside. A close-up view of the Cl$^-$ pathway, as seen from the membrane perspective. Inner-gate residues S017 and Y445, along with outer-gate residue E$_{gate}$, are depicted as stick models. Cl$^-$ ions are represented as teal spheres. As E$_{gate}$ moves into the Cl$^-$ pathway, concomitant opening of the inner gate positions the transporter to allow Cl$^-$ entry from the intracellular side.

Figure 5: K131 mutation reduces Cl⁻ binding at S_cen but maintains Cl⁻/H⁺ coupling. (**a**) Comparison of K131A and WT CLC-ec1 cryo-EM structures. Top panels: Region around the S_{cen} site. The Cl⁻ signal at S_{cen} is substantially less distinct in K131 compared to WT (7.9 vs 11.1 sigma z-score [96]). Bottom panel: The overall structures are highly similar (RMSD = 0.52 Å). Structural alignment was done in ChimeraX using the matchmaker command aligning the dimer. (**b**) Cl⁻ binding measured by ITC for WT and K131A CLC-ec1. The top panels show representative primary data from ITC experiments. The summary table below shows results for experiments performed on 3 (WT) or 2 (K131A) separate protein preparations; for K131, no heat of binding was detected (n.h.d.) in either preparation. (**c**) Cartoon depiction of the functional assay to quantify H⁺/Cl⁻ transport rates and coupling stoichiometry. Extravesicular [Cl⁻] and [H⁺] were simultaneously measured using Ag/AgCl and pH electrodes, respectively. Because the Cl⁻/H⁺ exchange cycle nets three charges across the membrane per cycle, buildup of an electrical gradient means that measurable Cl⁻ or H⁺ transport does not occur until the addition of valinomycin, which dissipates the electrical gradient by shuttling K⁺ ions. The experimental setup involves a 2-fold gradient for H⁺, such that any leak will involve movement of H⁺ out of the vesicles, and any H⁺ movement into the vesicles must occur via CLC-ec1 actively transporting H⁺ via coupling to the downhill movement of Cl⁻. (**d**) Representative Cl⁻ and H⁺ traces (upper and lower panels, respectively) for flux assays performed on vesicles reconstituted with WT, K131A, K131M, or no CLC-ec1 protein. "Val" indicates the time of addition of valinomycin. The signals indicate net Cl⁻ release from vesicles and net H⁺ uptake into the vesicles. (**e**) Summary of transport rates for WT, K131A, and K131M CLC-ec1. The graph on the right shows the data using an expanded scale for the y-axis. (**f**) Summary of Cl⁻/H⁺ coupling stoichiometry of WT and K131 mutant transporters. P values were determined using Student's t-test. (**g**) Summary of flux assays on uncoupled transporters (E148A background) with K (WT), A, or M at the 131 position, showing that K131 mutations do not affect turnover rates of uncoupled transporters.

Figure 6: Model for reversible Cl⁻/H⁺ exchange by CLC transporters. (**a**) The 2:1 stoichiometric transport mechanism involves conformational states identified in structural studies (labeled "up", "out", "down", and "middle", shown in solid boxes) along with transient states observed in simulations (illustrated within dashed boxes). For clarity, water wires are shown only at key H⁺-transport steps. The cycle, which can proceed in either direction, facilitates the transmembrane exchange of two Cl⁻ ions for one proton (H⁺). A detailed description of the model can be found in the main text. (**b**) In K131 mutant transporters that have weakened binding at S_{ext} and S_{cen}, off-cycle transitions allow Cl⁻ transport independent of H⁺ transport. The mild uncoupling observed experimentally (3:1 Cl⁻/H⁺) can be accounted for by one off-cycle event occurring during each regular transport cycle.

	CLC-ec1 pH 7.5	CLC-ec1 pH 4	CLC-ec1 pH 3	CLC-ec1 K131A pH 7.5
Data collection and processing				
Magnification	96,000	105,000	165,000	130,000
Voltage (kV)	300	300	300	300
Electron exposure (e–/Å^2)	80	60	60	60
Defocus range (μm)	-0.8 to -2	-0.8 to -2	-0.8 to -2	-0.8 to -2
Pixel size (Å)	0.82	0.8642	0.741	0.68
Symmetry imposed	C2	C2	C2	C2
Initial particle images (no.)	971,482	871,809	1,556,961	591,019
Final particle images (no.)	72,103	261,507	66,046	307,722
Map resolution (Å)	3.2	3.2	3.1	3.3
FSC threshold	0.143	0.143	0.143	0.143
Map resolution range (Å)	2 to 7.9	2.1 to 9.5	1.8 to 7.2	2.1 to 7.8
Refinement				
Initial model used (PDB code)	1OTS	1OTS	1OTS	1OTS
Model resolution (Å)	3.1	3.1	3.1	3.3
FSC threshold	0.143	0.143	0.143	0.143
Map sharpening B factor (Å^2)	-147.2	-115.9		
Model composition				
Non-hydrogen atoms	6668	6670	6486	6660
	888	888	870	888
Protein residues	2	2	4	2
Ligands				
B factors (Å^2)				
Protein	50.12	49.39	59.92	43.67
Ligand	70.43	59.92	86.14	88.32
R.m.s. deviations				
Bond lengths (Å)	0.003	0.002	0.002	0.003
Bond angles (°)	0.511	0.535	0.529	0.541
Validation				
MolProbity score	1.01	0.97	1.22	1.07
Clashscore	2.35	1.98	3.17	1.98
Poor rotamers (%)	0.30	0.60	0.62	0.60
Ramachandran plot				
Favored (%)	98.42	97.96	97.46	97.51
Allowed (%)	1.58	2.04	2.54	2.49
Disallowed (%)	0.00	0.00	0.00	0.00

Table 1: Cryo-EM data collection, refinement and validation statistics

CLC-ec1	Transport rate (s^{-1})		Cl^-/H^+ stoichiometry	n
	Cl^-	H^+		
WT	2300 ± 100	1100 ± 40	2.0 ± 0.1	6
K131A	110 ± 30	40 ± 10	3.0 ± 0.5	4
K131M	130 ± 30	40 ± 10	3.4 ± 0.5	3
E148A	470 ± 40	n/a	n/a	5
K131A/E148A	340 ± 30	n/a	n/a	4
K131M/E148A	460 ± 150	n/a	n/a	3

Table 2: Turnover rates and stoichiometry of the proteins in this study. Transport rates and stoichiometry are displayed as the mean ± SEM. The number of flux assays for each experiment is shown by n and stem from at least two independent protein preparations. The stoichiometry is calculated from the ratio of the Cl^- and H^+ transport rates.

Extended Data Figure 1. Sequence coverage map of CLC-ec1 by peptides providing HDX data. Online proteolysis of CLC-ec1 using co-immobilized Nepenthesin-2/Pepsin column (bed volume 70 µl). Digestion conditions: 19°C, 0.4% formic acid, flow rate of 200 µL min⁻¹. Each peptide is represented by a blue bar. Secondary structure elements are shown above the sequence. Almost complete (89%) sequence coverage was obtained through 237 peptides with average peptide length of 11.4 amino acids and average redundancy of 6.4.

Extended Data Figure 2: CLC-ec1 stability at pH 3.0. Size exclusion chromatography of CLC-ec1 using EnrichTM SEC 650 10x300 24 mL (Biorad), flow rate 1 mL/min. The top panel shows CLC-ec1 prepared at pH 7.5. The lower panels show CLC-ec1 assessed using size exclusion chromatography immediately after preparation at pH 3.0 and at several time points, demonstrating stability over an 8-hour period.

https://drive.google.com/file/d/1ehym3xH6S3V6jqn-heWvTFHm0tO6SJ-V/view?usp=sharing

Extended Data Figure 3: Deuterium uptake plots for all CLC-ec1 peptides providing HDX data. Five experimental conditions were followed: pH 3.0 (black), pH 3.5 (grey), pH 4.0 (blue), pH 4.5 (cyan) and 6.5 (orange). Exchange times are normalized to pH 3.0 to compensate for the difference in the exchange rate. For replicated time points (20s, 633s, and 20000s) average values with respective standard deviations are shown in plots. Data were corrected for back-exchange. Peptide limits and charge states are shown above each plot.

Extended Data Fig. 4: Cryo EM workflow and validation data for CLC-ec1 at pH 7.5.
(**a**) Cryo-EM data processing workflow. (**b**) Gold-standard FSC curve. The resolution is estimated based on FSC at 0.143. (**c**) Angular distribution plot (**d**) Per-residue Q-score as a function of residue number. The expected Q-score at the map resolution is indicated by the dotted line. (**e**) The cryo-EM density and molecular model overlay for central Cl⁻ binding site. The key residues and bound Cl⁻ and their corresponding Q-score are annotated. (**f**) The overlay of our cryo-EM structure (salmon color) with the previous crystal structure (green) (PDB: 1OTS).

Extended Data Fig. 5. Cryo EM workflow and validation data for CLC-ec1 at pH 4.0
(**a**) Cryo-EM data processing workflow. (**b**) Gold-standard FSC curve. The resolution is estimated based on FSC at 0.143. (**c**) Angular distribution plot (**d**) Per-residue Q-score as a function of residue number. The expected Q-score at the map resolution is indicated by the dotted line. (**e**) The cryo-EM density and molecular model overlay for central Cl⁻ binding site. The key residues and bound Cl⁻ and their corresponding Q-score are annotated.

Extended Data Figure 6: Cryo EM workflow and validation data for CLC-ec1 at pH 3.0.
(**a**) Cryo-EM data processing workflow. (**b**) Gold-standard FSC curve. The resolution is estimated based on FSC at 0.143. (**c**) Angular distribution plot (**d**) Per-residue Q-score as a function of residue number. The expected Q-score at the map resolution is indicated by the dotted line. (**e**) The cryo-EM density and molecular model overlay for central Cl⁻ binding site. Key residues and bound Cl⁻ and their corresponding Q-score are annotated. (**f**) Overlay of our cryo-EM structure (cyan) with the previous QQQ crystal structure (dark blue) (PDB: 6V2J).

Protonated Glu_{gate}

Extended Data Figure 7: Cl⁻ remains in the anion pathway during simulations with E_{gate} protonated. The presence of Cl⁻ within the anion pathway during simulations was evaluated by plotting the distance from the Cα atom of inner-gate residue Y445 to the nearest Cl⁻ ion.

Extended Data Figure 8: Water wires occur in the presence and absence of Cl⁻.
(**a**) Example snapshot of water wires formed in the presence of Cl⁻ in the permeation pathway.
(**b**) Example snapshot of water wires formed in the absence of Cl⁻ in the permeation pathway

Extended Data Figure 9. Cryo EM validation data for K131A CLC-ec1. (**a**) Cryo-EM data processing workflow. (**b**) Gold-standard FSC curve. The resolution is estimated based on FSC at 0.143. (**c**) Angular distribution plot (**d**) Per-residue Q-score as a function of residue number. The expected Q-score at the map resolution is indicated by dotted line. (**e**) The cryo-EM density and molecular model overlay in the region of K131 (left panel) and A131 (right panel). (**f**) The cryo-EM density and molecular model overlay for central Cl⁻ binding site. The key residues and bound Cl⁻ and their corresponding Q-score are annotated.