**Enteropancreatic neurons drive the glucoregulatory response to ingested lipid**

**Authors:** Anna G. Roberts[1]*, Leah Meyer[1]*, Mariana Norton[1], Phyllis Phuah[1], Aldara

Martin Alonso[1], Georgina K.C. Dowsett[2], Sijing Cheng[1], Cecilia Dunsterville[1], Jieruo Liu[1],

Pei-En Chung[1], Yuxuan Tao[1], Tobias Smitherman-Cairns[1], Alexandra B. Deutsch[1], Abhisekh

Chatterjee[1], Brian Y.H. Lam[2], Aylin C. Hanyaloglu[1], Ben Jones[1], Giles S.H. Yeo[2], Victoria

Salem[1], Kevin G. Murphy[1]

[1]Division of Diabetes, Endocrinology and Metabolism, Department of Metabolism,

Digestion, and Reproduction, Faculty of Medicine, Imperial College London, 6th Floor

Commonwealth Building, Hammersmith Hospital, London W12 0NN, UK.

[2]MRC Metabolic Diseases Unit, Institute of Metabolic Science Metabolic Research

Laboratories, University of Cambridge, Cambridge, CB2 0QQ, UK

*These authors contributed equally to this work.

**Graphical Abstract**

**Summary**

**Enteropancreatic neurons project from the small intestinal wall to the pancreas.
Though well positioned to mediate the effects of ingested nutrients on pancreatic
function, the metabolic role of these neurons is unclear. Diets rich in olive oil promote
weight loss and improve remission rates in patients with T2DM. Here, we show that
olive oil improves acute glucose tolerance by stimulating insulin release via neurotensin
receptor type 1 (NTSR1)-expressing enteropancreatic neurons. These neurons are
necessary for the effects of olive and neurotensin on glucose tolerance, and their
activation is sufficient to improve glucose tolerance. These findings suggest a
mechanism by which dietary olive oil regulates blood glucose levels and present a novel
functional role for enteropancreatic neurons in regulating glucose homeostasis.**

**Introduction**

The Mediterranean Diet, rich in vegetables, nuts, fruit, whole grains and olive oil, reduces
metabolic syndrome indicators, including reduced risk of cardiovascular disease, alongside
improvements in weight management and T2DM remission[1-3]. Evidence suggests that olive
oil may be the crucial component for improved glucoregulation. A Mediterranean Diet
supplemented with additional olive oil further reduced the incidence of T2DM and chronic
adherence to olive-oil enriched diets lowers blood glucose levels[4,5].

Oleic acid is the main monounsaturated fatty acid (MUFA) within the Mediterranean Diet[6]
and is present at high levels in olive oil. Previous work has shown both olive oil- and oleic
acid induce secretion of the gut hormone neurotensin from enteroendocrine cell models and
in vivo in a range of species[7-10]. Neurotensin is a 13-amino acid peptide synthesized in the
gastrointestinal tract and the central nervous system. Most known biological effects of

neurotensin are mediated by the neurotensin receptor type 1 (NTSR1). Centrally,

neurotensin-NTSR1 signalling plays roles in thermoregulation[11], analgesia[12], addiction[13],

food intake[14] and blood pressure regulation[15], and peripherally has been reported to modulate

gastric emptying[16,17] and lipid absorption[18-20]. However, the role of the neurotensin-NTSR1

system in glucose homeostasis is unclear[21-27]. Here we show that the acute glucoregulatory

effects of olive oil are mediated through NTSR1-expressing enteropancreatic neurons that

extend directly from the wall of the duodenum to the pancreas and represent a novel pathway

by which dietary intake regulates glucose homeostasis. This is the first report of a

physiological role for a population of enteropancreatic neurons.

**Results**

**Olive oil increases insulin secretion and improves acute glucose tolerance via a NTSR1-**

**mediated pathway**

We investigated the effects of olive oil on acute glucose tolerance in mice. Olive oil

significantly improved glucose tolerance 15-minutes following intraperitoneal or oral glucose

administration in lean mice (Fig.1A, Extended Data Fig.1A). Olive oil also increased

circulating insulin levels (Fig.1B). Olive oil is rich in MUFAs which have been suggested to

mediate its metabolic effects. A control oil rich in saturated fatty acids rather than MUFAs

did not improve glucose tolerance in mice (Extended Data Fig.1B). Of the MUFAs in virgin

olive oil, oleic acid comprises approximately 60-80%[28]. We therefore investigated whether

oleic acid might be responsible for the glucoregulatory effects of olive oil. Oral co-

administration of the lipase inhibitor Orlistat, which prevents the breakdown of fats into fatty

acids, blocked the improvement in glucose tolerance seen with olive oil (Fig.1C), and oral

administration of oleic acid alone significantly improved glucose tolerance (Fig.1D). Within

the gastrointestinal tract, MUFAs are predominantly sensed by the free fatty acid receptor 4

(FFAR4). We found that a specific FFAR4 antagonist also attenuated the glucoregulatory

effect of olive oil (Fig.1E). Together these data suggest that the acute effects of olive oil on

glucose tolerance are mediated by the sensing of its constituent fatty acids through the

FFAR4.

Previous work has shown that both oral and intra-luminal infusions of olive oil and oleic acid

increase circulating neurotensin levels[7-10]. We also found that plasma neurotensin levels

increased in mice following oral administration of olive oil (Fig.1F). We therefore

investigated the role of the NTSR1 in the glucoregulatory response to olive oil. Pretreatment

with an NTSR1 antagonist significantly attenuated the glucose-lowering effects of olive oil,

suggesting that a significant part of olive oil's acute glucoregulatory response is controlled by

activation of the NTSR1 (Fig 1G). The role of NTSR1 was nutrient specific, as the same

NTSR1 antagonist did not block the glucoregulatory effects of an oral protein bolus

(Extended Data Fig.1C). Olive oil has also been reported to stimulate the release of the

incretin glucagon-like peptide-1 (GLP-1)[29]. To confirm that the majority of the effects of

olive oil on glucose tolerance were not mediated by GLP-1, we investigated the effect of a

GLP-1 receptor antagonist and conducted glucose tolerance tests in a Pdx1CreERT-GLP-

1Rfl/fl mouse model in which the GLP-1R was knocked out of pancreatic beta cells. We

observed that olive oil administration still improved glucose tolerance in mice treated with a

GLP-1 antagonist, or lacking GLP-1R in beta cells, suggesting GLP-1 signalling is not

required for this effect (Extended Data Fig.2G&H).

To assess whether NTSR1 mediates the effects of olive oil in a physiological setting, mice

were fed olive-oil enriched chow following an overnight fast. Mice administered an NTSR1

antagonist prior to refeed had higher blood glucose values 1h later, suggesting that olive oil

restrains postprandial blood glucose levels in a NTSR1-dependent manner (Fig.1H).

We then investigated the glucoregulatory effects of exogenous neurotensin administration.

We found intraperitoneal neurotensin improved acute glucose tolerance in lean mice (Fig.1I).

We also found that administration of neurotensin was able to improve glucose tolerance in

diet-induced obese (DIO) mice (Fig.1J). Neurotensin administration was also associated with

an increase in circulating insulin levels (Fig.1K).

GLP-1R agonists are currently important gut hormone-based pharmacotherapies for T2DM,

and there is much interest in combining GLP-1R agonism with other glucoregulatory agents,

to increase effectiveness and reduce side effects. Previous work has suggested potential for

neurotensin/GLP-1 co-agonism to lower blood glucose levels[26,27]. We similarly confirmed

that low-dose neurotensin had an additive effect on glucose tolerance when co-administered

with a low-dose of a GLP-1R agonist (Extended Data Fig. 2A).

Neurotensin has been shown to play an important role in controlling lipid absorption in the

gut[18-20]. We therefore investigated whether the glucoregulatory effects of peripheral

neurotensin were driven by changes in circulating lipid levels that might be sensed at the

level of the beta-cell[30]. We found no difference in circulating triglyceride levels in a glucose

tolerance test at the time point at which neurotensin or olive oil have glucoregulatory effects

(Extended Data Fig.1D&E). Having established that olive oil improves glucose tolerance

through a neurotensin-NTSR1-dependent mechanism, we then investigated the specific

pathway responsible.

**Neurotensin activates neurons in the myenteric plexus through NTSR1, and NTSR1 is**

**expressed in enteropancreatic neurons**

In accord with previous work reporting that NTSR1 mRNA is not expressed in the pancreas

itself[31,32], we found negligible expression in murine pancreatic islets (Fig.2B), suggesting

neurotensin does not act directly on the pancreas to improve glucose tolerance. We therefore

investigated whether neurotensin might stimulate insulin release through a neural mechanism.

The major neural pathway stimulating insulin release acts via the muscarinic-3 acetylcholine

receptor (m3R), which is highly expressed on pancreatic beta cells[33,34]. We found that

administration of an m3R-antagonist blocked the glucoregulatory effects of both oral olive

and peripheral neurotensin (Fig.1L&M). The pancreatic ganglia provide significant direct

cholinergic innervation of the pancreatic islets[35], with the activity of these ganglia modulated

by external neuronal populations including vagal neurons extending from the dorsal motor

complex (DMC), sympathetic efferent neuronal populations, and neuronal projections that

extend directly from the gastrointestinal tract. We were therefore interested to determine

whether neurotensin acted centrally, indirectly via afferent vagal signalling, or via

enteropancreatic neurons to control glucoregulation.

We confirmed the NTSR1 to be highly expressed in the murine nodose ganglia, arcuate

nucleus and myenteric plexus neurons isolated from the proximal duodenum (Fig.2B). Mice

in which NTSR1-expressing neurons were ablated in the nodose ganglia, where the cell

bodies of afferent vagal neurons reside, showed improved glucose tolerance in response to

neurotensin, suggesting that neurotensin does not drive its glucoregulatory effects via direct

activation of vagal-NTSR1 (Extended Data Fig.3). Central administration of neurotensin at a

dose sufficient to lower body temperature did not improve glucose tolerance, and central

blockade of NTSR1 sufficient to block the anorectic effects of peripheral neurotensin did not

block the effects of peripheral neurotensin on glucose tolerance. These data suggest

peripherally administered neurotensin does not act on the brain to regulate glucose

homeostasis (Extended Data Fig.4). In accord with previous reports[36], we identified NTSR1-

expression in the myenteric plexus of the proximal duodenum (Fig.2A&B). It has been

reported that NTSR1 immunoreactivity is present in neurons in the pancreas[37], though

NTSR1 mRNA has not been detected[31,32]. We therefore hypothesised that NTSR1 was

expressed on enteropancreatic neurons that originate in the wall of the duodenum and that

extend to the pancreas to regulate pancreatic function. Enteropancreatic neurons were first

reported in both dogs and humans[38] and later identified in mice[39-41]. However, there has been

little study of their physiological role in vivo.

Having identified that the NTSR1 is expressed in neurons of the myenteric plexus of the

proximal duodenum, we assessed whether neurotensin could activate these neurons in a

NTSR1-dependent manner. We isolated myenteric neurons (MENs) from the proximal

duodenum of *Ntsr1-cre:tdTomato* mice.  We found cultured MENs expressing tdTomato

(NTSR1+) to be activated by neurotensin; this effect was blocked by an NTSR1 antagonist

(Fig.2C&D).   Using *Ntsr1-cre:tdTomato* reporter mice we found that NTSR1-expressing

neurons extended from the myenteric plexus in the duodenum and extended in fibre bundles

into the pancreas (Fig. 2E), where individual NTSR1-expressing processes could be

visualised close to pancreatic islets (Fig. 2F). Following injection of a retrograde AAV

encoding enhanced green fluorescent protein (eGFP) into the pancreas, we were able to detect

eGFP expression in the myenteric plexus of the duodenum (Fig. 2G).

**An intact enteropancreatic connection is required for the glucoregulatory effects of**

**neurotensin and olive oil**

To determine the necessity of these enteropancreatic neurons for the effects of olive oil and

neurotensin on glucose tolerance, the proximal duodenum was partially surgically separated

from its adjacent pancreatic section (Fig.3A). An eGFP-encoding AAV

(AAV5.hSyn.eGFP.WPRE.bGH) was injected into the pancreas adjacent to the proximal

duodenum to allow verification that enteropancreatic neuronal innervation of the pancreas

was reduced; a significant reduction in eGFP expression was observed in the duodenum of

mice with a separated enteropancreatic connection (Extended Data Fig.5B). Following

recovery, mice underwent a series of glucose tolerance tests. All mice exhibited significantly

improved glucose tolerance when treated with the GLP-1R agonist exendin-4 or with an oral

bolus of whey protein, suggesting the pancreas remained functional and responsive to a non-

lipid nutrient stimulus and a systemic gut hormonal signal (Extended Data Fig.5C-F). In

contrast, both neurotensin and olive oil significantly improved glucose tolerance in mice with

an intact enteropancreatic connection (Fig.3B-E) but not in mice in which the

enteropancreatic neurons were disrupted. When fasted mice were refed olive–oil enriched

chow, mice with a disrupted enteropancreatic connection had a significantly higher

postprandial glucose excursion compared to control mice (Fig.3F). Together, these data

suggest that the glucose lowering effects of olive oil ingestion and neurotensin administration

are reliant on enteropancreatic neurons originating in the proximal duodenum.

**NTSR1-expressing enteropancreatic neurons are sufficient to improve glucose tolerance**

**and necessary for the glucoregulatory effect of neurotensin and olive oil**

We next investigated whether activating NTSR1-expressing enteropancreatic neurons could

modulate glucose tolerance. A cre-dependent retrograde stimulatory DREADD-encoding

AAV, (AAVrg-hSyn-DIO-hM3D(Gq)-mCherry) was injected into the head of the pancreas in

*NTSR1-cre* mice (Fig.4A). Post-experimental dissection of the longitudinal muscle of the

myenteric plexus (LMMP) confirmed mCherry expression in the neurons of the myenteric

plexus. Two weeks later, to allow for recovery and sufficient gene expression, intraperitoneal

administration of the DREADD ligand compound 21 (C21) 15 minutes prior to an

intraperitoneal glucose tolerance test (IPGTT) significantly improved glucose tolerance

(Fig.4B&C). Activation of these neurons caused a trend for increased circulating insulin

levels (Extended Data Fig.6A-C). In a separate experiment in these mice, administration of

C21 did not alter core body temperature, suggesting it was not engaging major central

NTSR1 circuits (Extended Data Fig. 6D). These data suggest that activation of NTSR1-

expressing neurons that extend to the pancreas from the myenteric plexus of the proximal

duodenum is sufficient to improve glucose tolerance.

To assess whether NTSR1-expressing enteropancreatic neurons are necessary for the

glucoregulatory effects of peripheral neurotensin and olive oil, a cre-dependent Diphtheria

Toxin Receptor-encoding AAV (AAV2-FLEX-DTR-GFP) was injected into the head of the

pancreas of NTSR1-cre mice (Figure 4D). As a viral control, the same cre-dependent DTR-

encoding AAV (AAV2-FLEX-DTR-GFP) was injected into the pancreas of NTSR1-WT

mice and as a genotype control, a cre-dependent GFP-encoding AAV (AAV2-pCAG-FLEX-

EGFP-WPRE) was injected into the pancreas of NTSR1-cre mice. Following a 2-week rest

period to allow sufficient viral expression, all mice were injected with diphtheria toxin

intraperitoneally, which specifically killed NTSR1-expressing enteropancreatic neurons in

NTSR1-cre mice injected with AAV-DTR (Extended Data Fig.7B). All mice exhibited

significantly improved glucose tolerance when treated with the GLP-1R agonist exendin-4 or

with an oral bolus of whey protein, suggesting the pancreas remained functional and

responsive to a non-lipid nutrient stimulus and a systemic gut hormonal signal (Extended

Data Fig.7C-H). In glucose tolerance tests, both peripheral neurotensin and olive oil retained

their glucoregulatory effects in viral and genotype controls (Fig.4E&H, Extended Data

Fig.7I&J). However, in NTSR1-cre mice injected with AAV2-FLEX-DTR-GFP, in which

enteropancreatic NTSR1-expressing cells had been ablated, neither neurotensin nor olive oil

improved glucose tolerance (Fig4.E-F,H-I). In accord with this lack of effect on glucose

levels, peripheral neurotensin administration significantly increased circulating insulin levels

in genotype control mice but not in NTSR1-cre mice injected with AAV2-FLEX-DTR-GFP

(Fig4J&K). NTSR1-cre mice injected with AAV2-FLEX-DTR-GFP also experienced an

enhanced glucose excursion compared to genotype control mice in a fast-refeed study where

mice were refed olive oil-enriched chow (Fig.4G). These data suggest that NTSR1-

expressing enteropancreatic neurons have a role in glucose regulation, stimulating insulin

secretion in response to both exogenous neurotensin and physiological neurotensin release

following ingestion of olive oil.

**Discussion**

Dietary olive oil has previously been linked to improvements in glycaemia and a reduced risk

of developing type 2 diabetes in mice and humans[5,42,43]. However, the underlying mechanism

for the glucoregulatory effects of olive oil is poorly characterised. Here, we have shown for

the first time that ingested olive oil drives an acute improvement in glucose tolerance in mice

through driving neurotensin release. We have further demonstrated that this acute

glucoregulatory effect of olive oil depends on the digestion of olive oil into its component

fatty acids, and is at least partially mediated by the FFAR4.

There is much current interest in agonists of the incretin gut hormones receptors: GLP-1R

and GIPR as pharmacotherapies for enhancing glucoregulation. Administration of oleic acid,

the major fatty acid constituent of olive oil, has been shown to stimulate release of the gut

hormone neurotensin in both rodents and humans[7,10]. Neurotensin has previously been

suggested to have a role in glucose regulation, though reported results vary and the

mechanisms are unclear[21,22,24]. Previous studies have found neurotensin to have both

inhibitory and stimulatory effects on insulin release in vitro and in vivo[21-23,25,44], though more

recent work has suggested a synthetic NTSR1 agonist and a PEGylated form of neurotensin

can improve glucose tolerance[26,27]. Here, we provide evidence that neurotensin acts as an

incretin and has the potential to improve glucoregulation in both lean and diet-induced obese

mice.

Our data shows that neurotensin works to promote insulin secretion via NTSR1-expressing

enteropancreatic neurons that originate in the proximal duodenum. Although enteropancreatic

neurons were first discovered nearly fifty years ago[45] they have been relatively little studied.

Serotonergic and NADPH diaphorase-expressing enteropancreatic neurons have been

reported to have an inhibitory and excitatory role in exocrine pancreas secretions,

respectively, and myenteric cholinergic neurons have been shown to provide excitatory inputs

to the pancreas[39,40,46]. Using surgical and genetic mouse models, we have demonstrated that

activation of NTSR1-expressing enteropancreatic neurons is sufficient to improve glucose

tolerance, and these neurons are necessary for the glucoregulatory effects of neurotensin and

olive oil. This represents the first functional role for a population of enteropancreatic neurons

in the modulation of the endocrine pancreas.

Ingestion and infusion of olive oil and oleate has previously been reported to stimulate

endogenous secretion of neurotensin[7,8,10], and we also found that oral gavage of olive oil

increased circulating levels of neurotensin. However, neurotensin has a short half-life in

circulation[47], and it is possible that other agonists of the NTSR1, such as xenin, also play a

role in the glucoregulatory effect of olive oil. Xenin is a molecule secreted from

enteroendocrine K cells, and exogenous xenin has been reported to potentiate the

insulinotropic actions of GIP and bind to the NTSR1 to promote gastric emptying[48]. There

are, however, no reports of a physiological role for endogenous xenin.

The NTSR1 has also been found to promote lipid absorption, and knockout of the NTSR1 has

been shown to protect rodents from diet-induced obesity[19]. Whilst we have demonstrated that

the glucoregulatory effects of exogenous neurotensin and olive oil occur before any increase

in circulating triglycerides is detected, chronically agonising the NTSR1 as a therapy for

impaired glucoregulation may result in weight gain and obesity. Further research into the

enteropancreatic NTSR1 neurons might identify other drug targets to facilitate the more

specific targeting of these neurons for glucoregulation. We have also shown that neurotensin

can have an additive effect on glucoregulation when administered in combination with the

GLP-1 receptor agonist exendin-4, further underpinning the potential of NTSR1-specific

therapeutics for the treatment of type 2 diabetes. Chronic studies in disease models are now

necessary to determine the therapeutic utility of the NTSR1 system.

This is the first characterisation of a functional role for a neuronal population connecting the

gut directly to the pancreas. The position of these neurons in the upper duodenum allows

them to respond rapidly to dietary nutrients to drive appropriate pancreatic responses. There

may also be additional physiological roles for this enteropancreatic system. Further work is

necessary to explore whether NTSR1 agonists and other agents targeting enteropancreatic

neurons may be useful in the treatment of metabolic disease.

**Main References**

1.      Babio N, Bulló M, Salas-Salvadó J. Mediterranean diet and metabolic syndrome: the

evidence. *Public health nutrition*. 2009;12(9A):1607-1617.

2.      Esposito K, Maiorino MI, Petrizzo M, Bellastella G, Giugliano D. The effects of a

Mediterranean diet on the need for diabetes drugs and remission of newly diagnosed type 2

diabetes: follow-up of a randomized trial. *Diabetes care*. 2014;37(7):1824-1830.

3.      Shai I, Schwarzfuchs D, Henkin Y, et al. Weight loss with a low-carbohydrate,

Mediterranean, or low-fat diet. *New England Journal of Medicine*. 2008;359(3):229-241.

4.    Ryan M, McInerney D, Owens D, Collins P, Johnson A, Tomkin GH. Diabetes and

the Mediterranean diet: a beneficial effect of oleic acid on insulin sensitivity, adipocyte

glucose transport and endothelium-dependent vasoreactivity. *QJM*. Feb 2000;93(2):85-91.

doi:10.1093/qjmed/93.2.85

5.    Jurado-Ruiz E, Álvarez-Amor L, Varela LM, et al. Extra virgin olive oil diet

intervention improves insulin resistance and islet performance in diet-induced diabetes in

mice. *Sci Rep*. Aug 05 2019;9(1):11311. doi:10.1038/s41598-019-47904-z

6.    Kris-Etherton PM. Monounsaturated fatty acids and risk of cardiovascular disease.

*Circulation*. 1999;100(11):1253-1258.

7.    Ferris CF, Armstrong MJ, George JK, Stevens CA, Carraway RE, Leeman SE.

Alcohol and fatty acid stimulation of neurotensin release from rat small intestine.

*Endocrinology*. Mar 1985;116(3):1133-8. doi:10.1210/endo-116-3-1133

8.    Fujimura M, Khalil T, Sakamoto T, et al. Release of neurotensin by selective

perfusion of the jejunum with oleic acid in dogs. *Gastroenterology*. Jun 1989;96(6):1502-5.

doi:10.1016/0016-5085(89)90518-0

9.    Dumoulin Vr, Moro Fdr, Barcelo A, Dakka T, Cuber J-C. Peptide YY, glucagon-like

peptide-1, and neurotensin responses to luminal factors in the isolated vascularly perfused rat

ileum. *Endocrinology*. 1998;139(9):3780-3786.

10.    Drewe J, Mihailovic S, D'Amato M, Beglinger C. Regulation of fat-stimulated

neurotensin secretion in healthy subjects. *J Clin Endocrinol Metab*. May 2008;93(5):1964-70.

doi:10.1210/jc.2007-2238

11.    Bissette G, Nemeroff CB, Loosen PT, Prange AJ, Lipton MA. Hypothermia and

intolerance to cold induced by intracisternal administration of the hypothalamic peptide

neurotensin. *Nature*. Aug 12 1976;262(5569):607-9. doi:10.1038/262607a0

12.    Dobner PR. Neurotensin and pain modulation. *Peptides*. 2006;27(10):2405-2414.

13.    Rodríguez FD, Sánchez ML, Coveñas R. Neurotensin and Alcohol Use Disorders:

Towards a Pharmacological Treatment. *Int J Mol Sci*. May 12

2023;24(10)doi:10.3390/ijms24108656

14.    Ramirez-Virella J, Leinninger GM. The Role of Central Neurotensin in Regulating

Feeding and Body Weight. *Endocrinology*. 2021-05-01

2021;162(5)doi:10.1210/endocr/bqab038

15.    Osadchii OE. Emerging role of neurotensin in regulation of the cardiovascular

system. *European journal of pharmacology*. 2015;762:184-192.

16.    Blackburn AM, Fletcher DR, Bloom SR, et al. Effect of neurotensin on gastric

function in man. *Lancet*. May 10 1980;1(8176):987-9. doi:10.1016/s0140-6736(80)91434-8

17.    Hellström P, Nylander G, Rosell S. Effects of neurotensin on the transit of

gastrointestinal contents in the rat. *Acta Physiologica Scandinavica*. 1982;115(2):239-243.

18.    Gui X, Carraway RE. Enhancement of jejunal absorption of conjugated bile acid by

neurotensin in rats. *Gastroenterology*. Jan 2001;120(1):151-60. doi:10.1053/gast.2001.20876

19.    Li J, Song J, Zaytseva YY, et al. An obligatory role for neurotensin in high-fat-diet-

induced obesity. *Nature*. 05 19 2016;533(7603):411-5. doi:10.1038/nature17662

20.    Barchetta I, Cimini FA, Capoccia D, et al. Neurotensin Is a Lipid-Induced

Gastrointestinal Peptide Associated with Visceral Adipose Tissue Inflammation in Obesity.

*Nutrients*. Apr 23 2018;10(4)doi:10.3390/nu10040526

21.    Carraway RE, Demers LM, Leeman SE. Hyperglycemic effect of neurotensin, a

hypothalamic peptide. *Endocrinology*. Dec 1976;99(6):1452-62. doi:10.1210/endo-99-6-1452

22.    Kaneto A, Kaneko T, Kajinuma H, Kosaka K. Effects of substance P and neurotensin

infused intrapancreatically on glucagon and insulin secretion. *Endocrinology*. Feb

1978;102(2):393-401. doi:10.1210/endo-102-2-393

23.     Dolais-Kitabgi J, Kitabgi P, Freychet P. Glucose and glucagon do stimulate

somatostatin release from isolated pancreatic islets. *Diabetologia*. Sep 1981;21(3):238.

doi:10.1007/BF00252662

24.     Blackburn AM, Bloom SR, Edwards AV. Pancreatic endocrine responses to

physiological changes in plasma neurotensin concentration in the calf. *J Physiol*. Sep

1981;318:407-12. doi:10.1113/jphysiol.1981.sp013874

25.     Béraud-Dufour S, Abderrahmani A, Noel J, et al. Neurotensin is a regulator of insulin

secretion in pancreatic beta-cells. *Int J Biochem Cell Biol*. Oct 2010;42(10):1681-8.

doi:10.1016/j.biocel.2010.06.018

26.     Craig S, Gault VA, Shiels C, Hamscher G, Irwin N. Comparison of independent and

combined effects of the neurotensin receptor agonist, JMV-449, and incretin mimetics on

pancreatic islet function, glucose homeostasis and appetite control. *Biochimica et Biophysica*

*Acta (BBA)-General Subjects*. 2021;1865(8):129917.

27.     Ratner C, He Z, Grunddal KV, et al. Long-Acting Neurotensin Synergizes With

Liraglutide to Reverse Obesity Through a Melanocortin-Dependent Pathway. *Diabetes*. 2019-

06-01 2019;68(6):1329-1340. doi:10.2337/db18-1009

28.     Revelou PK, Xagoraris M, Alexandropoulou A, et al. Chemometric Study of Fatty

Acid Composition of Virgin Olive Oil from Four Widespread Greek Cultivars. *Molecules*. Jul

08 2021;26(14)doi:10.3390/molecules26144151

29.     Valverde I, Moreno P, Cancelas J, Prieto PG, Villanueva-Peñacarrillo ML, Malaisse

WJ. Effects of an Olive Oil-enriched Diet on Glucagon-like Peptide-1. *Olives and Olive Oil*

*in Health and Disease Prevention*. Elsevier; 2010:1213-1220.

30.     Komatsu M, Sharp G. Palmitate and myristate selectively mimic the effect of glucose

in augmenting insulin release in the absence of extracellular Ca2+. *Diabetes*. 1998;47(3):352-

357.

31.     Reubi JC, Waser B, Friess H, Büchler M, Laissue J. Neurotensin receptors: a new

marker for human ductal pancreatic adenocarcinoma. *Gut*. Apr 1998;42(4):546-50.

doi:10.1136/gut.42.4.546

32.     Wang L, Friess H, Zhu Z, et al. Neurotensin receptor-1 mRNA analysis in normal

pancreas and pancreatic disease. *Clin Cancer Res*. Feb 2000;6(2):566-71.

33.     Duttaroy A, Zimliki CL, Gautam D, Cui Y, Mears D, Wess J. Muscarinic stimulation

of pancreatic insulin and glucagon release is abolished in M3 muscarinic acetylcholine

receptor–deficient mice. *Diabetes*. 2004;53(7):1714-1720.

34.     Renuka T, Robinson R, Paulose C. Increased insulin secretion by muscarinic M1 and

M3 receptor function from rat pancreatic islets in vitro. *Neurochemical research*.

2006;31:313-320.

35.     Li W, Yu G, Liu Y, Sha L. Intrapancreatic Ganglia and Neural Regulation of

Pancreatic Endocrine Secretion. *Front Neurosci*. 2019;13:21. doi:10.3389/fnins.2019.00021

36.     Zhang S, Hyrc K, Wang S, Wice BM. Xenin-25 increases cytosolic free calcium

levels and acetylcholine release from a subset of myenteric neurons. *Am J Physiol*

*Gastrointest Liver Physiol*. Dec 15 2012;303(12):G1347-55. doi:10.1152/ajpgi.00116.2012

37.     Chowdhury S, Wang S, Patterson BW, Reeds DN, Wice BM. The combination of GIP

plus xenin-25 indirectly increases pancreatic polypeptide release in humans with and without

type 2 diabetes mellitus. *Regul Pept*. Nov 10 2013;187:42-50.

doi:10.1016/j.regpep.2013.10.003

38.     Tiscornia OM. The neural control of exocrine and endocrine pancreas. *Am J*

*Gastroenterol*. Jun 1977;67(6):541-60.

39.     Kirchgessner AL, Gershon MD. Innervation of the pancreas by neurons in the gut. *J*

*Neurosci*. May 1990;10(5):1626-42. doi:10.1523/JNEUROSCI.10-05-01626.1990

40.    Kirchgessner AL, Gershon MD. Presynaptic inhibition by serotonin of nerve-

mediated secretion of pancreatic amylase. *Am J Physiol*. Feb 1995;268(2 Pt 1):G339-45.

doi:10.1152/ajpgi.1995.268.2.G339

41.    Kirchgessner AL, Liu MT, Gershon MD. In situ identification and visualization of

neurons that mediate enteric and enteropancreatic reflexes. *J Comp Neurol*. Jul 22

1996;371(2):270-86. doi:10.1002/(SICI)1096-9861(19960722)371:2<270::AID-

CNE7>3.0.CO;2-#

42.    Salas-Salvadó J, Bulló M, Estruch R, et al. Prevention of diabetes with Mediterranean

diets: a subgroup analysis of a randomized trial. *Ann Intern Med*. Jan 07 2014;160(1):1-10.

doi:10.7326/M13-1725

43.    Violi F, Loffredo L, Pignatelli P, et al. Extra virgin olive oil use is associated with

improved post-prandial blood glucose and LDL cholesterol in healthy subjects. *Nutr*

*Diabetes*. Jul 20 2015;5(7):e172. doi:10.1038/nutd.2015.23

44.    Blackburn AM, Bloom SR, Edwards AV. Pancreatic endocrine responses to

exogenous neurotensin in the conscious calf. *J Physiol*. May 1981;314:11-21.

doi:10.1113/jphysiol.1981.sp013686

45.    Tiscornia OM, Martinez JL, Sarles H. Some aspects of human and canine

macroscopic pancreas innervation. *Am J Gastroenterol*. Oct 1976;66(4):353-61.

46.    Kirchgessner AL, Liu MT, Gershon MD. NADPH diaphorase (nitric oxide synthase)-

containing nerves in the enteropancreatic innervation: sources, co-stored neuropeptides, and

pancreatic function. *J Comp Neurol*. Apr 01 1994;342(1):115-30.

doi:10.1002/cne.903420111

47.    Aronin N, Carraway RE, Ferris CF, Hammer RA, Leeman SE. The stability and

metabolism of intravenously administered neurotensin in the rat. *Peptides*. 1982;3(4):637-

642.

48.      Chowdhury S, Reeds DN, Crimmins DL, et al. Xenin-25 delays gastric emptying and

reduces postprandial glucose levels in humans with and without type 2 diabetes. *Am J Physiol*

*Gastrointest Liver Physiol*. Feb 15 2014;306(4):G301-9. doi:10.1152/ajpgi.00383.2013

**Contact for reagent and resource sharing**

413  Further information and requests for reagents may be directed to and will be fulfilled by the

414  Lead Contact, Kevin G Murphy (k.g.murphy@ic.ac.uk). The data that support the findings of

415  this study are available from the corresponding author upon reasonable request.

**Experimental model and subject details**

**Mice**

Six- to twelve-week-old male and female C57bl/6 mice were used in studies, unless

otherwise stated. $Ntsr1^{\Delta NeoCre}$ (stock no. 033365) were bought in from Jax and breeding pairs

set up as hemizygotes x littermate control. $Ntsr1^{\Delta NeoCre}$ mice hemizygous for cre recombinase

are noted as *NTSR1-cre*. $Ntsr1^{\Delta NeoCre}$ mice without cre recombinase expression are notated as

*NTSR1-WT*. *Ntsr1-cre:tdTomato* mice were generated by crossing an *Ai9* female (stock no.

007909) with a *NTRS1-cre* male. The $NTSR1^{\Delta NeoCre}$ allele was detected by PCR genotyping

with primers 5'-GCTAGAGGGAAACCGTTGTG-3', 5'-GTTCTCCAGGAAGCCAAACA-

3' and 5'-GAGAGGGATAAGGGGCAAGA-3' where WT and mutant alleles produced PCR

products of 288 bp and 202 bp, respectively. The *Ai9* allele was detected by PCR genotyping

with primers 5'-AAGGGAGCTGCAGTGGAGTA-3', 5'-CCGAAAATCTCTGGGAAGTC-

3', 5'-GGCATTAAAGCAGCGTATCC-3' and 5'-CTGTTCCTGTACGGCATGG-3' where

WT and mutant alleles produced PCR products of 297 bp and 196 bp, respectively.

GLP-1R fl/fl mice were kindly gifted by Professor Randy Seeley at University of Michigan

and crossed with Pdx1-Cre/Esr1 mice (JAX strain: #024968). Tamoxifen (100mg/kg) was

injected intraperitoneally for 4-days to induce beta cell knockout of the GLP-1R.

**Primary Cell Culture**

Murine myenteric enteric neurons (MENs) were isolated from the proximal duodenum of

mice for use in activation studies. 6cm of proximal duodenum were dissected and

longitudinal muscle/myenteric plexus (LMMP) carefully peeled off and placed in ice-cold

digestion liquid (0.166mg/ml liberase in filtered Krebs-Ringer Bicarbonate Buffer (HKRB).

LMMP were next digested for 45 minutes at $37^{oC}$, washed in PBS and placed in complete

enteric neuronal media (Neurobasal A Media with B-27 supplement, 2mM L-glutamine, 10%

FBS, 1% penicillin/streptomycin and 10ng/ml GDNF). MENs were dispersed by trituration

using a p200 filter tip and pipette and filtered through a 70µM filter. Released myenteric

neurons were then resuspended in complete enteric neuronal media and maintained at $37^{oC}$,

5% $CO_2$ on laminin- and poly-D-lysine-coated slides.

Murine islets were isolated following injection of Collagenase NB 8 broad range (1mg/ml;

Universal Biologicals Ltd) into the bile duct of mice. Inflated pancreata were then digested at

$37^{oC}$ following which pancreata underwent a series of washes and manual digestions using

cold RPMI (RPMI 1640 medium; Merck, USA) and a Rotina 380R, Hettich Centrifuge. A

density gradient was next created using histopaque 1119 (Merck, USA), histopaque 1083

(Merck, USA) and RPMI medium no additives. Isolated islets were then picked using a

binocular microscope and maintained at $37^{oC}$, 5% $CO_2$ in complete media (RPMI media, 10%

FBS, 1% penicillin/streptomycin).

## Methods

### Animal Experiments

All mice were kept in individually ventilated cages with *ad libitum* access to water and standard laboratory RM1 chow in a temperate, humidity-controlled environment (21-23$^{oC}$) with constant 12h-light/dark cycle (lights on at 7am; lights off at 7pm), unless otherwise stated. Cages and bedding were changed once every week. To create a diet-induced obese model (DIO), mice were fed a high fat diet (60% fat, 20% protein, 20% carbohydrate; D12492, Research Diets, USA) from 6 weeks of age and studies were performed when they were aged 36 weeks. Mice were monitored daily for their health status. All procedures were carried out according to the Animals (Scientific Procedure) Act, 1986 and Amendment Regulations, 2012 and approved by the Imperial College London Animal Welfare and Ethical Review Body.

### Nodose Ganglion Surgery

6-week-old C5BL6/J mice were purchased from Charles River, U.K. and left to acclimatise for 1 week. Following this, mice were injected with either NTS-saporin [IT-56] or BLANK-saporin [IT-21] (1.5µg/µl; 0.5µl / nodose ganglia; Advanced Targeting Systems, CA) into the nodose ganglia. Injections were performed unilaterally with one week's rest before the second ganglion was injected. All injections were performed under isoflurane anaesthesia and mice were administered buprenorphine (0.12mg/kg) and carprofen (4mg/kg) as the perioperative analgesia. Mice were also administered atropine sulphate (2µg/kg) perioperatively to help prevent cardiac arrest caused by vagal manipulation during surgery. Postoperatively, mice were maintained on a liquid diet for 3 days (Ensure Plus Vanilla,

1.5kCal/ml) to minimise risk of unexpected gastrointestinal disturbances following vagal

manipulation and carprofen was added to their drinking water (2.72mg/100ml).

**Intracerebroventricular (ICV) Cannulation**

6-week-old C57bL6/J mice were purchased from Charles River, U.K. and left to acclimatise

for 1 week. Following this, mice were implanted with a stainless-steel cannula into the 3$^{rd}$

ventricle (-0.8mm anterior/posterior, +0.1mm lateral, -2.5mm dorsal/ventral). All

implantations were performed under isoflurane anaesthesia and mice were administered

buprenorphine (0.12mg/kg) and carprofen (4mg/kg) as the perioperative analgesia.

Postoperatively, carprofen was added to drinking water (2.72mg/100ml) for 3 days.

All injections (0.5µl/injection) were performed at a maximum infusion rate of 30µl/hour

using a locking internal cannula with 0.5mm projection in conscious animals. Cannula were

left indwelling for an additional minute post-injection to prevent backflow with withdrawal.

*In Vivo* **AAV Injections**

8-week-old *NTRS1-cre* and *NTSR1-WT* mice were injected with AAVrg-hSyn-DIO-

hM3D(Gq)-mCherry ($7x10^{12}$ vg/ml; 1µl/injection), AAV2-FLEX-DTR-GFP ($5x10^{12}$ vg/ml**;**

1µl/injection),  AAV2-pCAG-FLEX-EGFP-WPRE ($7x10^{12}$ vg/ml; 1µl/injection) or

AAV5.hSyn.eGFP.WPRE.bGH ($2.2x10^{13}$ vg/ml; 1µl/injection) in three separate areas in the

head of the pancreas. All injections were performed under isoflurane anaesthesia and mice

were administered buprenorphine (0.12mg/kg) and carprofen (4mg/kg) as the perioperative

analgesia. Postoperatively, carprofen was added to drinking water (2.72mg/100ml) for 3 days

and jelly infused with buprenorphine (0.1mg/ml) was placed in cages twice daily.

**Diphtheria Toxin-Mediated Ablation of NTSR1-Expressing Cells**

*NTRS1-cre* mice injected with AAV2-FLEX-DTR-GFP or AAV2-pCAG-FLEX-EGFP-WPRE and *NTSR1-WT* injected with AAV2-pCAG-FLEX-EGFP-WPRE received diphtheria toxin (30ng/g, Sigma, UK) via intraperitoneal injection once daily for two days. A week later, mice underwent intraperitoneal glucose tolerance tests.

**Surgical Separation of the Enteropancreatic Connection**

6-week-old C57bL6/J mice were purchased from Charles River and left to acclimatise for 1 week. To achieve surgical separation of the enteropancreatic network, the connection between the first cm of the proximal duodenum and adjacent pancreas was severed through blunt dissection using forceps, or left intact in control mice. Prior to separation, the pancreatic duct was located and care was taken to preserve it during surgery and all mice received pancreatic injection of AAV5.hSyn.eGFP.WPRE.bGH for model validation. All surgeries were performed under isoflurane anaesthesia and mice were administered buprenorphine (0.12mg/kg) and carprofen (4mg/kg) as perioperative analgesia. Postoperatively, carprofen was added to drinking water (2.72mg/100ml) for 3 days.

**Measurement of Olive Oil induced Neurotensin Secretion**

Mice were fasted for 5h with *ad libitum* access to water before being administered olive oil (200µl/mouse; Napolina) or control (dH20) by oral gavage. Mice were sacrificed 5 minutes after gavage and blood samples collected in EDTA-coated microvettes and spun at 3,600xg for 15 minutes. Plasma samples were then stored at $-80^{oC}$ until analysis by ELISA.

**Intraperitoneal Glucose Tolerance Tests**

Tests were performed on mice following a 5h fast with *ad libitum* access to water. Following

a baseline blood glucose measurement, mice were challenged with glucose (20%, 2g/kg) IP

and changes in blood glucose measured over the following 120 minutes. Blood glucose

readings were taken from a 0.5cm tail-tip venesection and measured using an AccuCheck

glucometer and glucose strips. Blood samples for plasma insulin and triglyceride analysis

were collected in EDTA-coated microvettes and spun at 3,600xg for 15 minutes. Plasma

samples were then stored at $-80^{oC}$ until analysed.

For assessment of the role of different types of oil on glucose tolerance, olive oil

(200µl/mouse; Napolina) and coconut oil (200µl/mouse; VITA COCA) or control (dH20)

were administered via oral gavage 15 minutes before glucose IP. The NTSR1 antagonist

SR48692 (80µg/kg, Tocris) or vehicle (1% DMSO) was administered intraperitoneally 60

minutes prior to glucose IP. Whey protein (10% w/v; 200µl/mouse; Pure Whey Isolate 90,

Bulk Powders, UK) or vehicle (50% PEG:dH20) was administered via oral gavage 15

minutes before glucose IP. Neurotensin acetate salt (6mg/kg, Bachem) or vehicle (dH20) was

administered intraperitoneally 15 minutes prior to glucose IP. Exendin-4 (0.6µg/kg, Wuxi

Apptec) was co-administered intraperitoneally with glucose. 4DAMP (1µg/kg) or vehicle

(1% DMSO) was administered intraperitoneally 15 minutes prior to glucose IP. Compound

21 (0.3mg/kg, HelloBio) or vehicle (saline) was administered intraperitoneally 15 minutes

prior to glucose IP. Orlistat (250mg/g of fat, Cambridge Biosciences), dissolved in olive oil,

or vehicles (olive oil, dH20) were administered via oral gavage 15 minutes prior to glucose IP

and AH-7614 (5mg/kg, Tocris) or vehicle (10% DMSO in corn oil) administered

subcutaneously 15 minutes prior to olive oil oral gavage.

**Fast-Refeeding Study**

Tests were performed on mice following a 16h fast with *ad libitum* access to water.

Following the fast, olive-oil enriched RM1 chow (1ml olive oil: 1 (~3g) pellet RM1 chow; 1

pellet / animal in cage) was given to the mice. Blood glucose readings were taken from a

0.5cm tail-tip venesection and measured using an AccuCheck glucometer and glucose strips

and change in blood glucose measured over the following 60 minutes. To investigate the role

of NTSR1, SR48692 (80µg/kg, Tocris) or vehicle (1% DMSO) were administered

intraperitoneally 60 minutes prior to the return of olive-oil enriched chow.

**Food Intake Study**

Tests were performed on individually caged mice following a 12h fast with *ad libitum* access

to water in the light phase. SR48692 (2.5µg; Tocris) or vehicle (1% DMSO; Sigma-Aldrich)

was administered ICV 20 minutes prior to intraperitoneal injection of neurotensin (6mg/kg,

Bachem) or vehicle (dH20). RM1 chow was weighed and returned to cages and food intake

measured over the following 1h.

**Core Temperature Study**

Tests were performed on mice following a 5h fast with *ad libitum* access to water. Rectal

temperature was taken using a RS PRO RS41 K handheld digital thermometer with external

K-type thermocouple bead probe and measured at baseline and 15 minutes post-treatment.

Neurotensin (6mg/kg, Bachem) or vehicle (dH20) were administered ICV into the 3[rd]

ventricle. Compound 21 (0.3mg/kg, HelloBio) or vehicle (saline) were administered

intraperitoneally.

**Metabolic Assays**

**Plasma Insulin Assay**

Plasma insulin concentration from collected plasma samples were assayed using the Mouse

Insulin ELISA (Mercodia, Sweden) according to the manufacturer's instructions.

**Plasma Triglyceride Assay**

Plasma triglyceride concentrations in collected plasma samples were assayed using the

Triglycerides TR210 Assay Kit (Randox, UK) according to the manufacturer's instructions.

**Plasma Neurotensin Assay**

Plasma neurotensin concentration from collected plasma samples were assayed using the

Mouse Neurotensin ELISA (E229Mo, BT Labs) according to the manufacturer's instructions.

**RNA Isolation and Gene Expression Analysis by qPCR**

**Preparation of Tissue Samples for RNA Extraction**

For the quantification of NTSR1 gene expression in different glucoregulatory tissues, punch

biopsies of arcuate nucleus were made using the optic chiasm and third ventricle as

landmarks and nodose ganglia were carefully dissected from mice and snap frozen at -80$^{oC}$.

Isolated islets and myenteric enteric neurons were isolated as described above and snap

frozen at -80$^{oC}$.

**RNA isolation**

Total RNA was isolated from tissues and cells using TRIsure (Bioline) and homogenised using a TissueLyser II (Qiagen, Hildon). Samples underwent a series of separation steps involving chloroform and centrifugation, following which RNA was precipitated and visualised using GlycoBlue (Invitrogen). The amount and purity of isolated RNA was assessed using a Nanodrop Lite Spectrophotometer (Thermofisher Scientific, Madison). Genomic DNA was removed from samples using RNase-free DNase (Promega) following which 1µg of RNA was reverse-transcribed using a High Capacity cDNA Reverse Transcription Kit (Applied Biosystems), according to the manufacturer's instructions.

**Quantitative Real-Time PCR (qPCR)**

qPCR was performed using Maxima Probe Master Mix (Life Technologies, Paisley) in a C1000 CFX384 Real-Time System Thermocycler using Taqman PCR core reagents (Applied Biosystems). To quantitatively analyse NTSR1 expression, the taqman probe for NTSR1 was used (assay ID: Mm00444459_m1; Applied Biosystems) and results analysed using the comparative $C_T$ method. Relative NTSR1 expression was normalised to the 'housekeeping' gene β-actin mRNA (assay ID: Mm02619580_g1; ThermoFisher).

**Tissue Clearing and Imaging**

NTSR1-cre:tdTomato mice were given an overdose of pentobarbital and transcardially perfused with 0.01M PBS followed by 4% PFA. The first cm of the proximal duodenum and its adjacent pancreatic section were dissected out and fixed overnight in 4% PFA at 4°C. Tissues were cleared using RTF tissue clearing[1] to retain endogenous fluorescence. Briefly, tissue was sequentially immersed in 3 different solutions constituted by triethanolamine

(TEA) (Sigma) and formamide (F) (ThermoFisher) and dH20 (W); RTF-R1

(30%TEA/40%F/30%W), RTF-R2 (60%TEA/25%F/15%W), RTF-R3

(70%TEA/15%F/15%W) over 1 day. The cleared sample was then imaged on an inverted

confocal Leica Stellaris 8 microscope (Facility for Imaging by Light Microscopy, Imperial

College London) and further processed in ImageJ. iDISCO whole-mount immunolabelling[2]

without methanol pre-treatment was conducted prior to tissue clearing and imaging. To label

insulin-secreting beta cells within the pancreas, a guinea pig anti-insulin antibody was used

(1:100; ab7842, Abcam) and immunofluorescence was identified using a goat anti-guinea pig

secondary IgG antibody conjugated to Alexa-Fluor 647 (1:500; ThermoFisher).

**LMMP Immunohistochemistry**

NTSR1-cre:tdTomato mice injected with AAV2-FLEX-DTR-GFP,

AAV5.hSyn.eGFP.WPRE.bGH or AAV2-pCAG-FLEX-EGFP-WPRE-injected mice were

killed by cervical dislocation and the LMMP of the proximal duodenum carefully dissected

out. LMMP was fixed overnight in 4% paraformaldehyde (PFA, Sigma-Aldrich, UK). To

identify NTSR1-expression in the myenteric plexus, LMMP were stained with the pan-

neuronal marker polyclonal rabbit anti-beta tubulin III (1:200; ab18207; Abcam, UK) or

mouse anti-beta tubulin III (1:500; ab78078; Abcam). All samples were blocked in 0.1M PBS

with 4% neonatal goat serum (New Zealand Origin; Gibco), 4% bovine albumin serum (BSA,

Sigma-Aldrich) and 0.5% triton-X (Sigma-Aldrich) prior to primary antibody, and then

overnight at $4^{oC}$. Immunofluorescence was identified using a goat anti-rabbit secondary IgG

antibody conjugated to AlexaFluor-488 (1:500; ThermoFisher), goat anti-mouse secondary

IgG antibody conjugated to AlexaFluor-488 (1:500; ThermoFisher) or goat anti-rabbit

secondary IgG antibody conjugated to AlexaFluor-633 (1:500; ThermoFisher). Nuclei were

stained using DAPI mounting media (ProLong™ Diamond Antifade Mountant with DAPI;

ThermoFisher) and images acquired using a Zeiss AxioObserver widefield microscope with

x5, x10 or x20 objective lens and analysed in ImageJ.

The neuronal nature of cultured myenteric enteric neurons was confirmed using the technique

as described above.

**Verification of Surgical Separation of the Proximal Duodenum and the Pancreas Model**

To validate enteropancreatic separation, the percentage eGFP reflecting virus

(intrapancreatically injected AAV5.hSyn.eGFP.WPRE.bGH) expression within total neurons

in the LMMP was quantified and compared to control mice. Dissected LMMPs were stained

with anti-beta tubulin III and imaged at 20x magnification. In ImageJ, a 'neuronal' (beta-

tubulin III-expressing) mask and a 'viral' (eGFP-expressing) mask was created. The masks

were overlaid and the percentage virus within the neuronal mask quantified. Thresholding

was kept consistent across channels between mice.  Each N value reflects quantification of 2-

8 averaged image quantifications for each mouse.

**Verification of AAV-DTR Ablation Model**

To validate ablation of the NTSR1-expressing enteropancreatic cells, the percentage

tdTomato, reflecting NTSR1, expressing neurons, within total neurons in the LMMP was

quantified and compared between *Ntsr1-cre:tdTomato* mice intrapancreatically injected with

AAV2-FLEX-DTR-GFP or AAV2-pCAG-FLEX-EGFP-WPRE. Dissected LMMPs were

stained with anti-beta tubulin III and imaged at 10x magnification. In ImageJ, a 'neuronal'

(beta-tubulin III-expressing) mask and a 'tdTomato' (eGFP-expressing) mask was created.

The masks were overlaid and the percentage tdTomato within the neuronal mask quantified.

Thresholding was kept consistent across channels between mice.  Each N value reflects

quantification of 2-6 averaged image quantifications for each mouse.

**Calcium activation studies**

**Signalling Assay**

To investigate treatment-stimulated cell activation, myenteric enteric neurons were cultured

as described above and imaged on day 5 when neuronal projections had begun to form.

Changes in intracellular calcium levels were measured using Fluo-4AM Direct labelling

(Invitrogen), according to manufacturer's protocol. Agonist-dependent calcium mobilization

was assessed using a Zeiss AxioObserver widefield microscope with a x20 objective lens,

capturing 1 frame per second.

Cells were incubated in Fluo-4 AM Direct at $37^{oC}$ for 30 minutes, followed by a 30-minute

incubation at room temperature. Baseline readings were then taken for 60 seconds prior to

addition of treatment in a 1:5 dilution (200pM or 100nM neurotensin, Bachem). SR48692

(10µM; Tocris) was added 20 minutes prior to baseline readings. KCl (50mM) was added 5

minutes post-treatment to confirm cell viability.

**Calcium Imaging Analysis**

Images were analysed using ImageJ and a macro package from Imperial FILM facility

(Intensity_2). Regions of interest (ROI) were drawn around all obvious cell bodies to capture

fluorescence (f). Average fluorescence for baseline readings of each cell were calculated,

using which fluorescent change for each cell was quantified (fluorescence at every frame –

baseline average / baseline = $\Delta f/f$). Max $\Delta f/f$ over the 2.5-minute period following addition of

treatment was calculated and cells stratified into responders and non-responders (responder:

max Δf/f of treatment > max Δf/f of baseline + 2x SD; Non-responder: max Δf/f of ligand

< max Δf/f of baseline + 2x SD).

**Quantification and Statistical Analysis**

Data were analysed using Prism 9.0 software (GraphPad Software) and expressed as mean ±

SEM. Analysis of data with one independent variable was analysed with one-way ANOVA

with Tukey's multiple comparison post hoc test. Data with only two independent variables

alongside a continuous outcome variable were analysed using a two-way ANOVA with

Sidak's multiple comparison post hoc test. In studies comparing three or more groups,

Tukey's multiple comparison post hoc test was used. Statistical analysis of area under the

curve data sets was conducted using a student's t-test (two-sided). p values <0.05 were

considered statistically significant.

**Method References**

704　1.　Yu T, Zhu J, Li Y, et al. RTF: a rapid and versatile tissue optical clearing method. *Sci*

705　*Rep*. 01 31 2018;8(1):1964. doi:10.1038/s41598-018-20306-3

706　2.　Renier N, Wu Z, Simon DJ, Yang J, Ariel P, Tessier-Lavigne M. iDISCO: a simple,

707　rapid method to immunolabel large tissue samples for volume imaging. *Cell*. Nov 06

708　2014;159(4):896-910. doi:10.1016/j.cell.2014.10.010

**Acknowledgements**

The Section of Endocrinology and Investigative Medicine is funded by grants from the MRC,

BBSRC, NIHR and is supported by the NIHR Biomedical Research Centre Funding Scheme

and the NIHR/Imperial Clinical Research Facility. The views expressed are those of the

authors and not necessarily those of the MRC, BBSRC, the NHS, the NIHR or the

Department of Health. KGM is supported by Diabetes UK (18/0005886, 20/0006295), the

BBSRC (BB/W001497/1, BB/X017273/1), the MRC (MR/Y013980/1) and the Wellcome

Trust (310835/Z/24/Z). BYHL and GSHY are supported by BBSRC Project Grant

(BB/S017593/1) and the MRC Metabolic Diseases Unit (MC_UU_00014/1). Graphical

abstract, Fig. 3A, 4A&D, Extended Data Fig. 2B, 3A, 4A, 5A, 7A were created using

BioRender (https://www.biorender.com). We thank Bryn Owen for his comments on the

manuscript.

**Author Contributions**

724    A.G.R., L.M., M.N., P.P., A.M-A., G.D., S.C., C.D., J.L., Y.T., T.S-C., A.B.D., A.C.,

725    K.G.M. performed experiments, analysed data and/or interpreted data. B.L., A.H., B.J., G.Y.,

726    V.S., K.G.M. contributed specific reagents or methods, interpreted data and provided input

727    on the experiments. A.G.R., L.M., K.G.M. wrote the manuscript with input from all co-

728    authors. K.G.M. directed and supervised the study.

**Competing Interests Declaration**

731    BYHL provides remunerated consultancy for Nuntius Therapeutics. GSHY receives grant

732    funding from Novo Nordisk and Amgen Inc; he also consults for both Novo Nordisk and Eli

733    Lilly and Company. The other authors declare no competing interests.

**Fig. 1: Olive oil improves glucose tolerance via activation of NTSR1.** A) Intraperitoneal glucose tolerance test (IPGTT) with olive oil or vehicle orally gavaged (OG)-15 minutes prior to glucose load. n=5/group. B). Plasma insulin levels measured using Mouse Insulin ELISA following olive oil or vehicle OG administration and intraperitoneal glucose load. n=8-9/group. C). IPGTT with olive oil, olive oil and orlistat or vehicle OG-15 minutes prior to glucose load. n=4,5,6/group. D). IPGTT with oleic acid or vehicle OG- 15 minutes prior to glucose load. n=10/group. E). IPGTT with subcutaneous administration of FFAR4 antagonist or vehicle 15 minutes prior to olive oil or vehicle OG-15 minutes prior to IP glucose load. n=11-15/group. F). Plasma neurotensin levels measured by ELISA 2 minutes following OG olive oil or vehicle. n=4/group. G). IPGTT with 1h pretreatment of an NTSR1-specific

antagonist IP followed by olive oil or vehicle OG. n=7-8/group. H). Fast-refeed study with 1h pretreatment of an NTSR1-specific antagonist IP. Mice were refed olive oil-enriched chow at t0 and blood glucose monitored for 1h. n =7-8/group. I). IPGTT with intraperitoneal (IP) neurotensin (0.2mg/kg – 6mg/kg) or vehicle 15 minutes prior to glucose load in lean mice. n=5-6/group. J). IPGTT with IP neurotensin or vehicle 15 minutes prior to glucose load in DIO mice. n=8/group. K). Plasma insulin levels measured using Mouse Insulin ELISA following peripheral neurotensin or vehicle administration and intraperitoneal glucose. n=9-10/group. L). IPGTT with IP 15-minute pretreatment of an M3R-specific antagonist or vehicle followed by IP of either neurotensin or vehicle. n=5/group. M). IPGTT with 15-minute pretreatment of an M3R-specific or vehicle IP followed by OG olive oil or vehicle n=5/group. All data analysed with either 1-way or 2-way ANOVA with Tukey's post-hoc analysis test. * indicates significance between different treatment groups, # indicates significance between olive oil and olive oil + orlistat (Fig.1C), $ indicates significance between vehicle + olive oil and FFAR-antagonist (Fig.1E) + olive oil or vehicle + olive oil and NTSR1-antagonist + olive oil (Fig.1G), ££ indicates significance between vehicle + NTS and M3R-antagonist + NTS (Fig.1L) or vehicle + olive oil and M3R-antagonist + olive oil (Fig.1M). In Fig 1I, £ indicates significance between 0.2mg/kg NTS and vehicle, $ indicates significance between 0.6mg/kg NTS and vehicle, # indicates significance between 2mg/kg NTS and vehicle, *indicates significance between 6mg/kg NTS and vehicle.

**Figure 2. NTSR1 is expressed in enteropancreatic neurons.** A). NTSR1 and beta-III tubulin expression in LMMP of the proximal duodenum. B). qPCR data showing relative NTSR1 expression in the murine arcuate nucleus (ARC), nodose ganglia (NG), primary myenteric enteric neurons (MENs) and isolated islets (ISLETS). n=5-6. C). Representative traces of Fura-4 AM-loaded myenteric enteric neurons (MENs) treated with either vehicle (HKRB), 100nM neurotensin or 10μM SR48692 + 100nM neurotensin. D). Maximum change in fluorescence measured in the 2.5 minutes following treatment stimulation. 3x technical replicates over 3x biological replicates. E). Expression of tdTomato in the NTSR1-cre:tdTomato mouse in the enteropancreatic connection at the proximal duodenum in cleared tissue. F). Expression of tdTomato and insulin expression in the pancreas adjacent to the proximal duodenum in a NTSR1-cre:tdTomato mouse. G). Expression of eGFP and beta-III tubulin in the LMMP of the proximal duodenum of mice injected into the pancreas with AAV5.hSyn.eGFP.WPRE.bGH . All scale bars are 100μm. Change in fluorescence data analysed using 1-way ANOVA with Tukey's post -hoc analysis.

**Fig. 3: The enteropancreatic axis is necessary for the glucoregulatory effects of neurotensin.** Intraperitoneal glucose tolerance tests (IPGTT) were performed in mice that had either undergone surgical separation of the connection between the proximal duodenum and pancreas, or sham surgery. A). Schematic depicting surgical separation of proximal duodenum and pancreas in mice. B). IPGTT with olive oil or vehicle oral gavaged (OG)-15 minutes prior to glucose load in mice with an intact connection between the proximal duodenum and pancreas. n=5/group. C). IPGTT with olive oil or vehicle OG-15 minutes prior to glucose load in mice with a severed connection between the proximal duodenum and pancreas. n=8/group. D). IPGTT with intraperitoneal (IP) neurotensin or vehicle 15 minutes prior to glucose load in mice with an intact connection between the proximal duodenum and pancreas. n=8/group. E). IPGTT with IP neurotensin or vehicle 15 minutes prior to glucose load in in mice with severed connection between the proximal duodenum and pancreas. n=8/group. F). Fast-refeed study with mice with an intact or severed connection between the proximal duodenum and pancreas. Mice were refed olive-oil enriched chow at t0 and blood glucose monitored for 1h. n=8/group. All IPGTT data were analysed with 2-way ANOVA with Tukey's post-hoc analysis test. AUC data were analysed with paired t-test.

**Fig. 4: Enteropancreatic-NTSR1 neurons are sufficient and necessary for the effects of olive oil and neurotensin on glucose tolerance.** *NTSR1-WT* and *NTSR1-cre* mice were injected with AAVrg -hSyn-DIO-hM3D(q) or of AAV2 -FLEX-DTR-GFP into the pancreas and intraperitoneal glucose tolerance tests (IPGTT) performed two weeks later or one week after injection of diphtheria toxin, respectively. A). Schematic depicting injection of AAVrg -hSyn-DIO-hM3D(q) into the pancreas. B). IPGTT with intraperitoneal (IP) vehicle or C21 IP 15 minutes prior to glucose load in *NTSR1-WT* mice. n=4/group. C). IPGTT with IP C21 or vehicle 15 minutes prior to glucose load in *NTSR1-cre* mice. n=17/group. D). Schematic depicting injection of AAV2 -FLEX-DTR-GFP into the pancreas. E). IPGTT with oral gavaged (OG) olive oil or vehicle 15 minutes prior to glucose load in *NTSR1-cre* mice injected with AAV2 -pCAG-FLEX-EGFP-WPRE. n=10/group. F). IPGTT with OG olive oil or vehicle 15 minutes prior to glucose load in *NTSR1-cre* mice injected with AAV2 -FLEX-DTR-GF. n=10/group. G). Fast -refeed study with *NTSR1-cre* mice injected with either

AAV2 -pCAG-FLEX-EGFP-WPRE or AAV2 -FLEX-DTR-GFP. Mice were refed olive -oil enriched chow at t0 and blood glucose monitored for 1h. n =11-12/group. H). IPGTT with IP neurotensin or vehicle 15 minutes before glucose load in *NTSR1-cre* mice injected with AAV2 -pCAG-FLEX-EGFP-WPRE. n=10/group. I). IPGTT with IP neurotensin or vehicle 15 minutes before glucose load in *NTSR1-cre* mice injected with AAV2 -FLEX-DTR-GFP. n=10/group. J). Plasma insulin levels measured 30 minutes post-IP neurotensin or vehicle administration and 15 minutes post-intraperitoneal glucose load (t30 on IPGTT) in *NTSR1-cre* mice injected with AAV2 -pCAG-FLEX-EGFP-WPRE. n=6/group. K). Plasma insulin levels measured 30 minutes post-IP neurotensin or vehicle administration and 15 minutes post-intraperitoneal glucose load (t15 on IPGTT) in *NTSR1-cre* mice injected with AAV2 -FLEX-DTR-GFP. n=6/group. All IPGTT and fast-refeed data was analysed with 2-way ANOVA with Sidak's post-hoc analysis test and embedded AUC data were analysed with paired t-test. Plasma insulin was analysed with a paired t-test.

**Extended Data Fig. 1: The glucoregulatory effects of olive oil are not replicated by coconut oil and blocking the NTSR1 does not prevent the glucoregulatory effects of another macronutrient.** A). Oral glucose tolerance test with IP neurotensin or vehicle 15 minutes prior to glucose load. n=8/group. B). Intraperitoneal glucose tolerance test (IPGTT) with olive oil, coconut oil or vehicle orally gavaged (OG)-15 minutes prior to glucose load. n=5/group. C). IPGTT with 1h IP pretreatment of an NTSR1-specific antagonist followed by OG whey protein (10% weight/vol) or vehicle. n=7-8/group. D). Serum triglyceride levels measured 15 minutes post-IP neurotensin or vehicle administration and intraperitoneal glucose load. n=3/group. E). Serum triglyceride levels measured 15 minutes post-OG olive oil or vehicle administration and intraperitoneal glucose load. n=6-7/group. All IPGTT data was analysed with 2-way ANOVA with Sidak's post-hoc analysis test. # indicates significance between vehicle and olive oil, $ indicates significance between vehicle + vehicle and vehicle + protein.

**Extended Data Fig. 2: The glucoregulatory effects of neurotensin and olive oil are not mediated by the GLP-1 receptor.** A). Intraperitoneal glucose tolerance test (IPGTT) with intraperitoneal (IP) neurotensin, exendin-4 (Ex-4) or neurotensin + Ex-4 15 minutes prior to glucose load. n=8/group. B). Schematic of tamoxifen-induced knock out of GLP-1R from pancreatic beta cells using Pdx1CreERT-GLP-1Rfl/fl mice. C). IPGTT in Pdx1CreERT-GLP-1Rfl/fl mice with Ex-4 or vehicle 15 minutes prior to glucose load pre-tamoxifen induction. n= 6-/group. D). IPGTT in Pdx1CreERT-GLP-1Rfl/fl mice with Ex-4 or vehicle 15 minutes prior to glucose load post-tamoxifen induction. n= 6-7/group. E). IPGTT in Pdx1CreERT-GLP-1Rfl/fl mice with IP neurotensin or vehicle 15 minutes prior to glucose load pre-tamoxifen induction. n= 9/group. F). IPGTT in Pdx1CreERT-GLP-1Rfl/fl mice with IP neurotensin or vehicle 15 minutes prior to glucose load post-tamoxifen induction. n= 16/group. G) IPGTT in PDX4 mice with olive oil or vehicle 15 minutes prior to glucose load pre-tamoxifen induction. n= 5-7/group. H). IPGTT in Pdx1CreERT-GLP-1Rfl/fl mice with olive oil or vehicle 15 minutes prior to glucose load post-tamoxifen induction. n= 5-8/group. All IPGTT data was analysed with 2-way ANOVA with Sidak's post-hoc analysis test. # indicates significance between vehicle + vehicle and vehicle + Ex-4, $ indicates significance between vehicle + vehicle and NTS + Ex-4.

**Extended Data Fig.3: Neurotensin does not improve glucose tolerance via activation of vagal-NTSR1.** A). Schematic depicting bilateral injection of NTS-saporin or BLANK-saporin into the vagal nodose ganglia of C57BL6/J mice. B). qPCR analysis of relative NTSR1 expression in the vagal nodose ganglia of NTS- and BLANK-saporin injected mice. n=4-5/group. C). IPGTT with 15-minute pretreatment of IP neurotensin or vehicle 15 minutes prior to glucose load in NTS- and BLANK-saporin injected mice. n=4-6/group. D). IPGTT with 15-minute oral gavage pretreatment of olive oil or vehicle 15 minutes prior to glucose load in NTS- and BLANK-saporin injected mice. n=4-6/group. qPCR data analysed using unpaired t-test. IPGTT data analysed using 2-way ANOVA with Tukey's post -hoc analysis. # indicates significance between NTS-SAP + vehicle and NTS-SAP + NTS or NTS-SAP + vehicle and NTS-SAP + olive oil.

**Extended Data Fig. 4: Neurotensin does not improve glucose tolerance via activation of central-NTSR1.** A). Schematic depicting cannulation into the 3rd ventricle of C57BL6/J mice. B). Change in core temperature following central infusion of neurotensin or vehicle into the 3rd ventricle. n=5/group. C). Intraperitoneal glucose tolerance test (IPGTT) with low dose (10ng) central administration of neurotensin or vehicle 5 minutes prior to glucose load. n=8-10/group. D). IPGTT with high dose (400ng) central neurotensin or vehicle 5 minutes prior to glucose load. n=8-10/group. E) Following a 12h fast, mice were centrally injected with an NTSR1 antagonist (SR48692, 2.5µg/mouse) or vehicle followed by peripheral neurotensin (6mg/kg) or vehicle. Food was returned after 10 minutes and food intake was measured 1h later. F). IPGTT with 1h pretreatment of central NTSR1-specific antagonist or vehicle followed by intraperitoneal neurotensin or vehicle 15 minutes prior to glucose load. n=4-5/group. Change in core temperature was analysed using an unpaired student's t-test. Food intake and IPGTT data were analysed using a 2-way ANOVA with Tukey's post-hoc analysis. # indicates significance between NTSR1-antagonist ICV + vehicle IP and NTSR1-antagonist ICV and NTS IP.

**Extended Data Fig. 5: Enteropancreatic separation model validation.** A). Schematic depicting surgical separation of proximal duodenum and pancreas in mice. B). Quantification of separation of the proximal duodenum and pancreas expressed as percentage eGFP expression within neurons in the LMMP of the proximal duodenum. C). Intraperitoneal glucose tolerance test (IPGTT) with 15-minute intraperitoneal (IP) pretreatment of Ex-4 or vehicle in mice with an intact connection between the proximal duodenum and pancreas. n=6/group. D). IPGTT with Ex-4 or vehicle in mice with a severed connection between the proximal duodenum and pancreas. n=7/group. E). IPGTT with 15-minute pretreatment of oral gavaged (OG) whey protein (10% weight/vol) or vehicle in mice with an intact connection between the proximal duodenum and pancreas. n=12/group. F). IPGTT with 15-minute pretreatment of OG 10% whey or vehicle in mice with a severed connection between the proximal duodenum and pancreas. n=8-10/group. Percentage area EGFP data was analysed using unpaired t-test. All IPGTT data were analysed using a 2-way ANOVA with Tukey's post-hoc analysis and embedded AUC data were analysed using paired t-test.

**Extended Data Fig. 6. Enteropancreatic chemogenetic model validation.** *NTSR1-WT* and *NTSR1-cre* mice were injected with AAVrg -hSyn-DIO-hM3D(q) into the pancreas. A). Plasma insulin levels measured using Mouse Insulin ELISA 15 minutes post-IP C21 or vehicle a (t0 on IPGTT). n= 13/group. B) Plasma insulin levels measured using Mouse Insulin ELISA 30 minutes post-IP C21 or vehicle administration and 15 minutes post-IP glucose load (t15 on IPGTT). n= 13/group. C). No change in core temperature following vehicle or C21 IP in *NTSR1-WT:Dq-DREADD* and *NTSR1-cre:Dq-DREADD*. n=3-4/group. Insulin data analysed using paired t-test. Change in core temperature between groups analysed using a 1-way ANOVA.

**Extended Data Fig. 7. Enteropancreatic DTR model validation.** A). Schematic depicting injection of AAV2 -FLEX-DTR-GRP into the pancreas. B). Quantification of ablation of NTSR1-expressing neurons expressed as percentage tdTomato expression within neurons in the LMMP of the proximal. C). IPGTT with intraperitoneal (IP) Ex-4 or vehicle 15 minutes prior to glucose load in *NTSR1-WT* mice injected with AAV2 -FLEX-DTR-GFP. n=6/group. D). IPGTT with IP Ex-4 or vehicle 15 minutes prior to glucose load in *NTSR1-cre* mice injected with AAV2 -pCAG-FLEX-EGFP-WPRE (viral control). n=12/group. E). IPGTT with

IP Ex-4 or vehicle 15 minutes prior to glucose load *in NTSR1-cre* mice injected with AAV2 -FLEX-DTR-GFP. n=10/group. F). IPGTT with oral gavaged (OG) 10% whey or vehicle 15 minutes prior to glucose load in *NTSR1-WT* mice injected with AAV2 -FLEX-DTR-GFP. n=7/group. G). IPGTT with OG 10% whey or vehicle 15 minutes prior to glucose load in *NTSR1-cre* mice injected with AAV2 -pCAG-FLEX-EGFP-WPRE. n=12/group. H). IPGTT with OG 10% whey or vehicle 15 minutes prior to glucose load in *in NTSR1-cre* mice injected with AAV2 -FLEX-DTR-GFP. n=11/group. I). IPGTT with olive oil or vehicle OG 15 minutes prior to glucose load in *NTSR1-WT* mice injected with AAV2 -FLEX-DTR-GFP. n=7/group. J). IPGTT with IP neurotensin or vehicle 15 minutes prior to glucose load in *NTSR1-WT* mice injected with AAV2 -FLEX-DTR-GFP. n=7/group. All IPGTT data were analysed using a 2-way ANOVA with Tukey's post -hoc analysis and embedded AUC data were analysed using paired t-test.