Modelling PRRS transmission between pig herds in Denmark and prediction of interventions impact

You Chang[1], Ana Rita Pinherio Marques[1], Mette Fertner[2], Nils Toft[3], Bjørn Lorenzen[3], Mossa Merhi Reimert[1], Hans Houe[1], Beate Conrady[1]

[1] Department of Veterinary and Animal Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, Grønnegårdsvej 8, DK-1870 Frederiksberg C, Denmark

[2] Livestock Innovation, SEGES Innovation P/S, Meldahlsgade 3, DK-1613 Copenhagen V, Denmark

[3] Danish Agriculture & Food Council, Axeltorv 3, 1609 København V

Corresponding author: You Chang; you.chang@sund.ku.dk

Highlights (85 characters):

- We generated the first between-herd R_0 map of PRRS transmission in Denmark

- 17% of undetected herds are responsible for 60% of total PRRS transmission

- Positive stable herds are 50 times less infectious than positive unstable herds

- Local spread drives the spatial heterogeneity of PRRS transmission

- Eradication requires multiple controls e.g., local controls, stricter risk-based trading and more frequent testing

Abstract

Porcine reproductive and respiratory syndrome (PRRS) is an endemic viral disease in most pig-producing countries, including Denmark. In 2022, Denmark launched a control program to reduce PRRS prevalence, with legislative changes in 2023 making testing and status reporting mandatory. The program also enforces the loss of PRRS-free status for farms that purchase pigs from non-PRRS-free sources and implements region-specific control measures to coordinate PRRS elimination within herds. However, the effectiveness of these interventions remains uncertain and requires thorough evaluation through transmission modelling and analysis of data before and after legislation changes.

To understand PRRS transmission prior to legislative changes in 2023 and predict the impact of control measures, we developed a between-herd stochastic compartmental model. This model includes compartments for susceptible (S), highly infectious (I_h), lowly infectious

(I_l) and detected (D) pig herds, using data from 2020-2021. The model (i) quantifies the relative

contributions of pig movements and local transmission to the spread of PRRS; (ii) generate

herd-level maps of the basic reproduction (R_0); and (iii) assess the effectiveness of targeted

interventions for eradicating PRRS in Denmark.

Model results indicated that more than 50% of herds had an R_0 greater than 1,

suggesting a potential for sustained transmission if no interventions had been implemented

after 2022. Both local spread and movement-mediated transmission play important roles, but

local transmission drives the spatial heterogeneity in observed PRRS prevalence across

Denmark. Although only 17% of infectious herds remain undetected under current

surveillance, they are responsible for 60% of total transmission. Local control via depopulation

and repopulation, is the fastest measure to reduce the observed prevalence of PRRS, but it has

a lower effect on true transmission due to the hidden infections. Therefore, achieving

eradication may require a combination of more frequent testing, targeted within-herd PRRS

elimination and stricter risk-based trading. This study identifies PRRS hotspots and

transmission routes, offering evidence-based recommendations for control.

1. Introduction
55
Porcine reproductive and respiratory syndrome (PRRS) is a viral disease that affects pig

populations worldwide (Lunney et al., 2010). The disease causes significant production losses

due to reproductive failure in sows, respiratory distress in younger pigs, and increased mortality

rates (OIE, 2008). PRRS is estimated to cause production losses ranging between €4 to €139

per sow per year in Denmark (Rathkjen & Dall, 2017), €126 per sow in the Netherlands

(Nieuwenhuis et al., 2012) and $664 million in losses annually for the pig industry in the U.S.

(Holtkamp et al., 2013). The infection spreads through multiple routes, including animal

movement, indirect contacts such as contaminated materials, and airborne transmission (Cho

& Dee, 2006; Corzo et al., 2010; Rahe & Murtaugh, 2017; Rowland & Morrison, 2012; Valdes-

Donoso et al., 2018). The etiological agent of PRRS is an RNA virus, the porcine reproductive

and respiratory syndrome virus (PRRSV), which is divided into type 1 and type 2 genotypes,

known as European and North American strains, respectively (Murtaugh et al., 1995; Rahe &

Murtaugh, 2017).

Denmark has faced endemic PRRS transmission since its first detection in March 1992

(Bøtner et al., 1994). The infections were widespread by mid-1990s, with a herd-level

prevalence of approximately 25% in sow herds and 33% in finisher herds, with an annual herd-

level incidence of 5-10% (Mortensen, 1996; Mortensen et al., 2002; Mortensen &
Strandbygaard, 1996). Denmark's PRRS control efforts have evolved over time, with the
Specific Pathogen Free (SPF) system forming the foundation of PRRS management since 1993
(Fertner et al., 2025). The SPF system is an industry-driven voluntary system. Enrolled herds
are routinely tested for the absence of seven pathogens, including PRRS, as part of the health
declaration system. To sustain the SPF status of herds, herds must undergo annual biosecurity
evaluations and follow a certain set of rules regarding biosecurity (Lopes Antunes et al., 2019;
Conrady et al., 2023; SPF, 2024), including investments in changing farm facilities, routine
serological testing, herd status classification, and restrictions on animal movement. Despite of
these efforts, national PRRS prevalence remained around 35% in 2020 (Kristensen et al., 2020;
Sundhedsstyringen, 2020).

In May 2022, Denmark launched national PRRS control program that extended PRRS
control beyond the SPF program. A key legislative change came in 2023 when a government
order was issued, obligating all herds to officially declare their PRRS status (Danish Ministry
of Food, Agriculture and Fisheries, 2023; State Serum Institut, 2023). In addition, local control
measures have been introduced, encouraging herd owners to collaborate and coordinate actions
to eliminate PRRS in positive herds within specific local areas. The final decision on whether,
and when, to eliminate PRRS within a herd lies with the herd owners. These decisions can be
influenced by within-herd outbreak size and economic considerations, rather than a compulsory
standard control. While rolling out these initiatives, the effectiveness of interventions after new
legislation remains uncertain.

To predict the effectiveness of interventions targeting different transmission routes, it
is essential to understand the relative contributions of each route, including pig movements,
and local spread. Local spread occurs when virus is transmitted through contact with virus-
laden fomites (such as shared equipment, vehicles, and workers) or virus-laden aerosols in the
vicinity of an infectious herd, often referred to as airborne transmission. Several mechanistic
modeling studies in the U.S. (Galvis, Corzo, & Machado, 2022; Galvis, Corzo, Prada, et al.,
2022) and Canada (Thakur, Revie, et al., 2015; Thakur, Sanchez, et al., 2015) have highlighted
the importance of these routes in the between-herd spread. Herd-level risk factor analyses have
identified regional swine density and the purchase of PRRSV-positive animals as primary risk
factors (Baysinger et al., 1997; Firkins & Weigel, 2004; Mortensen et al., 2002; Velasova et
al., 2012). However, these findings cannot be directly applied to other regions or situations
where interventions alter disease dynamics. The relative contribution of local spread and
movement may vary between countries due to demographic and environmental heterogeneity,

such as livestock density, frequencies of livestock movements, herd size, farm biosecurity
standards, and farm structures (Conrady et al., 2023, 2024). Therefore, a mechanistic model
specific to Denmark is needed to understand transmission and predict the impact of new
legislative changes (after 2023) on PRRS control.

The present study aims to develop a stochastic compartmental model to *(i)* generate
herd-level maps of the basic reproduction ratio (R_0); *(ii)* quantify the relative contributions of
pig movements and local transmission to the spread of PRRS; and *(iii)* assess the effectiveness
of targeted interventions for PRRS control in Denmark.

2. Material and method

We developed a stochastic between-herd compartmental model to represent the PRRS
transmission dynamics using the SimInf package (Widgren et al., 2016; Widgren et al., 2019).
The data were described in Section 2.1, followed by the model description in Section 2.2. In
addition, parameters were explained in Section 2.3, and statistical method was described in
Section 2.4, and the calculation of the basic reproduction ratio (R0) in Section 2.5. In Section
2.6, uncertainty in the parameters was investigated by sensitivity analysis, and in Section 2.7,
several interventions that target different transmission routes were explained.

2.1. Data

Data used in this study covered the period from January 1, 2020, to December 31, 2021, and
consisted of 4 main datasets: (1) herd-level PRRS status and biosecurity level extracted from
the Danish SPF health dataset; (2) production type; (3) the number of moved pigs between
premises considered as discrete herd-to-herd pairs and (4) geographic information of the herd
locations (Figure 1).

Figure 1. Example of four synthetic datasets, illustrating data structures used in this study.

The study population consists of herds with reported PRRS status during the study period (2021-2022). In total, 3,420 herds were included, representing approximately 60% of all pig herds in Denmark during this period. For each herd, we extract the associated herd identification number, biosecurity level, and the PRRS status.

In SPF system, herds' biosecurity level is classified into green, blue and red level and non-SPF herds (Lopes Antunes et al., 2015):

1. Red herds: These herds undergo monthly testing and demonstrate a high commitment to eliminating PRRS in the event of an outbreak.

2. Blue and green herds: These herds undergo annual testing and show a moderate commitment to achieving PRRS elimination.

3. Non-SPF herds: These herds have the lowest biosecurity measures and typically do not report their status. However, some non-SPF herds had data entries in the surveillance system, which were also included in this study.

For each testing, samples from 10 animals (5 gilts and 5 sows) from red herds and 20 animals from blue herds (selected randomly within the herd) are collected and tested by serological tests. The current SPF health system automatically assigns a PRRS-positive status to herds receiving pigs from PRRS-positive herds without additional testing. While this system is designed to reduce risk, it increases the proportion of false positive test results in the official control system.

2.2 Model description

A stochastic between-herd compartmental model was developed using R package SimInf (Widgren et al., 2016; Widgren et al., 2019). In this model, herds are considered as epidemiological units and are classified into four compartments (Figure 2): Susceptible (S), highly infectious (I_h), lowly infectious (I_l) and detected (D). When a susceptible herd is introduced to PRRSV, it typically enters an acute epidemic phase, during which most pigs become infected in a short period. This phase is referred to as the highly infectious herds, characterized by significant viral shedding and a high risk of infecting other herds. This corresponds to "positive unstable" herds in PRRS terminology (Holtkamp et al., 2011). Over time, typically after approximately three months (expert's opinion), the infection stabilizes and becomes endemic within the herd, but herds remain seropositive. Herds in this stable phase, referred to as the lowly infectious herds, have much lower levels of viral shedding and a lower risk of infecting other herds. This corresponds to "positive stable" herds in PRRS terminology (Holtkamp et al., 2011). The transition from the highly infectious phase to the lowly infectious

166 phase is modelled as γI_h. Both infectious phases can contribute to the local spread and
167 movement-mediated transmission.

168
Figure 2. A conceptual diagram of between-herd PRRS transmission. Transmission occurs
through local spread and movement. Local spread occurs within a 5 km radius due to indirect
contact and airborne transmission at a rate of $\lambda_{local\ spread}$. Movement-mediated transmission,
modelled using trading data, occurs at a rate of $\lambda_{movement}$. Highly infectious herds can
transition to lowly infectious (seropositive) herds at a rate of γI_h. Both highly and lowly
infectious herds can be detected as positive through blood testing and clinical checkups. Upon
detection, herds can implement within-herd PRRS elimination at rate of αD.

176
The transmission rate via movement ($\lambda_{movement}$) is modelled as $P_{mov} \cdot I_{h_{mov}} + P_{mov} \cdot k_{mov} \cdot$
$\left(I_{l_{mov}} + D_{mov}\right)$,
where:
- P_{mov} represents the infection probability per risky movement from a highly infectious
herd
- $I_{h_{mov}}$ represents the number of risky movement contacts with the highly infectious
herds per day
- k_{mov} is a scaling factor representing the relative infectiousness of lowly infectious
compared to highly infectious herd via movement-mediated transmission
- $I_{l_{mov}}$ represents the number of movement contacts with lowly infectious herds
- D_{mov} represents number of movement contacts with detected herds.
Under the SPF system, risk-based trading is encouraged and practiced on a voluntary way.
Herds with a higher biosecurity level (i.e. red herds) can sell pigs to those with a lower
biosecurity level (i.e. blue herds), but not vice versa (Supplement Figure S1). Additionally,
herds lose their PRRS-negative status if they acquire pigs from PRRS-positive herds. This

means that risky movements - buying pigs from positive herds- can still occur, as herd owners
decide whether to find a new trade partner with PRRS-negative status or continue to trade while
accepting the loss of their PRRS-negative status. Therefore, in the default scenario,
representing the period from 2020 to 2021, we model a proportion of herds complying with
risk-based trading rule by purchasing pigs only from PRRS-negative herds. The level of
compliance is determined using surveillance data.

Transmission via local spread is caused by contact with virus-laden fomite (shared
equipment, vehicle, and workers), or virus-laden aerosols in the vicinity of an infectious herd
(often referred to as airborne transmission). In this study, a local area is defined as a 5 km radius
area, within which the transmission rate is assumed to be density-dependent; hence the
transmission rate ($\lambda_{local\ spread}$) is modelled as $\beta_{ls} \cdot I_{h_{ls}} + \beta_{ls} \cdot k_{ls} \cdot (I_{l_{ls}} + D_{ls})$,

Where:

- β_{ls} represents the transmission rate parameter for local spread of highly infectious herds

- $I_{h_{ls}}$ represents the number of highly infectious neighbor herds (within 5 km of the
recipient herd)

- k_{ls} is a scaling factor representing the relative infectiousness of lowly infectious
compared to highly infectious herd via local spread

- $I_{l_{ls}}$ represents the number of lowly infectious neighbor herds (within 5 km of the
recipient herd)

- D_{ls} represents the number of detected PRRS positive herds (within 5 km of the recipient
herd).

Both types of infectious herds can be detected via blood testing (active surveillance) or
clinical checks by veterinarians (passive surveillance). Regular blood sample testing is
conducted, with the frequency of sampling determined by herds' biosecurity level. In addition,
herds are visited by veterinarians regularly, with the frequency and sensitivity of clinical
checkups determined by herds' production types, as sow herds have more severe clinical signs
than finisher and weaner herds. Therefore, the detection of PRRS is modelled as
$blood_{sampling} \cdot SE_b + clinical_{visit} \cdot SE_{clinic}$, where:

- $blood_{sampling}$ is the rate of regular blood sampling

- $clinical_{visit}$ is the rate of the vet visiting for clinical check up

- SE_b and SE_{clinic} are the sensitivities of blood sampling and clinical checkups.

After detection, PRRS positive herds can eliminate the virus through control measures (see
Section 2.7). The time required for elimination depends on when farmers implement control
measures and the specific strategies they choose. Once PRRS is eliminated, herds are assumed
to become immediately susceptible. The elimination is modelled as αD, where α represents the
elimination rate determined by the SPF types (see Table 2).
231

232   2.3 Parameters
233

234 The model includes two types of parameters: local and regional. Local parameters are herd-
235 specific and reflect characteristics such as biosecurity levels, production type, trading pattern
236 matrix, and proximity to neighboring farms, as well as farm management decisions related to
237 interventions (Table 1).

238 Table 1. Local parameters of the model

Local parameter	Type	Value/Structure	Source
Biosecurity level	Categorical	Red SPF, Blue SPF, non-SPF	Surveillance data
Production type	Categorical	Sow herd, weaner herd, fattener herd	Surveillance data
Trading pattern matrix	List of tuples	{Seller ID 1, trading frequency}, {Seller ID 2, trading frequency} …	Surveillance data
Distance matrix	List of tuples	{Neighbor herd ID 1, distance}, {Neighbor herd ID 2, distance} ...	Geographic data
Risk-based trading	Binary	1 = avoid buying from PRRS-positive herds 0 = No such restriction	Surveillance data
Biosecurity improvement	Continuous	Level of reduction in herd susceptibility due to improved biosecurity (0 -1 scale)	Assumed as 1 under default scenario
Depopulation-repopulation	Binary	1 = implements depopulation-repopulation once herd is detected as PRRS-positive 0 = Does not implement	Assumed as 0 under default scenario

Regional parameters remain consistent across all areas and include transmission rate parameter for local spread and infection probability via movement, the infectious period of herds, and sensitivity for blood testing and veterinary clinical checks (Table 2). Due to discrepancies between two references regarding to herd-level sensitivity of clinical checkup, the impact of using the alternative herd-sensitivity is analyzed and presented in Supplement Figure S2.

Table 2. Regional parameters of the model

Regional parameter	Definition	Point estimation	Source
β_{ls}	Transmission rate parameter for local spread from highly infectious herds	0.0015 herd^{-1}Day^{-1}	Estimated (Section 2.4)
k_{ls}	Scaling parameters for lowly infectious herd (local spread route)	0.016	Estimated (Section 2.4)
P_{mov}	Infection probability per risky contact from highly infectious herds	0.26	(Galvis, Corzo, Prada, et al., 2022)
k_{mov}	Scaling parameters for lowly infectious herd (movement route)	1	(Galvis, Corzo, Prada, et al., 2022)
γ	Highly-to-lowly infection transition rate	1/90 Day^{-1}	Expert opinion
SE_b	Sensitivity for blood sampling test	0.85	(Willkan et al., 2025)
SE_{clinic}	Sensitivity for clinical check up by veterinarians	Sow herds: 0.3 (alternative: 0.07) Finisher and weaner herds: 0.05 (alternative: 0.015)	(Willkan et al., 2025; SEGES Innovation, 2025)
$blood\_sampling$	Sampling rate for the blood testing	Blue SPF and non-SPF herds: 1/365 Day^{-1} Red SPF: 1/30 Day^{-1}	(SPF, 2024)
$clinical\_visit$	Clinical visit rate by veterinarians	Sow herd: 1/30 Day^{-1} Weaner herd: 1/30 Day^{-1} Finisher herd: 1/60 Day^{-1}	(SPF, 2024)
α	Elimination rate	Red SPF: 1/90 Day^{-1} Blue SPF: 1/700 Day^{-1} Non-SPF: 1/800 Day^{-1}	Estimated (Section 2.4)
$complaince$	The proportion of herds complying with risk-based trading	0.8	Estimated (Section 2.4)

2.4 Statistical method

We estimate transmission rate parameters for local spread by fitting time-series infection data into the model. This method relates the exposure to the infection sources to the infection probability. The number of herds becoming infected over each test interval $(t, t + \Delta t)$ follows a binomial distribution with a binomial total susceptible herd at each time interval (which is the time difference between two data entries in surveillance data). The probability of getting infected due to local spread can be derived as:

$$P\left(\frac{case}{S}\right) = 1 - e^{-(\beta_{ls} \cdot I_{h_{ls}} \cdot \Delta t + \beta_{ls} \cdot k \cdot I_{l_{ls}} \cdot \Delta t)} \qquad \text{Eq(1)}$$

Where $P\left(\frac{case}{S}\right)$ is a binary value representing whether a new case occurs during an observation time interval. $\beta_{ls} \cdot I_{h_{ls}} \cdot \Delta t$ represents the transmission rate contributed by highly infectious herds within a 5 km during a time interval Δt while $\beta_{ls} \cdot k \cdot I_{l_{ls}} \cdot \Delta t$ accounts for the transmission rate from lowly infectious herds within the same radius and time interval.

The likelihood function, as a function of transmission rate parameters, is given by:

$$L(P(\theta)) = \prod_x (P)^{cases_x} (1 - P)^{(S_x - cases_x)} \qquad \text{Eq(2)}$$

Where P is derived from Eq(1). θ represents the two parameters β_{ls} and k_{ls}. The index x donates the number of data lines in the infection dataset. The confidence interval of estimation was listed in the Supplement Table S3.

The elimination rate α per SPF types is estimated by calculating the average time between the detected and the return to negative herd. Similarly, the compliance to risk-based trading is calculated as the proportion of herds that did not receive pigs from positive herds.

2.5 R_0 calculation

In this study, we calculate two types of R_0, including between-herd R_0 for each herd and a national R_0 (Chang et al., 2025). Between-herd R_0 represents the number of herds that an

infectious herd can infect in a population where all other herds are susceptible, which is used to present the spatial heterogeneity. We obtain the between-herd R_0 using Next Generation Matrix method (Diekmann et al., 2010) (Supplement Text S4). The between-herd R_0 consists of two components: the local spread route $R_{local\ spread}$ and movement route $R_{movement}$:

$$R_0 = R_{local\ spread} + R_{movement} \qquad\qquad Eq(3)$$

For local spread route, the $R_{local\ spread}$ can be calculated as:

$$R_{local\ spread} = \frac{\beta_{ls} \cdot N_{ls}}{(\gamma + \omega)} + \frac{k \cdot \beta_{ls} \cdot N_{ls}}{\omega} \frac{\gamma}{(\gamma + \omega)} + \frac{k \cdot \beta_{ls} \cdot N_{ls}}{\alpha} \qquad Eq(4)$$

Where the first part of the equation represents the contribution from highly infectious herds while the second part and the third part represent the contribution from lowly infectious herds (undetected or detected part respectively). N_{ls} is the total number of herds in a 5 km neighborhood. γ is the rate at which herds transition from highly infectious to less infectious but seropositive phases. ω represents the average detection rate, which varies by biosecurity and production type. It is calculated based on the frequency and sensitivity of test or clinical checkups as below:

$$1 - e^{\frac{w_b}{blood\_sampling}} = SE_b \qquad\qquad Eq(5)$$

$$1 - e^{\frac{w_c}{clinical\ visit}} = SE_c$$

$$\omega = \omega_b + \omega_c$$

Table 3. Detection rate based for herds categorized with different production types and SPF types

Parameter	Definition	Value
ω_b	The average detection rate for blood sampling	Red SPF: 0.0630 Day^{-1} Blue SPF: 0.0052 Day^{-1} Non-SPF: 0.0052 Day^{-1}
ω_c	The average detection rate for clinical checkup	Sow herds: 0.0120 Day^{-1} Weaner herds: 0.0017 Day^{-1} Finisher herds: 0.00085 Day^{-1}

294   In the Eq (4), the $\frac{1}{\gamma + \omega}$ represents the average time herds stay in a highly infectious phase. The

295    $\frac{1}{\omega}$ represents the period that herds remain in the lowly infectious phase. The ratio

296    $\frac{\gamma}{(\gamma + \omega)}$ represent the proportion of herds that reach the lowly infectious phase before detection.

297   Similarly, $R_{movement}$ can be calculated as:

$$R_{movement} = \frac{P_{mov}c}{(\gamma + \omega)} + \frac{k_{mov}P_{mov}c}{\omega} \frac{\gamma}{(\gamma + \omega)} + \frac{k_{mov}P_{mov}c}{\alpha} \qquad Eq(5)$$

where c is contact rate with the outward trading partners.

The national level R_0 represents the average number of herds that one infectious herd can infect

when all the other herds are susceptible in Denmark. National level R_0 is calculated using the

simulated herd prevalence once transmission dynamics reach equilibrium ($National\ R_0 =$

$\frac{1}{(1 - herd\ prevalence)}$) (Chang et al., 2025). For each intervention strategy, a corresponding

national R_0 can be estimated. If simulated herd prevalence decreases to 0, it indicates that

PRRS can be eradicated by the intervention ($R_0 < 1$). However, if herd prevalence decreases

but reaches a new equilibrium, the intervention can reduce transmission but cannot eradicate

PRRS (national $R_0 > 1$).

2.6 Sensitivity analysis

Under the current surveillance system, herds lose their PRRS-negative status when they acquire

pigs from PRRS-positive herds, assuming a 100% infection risk from such movements without

testing. This assumption in the surveillance data prevents estimation of actual the infection

probability per movement (P_{mov}) via pig movements and surveillance data. To address this, we

adopt literature-based estimate for P_{mov} in our default scenario (Scenario S1). In Scenario S2,

we apply expert opinion from the Danish pig industry, setting P_{mov} to 1 and assuming scaling

factor k_{mov} is the same as the k_{ls} (Table 4; Scenario S2). In Scenario S3, reflects the current

surveillance assumption, with both P_{mov} and k_{mov} set to 1, while halving the estimated local

transmission parameters estimated from Eq. (2). The values used in all three scenarios are

summarized in Table 4.

Table 4. Transmission rate parameters and infection probability for sensitivity analysis

Parameter(unit)	Scenario 1	Scenario 2	Scenario 3
P_{mov} (unitless)	0.26	1	1
k_{mov} (unitless)	1	0.016	1
β_{ls} (per herd^{-1}Day^{-1})	1.5e-3	1.5e-3	7.5e-4
k_{ls} (unitless)	0.016	0.016	0.016

325
326

2.7 Interventions

Interventions can be classified into several groups, including those targeting internal
biosecurity, external biosecurity, movement and testing frequency. Improving internal
biosecurity can shorten a herd's infectious period. For instance, depopulation-repopulation
(DPRP) is an effective but costly method of PRRS elimination within herds and is mainly used
in herds with weaners and finishers. This approach involves fully depopulating a production
unit, thoroughly cleaning and disinfecting the facilities, and then introducing new PRRS-free
pigs after seven days of being empty. This strategy can return a sow herd to susceptible status
in approximately 5 months, a weaner herd in 2 months, and a finisher herd in 3 months,
depending on the production schedule. To model this intervention, each herd has a binary
depopulation-repopulation parameter indicating the herd owners' decision (See Table 1).

In comparison, external biosecurity, such as quarantine facilities, disinfecting trucks
and controlled ventilation, can reduce the transmission risk between herds. The SPF system
provides a rough classification of farm biosecurity levels. However, since we cannot find
significant difference in the transmission rate parameters among different SPF system herds,
we assume that the biosecurity level is consistent across herds. Given that we do not have the
quantified data on the impact of each biosecurity measure on transmission risk, nor the
percentage of farms that comply with each measure, we model the effect of improved external
biosecurity on transmission rate parameter by a factor between 0 and 1 (Table 1), and the
number of herds adopting improved external biosecurity is modelled to vary from 0% ~ 100%.

Herds are recommended not to purchase pigs from PRRS-positive herds. Therefore,
each herd is assigned a binary local parameter indicating whether it follows this
recommendation (Table 1). During the study period, about 80% of herd owners complied with
the risk-based trading rule and 20% of herds received pigs from PRRS-positive herds, which
is modelled in the default scenario. Lastly, the active and passive surveillance system is
modelled with sensitivity and frequency presented in Table 2.

In addition to the default scenario, we investigate several additional interventions that

target different routes:

Interventions at movement route:

       1) Default scenario: risk-based trading with 80% of compliance

       2) Risk-based trading with 100% compliance: Herds only buy pigs from herds that have

358   not been detected as positive. However, they might still acquire pigs from infectious

359   but undetected herds (I_h and I_l)

360   3) Movement ban: The scenario is for exploration

361   Interventions reduce the external biosecurity, hence reducing susceptibility of herds:

362   1) Default scenario: under the default biosecurity (red and blue)

363   2) 50% of herds are randomly selected to enhance external biosecurity, reducing their

364   susceptibility to PRRS infection by 20%

365   3) All the herds in high-risk herds will improve their external biosecurity, reducing the

366   herds susceptibility by 20%

367   Interventions improve the internal biosecurity via DPRP (i.e. depopulation-repopulation):

368   1) Implement DPRP in 50% of herds, selected randomly

369   2) Implement DPRP in the high-density areas (herds with > 10 neighboring herds within

370   5km)

371   3) Implement DPRP in all herds

372   Interventions at surveillance system:

373   1) Increase testing frequency to every half year

374   2) Increase testing frequency to every month

375 3. Results

376 3.1 Descriptive analysis of the data

A total of 3,420 herds were registered with PRRS status during the study period, representing

approximately 60% of the entire population. Herds are classified into several production types

based on the ratio between sows and finishers (Schulz, 2019). Information on production type

was missing for 277 herds. Among herds with known production types, finisher herds were the

most common (69%), followed by sow herds (24%) and weaners (7%) (See Figure 3A). Among

the study population, 1,804 herds were classified as SPF blue, 5 as SPF green, and 128 as SPF

red, and 1,483 of non-SPF herds. Since SPF blue and SPF green herds follow the same

detection process, they were treated identically in the model.

Figure 3. Composition of the study population: A) The percentage distribution of the three production types; B) the distribution of herds across production types and SPF status, showing the flow of herds into different biosecurity level (Green, blue, red, non-SPF). The thickness of the connecting lines represents the relative proportion of herds in each category.

The movement dataset consists of 100,464 records of movements, involving 1,682 sellers and 2,596 buyers between January 2020 and December 2021, with 15% (n = 498) of farms reporting no between-farm movement activities. Throughout the study period, 78 % of the pig movement relationships between pig farms persisted, and one fifth of the seller herds (n = 336) generated half of all pig movements. Most pig movements, among known production categories, occurred between sows and finisher facilities (34%) and between sows and weaner farms (15%). Overall, there was little variation in the distance traveled by pigs between farms across production types, with a median of 10.5 km (mean 35.6 km with a range 0–386.2 km). On average, herds have 11 neighboring herds in the local area (within 5km radius), with 25% percentile at 6 and 75% percentile at 16. Different production types show no significant differences in local density, though weaner herds are slightly more concentrated. Finisher herds have the fewest outward trading partners, averaging 2.26, while sow and weaner herds have more outward trading partners, averaging 2.57 and 3.14, respectively.

3.2 Between herd R_0 map

Using the estimated parameters (Section 2.4), we calculate the between-herd basic reproduction ratio R_0 and stratify it by the movement route $R_{movement}$ and local spread $R_{local\ spread}$. Herds with $R_0 > 1$ are expected to transmit the infection to more than one other herd during their infectious period, while herds with $R_0 < 1$ are expected to transmit to fewer than one herd. The R_0 map (Figure 4) shows that high risk herds are predominantly clustered in North Denmark,

Central Denmark (especially along the coast), and certain areas of South Denmark. Our R_0
map shows a strong alignment with observed prevalence. The distribution of between herd R_0
is presented in Supplement Figure S5.

Figure 4. Between herd R_0 map for the PRRS transmission. A) the observed prevalence in
different regions in Denmark (Source from Landmand.dk); B) the between-herd R_0 map; C) the
$R_{local\ spread}$ map. Dots in red represent herds with $R_0 > 1$ and dots in green and yellow
represent herds with $R_0 < 1$; D) $R_{movment}$ map under 80% compliance with risk-based trading.

3.3 Relative contribution

Infectious herds progress through three phases, highly infectious (I_h), lowly infectious phase
(I_l), and detected phase (D). In Section 2.5, we derive the between-herd R_0 for each infectious
phase, allowing us to express their relative contribution. For example, for the local spread, the
three phases contribute to transmission with $\frac{1}{(\gamma+\omega)} : \frac{k_{ls}}{\omega}\frac{\gamma}{(\gamma+\omega)} : \frac{k_{ls}}{\alpha}$, where herd production type
and biosecurity level influence these ratios (Table 5). On average, detected herds contribute to
around 40% of the total transmission. Parameters from Scenario 2 - 3 have only a marginal
impact on this distribution, with contribution of detected herds ranging from 35% to 50%.

In general, 59-79 % of the new infections are caused by the local spread, with an overall average contribution of 68%. The contribution tends to be highest in the sow herds, followed by the finisher herds and weaner herds. Transmission rate parameters can influence the relative contribution of the two routes: local spread contributes to approximately 70% of infections under Scenario 2, and 58% under Scenario 3.

Table 5. Relative contributions of infectious phases, detection methods, and transmission routes across different production types and biosecurity levels using Scenario 1 and default herd sensitivity value. The "% of herds" column represents the proportion of herds for each production type and biosecurity level, summing to 100% across all categories. For each row, the relative contributions of I_h (highly infectious), I_l (lowly infectious), and D (detected) sum to 100%. The "% detected by blood testing" column reflects the proportion of detected herds identified by blood testing, with the remainder detected via clinical notification (not shown in the table), together summing to 100%. The "% local spread" column represents the contribution of local transmission, with the remainder attributed to movements (not shown in the table), together summing to 100%.

Production type	Biosecurity level	% of herds	% of I_h contribution	% of I_l contribution	% of D contribution	% of local spread
Sow	Red	1%	58%	2%	40%	79%
Sow	Blue	15%	45%	7%	48%	65%
Sow	Non-SPF	5%	33%	8%	59%	70%
Weaner	Red	0.2%	59%	4%	37%	65%
Weaner	Blue	5%	39%	22%	39%	59%
Weaner	Non-SPF	2%	33%	22%	45%	67%
Finisher	Red	2%	63%	3%	34%	65%

Finisher	Blue	33%	41%	24%	35%	63%
Finisher	Non-SPF	37%	37%	24%	39%	70%

The model includes both active surveillance (via blood sample testing) and passive surveillance (via clinical checkups). The proportion detected by blood sampling is given by competing rates of detection: $\frac{\omega_b}{\omega_b + \omega_c}$, where two detection rates ω_b and ω_c are functions of the herd-level sensitivities of the clinical and blood-testing, as well as their frequency (See Eq. (5)). For red herds, which have monthly blood sampling, most of them are detected by blood testing ($> 98\%$).

Using the herd-level sensitivities of clinical checkups from a recent study by Willkan et al. (2025), the model estimates that 35% of infected sow herds are detected by blood sampling and 66% by clinical checkups, while 79-88% of infections in weaner and finisher herds are detected by blood sampling with only 12%-21% detected by clinical checkups. However, if we match detection ratios similar to observed ratios from a report from SEGES innovation (2025), with approximately 70% of sow herd infections and 95% of weaner/finisher herd infections detected by blood testing, the herd level sensitivity for clinical checkups should be 3-4 fold lower (Table 2). However, the overall effect of herd-level sensitivity of clinical checkups on model output is limited, as shown in Supplement Figure S2.

3.4 Impact of Interventions

Observed herd prevalence stabilizes around 30% without additional interventions under the default control strategy in three scenarios (Figure 5). Despite variations in parameter assumptions, Scenarios 1-3 shows consistent trends in the impact of interventions.

With improving risk-based trading to full compliance, observed prevalence reduces to around 22% under Scenario 1 and 3 (Figure 5A&C, line 03), but increase to 28 % in Scenario 2. Risk-based trading cannot entirely prevent movement-mediated spread because of undetected infectious herds, as a hypothetic scenario complete movement ban can reduce prevalence to around 4% in Scenario 1 and 2 (Figure 5A & B, line 11) and 0.4% in Scenario 3 (Figure 5C, line 11).

Improving biosecurity by reducing herd susceptibility by 20% has only a modest impact with observed prevalence stabilizing at 23% when applied to half of the herds and 18% when

implemented in all herds – consistently across three scenarios (Figure 5A-C, line 02 and 04).

Increasing testing frequency to twice per year initially leads to a rise in observed prevalence

and then reduce to around 20% (Figure 5A-C, line 05). Monthly testing has a much stronger

effect, ultimately lowering prevalence to 2%-3% (Figure 5A-C, line 06).

Among all the interventions, depopulation-repopulation (DPRP) has the most

immediate impact within the first two years, with prevalence declining to 10%-20% (Figure 5,

line 07-09), depending on the proportion of herds complying with the measure. The

combination of depopulation-repopulation, stricter risk-based trading and improving blood

testing frequency to twice per year proves to be sufficient in eradicating PRRS within 20 years

on average (Figure 5A & C, line 13), but not in Scenario 2 (Figure 5B, line 13). These results

highlight that while individual interventions can reduce transmission to some extent, a

comprehensive approach might be needed for eradication.

Figure 5. Impact of different intervention strategies on the observed prevalence across three scenarios: A) Scenario 1, B) Scenario 3. Each line represents a specific intervention on combination of interventions, with colors corresponding to the legend on the right. The y-axis shows the percentage of positive herds, and the x-axis represents time in years. Shaded areas indicate variation across 100 model simulations. Interventions include biosecurity improvements, risk-based trading, depopulation-repopulation (DPRP) strategies, and combinations. All interventions are implemented starting from year 0.

Using the simulated true prevalence (see Supplement Figure S6) and the equation in
Section 2.5, the national-level basic reproduction number R_0 are estimated (Table 6). Whether
PRRS can persist for each control strategy depends on whether R_0 is below 1. Either local
spread or movement alone may be sufficient to sustain PRRS transmission, as indicated by $R_0 >$
1 under control strategies 11 and 12 (Table 6).

Controlling local spread via depopulation-repopulation (DPRP) might seem effective
in reducing the observed prevalence (Figure 5 line 09), but its impact on the true prevalence
and R_0 are limited (Table 6 line 09). In comparison, increasing the frequency of blood testing
frequency has the strongest effect in reducing transmission (Table 6 line 06), even though it
can initially lead to an increase in observed prevalence due to improved detection (Figure 5
line 06).

Table 6. National R_0 under different control strategies and scenarios

Control strategy	Scenario 1	Scenario 2	Scenario 3
01 default	1.56	1.51	1.52
02 improve biosecurity in 50 % herds	1.41	1.36	1.38
03 risk-based trading with 100% compliance	1.37	1.54	1.35
04 improve biosecurity in 100 % herds	1.30	1.22	1.25
05 blood testing frequency 182 days	1.33	1.33	1.25
06 blood testing frequency 30 days	1.03	1.01	1.02
07 DPRP in 50% of the random herds	1.45	1.42	1.50
08 DPRP in high density areas	1.43	1.37	1.46
09 DPRP in all herds	1.30	1.24	1.32
10 DPRP and risk-based trading in all herds	1.22	1.26	1.23
11 movement bans	1.05	1.05	1.00 (>1)
12 no local spread	1.01	1.00 (> 1)	1.02
13 DPRP, risk-based trading in all herds and blood testing frequency 182 days	1.00 (>1)	1.07	< 1

4. Discussion

PRRS has been endemic in Denmark for over 30 years, causing significant economic losses, with the estimated cost ranging between 4 to 139 euro per sow per year (Rathkjen & Dall, 2017). In 2022, a reduction plan was introduced, focusing on compulsory PRRS-status reporting, elimination of PRRS in positive herds and recommended risk-based trading under SPF system. This raises questions about the effectiveness of these measures. To address this, we developed a between-herd transmission model accounting for both local spread and movement-mediated transmission, replicating dynamics before the legislative change and predicting control impacts.

A key innovation of this study is its modeling of PRRS spread, considering the distinct roles of highly infectious and lowly positive herds and explicitly incorporating the Danish surveillance system. The model estimated that highly infectious herds (positive unstable) are 50 times more infectious than lowly infectious herds (positive stable), highlighting their important role in PRRS transmission and control. In addition, we assessed that the current surveillance program misses 16% of infectious herds, which may contribute up to 60% of the overall transmission. These undetected infections can be reduced by more frequent testing, aligning with recent findings on benefits of increased sampling frequency for early detection (Willkan et al., 2025). Since herds are most infectious during the first three months after infection, testing at intervals of three months or shorter can substantially reduce the number of hidden infections, as illustrated by the comparison between lines 10 and 13 in Figure 5.

This study generates the first between-herd R_0 map for the PRRS, explaining the spatial heterogeneity in PRRS transmission. The findings are consistent with observed high prevalence in certain Danish regions including North Demark, Central Denmark (especially along the coast), and parts of South Denmark (Figure 4). A previous study also identified significant spatial clusters, suggesting PRRS endemic states in these areas (Lopes Antunes et al., 2015). Stratification of R_0 reveals distinct spatial pattern: R_{mov} is relatively homogenous across the country, whereas $R_{local\ spread}$ shows strong spatial clustering, reflecting spatial patterns in observed surveillance data (Figure 4). This suggests that local spread is a key driver of PRRS transmission in Denmark. This is further supported by the estimates of approximately 60-70% of new infections caused by local spread (Table 5). Although scenarios 2 and 3 increase the relative contribution of movement-related transmission to 40%-50%, local spread remains a dominant role. However, it is important to note that our study includes all the herds that report

their PRRS status, representing 60% of the total population. Consequently, some local spread may in fact reflect movements involving those non-included herds. With the eradication plan requiring compulsory PRRS status reporting for herds with more than 10 sows or 100 pigs, future analyses should include a broader range of herds when available and could help clarify this uncertainty and improve our understanding.

Several risk factor analyses have identified high density as a main risk factor, suggesting that local spread – potentially through airborne transmission– plays an important role (Arruda et al., 2017; Baysinger et al., 1997; Firkins & Weigel, 2004; Mortensen et al., 2002). Modelling studies in the U.S. have estimated that pig movement explained 15 to 20% of the new infections, while 50%-60% due to local spread in sow and finisher herds (Galvis, Corzo, & Machado, 2022; Galvis, Corzo, Prada, et al., 2022). These studies also found that pig movements played a greater role in weaner herds but were less in finisher and sow herds. The pattern aligns with our modeling results. Approximately 70% of Danish herds are finishers, and since they primarily sell pigs to slaughterhouses, their role in movement-mediated transmission is limited. Compared to these studies, our model simplifies indirect contact as part of local spread due to data limitations. However, this simplification is unlikely to have a significant impact, as indirect contact accounts for only 0.3% to 2.5% of transmission (Galvis, Corzo, & Machado, 2022). In contrast, studies in Canada have attributed PRRS transmission primarily to direct or indirect contact between herds under shared ownership as they found no evidence of airborne transmission (Rosendal et al., 2014). As a result, modeling efforts in Canada have focused solely on movement-mediated transmission (Thakur et al., 2015). These cross-country differences may be explained by variations in herd density, herd structure, and control measures, such as risk-based trading in Denmark. Despite multiple studies highlighting the role of animal movement in PRRS spread (Amirpour Haredasht et al., 2017; Pamornchainavakul et al., 2023; Thakur, Sanchez, et al., 2015), our model is the first to assess the impact of risk-based trading. We showed that with 80% of compliance, risk-based trading reduced the observed prevalence from 40% to 30% and can further reduce it to 20% with 100% of compliance.

High density of herds, complex trade patterns, infrequent active surveillance and low sensitivity of passive surveillance all contribute to the persistence of PRRS (as shown in Table 5). Depopulation-repopulation shows an immediate and strong effect on observed prevalence, but its impact on true underlying transmission is limited as it does not influence hidden infections. In contrast, increasing testing frequency has the greatest effect on the reducing the real transmission, although it initially raises observed prevalence. Our results suggest that

interventions targeting different transmission routes can have an additive effect. A

comprehensive control should include measures to reduce hidden infections via more frequent

testing, control the local spread in high-density areas via improving biosecurity measures,

depopulation-repopulation, stricter risk-based trading to limit the introduction of PRRS (Figure

5, line 03) in free areas (Figure 5, line 05 and 13). Similar suggestions have been made in

Germany (Fahrion et al., 2014).

An overestimation of R_0 and an underestimation of intervention effectiveness in the

present study could be caused by the higher proportion of false-positive test results in the

current surveillance system. Specifically, herds that purchase pigs from PRRS-positive herds

are classified as PRRS-positive, introducing uncertainty in estimating the true infection

probability from movement. False-positive herds in the surveillance data do not contribute to

further PRRS transmission, whereas assuming an infection probability of 1 artificially

increases the transmission potential. To assess the impact of assuming a 100% infection

probability via risky movement for both highly infectious and lowly infectious herds, we tested

this scenario, which led to an overestimation of PRRS prevalence at around 43%, compared to

the observed prevalence of 30%. This suggests that the infection probability via risky

movement is likely lower than 1.

Further, a characteristic of the Danish pig industry is the establishment of joint

operations to reduce testing costs and mitigate transmission risks (SPF, 2024). Under this

system, multiple herds submit samples together, and their PRRS status is updated

simultaneously in the surveillance system. Herds within a joint operation can be automatically

classified as positive if any other herd in the group tests positive. A similar issue exists in the

Danish cattle industry for Salmonella surveillance, where classifications are updated at the

business level rather than for individual farms (Conrady et al., 2024). Pig herds within a joint

operation typically share the same owner, engage in regular trade (e.g., from sow herds to

weaner herds and then to finisher herds), or are located in close proximity. Given the higher

likelihood of transmission between these connected herds, it is reasonable to treat them as a

single epidemiological unit. Despite functioning as a unified unit epidemiologically, each herd

within the joint operation still maintains individual entries in the surveillance system. This

leads to an overestimation of the observed prevalence. Relying on this biased surveillance data

to assess transmission dynamics can result in an overestimation of R_0, making PRRS appear

more difficult to eradicate in the model than it is.

This model was calibrated with PRRS-1 surveillance data, as PRRS-1 is more prevalent

in Denmark (Strategy for the Reduction of PRRS in Pigs in Denmark, 2022), while PRRS-2

can be associated with the vaccination strain was originally introduced with a vaccination program (Balka et al., 2015; Kvisgaard, 2017). Experimental studies suggest that vaccination can protect animals from re-infection (Canelli, 2018; Lunney et al., 2010), but its efficacy in preventing transmission at herd level remains unclear and controversial. Vaccine strains have been shown to spread between farms (Bøtner et al., 1997, 1999; Grosse Beilage et al., 2009; Kiss et al., 2006), raising concerns about their role in transmission. Modelling studies assumed 1% of vaccine efficacy at the herd level, leading to only a marginal decrease in incidence (Galvis, Corzo, Prada, et al., 2022; Valdes-Donoso & Jarvis, 2022). In Denmark, modified live viral vaccines for PRRS-1 and PRRS-2 are used exclusively in PRRS-positive herds to mitigate clinical symptoms and are not considered a tool for reducing between herd transmission. Therefore, we did not model the impact of vaccination.

While our model sufficiently captures spatial heterogeneity at a national scale, we acknowledge that the model and data have limitations in representing all aspect of spatial heterogeneity. As a herd-level transmission model, it cannot assess the impact of herd size or internal biosecurity measures on the within-herd transmission dynamics, such as the number of pigs getting infectious or the duration of outbreak in a herd (Evans et al., 2010). In addition, while the model explicitly accounts for herd owners' decisions on depopulation-repopulation and improving biosecurity level, the lack of data at herd level requires assuming these decisions. Although SPF herds in Denmark are recommended to quarantine incoming gilts for 42 days—potentially reducing infection risk via movements - there is no available data to quantify transmission probability through risky movements (P_{mov}). As a result, we are unable to assess the effectiveness of quarantine in reducing PRRS transmission with the current surveillance data. Future studies that monitor herd status changes after risky movement, both with and without quarantine, could improve our understanding of how quarantine impact the control of PRRS. Nonetheless, our study suggests that achieving eradication may require a combination of more frequent testing, targeted within-herd PRRS elimination, and stricter risk-based trading practices. It also identifies PRRS hotspots and transmission routes, providing evidence-based recommendations for effective control.

Conclusion and recommendation

This study identifies high-risk areas for PRRS transmission presented in R_0 map. In the study population, current surveillance misses about 17% of the infectious herds that contribute to

60% of the transmission. While PRRS elimination within herds via depopulation-repopulation
seems the most effective control measure on reducing the observed prevalence, it might have
limited impact on reducing the true transmission, as large part of the transmission is attributed
to hidden infection. A comprehensive control strategy is essential for nationwide eradication.
Improved active surveillance through more frequent testing, prioritizing local control in high-
risk areas, stricter risk-based trading to protect PRRS-free areas are key to achieving
eradication.

Acknowledgement

646
647
648 We would like to thank Stefan Widgren, Pauline Ezanno, Nicolai Rosager Weber, Sara
649 Dalsgaard, and Kristian Møller for their invaluable suggestions contributed to the improvement
650 of this study. This work received co-funding from European Union's Horizon Europe under
651 Grant Agreement No 101136346: European Partnership Animal Health &Welfare, Joint
652 Internal Project SOA10, SOA 22 and SOA 15 and the pigs levy fund.

References

References

Amirpour Haredasht, S., Polson, D., Main, R., Lee, K., Holtkamp, D., & Martínez-López,

B. (2017). Modeling the spatio-temporal dynamics of porcine reproductive & res-

piratory syndrome cases at farm level using geographical distance and pig trade

network matrices. *BMC Veterinary Research*, *13*(1), 163.

https://doi.org/10.1186/s12917-017-1076-6

Arruda, A. G., Vilalta, C., Perez, A., & Morrison, R. (2017). Land altitude, slope, and cov-

erage as risk factors for Porcine Reproductive and Respiratory Syndrome (PRRS)

outbreaks in the United States. *PloS One*, *12*(4), e0172638.

Balka, G., Wang, X., Olasz, F., Bálint, Á., Kiss, I., Bányai, K., Rusvai, M., Stadejek, T., Mar-

thaler, D., Murtaugh, M. P., & Zádori, Z. (2015). Full genome sequence analysis of

a wild, non-MLV-related type 2 Hungarian PRRSV variant isolated in Europe. *Virus*

Research, *200*, 1–8. https://doi.org/10.1016/j.virusres.2015.01.014

Baysinger, A. K., Dewey, C. E., Straw, B. E., Brumm, M. C., Schmitz, J., Doster, A., & Kel-

ling, C. (1997). Risk factors associated with endemic reproductive deficiencies

caused by PRRSV infection. *Journal of Swine Health and Production*, *5*(5), 179–

187.

Bøtner, A., Nielsen, J., & Bille-Hansen, V. (1994). Isolation of porcine reproductive and

respiratory syndrome (PRRS) virus in a Danish swine herd and experimental in-

fection of pregnant gilts with the virus. *Veterinary Microbiology*, *40*(3–4), 351–

360. https://doi.org/10.1016/0378-1135(94)90122-8

Bøtner, A., Nielsen, J., Oleksiewicz, M. B., & Storgaard, T. (1999). Heterologous chal-

lenge with porcine reproductive and respiratory syndrome (PRRS) vaccine virus:

No evidence of reactivation of previous European-type PRRS virus infection. *Vet-*

erinary Microbiology, *68*(3), 187–195. https://doi.org/10.1016/S0378-

1135(99)00053-X

Bøtner, A., Strandbygaard, B., Sørensen, K. J., Have, P., Madsen, K. G., Madsen, E. S., &

Alexandersen, S. (1997). Appearance of acute PRRS-like symptoms in sow herds

after vaccination with a modified live PRRS vaccine. *Veterinary Record*, *141*(19),

Article 19. https://doi.org/10.1136/vr.141.19.497

Canelli, E. (2018). Efficacy of a modified-live virus vaccine in pigs experimentally in-

fected with a highly pathogenic porcine reproductive and respiratory syndrome

virus type 1 (HP-PRRSV-1). *Veterinary Microbiology*.

Chang, Y., Widgren, S., de Jong, M. C., Tratalos, J. A., More, S. J., & Hartemink, N. (2025).

Evaluating the effectiveness of badger vaccination combined with cattle test-

and-removal in managing Bovine Tuberculosis: Insights from a two-host and

multi-route transmission model. *Preventive Veterinary Medicine*, *235*, 106386.

Cho, J. G., & Dee, S. A. (2006). Porcine reproductive and respiratory syndrome virus.

Theriogenology, *66*(3), Article 3. https://doi.org/10.1016/j.theriogen-

ogy.2006.04.024

Conrady, B., Dervic, E. H., Klimek, P., Pedersen, L., Reimert, M. M., Rasmussen, P.,

Apenteng, O. O., & Nielsen, L. R. (2024). Social network analysis reveals the fail-

ure of between-farm movement restrictions to reduce Salmonella transmission.

Journal of Dairy Science, *107*(9), 6930–6944. https://doi.org/10.3168/jds.2023-

24554

Conrady, B., Mortensen, S., Nielsen, S. S., Houe, H., Calvo-Artavia, F. F., Ellis-Iversen, J.,

& Boklund, A. (2023). Simulation of Foot-and-Mouth Disease Spread and Effects

701   of Mitigation Strategies to Support Veterinary Contingency Planning in Denmark.

702   *Pathogens*, *12*(3), 435. https://doi.org/10.3390/pathogens12030435

703 Corzo, C. A., Mondaca, E., Wayne, S., Torremorell, M., Dee, S., Davies, P., & Morrison, R.

704   B. (2010). Control and elimination of porcine reproductive and respiratory syn-

705   drome virus. *Virus Research*, *154*(1–2), Article 1–2. https://doi.org/10.1016/j.vi-

706   rusres.2010.08.016

707 Danish Ministry of Food, Agriculture and Fisheries. (2023). *Executive Order on Porcine*

708   *Reproductive and Respiratory Disease (PRRS)*. https://www.retsinfor-

709   mation.dk/eli/lta/2023/997

710 Diekmann, O., Heesterbeek, J., & Roberts, M. G. (2010). The construction of next-gener-

711   ation matrices for compartmental epidemic models. *Journal of the Royal Society*

712   *Interface*, *7*(47), 873–885.

713 Evans, C. M., Medley, G. F., Creasey, S. J., & Green, L. E. (2010). A stochastic mathemati-

714   cal model of the within-herd transmission dynamics of porcine reproductive and

715   respiratory syndrome virus (PRRSV): Fade-out and persistence. *Preventive Veter-*

716   *inary Medicine*, *93*(4), 248–257. https://doi.org/10.1016/j.prevet-

717   med.2009.11.001

718 Fahrion, A. S., Beilage, E. G., Nathues, H., Dürr, S., & Doherr, M. G. (2014). Evaluating

719   perspectives for PRRS virus elimination from pig dense areas with a risk factor

720   based herd index. *Preventive Veterinary Medicine*, *114*(3–4), 247–258.

721   https://doi.org/10.1016/j.prevetmed.2014.03.002

722 Fertner, M., Conrady, B., Røgind, A. S., Nielsen, E. O., & Boklund, A. (2025). Indirect con-

723   tacts between Danish pig farms–what are the frequencies and risk-reducing

measures, and how can they be used in simulation models? *Acta Veterinaria*

Scandinavica, *67*(1), 7.

Firkins, L. D., & Weigel, R. M. (2004). A retrospective study of risk factors for porcine re-

productive and respiratory syndrome virus infection and clinical disease in swine

herds in Illinois during the early years of the pandemic. *Journal of Swine Health*

and Production, *12*(1), 23–28.

Galvis, J. A., Corzo, C. A., & Machado, G. (2022). Modelling and assessing additional

transmission routes for porcine reproductive and respiratory syndrome virus: Ve-

hicle movements and feed ingredients. *Transboundary and Emerging Diseases*,

69(5), e1549–e1560. https://doi.org/10.1111/tbed.14488

Galvis, J. A., Corzo, C. A., Prada, J. M., & Machado, G. (2022). Modelling the transmission

and vaccination strategy for porcine reproductive and respiratory syndrome vi-

rus. *Transboundary and Emerging Diseases*, *69*(2), 485–500.

https://doi.org/10.1111/tbed.14007

Grosse Beilage, E., Nathues, H., Meemken, D., Harder, T. C., Doherr, M. G., Grotha, I., &

Greiser-Wilke, I. (2009). Frequency of PRRS live vaccine virus (European and

North American genotype) in vaccinated and non-vaccinated pigs submitted for

respiratory tract diagnostics in North-Western Germany. *Preventive Veterinary*

Medicine, *92*(1–2), 31–37. https://doi.org/10.1016/j.prevetmed.2009.07.010

Holtkamp, D., Kliebenstein, J., Neumann, E., Zimmerman, J., Rotto, H., Yoder, T., Wang,

C., Yeske, P., Mowrer, C., & Haley, C. (2013). Assessment of the economic impact

of porcine reproductive and respiratory syndrome virus on United States pork

producers. *Journal of Swine Health and Production*, *21*(2), 72–84.

https://doi.org/10.54846/jshap/754

Holtkamp, D., Polson, D. D., Torremorell, M., Classen, D. M., Becton, L., Henry, S., Rodi-

baugh, M. T., Rowland, R. R., Snelson, H., & Straw, B. (2011). Terminology for

classifying swine herds by porcine reproductive and respiratory syndrome virus

status. *Journal of Swine Health and Production*, *19*(1), 44–56.

Kiss, I., Sámi, L., Kecskeméti, S., & Hanada, K. (2006). Genetic variation of the prevailing

porcine respiratory and reproductive syndrome viruses occurring on a pig farm

upon vaccination. *Archives of Virology*, *151*, 2269–2276.

Kristensen, C. S., Christiansen, M. G., Pedersen, K., & Larsen, L. E. (2020). Production

losses five months after outbreak with a recombinant of two PRRSV vaccine

strains in 13 Danish sow herds. *Porcine Health Management*, *6*(1), 26.

https://doi.org/10.1186/s40813-020-00165-z

Kvisgaard, L. K. (2017). Genetic and biological characterization of a Porcine Reproduc-

tive and Respiratory Syndrome Virus 2 (PRRSV-2) causing significant clinical dis-

ease in the field. *Veterinary Microbiology*.

Lopes Antunes, A. C., Halasa, T., Lauritsen, K. T., Kristensen, C. S., Larsen, L. E., & Toft,

N. (2015). Spatial analysis and temporal trends of porcine reproductive and res-

piratory syndrome in Denmark from 2007 to 2010 based on laboratory submis-

sion data. *BMC Veterinary Research*, *11*(1), 303. https://doi.org/10.1186/s12917-

015-0617-0

Lopes Antunes, A. C., Jensen, V. F., & Toft, N. (2019). Outcomes from using mortality, an-

timicrobial consumption, and vaccine use data for monitoring endemic diseases

in danish swine herds. *Frontiers in Veterinary Science*, *6*, 41.

Lunney, J. K., Benfield, D. A., & Rowland, R. R. R. (2010). Porcine reproductive and respir-

atory syndrome virus: An update on an emerging and re-emerging viral disease of

swine. *Virus Research*, *154*(1–2), 1–6. https://doi.org/10.1016/j.vi-

rusres.2010.10.009

Mortensen, S. (1996). PRRS antibodies in Danish swine. *Vet. Inform*, *1*, 35–36.

Mortensen, S., & Strandbygaard, B. (1996). Spread of PRRS slowed down. *Vet. Inform*, *4,7*.

Mortensen, S., Stryhn, H., Søgaard, R., Boklund, A., Stärk, K. D. C., Christensen, J., & Willeberg, P. (2002). Risk factors for infection of sow herds with porcine repro- ductive and respiratory syndrome (PRRS) virus. *Preventive Veterinary Medicine*, *53*(1), 83–101. https://doi.org/10.1016/S0167-5877(01)00260-4

Murtaugh, M. P., Elam, M. R., & Kakach, L. T. (1995). Comparison of the structural pro- tein coding sequences of the VR-2332 and Lelystad virus strains of the PRRS vi- rus. *Archives of Virology*, *140*(8), 1451–1460. https://doi.org/10.1007/BF01322671

Nieuwenhuis, N., Duinhof, T. F., & Van Nes, A. (2012). Economic analysis of outbreaks of porcine reproductive and respiratory syndrome virus in nine sow herds. *Veteri- nary Record*, *170*(9), Article 9. https://doi.org/10.1136/vr.100101

OIE. (2008). *The OIE AD HOC group on porcine reproductive respiratorysyndrome. PRRS: the disease, its diagnosis, prevention and control[Internet]. Group 2008 [cited 2021 Oct 15]. Available from*. https://www.woah.org/filead- min/Home/eng/Our_scientific_expertise/docs/pdf/PRRS_guide_web_bulletin.pdf

Pamornchainavakul, N., Makau, D. N., Paploski, I. A. D., Corzo, C. A., & VanderWaal, K. (2023). Unveiling invisible farm-to-farm PRRSV-2 transmission links and routes through transmission tree and network analysis. *Evolutionary Applications*, *16*(10), 1721–1734. https://doi.org/10.1111/eva.13596

Rahe, M. C., & Murtaugh, M. P. (2017). Effector mechanisms of humoral immunity to

porcine reproductive and respiratory syndrome virus. *Veterinary Immunology*

and Immunopathology, *186*, 15–18.

https://doi.org/10.1016/j.vetimm.2017.02.002

Rathkjen, P. H., & Dall, J. (2017). Control and eradication of porcine reproductive and

respiratory syndrome virus type 2 using a modified-live type 2 vaccine in combi-

nation with a load, close, homogenise model: An area elimination study. *Acta*

Veterinaria Scandinavica, *59*(1), Article 1. https://doi.org/10.1186/s13028-016-

0270-z

Rowland, R. R. R., & Morrison, R. B. (2012). Challenges and Opportunities for the Con-

trol and Elimination of Porcine Reproductive and Respiratory Syndrome Virus:

PRRS Control and Elimination. *Transboundary and Emerging Diseases*, *59*, 55–

59. https://doi.org/10.1111/j.1865-1682.2011.01306.x

Schulz, J. (2019). *Simulation modelling of LA-MRSA dispersal and control between*

swine herds.

SEGES Innovation. (2025). *Causes of PRRS status decline in 2019–2023*.

https://www.landbrugsinfo.dk/-/media/landbrugsinfo/public/4/3/0/ar-

sager_prrs_statusfald_2019_2023_notat2503.pdf

SPF. (2024, July 18). *Specific Pathogen Free System (SPF - Danmark)*.

https://spfsus.dk/en

State Serum Institut. (2023). *News regarding PRRS investigations*.

https://www.vetssi.dk/vet-nyheder/2023/nyt-vedrorende-prrs-undersogelse

Strategy for the reduction of PRRS in pigs in Denmark (p. 11). (2022). Danish Agrticulture

and Food Council.

Sundhedsstyringen. (2020). *The register of PRRS-infected swine herds in Denmark*.

Thakur, K. K., Revie, C. W., Hurnik, D., Poljak, Z., & Sanchez, J. (2015). Simulation of be-

tween-farm transmission of porcine reproductive and respiratory syndrome virus

in Ontario, Canada using the North American Animal Disease Spread Model. *Pre-*

ventive Veterinary Medicine, *118*(4), 413–426. https://doi.org/10.1016/j.prevet-

med.2015.01.006

Thakur, K. K., Sanchez, J., Hurnik, D., Poljak, Z., Opps, S., & Revie, C. W. (2015). Develop-

ment of a network based model to simulate the between-farm transmission of

the porcine reproductive and respiratory syndrome virus. *Veterinary Microbiol-*

ogy, *180*(3–4), 212–222. https://doi.org/10.1016/j.vetmic.2015.09.010

Valdes-Donoso, P., Alvarez, J., Jarvis, L. S., Morrison, R. B., & Perez, A. M. (2018). Produc-

tion Losses From an Endemic Animal Disease: Porcine Reproductive and Respir-

atory Syndrome (PRRS) in Selected Midwest US Sow Farms. *Frontiers in Veteri-*

nary Science, *5*, 102. https://doi.org/10.3389/fvets.2018.00102

Valdes-Donoso, P., & Jarvis, L. S. (2022). Combining epidemiology and economics to as-

sess control of a viral endemic animal disease: Porcine Reproductive and Respir-

atory Syndrome (PRRS). *PLOS ONE*, *17*(9), Article 9. https://doi.org/10.1371/jour-

nal.pone.0274382

Velasova, M., Alarcon, P., Williamson, S., & Wieland, B. (2012). Risk factors for porcine

reproductive and respiratory syndrome virus infection and resulting challenges

for effective disease surveillance. *BMC Veterinary Research*, *8*(1), 184.

https://doi.org/10.1186/1746-6148-8-184

Willkan, M. (2025). *Quantification of early detection surveillance in PRRS-free regions*.

Willkan, M., Chapot, L., Cameron, A., Boldsen, S. K., & Fertner, M. (2025). Quantification

of early detection surveillance in PRRS-free regions. *Preventive Veterinary Medi-*

cine, 106461. https://doi.org/10.1016/j.prevetmed.2025.106461

Supplementary Materials

S1. Illustration of the SPF system movement restriction

850
851 Figure S1. Illustration of the movement restriction in the SPF system. Herd owners can choose
852 to follow this scheme to maintain their PRRS status. However, purchasing pigs from herds with
853 a lower status will result in losing their PRRS status.

S2. Sensitivity analysis for the herd sensitivity of the clinical checkups

857
858 Figure S2. Sensitivity analysis of herd-level sensitivity for clinical checkups. The default herd

859 sensitivity is 0.30 for sow herds and 0.05 for weaner/finisher herds (Willkan et al., 2025). In

860 the alternative scenario, herd sensitivity is reduced to 0.07 for sow herds and 0.015 for

861 weaner/finisher herds (SEGES Innovation, 2025). The lower sensitivity values result in a

862 slightly higher proportion of positive herds detected compared to the default setting.

S3. Confidence interval of estimated transmission rate parameters

866 *Table S1. Estimation of transmission rate parameter for local spread*

Parameter	Definition	Point estimation	95% confidence interval bound
$\boldsymbol{\beta_{ls}}$	Local spread transmission rate parameter for highly infectious herds	1.58e-3	(3.1e-4, 3.0e-3)
$\boldsymbol{K_{ls}\beta_{ls}}$	Local spread transmission rate parameter for lowly infectious herds	1.74e-5	(2.53e-4, 3.67e-4)

S4. Calculation of R_0

For each transmission route, the basic reproduction ratio can be calculated using the
transmission matrix, using $R_{local\ spread}$ as an example.

The transmission matrix is $T = \begin{bmatrix} \beta S & \beta k S & \beta k S \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

The transition matrix is $\Sigma = \begin{bmatrix} -\gamma - \omega & 0 & 0 \\ \gamma & -\omega & 0 \\ \omega & \omega & -\alpha \end{bmatrix}$

The inverse of the transition matrix is $\Sigma^{-1} = \begin{bmatrix} \frac{-1}{(\gamma+\omega)} & 0 & 0 \\ \frac{-\gamma}{\omega(\gamma+\omega)} & -\frac{1}{\omega} & 0 \\ -\frac{1}{\alpha} & -\frac{1}{\alpha} & -\frac{1}{\alpha} \end{bmatrix}$

$$R_{local\ spread} = -T\Sigma^{-1} \begin{bmatrix} \beta S & \beta k S & \beta k S \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{(\gamma+\omega)} & 0 & 0 \\ \frac{\gamma}{\omega(\gamma+\omega)} & \frac{1}{\omega} & 0 \\ \frac{1}{\alpha} & \frac{1}{\alpha} & \frac{1}{\alpha} \end{bmatrix}$$

The next generation matrix to calculate $R_{local\ spread} = \frac{\beta N}{(\gamma+\omega)} + \frac{k\beta N}{\omega}\frac{\gamma}{(\gamma+\omega)} + \frac{k\beta N}{\alpha}$. Similarly,

for the movement route, $R_{movement} = \frac{P_{mov}c}{(\gamma+\omega)} + \frac{k_{mov}P_{mov}c}{\omega}\frac{\gamma}{(\gamma+\omega)} + \frac{k_{mov}P_{mov}c}{\alpha}$.

S5. Between herd R_0 distribution

Figure S5. Distribution of the between-herd basic reproduction ratio R_0 for all herds in the study population. A) total between-herd R_0; B) $R_{local\ spread}$; C) $R_{movement}$

In the study population, 57% of herds have a between-herd reproduction ratio R_0 greater than 1. Local spread alone results in 47% of herds infecting more than one herds ($R_{local\ spread} > 1$). In comparison, fewer than 0.2% of herds have $R_{movment} > 1$, as the average number of trading partners per herd is below 1 under current risk-based trading compliance.

S6. Simulated true prevalence under three scenarios

Figure S6. Impact of different intervention strategies on the true prevalence across three scenarios: A) Scenario 1, B) of Scenario 2, C) Scenario 3. Each line represents a specific intervention on combination of interventions, with colors corresponding to the legend on the right. The y-axis shows the percentage of positive herds, and the x-axis represents time in years. Shaded areas indicate variation across 100 model simulations. Interventions include biosecurity improvements, risk-based trading, depopulation-repopulation (DPRP) strategies, and combinations. All interventions are implemented starting from year 0.