**Title**: RNA decay regulates RNA abundances and transcriptome homeostasis using different mechanisms

Authors: Reed S. Sorenson and Leslie E. Sieburth*
School of Biological Sciences, University of Utah, Salt Lake City, Utah 84112
* correspondence: sieburth@biology.utah.edu

## SUMMARY

RNA decay is essential for maintenance of normal RNA abundances, however RNA abundances are regulated by two opposing processes, transcription and decay, yet how decay contributes to abundance change is poorly understood. Here, we measured rates of RNA abundance change, RNA half-lives ($t_{1/2}$s), and transcription rates in stimulated Arabidopsis leaf cells. This revealed two mechanisms in which decay influenced RNA abundance changes. The biggest changes in RNA abundances resulted from $t_{1/2}$ changes that reinforced transcriptional regulation (synergistic). Smaller RNA abundance changes arose from a second mechanism in which $t_{1/2}$ changes opposed transcriptional regulation (oppositional). RNA decay also contributed to stimulus-induced buffering of RNA abundances. Oppositional and buffering regulation shared excessive rate changes where changes in decay rates were commensurate and opposing transcriptional changes. We call this common mechanism countercyclical regulation. Its mirroring of rate changes implies coordination between nuclear and cytoplasmic processes, and it is widespread, accounting for 90% of the RNA $t_{1/2}$ regulation.

**Keywords**: RNA buffering, RNA decay, transcription, RNA $t_{1/2}$, countercyclical, nuclear-cytoplasmic coordination, procambium, vascular development, vascular transdifferentiation

**INTRODUCTION**

Cells respond to developmental and environmental perturbations in part by altering messenger RNA (mRNA) abundances. The identity of responsive RNAs can be highly informative; they can indicate developmental status or perception of environmental stress, and can be used to test hypotheses, for example in working out gene regulatory networks. However, despite RNA abundances being governed by just two processes, transcription and decay, we know relatively little about RNA decay's contributions. An implicit assumption is that mRNA abundances go up because transcription rates (TRs) increase and perhaps RNA decay rates ($t_{1/2}$s) diminish, and that mRNA abundances go down because TRs diminish and perhaps RNA decay rates increase. However, whether transcription and decay work together in this way, and how frequently RNA stabilities are regulated to contribute to RNA abundance regulation, is largely unknown.

Decay of RNA is a highly regulated process; it initiates by removal of 5' and 3' structures that confer stability (5' cap and 3' tail) or through internal cleavage. The remaining terminal nucleotides are then vulnerable to exoribonucleases such as XRN1/XRN4[1,2] or the RNA exosome[3]. RNA stability is regulated by a variety of mechanisms, e.g., signal transduction can activate decay machinery by phosphorylation[4,5]; DNE1 can initiate decay by internal cleavage [6–8]; and RNA binding proteins or RNA modification can influence recruitment of decay machinery[9–13]. In addition, RNA localization such as sequestration in processing bodies or stress granules[14–16] can also affect stability. However, despite understanding these mechanisms of regulation, the extent and manner in which RNA decay contributes to changing RNA abundances is still poorly understood.

A few studies have addressed the contributions of RNA decay regulation to gene expression. In animals, 20-30% of RNAs are regulated by the combined influences of transcription and decay, whereas in yeast decay contributes much more widely[17–23]. Studies using plants, though typically focused on small sets of genes, have demonstrated decay-influenced abundance regulation in response to time-of-day[24], iron-mediated oxidative stress[25], boron treatment[26], osmotic stress [5,27], and temperature[28–31]. However, a systems-level view of transcription and decay interactions in RNA abundance regulation is still lacking.

A challenge for understanding RNA abundance regulation is that its two regulatory processes, transcription and decay, mostly occur in separate compartments. Nevertheless, there is evidence that these processes can be coordinated. Best known is RNA buffering, which was discovered in a *Saccharomyces cerevisiae* mutant with an RNA polymerase II defect. This mutant was found to have normal RNA abundances because of compensatory changes in mRNA decay rates[32]. RNA buffering is now known to be widespread in yeast genotypes with defects in transcription and decay[33,34]. The mechanism underlying RNA buffering is not fully understood, but shuttling of RNA decay enzymes between the nucleus and cytoplasm, especially XRN1, is a central component[33,35–38]. RNA buffering has also been identified in Arabidopsis[39] and in animal cells[35,36,40,41]. For example,

in Arabidopsis, dysfunction of SUPPRESSOR OF VARICOSE (SOV)/DIS3L2 evoked RNA buffering; ~ 30% of the transcriptome underwent significant shifts in transcription and decay which allowed normal abundances to be maintained[39]. However, whether buffering-like coordination between transcription and decay extends to other modes of gene expression, including those leading to differential RNA abundances, remains unknown.

Here, we investigated the roles of RNA decay in abundance regulation using leaf cells as they responded to a developmental stimulus. We found widespread changes in RNA $t_{1/2}$s; 22% of the analyzed RNAs had differing mock and stimulus $t_{1/2}$. These $t_{1/2}$ changes affected RNA abundances using two distinct mechanisms. One was consistent with implicit assumptions about gene regulation; RNA $t_{1/2}$ changes worked alongside changes in transcription to bring about large changes in RNA abundances. We call this synergistic regulation, and it was rare. The other mechanism was widespread and featured $t_{1/2}$ changes that opposed the regulatory effect of transcription. These rate changes were excessive, sometimes dramatically so, and typically led to modest transcription-driven changes in abundance. We call this oppositional regulation. Similar and balanced excessive rate changes were observed for a set of RNAs whose abundances were steady and so appeared to be buffered. Oppositionally regulated and buffered RNAs were only distinguishable by their impact on RNA abundance, suggesting a common mechanism. Their reciprocal gene-specific regulation of transcription and decay suggested coordination between nuclear and cytoplasmic activities in a widespread mechanism that we call countercyclical.

## RESULTS

### A dynamic system for study of mRNA abundance regulation

To determine how RNA decay contributes to changes in mRNA abundances, we used a well-characterized leaf vascular transdifferentiation system[42–45] in which many mRNAs change abundances in response to a stimulus (Supplemental Data File 1). Developmental progression of the stimulus response was monitored by following expression of the preprocambial cell reporter *proAtHB8:4xYFP*[46–48]. *AtHB8* expression is normally activated in narrow cell files that later differentiate into organized leaf veins[49,50]. In stimulus-treated leaves, *proAtHB8:4xYFP* expression had expanded beyond these narrow cell files and by 72 h cells of interveinal regions had thick and lignified cell walls that resembled xylem tracheary elements (Supplemental Figure S1A, S1B, S1C). These observations were consistent with the system's described developmental progression and demonstrated reliable elevated *proAtHB8:4xYFP* fluorescence after 12 h of stimulus (Figure 1A). Vascular induction was confirmed by qPCR analysis of *AtHB8* and *DOF5.8*, another early vascular transcription factor (Figure 1B), while a later-expressing phloem-specific gene, *SMXL5*, showed peak expression later (between 12 and 16 h)[51–55]. These observations indicated that vascular-related gene expression was ectopically activated and proceeding along a

stereotypical timeline. Because the molecular signature of vascular induction was apparent at 8 h while cell morphologies had not fully differentiated (Supplemental Figure S1B) this timepoint's many mRNAs with changing abundances seemed suitable for analysis of RNA decay's roles in abundance regulation.

To characterize the developmental status of the 8 h stimulus treatment, we quantified mRNA abundances of stimulus and mock-treated leaf tissue using RNA-seq (Supplemental Data File 1). This yielded relative abundances of 18,152 nuclear-encoded mRNAs, 55% of which differed significantly in the stimulus and mock samples (FDR < 0.05) (Figure 1C, Supplemental Figure S1D). These abundance changes resembled early transdifferentiation responses in cotyledons as captured by a microarray time course[45], and differed from expression of mature vascular tissues of late stimulus responses[44] (Supplemental Figure S1E-S1F). A Gene Ontology analysis supported 8 h as early in vascular development because stimulus up-regulated RNAs were not enriched in genes of later vascular cell type differentiation (*phloem or xylem histogenesis* (GO:0010087). Up-regulated RNAs were also not enriched in *auxin−activated signaling* (GO:0009734), suggesting that the treated leaves were not simply responding to the stimulus cocktail (Figure 1D). Moreover stimulus down-regulated RNAs were enriched in the *response to light stimulus* (GO:0009416) and *thylakoid* (GO:0009579) GOs (73/209 and 143/148, respectively), consistent with mesophyll cell dedifferentiation (Supplemental Data File 2)[56].

We also compared 8 h RNA abundances to leaf cell types classified from single-cell RNA sequencing[57] (see methods, Supplemental Figure S1G, Supplemental Data Files 3 and 4). Stimulus-induced abundance changes included an increase in positive markers for the procambium cluster (no. 10) and a decrease in negative markers (Figure 1E), indicating an increase in procambium cells. Additionally, a strong reduction in negative markers of differentiating and mature vascular cell types (10, 18, 15, and 4) was observed. Moreover, positive markers of mesophyll cell-type clusters (clusters 2, 3, 5, 6) were lower in stimulus while negative markers were unchanged or higher (Figure 1F), consistent with the loss of mesophyll identity as the stimulus redirects development toward vascular cell types. Surprisingly, positive markers of mesophyll cell clusters 11 and 14 were as high or higher than procambium cell cluster 10, suggesting that these mesophyll cell subtypes might be procambial cell precursors. Together, the changes in cell identity with their accompanying shifts in gene expression at 8 h make this an attractive system for teasing out roles of mRNA decay in RNA abundance regulation.

### Stimulus induces widespread changes in mRNA $t_{1/2}$

To determine mRNA $t_{1/2}$s, we used the 8 h RNA-seq data described above along with RNA-seq abundances from co-collected tissue treated with the transcription inhibitor cordycepin for 15, 60, or 240 min (Figure 1G). Data were processed through our previously described RNA decay modeling pipeline, which accounts for potential feedback from transcription arrest using a modeling-based approach to estimate initial decay rates and.

also provided statistical support for gene-specific $t_{1/2}$ differences in stimulus and mock conditions[39]. We obtained RNA decay rate estimates for 15,083 transcripts. As seen in previous Arabidopsis RNA decay studies, $t_{1/2}$s were spread across 2.5 orders of magnitude[39,58–61] and $t_{1/2}$s ranged from 5.0 min (*SAUR21* in mock) and 4.6 min (*SAUR14* in stimulus) to ≥ 1,440 min, our capped maximum (Supplemental Data File 1).

We found that 22% of mRNAs (3,329) responded to the stimulus by changing RNA $t_{1/2}$ (Figure 1H). These $t_{1/2}$ shifts varied in their magnitude; nearly half (1,571) were greater than 2-fold, with120 RNAs greater than 10-fold (Figure 1I). Some $t_{1/2}$ changes were extraordinarily large. For example, $t_{1/2}$ lengthened 23-fold for RNAs of *CIP1* ($t_{1/2}$ mock = 63 min , $t_{1/2}$ stimulus = 1440 min) and AT5G42370 ($t_{1/2}$ mock = 63 min, $t_{1/2}$ stimulus = 1440 min) (Figure 1J), and shortened 33-fold for *CPK29* ($t_{1/2}$ mock ≥ 1,440 min, $t_{1/2}$ stimulus = 43 min) and 21-fold for *SAP1* ($t_{1/2}$ mock ≥ 1,440 min, $t_{1/2}$ stimulus = 70 min) (Figure 1K).

To assess trends in RNA $t_{1/2}$ shifts, we separated RNAs into two groups: one group included RNAs with $t_{1/2}$s in the standard range (2,818, mock and stimulus $t_{1/2}$ ≤ 1440 min) and a second group had least one extraordinarily long $t_{1/2}$ (511, $t_{1/2}$ ≥ 1440 min). RNAs in the standard $t_{1/2}$ range showed similar shifts to shorter and longer $t_{1/2}$ (shift to shorter: mean fold change 2.2, median 1.9; shifts to longer: mean fold change of 2.0, median of 1.8) (Figure 1L, Supplemental Figure S2A). Some of these RNAs had very different mock and stimulus $t_{1/2}$s; e.g., AT5G66580 shifted $t_{1/2}$ from 178 to 9 min, and AT4G33890, shifted from 60 to 1295 min. The smooth curvature of these plots indicated that RNAs across the full range of mock $t_{1/2}$s (2.5 orders of magnitude) underwent $t_{1/2}$ shifts in both directions, with exceptions being extremely short-lived RNAs, which tended to shift to longer $t_{1/2}$s (75 of the 100 shortest-lived RNAs made this shift) and extremely long-lived RNAs, which tended to shift to shorter $t_{1/2}$s (87 of the 100 longest-lived RNAs) (Supplemental Figure S2B). These patterns suggested that our data approached the upper and lower limits of Arabidopsis RNA $t_{1/2}$s.

The group of RNAs with at least one extraordinarily long $t_{1/2}$ included 435 RNAs whose standard mock $t_{1/2}$ shifted to ≥ 1,440 min, and 76 RNAs with a mock $t_{1/2}$ ≥ 1,440 min that shifted into the standard $t_{1/2}$ range (Figure 1J-1K and Supplemental Figure S2A and S2C). The magnitudes of these $t_{1/2}$ shifts were much larger than for RNAs in the standard range (mean fold change of 3.9, median of 2.8) and so might be entering or exiting sites of RNA sequestration, e.g., condensates such as stress granules or processing bodies[16,62–64].

Previously we noted that RNAs with rapid turnover were enriched in GOs termed responsive, e.g., to stress or signaling[39]. This suggested that short-lived RNAs might be likely to change abundances by altering their $t_{1/2}$, e.g., by shifting to longer $t_{1/2}$s. We addressed this idea by comparing $t_{1/2}$ changes across five quintiles of mock $t_{1/2}$. We found that most $t_{1/2}$ changes occurred among long-lived RNAs (Supplemental Figure S2D), leaving the purpose for short mRNAs $t_{1/2}$s an open question.

## Analytical approach refinement using an early vascular regulatory network

To sort out how best to determine the impact of RNA decay on RNA abundances, we compared two approaches using a subset of the data, a 65-gene early vascular regulatory network (EVR network) (Supplemental Table 1). We obtained RNA abundance and $t_{1/2}$ data for 57 of the EVR network genes, 46 (81%) of which were differentially expressed (Figure 2A and 2B). We first used abundance and $t_{1/2}$ to infer regulation (Supplemental Figure S3A). Thirty-nine RNAs had higher 8-h abundances in stimulus; 30 of these had unchanged $t_{1/2}$, suggesting that their abundance change was driven by transcription. The remaining included three RNAs that shifted to longer $t_{1/2}$s (e.g., *LBD4*) and six that shifted to shorter $t_{1/2}$s. The shift to longer $t_{1/2}$ likely contributed to the RNA's abundance change, but the more common outcome, shifts to shorter $t_{1/2}$ would have worked against their increased abundances (Figure 2C). For example, the abundance of *SMXL4* mRNA increased nearly five-fold while its $t_{1/2}$ shortened. Seven RNAs had lower abundances; six of these had unchanged $t_{1/2}$s (suggesting transcription regulation), and one RNA shifted to a longer $t_{1/2}$, working against its lowered abundance (Supplemental Figure S3A, Supplemental Data File 5). A shortcoming of this approach was that the 8 h $t_{1/2}$ measurement was being compared to an abundance change that could have occurred at any time prior to 8 h, limiting regulatory inferences.

Our second approach identified RNAs with actively changing abundances across the 8 h timepoint; this used RNA abundances (RNA-seq) from 6, 7, 8, 9 and 10 h (Figure 2D) and linear regression to estimate slopes of RNA abundance change. These additional data allowed us to focus on the RNAs whose abundances were actively changing and also allowed us to calculate TR for each RNA as the amount of synthesis that was required to sustain the rate of abundance change (RoAC) occurring at 8 h, given the $t_{1/2}$ [19,65,66] (Supplemental Figure S3B, S3C, S4A). These data showed that about a third of the RNAs had significant rates of abundance change (i.e., not 0, $p < 0.05$). These rates were broadly distributed with most mRNAs changing abundance in a single condition (85%), and 15% (751) changing in both mock and stimulus (Figure 2E-2G).

We re-evaluated regulation of the 57 EVR network genes using these data; 23 of these RNAs had actively changing abundances ($p<0.05$)(Figure 2H). Among these 23, 19 had indistinguishable mock & stimulus $t_{1/2}$s, indicating that transcription regulation alone was driving their changing abundances. This defined a transcriptional regulatory modality (e.g., *SMXL5* and TMO6) (Figure 2I). Another gene, *PXC1*, defined an RNA decay regulatory modality; its RNA much shorter stimulus $t_{1/2}$ drove its actively changing abundance as its stimulus and mock TRs were similar (i.e., below our difference cutoff, Supplemental Figure S3B, S4B, Figure 2J). Two additional regulatory modalities featured changes in both RNA $t_{1/2}$s and TRs. *LBD4* defined a synergistic regulatory modality in which faster TR and longer $t_{1/2}$ both contributed to its increasing abundance (Figure 2K). *ATHB8* and *SMXL4* defined an oppositional regulatory modality. These RNAs had increasing abundances and faster TRs, however unlike synergistic, their RNA $t_{1/2}$s became shorter (Figure 2L) indicating the $t_{1/2}$ change opposed transcription regulation. Oppositional regulation could be quite inefficient, e.g., *SMXL4*'s increasing abundance required a 1.66-fold faster TR to counter its shorter $t_{1/2}$ (17.1 RPKM h$^{-1}$ vs 10.3 RPKM h$^{1}$ without the change in RNA $t_{1/2}$). Because

including RoAC data provided greater insight into the roles of RNA decay, we extended this second analytical strategy to the whole transcriptome, using the 14,600 genes with TRs, $t_{1/2}$s and RoAC data.

## Transcriptome-wide Analysis reveals RNA buffering as an additional regulatory modality

Because the EVR network analysis identified several regulatory modalities contributing to actively changes abundances (transcription, decay, synergistic, and oppositional), we next determined their transcriptome-wide usage. Of the 4,268 RNAs with actively changing 8 h abundances, 3,108 had indistinguishable mock and stimulus $t_{1/2}$s, indicating that their RNA abundances were  transcriptionally regulated (Figure 3A). Transcription alone was the major modality of changing abundances (21% of the full dataset) and its regulation led to both faster (1,305 RNAs) and slower (1,803) rates of abundance change (Figure 3B). Only 64 genes encoded RNAs whose abundances were regulated by decay alone (Figure 3C) making it the least used regulatory modality. The remaining 1,096 genes had RNA abundances regulated by the combined influence of transcription and decay. Synergistic regulation was found for a small number of RNAs (106) (Figure 3D) whereas oppositional regulation was far more common (990 RNAs) (Figure 3E).

We also found $t_{1/2}$ regulation among RNAs with steady abundances (p > 0.05). One group of RNAs had differing $t_{1/2}$s and abundances, which we called steady 1 (1126 RNAs)(Figure 3F). Another group had differing $t_{1/2}$s but abundances were indistinguishable (steady 2, 920 RNAs (Figure 3G and Supplemental Figure S4C). These combined features – changes in TR and $t_{1/2}$ but not abundance – fit the operational definition of RNA buffering, so we called steady 2 buffered. This was unexpected because RNA buffering has typically been observed in systems with dysfunction in transcription or decay machinery[33–36,40,41].

In total, this analysis led to recognition of 9,714 RNAs with steady abundances (p > 0.05). In addition to the two modalities described above, we also found RNAs for which $t_{1/2}$s and abundances were indistinguishable (steady 3, 3455 RNAs)(Figure 3H) and RNAs for which $t_{1/2}$s were indistinguishable but mock and stimulus abundances differed (steady 4, 4213 RNAs)(Figure 3I). Because steady 4's abundances were not actively changing, the regulation leading to these abundance changes occurred prior to our experimental window. RNAs with differing abundances were maintained through different TRs (steady 4, Figure 3I) or by different RNA $t_{1/2}$s with or without TR changes (steady 1, Figure 3F).

## Regulatory modalities and rate specialization

As a step towards discovering whether any modalities had unique properties, we compared dynamic ranges of mock and stimulus TRs and $t_{1/2}$ for each modality. In all modalities, TRs and $t_{1/2}$s were distributed across their full dynamic range (Supplemental Figure S5A-S5E). This indicated that none of the modalities were specialized for fast or slow rates (TR and/or $t_{1/2}$).

To examine whether modalities differed in the extent to which TRs and $t_{1/2}$s changed, we determined the mean TR rate change and $t_{1/2}$ change for each (Table 1 and Supplemental Figure S5A-S5E). Comparing modalities, we found similar rate changes for transcription and decay. This indicated that these two processes had similar capacity to impact RNA abundance. Four modalities featured changes in both transcription and decay: synergistic, oppositional, steady 1, and buffered (steady 2). Synergistically regulated RNAs underwent small changes in TR and $t_{1/2}$. By contrast, oppositionally regulated, steady 1 and buffered (steady 2) RNAs underwent large changes in TR and $t_{1/2}$. Thus, modalities do appear to have specializations for rate changes.

**Synergistic and oppositional modalities use distinct mechanisms**

We next explored gene-specific relationships between $t_{1/2}$ and TR across the modalities. The simplest were transcription and decay; both of these modalities had rate changes that ranged from modest to large (>10-fold for TR and >8-fold for $t_{1/2}$) (Figure 4A-4B).

For modalities with changes to both TRs and $t_{1/2}$s, the gene-specific ranges of TR and $t_{1/2}$ were represented by stacked bar plots. In the synergistic modality, the cooperativity of changes in TR and $t_{1/2}$ was represented by bars stacked on the same side of the X axis (Figure 4C). Despite the cooperative directionality of rate changes, magnitudes of TR and $t_{1/2}$ changes differed, suggesting that these processes were regulated independently. Changes in $t_{1/2}$ provided 65% of the total regulatory changes; that is, RNA abundance regulation of most synergistically regulated genes was driven predominantly by gene-specific change in $t_{1/2}$s.

The oppositional modality's stacked bar plots had a strikingly different appearance. The gene-specific ranges of TR and $t_{1/2}$ showed high gene-to-gene variability. These changes were often very large, far exceeding that required for net regulation (the sum of $\log_2$ relative $t_{1/2}$ and TR, indicated as black dots) (Figure 4D). In addition, for each gene, TR and $t_{1/2}$ underwent very similar rate changes (i.e., mirrored). In other words, oppositional decay was remarkable for its frequent large and opposing changes in transcription and decay that resulted in small net regulation. Mirroring of gene-specific rate changes was most obvious in this plot's central region where net regulation was smallest (Supplemental Figure S5F-S5H). In the upper and lower 10% of net regulation, 78% of RNAs had larger changes in TR than $t_{1/2}$. In other words, in oppositional regulation changing abundances can be driven by transcription or decay, but in this study, transcription was the more frequent driver.

The synergistic and oppositional modalities were similar in that both brought about changes in RNA abundances. However, they appear to do it in distinct ways. $t_{1/2}$ regulation was independent of transcription for RNAs in the synergistic modality. By contrast, $t_{1/2}$ regulation was tightly linked to regulation of TR in the oppositional modality. These features suggest that synergistic and oppositional regulation are mechanistically distinct.

## Opposing transcription and decay regulation is a common strategy for transcriptome homeostasis

Two additional modalities, buffered (steady 2) and steady 1, had steady abundances and both TR and $t_{1/2}$ differed in the mock and stimulus. Comparison of magnitudes of TR and $t_{1/2}$ change in Table 1 suggested that these modalities were similar to oppositional. To assess whether buffered (steady 2) and steady 1 shared oppositional's other features, we extended the analysis of gene-specific rate changes (TR and $t_{1/2}$). Similar to oppositional, we found large ranges, with high gene-to-gene variability and commensurate changes in TR and $t_{1/2}$ (Figures 4E-4F). This suggests that these three modalities, oppositional, buffered (steady 2), and steady 1, share a common mechanism.

To assess whether buffered RNAs in general might share these same features, we re-analyzed a set of buffered RNAs previously identified in the Arabidopsis *sov/dis3l2* mutant[39]. We examined these data using stacked bar plots and observed that magnitudes of rate changes showed high gene-to-gene variability (Figure 4G). In addition, gene-specific changes in TR and $t_{1/2}$ had similar magnitudes. This supports buffering being part of a larger shared mechanism that includes regulation leading to altered RNA abundances.

RNAs buffered in *sov* mutants showed the kinetic features of an overcompensating mechanism[39]; buffered RNAs that had shorter $t_{1/2}$s also had slightly higher abundances (insignificant) and RNAs that had longer $t_{1/2}$s had slightly lower abundances (insignificant) (Supplemental Figure S5I). Overcompensation was also observed for mouse RNAs buffered following chromatin disruption[67]. To assess whether this study's buffered RNAs also exhibited overcompensating feedback, we examined the pattern of $t_{1/2}$ changes for RNAs with small (insignificant) abundance changes. Similar numbers of the buffered (steady 2) RNAs shifted to shorter and longer half-lives in the higher and lower abundance groups (Supplemental Figure S5J). The apparent lack of over-compensating feedback in buffered (steady 2) RNAs raises the possibility of distinct regulatory mechanisms for RNAs buffered as a consequence of regulatory dysfunction versus RNAs buffered as part of a biological process.

## Functional roles of regulatory modalities

We considered whether modalities might be used for specific functions, for example, during stages of a wave of differential expression. At the beginning of a wave of expression abundances increase, and after peaking abundances decrease, and waves of expression occur broadly, including during vascular transdifferentiation[45,51,52]. We identified RNAs in these three phases by clustering the 4,268 RNAs with changing abundances based on abundance profiles across the 6, 7, 8, 9, 10 h timepoints. We sorted 24 clusters into three groups: diverging abundances (1,366-genes in 11 clusters), parallel abundances (1,220 genes, 6 clusters) and converging abundances (1,682-genes, 7 clusters)(Figure 5A). We found strong enrichment of synergistically regulated RNAs in the diverging clusters, and strong depletion of oppositionally regulated RNAs in the converging clusters (Figure 5B).

This suggests roles for both synergistic and oppositional modalities in RNAs with diverging abundances.

If these modalities are important for diverging abundances, RNAs in the synergistic and oppositional modalities might show substantial changes in abundance. Although this analysis was complicated by abundances that were in the process of changing, on average RNAs in the synergistic modality showed the largest abundance changes (Figure 5C). Together, these data support the synergistic modality as the most important in effecting large changes in RNA abundance. Furthermore, because abundances of synergistic regulated RNAs were primarily driven by decay, RNA decay emerges as an important regulator of RNA abundances.

We also used GO enrichment analysis to assess specialized functions of the modalities[68]. The strongest of these were for RNAs in steady 4, which had differing stimulus and mock abundances that arose prior to the experimental window and maintained the same $t_{1/2}$ (Supplemental Figure S5K). Enriched in the RNAs related to ribosome constituents and translation, suggesting that ribosomal protein RNA abundances were adjusted early and without contributions by decay regulation.

RNA decay pathways are known to have specialized function[39,59,69], and we considered that there might be specialized contributions to modality regulation. To test this idea, we used the metric of $t_{1/2}$ dependence on VCS/EDC4, the scaffold of the mRNA decapping complex (previously determined[39]). We considered enrichment of three sets of genes: sets with high (stabilized > 2-fold in *vcs* mutants) and moderate (stabilized < 2-fold in *vcs* mutants) VCS dependence, and a set which were strongly destabilized in *vcs* mutants which we referred to as feedback regulated[39]. The RNAs in steady 4 were enriched in both the high and moderate VCS-dependent RNAs, indicating an important role for mRNA decapping in facilitating maintenance of steady RNA abundances as TRs change (Figure 5D). The strongest enrichment was for buffered (steady 2) RNAs; these were enriched in *vcs* feedback-regulated RNAs. Thus, this group includes RNAs that appear to have a propensity for feedback regulation.

## DISCUSSION

Measurable changes in mRNA abundances, commonly referred to as gene expression, are the outcomes of changes in rates of transcription and RNA decay. Here, we demonstrated that regulated RNA decay rates played important roles in RNA abundance control by driving changing abundances in the synergistic and oppositional modalities, and in buffering RNA abundances during a stimulus response. For the synergistic modality, RNA $t_{1/2}$ shifts reinforced transcription. Decay was the main driver of changing abundances and resulted in the largest abundance changes. For the oppositional modality, RNA $t_{1/2}$ shifts opposed transcriptional regulation; transcription was the main driver of RNA abundance

change but decay-regulated RNAs were also observed. That is, typically changes in decay tempered large changes in TRs and resulted in relatively small changes in abundances. Because magnitudes of transcription and decay rate changes were mirrored, these two processes appeared linked. Two additional regulatory modalities, buffered (steady 2) and steady 1, showed similar features, suggesting use of a widespread shared mechanism. Together, these accounted for 90% of the RNAs with regulated $t_{1/2}$s; we call this shared mechanism countercyclical regulation.

Synergistic and oppositional relationships between rates of transcription and decay appear to have specialized usage. In yeast, these modalities showed stress-specific roles; response to DNA damage included strong use of synergistic regulation while responses to osmotic and oxidative stress predominantly used oppositional regulation[17,70,19]. Similarly, studies using drosophila and mouse cells found widespread use of oppositional regulation while synergistic appeared to have more specialized functions, e.g., in cell-type specific regulation[17,19,71,72]. Here, the synergistic modality was used for large changes in expression, including in genes whose RNAs were becoming differentially regulated. Consistent the idea that modalities are utilized during specific responses, oppositional regulation is used as RNAs change in response to specific stimuli in yeast and animals cells[19,73]. In our study, synergistic and oppositional regulation also played roles in distinct phases of gene expression, such as the sequential waves of gene expression associated with vascular transdifferentiation[74,50,75,44,51,52]. RNAs with diverging abundances were strongly enriched in both synergistically and oppositionally regulated RNAs, in contrast to RNAs with converging abundances. Whether these observations of stress- and phase-specific use of regulatory modalities are general awaits further study.

Countercyclical regulation of RNAs was apparent in the oppositional, buffered (steady 2) and steady 1 regulatory modalities. We initially focused on RNA abundance change and placed oppositional and buffered RNAs in distinct groups. However, by prioritizing their regulation, they could be arrayed across a continuum, with the changing abundances of oppositionally-regulated RNAs defining the ends and with buffered RNAs occupying the center. Steady 1 RNAs appeared to have similar regulation because of their large and opposing changes in transcription and decay. This suggests that their different abundances might have arisen from a countercyclical mechanism, and that these changes were durable. Countercyclical regulation appeared to be gene specific; magnitudes of TR and $t_{1/2}$ changes differed between regulated genes and only a portion of the transcriptome was regulated in this manner. Gene-specific buffering also occurred in *sov* mutants, where 42% of the transcriptome was affected[39]. This contrasts with the global nature of RNA buffering in yeast[76] and is consistent with the recent report of gene-specific buffering in animals[67]. It also contrasts with synergistic regulation, which showed gene-specific changes in decay rates that were unlinked to transcription, and so mechanistically distinct from countercyclical.

A future challenge is determining the mechanism of countercyclical regulation, which appears to require nuclear-cytoplasmic coordination. One way that nuclear and

cytoplasmic processes are coordinated is through shuttling of proteins between these compartments. In yeast, XRN1 shuttling plays an important role in RNA buffering[33,35–37,77]. Cytoplasmic XRN1 degrades decapped and fragmented mRNAs[36] and shuttles into the nucleus in a decay-dependent manner. Nuclear XRN1 binds DNA and contributes to gene expression[77]. However, because plants lack a cognate homolog of XRN1 and instead use the XRN4 paralog which lacks key nuclear localization signals[61,77,78], this model is unlikely to apply to plants. Other potential mechanisms, but unexplored in plants, include shuttling of signals dependent on 3' to 5' decay[41], mRNA imprinting (e.g., with Rpb4 and Rpb7)[32,70,79–81], gene-specific mRNA methylation linked to transcription elongation rates[72], or other undiscovered mechanisms of coordination such as those proposed by Wang and Lin[82].

Homeostasis of mRNAs is essential for normal cellular function, as underscored by its exquisite regulation during mammalian cell expansion[40,83–85]. Countercyclical regulation is a mechanism for transcriptome homeostasis because it reins in RNAs whose TRs and/or $t_{1/2}$s are excessive and places regulated decay at the center of transcriptome homeostasis. In yeast, RNA buffering is the key homeostatic mechanism; it maintains stereotypical mRNA abundances when components of the transcription or decay machinery are dysfunctional[33,34,86–89]. Yeast also shows oppositional regulation, however whether it is mechanistically similar to yeast RNA buffering is unknown. Countercyclical-like regulation has also been documented in drosophila and mammalian cells[33,34,71,72,86–89] and so appears to be an evolutionarily conserved mechanism for transcriptome homeostasis.

## Acknowledgements

We thank the Arabidopsis Biological Resource Center for maintaining and sharing genetic material. We thank Dominique Gagliardi and members of the RNA degradation group at the Institut de Biologie Moléculaire des Plantes (IBMP) in Strasbourg, France and Sally Assman from the Pennsylvania State University Huck Institute of the Life Sciences for critical suggestions. This work was supported by the National Science Foundation under award numbers 1838345 and 1616779. R.S.S. was supported by an NIH Developmental Biology Training Grant (5T32 HD07491). Research reported here used the High-Throughput Genomics and Cancer Bioinformatics Shared Resource at Huntsman Cancer Institute at the University of Utah, supported by the National Cancer Institute of the National Institutes of Health under award number P30CA042014. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

## Data availability

Sequencing data for this study can be accessed at the Gene Expression Omnibus (https://www.ncbi.nlm.nih.gov/geo/) under series accession number GSE278544.

## Author contributions

Conceptualization, Funding Acquisition, Methodology, Writing, R.S.S. and L.E.S.; Investigation, Formal Analysis, Data Curation, Visualization, R.S.S.; Supervision, L.E.S.

**Limitations of the study**

RNA $t_{1/2}$ estimates could not distinguish cytoplasmic RNA decay from decay that occurs in the nucleus and so the extent to which symmetric changes in transcription and decay rates reflected coordination between cytoplasmic and nuclear compartments is unknown. $t_{1/2}$ measurements used leaf tissue, which resulted in the averaging over multiple cell types. Rates of abundance change were based on best linear fit across three to five timepoints, and a non-parametric fit might have yielded different estimates with potential impacts extending to TR calculations. While this study identifies transcriptome homeostasis as an important role of RNA decay, its potential link to scaling of mRNA abundance to cell size was not considered. Finally, this study's conclusions are based on analysis of a single timepoint of a response to a single stimulus which may limit the generality of our conclusions.

## METHODS

### Plant Genetic Material and Growth Conditions

*AtHB8::4xYFP* (Columbia-0 genetic background) [48] were received from Arabidopsis Biological Resource Center (CS2106110, Ohio State University). Seeds were sown on moistened soil (Sungro Redi-Earth Plug and Seedling propagation mix) and imbibed in the dark at 4°C. After 2-4 d, pots were grown under continuous illumination with 75 µmol photons $m^{-2} s^{-1}$ at 21°C for 10 d. Leaves 1 and 2 were used in all experiments unless otherwise indicated.

### Phloroglucinol Staining & Microscopy

Lignin staining was performed after clearing leaves in warm 70% ethanol. Cleared tissue was stained and imaged in phloroglucinol stain (prepared as a fresh 2:1 mixture of 3% w/v phloroglucinol in ethanol with concentrated hydrochloric acid (37 N))[90]. For DIC imaging, treated leaves were cleared using 30% acetic acid 70% ethanol overnight to 7 d, rinsed 3 times in 70% ethanol and transferred to ClearSee solution (10% w/v xylitol, 15% w/v sodium deoxycholate, 25% w/v urea)[91], incubated overnight and transferred to ClearSee with 12.5% glycerol for mounting and imaging. Microscopy was performed on an Olympus BX-50 compound microscope. DIC images were taken with the Olympus DP-74 camera. YFP fluorescent images were captured from leaves mounted in water and using the U-MWIB filter cube by the Olympus DP-70 camera.

### Developmental Stimulus Treatment and RNA Decay Analysis

For the developmental stimulus treatment, we used the protocol of Kondo et al.[43], and modified it for treatment to whole leaves. Detached leaves (~18-20/sample) were floated on distilled water during collection, followed by incubation in 10% bleach 0.1% Silwet L-77 for 3-5 min, and rinsing 3 times in sterile distilled water to surface sterilize. Sterile leaves were transferred to incubation buffer (0.22% Murashigie and Skoog salts (Caisson Laboratories) [pH 5.7], 5.0% glucose, 0.2% dimethyl sulfoxide) with (stimulus) or without (mock) 20 µM bikinin (Calbiochem), 2.5 µg mL$^{-1}$ 2,4-Dichlorophenoxyacetic acid (Sigma), and 0.5 µg mL$^{-1}$ kinetin (Sigma). Samples were incubated at 21°C with shaking (120 RPM) for designated times and collected. To inhibit transcription, leaves were incubated in the same buffer with 1 mM cordycepin (Chengdu Biopurify Phytochemicals) and incubated with shaking for the additional designated time. Time points were selected to be equidistant on a log scale that included a short time point (15 min) for higher accuracy of short $t_{1/2}$s. Following incubation, excess moisture was removed, and tissues were flash frozen, ground to powder and stored at -80°C until RNA isolation.

## RNA Sample processing

RNA was extracted using the Quick-RNA Plant mini-prep kit (Zymo) and Treated with DNase I. For quantitative polymerase chain reaction measurement of specific mRNA abundances, RNA was primed with random hexamers and reverse transcribed using the High Capacity cDNA Reverse Transcription (Applied Biosystems) kit. qPCR was performed on a C1000 Touch Thermal Cycler with the CFX-96 Real-Time Detection System (Biorad) using primers UBC10_qF: tcactggcaggcaacgataatgg, UBC10_qR: aagatgtcgaggcaaatgctaccg, DOF5.8_qF:cggtggtgtttcccgtaaaagc, DOF5.8_qR:cgtggcgggtatttgagcaatg, AtHB8_qF:caagggtcgcggagatcctaaag, AtHB8_qR:tctctgccctaacgaaatgaggaga, SMXL5_qF:taagccttcatgcgacaagtggac, SMXL5_qR:agaacgccccaaatcaagagtgac.

For RNA sequencing, total RNA was submitted to the Huntsman Cancer Institute High Throughput Genomic Core for library construction and sequencing. rRNAs were depleted from the samples using Ribo-Zero Plant. Stranded RNA sequencing libraries were prepared using the TruSeq Stranded Total RNA Kit (Illumina). For the RNA decay samples, purified libraries were evaluated using an Agilent Technologies ScreenTape assay. The molarity of adapter-modified molecules was defined by quantitative PCR using the Kapa Library Quant Kit (Kapa Biosystems). Individual libraries were normalized to 10 nM and equal volumes were pooled in preparation for Illumina sequence analysis. Sequencing libraries (25 pM) were chemically denatured and, for the RNA decay experiment samples, , applied to an Illumina HiSeq v4 single read flow cell. Hybridized molecules were clonally amplified and annealed to sequencing primers with reagents from an Illumina HiSeq SR Cluster Kit v4-cBot. Following transfer of the flowcell of an Illumina HiSeq 2500 instrument, a 50 cycle single-read sequence run was performed using HiSeq SBS Kit v4 sequencing reagents. For the time course experiment, libraries were transferred to the flowcell of an Illumina NovaSeq 6000 instrument, a 151 x 151 cycle paired end sequence run was performed using a NovaSeq 6000 S4 reagent Kit v1.5 (20028312).

## Bioinformatic Analysis

For analysis of RNA sequencing data, steps were performed within R using the *systemPipeR* package[92]. 50 bp reads were trimmed of adaptor sequences (Read 1 Adaptor: AGATCGGAAGAGCACACGTCTGAACTCCAGTCA; Read 2 Adaptor: AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT) using the *trimLRPatterns* function from the *Biostrings* package[93], and aligned to the TAIR10 genome sequence using *Tophat Bowtie2* allowing a single base mismatch[94,95]. Read counting was performed using the *summarizeOverlaps* function of the *GenomicAlignments* package[96] using gene models of Araport11_GFF3_genes_transposons.201606.gff (accessed at TAIR: www.arabidopsis.org/download). Differential Expression analysis was performed using the *edgeR* package[97], significant RNA abundance differences were filtered based on an FDR < 0.05. Gene alias (gene_aliases_20201231.txt ) and functional descriptions (Araport11_functional_descriptions_20201231.txt ), and gene ontology assignment (ATH_GO_GOSLIM.txt) files were also downloaded from TAIR. The gene ontology annotations file (go-basic.obo) was accessed at geneontology.org. Gene ontology enrichment was calculated using the *topGO* package[98]. Visualization of data used the *ggplot2* package[99].

## RNA Kinetic Analyses

RNA decay rates: RNA decay rates were estimated by modeling the measured RNA abundance decrease following treatment with cordycepin to inhibit transcription[39]. RNA counts were first normalized to the total number of read counts in the library and scaled to 1 million. Scaled read counts were further scaled to a normalized gene length of 1000 nt (RPKM values). Prior to modeling, low abundance RNAs were removed (sum of the mean T0 abundances (RPKMs) of mock and stimulus were less than 2 were removed. The read counts from libraries from RNA from samples collected after cordycepin treatment were also scaled so that the mean RNA abundance of 30 stable RNAs (ATCG00490, ATCG00680, ATMG00280, ATCG00580, ATCG00140, AT4G38970, AT2G07671, ATCG00570, ATMG00730, AT2G07727, AT2G07687, ATMG00160, AT3G11630, ATMG00060, ATCG00600, ATMG00220,   ATMG01170, ATMG00410, AT1G78900, AT3G55440, ATMG01320, AT2G21170, AT5G08670, AT5G53300, ATMG00070, AT1G26630, AT5G48300, AT2G33040, AT5G08690, AT1G57720) was the same. Decay rates were modeled from normalized data using the *RNAdecay* package[39,100]. We interpreted the initial decay rate estimate as representative of cellular RNA stability (in fraction of RNAs degraded $min^{-1}$), and these were converted to $t_{1/2}$s (min $^{-1}$). For identification of shifted RNA decay rates between mock and stimulus, we compared distinct models in which treatment parameters were constrained or in which they could vary independently. These models were compared using the corrected Akaike Information Criterion (AICc) metric. The model with the lowest AICc was selected and parameter estimates were used for downstream analysis. Following modeling, RNAs with high modeled variance were filtered ($\sigma^2 > 0.0625$).

Rates of RNA Abundance Change:  The rate of RNA abundance change was modeled from RNA abundances measured by RNA-seq from samples treated in mock and stimulus conditions for 6, 7, 8, 9, or 10 h. Linear regression was applied using RPKM values of the 7, 8, 9 h; 6, 7, 8, 9 h; 6, 7, 8, 9, 10 h; or 7, 8, 9, 10 h. The model with the smallest stimulus slope p-value was used for downstream analysis. Modeled slopes of RNAs with slope p-values < 0.05 were considered significant and were interpreted as the rate of RNA abundance change in units of RPKMs h$^{-1}$. Abundances of other RNAs (p > 0.05) were considered unchanging.

Transcription Rates: The TR at 8 h treatment was calculated using the measured abundance at that time point ($m$), modeled decay rate ($a$), and rate of abundance change ($\frac{dm}{dt}$) using the following equation:

$$TR = \frac{dm}{dt} + \alpha m$$

(described in Pérez-Ortín et al.[65] and used by Xu et al.[66]).

To identify RNAs that were regulated predominantly through decay regulation, we required stimulus and mock TRs to be very close by applying a stringent filter (|log$_2$ relative TR stimulus versus mock| < 0.138), which was selected based on the distribution of log$_2$ relative TR (stimulus vs mock) (Supplemental Figure S4B). This also resulted in consideration of relative TR differences that were small in other modalities. 206 RNAs with calculated TRs that were negative (indicating a faster decrease in abundance in the time course experiment than could be accounted for by the measured $t_{1/2}$) were removed. For calculating sov TR, we did not have rate of abundance change data, so we calculated the TR with the assumption that all RNAs were at a steady state (i.e, dm/dt = 0).

For clustering time course abundance profiles, abundances were normalized to the sum of the all time points in the mock and stimulus for each gene. RNAs were then clustered using the normalized values using k-means (the kmeans function from the *stats* package) clustering in R. All other statistical tests were performed using R functions.

## Identification of cell-type cluster markers from whole leaf single-cell RNA-seq

We used the dataset of whole leaf single cell transcriptomes generated Kim et al.[57] to characterize the RNA abundance differences observed at 8 h stimulus treatment. Kim et al.[57] characterized 19 cell clusters among 5,230 single cell transcriptomes and associated them with distinct cell-type identities. We performed UMAP dimensional reduction and clustering analysis on their dataset using the *Seurat* package version 4.0.5[101] to reconstitute their clusters and identify cluster markers (accessed Aug. 1, 2021 from https://www.ncbi.nlm.nih.gov/geo/: accession number GSE161332; files: GSE161332_features.tsv, GSE161332_barcodes.tsv, leaf_cell_info.txt, GSE161332_matrix.mtx). Cluster gene membership after reanalysis was not identical to

those of Kim et al.[57] but yielded a similar UMAP plot (Figure S1G) and 18 clusters which we numbered based on membership similarity with Kim et al.[57]. In our analysis, cluster 7 and 12 from Kim et al.[57] merged, but on average clusters shared 95% membership (Supplemental Data File 3). We used the Seurat package *FindAllMarkers* function to apply the Wilcoxon Rank Sum test to RNAs present in at least 25% of cells, and with |log$_2$FC| > 0.25 in the cell cluster compared with all others. We identified 7,996 RNAs as positive (enriched) or negative (depleted) cluster markers (Supplemental Data File 4).

## REFERENCES

1.  Hsu, C.L., and Stevens, A. (1993). Yeast Cells Lacking 5'→3' Exoribonuclease 1 Contain mRNA Species That are Poly(A) Deficient and Partially Lack the 5' Cap Structure. Mol. Cell. Biol. *13*, 4826–4835. https://doi.org/10.1128/mcb.13.8.4826-4835.1993.

2.  Kastenmayer, J.P., and Green, P.J. (2000). Novel features of the XRN-family in Arabidopsis: Evidence that AtXRN4, one of several orthologs of nuclear Xrn2p/Rat1p, functions in the cytoplasm. Proc. Natl. Acad. Sci. *97*, 13985–13990.

3.  Anderson, J.S., and Parker, R.P. (1998). The 3' to 5' degradation of yeast mRNAs is a general mechanism for mRNA turnover that requires the SKI2 DEVH box protein and 3' to 5' exonucleases of the exosome complex. EMBO J. *17*, 1497–1506. https://doi.org/10.1093/emboj/17.5.1497.

4.  Xu, J., and Chua, N.-H. (2012). Dehydration stress activates Arabidopsis MPK6 to signal DCP1 phosphorylation. EMBO J. *31*, 1975–1984. https://doi.org/10.1038/emboj.2012.56.

5.  Soma, F., Mogami, J., Yoshida, T., Abekura, M., Takahashi, F., Kidokoro, S., Mizoi, J., Shinozaki, K., and Yamaguchi-Shinozaki, K. (2017). ABA-unresponsive SnRK2 protein kinases regulate mRNA decay under osmotic stress in plants. Nat. Plants *3*, nplants2016204. https://doi.org/10.1038/nplants.2016.204.

6.  Schiaffini, M., Chicois, C., Pouclet, A., Chartier, T., Ubrig, E., Gobert, A., Zuber, H., Mutterer, J., Chicher, J., Kuhn, L., et al. (2022). A NYN domain protein directly interacts with DECAPPING1 and is required for phyllotactic pattern. Plant Physiol. *188*, 1174–1188. https://doi.org/10.1093/plphys/kiab529.

7.  Nagarajan, V.K., Stuart, C.J., DiBattista, A.T., Accerbi, M., Caplan, J.L., and Green, P.J. (2023). RNA degradome analysis reveals DNE1 endoribonuclease is required for the turnover of diverse mRNA substrates in Arabidopsis. Plant Cell *35*, 1936–1955. https://doi.org/10.1093/plcell/koad085.

8.  Pouclet, A., Pflieger, D., Merret, R., Carpentier, M.-C., Schiaffini, M., Zuber, H., Gagliardi, D., and Garcia, D. (2024). Multi-transcriptomics identifies targets of the

endoribonuclease DNE1 and highlights its coordination with decapping. Plant Cell, koae175. https://doi.org/10.1093/plcell/koae175.

9.  Molle, C., Zhang, T., Ysebrant de Lendonck, L., Gueydan, C., Andrianne, M., Sherer, F., Van Simaeys, G., Blackshear, P.J., Leo, O., and Goriely, S. (2013). Tristetraprolin regulation of interleukin 23 mRNA stability prevents a spontaneous inflammatory disease. J. Exp. Med. *210*, 1675–1684. https://doi.org/10.1084/jem.20120707.

10. Glasmacher, E., Hoefig, K.P., Vogel, K.U., Rath, N., Du, L., Wolf, C., Kremmer, E., Wang, X., and Heissmeyer, V. (2010). Roquin binds inducible costimulator mRNA and effectors of mRNA decay to induce microRNA-independent post-transcriptional repression. Nat. Immunol. *11*, 725–733. https://doi.org/10.1038/ni.1902.

11. Jiao, X., Doamekpor, S.K., Bird, J.G., Nickels, B.E., Tong, L., Hart, R.P., and Kiledjian, M. (2017). 5' End Nicotinamide Adenine Dinucleotide Cap in Human Cells Promotes RNA Decay through DXO-Mediated deNADding. Cell *168*, 1015-1027.e10. https://doi.org/10.1016/j.cell.2017.02.019.

12. Anderson, S.J., Kramer, M.C., Gosai, S.J., Yu, X., Vandivier, L.E., Nelson, A.D.L., Anderson, Z.D., Beilstein, M.A., Fray, R.G., Lyons, E., et al. (2018). N6-Methyladenosine Inhibits Local Ribonucleolytic Cleavage to Stabilize mRNAs in Arabidopsis. Cell Rep. *25*, 1146-1157.e3. https://doi.org/10.1016/j.celrep.2018.10.020.

13. Wang, Y., Li, S., Zhao, Y., You, C., Le, B., Gong, Z., Mo, B., Xia, Y., and Chen, X. (2019). NAD+-capped RNAs are widespread in the Arabidopsis transcriptome and can probably be translated. Proc. Natl. Acad. Sci., 201903682. https://doi.org/10.1073/pnas.1903682116.

14. Brengues, M., Teixeira, D., and Parker, R. (2005). Movement of eukaryotic mRNAs between polysomes and cytoplasmic processing bodies. Science *310*, 486–489. https://doi.org/10.1126/science.1115791.

15. Liu, C., Mentzelopoulou, A., Hatzianestis, I.H., Tzagkarakis, E., Skaltsogiannis, V., Ma, X., Michalopoulou, V.A., Romero-Campero, F.J., Romero-Losada, A.B., Sarris, P.F., et al. (2024). A proxitome-RNA-capture approach reveals that processing bodies repress coregulated hub genes. Plant Cell *36*, 559–584. https://doi.org/10.1093/plcell/koad288.

16. Sorenson, R., and Bailey-Serres, J. (2014). Selective mRNA sequestration by OLIGOURIDYLATE-BINDING PROTEIN 1 contributes to translational control during hypoxia in Arabidopsis. Proc. Natl. Acad. Sci. *111*, 2373–2378. https://doi.org/10.1073/pnas.1314851111.

17. Shalem, O., Dahan, O., Levo, M., Martinez, M.R., Furman, I., Segal, E., and Pilpel, Y. (2008). Transient transcriptional responses to stress are generated by opposing effects

of mRNA production and degradation. Mol. Syst. Biol. *4*, 4.
https://doi.org/10.1038/msb.2008.59.

18. Rabani, M., Levin, J.Z., Fan, L., Adiconis, X., Raychowdhury, R., Garber, M., Gnirke, A., Nusbaum, C., Hacohen, N., Friedman, N., et al. (2011). Metabolic labeling of RNA uncovers principles of RNA production and degradation dynamics in mammalian cells. Nat. Biotechnol. *29*, 436–442. https://doi.org/10.1038/nbt.1861.

19. Miller, C., Schwalb, B., Maier, K., Schulz, D., Dümcke, S., Zacher, B., Mayer, A., Sydow, J., Marcinowski, L., Dölken, L., et al. (2011). Dynamic transcriptome analysis measures rates of mRNA synthesis and decay in yeast. Mol. Syst. Biol. *7*, 458. https://doi.org/10.1038/msb.2010.112.

20. Paulsen, M.T., Veloso, A., Prasad, J., Bedi, K., Ljungman, E.A., Tsan, Y.-C., Chang, C.-W., Tarrier, B., Washburn, J.G., Lyons, R., et al. (2013). Coordinated regulation of synthesis and stability of RNA during the acute TNF-induced proinflammatory response. Proc. Natl. Acad. Sci. *110*, 2240–2245. https://doi.org/10.1073/pnas.1219192110.

21. Rabani, M., Raychowdhury, R., Jovanovic, M., Rooney, M., Stumpo, D.J., Pauli, A., Hacohen, N., Schier, A.F., Blackshear, P.J., Friedman, N., et al. (2014). High-Resolution Sequencing and Modeling Identifies Distinct Dynamic RNA Regulatory Strategies. Cell *159*, 1698–1710. https://doi.org/10.1016/j.cell.2014.11.015.

22. Tiana, M., Acosta-Iborra, B., Hernández, R., Galiana, C., Fernández-Moreno, M.Á., Jimenez, B., and Peso, L. del (2020). Metabolic labeling of RNA uncovers the contribution of transcription and decay rates on hypoxia-induced changes in RNA levels. RNA *26*, 1006–1022. https://doi.org/10.1261/rna.072611.119.

23. Tan, K., and Wilkinson, M.F. (2022). Regulation of both transcription and RNA turnover contribute to germline specification. Nucleic Acids Res. *50*, 7310–7325. https://doi.org/10.1093/nar/gkac542.

24. Lidder, P., Gutiérrez, R.A., Salomé, P.A., McClung, C.R., and Green, P.J. (2005). Circadian control of messenger RNA stability. Association with a sequence-specific messenger RNA decay pathway. Plant Physiol. *138*, 2374–2385. https://doi.org/10.1104/pp.105.060368.

25. Ravet, K., Reyt, G., Arnaud, N., Krouk, G., Djouani, E., Boucherez, J., Briat, J., and Gaymard, F. (2011). Iron and ROS control of the DownSTream mRNA decay pathway is essential for plant fitness. EMBO J. *31*, 175–186. https://doi.org/10.1038/emboj.2011.341.

26. Tanaka, M., Takano, J., Chiba, Y., Lombardo, F., Ogasawara, Y., Onouchi, H., Naito, S., and Fujiwara, T. (2011). Boron-Dependent Degradation of NIP5;1 mRNA for Acclimation

to Excess Boron Conditions in Arabidopsis. Plant Cell *23*, 3547–3559. https://doi.org/10.1105/tpc.111.088351.

27. Prasetyaningrum, P., Litthauer, S., Vegliani, F., Battle, M.W., Wood, M.W., Liu, X., Dickson, C., and Jones, M.A. (2023). Inhibition of RNA degradation integrates the metabolic signals induced by osmotic stress into the Arabidopsis circadian system. J. Exp. Bot. *74*, 5805–5819. https://doi.org/10.1093/jxb/erad274.

28. Chiba, Y., Mineta, K., Hirai, M.Y., Suzuki, Y., Kanaya, S., Takahashi, H., Onouchi, H., Yamaguchi, J., and Naito, S. (2013). Changes in mRNA Stability Associated with Cold Stress in Arabidopsis Cells. Plant Cell Physiol. *54*, 180–194. https://doi.org/10.1093/pcp/pcs164.

29. Merret, R., Descombin, J., Juan, Y., Favory, J.-J., Carpentier, M.-C., Chaparro, C., Charng, Y., Deragon, J.-M., and Bousquet-Antonelli, C. (2013). XRN4 and LARP1 Are Required for a Heat-Triggered mRNA Decay Pathway Involved in Plant Acclimation and Survival during Thermal Stress. Cell Rep. *5*, 1279–1293. https://doi.org/10.1016/j.celrep.2013.11.019.

30. Sidaway-Lee, K., Costa, M.J., Rand, D.A., Finkenstadt, B., and Penfield, S. (2014). Direct measurement of transcription rates reveals multiple mechanisms for configuration of the Arabidopsis ambient temperature response. Genome Biol. *15*, R45. https://doi.org/10.1186/gb-2014-15-3-r45.

31. Arae, T., Isai, S., Sakai, A., Mineta, K., Yokota Hirai, M., Suzuki, Y., Kanaya, S., Yamaguchi, J., Naito, S., and Chiba, Y. (2017). Co-ordinated Regulations of mRNA Synthesis and Decay during Cold Acclimation in Arabidopsis Cells. Plant Cell Physiol. *58*, 1090–1102. https://doi.org/10.1093/pcp/pcx059.

32. Goler-Baron, V., Selitrennik, M., Barkai, O., Haimovich, G., Lotan, R., and Choder, M. (2008). Transcription in the nucleus and mRNA decay in the cytoplasm are coupled processes. Genes Dev. *22*, 2022–2027. https://doi.org/10.1101/gad.473608.

33. Haimovich, G., Medina, D.A., Causse, S.Z., Garber, M., Millán-Zambrano, G., Barkai, O., Chávez, S., Pérez-Ortín, J.E., Darzacq, X., and Choder, M. (2013). Gene expression is circular: factors for mRNA degradation also foster mRNA synthesis. Cell *153*, 1000–1011. https://doi.org/10.1016/j.cell.2013.05.012.

34. Sun, M., Schwalb, B., Pirkl, N., Maier, K.C., Schenk, A., Failmezger, H., Tresch, A., and Cramer, P. (2013). Global Analysis of Eukaryotic mRNA Degradation Reveals Xrn1-Dependent Buffering of Transcript Levels. Mol. Cell *52*, 52–62. https://doi.org/10.1016/j.molcel.2013.09.010.

35. Singh, P., James, R.S., Mee, C.J., and Morozov, I.Y. (2019). mRNA levels are buffered upon knockdown of RNA decay and translation factors via adjustment of transcription

rates in human HepG2 cells. RNA Biol. *16*, 1147–1155.
https://doi.org/10.1080/15476286.2019.1621121.

36. Abernathy, E., Gilbertson, S., Alla, R., and Glaunsinger, B. (2015). Viral Nucleases Induce an mRNA Degradation-Transcription Feedback Loop in Mammalian Cells. Cell Host Microbe *18*, 243–253. https://doi.org/10.1016/j.chom.2015.06.019.

37. Gilbertson, S., Federspiel, J.D., Hartenian, E., Cristea, I.M., and Glaunsinger, B. (2018). Changes in mRNA abundance drive shuttling of RNA binding proteins, linking cytoplasmic RNA degradation to transcription. eLife *7*, e37663. https://doi.org/10.7554/eLife.37663.

38. Blasco-Moreno, B., de Campos-Mata, L., Böttcher, R., García-Martínez, J., Jungfleisch, J., Nedialkova, D.D., Chattopadhyay, S., Gas, M.-E., Oliva, B., Pérez-Ortín, J.E., et al. (2019). The exonuclease Xrn1 activates transcription and translation of mRNAs encoding membrane proteins. Nat. Commun. *10*, 1298. https://doi.org/10.1038/s41467-019-09199-6.

39. Sorenson, R.S., Deshotel, M.J., Johnson, K., Adler, F.R., and Sieburth, L.E. (2018). Arabidopsis mRNA decay landscape arises from specialized RNA decay substrates, decapping-mediated feedback, and redundancy. Proc. Natl. Acad. Sci., 201712312. https://doi.org/10.1073/pnas.1712312115.

40. Berry, S., and Pelkmans, L. (2022). Mechanisms of cellular mRNA transcript homeostasis. Trends Cell Biol. *32*, 655–668. https://doi.org/10.1016/j.tcb.2022.05.003.

41. Duncan-Lewis, C., Hartenian, E., King, V., and Glaunsinger, B.A. (2021). Cytoplasmic mRNA decay represses RNA polymerase II transcription during early apoptosis. eLife *10*, e58342. https://doi.org/10.7554/eLife.58342.

42. Kondo, Y., Ito, T., Nakagami, H., Hirakawa, Y., Saito, M., Tamaki, T., Shirasu, K., and Fukuda, H. (2014). Plant GSK3 proteins regulate xylem cell differentiation downstream of TDIF–TDR signalling. Nat. Commun. *5*, 3504. https://doi.org/10.1038/ncomms4504.

43. Kondo, Y., Fujita, T., Sugiyama, M., and Fukuda, H. (2015). A Novel System for Xylem Cell Differentiation in Arabidopsis thaliana. Mol. Plant *8*, 612–621. https://doi.org/10.1016/j.molp.2014.10.008.

44. Kondo, Y., Nurani, A.M., Saito, C., Ichihashi, Y., Saito, M., Yamazaki, K., Mitsuda, N., Ohme-Takagi, M., and Fukuda, H. (2016). Vascular Cell Induction Culture System Using Arabidopsis Leaves (VISUAL) Reveals the Sequential Differentiation of Sieve Element-Like Cells. Plant Cell *28*, 1250–1262. https://doi.org/10.1105/tpc.16.00027.

45. Furuya, T., Saito, M., Uchimura, H., Satake, A., Nosaki, S., Miyakawa, T., Shimadzu, S., Yamori, W., Tanokura, M., Fukuda, H., et al. (2021). Gene co-expression network

analysis identifies BEH3 as a stabilizer of secondary vascular development in Arabidopsis. Plant Cell *33*, 2618–2636. https://doi.org/10.1093/plcell/koab151.

46. Baima, S., Possenti, M., Matteucci, A., Wisman, E., Altamura, M.M., Ruberti, I., and Morelli, G. (2001). The arabidopsis ATHB-8 HD-zip protein acts as a differentiation-promoting transcription factor of the vascular meristems. Plant Physiol. *126,* 643–655. https://doi.org/10.1104/pp.126.2.643.

47. Scarpella, E., Francis, P., and Berleth, T. (2004). Stage-specific markers define early steps of procambium development in Arabidopsis leaves and correlate termination of vein formation with mesophyll differentiation. Development *131*, 3445–3455. https://doi.org/10.1242/dev.01182.

48. Marquès-Bueno, M.D.M., Morao, A.K., Cayrel, A., Platre, M.P., Barberon, M., Caillieux, E., Colot, V., Jaillais, Y., Roudier, F., and Vert, G. (2016). A versatile Multisite Gateway-compatible promoter and transgenic line collection for cell type-specific functional genomics in Arabidopsis. Plant J. Cell Mol. Biol. *85*, 320–333. https://doi.org/10.1111/tpj.13099.

49. Sawchuk, M.G., Head, P., Donner, T.J., and Scarpella, E. (2007). Time-lapse imaging of Arabidopsis leaf development shows dynamic patterns of procambium formation. New Phytol. *176*, 560–571. https://doi.org/10.1111/j.1469-8137.2007.02193.x.

50. Donner, T.J., Sherr, I., and Scarpella, E. (2009). Regulation of preprocambial cell state acquisition by auxin signaling in Arabidopsis leaves. Development *136*, 3235–3246. https://doi.org/10.1242/dev.037028.

51. Konishi, M., and Yanagisawa, S. (2007). Sequential activation of two Dof transcription factor gene promoters during vascular development in Arabidopsis thaliana. Plant Physiol. Biochem. *45*, 623–629. https://doi.org/10.1016/j.plaphy.2007.05.001.

52. Konishi, M., and Yanagisawa, S. (2015). Transcriptional repression caused by Dof5.8 is involved in proper vein network formation in Arabidopsis thaliana leaves. J. Plant Res. *128*, 643–652. https://doi.org/10.1007/s10265-015-0712-0.

53. Konishi, M., Donner, T.J., Scarpella, E., and Yanagisawa, S. (2015). MONOPTEROS directly activates the auxin-inducible promoter of the Dof5.8 transcription factor gene in Arabidopsis thaliana leaf provascular cells. J. Exp. Bot. *66*, 283–291. https://doi.org/10.1093/jxb/eru418.

54. Wallner, E.-S., López-Salmerón, V., Belevich, I., Poschet, G., Jung, I., Grünwald, K., Sevilem, I., Jokitalo, E., Hell, R., Helariutta, Y., et al. (2017). Strigolactone- and Karrikin-Independent SMXL Proteins Are Central Regulators of Phloem Formation. Curr. Biol. *27*, 1241–1247. https://doi.org/10.1016/j.cub.2017.03.014.

55. Wallner, E.-S., Tonn, N., Shi, D., Jouannet, V., and Greb, T. (2020). SUPPRESSOR OF MAX2 1-LIKE 5 promotes secondary phloem formation during radial stem growth. Plant J. *102*, 903–915. https://doi.org/10.1111/tpj.14670.

56. Sawchuk, M.G., Donner, T.J., Head, P., and Scarpella, E. (2008). Unique and Overlapping Expression Patterns among Members of Photosynthesis-Associated Nuclear Gene Families in Arabidopsis. Plant Physiol. *148*, 1908–1924. https://doi.org/10.1104/pp.108.126946.

57. Kim, J.-Y., Symeonidi, E., Pang, T.Y., Denyer, T., Weidauer, D., Bezrutczyk, M., Miras, M., Zöllner, N., Hartwig, T., Wudick, M.M., et al. (2021). Distinct identities of leaf phloem cells revealed by single cell transcriptomics. Plant Cell. https://doi.org/10.1093/plcell/koaa060.

58. Narsai, R., Howell, K.A., Millar, A.H., O'Toole, N., Small, I., and Whelan, J. (2007). Genome-wide analysis of mRNA decay rates and their determinants in Arabidopsis thaliana. Plant Cell *19*, 3418–3436. https://doi.org/10.1105/tpc.107.055046.

59. Chantarachot, T., Sorenson, R.S., Hummel, M., Ke, H., Kettenburg, A.T., Chen, D., Aiyetiwa, K., Dehesh, K., Eulgem, T., Sieburth, L.E., et al. (2020). DHH1/DDX6-like RNA helicases maintain ephemeral half-lives of stress-response mRNAs. Nat. Plants *6*, 675–685. https://doi.org/10.1038/s41477-020-0681-8.

60. Szabo, E.X., Reichert, P., Lehniger, M.-K., Ohmer, M., Amorim, M. de F., Gowik, U., Schmitz-Linneweber, C., and Laubinger, S. (2020). Metabolic Labeling of RNAs Uncovers Hidden Features and Dynamics of the Arabidopsis Transcriptome. Plant Cell. https://doi.org/10.1105/tpc.19.00214.

61. Carpentier, M.-C., Receveur, A.-E., Boubegtitene, A., Cadoudal, A., Bousquet-Antonelli, C., and Merret, R. (2024). Genome-wide analysis of mRNA decay in Arabidopsis shoot and root reveals the importance of co-translational mRNA decay in the general mRNA turnover. Nucleic Acids Res., gkae363. https://doi.org/10.1093/nar/gkae363.

62. Jang, G.-J., Yang, J.-Y., Hsieh, H.-L., and Wu, S.-H. (2019). Processing bodies control the selective translation for optimal development of Arabidopsis young seedlings. Proc. Natl. Acad. Sci. *116*, 6451–6456. https://doi.org/10.1073/pnas.1900084116.

63. Legen, J., Lenzen, B., Kachariya, N., Feltgen, S., Gao, Y., Mergenthal, S., Weber, W., Klotzsch, E., Zoschke, R., Sattler, M., et al. (2024). A prion-like domain is required for phase separation and chloroplast RNA processing during cold acclimation in Arabidopsis. Plant Cell, koae145. https://doi.org/10.1093/plcell/koae145.

64. Li, Q., Liu, Y., and Zhang, X. (2024). Biomolecular condensates in plant RNA silencing: insights into formation, function, and stress responses. Plant Cell *36*, 227–245. https://doi.org/10.1093/plcell/koad254.

65. Pérez-Ortín, J.E., Alepuz, P.M., and Moreno, J. (2007). Genomics and gene transcription kinetics in yeast. Trends Genet. *23*, 250–257. https://doi.org/10.1016/j.tig.2007.03.006.

66. Xu, Z., Sziraki, A., Lee, J., Zhou, W., and Cao, J. (2023). Dissecting key regulators of transcriptome kinetics through scalable single-cell RNA profiling of pooled CRISPR screens. Nat. Biotechnol., 1–6. https://doi.org/10.1038/s41587-023-01948-9.

67. Forouzanfar, F., Moreno, D.F., Plassard, D., Furst, A., Oliveira, K.A., Reina-San-Martin, B., Tora, L., Molina, N., and Mendoza, M. (2025). Gene-specific transcript buffering revealed by perturbation of coactivator complexes. Sci. Adv. *11*, eadr1492. https://doi.org/10.1126/sciadv.adr1492.

68. Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., et al. (2000). Gene Ontology: tool for the unification of biology. Nat. Genet. *25*, 25–29. https://doi.org/10.1038/75556.

69. Lange, H., Ndecky, S.Y.A., Gomez-Diaz, C., Pflieger, D., Butel, N., Zumsteg, J., Kuhn, L., Piermaria, C., Chicher, J., Christie, M., et al. (2019). RST1 and RIPR connect the cytosolic RNA exosome to the Ski complex in Arabidopsis. Nat. Commun. *10*, 1–12. https://doi.org/10.1038/s41467-019-11807-4.

70. Shalem, O., Groisman, B., Choder, M., Dahan, O., and Pilpel, Y. (2011). Transcriptome Kinetics Is Governed by a Genome-Wide Coupling of mRNA Production and Degradation: A Role for RNA Pol II. PLOS Genet. *7*, e1002273. https://doi.org/10.1371/journal.pgen.1002273.

71. Faucillion, M.-L., Johansson, A.-M., and Larsson, J. (2022). Modulation of RNA stability regulates gene expression in two opposite ways: through buffering of RNA levels upon global perturbations and by supporting adapted differential expression. Nucleic Acids Res. *50*, 4372–4388. https://doi.org/10.1093/nar/gkac208.

72. Slobodin, B., Bahat, A., Sehrawat, U., Becker-Herman, S., Zuckerman, B., Weiss, A.N., Han, R., Elkon, R., Agami, R., Ulitsky, I., et al. (2020). Transcription Dynamics Regulate Poly(A) Tails and Expression of the RNA Degradation Machinery to Balance mRNA Levels. Mol. Cell *78*, 434-444.e5. https://doi.org/10.1016/j.molcel.2020.03.022.

73. Rabani, M., Raychowdhury, R., Jovanovic, M., Rooney, M., Stumpo, D.J., Pauli, A., Hacohen, N., Schier, A.F., Blackshear, P.J., Friedman, N., et al. (2014). High-Resolution Sequencing and Modeling Identifies Distinct Dynamic RNA Regulatory Strategies. Cell *159*, 1698–1710. https://doi.org/10.1016/j.cell.2014.11.015.

74. Scarpella, E., Marcos, D., Friml, J., and Berleth, T. (2006). Control of leaf vascular patterning by polar auxin transport. Genes Dev. *20*, 1015–1027. https://doi.org/10.1101/gad.1402406.

75. Gardiner, J., Sherr, I., and Scarpella, E. (2010). Expression of DOF genes identifies early stages of vascular development in Arabidopsis leaves. Int. J. Dev. Biol. *54*, 1389–1396. https://doi.org/10.1387/ijdb.093006jg.

76. Garcia-Martinez, J., Medina, D.A., Bellvis, P., Sun, M., Cramer, P., Chavez, S., and Perez-Ortin, J.E. (2021). The total mRNA concentration buffering system in yeast is global rather than gene-specific. RNA, rna.078774.121. https://doi.org/10.1261/rna.078774.121.

77. Chattopadhyay, S., Garcia-Martinez, J., Haimovich, G., Fischer, J., Khwaja, A., Barkai, O., Chuartzman, S.G., Schuldiner, M., Elran, R., Rosenberg, M.I., et al. (2022). RNA-controlled nucleocytoplasmic shuttling of mRNA decay factors regulates mRNA synthesis and a novel mRNA decay pathway. Nat. Commun. *13*, 7184. https://doi.org/10.1038/s41467-022-34417-z.

78. Rymarquis, L.A., Souret, F.F., and Green, P.J. (2011). Evidence that XRN4, an Arabidopsis homolog of exoribonuclease XRN1, preferentially impacts transcripts with certain sequences or in particular functional categories. RNA *17*, 501–511. https://doi.org/10.1261/rna.2467911.

79. Schulz, D., Pirkl, N., Lehmann, E., and Cramer, P. (2014). Rpb4 Subunit Functions Mainly in mRNA Synthesis by RNA Polymerase II♦. J. Biol. Chem. *289*, 17446–17452. https://doi.org/10.1074/jbc.M114.568014.

80. Richard, S., Gross, L., Fischer, J., Bendalak, K., Ziv, T., Urim, S., and Choder, M. (2021). Numerous Post-translational Modifications of RNA Polymerase II Subunit Rpb4/7 Link Transcription to Post-transcriptional Mechanisms. Cell Rep. *34*, 108578. https://doi.org/10.1016/j.celrep.2020.108578.

81. Kelbert, M., Jordán-Pla, A., de Miguel-Jiménez, L., García-Martínez, J., Selitrennik, M., Guterman, A., Henig, N., Granneman, S., Pérez-Ortín, J.E., Chávez, S., et al. (2024). The zinc-finger transcription factor Sfp1 imprints specific classes of mRNAs and links their synthesis to cytoplasmic decay. eLife *12*, RP90766. https://doi.org/10.7554/eLife.90766.

82. Wang, Q., and Lin, J. (2024). Homeostasis of mRNA concentrations through coupling transcription, export, and degradation. iScience *27*. https://doi.org/10.1016/j.isci.2024.110531.

83. Berry, S., Müller, M., Rai, A., and Pelkmans, L. (2022). Feedback from nuclear RNA on transcription promotes robust RNA concentration homeostasis in human cells. Cell Syst. *13*, 454-470.e15. https://doi.org/10.1016/j.cels.2022.04.005.

84. Sun, X.-M., Bowman, A., Priestman, M., Bertaux, F., Martinez-Segura, A., Tang, W., Whilding, C., Dormann, D., Shahrezaei, V., and Marguerat, S. (2020). Size-Dependent Increase in RNA Polymerase II Initiation Rates Mediates Gene Expression Scaling with Cell Size. Curr. Biol. *30*, 1217-1230.e7. https://doi.org/10.1016/j.cub.2020.01.053.

85. Padovan-Merhar, O., Nair, G.P., Biaesch, A.G., Mayer, A., Scarfone, S., Foley, S.W., Wu, A.R., Churchman, L.S., Singh, A., and Raj, A. (2015). Single Mammalian Cells Compensate for Differences in Cellular Volume and DNA Copy Number through Independent Global Transcriptional Mechanisms. Mol. Cell *58*, 339–352. https://doi.org/10.1016/j.molcel.2015.03.005.

86. Rodríguez-Molina, J.B., Tseng, S.C., Simonett, S.P., Taunton, J., and Ansari, A.Z. (2016). Engineered Covalent Inactivation of TFIIH-Kinase Reveals an Elongation Checkpoint and Results in Widespread mRNA Stabilization. Mol. Cell *63*, 433–444. https://doi.org/10.1016/j.molcel.2016.06.036.

87. Baptista, T., Grünberg, S., Minoungou, N., Koster, M.J.E., Timmers, H.T.M., Hahn, S., Devys, D., and Tora, L. (2017). SAGA Is a General Cofactor for RNA Polymerase II Transcription. Mol. Cell *68*, 130-143.e5. https://doi.org/10.1016/j.molcel.2017.08.016.

88. Chappleboim, A., Joseph-Strauss, D., Gershon, O., and Friedman, N. (2022). Transcription feedback dynamics in the wake of cytoplasmic mRNA degradation shutdown. Nucleic Acids Res. *50*, 5864–5880. https://doi.org/10.1093/nar/gkac411.

89. Bryll, A.R., and Peterson, C.L. (2022). Functional interaction between the RNA exosome and the sirtuin deacetylase Hst3 maintains transcriptional homeostasis. Genes Dev. *36*, 17–22. https://doi.org/10.1101/gad.348923.121.

90. Mitra, P.P., and Loqué, D. (2014). Histochemical Staining of Arabidopsis thaliana Secondary Cell Wall Elements. JoVE J. Vis. Exp., e51381. https://doi.org/10.3791/51381.

91. Kurihara, D., Mizuta, Y., Sato, Y., and Higashiyama, T. (2015). ClearSee: a rapid optical clearing reagent for whole-plant fluorescence imaging. Development *142*, 4168–4179. https://doi.org/10.1242/dev.127613.

92. Backman, T.W.H., and Girke, T. (2016). systemPipeR: NGS workflow and report generation environment. BMC Bioinformatics *17*, 388. https://doi.org/10.1186/s12859-016-1241-0.

93. Pages, H., Aboyoun, P., Gentleman, R., and DebRoy, S. Biostrings: String objects representing biological sequences, and matching algorithms.

94. Trapnell, C., Pachter, L., and Salzberg, S.L. (2009). TopHat: discovering splice junctions with RNA-Seq. Bioinforma. Oxf. Engl. *25*, 1105–1111. https://doi.org/10.1093/bioinformatics/btp120.

95. Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. Genome Biol. *10*, R25. https://doi.org/10.1186/gb-2009-10-3-r25.

96. Lawrence, M., Huber, W., Pagès, H., Aboyoun, P., Carlson, M., Gentleman, R., Morgan, M.T., and Carey, V.J. (2013). Software for Computing and Annotating Genomic Ranges. PLOS Comput. Biol. *9*, e1003118. https://doi.org/10.1371/journal.pcbi.1003118.

97. Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics *26*, 139–140. https://doi.org/10.1093/bioinformatics/btp616.

98. Alexa, A., and Rahnenfuhrer, J. (2020). topGO: Enrichment Analysis for Gene Ontology. Version 2.40.0.

99. Wickham, H. (2009). ggplot2: elegant graphics for data analysis (Springer New York).

100. Sorenson, R.S., Johnson, K., Adler, F.R., and Sieburth, L.E. (2018). RNAdecay: Maximum Likelihood Decay Modeling of RNA Degradation Data. Version 1.0.1 (DOI: 10.18129/B9.bioc.RNAdecay).

101. Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J., Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. Cell *184*, 3573-3587.e29. https://doi.org/10.1016/j.cell.2021.04.048.

102. Oda, Y., and Fukuda, H. (2012). Secondary cell wall patterning during xylem differentiation. Curr. Opin. Plant Biol. *15*, 38–44. https://doi.org/10.1016/j.pbi.2011.10.005.

**FIGURE LEGENDS**

**Figure 1. Vascular developmental stimulus induces active cellular reprogramming**

(A) Effect of stimulus on an early procambial marker. YFP fluorescence was measured as mean pixel intensity x $10^{-3}$ from 2 mm diameter leaf disks of proAtHB8::4xYFP plants over 24 h treatment.

(B) Effect of stimulus on relative RNA abundance of vascular differentiation markers. RNA abundance was measured by qRT-PCR over 20 h of treatment and normalized to the *UBC10* reference RNA.

(C) Pie chart showing the proportions of RNAs with increased or decreased RNA abundance (FDR < 0.05) by RNA-seq at 8 h stimulus treatment compared with mock.

(D) Effect of stimulus on RNA abundances of members of the *phloem or xylem histogenesis*, *auxin activated signaling*, *thylakoid*, and *response to light stimulus* gene ontologies. Color squares represent $\log_2$ relative RNA abundance (LFC) of GO members. BP, biological process; CC, cellular component.

(E, F) Stimulus-induced changes in leaf vascular cell-identity (E) and mesophyll (F) markers. Box plot of $\log_2$ relative RNA abundance change (8 h stimulus vs mock) of markers identified by leaf single cell RNA-seq clusters generated by Kim et al.[57].

(G) Experimental design schematic for RNA abundance and RNA decay analysis.

(H) Pie chart showing proportions of RNAs with increased, decreased, or unchanged $t_{1/2}$s.

(I) Distribution of $t_{1/2}$ changes.

(J,K) Decay plots showing RNA abundance changes after inhibition of transcription (thin lines, mean±SE) and $t_{1/2}$ estimates based on modeled decay (thick lines) of 8 h stimulus- (blue and red) and mock-treated (black) leaves.

(L) Distribution of $t_{1/2}$ (8 h mock, black dots) and shift in $t_{1/2}$ (line segments) for 2,818 RNAs with shifted $t_{1/2}$s in the standard range. Line segment endpoints are positioned at the mock and stimulus $t_{1/2}$s. Line color indicates $\log_2$ relative abundance (LFC). See Figure S5A for enlargement of the 100 shortest and longest mock $t_{1/2}$s. Numbered line segments indicate RNAs that have much shorter or longer $t_{1/2}$s in stimulus: (1) Protein kinase family protein with leucine-rich repeat (AT1G35710), (2) TIR-NBS-LRR Disease resistance protein (AT4G36150), (3) CCCH-type zinc finger (AT5G58620), (4) hypothetical protein (AT2G07779), (5) hypothetical protein (AT5G66580), (6) snoRNA (AT1G32385), (7) FTSH11 (AT5G53170), (8) BTB/POZ domain-containing protein (AT5G59140), (9) CID4 (AT3G14010), (10) TRM8 (AT5G26910), (11) transcriptional regulator of RNA PolII, SAGA, subunit (AT4G33890), (12) Disease resistance protein (TIR-NBS-LRR class) (AT4G16900).

**Figure 2. Broad regulation of RNA decay and RoAC  occurs during stimulus response.**

(A) Effect of stimulus response on members of the early vascular regulator gene network. Heatmap of relative RNA abundance (8h stimulus vs. mock). Color and numbers indicate $\log_2$ fold change and fold change, respectively. Bold numbers indicate significant changes (FDR < 0.05). *, *ATHB8* is represented in two locations on the map. ‡, TMO6 is also represented twice.

(B) $t_{1/2}$ regulation in the EVR network (as in (A)). Color indicates $t_{1/2}$. 11 RNAs shifted $t_{1/2}$s and are indicated by a black outline and label.

(C) Schematic of RNA regulation occurring for EVR network RNAs with higher (top) or lower (bottom) abundance (FDR < 0.05).

(D) Schematic for determining rates of abundance change. Mock- and stimulus-treated tissue were collected after 6, 7, 8, 9, or 10 h to model the RoAC across the 8 h time point for each RNA.

(E) Distributions of RoAC for RNAs that were decreasing (left) or increasing (right) abundance at 8 h mock (top) or stimulus (bottom).

(F) Venn diagram showing RNAs with changing abundances in mock or stimulus.

(G) Pie chart showing the effect of stimulus on RoAC; rates increased (faster), decreased (slower), or were the same.

(H) Net effect of RNA regulatory mechanisms on RoAC (8 h stimulus vs. mock) in the EVR network (as in (A)). Positive or negative influence was found on rates of the 23 labeled RNAs.

(I-L) RNAs of the EVR network reveal distinct regulatory modalities through changes in transcription and decay rates as indicated by arrows: red, faster; gray, unchanged; blue, slower.

**Figure 3. Transcriptome-wide snapshot of 8h stimulus response includes active regulatory modalities and steady abundances maintained through changes in flux.**

(A) Regulatory modalities are defined by change or lack of change in synthesis and decay and the net effect on RoAC. Counts (and proportions) of RNAs exhibiting behavior of each regulatory modality are presented as a stacked bar plot.

(B-I) Regulation patterns of faster or slower synthesis and/or decay are demonstrated for each regulatory modality and are represented pictographically as larger red (faster TR/shorter $t_{1/2}$) or smaller blue (slower TR/longer $t_{1/2}$) arrows on either side of an oval. Ovals represent RNA abundance. Green arrows around the ovals represent increasing or decreasing RoAC. For each example RNA, abundance measurements (RPKM mean±SE) at 6, 7, 8, 9, 10 h of mock (thin light gray line) and stimulus (thin dark gray line) along with linear models of RoAC are shown for mock (thick black line) and stimulus (thick green line). Model parameters label the graphs: s, slope of the linear model (RPKM h-1); p, slope p-value; *, indicates a significant difference at the given time point, FDR < 0.05. TRs and $t_{1/2}$s are also shown in bar plots to the right for mock (m), and stimulus (s).

**Figure 4. Dynamic range in transcription and $t_{1/2}$ regulation are modality dependent.**

(A-G) Contributions of transcription and decay regulation for individual RNAs. Stacked bar plots show direct comparison of regulation of TR and $t_{1/2}$ for each RNA of regulatory modalities; bars indicate direction and size of $\log_2$ relative TR (purple) and $\log_2$ relative $t_{1/2}$ (orange). Black dots indicate net regulation (sum of the LFC rel. TR and LFC rel. $t_{1/2}$). Genes within each regulatory modality are organized by increasing net regulation. The net regulation extremes that slightly deviate from 0 in the buffered RNAs reflects abundance variance below the cutoff.

**Figure 5. Synergistic and oppositional modalities drive responsive RNA abundance.**

(A) RNA abundance response profiles. k-means clustering of 4,268 RNAs of the transcription, decay, synergistic, and oppositional modalities. Abundance data of each RNA was normalized to the total (sum) abundance of all time points so clustering would group RNAs with similar abundance response profiles independent of level of expression. Clusters were numbered based on the relative abundance difference between mock (gray lines) and stimulus (green lines). Cluster number and size label each plot; label color indicates abundance trajectories at 8 h: diverging (blue), converging (pink), or other (black).

(B) Modality enrichment among responsive abundance. RNAs of diverging or converging abundance clusters in (A) were grouped and evaluated for enrichment (blue) or depletion (yellow) of the synergistic and oppositional regulatory modalities by the hypergeometric test; enrichment/depletion are displayed as a heatmap of and labeled with $-\log_{10}$ p-values.

(C) Box and whisker plot of the abundance change magnitudes (|L2FC abundance|) by regulatory modality. Abundances of all stimulus time points were compared to those of all the mock time points.

(D) Enrichment analysis of RNA groups with distinct changes in $t_{1/2}$ in *vcs* mutants: *vcs* feedback decay (rel. $t_{1/2}$ *vcs*/WT < 1), moderate *VCS*−dependence (rel. $t_{1/2}$ *vcs*/WT > 1 & < 2), high *VCS*−dependence (rel. $t_{1/2}$ *vcs*/WT > 2). Significant enrichments are labeled with $-\log_{10}$ p-values.

**Supplemental Figure S1. Stimulus induces developmental, molecular and biochemical changes to cells.**

(A) YFP fluorescence of rosette leaves 1 and 2 of *proATHB8::4xYFP* plants incubated in mock and stimulus treatments for 24 and 72 h (bar = 1 mm).

(B) Development of xylem element characteristics in cells outside of vascular bundles after 72 h of stimulus treatment but not 72 h mock treatment. DIC micrograph of ectopic formation of spiral patterned secondary cell walls[102] can be seen (bar = 50 µm).

(C) Lignification of cell walls 72 h after stimulus treatment. Phloroglucinol was used to stain lignin (pink color). Box (dashed line) in the left column are enlarged to the right.

(D) Volcano plot overlayed on a histogram of 8 h RNA-seq comparison of stimulus vs mock.

(E) Heatmap of Pearson correlation values comparing similarity of abundance change of 460 RNA detected in both RNA seq data of this study and those measured by microarray by Furuya et al.[45].

(F) Venn diagram comparing RNA identities of 8 h stimulus-upregulated RNAs from this study and stimulus-upregulated xylem- and phloem-specific RNAs after 72 h[44].

(G) Leaf single cell RNA-seq UMAP clusters based on RNA expression measured by Kim et al.[57]. PC, procambium; PP, Phloem parenchyma; XP, xylem parenchyma. Compare Figure 1E, 1F.

**Supplemental Figure S2. RNA t1/2 extremes have unique features.**

(A) Dynamic ranges of RNAs that shift to or from extremely long $t_{1/2}$s (>1440 min) were separated from other RNAs. Line segment ends indicate $t_{1/2}$s; black dots indicate the mock ends.

(B) At $t_{1/2}$ extremes, short short-lived RNAs tended to become longer-lived, and long-lived RNAs tended to become shorter-lived. The 100 shortest and 100 longest $t_{1/2}$s in mock are shown. Line segments represent shifts in $t_{1/2}$ and are positioned at the mock $t_{1/2}$s (black dots). Line color indicates $\log_2$ relative abundance (LFC); compare with (B) and Figure 4A.

(C) Dynamics of long-lived RNAs ($t_{1/2} \geq 1440$ min). Some RNAs are long-lived in mock but move out of this group in stimulus (blue circle), others become long-lived in stimulus (green circle), and others remain at $t_{1/2} \geq 1440$ min in stimulus and mock.

(D) Bar plot comparing proportion of mock $t_{1/2}$ quintiles with shifts in decay.

**Supplemental Figure S3. Analysis flowchart for comparing abundances, $t_{1/2}$s, and TRs of mock and stimulus treatments.**

(A) Initial EVR network analysis by comparison of 8 h abundances with $t_{1/2}$s.

(B) EVR network analysis using RoAC, $t_{1/2}$ regulation, and calculated TRs.

(C) Whole transcriptome regulation analysis as in (B). See (B) for modality assignment criteria.

**Supplemental Figure S4. Early stimulus transcriptional response is varied and dynamic and includes buffered RNAs.**

(A) Distributions of TRs for mock and stimulus RNAs with changing abundances presented as histograms.

(B) Distribution of $\log_2$ relative TRs (stimulus/mock) as a histogram. Solid light blue lines indicate the cutoffs for unchanged TR (1 +/- 0.18).

(C) Examples of RNA buffering from Steady group 4. Buffered RNAs have steady abundances but their TRs and $t_{1/2}$s differ. For each example RNA, abundance measurements (RPKM mean±SE) at 6, 7, 8, 9, 10 h of mock (thin light gray line) and stimulus (thin dark gray line) along with linear models of rate of abundance change are shown for mock (thick black line) and stimulus (thick green line). Model p-values label the graphs. TRs and $t_{1/2}$s are also shown in bar plots to the right for mock (m), and stimulus (s).

**Supplemental Figure S5. Dynamic range extended analysis.**

(A,C-E) Dynamic range of decay rate shifts (orange line segments, left column) in regulatory modalities. Line segment endpoints indicate mock and stimulus $t_{1/2}$s. RNAs are ordered on the y-axis by increasing $t_{1/2}$ mean of stimulus and mock treatments (indicated by black circles). Inset histograms show distribution of $\log_2$ relative $t_{1/2}$.

(B,C-E) Dynamic range of TR changes (purple line segments, right column) in regulatory modalities. Line segment endpoints indicate mock and stimulus TRs. RNAs are ordered on the y-axis by increasing mean TR of stimulus and mock treatments (indicated by black circles). Inset histograms show distribution of $\log_2$ relative TR.

(F) Oppositional and Steady 2: Buffered stacked bar plots with dashed-line boxes indicating expanded regions in (G) and (H), respectively. Asterisks highlight exceptionally large and nearly symmetric changes in TR and $t_{1/2}$.

(I,J) Comparison of abundances and $t_{1/2}$ shifts of RNA buffered in *sov* mutants and buffered (Steady 2).

(K) Gene ontology enrichment analysis of regulatory modality RNAs. The decay, synergistic, and steady 3 modalities did not have enriched GOs (p-adj, adjusted p-value).

Table 1. Magnitudes of transcription rate and half-life change by regulatory modality.

| Modality | $t_{1/2}$ shift | | transcription rate change | |
|---|---|---|---|---|
| | fold change mean* | fold change median** | fold change mean* | fold change median** |
| transcription | 1.0 | 1.0 | 1.8 | 1.5 |
| decay | 1.9 | 1.7 | 1.0 | 1.1 |
| synergistic | 2.3 | 1.9 | 1.6 | 1.4 |
| oppositional | 2.7 | 2.0 | 3.1 | 2.3 |
| steady 1 | 2.8 | 2.0 | 3.2 | 2.6 |
| buffered steady 2 | 2.7 | 2.0 | 2.8 | 2.0 |
| steady 3 | 1.0 | 1.0 | 1.1 | 1.1 |
| steady 4 | 1.0 | 1.0 | 1.5 | 1.4 |

* 2^mean(|log2(stimulus/mock)|); ** 2^median(|log2(stimulus/mock)|)

**Supplemental Table 1. Early vascular network members and source literature.**

| name | geneid | source |
|---|---|---|
| ACL5 | AT5G19530 | Jouannet et al. 2015 [1] |
| AHP6/ARF17 | AT1G80100 | transcriptional regulatory network of Smit et al. 2019 ; Ohashi-ito et al. 2014 [2,3] |
| ATHB8 | AT4G32880 | Jouannet et al 2015; PC-associated genes of Kondo et al. 2015 [1,4] |
| ATSK11 | AT5G26751 | transcriptional regulatory network of Smit et al. 2019 [2] |
| ATSK13 | AT5G14640 | transcriptional regulatory network of Smit et al. 2019 [2] |
| BIL1/SK2-2/SK23/SK-D-zeta | AT2G30980 | transcriptional regulatory network of Smit et al. 2019 [2] |
| BIL2 | AT1G06390 | transcriptional regulatory network of Smit et al. 2019 [2] |
| BIN1/BRI1/CBB2/DWF2 | AT4G39400 | transcriptional regulatory network of Smit et al. 2019 [2] |
| BRL1 | AT1G55610 | transcriptional regulatory network of Smit et al. 2019 [2] |
| BRL2/VH1 | AT2G01950 | PC-associated genes of Kondo et al. 2015 [4] |
| BZR1 | AT1G75080 | transcriptional regulatory network of Smit et al. 2019 [2] |
| CLE41 | AT3G24770 | transcriptional regulatory network of Smit et al. 2019 [2] |
| CLE44 | AT4G13195 | transcriptional regulatory network of Smit et al. 2019 [2] |
| CVL1 | AT2G32010 | transcriptional regulatory network of Smit et al. 2019 [2] |
| CVP2 | AT1G05470 | transcriptional regulatory network of Smit et al. 2019 [2] |
| DOF2.1 | AT2G28510 | PC-associated genes of Kondo et al. 2015; transcriptional regulatory network of Smit et al. 2019 [2,4] |
| DOF5.8 | AT5G66940 | PC-associated genes of Kondo et al. 2015; Konishi et al. 2015; Larrieu et al. 2022 [4–6] |
| DOF6 | AT3G45610 | Miyashima et al. 2019 [7] |
| FKD1 | AT3G63300 | PC-associated genes of Kondo et al. 2015 [4] |
| GNOM/MIZ2/VAN7 | AT1G13980 | Geldner et al. 2003 [8] |
| HB15/CAN/ICU4 | AT1G52150 | PC-associated genes of Kondo et al. 2015 [4] |
| HCA2/DOF5.6 | AT5G62940 | transcriptional regulatory network of Smit et al. 2019 [2] |
| IAA29 | AT4G32280 | transcriptional regulatory network of Smit et al. 2019 [2] |
| IAA30 | AT3G62100 | transcriptional regulatory network of Smit et al. 2019 [2] |
| LAX2 | AT2G21050 | PC-associated genes of Kondo et al. 2015 [4] |
| LBD1 | AT1G07900 | transcriptional regulatory network of Smit et al. 2019 [2] |
| LBD3/ASL9 | AT1G16530 | transcriptional regulatory network of Smit et al. 2019 [2] |
| LBD4 | AT1G31320 | transcriptional regulatory network of Smit et al. 2019 [2] |
| LHL1 | AT1G06150 | transcriptional regulatory network of Smit et al. 2019 [2] |
| LHW | AT2G27230 | Jouannet et al. 2015 [1] |
| LOG3 | AT2G37210 | transcriptional regulatory network of Smit et al. 2019 [2] |
| LOG4 | AT3G53450 | transcriptional regulatory network of Smit et al. 2019 [2] |
| MP/ARF5 | AT1G19850 | PC-associated genes of Kondo et al. 2015 [4] |
| OBP1/DOF3.4 | AT3G50410 | Larrieu et al. 2022 [6] |
| OBP2 | AT1G07640 | Miyashima et al. 2019 [7] |
| OPL2 | AT2G38070 | Ruiz Sola et al., 2017 [9] |
| OPS | AT3G09070 | PC-associated genes of Kondo et al. 2015 [4] |
| PEAR1/DOF2.4 | AT2G37590 | PC-associated genes of Kondo et al. 2015 [4] |

| | | |
|---|---|---|
| PEAR2/DOF5.1 | AT5G02460 | Miyashima et al. 2019; Kim et al. 2010 [7,10] |
| PHB/HB14 | AT2G34710 | transcriptional regulatory network of Smit et al. 2019 [2] |
| PHV/HB9 | AT1G30490 | transcriptional regulatory network of Smit et al. 2019 [2] |
| PIN1 | AT1G73590 | PC-associated genes of Kondo et al. 2015 [4] |
| PIN6 | AT1G77110 | Nisar et al., 2014 [11] |
| PNH/AGO10/ZWILLE | AT5G43810 | Nagawa et al. 2006, Table 2 [12] |
| PXC1 | AT2G36570 | Wang et al., 2013 [13] |
| PXC2 | AT5G01890 | Wang et al., 2013 [13] |
| PXC3 | AT2G41820 | Wang et al., 2013 [13] |
| REV/AVB1/IFL1/ | AT5G60690 | transcriptional regulatory network of Smit et al. 2019 [2] |
| SCR | AT3G54220 | Wysocka-Diller et al. 2000 [14] |
| SERK1 | AT1G71830 | PC-associated genes of Kondo et al. 2015 [4] |
| SFC/FKD2/VAN3 | AT5G13300 | Sieburth et al. 2006 [15] |
| SHR/SGR7 | AT4G37650 | PC-associated genes of Kondo et al. 2015 [4] |
| SMXL3 | AT3G52490 | Miyashima et al. 2019 [7] |
| SMXL4 | AT4G29920 | Nagawa et al. 2006, Table 2 [12] |
| SMXL5 | AT5G57130 | Kondo et al. 2016 [16] |
| SUB1 | AT4G08810 | PC-associated genes of Kondo et al. 2015 [4] |
| T5L1/ABS5 | AT1G68810 | transcriptional regulatory network of Smit et al. 2019 [2] |
| TDR/PXY | AT5G61480 | PC-associated genes of Kondo et al. 2015 [4] |
| TMO5/AIG1/BHLH32 | AT3G25710 | transcriptional regulatory network of Smit et al. 2019 [2] |
| TMO6/DOF5.3 | AT5G60200 | PC-associated genes of Kondo et al. 2015 [4] |
| VDOF1/DOF4.6 | AT4G24060 | PC-associated genes of Kondo et al. 2015 [4] |
| VDOF2/DOF1.8 | AT1G64620 | transcriptional regulatory network of Smit et al. 2019 [2] |
| WOL/AHK4/CRE1 | AT2G01830 | transcriptional regulatory network of Smit et al. 2019 [2] |
| WOX14 | AT1G20700 | transcriptional regulatory network of Smit et al. 2019 [2] |
| WOX4 | AT1G46480 | transcriptional regulatory network of Smit et al. 2019 [2] |

1.  Jouannet, V., Brackmann, K., and Greb, T. (2015). (Pro)cambium formation and proliferation: two sides of the same coin? Curr. Opin. Plant Biol. *23*, 54–60. 10.1016/j.pbi.2014.10.010.

2.  Smit, M., McGregor, S., Sun, H., Gough, C., Bågman, A.-M., Soyars, C.L., Kroon, J.T.M., Gaudinier, A., Williams, C.J., Yang, X., et al. (2019). A PXY-Mediated Transcriptional Network Integrates Signaling Mechanisms to Control Vascular Development in Arabidopsis. Plant Cell. 10.1105/tpc.19.00562.

3.  Ohashi-Ito, K., Saegusa, M., Iwamoto, K., Oda, Y., Katayama, H., Kojima, M., Sakakibara, H., and Fukuda, H. (2014). A bHLH Complex Activates Vascular Cell Division via Cytokinin Action in Root Apical Meristem. Curr. Biol. *24*, 2053–2058. 10.1016/j.cub.2014.07.050.

4.  Kondo, Y., Fujita, T., Sugiyama, M., and Fukuda, H. (2015). A Novel System for Xylem Cell Differentiation in Arabidopsis thaliana. Mol. Plant *8*, 612–621. 10.1016/j.molp.2014.10.008.

5. Konishi, M., and Yanagisawa, S. (2015). Transcriptional repression caused by Dof5.8 is involved in proper vein network formation in Arabidopsis thaliana leaves. J. Plant Res. *128*, 643–652. 10.1007/s10265-015-0712-0.

6. Larrieu, A., Brunoud, G., Guérault, A., Lainé, S., Hennet, L., Stigliani, A., Gildea, I., Just, J., Soubigou-Taconnat, L., Balzergue, S., et al. (2022). Transcriptional reprogramming during floral fate acquisition. iScience *25*, 104683. 10.1016/j.isci.2022.104683.

7. Miyashima, S., Roszak, P., Sevilem, I., Toyokura, K., Blob, B., Heo, J., Mellor, N., Help-Rinta-Rahko, H., Otero, S., Smet, W., et al. (2019). Mobile PEAR transcription factors integrate positional cues to prime cambial growth. Nature *565*, 490–494. 10.1038/s41586-018-0839-y.

8. Geldner, N., Anders, N., Wolters, H., Keicher, J., Kornberger, W., Muller, P., Delbarre, A., Ueda, T., Nakano, A., and Jürgens, G. (2003). The Arabidopsis GNOM ARF-GEF mediates endosomal recycling, auxin transport, and auxin-dependent plant growth. Cell *112*, 219–230. 10.1016/s0092-8674(03)00003-5.

9. Ruiz Sola, M.A., Coiro, M., Crivelli, S., Zeeman, S.C., Schmidt Kjølner Hansen, S., and Truernit, E. (2017). OCTOPUS-LIKE 2, a novel player in Arabidopsis root and vascular development, reveals a key role for OCTOPUS family genes in root metaphloem sieve tube differentiation. New Phytol. *216*, 1191–1204. 10.1111/nph.14751.

10. Kim, H.-S., Kim, S.J., Abbasi, N., Bressan, R.A., Yun, D.-J., Yoo, S.-D., Kwon, S.-Y., and Choi, S.-B. (2010). The DOF transcription factor Dof5.1 influences leaf axial patterning by promoting Revoluta transcription in Arabidopsis. Plant J. *64*, 524–535. 10.1111/j.1365-313X.2010.04346.x.

11. Nisar, N., Cuttriss, A.J., Pogson, B.J., and Cazzonelli, C.I. (2014). The promoter of the Arabidopsis PIN6 auxin transporter enabled strong expression in the vasculature of roots, leaves, floral stems and reproductive organs. Plant Signal. Behav. *9*, e27898. 10.4161/psb.27898.

12. Nagawa, S., Sawa, S., Sato, S., Kato, T., Tabata, S., and Fukuda, H. (2006). Gene Trapping in Arabidopsis Reveals Genes Involved in Vascular Development. Plant Cell Physiol. *47*, 1394–1405. 10.1093/pcp/pcl009.

13. Wang, J., Kucukoglu, M., Zhang, L., Chen, P., Decker, D., Nilsson, O., Jones, B., Sandberg, G., and Zheng, B. (2013). The Arabidopsis LRR-RLK, PXC1, is a regulator of secondary wall formation correlated with the TDIF-PXY/TDR-WOX4 signaling pathway. BMC Plant Biol. *13*, 94. 10.1186/1471-2229-13-94.

14. Wysocka-Diller, J.W., Helariutta, Y., Fukaki, H., Malamy, J.E., and Benfey, P.N. (2000). Molecular analysis of SCARECROW function reveals a radial patterning mechanism common to root and shoot. Dev. Camb. Engl. *127*, 595–603. 10.1242/dev.127.3.595.

15. Sieburth, L.E., Muday, G.K., King, E.J., Benton, G., Kim, S., Metcalf, K.E., Meyers, L., Seamen, E., and Van Norman, J.M. (2006). SCARFACE Encodes an ARF-GAP That Is Required for Normal Auxin Efflux and Vein Patterning in Arabidopsis. Plant Cell *18*, 1396–1411. 10.1105/tpc.105.039008.

16. Kondo, Y., Nurani, A.M., Saito, C., Ichihashi, Y., Saito, M., Yamazaki, K., Mitsuda, N., Ohme-Takagi, M., and Fukuda, H. (2016). Vascular Cell Induction Culture System Using Arabidopsis Leaves

(VISUAL) Reveals the Sequential Differentiation of Sieve Element-Like Cells. Plant Cell *28*, 1250–1262. 10.1105/tpc.16.00027.

Figure 1. Vascular developmental stimulus induces active cellular reprogramming at 8 h.

(A) Effect of stimulus on an early procambial marker. YFP fluorescence was measured as mean pixel intensity x 10⁻³ from 2 mm diameter leaf disks of proAtHB8::4xYFP plants over 24 h treatment. (B) Effect of stimulus on relative RNA abundance of vascular differentiation markers. RNA abundance was measured by qRT-PCR over 20 h of treatment and normalized to the UBC10 reference RNA. (C) Pie chart showing the proportions of RNAs with increased or decreased RNA abundance (FDR < 0.05) by RNA-seq at 8 h stimulus treatment compared with mock. (D) Effect of stimulus on RNA abundances of members of the 'phloem or xylem histogenesis', 'auxin activated signaling', 'thylakoid', and 'response to light stimulus' gene ontologies . Color squares represent log2 relative RNA abundance (LFC) of GO members. BP, biological process; CC, cellular component. (E, F) Stimulus-induced changes in leaf vascular cell-identity (E) and mesophyll (F) markers. Box plot of log2 relative RNA abundance change (8 h stimulus vs mock) of markers identified by leaf single cell RNA-seq clusters generated by Kim et al (2021). (G) Experimental design schematic for RNA abundance and RNA decay analysis. (H) Pie chart showing proportion of RNAs with increased, decreased, or unchanged half-lives. (I) Distribution of $t_{1/2}$ changes. (J,K) Decay plots showing RNA abundance changes after inhibition of transcription (thin lines, mean±SE) and $t_{1/2}$ estimates based on modeled decay (thick lines) of 8 h stimulus- (blue and red) and mock-treated (black) leaves. (L) Distribution of $t_{1/2}$ (8 h mock, black dots) and shift in t1/2 (line segments) for 2,818 RNAs with shifted $t_{1/2}$s in the standard range. Line segment endpoints are positioned at the mock and stimulus $t_{1/2}$s. Line color indicates log2 relative abundance (LFC). See Figure S5A for enlargement of the 100 shortest and longest mock half-lives. Numbered line segments indicate RNAs that have much shorter or longer half-lives in stimulus: (1) Protein kinase family protein with leucine-rich repeat (AT1G35710), (2) TIR-NBS-LRR Disease resistance protein (AT4G36150), (3) CCCH-type zinc finger (AT5G58620), (4) hypothetical protein (AT2G07779), (5) hypothetical protein (AT5G66580), (6) snoRNA (AT1G32385), (7) FTSH11 (AT5G53170), (8) BTB/POZ domain-containing protein (AT5G59140), (9) CID4 (AT3G14010), (10) TRM8 (AT5G26910), (11) transcriptional regulator of RNA PolII, SAGA, subunit (AT4G33890), (12) Disease resistance protein (TIR-NBS-LRR class) (AT4G16900).

Figure 2. Broad regulation of RNA decay and rate of abundance change (RoAC) occurs during stimulus response.

(A) Effect of stimulus response on members of the early vascular regulator gene network. Heatmap of relative RNA abundance (8h stimulus vs. mock). Color and numbers indicate log fold change and fold change, respectively. Bold numbers indicate significant changes (FDR < 0.05). ‡, TMO6 is represented in two locations on the map. *, ATHB8 is represented twice.

(B) t$_{1/2}$ regulation in the EVR network (as in (A)). Color indicates t1/2. 11 RNAs shifted t1/2 and are indicated by a black outline and label.

(C) Schematic of RNA regulation occurring for EVR network RNAs with higher (top) or lower (bottom) abundance (FDR < 0.05).

(D) Schematic for determining rates of abundance change. Mock- and stimulus-treated tissue were collected after 6, 7, 8, 9, or 10 h to model the RoAC across the 8 h time point for each RNA.

(E) Distributions of RoAC for RNAs that were decreasing (left) or increasing (right) abundance at 8 h mock (top) or stimulus (bottom).

(F) Venn diagram showing RNAs with changing abundances in mock or stimulus.

(G) Pie chart showing the effect of stimulus on RoAC; rates increased (faster), decreased (slower), or were the same.

(H) Net effect of RNA regulatory mechanisms on RoAC (8 h stimulus vs. mock) in the EVR network (as in (A)). Positive or negative influence was found on rates of the 23 labeled RNAs.

(I-L) RNAs of the EVR network reveal distinct regulatory modalities through changes in transcription and decay rates as indicated by arrows: red,

Figure 3. Transcriptome-wide snapshot of 8h stimulus response includes active regulatory modalities and steady abundances maintained through changes in flux.

(A) Regulatory modalities are defined by change or lack of change in synthesis and decay and the net effect on rate of RNA abundance change (RoAC). Counts (and proportions) of RNAs exhibiting behavior of each regulatory modality are presented as a stacked bar plot.

(B-I) Regulation patterns of faster or slower synthesis and/or decay are demonstrated for each regulatory modality and are represented pictographically as larger red (faster TR/shorter t1/2) or smaller blue (slower TR/longer t1/2) arrows on either side of an oval. Ovals represent RNA abundance. Green arrows around the ovals represent increasing or decreasing RoAC. For each example RNA, abundance measurements (RPKM mean±SE) at 6, 7, 8, 9, 10 h of mock (thin light gray line) and stimulus (thin dark gray line) along with linear models of RoAC are shown for mock (thick black line) and stimulus (thick green line). Model parameters label the graphs: s, slope of the linear model (RPKM h-1); p, slope p-value; *, indicates a significant difference at the given time point, FDR < 0.05. TR and t1/2 are also shown in bar plots to the right for mock (m), and stimulus (s).

Figure 4. Dynamic range in transcription and t1/2 regulation are modality dependent.
(A-G) Contributions of transcription and decay regulation for individual RNAs. Stacked bar plots show direct comparison of regulation of TR and $t_{1/2}$ for each RNA of regulatory modalities; bars indicate direction and size of $\log_2$ relative TR (purple) and log2 relative $t_{1/2}$ (orange). Black dots indicate net regulation (sum of the LFC rel. TR and LFC rel. $t_{1/2}$). Genes within each regulatory modality are organized by increasing net regulation. The net regulation extremes that slightly deviate from 0 in the buffered RNAs reflects abundance variance below the cutoff.

Figure 5. Synergistic and oppositional modalities drive responsive RNA abundance.

(A) RNA abundance response profiles. k-means clustering of 4,268 RNAs of the transcription, decay, synergistic, and oppositional modalities. Abundance data of each RNA was normalized to the total (sum) abundance of all time points so clustering would group RNAs with similar abundance response profiles independent of level of expression. Clusters were ordered and numbered based on the relative abundance difference between mock (gray lines) and stimulus (green lines). Cluster number and size label each plot; label color indicates abundance trajectories at 8 h: diverging (blue), converging (pink), or other (black).

(B) Modality enrichment for RNAs of diverging or converging abundance clusters in (A) were grouped and evaluated for enrichment (blue) or depletion of the synergistic and oppositional regulatory modalities by the hypergeometric test; enrichment/depletion are displayed as a heatmap and labeled with -log10 p-values.

(C) Box and whisker plot of the abundance change magnitudes (|L2FC abundance|) by regulatory modality. Abundances of all stimulus time points were compared to those of all the mock time points.

(D) Enrichment analysis of RNA groups with distinct changes in $t_{1/2}$ in vcs mutants: vcs feedback decay (rel. $t_{1/2}$ vcs/WT < 1), moderate VCS–dependence (rel. $t_{1/2}$ vcs/WT > 1 & < 2), high VCS–dependence (rel. $t_{1/2}$ vcs/WT > 2). Significant enrichments are labeled with -log10 p-values.

Supplemental Figure S1. Stimulus induces developmental, molecular and biochemical changes to cells.

(A) YFP fluorescence of rosette leaves 1 and 2 of proATHB8::4xYFP plants incubated in mock and stimulus treatments for 24 and 72 h (bar = 1 mm).

(B) Development of xylem element characteristics in cells outside of vascular bundles after 72 h of stimulus treatment but not 72 h mock treatment. DIC micrograph of ectopic formation of spiral patterned secondary cell walls (Oda and Fukuda, 2012) can be seen (bar = 50 μm).

(C) Lignification of cell walls 72 h after stimulus treatment. Phloroglucinol was used to stain lignin (pink color). Box (dashed line) in the left column are enlarged to the right.

(D) Volcano plot overlayed on a histogram of 8 h RNA-seq comparison of stimulus vs mock.

(E) Heatmap of Pearson correlation values comparing similarity of abundance change of 460 RNA detected in both RNA seq data of this study and those measured by microarray by Furuya et al. (2021).

(F) Venn diagram comparing RNA identities of 8 h stimulus-upregulated RNAs from this study and stimulus-upregulated xylem- and phloem-specific RNAs after 72 h (Kondo et al., 2016).

(G) Leaf single cell RNA-seq UMAP clusters based on RNA expression measured by Kim et al. (2021). PC, procambium; PP, Phloem parenchyma; XP, xylem parenchyma. Compare Figure 1E, 1F.

Supplemental Figure S2. RNA $t_{1/2}$ extremes have unique features.

(A) Dynamic ranges of RNAs that shift to or from extremely long half-lives (>1440 min) were separated from other RNAs.

(B) At $t_{1/2}$ extremes, short short-lived RNAs tended to become longer-lived, and long-lived RNAs tended to become shorter-lived. The 100 shortest and 100 longest $t_{1/2}$s in mock are shown. Line segments represent shift in $t_{1/2}$ and are positioned at the mock $t_{1/2}$ (black dots). Line color indicates $\log_2$ relative abundance (LFC).

(C) Dynamics of long-lived RNAs ($t_{1/2} \geq 1440$ min). Some RNAs are long-lived in mock but move out of this group in stimulus (blue circle), others become long-lived in stimulus (green circle), and others remain at $t_{1/2} \geq 1440$ min in stimulus and mock.

(D) Bar plot comparing proportion of mock $t_{1/2}$ quintiles with shifts in decay.

.

# A
## EVR Network (65 RNAs)

**RNA abundances**
RNA-seq
8 h stimulus vs mock

**62 RNAs with abundance data**
8 lower, 43 higher, 11 unchanged

**RNA $t_{1/2}$**
8 h stimulus & mock;

**57 RNAs with measured $t_{1/2}$**
4 longer, 7 shorter, 46 unchanged
in stimulus vs mock

## Regulatory Conclusions

Δ abundances (51)    $t_{1/2}(s) \neq t_{1/2}(m)$ (11)

5 | 36 | 10 | 1

no $t_{1/2}$ data

**transcription (36):**
Δ abundances & $t_{1/2}(s) = t_{1/2}(m)$

**regulation includes change in $t_{1/2}$ (10):**
Δ abundances & $t_{1/2}(s) \neq t_{1/2}(m)$

39 higher abund. | 7 lower abund.
30 | 6
1 longer $t_{1/2}$ | 6 | 1 shorter $t_{1/2}$ | 4 longer $t_{1/2}$ | 6 shorter $t_{1/2}$

**cooperative** $t_{1/2}$    **antagonistic** $t_{1/2}$

# B
## EVR Network (65 RNAs)

**rate of abundance change (RoAC)**
RNA-seq
6, 7, 8, 9, 10 h stimulus vs mock

**RNA $t_{1/2}$ (57)**
8 h stimulus & mock

**57 RNAs with RoAC data**
34 steady in mock and stimulus,
23 with active rates of change

**transcription rate**
calculation based on rate of
abundance changes and $t_{1/2}$

**57 transcription rates**
42 faster in stimulus
10 slower in stimulus
5 unchanged

## Regulatory Conclusions

**transcription (19)**
TR changes, $t_{1/2}(s) = t_{1/2}(m)$ -->
RoAC changes

**decay (1)**
TR(s) = TR(m), $t_{1/2}(s) < t_{1/2}(m)$ -->
lower RoAC

**synergistic (1)**
TR(s) > TR(m), $t_{1/2}(s) > t_{1/2}(m)$ -->
faster RoAC

**oppositional (2)**
TR(s) >> TR(m), $t_{1/2}(s) < t_{1/2}(m)$ -->
faster RoAC

**[steady (34)]**
RoACs remain steady

# C
## Transcriptome (18,563 RNAs)

**rate of abundance change (RoAC)**
RNA-seq
6, 7, 8, 9, 10 h stimulus vs mock

**RNA $t_{1/2}$ (15,083)**
8 h stimulus & mock

**14,841 RNAs with RoAC data**
9,714 steady in mock and stimulus,
5,127 with active rates of change

**transcription rate**
calculation based on rate of
abundance changes and $t_{1/2}$

**14,841 rates**
241 quality filtered

**14,600 transrcription rates**
5,639 faster in stimulus
6,302 slower in stimulus
2,659 unchanged

## Regulatory Conclusions

**transcription (3108)**

**decay (64)**

**synergistic (106)**

**oppositional (990)**

**steady (9,714)**

**other (618)**

Supplemental Figure S3. Analysis flowchart for comparing abundances, t1/2s, and TRs of mock and stimulus treatments.
(A) Initial EVR network analysis by comparison of 8 h abundance with t1/2.
(B) EVR network analysis using rate of abundance change (RoAC), t1/2 regulation, and calculated TR analysis.
(C) Whole transcriptome regulation analysis as in (B). See (B) for modality assignment criteria.

Supplemental Figure S4. Early stimulus transcriptional response is varied and dynamic and includes buffered RNAs.

(A) Distributions of TR for mock and stimulus RNAs with changing abundance presented as histograms.

(B) Distribution of log2 relative TR (stimulus/mock) as a histogram. Solid light blue lines indicate the cutoffs for unchanged TR (1 +/- 0.18).

(C) Examples of RNA buffering from steady group 2. Buffered RNAs have steady abundances but their TR and $t_{1/2}$ differ. For each example RNA, abundance measurements (RPKM mean±SE) at 6, 7, 8, 9, 10 h of mock (thin light gray line) and stimulus (thin dark gray line) along with linear models of rate of abundance change are shown for mock (thick black line) and stimulus (thick green line). Model p-values label the graphs. TR and $t_{1/2}$ are also shown in bar plots to the right for mock (m), and stimulus (s).

Supplemental Figure S5. Stacked bar-plot extended analysis.

(A,C-E) Dynamic range of decay rate shifts (orange line segments, left column) in regulatory modalities. Line segment endpoints indicate mock and stimulus half-lives. RNAs are ordered on the y-axis by increasing $t_{1/2}$ mean of stimulus and mock treatments (indicated by a black circles). Inset histograms show distribution of $\log_2$ relative $t_{1/2}$.

(B,C-E) Dynamic range of TR changes (purple line segments, right column) in regulatory modalities. Line segment endpoints indicate mock and stimulus TRs. RNAs are ordered on the y-axis by increasing mean TR of stimulus and mock treatments (indicated by black circles). Inset histograms show distribution of log2 relative TR.

(F) Oppositional and steady 2: Buffered stacked bar plots with dashed-line boxes indicating expanded regions in (H) and (I), respectively. Asterisks highlight exceptionally large and nearly symmetric changes in TR and $t_{1/2}$.

(I,J) Comparison of abundances and $t_{1/2}$ shifts of RNA buffered in sov mutants and buffered (steady 2).

(K) Gene ontology enrichment analysis of regulatory modality RNAs. The decay, synergistic, and steady 1 modalities did not have enriched GOs (p-adj, adjusted p-value).