Unreliable homeostatic action potential broadening in cultured dissociated

neurons

Andreas Ritzau-Jost[1,#], Salil Rajayer[2,3], Jana Nerlich[1], Filip Maciag[4], Alexandra

John[1], Michael Russier[5], Victoria Gonzalez Sabater[6,7], Luke J. Steiger[2,3], Jacques-

Olivier Coq[5], Jens Eilers[1], Maren Engelhardt[8,9,10], Juan Burrone[6,7], Dominique

Debanne[5], Martin Heine[4], Stephen M Smith[2,3,#], Stefan Hallermann[1,#]

[1] Carl-Ludwig-Institute of Physiology, Faculty of Medicine, Leipzig University, Leipzig, 04103, Germany

[2] Division of Pulmonary and Critical Care Medicine, Department of Medicine, Oregon Health and

Science University, Portland, Oregon, 97239-3098, USA

[3] Section of Pulmonary and Critical Care Medicine, VA Portland Health Care System, Portland, Oregon,

97239-2964, USA

[4] Institute for Developmental Biology and Neurobiology, Johannes Gutenberg University, Mainz,

55128, Germany

[5] Unité de Neurobiologie des canaux Ioniques et de la Synapse (UNIS), UMR_S 1072, INSERM, Aix-

Marseille Université, Marseille, 13015, France

[6] MRC Centre for Neurodevelopmental Disorders, Institute of Psychiatry, Psychology and

Neuroscience, King's College London, London, SE1 1UL, United Kingdom

[7] Centre for Developmental Neurobiology, Institute of Psychiatry, Psychology and Neuroscience,

King's College London, London, SE1 1UL, United Kingdom

[8] Institute of Neuroanatomy, Mannheim Center for Translational Neuroscience, Medical Faculty

Mannheim, Heidelberg University, Mannheim, 68167, Germany

[9] Institute of Anatomy and Cell Biology, Johannes Kepler University, Linz, 4020, Austria

[10] Clinical Research Institute for Neurosciences, Johannes Kepler University, Linz, 4020, Austria

[#] Correspondence: andreas.ritzau.jost@gmail.com (A.R-J.), smisteph@ohsu.edu (S.M.S.), and

hallermann@medizin.uni-leipzig.de (S.H.)

Summary

Homeostatic plasticity preserves neuronal activity against perturbations. Recently,

somatic action potential broadening was proposed as a key homeostatic adaptation to

chronic inactivity in neocortical neurons. Since action potential shape critically controls

calcium entry and neuronal function, broadening provides an attractive homeostatic

feedback mechanism to regulate activity. Here, we report that chronic inactivity

induced by sodium channel block does not broaden action potentials in neocortical

neurons under a wide range of conditions. In contrast, action potentials were

broadened in CA3 neurons of organotypic hippocampal cultures by chronic sodium

channel block and in hippocampal dissociated cultures by chronic synaptic block.

Mechanistically, BK-type potassium channels were proposed to underly inactivity-

induced action potential broadening. However, BK channels did not affect action

potential duration in our recordings. Our results indicate that action potential

broadening can occur in specific neurons and conditions but is not a general

mechanism of homeostatic plasticity in cultured neurons.

Introduction

Action potentials (APs) control various functions including neurotransmitter release from axonal boutons[1], calcium signals in the soma[2], and synaptic plasticity in dendrites[3]. AP broadening during short-term plasticity on the time scale of milliseconds to seconds is well-established and occurs in many types of neuronal cell bodies[4,5] and nerve terminals[6,7]. There are also indications that APs change during long-term plasticity on the time scale of minutes to hours. When examining Hebbian long-term plasticity (i.e., positive feedback-based and input-specific long-term plasticity[8]), some studies reported AP broadening. For example, presynaptic APs of hippocampal mossy fiber boutons broaden during depolarization-induced potentiation[9] and somatic APs of neurons in the amygdala broaden during fear conditioning[10]. In contrast, the role of the AP duration is less clear for non-Hebbian homeostatic long-term plasticity (i.e., negative feedback-based and non-input specific long-term plasticity[8]). Specifically, homeostatic plasticity induced by neuronal inactivity in the presence of the sodium channel blocker tetrodotoxin (TTX) has been studied for decades, but AP broadening was initially not described[11,12]. A recent study however reports prominent AP broadening during TTX-induced homeostatic plasticity at the soma of cultured neocortical neurons[13], which was not observed at presynaptic boutons of cultured neocortical neurons[14]. Compartmentalized AP broadening at the neuronal cell body would provide a strikingly simple and previously overlooked mechanism by which neuronal inactivity could increase subsequent excitability[13]. This mechanism involves decreased calcium entry during TTX application causing changes in gene expression that would result in broadened APs and more calcium entry after TTX removal. In this model, AP broadening is a key link in the homeostatic feedback loop underlying homeostatic plasticity. In a collaborative effort across several laboratories, we therefore systematically investigated AP broadening in cultured neurons during homeostatic plasticity under various experimental conditions.

Results

Cell type- and model-specific AP broadening in hippocampal neurons

APs in CA3 neurons of hippocampal organotypic slice cultures broaden homeostatically upon chronic inactivity using both, synaptic block by glutamate receptor antagonists[15] or sodium channel block by TTX[16] (see also ref. 17 for a similar trend). We first aimed to confirm homeostatic AP broadening in CA3 neurons in

organotypic slice cultures. APs were elicited by depolarizing currents from the resting

membrane potential during perfusion with TTX-free artificial cerebrospinal fluid. APs

recorded in CA3 pyramidal neurons treated with TTX (48 hours) had longer durations

than those in control cells (Fig. 1A, B; median [IQR] 1.75 [1.69–1.86] ms and 2.13

[2.07–2.22] ms for control and TTX-treatment, respectively, 14 cells each, $P < 0.001$),

confirming earlier reports of homeostatic AP broadening in these neurons[15,16].

Next, we tested whether homeostatic AP broadening occurred in dissociated

hippocampal cultured neurons, a model widely used to study homeostatic plasticity.

We inhibited neuronal activity for 48 hours to induce homeostatic plasticity by either

blocking AP firing using TTX (Fig. 1C) or by blocking synaptic transmission using the

low-affinity glutamate receptor blocker kynurenic acid (Kyn) or the high-affinity

glutamate receptor blocker 2,3-Dioxo-6-nitro-1,2,3,4-tetrahydrobenzo [f]quinoxaline-7-

sulfonamide (NBQX). Surprisingly, AP duration was unaffected by TTX-treatment in

primary hippocampal cultured neurons (Fig. 1D; median [IQR] 2.07 [1.79–2.34] ms and

2.08 [1.32–2.61] ms for control and TTX, 25 and 24 cells, respectively, $P = 0.96$). In

contrast, Kyn-treatment significantly increased AP duration in the same set of

interleaved experiments (Fig. 1D; median [IQR] 2.52 [2.31–2.72] ms, 22 cells, $P =$

0.014) and NBQX-treatment led to a small trend towards broader APs (Fig. 1D; $P =$

0.26). Importantly, sodium channel block and synaptic block both increased

spontaneous AP firing frequency, indicating that homeostatic adaptations other than

AP broadening were shared between both induction methods (Fig. 1E, F; $P = 0.008$

for control against Kyn, $P = 0.005$ for control against NBQX, $P = 0.013$ for control

against TTX). These data indicate that AP broadening occurred upon sodium channel

block in CA3 neurons in organotypic slice cultures and upon synaptic block in

dissociated hippocampal cultures, but not in the commonly-studied model of

homeostatic plasticity using dissociated hippocampal cultured neurons and 48 hours

of TTX application.

Stable AP duration in primary dissociated neurons during TTX-induced

homeostatic plasticity

In contrast to our finding in dissociated cultured neurons of the hippocampus, a recent

study described TTX-induced AP broadening in dissociated cultured neurons of the

neocortex[13]. To resolve this discrepancy, we set out to replicate AP broadening in

neocortical cultured neurons. Given that exact reproduction of experimental conditions

is challenging and neuronal plasticity depends on brain region[18], animal species[19], and

animal strain[20], we probed homeostatic AP broadening under different experimental

conditions varying: 1) experimental animal species (rat and mouse), 2) strain (C57BL/6

and CD1 mice), 3) sub-strain (C57BL/6N and C57BL/6J mice), 4) brain area (cortex

and hippocampus), 5) cortical subdomain (prefrontal cortex), 6) culture duration (10–

14, 15–21, and 32 days *in vitro*), 7) culture growth medium, 8) recording solutions, and

9) laboratory and experimenter performing recordings (conditions I–X in Fig. 2B,

condition XI are recordings from Fig. 1; see also Fig. 2 – figure supplement 1). Even

though AP durations varied up to 2-fold between conditions, statistically significant

homeostatic AP broadening was not detectable in any of the tested conditions (Fig.

2B). To minimize type II errors (false negative) we intentionally did not apply a

correction for multiple comparisons. The only significance was observed in condition

III but in an opposite direction (i.e. AP narrowing with TTX, P=0.026; Fig. 2B). However,

this is likely a false positive because application of corrections for false discovery rate

results in P=0.268 for both Benjamini–Hochberg and Bonferroni correction. To detect

cross-conditional AP broadening by TTX, we merged all conditions based on their

difference in AP duration after TTX-treatment (Fig. 2C). However, AP duration was the

same in control and TTX-treated cells (217 and 197 cells, respectively, P = 0.83).

We next tested whether AP broadening was concealed by systematic differences in

neuronal health or recording quality between control and treatment group. Because

unhealthy neurons tend to have small and slow APs, possibly due to changes in resting

membrane potential or expression of voltage-gated sodium and potassium channels,

we first analyzed AP amplitude as a measure of neuronal viability. AP amplitudes were

not affected following TTX treatment in any of the eleven recording conditions (Fig. 2D)

or a cross-conditional comparison (Fig. 2E). In addition, we focused on the culture

protocol or cortical subdomain previously associated with AP broadening[13] (conditions

I, VII, and VIII) and applied further inclusion criteria to rigorously discriminate healthy

excitatory neurons: the occurrence of multiple APs upon 200 ms-current injections (≥

2 APs), a resting membrane potential below − 55 mV, and a maximum AP

depolarization rate between 100 and 337 V/s (includes 96% of excitatory and exclude

>80% of the inhibitory neurons; ref. 21). Applying any or all three of these criteria did

not reveal AP broadening upon TTX treatment (Fig. 2 – figure supplement 2). These

data indicate that stricter exclusion criteria result in smaller variability in action potential

duration but do not uncover AP broadening in these datasets.

Instead of depolarization-evoked APs, Li *et al.*[13] recorded hyperpolarization-evoked

'rebound APs' (Fig. 2F), possibly reducing the impact of different resting membrane

potentials and depolarizing current injection on the AP waveform. We determined that

the duration of rebound APs was similarly unaffected by TTX treatment across all

recorded conditions (Fig. 2G, H). These data show that TTX-induced AP broadening

does not robustly occur in dissociated cultured neurons.

Homeostatic plasticity increases network activity and excitatory synaptic

transmission

To verify successful induction of homeostatic plasticity under our experimental

conditions, we quantified changes in neuronal activity and synaptic strength following

TTX-induced homeostatic plasticity. TTX-treatment increased the number of

spontaneously active neurons (Fig. 3A; merge of conditions I–IV and VII; control: 23

out of 87 cells, TTX-treatment: 58 out of 74 cells, $P < 0.001$ with Fisher's exact test)

and the number of APs per spontaneous burst (Fig. 3B, C; median [IQR] 1.0 [1.0–1.12]

APs and 1.67 [1.0–2.71] APs for control and for TTX-treatment, 23 and 58 cells,

respectively, $P < 0.001$). These results indicate an adaptive increase in neuronal

activity following TTX-treatment, in line with previous findings on homeostatic

plasticity[16,22,23]. Furthermore, spontaneous excitatory postsynaptic currents (sEPSCs;

spontaneous synaptic currents recorded in the absence of TTX) were recorded as a

measure of synaptic strength (Fig. 3D). sEPSC frequency and amplitude increased

upon TTX-treatment (Fig. 2E, merge of recordings from condition I and II; median [IQR]

frequency 1.11 [0.42–1.47] Hz and 5.07 [1.95–5.52] Hz in control and TTX-treated

cells, 16 and 11, respectively, $P = 0.027$; median [IQR] amplitude 12.4 [10.6–14.4] pA

and 15.5 [14.1–17.4] pA in control and TTX-treated cells, 16 and 11 cells, respectively,

$P = 0.042$), in accord with known homeostatic synaptic adaptations to inactivity[24–27].

These changes in sEPSC amplitude and frequency are not specific for somatic, pre-

or postsynaptic adaptations. However, the results show that blocking AP firing with

TTX successfully induced homeostatic plasticity under our experimental conditions.

Lack of BK channel-dependent AP broadening in dissociated cultured neurons

Various potassium channel subtypes regulate AP repolarization and could therefore underlie AP broadening. Li *et al.*[13] reported that downregulation of BK-type potassium channels led to homeostatic AP broadening in neocortical cultured neurons. We tested the role of BK channels in AP repolarization (neurons grown under condition I). However, blocking BK channels with Iberiotoxin (IbTx, 300 nM) did not lead to broadening of depolarization-evoked APs in control and TTX-treated cells (Fig. 4A; P = 0.10 and 0.48, respectively), or a merge of both conditions (Fig. 4B; P = 0.10). Rebound APs were similarly unaffected by BK channel block (Fig. 4 – figure supplement 1). Next, we tested whether recurrent neuronal activity, which increases cytosolic calcium and boosts BK-channel activity, enhanced AP broadening following TTX treatment. While APs significantly broadened during repetitive firing, AP durations were similar in control and TTX-treated neurons (Fig. 4C and D), suggesting no differential regulation of BK-channels following TTX-treatment in these recordings.

In electrical recordings, intracellular recording solution may have disturbed cytosolic calcium dynamics and therefore BK- and thus calcium-dependent AP repolarization. As an independent approach, we recorded the AP duration in hippocampal cultured neurons using the genetically encoded voltage indicator (GEVI) Ace-mNeon[28] before and after acute IbTx perfusion (100 nM). BK channel block did not affect AP duration in the soma or along the axon (Fig. 4E). Using the GEVI, we again tested the role of BK channels in AP broadening during a train of APs (5 APs at 20 Hz, Fig. 4F). The broadening of somatic, proximal axonal, and distal axonal APs during trains was similar before and after BK channel block, indicating that AP duration was not modulated by BK channels. To confirm that IbTx was indeed pharmacologically active in Ace-mNeon recordings, we preconditioned cells with 4-AP (30 µM) prior to IbTx perfusion (Fig. 4G). Neurons preconditioned with 4-AP had broader somatic APs which were further broadened by IbTx, presumably due to increased calcium influx following 4-AP treatment, in turn activating more BK channels[29]. These data indicate that IbTx application was pharmacologically active but BK channels did not contribute to AP repolarization in cultured neocortical neurons.

Discussion

Homeostatic AP broadening has been reported in some systems, but it has not been tested systematically across experimental models so far. Because BK channels that were proposed to underlie AP broadening are activated by depolarization and calcium[30,32], we studied APs elicited with depolarizations from the RMP where BK activity was presumably increased, as well as anode break ('rebound') APs thought to obviate variance from ion channel inactivation. We found that broadening of rebound and depolarization-evoked APs did not occur following the induction of homeostatic plasticity with TTX in the most commonly studied model system (primary dissociated neuronal cultures). However, AP broadening was observed in cultured hippocampal slices. Furthermore, blocking BK channels, which were reported to mediate AP broadening during homeostatic plasticity[13], did not impact AP duration in dissociated cultured neurons. Our results therefore indicate that AP broadening in cultured dissociated neurons is not a required component of homeostatic plasticity.

TTX-induced homeostatic AP regulation in organotypic hippocampal cultured neurons

AP broadening was previously shown to occur in CA3 neurons of organotypic hippocampal slice cultures following TTX-induced homeostatic plasticity[16]. Consistently, we recorded broader APs in CA3 neurons following neuronal inactivity (Fig. 1). In contrast to CA3 neurons, granule cells of the dentate gyrus did not show TTX-induced AP broadening[17]. Similarly, CA1 neurons did not show AP broadening following synaptic block-induced homeostatic plasticity[31]. Therefore, we conclude that homeostatic AP broadening in organotypic hippocampal cultures is most likely cell type-specific.

TTX-induced homeostatic AP regulation in primary dissociated cultured neurons

To the best of our knowledge, only three studies so far reported on the duration of APs following TTX-induced homeostatic plasticity in dissociated cultured neurons[13,33,34]. While Li *et al.*[13] observed TTX-induced broadening, the studies by Lee, Chung, and others[33,34] reported stable AP duration, consistent with our findings (Fig. 2). Furthermore, in other studies investigating TTX-induced homeostatic plasticity in dissociated cultured neurons, prominent AP broadening was either missed or absent (see for example refs. 23,35–37). One possible explanation for this inconsistency could

be the cell type-specificity of AP broadening. We focused on large, presumably

excitatory neurons. However, we and others could have selected different neuron

types compared to Li *et al.*[13]. In an attempt to reduce bias, we incorporated multi-lab,

randomization, and blinding strategies across many of our experiments. Only

recordings with leaky seal formation or inadequate capacitance compensation[38] were

excluded. However, in a subset of experimental conditions we applied additional

exclusion criteria to ensure that variability in neuron type, health or recording quality

were not causing a type II (false negative) statistical error. Application of these

exclusion criteria did not reveal AP broadening in our data (Fig. 2 – figure supplement

2), confirming that AP broadening was not associated with homeostatic plasticity in

neocortical neurons following TTX treatment.

The sensitivity to chronic TTX treatment might depend on baseline neuronal activity,

which is in part related to neuronal culture density[39]. However, TTX did not induce AP

broadening despite different baseline activities (Fig. 3A) and a nearly threefold

variation in the number of plated cells per cover slip between conditions (25k – 70k

cells per cover slip). Although we mainly focus on neocortical cultured neurons

(condition I to VIII, Fig. 2) because Li *et al.* used neocortical neurons, the absence of

AP broadening in hippocampal neurons (group IX to XI) could in principle be explained

by the selective loss of CA3 neurons, which show AP broadening in organotypic

cultured neurons (Fig. 1A and B). However, CA3 neurons were shown to survive in

dissociated cultures following region-specific microdissection[40], and CA1 neurons are

generally more stress-sensitive to excitotoxicity with glutamate or NMDA than CA3 and

DG neurons[42], arguing against a general selective loss of CA3 neuron in dissociated

cultures. Altered cell-cell interactions with glia and neurons in organotypic and

dissociated neuronal cultures could instead contribute to the different findings in

various hippocampal preparations.

Synaptic block-induced homeostatic AP regulation in primary dissociated

cultured neurons

Homeostatic AP broadening was shown in CA3 neurons upon activity block by the low-

affinity glutamate receptor blocker Kyn[15]. We similarly observed Kyn-induced AP

broadening in dissociated cultured hippocampal neurons (Fig. 1C, D). In contrast, APs

were not significantly broader following synaptic block by NBQX (Fig. 1C, D), in accord

with recordings from CA1 neurons in organotypic cultures using CNQX. TTX-induced

broadening may therefore be cell-type specific or due to a differential effect of the

glutamate receptor blockers on NMDA receptors which are blocked by Kyn but not

NBQX/CNQX or TTX and which have recently been demonstrated to be important for

the induction of synaptic homeostatic plasticity[41]. Further studies are needed to

investigate the role of subcellular calcium signals in the regulation of AP duration in

hippocampal neurons during block of glutamate receptors.

Absence of BK channel-dependent AP broadening in dissociated cultured

neurons

Li *et al.*[13] proposed that BK channels hasten AP repolarization and that this effect is

lost during TTX-induced homeostatic plasticity, leading to AP broadening. We did not

detect AP broadening when blocking BK channels in either control or TTX-treated cells

(Fig. 4). Furthermore, because calcium accumulation following multiple APs has been

shown to enhance intraspike BK contribution in some neurons[43] we compared how 1st,

2nd, or 3rd APs were impacted by prior TTX-exposure. AP broadening did occur in the

later APs but the changes were unaffected by TTX exposure (Fig. 4D), suggesting no

activity-dependent effect of TTX or BK channels on AP duration.

Voltage imaging using Ace-mNeon similarly showed that duration of single APs and

broadening during AP trains in the soma and axon were unaffected by BK channel

block. Blocking BK channels only affected AP repolarization when cells were

preconditioned with 4-AP. These results are consistent with previous reports showing

that in some neurons BK channels do not contribute to repolarization[43,44] while doing

so in others[44–47]. Differences in the abundance of other types of potassium channels[44]

and intracellular calcium buffering[46] are known to affect the contribution of BK channels

to AP repolarization and may account for the lack of AP broadening upon BK channel

block in our dissociated cultured neurons. Alternatively, BK activity may occur only

under conditions associated with elevated intracellular[30,32]. The lack of BK channel

contribution to AP repolarization in our dissociated neurons is another possible

explanation (in addition to different selection of cells) for why we and others have not

observed the homeostatic AP broadening described by Li *et al.*[13].

Inactivity-induced homeostatic AP regulation *in vivo*

Homeostatic plasticity has also been studied using *in vivo* recordings from the activity-
suppressed visual cortex[48,49]. The majority of these studies reported stable AP duration
upon monocular deprivation or chemogenetic activity suppression[50–52]. In contrast, Li
et al.[13] found homeostatically broadened APs in the visual cortex upon monocular
deprivation. Furthermore, stable AP duration was also observed in other brain areas
following activity suppression such as (1.) the somatosensory cortex upon unilateral
whisker trimming[53–58] or forepaw denervation[59], (2.) the auditory system upon cochlear
ablation[60,61], hearing loss[62], or congenital deafness[63], (3.) the vestibular nucleus upon
vestibular ablation[64], and (4.) the olfactory mitral cells upon naris occlusion[65] or
senescent hippocampal deafferentiation[66]. Besides somatic APs, APs in excitatory
presynaptic terminals of the somatosensory thalamus were also unaffected by whisker
deprivation[67]. Thus, AP broadening does not appear to occur robustly upon
suppression of neuronal activity in various neuronal pathways.

Hyperactivity-induced homeostatic AP regulation
Homeostatic plasticity occurs not only upon activity block but also upon hyperactivity,
generally involving opposite adaptations. In cultured neurons, AP duration was
unaffected following hyperactivity induced by chronic depolarization[68], block of
potassium channels[69], block of inhibitory synaptic transmission[31,33,34], or chronic ChR2-
induced photostimulation[70]. In contrast, Li *et al.*[13] reported similar BK channel
downregulation upon both, TTX-induced activity block and depolarization-induced
hyperactivity.

As in cultured neurons, AP duration was mostly unaffected in brain slice recordings
from animals with increased neuronal activity due to an enriched environment[58,71,72],
psychosocial stress[73], or febrile seizures[74]. AP shortening occurred in cerebellar
granule cells after exposure to an enriched environment[75], in auditory neurons
following chronic noise exposure[76], and in hippocampal neurons following age-
related[77] or pilocarpine-induced hyperactivity[78]. Thus, in contrast to Li *et al.*[13], AP
duration was either unaffected or, if anything, shortened during hyperactivity-induced
homeostatic plasticity in previous studies.

In this study, several laboratories collaborated to determine the robustness of
inactivity-induced AP broadening. AP broadening did not occur upon TTX-treatment in

dissociated hippocampal or neocortical neurons. Furthermore, BK channel block did

not affect AP duration in dissociated neurons, arguing against BK channels as

mediators of a general homeostatic AP regulation.

Despite the lack of homeostatic, TTX-induced AP broadening in dissociated cultures,

AP duration was broadened upon Kyn-treatment in dissociated cultures and using TTX

in CA3 neurons in organotypic cultures. Because BK-channels control AP duration in

CA3 neurons of organotypic cultures[79], homeostatic BK-channel downregulation as

proposed by Li *et al.* may be involved in AP broadening in this specific cell type. While

the reasons for the variable occurrence of homeostatic AP broadening remain

unknown, this may render neuronal circuitries more robust to perturbations. The

regulation of AP duration therefore might represent one element in the repertoire of

neuronal plasticity that is, similar to other plasticity mechanisms, not generally shared,

but specifically expressed in some cell types and neuronal compartments.

METHODS

Ethics

At all times animal procedures were in accordance with the relevant international,

national and institutional guidelines for the care and use of laboratory animals

(European Council Directive 86/609/EEC, U.S. Public Health Service Policy on

Humane Care and Use of Laboratory Animals, U.S. National Institutes of Health Guide

for the Care and Use of Laboratory Animals, German Tierschutzgesetz, French

National Research Council guidelines, and Leipzig and Mainz University guidelines).

All animal procedures were approved in advance by the local health authorities (Ethics

Committees at Leipzig University and Mainz University, federal Saxonian Animal

Welfare Committee, the District administration Mainz-Bingen (41a/177-5865-§11

ZVTE), the Committee of the Préfecture des Bouches-du-Rhône (D13055-08), V.A.

Portland Health Care System Institutional Animal Care and Use Committee).

Animal models

Organotypic hippocampal cultures were generated from postnatal day 5–7 Wistar rats.

Dissociated neocortical cultures were generated from postnatal day 0 or 1 Sprague-

Dawley rats (conditions I, II and VII), C57BL/6N mice (conditions III and IV), C57BL/6J

mice (conditions V and VIII), or CD1 mice (condition VI). Dissociated hippocampal

cultures were generated from postnatal day 0 or 1 C57BL/6N mice (conditions IX),

C57BL/6J mice (condition X), or Sprague-Dawley rats (condition XI). Generally,

animals of both sexes were used to generate cultures. Mother animals were kept in

individually ventilated cages under a 12h/12h light-dark cycle and received water and

food *ad libitum*.

Induction of Homeostatic Plasticity

Homeostatic plasticity was induced by inclusion of one of the respective blockers in

culture medium for 48–56 hours prior to recordings. All blockers were first diluted in

100 µL of fresh medium per treated well and subsequently added to the respective

wells containing 1 mL of medium. For control conditions, only 100 µL of fresh medium

without blockers was added to the respective wells. The blockers had final

concentrations of (in mM): 0.002 tetrodotoxin (TTX; Tocris, Wiesbaden-Nordenstadt,

Germany), 2 kynurenic acid (Kyn), or 0.01 2,3-Dioxo-6-nitro-1,2,3,4-tetrahydrobenzo-

[f]quinoxaline-7-sulfonamide (NBQX). To avoid bias, coverslips were randomized to

treatment (control or blocker) and recordings were made and analyzed in a blinded

fashion (except for condition IV and VI). Blinding was performed by an independent

investigator. Control or TTX-treatment groups were assigned using an 8-block

randomization approach for experiments of condition VIII. Unblinding of the

experimental groups was performed after analysis and disclosure of the data by the

experimenter.

Preparation of dissociated neocortical cultures

Conditions I and II (Hallermann Lab, Leipzig, Germany)

Neocortical cultures were generated as described by Li *et al.*[13]. Shortly, neocortices

were isolated and dissected into 5–10 pieces per hemisphere. Samples were washed

twice in ice-cold modified HBSS (4.2 mM $NaHCO_3$ and 1 mM HEPES, pH 7.35, 300

mOsm) supplemented with 20% fetal bovine serum (FBS) and digested at 37 °C for 30

min using papain. After stopping digestion with 20% FBS-containing modified HBSS,

samples were washed and dissociated using Pasteur pipettes, pelleted twice, and

filtered using a 70 nm nylon strainer. 30.000–50.000 vital dissociated cells were plated

on poly-D-lysine-coated 14 mm coverslips and cultured in NbActive4 medium

(Transnetyx, Cordova, TN, USA) in an incubator until use (37 °C, 5% CO_2 in room air,

water vapor saturated).

Conditions III and VII (Hallermann Lab, Leipzig, Germany)

Cultures were generated as previously described by Moutin *et al.*[80]. In brief, mice were

decapitated and neocortices (for condition III) or the frontal third of neocortices (for

condition VII) were isolated in ice-cold Hibernate-A medium (Thermo Fisher Scientific,

Waltham, MA, USA) and cut into 5–10 chunks. Chunks were first digested by papain

(dissolved in Hibernate-A; 10 min at 37 °C) and subsequently by papain + DNAse (125

µL of 20 mg/mL DNAse stock; 5 min at 37 °C). Digestion was stopped by CM+ medium

(Neurobasal A 86.5 %, B27 2 %, Glutamax 0.25 %, L-Glutamine 0.25%, penicillin

G/streptomycin 1 %, heat-inactivated FBS 10 %) and chunks were mechanically

dissociated in two steps by 1250 µL pipette tips with gravity decantation and collection

of the supernatant between dissociation steps and after dissociation. Supernatant was

passed through 2 mL of frothed bovine serum albumin, centrifuged (7 min, 300 x g),

the resulting pellet was resuspended in 1–2 mL of CM+ medium and 25.000–30.000

resuspended vital cells were plated onto poly-L-lysine-coated coverslips. After 30 min,

0.5 mL of CM+ medium was added to the wells and after 2 days *in vitro* (DIV), cytosine

arabinoside (ARA-C, 1 µM final concentration) was added to the wells for 8–18 h.

Subsequently, 0.4 mL medium per well was replaced by fresh CM- medium (BrainPhys

96.75% by Stemcell Technologies, Vancouver, Canada, B27 2%, Glutamax 0.25%,

penicillin G/streptomycin 1%) and further 0.1 mL of medium were exchanged every 6–

7 days with fresh CM- medium until cells were used.

Condition IV (Hallermann Lab, Leipzig, Germany)

Cultures were generated following the protocol by Ritzau-Jost *et al.*[14]. In short, mice

were decapitated and cerebral cortices were removed, dissected in ice-cold Hank's

balanced saline solution, and cut into 5–10 pieces per hemisphere. Samples were

digested by Trypsin (5 mg, dissolved in digestion solution; for composition see below)

for 5 minutes at 37 °C. Trypsin was stopped by ice-cold MEM growth medium

(composition see below), supernatant was discarded, and samples were mechanically

and enzymatically dissociated using Pasteur pipettes of narrowing tip diameters in

DNAse-supplemented MEM growth medium (10 mg/mL). The supernatant was

centrifuged and the resulting pellet was resuspended in MEM growth medium for

plating onto Matrigel-coated 14 mm coverslips (50.000 vital cells/coverslip). Forty-eight

hours after plating, the medium was fully replaced by MEM growth medium containing

4 µM ARA-C to limit glial growth. After another 48 hours, the medium was fully replaced

by MEM growth medium. Cells were incubated at 37 °C, 93% humidity, and room air

plus 5% CO_2 until use. Digestion solution contained the following (in mM): 137 NaCl,

5 KCl, 7 Na_2HPO_4, 25 HEPES, pH adjusted to 7.2 by NaOH. MEM growth medium was

prepared from 1l MEM (Earle's salts + L-Glutamine) supplemented with 5 g Glucose,

0.2 g $NaHCO_3$, 0.1 g bovine holo-Transferrin, 0.025 g bovine insulin, 50 mL FBS and

10 mL B-27.

Conditions V, VI and VIII (Smith Lab, Portland, OR, USA)

Cultures were generated following the protocol by Martiszus *et al.*[81]. Briefly, 1–2-day

old mice were decapitated following general anesthesia with isoflurane, and their

cerebral cortices (for conditions V and VI) or the frontal third of neocortices (for

condition VIII) were dissected out while bathed in ice-cold modified Hank's balanced

saline solution (modified HBSS; Hyclone laboratories, Logan, UT, USA) and cut into

5–10 pieces per hemisphere. Samples were digested by Trypsin (10 mg, dissolved in

digestion solution as mentioned in Condition IV) for 5 minutes at 37 °C. Trypsin was

de-activated by ice-cold HBSS with 20% FBS, supernatant was discarded, and

samples were enzymatically and mechanically dissociated using Pasteur pipettes with

narrowing tip diameters in DNAse-supplemented HBSS. The supernatant was

centrifuged and the resulting pellet was resuspended in 5% MEM growth medium (see

composition above in Condition IV) for plating 65.000 vital cells onto Matrigel-coated

12 mm glass coverslips. MEM growth medium containing 4 µM ARA-C was added 48

hours later to limit glial proliferation, and subsequently replaced with MEM growth

medium following another 48 hours. Cells were incubated at 37 °C, 93% humidity, and

room air plus 5% CO_2 until use between days 11–14 *in vitro.*

Preparation of dissociated hippocampal cultures

Condition IX and X (Heine Lab, Mainz, Germany)

In short, mice pups were decapitated and brains were removed. Hippocampi were

dissected in ice-cold Hank's balanced saline solution and digested by trypsin for 15

minutes at 37 °C. Next, the tissue was washed with cold HBSS and mechanically

dissociated using Pasteur pipettes in DNAse-, glutamate- and FBS-supplemented

DMEM growth medium. 100 µL of cell suspension containing approximately 70.000

cells was placed on a glass coverslip and incubated at 37 °C, 5% CO_2 for 1 hour.

Coverslips were then transferred to 12-well plates with glutamate-, B27- and sodium

pyruvate-supplemented Neurobasal A medium (1 mL per well) and cultivated until use.

Double distilled water was added in between the wells of the 12-well plate to prevent

evaporation of the medium during cultivation.

Condition XI (Hallermann Lab, Leipzig, Germany)

Cultures were generated as described for Condition III, except that hippocampi from

postnatal day 0–1 Sprague-Dawley rats were used for preparation. 25.000–30.000 vital

cells were plated per coverslip.

Recordings from dissociated cultures

Within conditions, recordings were performed blinded (except conditions IV and VI)

and interleaved from randomized sister coverslips of the same culture, at the same

recording setup, and by the same experimenter.

Conditions I – IV, VII and XI (Hallermann Lab, Leipzig, Germany)

Voltage- and current-clamp recordings were performed with borosilicate pipettes using

a HEKA EPC10/2 amplifier (HEKA Elektronik, Lambrecht/Pfalz, Germany).

Borosilicate glass tubes (Science Products, Hofheim, Germany) were pulled by a DMZ

Universal Electrode Puller with filament heating (Zeitz Instruments, Martinsried,

Germany) to 3–6.5 MΩ pipette resistance. Pipettes were filled with internal solution

(see *Solutions and reagents* section) and mounted on a micromanipulator (Kleindiek

Nanotechnik, Reutlingen, Germany) using a custom-built pipette holder. Recordings

were performed at room temperature (22–25 °C). All current-clamp recordings were

filtered with the internal 10 kHz 8-pole Bessel filter of the HEKA EPC10/2 amplifier and

subsequently digitized (200 kHz) with the HEKA EPC10/2 using Patchmaster software

(HEKA Elektronik, Lambrecht/Pfalz, Germany). Built-in compensation was used for

voltage offset and pipette capacitance. Membrane potentials were not corrected for

liquid junction potential and bridge compensation was adjusted to minimize the current

injection artifact. APs were evoked by brief 1 ms depolarizing current injections of 500–

1500 pA or 500 ms hyperpolarizing pulses (300–1000 pA) leading to APs upon return

to resting membrane potential ('rebound APs'). Both, depolarization-evoked and

rebound APs were first recorded at resting membrane potential without holding current.

While depolarization readily evoked APs in all conditions, rebound AP did not occur in

all conditions at their resting membrane potential. In these cells, holding currents were

used to depolarize cells until rebound APs were observed upon long hyperpolarizing

pulses. Spontaneous excitatory postsynaptic currents were recorded in voltage-clamp

mode at -70 mV membrane potential in neurons grown in conditions I and II. Currents

were filtered with the internal 3 kHz 8-pole Bessel filter of the HEKA EPC10/2 amplifier,

digitized (200 kHz) and detected using the template matching algorithm [82] implemented

in NeuroMatic (Version 2.00, 15.09.2008; ref. [83]).

Conditions V, VI and VIII (Smith Lab, Portland, OR, USA)

Current-clamp recordings were performed with borosilicate pipettes using a HEKA

EPC10 amplifier (HEKA Elektronik, Lambrecht/Pfalz, Germany). Patch pipettes (5–10

MΩ resistance) were pulled from thin-walled borosilicate glass tubes (Warner

Instruments, Holliston, MA, USA) using a P-87 electrode puller (Sutter Instruments,

Novato, CA, USA). Pipettes were filled with an internal solution (see *Solutions and*

reagents section) and mounted on a micromanipulator (ROE-200, Sutter Instruments,

Novato, CA, USA). Recordings were performed at room temperature (22–25 °C).

Voltage signals were filtered at 2.9 kHz and digitized at 20 kHz. Membrane potentials

were corrected for liquid junction potential and bridge balance was performed manually

prior to data acquisition. All APs were evoked by a sequence of transient injections

relative to the resting membrane potential and measurements taken from the APs

elicited with the minimal stimulus using either depolarizing current injections (500–

1000 pA in 100 pA increments for 1–3 ms in 0.5 ms increments in conditions V and VI

or 20–140 pA in 20 pA increments for 0.5 s in condition VIII) or following rebound from

longer hyperpolarizing pulses (300–1200 pA in 100 pA increments for 500 ms).

Conditions IX & X (Heine Lab, Mainz, Germany)

Voltage- and current-clamp recordings were performed with borosilicate pipettes using

a HEKA EPC9 amplifier (HEKA Elektronik, Lambrecht/Pfalz, Germany). Filamented

glass tubes (Science Products, Hofheim, Germany) were pulled by a Narishige PC-

100 electrode puller. Resistance of the pipettes was 2–6 MΩ. Pipettes were filled with

internal solution (see *Solutions and reagents* section) and mounted on a

micromanipulator (Luigs & Neumann, Ratingen, Germany). Recordings were

performed at room temperature (21–24 °C). Signals were filtered at 2.9 kHz and

digitized at 20 kHz using Patchmaster software (HEKA Elektronik, Lambrecht/Pfalz,

Germany). Built-in compensation was used for voltage offset and pipette capacitance.

Membrane potentials were not corrected for liquid junction potential. APs were evoked

by brief 1 ms depolarizing current injection of 50–300 pA or 500 ms hyperpolarizing

pulses (300–400 pA) leading to APs upon return to resting membrane potential

('rebound APs').

Preparation of organotypic hippocampal cultures

Cultures were prepared as described previously[84]. In brief, young Wistar rats (P5–P7)

were anesthetized with isoflurane and killed by decapitation, the brain was removed,

and each hippocampus was dissected. Hippocampal slices (350 µm) were obtained

using a Vibratome (Leica, VT1200S, Wetzlar, Germany). They were placed on 20 mm

latex membranes (Millicell; Merck, Darmstadt, Germany) inserted into 35 mm Petri

dishes containing 1 mL of culture medium and maintained for up to 8 days in an

incubator at 34 °C, 95% O_2, 5% CO_2. The culture medium contained 25 mL MEM, 12.5

mL HBSS, 12.5 mL horse serum, 0.5 mL penicillin/streptomycin, 0.8 mL glucose (1 M),

0.1 mL ascorbic acid (1 mg/mL), 0.4 mL HEPES (1 M), 0.5 mL B27, and 8.95 mL sterile

H_2O. To limit glial proliferation, 5 µM ARA-C was added to the culture medium starting

at DIV 4.

AP recordings from organotypic hippocampal cultures

Whole-cell recordings were obtained from CA3 neurons in organotypic cultures at DIV

7 to DIV 10. All recordings were made at 34 °C in a temperature-controlled recording

chamber (Luigs & Neumann, Ratingen, Germany) using 7–10 MΩ patch pipettes. CA3

pyramidal neurons were recorded in current-clamp with a Multiclamp 700B Amplifier

(Molecular Devices, San Jose, CA, USA). APs were elicited by delivering DC step

pulses (duration 1.0 s) at an interstimulus interval of 10–15 s. During the whole

recording, ionotropic glutamate and $GABA_A$ receptors were blocked with 2–4 mM

kynurenic acid and 100 µM picrotoxin, respectively.

Analysis of APs

Data were analyzed using IGOR PRO software (WaveMetrics, Lake Oswego, OR,

USA; Version 6.32A or 6.37), the Patcher's Power Tool (Version 2.19 Carbon,

15.03.2011) and NeuroMatic (Version 2.00, 15.09.2008; ref. [83]) extensions, as well as

self-written IGOR PRO scripts. AP durations were quantified as the duration at half-

maximal AP amplitude, amplitude being the difference between AP peak and the

membrane potential preceding stimulation. When AP durations were recorded at half-

maximal amplitude between AP threshold (defined as the membrane potential at which

the voltage rate of change first reaches 10 V/s) and AP peak, AP durations were still

unaffected by TTX treatment (data not shown). To exclude the effect of mis-balanced

bridge compensation on AP shape, only APs that occurred after the end of the second

current injection artifact (elicited by the end of current-injection) were analyzed. In

some recordings, current injection artifacts were blanked over 50 µs at the beginning

and the end of the current injection and only experiments in which blanking did not

affect AP amplitude and duration were subsequently analyzed.

Hippocampal dissociated cultures and transfection for GEVI imaging

Culture preparation and GEVI transfection were performed as previously described by

Sabater *et al.*[28]. In short, hippocampi were dissected from embryonic day 17.5 Wistar

rat pups of either sex, treated with trypsin (Worthington Biochemical Corporation,

Lakewood, NJ, USA) at 0.5 mg/mL, and mechanically dissociated using fire polished

Pasteur pipettes. Neurons were plated on 18 mm glass coverslips pre-treated with 100

µg/mL poly-L-lysine and coated with 10 µg/mL laminin (Thermo Fisher Scientific,

Waltham, MA, USA). Cultures were maintained in Neurobasal medium with B27 (1x)

and GlutaMAX (1x, all obtained from Thermo Fisher Scientific, Waltham, MA, USA),

supplemented with FBS (2%, Biosera, Nuaille, France) and Penicillin/Streptomycin

(1%, Sigma-Aldrich, St. Louis, MO, USA), at 37°C in a humidified incubator with 5%

CO_2. For transfections with Ace2N-mNeon-4AA (Ace-mNeon, ref. [85]; distributed by

Biolife Solutions Inc., Bothell, WA, USA), the Effectene transfection reagent (Qiagen,

Venlo, Netherlands) was used. The medium was changed at DIV 3 to culture medium

without antibiotic or FBS, and transfections with Effectene were performed at DIV 7

following the manufacturer's protocol. After transfection neurons were maintained in

serum-free media without antibiotics. For all experiments 30% of the medium was

changed weekly and neurons were imaged 7–14 days after transfection (14–21 DIV).

GEVI Imaging conditions

GEVI imaging was performed as previously described by Sabater *et al.*[28]. In short,

neurons were imaged using an inverted Olympus IX71 epifluorescence microscope

with a 60x 1.42 NA oil-immersion objective (Olympus, Tokyo, Japan). Coverslips were

mounted in a heated chamber (Warner instruments, Holliston, MA, USA; total volume

~500 µL) and placed on an IMTP microscope stage (Scientifica, Uckfield, United

Kingdom). Cells were maintained in external HEPES-buffered saline solution (HBS: 2

mM $CaCl_2$, 1.6 mM $MgCl_2$, 1.45 mM NaCl, 2.5 mM KCl, 10 mM Glucose, 10 mM

HEPES, pH 7.4, Osmolarity 290 mOsm) and experiments were carried out at near

physiological temperatures of 32–35 ºC.

In Ace-mNeon imaging experiments, ~505 nm excitation illumination was provided

using a 525 nm LED (Solis LED, Thorlabs, Newton, NJ, USA), a 500/20 nm excitation

filter, and a 510 nm long-pass dichroic mirror. Ace-mNeon emission transmitted

through the dichroic was filtered using a 520 nm long-pass filters (Chroma, Bellows

Falls, VT, USA). A power density of 10 mW/mm^2 was obtained at the specimen plane.

Images were acquired at 3.2 kHz with an ORCA-Flash4.0 V2 C11440-22CU scientific

CMOS camera (Hamamatsu Photonics, Hamamatsu, Japan) cooled to ~ -20 ºC with

the Exos2 water cooling system (Koolance, Auburn, WA, USA). Images were acquired

with HCImage software (Hamamatsu, Hamamatsu, Japan), binned to 4 × 4 and

cropped to a 16 × 512-pixel region of interest, necessary to achieve the high image

acquisition rates. Images were saved in CXD format.

Stimulation of APs was achieved with an extracellular tungsten parallel bipolar

electrode (FHC, Bowdoin, ME, USA) mounted on a PatchStar motorized micro-

manipulator (Scientifica, Uckfield, United Kingdom). To reliably induce an AP, we

delivered a 1 ms 10 mV pulse at the soma. The timing of the LED illumination, the

acquisition and stimulation were triggered externally through Clampex software

(pClamp 10, Molecular Devices, San Jose, CA, USA).

All GEVI Imaging recordings were performed in presence of NBQX (10 µM), Gabazine

(10 µM) and D-2-amino-5-phosphonovalerate (APV, 25 µM; all from Bio-Techne,

Wiesbaden-Nordenstadt, Germany) in order to block synaptic transmission and ensure

that the observed events were due to the stimulation only. IbTx was delivered using a

custom gravity-fed perfusion system and bath volume was maintained by a peristaltic

pump (Watson-Marlow 120s; Watson-Marlow, Rommerskirchen, Germany).

Image analysis

Image analysis was performed as previously described by Sabater *et al.*[28]. In short,

voltage imaging recordings were analyzed in ImageJ and custom scripts within

MATLAB software as follows. Prior to analysis, if drift had occurred in the x and y axis

during the recording, the images were aligned using the TurboReg ImageJ plugin[86].

Images were then imported into MATLAB and fluorescence intensity profiles were

obtained from regions of interest (ROIs) manually drawn around neuronal structures

identified visually on a maximum projection image of the time series by averaging

across the ROI pixels in each frame. The fluorescence profile was separated into single

trials and corrected for background camera noise by subtracting the average intensity

value of the timepoints where no illumination light was applied. Then, recordings were

reconstructed at a sampling rate of 100 kHz using cubic spline interpolation. Bleaching

was estimated by fitting a single exponential function to each individual trace smoothed

using an averaging filter with a window of 3 and interpolated. The resulting curve was

used to correct an unfiltered interpolated version of the recording. An accurate

representation of the AP waveform with an enhanced SNR was obtained by averaging

over the repeats and extracting the AP parameters from the resulting trace. The

fluorescence profile during the 20 ms preceding a response was used as baseline.

Solutions and reagents

Reagents were purchased from Sigma-Aldrich (St. Louis, MO, USA) if not indicated

otherwise.

Conditions I, II, VII and XI

The extracellular bath solution contained (in mM): 122 NaCl, 3 KCl, 10 D-Glucose, 26

$NaHCO_3$, 1.25 Na_2HPO_4, 1.3 $MgCl_2$, and 2.0 $CaCl_2$. After equilibration with 95% O_2–

5% CO_2, pH was adjusted to 7.35 if needed by NaOH or HCl. The intracellular pipette

solution contained (in mM): 130 K gluconate, 1 $MgCl_2$, 10 HEPES, 0.3 EGTA, 10 Tris-

Phosphocreatine, 4 Mg-ATP, 0.3 Na-GTP. When testing the contribution of BK

channels, 300 nM Iberiotoxin (Abcam, Cambridge, United Kingdom) was included in

the bath solution.

Conditions III and IV

The extracellular bath solution contained (in mM): 125 NaCl, 3 KCl, 25 Glucose, 25

NaHCO$_3$, 1.25 Na$_2$HPO$_4$, 1.1 MgCl$_2$, and 1.1 CaCl$_2$. After equilibration with 95% O$_2$–

5% CO$_2$, pH was adjusted to 7.35 if needed by NaOH or HCl. The intracellular pipette

solution contained (in mM): 150 K Gluconate, 10 K-HEPES, 3 Mg-ATP, 0.3 Na-GTP,

0.05 EGTA, 10 NaCl.

Conditions V and VI

The extracellular bath solution contained (in mM): 150 NaCl, 4 KCl, 10 HEPES, 10

glucose, 1.0 MgCl$_2$, and 2.0 CaCl$_2$, and was equilibrated with 95% O$_2$–5% CO$_2$. The

intracellular pipette solution contained (in mM): 135 K gluconate, 4 NaCl, 0.05 EGTA,

10 HEPES, 3 MgCl$_2$, 0.3 NaGTP, 3 Na$_2$ATP, 10 Na$_2$-Creatine phosphate with pH

normalized to 7.25 using KOH.

Condition VIII

The extracellular bath solution contained (in mM): 122 NaCl, 3 KCl, 1.25 NaH$_2$PO$_4$, 26

NaHCO$_3$, 10 glucose, 1.3 MgCl$_2$, and 2.0 CaCl$_2$, and was bubbled with 95% O$_2$–5%

CO$_2$. The intracellular pipette solution contained (in mM): 130 K gluconate, 0.3 EGTA,

10 HEPES, 2 MgCl$_2$, 0.3 NaGTP, 4 Na$_2$ATP (or Mg-ATP), 10 Na$_2$-Creatine phosphate

(or Tris-Creatinine phosphate).

Conditions IX and X

The extracellular bath solution contained (in mM): 145 NaCl, 2.5 KCl, 10 HEPES, 24

glucose, 1.3 MgCl$_2$, and 2.0 CaCl$_2$, and was equilibrated with 95% O$_2$–5% CO$_2$. The

intracellular pipette solution contained (in mM): 140 K gluconate, 2 NaCl, 0.5 EGTA,

20 HEPES, 3 MgCl$_2$, 0.3 NaGTP, 4 Na$_2$ATP, with pH adjusted to 7.25 using KOH.

Organotypic hippocampal neuron recordings

The external bath solution contained (in mM): 125 NaCl, 26 NaHCO$_3$, 3 CaCl$_2$, 2.5 KCl,

2 MgCl$_2$, 0.8 NaH$_2$PO$_4$, and 10 D-glucose, and was equilibrated with 95% O$_2$–5% CO$_2$.

The intracellular pipette solution contained (mM): 120 K gluconate, 20 KCl, 10 HEPES,

0.5 EGTA, 2 MgCl$_2$, 2 Na$_2$ATP, and 0.3 NaGTP (pH 7.4).

Statistics

If not otherwise noted, data are expressed as box plots with boxes reflecting 25^{th}–75^{th}

percentile. The median is indicated by a line within the box, whiskers indicate 10^{th} and

90^{th} percentile. Sample sizes (n) provide the number of individual somatic recordings.

All data sets were obtained from ≥ 2 independent cell cultures. Data in Fig. 3A were

tested for statistical difference using Fisher's exact test. Confidence limits for 95%

confidence intervals were calculated as Wilson interval[87]. Other experimental groups

were independent and tested by the Mann-Whitney *U* test. Tests were performed in

IGOR PRO or GraphPad Prism 8. Results were considered significant at $p < 0.05$.

Acknowledgment

This work was supported by a European Research Council Consolidator Grant (ERC

CoG 865634) the German Research Foundation (HA6386/10-2) to S.H. and by grants

awarded by the U.S. Department of Veterans Affairs (BX002547) and NIGMS

(GM134110) to S.M.S.

Declaration of interests

The authors declare no competing interests.

References

1. Borst, J.G.G., and Sakmann, B. (1999). Effect of changes in action potential shape on calcium currents and transmitter release in a calyx-type synapse of the rat auditory brainstem. Philosophical Transactions of the Royal Society B: Biological Sciences *354*, 347–355. https://doi.org/10.1098/RSTB.1999.0386.

2. Christie, B.R., Eliot, L.S., Ito, K.I., Miyakawa, H., and Johnston, D. (1995). Different Ca2+ channels in soma and dendrites of hippocampal pyramidal neurons mediate spike-induced Ca2+ influx. J. Neurophysiol. *73*, 2553–2557. https://doi.org/10.1152/jn.1995.73.6.2553.

3. Watanabe, S., Hoffman, D.A., Migliore, M., and Johnston, D. (2002). Dendritic K+ channels contribute to spike-timing dependent long-term potentiation in hippocampal pyramidal neurons. Proc. Natl. Acad. Sci. U. S. A. *99*, 8366–8371. https://doi.org/10.1073/PNAS.122210599.

4. Bean, B.P. (2007). The action potential in mammalian central neurons. Nat. Rev. Neurosci. *8*, 451–465. https://doi.org/10.1038/nrn2148.

5. Debanne, D., Campanac, E., Bialowas, A., Carlier, E., and Alcaraz, G. (2011). Axon physiology. Physiol. Rev. *91*, 555–602. https://doi.org/10.1152/PHYSREV.00048.2009.

6. Geiger, J.R.P., and Jonas, P. (2000). Dynamic control of presynaptic Ca2+ inflow by fast-inactivating K+ channels in hippocampal mossy fiber boutons. Neuron *28*, 927–939. https://doi.org/10.1016/S0896-6273(00)00164-1.

7. Jackson, M.B., Konnerth, A., and Augustine, G.J. (1991). Action potential broadening and frequency-dependent facilitation of calcium signals in pituitary nerve terminals. Proc. Natl. Acad. Sci. U. S. A. *88*, 380–384. https://doi.org/10.1073/PNAS.88.2.380.

8. Keck, T., Toyoizumi, T., Chen, L., Doiron, B., Feldman, D.E., Fox, K., Gerstner, W., Haydon, P.G., Hübener, M., Lee, H.K., et al. (2017). Integrating Hebbian and homeostatic plasticity: The current state of the field and future research directions. Philosophical Transactions of the Royal Society B: Biological Sciences *372*. https://doi.org/10.1098/rstb.2016.0158.

9. Carta, M., Lanore, F., Rebola, N., Szabo, Z., Da Silva, S.V., Lourenço, J., Verraes, A., Nadler, A., Schultz, C., Blanchet, C., et al. (2014). Membrane lipids tune synaptic transmission by direct modulation of presynaptic potassium channels. Neuron *81*, 787–799. https://doi.org/10.1016/J.NEURON.2013.12.028.

10. Senn, V., Wolff, S.B.E., Herry, C., Grenier, F.O., Ehrlich, I., Grü, J., Fadok, J.P., Müller, C., Letzkus, J.J., and Lüthi, A. (2014). Long-range connectivity defines behavioral specificity of amygdala neurons. Neuron *81*, 428–437. https://doi.org/10.1016/j.neuron.2013.11.006.

11. Turrigiano, G.G., and Nelson, S.B. (2004). Homeostatic plasticity in the developing nervous system. Nat. Rev. Neurosci. *5*, 97–107. https://doi.org/10.1038/nrn1327.

12. Pozo, K., and Goda, Y. (2010). Unraveling mechanisms of homeostatic synaptic plasticity. Neuron *66*, 337–351. https://doi.org/10.1016/J.NEURON.2010.04.028.

13. Li, B., Suutari, B.S., Sun, S.D., Luo, Z., Wei, C., Chenouard, N., Mandelberg, N.J., Zhang, G., Wamsley, B., Tian, G., et al. (2020). Neuronal inactivity co-opts LTP machinery to drive potassium channel splicing and homeostatic spike widening. Cell *181*, 1547-1565.e15. https://doi.org/10.1016/j.cell.2020.05.013.

14. Ritzau-Jost, A., Tsintsadze, T., Krueger, M., Ader, J., Bechmann, I., Eilers, J., Barbour, B., Smith, S.M., and Hallermann, S. (2021). Large, stable spikes

exhibit differential broadening in excitatory and inhibitory neocortical boutons.
Cell Rep. *34*, 108612. https://doi.org/10.1016/j.celrep.2020.108612.

15. Zbili, M., Rama, S., Benitez, M.J., Fronzaroli-Molinieres, L., Bialowas, A.,
Boumedine-Guignon, N., Garrido, J.J., and Debanne, D. (2021). Homeostatic
regulation of axonal Kv1.1 channels accounts for both synaptic and intrinsic
modifications in the hippocampal CA3 circuit. Proc. Natl. Acad. Sci. U. S. A.
118. https://doi.org/10.1073/pnas.2110601118.

16. Kim, J., and Tsien, R.W. (2008). Synapse-specific adaptations to inactivity in
hippocampal circuits achieve homeostatic gain control while dampening
network reverberation. Neuron *58*, 925–937.
https://doi.org/10.1016/j.neuron.2008.05.009.

17. Mitra, A., Mitra, S.S., and Tsien, R.W. (2012). Heterogeneous reallocation of
presynaptic efficacy in recurrent excitatory circuits adapting to inactivity. Nat.
Neurosci. *15*, 250–257. https://doi.org/10.1038/nn.3004.

18. Castro-Alamancos, M.A., and Connors, B.W. (1997). Distinct forms of short-
term plasticity at excitatory synapses of hippocampus and neocortex. Proc.
Natl. Acad. Sci. U. S. A. *94*, 4161–4166.
https://doi.org/10.1073/pnas.94.8.4161.

19. Manahan-Vaughan, D. (2018). Special considerations when using mice for In
vivo electrophysiology and long-term studies of hippocampal synaptic plasticity
during behavior. Handb. Behav. Neurosci. *28*, 63–84.
https://doi.org/10.1016/B978-0-12-812028-6.00003-3.

20. Nguyen, P. V., Duffy, S.N., and Young, J.Z. (2000). Differential maintenance
and frequency-dependent tuning of LTP at hippocampal synapses of specific
strains of inbred mice. J. Neurophysiol. *84*, 2484–2493.
https://doi.org/10.1152/jn.2000.84.5.2484.

21. Prestigio, C., Ferrante, D., Valente, P., Casagrande, S., Albanesi, E.,
Yanagawa, Y., Benfenati, F., and Baldelli, P. (2019). Spike-Related
Electrophysiological Identification of Cultured Hippocampal Excitatory and
Inhibitory Neurons. Mol. Neurobiol. *56*, 6276–6292.
https://doi.org/10.1007/s12035-019-1506-5.

22. Maffei, A., and Turrigiano, G.G. (2008). Multiple modes of network homeostasis
in visual cortical layer 2/3. Journal of Neuroscience *28*, 4377–4384.
https://doi.org/10.1523/JNEUROSCI.5298-07.2008.

23. Desai, N.S., Rutherford, L.C., and Turrigiano, G.G. (1999). Plasticity in the
intrinsic excitability of cortical pyramidal neurons. Nat. Neurosci. *2*, 515–520.
https://doi.org/10.1038/9165.

24. Turrigiano, G.G., Leslie, K.R., Desai, N.S., Rutherford, L.C., and Nelson, S.B.
(1998). Activity-dependent scaling of quantal amplitude in neocortical neurons.
Nature *391*, 892–896. https://doi.org/10.1038/36103.

25. Hanes, A.L., Koesters, A.G., Fong, M.-F., Altimimi, H.F., Stellwagen, D.,
Wenner, P., and Engisch, K.L. (2020). Cellular/molecular divergent synaptic
scaling of miniature EPSCs following activity blockade in dissociated neuronal
cultures. *40*, 4090–4102. https://doi.org/10.1523/JNEUROSCI.1393-19.2020.

26. Frank, C.A., Kennedy, M.J., Goold, C.P.P., Marek, K.W., and Davis, G.W.W.
(2006). Mechanisms underlying the rapid induction and sustained expression of
synaptic homeostasis. Neuron *52*, 663–677.
https://doi.org/10.1016/J.NEURON.2006.09.029.

27. Aoto, J., Nam, C.I., Poon, M.M., Ting, P., and Chen, L. (2008). Synaptic
Signaling by All-Trans Retinoic Acid in Homeostatic Synaptic Plasticity. Neuron
60, 308–320. https://doi.org/10.1016/j.neuron.2008.08.012.

28.   Gonzalez Sabater, V., Rigby, M., and Burrone, J. (2021). Voltage-gated
842   potassium channels ensure action potential shape fidelity in distal axons. The
843   Journal of Neuroscience *41*, 5372–5385.
844   https://doi.org/10.1523/jneurosci.2765-20.2021.
845   29.   Hu, H., Shao, L.-R., Chavoshy, S., Gu, N., Trieb, M., Behrens, R., Laake, P.,
846   Pongs, O., Knaus, H.G., Ottersen, O.P., et al. (2001). Presynaptic Ca2+-
847   activated K+ channels in glutamatergic hippocampal terminals and their role in
848   spike repolarization and regulation of transmitter release. The Journal of
849   Neuroscience *21*, 9585–9597. https://doi.org/10.1523/JNEUROSCI.21-24-
850   09585.2001.
851   30.   Lorenzo-Ceballos, Y., Carrasquel-Ursulaez, W., Castillo, K., Alvarez, O., and
852   Latorre, R. (2019). Calcium-driven regulation of voltagesensingdomains in BK
853   channels. Elife *8*, 1–24. https://doi.org/10.7554/eLife.44934.
854   31.   Karmarkar, U.R., and Buonomano, D. V. (2006). Different forms of homeostatic
855   plasticity are engaged with distinct temporal profiles. European Journal of
856   Neuroscience *23*, 1575–1584. https://doi.org/10.1111/j.1460-
857   9568.2006.04692.x.
858   32.   Horrigan, F.T., and Aldrich, R.W. (2002). Coupling between voltage sensor
859   activation, Ca2+ binding and channel opening in large conductance (BK)
860   potassium channels. Journal of General Physiology *120*, 267–305.
861   https://doi.org/10.1085/jgp.20028605.
862   33.   Lee, K.Y., and Chung, H.J. (2014). NMDA receptors and L-type voltage-gated
863   Ca2+ channels mediate the expression of bidirectional homeostatic intrinsic
864   plasticity in cultured hippocampal neurons. Neuroscience *277*, 610–623.
865   https://doi.org/10.1016/j.neuroscience.2014.07.038.
866   34.   Lee, K.Y., Royston, S.E., Vest, M.O., Ley, D.J., Lee, S., Bolton, E.C., and
867   Chung, H.J. (2015). N-methyl-D-aspartate receptors mediate activity-
868   dependent down-regulation of potassium channel genes during the expression
869   of homeostatic intrinsic plasticity. Mol. Brain *8*, 1–16.
870   https://doi.org/10.1186/s13041-015-0094-1.
871   35.   Zhao, C.J., Dreosti, E., and Lagnado, L. (2011). Homeostatic synaptic plasticity
872   through changes in presynaptic calcium influx. Journal of Neuroscience *31*,
873   7492–7496. https://doi.org/10.1523/JNEUROSCI.6636-10.2011.
874   36.   Desai, N.S., Rutherford, L.C., and Turrigiano, G.G. (1999). BDNF regulates the
875   intrinsic excitability of cortical neurons. Learning and Memory *6*, 284–291.
876   https://doi.org/10.1101/lm.6.3.284.
877   37.   Rutherford, L.C., DeWan, A., Lauer, H.M., and Turrigiano, G.G. (1997). Brain-
878   derived neurotrophic factor mediates the activity-dependent regulation of
879   inhibition in neocortical cultures. Journal of Neuroscience *17*, 4527–4535.
880   https://doi.org/10.1523/jneurosci.17-12-04527.1997.
881   38.   Ritzau-Jost, A., Nerlich, J., Kaas, T., Krueger, M., Tsintsadze, T., Eilers, J.,
882   Barbour, B., Smith, S.M., and Hallermann, S. (2023). Direct whole-cell patch-
883   clamp recordings from small boutons in rodent primary neocortical neuron
884   cultures. STAR Protoc. *4*, 102168. https://doi.org/10.1016/j.xpro.2023.102168.
885   39.   Ivenshitz, M., and Segal, M. (2010). Neuronal density determines network
886   connectivity and spontaneous activity in cultured hippocampus. J.
887   Neurophysiol. *104*, 1052–1060. https://doi.org/10.1152/jn.00914.2009.
888   40.   Mattson, M.P., and Kater, S.B. (1989). Development and selective
889   neurodegeneration in cell cultures from different hippocampal regions. Brain
890   Res. *490*, 110–125. https://doi.org/https://doi.org/10.1016/0006-
891   8993(89)90436-8.

41. Chipman, P.H., Fetter, R.D., Panzera, L.C., Bergerson, S.J., Karmelic, D.,
Yokoyama, S., Hoppa, M.B., and Davis, G.W. (2022). NMDAR-dependent
presynaptic homeostasis in adult hippocampus: Synapse growth and cross-
modal inhibitory plasticity. Neuron *110*, ki.
https://doi.org/10.1016/j.neuron.2022.08.014.
42. Vinet, J., van Weering, H.R.J., Heinrich, A., Kälin, R.E., Wegner, A., Brouwer,
N., Heppner, F.L., van Rooijen, N., Boddeke, H.W.G.M., and Biber, K. (2012).
Neuroprotective function for ramified microglia in hippocampal excitotoxicity. J.
Neuroinflammation *9*, 1–15. https://doi.org/10.1186/1742-2094-9-27.
43. Niday, Z., and Bean, B.P. (2021). BK channel regulation of afterpotentials and
burst firing in cerebellar purkinje neurons. Journal of Neuroscience *41*, 2854–
2869. https://doi.org/10.1523/JNEUROSCI.0192-20.2021.
44. Gittis, A.H., Moghadam, S.H., and Du Lac, S. (2010). Mechanisms of sustained
high firing rates in two classes of vestibular nucleus neurons: Differential
contributions of resurgent Na, Kv3, and BK currents. J. Neurophysiol. *104*,
1625–1634. https://doi.org/10.1152/jn.00378.2010.
45. Kimm, T., Khaliq, Z.M., and Bean, B.P. (2015). Differential regulation of action
potential shape and burst-frequency firing by BK and Kv2 channels in
substantia nigra dopaminergic neurons. Journal of Neuroscience *35*, 16404–
16417. https://doi.org/10.1523/JNEUROSCI.5291-14.2015.
46. Müller, A., Kukley, M., Uebachs, M., Beck, H., and Dietrich, D. (2007).
Nanodomains of single Ca2+ channels contribute to action potential
repolarization in cortical neurons. Journal of Neuroscience *27*, 483–495.
https://doi.org/10.1523/JNEUROSCI.3816-06.2007.
47. Bock, T., and Stuart, G.J. (2016). The impact of BK channels on cellular
excitability depends on their subcellular location. Front. Cell. Neurosci. *10*, 1–8.
https://doi.org/10.3389/fncel.2016.00206.
48. Desai, N.S., Cudmore, R.H., Nelson, S.B., and Turrigiano, G.G. (2002). Critical
periods for experience-dependent synaptic scaling in visual cortex. Nat.
Neurosci. *5*, 783–789. https://doi.org/10.1038/nn878.
49. Wiesel, T.N., and Hubel, D.H. (1963). Single-cell responses in striate cortex of
kittens deprived of vision in one eye. J. Neurophysiol. *26*, 1003–1017.
50. Wen, W., and Turrigiano, G.G. (2021). Developmental regulation of
homeostatic plasticity in mouse primary visual cortex. The Journal of
Neuroscience *41*, JN-RM-1200-21. https://doi.org/10.1523/jneurosci.1200-
21.2021.
51. Trojanowski, N.F., and Turrigiano, G.G. (2021). CaMKIV signaling is not
essential for the maintenance of intrinsic or synaptic properties in mouse visual
cortex. eNeuro *8*, 1–12. https://doi.org/10.1523/ENEURO.0135-21.2021.
52. Griffen, T.C., Haley, M.S., Fontanini, A., and Maffei, A. (2017). Rapid plasticity
of visually evoked responses in rat monocular visual cortex. PLoS One *12*, 1–
12. https://doi.org/10.1371/journal.pone.0184618.
53. Gainey, M.A., Aman, J.W., and Feldman, D.E. (2018). Rapid disinhibition by
adjustment of PV intrinsic excitability during whisker map plasticity in mouse
S1. Journal of Neuroscience *38*, 4749–4761.
https://doi.org/10.1523/JNEUROSCI.3628-17.2018.
54. Maravall, M., Stern, E.A., and Svoboda, K. (2004). Development of intrinsic
properties and excitability of layer 2/3 pyramidal neurons during a critical period
for sensory maps in rat barrel cortex. J. Neurophysiol. *92*, 144–156.
https://doi.org/10.1152/jn.00598.2003.

55.   Breton, J.D., and Stuart, G.J. (2009). Loss of sensory input increases the
intrinsic excitability of layer 5 pyramidal neurons in rat barrel cortex. J. Physiol.
587, 5107–5119. https://doi.org/10.1113/JPHYSIOL.2009.180943.

56.   Sun, Q.Q. (2009). Experience-dependent intrinsic plasticity in interneurons of
barrel cortex layer IV. J. Neurophysiol. *102*, 2955–2973.
https://doi.org/10.1152/jn.00562.2009.

57.   Li, L., Gainey, M.A., Goldbeck, J.E., and Feldman, D.E. (2014). Rapid
homeostasis by disinhibition during whisker map plasticity. Proc. Natl. Acad.
Sci. U. S. A. *111*, 1616–1621. https://doi.org/10.1073/pnas.1312455111.

58.   Jamann, N., Dannehl, D., Lehmann, N., Wagener, R., Thielemann, C., Schultz,
C., Staiger, J., Kole, M.H.P., and Engelhardt, M. (2021). Sensory input drives
rapid homeostatic scaling of the axon initial segment in mouse barrel cortex.
Nat. Commun. *12*, 1–14. https://doi.org/10.1038/s41467-020-20232-x.

59.   Hickmott, P.W. (2005). Changes in intrinsic properties of pyramidal neurons in
adult rat S1 during cortical reorganization. J. Neurophysiol. *94*, 501–511.
https://doi.org/10.1152/jn.00924.2004.

60.   Francis, H.W., and Manis, P.B. (2000). Effects of deafferentation on the
electrophysiology of ventral cochlear nucleus neurons. Hear. Res. *149*, 91–
105. https://doi.org/10.1016/S0378-5955(00)00165-9.

61.   Lu, Y., Monsivais, P., Tempel, B.L., and Rubel, E.W. (2004). Activity-dependent
regulation of the potassium channel subunits Kv1.1 and Kv3.1. Journal of
Comparative Neurology *470*, 93–106. https://doi.org/10.1002/cne.11037.

62.   Wang, Y., and Manis, P.B. (2006). Temporal coding by cochlear nucleus bushy
cells in DBA/2J mice with early onset hearing loss. JARO - Journal of the
Association for Research in Otolaryngology *7*, 412–424.
https://doi.org/10.1007/s10162-006-0052-9.

63.   Leao, R.N., Berntson, A., Forsythe, I.D., and Walmsley, B. (2004). Reduced
low-voltage activated K+ conductances and enhanced central excitability in a
congenitally deaf (dn/dn) mouse. J. Physiol. *559*, 25–33.
https://doi.org/10.1113/JPHYSIOL.2004.067421.

64.   Him, A., and Dutia, M.B. (2001). Intrinsic excitability changes in vestibular
nucleus neurons after unilateral deafferentation. Brain Res. *908*, 58–66.
https://doi.org/10.1016/S0006-8993(01)02600-2.

65.   George, N.M., Gentile Polese, A., Merle, L., Macklin, W.B., and Restrepo, D.
(2022). Excitable axonal domains adapt to sensory deprivation in the olfactory
system. The Journal of Neuroscience *42*, 1491–1509.
https://doi.org/10.1523/JNEUROSCI.0305-21.2021.

66.   Barnes, B.Y.C.A., and McNaughton, B.L. (1980). Physiological compensation
for loss of afferent synapses in rat hippocampal granule cells during
senescence. J. Physiol *309*, 473–485.
https://doi.org/10.1113/jphysiol.1980.sp013521ch.

67.   Midorikawa, M., and Miyata, M. (2021). Distinct functional developments of
surviving and eliminated presynaptic terminals. Proc. Natl. Acad. Sci. U. S. A.
118. https://doi.org/10.1073/pnas.2022423118.

68.   O'Leary, T., van Rossum, M.C.W., and Wyllie, D.J.A. (2010). Homeostasis of
intrinsic excitability in hippocampal neurones: Dynamics and mechanism of the
response to chronic depolarization. Journal of Physiology *588*, 157–170.
https://doi.org/10.1113/jphysiol.2009.181024.

69.   Pozzi, D., Lignani, G., Ferrea, E., Contestabile, A., Paonessa, F.,
D'Alessandro, R., Lippiello, P., Boido, D., Fassio, A., Meldolesi, J., et al.
(2013). REST/NRSF-mediated intrinsic homeostasis protects neuronal

networks from hyperexcitability. EMBO Journal *32*, 2994–3007.
https://doi.org/10.1038/emboj.2013.231.

70.   Wefelmeyer, W., Cattaert, D., and Burrone, J. (2015). Activity-dependent
mismatch between axo-axonic synapses and the axon initial segment controls
neuronal output. Proc. Natl. Acad. Sci. U. S. A. *112*, 9757–9762.
https://doi.org/10.1073/pnas.1502902112.

71.   Malik, R., and Chattarji, S. (2012). Enhanced intrinsic excitability and EPSP-
spike coupling accompany enriched environment-induced facilitation of LTP in
hippocampal CA1 pyramidal neurons. J. Neurophysiol. *107*, 1366–1378.
https://doi.org/10.1152/jn.01009.2011.

72.   Valero-Aracama, M.J., Sauvage, M.M., and Yoshida, M. (2015). Environmental
enrichment modulates intrinsic cellular excitability of hippocampal CA1
pyramidal cells in a housing duration and anatomical location-dependent
manner. Behavioural Brain Research *292*, 209–218.
https://doi.org/10.1016/j.bbr.2015.05.032.

73.   Kole, M.H.P., Czéh, B., and Fuchs, E. (2004). Homeostatic maintenance in
excitability of tree shrew hippocampal CA3 pyramidal neurons after chronic
stress. Hippocampus *14*, 742–751. https://doi.org/10.1002/hipo.10212.

74.   Kamal, A., Notenboom, R.G.E., De Graan, P.N.E., and Ramakers, G.M.J.
(2006). Persistent changes in action potential broadening and the slow
afterhyperpolarization in rat CA1 pyramidal cells after febrile seizures.
European Journal of Neuroscience *23*, 2230–2234.
https://doi.org/10.1111/j.1460-9568.2006.04732.x.

75.   Eshra, A., Hirrlinger, P., and Hallermann, S. (2019). Enriched environment
shortens the duration of action potentials in cerebellar granule cells. Front. Cell.
Neurosci. *13*, 1–9. https://doi.org/10.3389/fncel.2019.00289.

76.   Ngodup, T., Goetza, J.A., McGuire, B.C., Sun, W., Lauer, A.M., and Xu-
Friedman, M.A. (2015). Activity-dependent, homeostatic regulation of
neurotransmitter release from auditory nerve fibers. Proc. Natl. Acad. Sci. U. S.
A. *112*, 6479–6484. https://doi.org/10.1073/PNAS.1420885112/-
/DCSUPPLEMENTAL/PNAS.201420885SI.PDF.

77.   Simkin, D., Hattori, S., Ybarra, N., Musia, T.F., Buss, E.W., Richter, H.,
Matthew Oh, M., Nicholson, D.A., and Disterhoft, J.F. (2015). Aging-related
hyperexcitability in CA3 pyramidal neurons is mediated by enhanced A-type K+
channel function and expression. Journal of Neuroscience *35*, 13206–13218.
https://doi.org/10.1523/JNEUROSCI.0193-15.2015.

78.   Mehranfard, N., Gholamipour-Badie, H., Motamedi, F., Janahmadi, M., and
Naderi, N. (2015). Long-term increases in BK potassium channel underlie
increased action potential firing in dentate granule neurons following
pilocarpine-induced status epilepticus in rats. Neurosci. Lett. *585*, 88–91.
https://doi.org/10.1016/j.neulet.2014.11.041.

79.   Raffaelli, G., Saviane, C., Mohajerani, M.H., Pedarzani, P., and Cherubini, E.
(2004). BK potassium channels control transmitter release at CA3-CA3
synapses in the rat hippocampus. Journal of Physiology *557*, 147–157.
https://doi.org/10.1113/jphysiol.2004.062661.

80.   Moutin, E., Hemonnot, A.L., Seube, V., Linck, N., Rassendren, F., Perroy, J.,
and Compan, V. (2020). Procedures for culturing and genetically manipulating
murine hippocampal postnatal neurons. Front. Synaptic Neurosci. *12*, 1–16.
https://doi.org/10.3389/fnsyn.2020.00019.

81.   Martiszus, B.J., Tsintsadze, T., Chang, W., and Smith, S.M. (2021). Enhanced
excitability of cortical neurons in low-divalent solutions is primarily mediated by

altered voltage-dependence of voltage-gated sodium channels. Elife *10*, 1–26.
https://doi.org/10.7554/eLife.67914.

82. Clements, J.D., and Bekkers, J.M. (1997). Detection of spontaneous synaptic events with an optimally scaled template. Biophys. J. *73*, 220–229.
https://doi.org/10.1016/S0006-3495(97)78062-7.

83. Rothman, J.S., and Silver, R.A. (2018). Neuromatic: An integrated open-source software toolkit for acquisition, analysis and simulation of electrophysiological data. Front. Neuroinform. *12*, 1–21. https://doi.org/10.3389/fninf.2018.00014.

84. Debanne, D., Boudkkazi, S., Campanac, E., Cudmore, R.H., Giraud, P., Fronzaroli-Molinieres, L., Carlier, E., and Caillard, O. (2008). Paired-recordings from synaptically coupled cortical and hippocampal neurons in acute and cultured brain slices. Nat. Protoc. *3*, 1559–1568.
https://doi.org/10.1038/nprot.2008.147.

85. Gong, Y., Huang, C., Li, J.Z., Grewe, B.F., Zhang, Y., Eismann, S., and Schnitzer, M.J. (2015). High-speed recording of neural spikes in awake mice and flies with a fluorescent voltage sensor. Science (1979). *350*, 1361–1366.
https://doi.org/10.1126/science.aab0810.

86. Thevenaz, P., Ruttimann, U.E., and Unser, M. (1998). A pyramid approach to subpixel registration based on intensity. IEEE Transactions on Image Processing *7*, 27–41. https://doi.org/10.1109/83.650848.

87. Brown, L.D., Cai, T.T., and DasGupta, A. (2001). Interval estimation for a binomial proportion. Statistical Science *16*, 101–133.

1068 **Figures and legends**

1069

1070

1071

1072

Figure 1 Cell type- and model-specific AP broadening in hippocampal neurons

(A) Example evoked APs recorded from CA3 pyramidal neurons in organotypic

hippocampal cultures under control condition (black) and following 48 hours of

TTX-treatment (blue).

(B) AP durations recorded under conditions in A.

(C) Example depolarization-evoked APs (500 pA for 1 ms) recorded from neurons in

primary dissociated hippocampal cultures under control condition (black) and

following 48 hours of TTX- (blue), Kyn- (purple), or NBQX-treatment (turquoise).

(D) AP durations recorded under conditions in C.

(E) Example spontaneous APs recorded under the conditions in C.

(F) Spontaneous AP frequency under the conditions in C.

Numbers in brackets reflect number of recorded neurons. Box plots as median ±

interquartile range, whiskers indicate 10–90 percentile. P-values above graphs,

significance tested by Mann-Whitney-*U* test.

Figure 2 Stable AP duration in primary dissociated neurons during TTX-induced homeostatic plasticity

(A) Example current injection-evoked APs (500 pA for 1 ms) under control condition (black) and following 48 hours of TTX-treatment (blue). AP amplitudes are recorded between the membrane potential preceding the AP and AP peak (arrows), AP durations are recorded at half-maximal amplitude (arrow heads).

(B) AP duration in control and TTX-treated cells (color code as in A) for eleven different experimental conditions (I–XI). Specification of individual conditions is provided in Table S1. Condition XI adopted from Fig. 1D.

(C) Merge of AP durations in control and TTX-treated cells across conditions I–IX normalized to the corresponding condition's control median duration.

(D) AP amplitudes in control and TTX-treated cells as in B.

(E) Cross-conditional merge of AP amplitudes in control and TTX-treated cells as in C.

(F) Example hyperpolarization-evoked "rebound APs" (-150 pA or -60 pA for 500ms) in control and TTX-treated cells (color code as in A).

(G) Rebound AP duration in control and TTX-treated cells as in B.

1105   (H) Cross-conditional merge of normalized durations as in C for rebound APs.

1106   Numbers in brackets reflect number of recorded neurons. Box plots as median ±

1107   interquartile range, whiskers indicate 10–90 percentile. P-values provided above

1108   graphs; significance tested using Mann-Whitney-*U* tests.

Figure 3 Homeostatic plasticity increases network activity and excitatory synaptic transmission

(A) *Left:* Spontaneously active control (black) and TTX-treated cells (blue) in condition I–IV and VII. *Right*: Active neurons merged across conditions I–IV and VII.

(B) Example spontaneous AP bursts in control and TTX-treated cells. Asterisks mark individual APs.

(C) AP number per burst in control and TTX-treated cells across conditions I–IV and VII.

(D) Example spontaneous excitatory postsynaptic currents (sEPSCs) in control and TTX-treated cells.

(E) *Left:* sEPSC frequency and *Right:* sEPSC amplitude in control recordings and after TTX-treatment (merged data from conditions I and II).

1124   Numbers in brackets reflect number of recorded neurons. Bars in A reflect fraction
1125   of spontaneously active neurons. Whiskers in A reflect confidence limits for 95%
1126   confidence interval based on Wilson interval. Box plots as median ± interquartile
1127   range, whiskers indicate 10–90 percentile. P-values above graphs, significance
1128   tested by Fisher's exact test in A or Mann-Whitney-U test elsewhere.

Figure 4 Lack of BK channel-dependent AP broadening in dissociated cultured neurons

(A) *Left:* Example current injection-evoked APs (500 pA for 1 ms) in control solution (black) and solution containing 300 nM Iberiotoxin (IbTx, green). *Right:* AP duration in control and TTX-treated cells recorded IbTx-free (black and blue, respectively) or IbTx-containing solution (green). Data recorded under condition I.

(B) Merged AP duration across control and TTX-treated cells for recordings in IbTx-free and IbTx-containing solution (data in A normalized to the respective group's median duration in control solution).

(C) APs evoked by a 200 ms current injection (top) and overlay of the forst three evoked APs (bottom).

(D) Broadening of AP duration of the 3 first APs during 200 ms current injections in control (black, *left*) and TTX-treated cells (blue, *right*). Bars as mean ± SEM, lines with markers depict recordings from individual neurons.

(E) *Left:* Example image of neuron expressing the genetically encoded voltage indicator Ace-mNeon (scale bar 50 μm). *Middle:* Example Ace-mNeon-recorded somatic, proximal axonal, and distal axonal APs in a primary hippocampal cultured neuron in IbTx-free solution (black) and following BK channel block by IbTx (100

nM, green). *Right:* Ratio of somatic and axonal AP durations in IbTx-free and IbTx-

containing solution.

(F) *Left:* Example AP train (5 x 20 Hz) and overlay of 1^{st} and 5^{th} train AP at the soma,

proximal, and distal axon in control condition (grey and black) and following BK

channel block by IbTx (yellow and green). *Right:* AP broadening (ratio of 5^{th} over

1^{st} train AP duration) in the three compartments in control and IbTx-containing

solution.

(G)Somatic APs under control condition (grey), following treatment with 4-AP (black),

and following treatment with 30µM 4-AP + 100 nM IbTx (green).

Numbers in brackets reflect number of recorded neurons. Box plots as median ±

interquartile range, whiskers indicate 10–90 percentile. Plots in D–F as mean ± SEM.

P-values above graphs, significance tested by Mann-Whitney-*U* test.

Figure supplements

Group	Species	Strain	Brain area	Culture age (days *in vitro*)	Lab which recordings were performed at	External recording solution	External [Ca²⁺] (mM)	Internal recording solution	Growth medium	Protocol Reference
I	Rat	Sprague-Dawley	Cortex	10–14	Hallermann Lab, Leipzig (Germany)	1.)	2.0	5.)	NbActiv4	Li *et al.*, Cell 2020
II	Rat	Sprague-Dawley	Cortex	16–21	Hallermann Lab, Leipzig (Germany)	1.)	2.0	5.)	NbActiv4	Li *et al.*, Cell 2020
III	Mouse	C57BL/6N	Cortex	15–20	Hallermann Lab, Leipzig (Germany)	2.)	1.1	6.)	BrainPhys	Moutin *et al.*, Front. Synaptic Neurosci 2020
IV	Mouse	C57BL/6N	Cortex	16	Hallermann Lab, Leipzig (Germany)	2.)	1.1	6.)	MEM, 5% serum + supplements	Ritzau-Jost *et al.*, Cell Reports 2021
V	Mouse	C57BL/6J	Cortex	11–13	Smith Lab, Portland (OR, USA)	3.)	2.0	7.)	MEM, 5% serum + supplements	Martiszus BJ *et al.*, eLife 2021
VI	Mouse	CD1	Cortex	11–13	Smith Lab, Portland (OR, USA)	3.)	2.0	7.)	MEM, 5% serum + supplements	Martiszus BJ *et al.*, eLife 2021
VII	Rat	Sprague-Dawley	Pre-frontal cortex	10–14	Hallermann Lab, Leipzig (Germany)	1.)	2.0	5.)	BrainPhys	Moutin *et al.*, Front. Synaptic Neurosci 2020
VIII	Mouse	C57BL/6J	Pre-frontal cortex	10–14	Smith Lab, Portland (OR, USA)	1.)	2.0	8.)	MEM, 5% serum + supplements	Martiszus BJ *et al.*, eLife 2021
IX	Mouse	C57BL/6N	Hippo-campus	32	Heine Lab, Mainz (Germany)	4.)	2.0	9.)	Neurobasal A+ supplements	Li *et al.*, Cell 2020
X	Mouse	C57BL/6J	Hippo-campus	15	Heine Lab, Mainz (Germany)	4.)	2.0	9.)	Neurobasal A + supplements	Li *et al.*, Cell 2020
XI	Rat	Sprague-Dawley	Hippo-campus	16–20	Hallermann Lab, Leipzig (Germany)	1.)	2.0	5.)	BrainPhys	Moutin *et al.*, Front. Synaptic Neurosci 2020

1.) 122 NaCl, 3 KCl, 10 Glucose, 26 $NaHCO_3$, 2 $CaCl_2$, 1.3 $MgCl_2$, 1.25 NaH_2PO_4
2.) 125 NaCl, 3 KCl, 25 Glucose, 25 $NaHCO_3$, 1.1 $CaCl_2$, 1.1 $MgCl_2$, 1.25 Na_2HPO_4
3.) 150 NaCl, 4 KCl, 10 Glucose, 10 HEPES, 2 $CaCl_2$, 1 $MgCl_2$
4.) 145 NaCl, 2.5 KCl, 24 Glucose, 10 HEPES, 2 $CaCl_2$, 1.3 $MgCl_2$
5.) 130 K gluconate, 10 HEPES, 1 $MgCl_2$, 0.3 EGTA, 3 Mg-ATP, 0.3 Na-GTP, 10 Tris-Phosphocreatine
6.) 150 K gluconate, 10 K-HEPES, 10 NaCl, 0.05 EGTA, 3 Mg-ATP, 0.3 Na-GTP
7.) 135 K gluconate, 10 HEPES, 4 NaCl, 0.05 EGTA, 3 Na_2ATP, 0.3 NaGTP, 3 $MgCl_2$, 10 Na_2-Creatine phosphate
8.) 130 K gluconate, 10 HEPES, 0.3 EGTA, 4 Na_2ATP (or Mg-ATP), 0.3 NaGTP, 2 $MgCl_2$, 10 Na_2-Creatine phosphate (or Tris-Creatinine phosphate)
9.) 140 K gluconate, 20 HEPES, 2 NaCl, 0.5 EGTA, 4 Na_2ATP, 0.3 NaGTP, 3 $MgCl_2$

1161

1162

1163 **Figure 2 – figure supplement 1**

1164 **Additional information on recording conditions**

1165 The table lists additional relevant information on the experimental conditions I – XI.

1166

1167

1168

Figure 2 – figure supplement 2

Absence of TTX-induced AP broadening in further selected conditions

(A) Box plots of AP durations under control condition (black) or upon TTX-treatment

(blue) recorded in condition I either unselected or selected for parameters

reflecting unimpaired neuronal function.

(B) AP durations as in A for recordings in condition VII.

(C) AP durations as in A for recordings in condition VIII.

Numbers of recorded neurons provided in brackets. Box plots depict median ±

interquartile range, whiskers reflect 10–90 percentile. P-values calculated by Mann-

Whitney-*U* test.

1179

1180

1181

Figure 4 – figure supplement 1

BK channel block does not affect rebound AP duration

(A) Example rebound APs in control solution (black) and solution containing 300 nM

Iberiotoxin (IbTx, green).

(B) AP duration in control and TTX-treated cells in IbTx-free (black and blue,

respectively) or IbTx-containing solution (green). Data recorded in condition I.

(C) Merged AP duration across control and TTX-treated cells for recordings in IbTx-

free and IbTx-containing solution (data in A, normalized to the respective group's

median duration in IbTx-free solution).

Numbers of recorded neurons provided in brackets. Box plots depict median ±

interquartile range, whiskers reflect 10–90 percentile. P-values calculated by Mann-

Whitney-*U* test.