Unreliable homeostatic action potential broadening in cultured dissociated neurons

Andreas Ritzau-Jost[1,#], Salil Rajayer[2,3], Jana Nerlich[1], Filip Maciag[4], Alexandra John[1], Michael Russier[5], Victoria Gonzalez Sabater[6,7], Luke J. Steiger[2,3], Jacques-Olivier Coq[5], Jens Eilers[1], Maren Engelhardt[8,9], Juan Burrone[6,7], Dominique Debanne[5], Martin Heine[4], Stephen M Smith[2,3,#], Stefan Hallermann[1,#]

[1] Carl-Ludwig-Institute of Physiology, Faculty of Medicine, Leipzig University, Leipzig, 04103, Germany
[2] Division of Pulmonary and Critical Care Medicine, Department of Medicine, Oregon Health and Science University, Portland, Oregon, 97239-3098, USA
[3] Section of Pulmonary and Critical Care Medicine, VA Portland Health Care System, Portland, Oregon, 97239-2964, USA
[4] Institute for Developmental Biology and Neurobiology, Johannes Gutenberg University, Mainz, 55128, Germany
[5] Unité de Neurobiologie des canaux Ioniques et de la Synapse (UNIS), UMR_S 1072, INSERM, Aix-Marseille Université, Marseille, 13015, France
[6] MRC Centre for Neurodevelopmental Disorders, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, SE1 1UL, United Kingdom
[7] Centre for Developmental Neurobiology, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, SE1 1UL, United Kingdom
[8] Institute of Neuroanatomy, Mannheim Center for Translational Neuroscience, Medical Faculty Mannheim, Heidelberg University, Mannheim, 68167, Germany
[9] Institute of Anatomy and Cell Biology, Johannes Kepler University, Linz, 4020, Austria
[#] Correspondence: andreas.ritzau-jost@medizin.uni-leipzig.de (A.R-J.), smisteph@ohsu.edu (S.M.S.), and hallermann@medizin.uni-leipzig.de (S.H.)

Summary

Homeostatic plasticity preserves neuronal activity against perturbations. Recently, somatic action potential broadening was proposed as a key homeostatic adaptation to chronic inactivity in neocortical neurons. Since action potential shape critically controls calcium entry and neuronal function, broadening provides an attractive homeostatic feedback mechanism to regulate activity. Here, we report that chronic inactivity induced by sodium channel block does not broaden action potentials in neocortical neurons under a wide range of conditions. In contrast, action potentials were broadened in CA3 neurons of organotypic hippocampal cultures by chronic sodium channel block and in hippocampal dissociated cultures by chronic synaptic block. Mechanistically, BK-type potassium channels were proposed to underly inactivity-induced action potential broadening. However, BK channels did not affect action potential duration in our recordings. Our results indicate that action potential broadening can occur in specific neurons and conditions but is not a general mechanism of homeostatic plasticity in cultured neurons.

Introduction

Action potentials (APs) control various functions including neurotransmitter release from axonal boutons[1], calcium signals in the soma[2], and synaptic plasticity in dendrites[3]. AP broadening during short-term plasticity on the time scale of milliseconds to seconds is well-established and occurs in many types of neuronal cell bodies[4,5] and nerve terminals[6,7]. There are also indications that APs change during long-term plasticity on the time scale of minutes to hours. When examining Hebbian long-term plasticity (i.e., positive feedback-based and input-specific long-term plasticity[8]), some studies reported AP broadening. For example, presynaptic APs of hippocampal mossy fiber boutons broaden during depolarization-induced potentiation[9] and somatic APs of neurons in the amygdala broaden during fear conditioning[10]. In contrast, the role of the AP duration is less clear for non-Hebbian homeostatic long-term plasticity (i.e., negative feedback-based and non-input specific long-term plasticity[8]). Specifically, homeostatic plasticity induced by neuronal inactivity in the presence of the sodium channel blocker tetrodotoxin (TTX) has been studied for decades, but AP broadening was initially not described[11,12]. A recent study however reports prominent AP broadening during TTX-induced homeostatic plasticity at the soma of cultured neocortical neurons[13], which was not observed at presynaptic boutons of cultured neocortical neurons[14]. Compartmentalized AP broadening at the neuronal cell body would provide a strikingly simple and previously overlooked mechanism by which neuronal inactivity could increase subsequent excitability[13]. This mechanism involves decreased calcium entry during TTX application causing changes in gene expression that would result in broadened APs and more calcium entry after TTX removal. In this model, AP broadening is a key link in the homeostatic feedback loop underlying homeostatic plasticity. In a collaborative effort across several laboratories, we therefore systematically investigated AP broadening in cultured neurons during homeostatic plasticity under various experimental conditions.

Results

Cell type- and model-specific AP broadening in hippocampal neurons

APs in CA3 neurons of hippocampal organotypic slice cultures broaden homeostatically upon chronic inactivity using both, synaptic block by glutamate receptor antagonists[15] or sodium channel block by TTX[16] (see also ref. [17] for a similar

trend). We first aimed to confirm homeostatic AP broadening in CA3 neurons in

organotypic slice cultures. APs were elicited by depolarizing currents from the resting

membrane potential during perfusion with TTX-free artificial cerebrospinal fluid. APs

recorded in CA3 pyramidal neurons treated with TTX (48 hours) had longer durations

than those in control cells (Fig. 1A, B; median [IQR]: 1.75 [1.69–1.86] ms and 2.13

[2.07–2.22] ms for control and TTX-treatment, respectively, 14 cells each, P < 0.001),

confirming earlier reports of homeostatic AP broadening in these neurons[15,16].

Next, we tested whether homeostatic AP broadening occurred in dissociated

hippocampal cultured neurons, a model widely used to study homeostatic plasticity.

We inhibited neuronal activity for 48 hours to induce homeostatic plasticity by either

blocking AP firing using TTX (Fig. 1C) or by blocking synaptic transmission using the

low-affinity glutamate receptor blocker kynurenic acid (Kyn) or the high-affinity

glutamate receptor blocker 2,3-Dioxo-6-nitro-1,2,3,4-tetrahydrobenzo [f]quinoxaline-

7-sulfonamide (NBQX). Surprisingly, AP duration was unaffected by TTX-treatment in

primary hippocampal cultured neurons (Fig. 1D; median [IQR]: 2.07 [1.79–2.34] ms

and 2.08 [1.32–2.61] ms for control and TTX, 25 and 24 cells, respectively, P = 0.96).

In contrast, Kyn-treatment significantly increased AP duration in the same set of

interleaved experiments (Fig. 1D; 2.52 [2.31–2.72] ms, 22 cells, P = 0.014) and

NBQX-treatment led to a small trend towards broader APs (Fig. 1D; P = 0.26).

Importantly, sodium channel block and synaptic block both increased spontaneous

AP firing frequency, indicating that homeostatic adaptations other than AP

broadening were shared between both induction methods (Fig. 1E, F; P = 0.008 for

control against Kyn, P = 0.005 for control against NBQX, P = 0.013 for control against

TTX). These data indicate that AP broadening occurred upon sodium channel block

in CA3 neurons in organotypic slice cultures and upon synaptic block in dissociated

hippocampal cultures, but not in the commonly-studied model of homeostatic

plasticity using dissociated hippocampal cultured neurons and 48 hours of TTX

application.

Stable AP duration in primary dissociated neurons during TTX-induced

homeostatic plasticity

In contrast to our finding in dissociated cultured neurons of the hippocampus, a

recent study described TTX-induced AP broadening in dissociated cultured neurons

of the neocortex[13]. To resolve this discrepancy, we set out to replicate AP broadening

in neocortical cultured neurons. Given that exact reproduction of experimental

conditions is challenging and neuronal plasticity depends on brain region[18], animal

species[19], and animal strain[20], we probed homeostatic AP broadening under different

experimental conditions varying: 1) experimental animal species (rat and mouse), 2)

strain (C57BL/6 and CD1 mice), 3) sub-strain (C57BL/6N and C57BL/6J mice), 4)

brain area (cortex and hippocampus), 5) cortical subdomain (prefrontal cortex), 6)

culture duration (10–14, 15–21, and 32 days *in vitro*), 7) culture growth medium, 8)

recording solutions, and 9) laboratory and experimenter performing recordings

(conditions I–X in Fig. 2B, condition XI are recordings from Fig. 1; see also Fig. 2 –

figure supplement 1). Even though AP durations varied up to 2-fold between

conditions, homeostatic AP broadening was not detectable in any of the tested

conditions (Fig. 2B). To detect cross-conditional AP broadening by TTX, we merged

all conditions based on their difference in AP duration after TTX-treatment (Fig. 2C).

However, AP duration was the same in control and TTX-treated cells (217 and 197

cells, respectively, $P = 0.83$).

We next tested whether AP broadening was concealed by systematic differences in

neuronal health or recording quality between control and treatment group. We first

analyzed AP amplitude as a measure of neuronal viability. AP amplitudes were not

affected following TTX treatment in any of the eleven recording conditions (Fig. 2D)

or a cross-conditional comparison (Fig. 2E). Next, neurons grown using a culture

protocol or cortical subdomain previously associated with AP broadening were further

analyzed to avoid the inclusion of inhibitory or depolarized neurons (ref. 13;

conditions I, VII, and VIII). Inclusion criteria were the occurrence of multiple APs upon

200 ms-current injections (≥ 2 APs), a resting membrane potential below – 55 mV,

and a maximum AP depolarization rate between 100 and 337 V/s (includes 96% of

excitatory and exclude >80% of the inhibitory neurons; ref. 21). Applying any or all

three of these criteria did not reveal AP broadening upon TTX treatment (Fig. 2 –

figure supplement 2). These data indicate that stricter exclusion criteria result in

smaller variability in action potential duration but do not uncover AP broadening in

these datasets.

Instead of depolarization-evoked APs, Li *et al.*[13] recorded hyperpolarization-evoked

'rebound APs' (Fig. 2F), possibly reducing the impact of different resting membrane

potentials and depolarizing current injection on the AP waveform. We determined that

the duration of rebound APs was similarly unaffected by TTX treatment across

several conditions (Fig. 2G, H). These data show that TTX-induced AP broadening

does not robustly occur in dissociated cultured neurons.

Homeostatic plasticity increases network activity and excitatory synaptic

transmission

To verify successful induction of homeostatic plasticity under our experimental

conditions, we quantified changes in neuronal activity and synaptic strength following

TTX-induced homeostatic plasticity. Spontaneously active neurons (Fig. 3A; merge of

conditions I–IV and VII; control: 23 out of 87 cells, TTX-treatment: 58 out of 74 cells,

P < 0.001 with Fisher's exact test) and the number of APs per spontaneous burst

(Fig. 3B, C; median [IQR]: 1.0 [1.0–1.12] APs and 1.67 [1.0–2.71] APs for control and

for TTX-treatment, 23 and 58 cells, respectively, P < 0.001) were increased after

TTX-treatment. These results indicate an adaptive increase in neuronal activity

following TTX-treatment, in line with previous findings on homeostatic plasticity[16,22,23].

Furthermore, spontaneous excitatory postsynaptic currents (sEPSCs; i.e.,

spontaneous synaptic currents recorded in the absence of TTX) were recorded as a

measure of synaptic strength (Fig. 3D). sEPSC frequency and amplitude increased

upon activity block by TTX (Fig. 2E, merge of recordings from condition I and II;

median [IQR] frequency: 1.11 [0.42–1.47] Hz and 5.07 [1.95–5.52] Hz in control and

TTX-treated cells, 16 and 11, respectively, P = 0.027; median [IQR] amplitude: 12.4

[10.6–14.4] pA and 15.5 [14.1–17.4] pA in control and TTX-treated cells, 16 and 11

cells, respectively, P = 0.042), in accord with known homeostatic synaptic

adaptations to inactivity[24–27]. Taken together, blocking AP firing with TTX successfully

induced homeostatic plasticity of excitability and synaptic strength under our

experimental conditions.

Lack of BK channel-dependent AP broadening in dissociated cultured neurons

Various potassium channel subtypes regulate AP repolarization and could therefore

underlie AP broadening. Li *et al.*[13] reported that downregulation of BK-type potassium

channels led to homeostatic AP broadening in neocortical cultured neurons. We

tested the role of BK channels in AP repolarization (neurons grown under condition I).

However, blocking BK channels with Iberiotoxin (IbTx, 300 nM) did not lead to

broadening of depolarization-evoked APs in control and TTX-treated cells (Fig. 4A; P

= 0.10 and 0.48, respectively), or a merge of both conditions (Fig. 4B; P = 0.10).

Rebound APs were similarly unaffected by BK channel block (Fig. 4 – figure

supplement 1). Next, we tested whether recurrent neuronal activity, which increases

cytosolic calcium and boosts BK-channel activity, enhanced AP broadening following

TTX treatment.   While APs significantly broadened during repetitive firing, AP

durations were similar in control and TTX-treated neurons (Fig. 4C and D),

suggesting no differential regulation of BK-channels following TTX-treatment in these

recordings.

In electrical recordings, intracellular recording solution may have disturbed cytosolic

calcium dynamics and therefore BK- and thus calcium-dependent AP repolarization.

As an independent approach, we recorded the AP duration in hippocampal cultured

neurons using the genetically encoded voltage indicator (GEVI) Ace-mNeon[28] before

and after acute IbTx perfusion (100 nM). BK channel block did not affect AP duration

in the soma or along the axon (Fig. 4E). Using the GEVI, we again tested the role of

BK channels in AP broadening during a train of APs (5 APs at 20 Hz, Fig. 4F). The

broadening of somatic, proximal axonal, and distal axonal APs during trains was

similar before and after BK channel block, indicating that AP duration was not

modulated by BK channels. To confirm that IbTx was indeed pharmacologically

active in Ace-mNeon recordings, we preconditioned cells with 4-AP (30 µM) prior to

IbTx perfusion (Fig. 4G). Neurons preconditioned with 4-AP had broader somatic APs

which were further broadened by IbTx, presumably due to increased calcium influx

following 4-AP treatment, in turn activating more BK channels[29]. These data indicate

that IbTx application was pharmacologically active but BK channels did not contribute

to AP repolarization in cultured neocortical neurons.

Discussion

Homeostatic AP broadening has been reported in some systems, but it has not been

tested systematically across experimental models so far. We found that AP

broadening did not occur following the induction of homeostatic plasticity with TTX in

the most commonly studied model system (primary dissociated neocortical neurons).

However, AP broadening was observed in cultured hippocampal slices. Furthermore,

blocking BK channels, which were reported to mediate AP broadening during

homeostatic plasticity[13], did not impact AP duration in dissociated cultured neurons.

Our results therefore indicate that AP broadening in cultured dissociated neurons is

not a required component of homeostatic plasticity.

TTX-induced homeostatic AP regulation in organotypic hippocampal cultured

neurons

AP broadening was previously shown to occur in CA3 neurons of organotypic

hippocampal slice cultures following TTX-induced homeostatic plasticity[16].

Consistently, we recorded broader APs in CA3 neurons following neuronal inactivity

(Fig. 1). In contrast to CA3 neurons, granule cells of the dentate gyrus did not show

TTX-induced AP broadening[17]. Similarly, CA1 neurons did not show AP broadening

following synaptic block-induced homeostatic plasticity[30]. Therefore, homeostatic AP

broadening is cell type-specific.

TTX-induced homeostatic AP regulation in primary dissociated cultured

neurons

To the best of our knowledge, only three studies so far reported on the duration of

APs following TTX-induced homeostatic plasticity in dissociated cultured

neurons[13,31,32]. While Li *et al.*[13] observed TTX-induced broadening, the studies by

Lee, Chung, and others[31,32] reported stable AP duration, consistent with our findings

(Fig. 2). Furthermore, in other studies investigating TTX-induced homeostatic

plasticity in dissociated cultured neurons, prominent AP broadening as observed by

Li *et al.*[13] was either missed or absent, as in our recordings (see for example refs.

[23,33–35]). One possible explanation for this inconsistency could be the cell type-

specificity of AP broadening. We focused on large, presumably excitatory neurons.

However, we and others could have selected different neuron types compared to Li

et al.[13]. In an attempt to reduce bias, we incorporated multi-lab, randomization, and

blinding strategies across many of our experiments. Only recordings with leaky seal

formation or inadequate capacitance compensation[36] were excluded. AP amplitudes

were unaffected by TTX treatment (Fig. 2). We then applied additional exclusion

criteria to ensure that variability in neuronal type, health or recording quality were not

causing a type II statistical error. Again, AP amplitudes were unaffected by TTX

treatment after excluding recordings with low resting membrane potential, with peak

AP rates of depolarization outside the range for excitatory neurons, or those that did

not fire repetitively (Fig. 2 – figure supplement 2). Even when these three criteria

were applied together TTX failed to broaden AP duration, confirming that AP

broadening was not associated with homeostatic plasticity in neocortical neurons

following TTX treatment.

Synaptic block-induced homeostatic AP regulation in primary dissociated

cultured neurons

Homeostatic AP broadening was shown in CA3 neurons upon activity block by the

low-affinity glutamate receptor blocker Kyn[15]. We similarly observed Kyn-induced AP

broadening in dissociated cultured hippocampal neurons (Fig. 1C, D). In contrast,

APs were not significantly broader following synaptic block by NBQX (Fig. 1C, D). A

differential effect of the glutamate receptor blockers Kyn and NBQX may relate to

NMDA receptors, which are blocked by Kyn but not NBQX or TTX and which have

recently been demonstrated to be important for the induction of synaptic homeostatic

plasticity[37]. Further studies are needed to investigate the role of subcellular calcium

signals in the regulation of AP duration during block of glutamate receptors.

Absence of BK channel-dependent AP broadening in dissociated cultured

neurons

Li *et al.*[13] proposed that BK channels hasten AP repolarization and that this effect is

lost during TTX-induced homeostatic plasticity, leading to AP broadening. We did not

detect AP broadening when blocking BK channels in either control or TTX-treated

cells (Fig. 4). Furthermore, APs did not broaden differentially during repetitive firing in

control and TTX-treated neurons, suggesting no activity-dependent effect of TTX or

BK channels on AP duration. Voltage imaging using Ace-mNeon similarly showed

that width of single APs and broadening during AP trains in the soma and axon were

unaffected by BK channel block. Blocking BK channels only affected AP

repolarization when cells were preconditioned with 4-AP. These results are

consistent with previous reports showing that in some neurons BK channels do not

contribute to repolarization[38,39] while doing so in others[39–42]. Differences in the

abundance of other types of potassium channels[39] and intracellular calcium

buffering[41] are known to affect the contribution of BK channels to AP repolarization

and may account for the lack of AP broadening upon BK channel block in our dissociated cultured neurons. The lack of BK channel contribution to AP repolarization in our dissociated neurons is another possible explanation (in addition to different selection of cells) for why we and others have not observed the homeostatic AP broadening described by Li *et al.*[13].

Inactivity-induced homeostatic AP regulation *in vivo*

Homeostatic plasticity has also been studied using *in vivo* recordings from the activity-suppressed visual cortex[43,44]. The majority of these studies reported stable AP duration upon monocular deprivation or chemogenetic activity suppression[45–47]. In contrast, Li *et al.*[13] found homeostatically broadened APs in the visual cortex upon monocular deprivation. Furthermore, stable AP duration was also observed in other brain areas following activity suppression such as (1.) the somatosensory cortex upon unilateral whisker trimming[48–53] or forepaw denervation[54], (2.) the auditory system upon cochlear ablation[55,56], hearing loss[57], or congenital deafness[58], (3.) the vestibular nucleus upon vestibular ablation[59], and (4.) the olfactory mitral cells upon naris occlusion[60] or senescent hippocampal deafferentiation[61]. Besides somatic APs, APs in excitatory presynaptic terminals of the somatosensory thalamus were also unaffected by whisker deprivation[62]. Thus, AP broadening does not appear to occur robustly upon suppression of neuronal activity in various neuronal pathways.

Hyperactivity-induced homeostatic AP regulation

Homeostatic plasticity occurs not only upon activity block but also upon hyperactivity, generally involving opposite adaptations. In cultured neurons, AP duration was unaffected following hyperactivity induced by chronic depolarization[63], block of potassium channels[64], block of inhibitory synaptic transmission[30–32], or chronic ChR2-induced photostimulation[65]. In contrast, Li *et al.*[13] reported similar BK channel downregulation upon both, TTX-induced activity block and depolarization-induced hyperactivity.

As in cultured neurons, AP duration was mostly unaffected in brain slice recordings from animals with increased neuronal activity due to an enriched environment[53,66,67], psychosocial stress[68], or febrile seizures[69]. AP shortening occurred in cerebellar granule cells after exposure to an enriched environment[70], in auditory neurons

following chronic noise exposure[71], and in hippocampal neurons following age-
related[72] or pilocarpine-induced hyperactivity[73]. Thus, in contrast to Li *et al.*[13], AP
duration was either unaffected or, if anything, shortened during hyperactivity-induced
homeostatic plasticity in previous studies.

In this study, several laboratories collaborated to determine the robustness of
inactivity-induced AP broadening. AP broadening did not occur upon TTX-treatment
in dissociated hippocampal or neocortical neurons. Furthermore, BK channel block
did not affect AP duration in dissociated neurons, arguing against BK channels as
mediators of homeostatic AP regulation.

METHODS

Ethics

At all times animal procedures were in accordance with the relevant international,
national and institutional guidelines for the care and use of laboratory animals
(European Council Directive 86/609/EEC, U.S. Public Health Service Policy on
Humane Care and Use of Laboratory Animals, U.S. National Institutes of Health
Guide for the Care and Use of Laboratory Animals, German Tierschutzgesetz,
French National Research Council guidelines, and Leipzig and Mainz University
guidelines). All animal procedures were approved in advance by the local health
authorities (Ethics Committees at Leipzig University and Mainz University, federal
Saxonian Animal Welfare Committee, the District administration Mainz-Bingen
(41a/177-5865-§11 ZVTE), the Committee of the Préfecture des Bouches-du-Rhône
(D13055-08), V.A. Portland Health Care System Institutional Animal Care and Use
Committee).

Animal models

Organotypic hippocampal cultures were generated from postnatal day 5–7 Wistar
rats.

Dissociated neocortical cultures were generated from postnatal day 0 or 1 Sprague-
Dawley rats (conditions I, II and VII), C57BL/6N mice (conditions III and IV),
C57BL/6J mice (conditions V and VIII), or CD1 mice (condition VI). Dissociated
hippocampal cultures were generated from postnatal day 0 or 1 C57BL/6N mice
(conditions IX), C57BL/6J mice (condition X), or Sprague-Dawley rats (condition XI).

Generally, animals of both sexes were used to generate cultures. Mother animals
were kept in individually ventilated cages under a 12h/12h light-dark cycle and
received water and food *ad libitum*.

Induction of Homeostatic Plasticity
Homeostatic plasticity was induced by inclusion of one of the respective blockers in
culture medium for 48–56 hours prior to recordings. All blockers were first diluted in
100 µL of fresh medium per treated well and subsequently added to the respective
wells containing 1 mL of medium. For control conditions, only 100 µL of fresh
medium without blockers was added to the respective wells. The blockers had final
concentrations of (in mM): 0.002 tetrodotoxin (TTX; Tocris, Wiesbaden-Nordenstadt,
Germany), 2 kynurenic acid (Kyn), or 0.01 2,3-Dioxo-6-nitro-1,2,3,4-tetrahydrobenzo-
[f]quinoxaline-7-sulfonamide (NBQX). To avoid bias, coverslips were randomized to
treatment (control or blocker) and recordings were made and analyzed in a blinded
fashion (except for condition IV and VI). Blinding was performed by an independent
investigator. Control or TTX-treatment groups were assigned using an 8-block
randomization approach for experiments of condition VIII. Unblinding of the
experimental groups was performed after analysis and disclosure of the data by the
experimenter.

Preparation of dissociated neocortical cultures
Conditions I and II (Hallermann Lab, Leipzig, Germany)
Neocortical cultures were generated as described by Li *et al.*[13]. Shortly, neocortices
were isolated and dissected into 5–10 pieces per hemisphere. Samples were washed
twice in ice-cold modified HBSS (4.2 mM $NaHCO_3$ and 1 mM HEPES, pH 7.35, 300
mOsm) supplemented with 20% fetal bovine serum (FBS) and digested at 37 °C for
30 min using papain. After stopping digestion with 20% FBS-containing modified
HBSS, samples were washed and dissociated using Pasteur pipettes, pelleted twice,
and filtered using a 70 nm nylon strainer. Dissociated cells were plated on poly-D-
lysine-coated 14 mm coverslips and cultured in NbActive4 medium (Transnetyx,
Cordova, TN, USA) in an incubator until use (37 °C, 5% CO_2 in room air, water vapor
saturated).

Conditions III and VII (Hallermann Lab, Leipzig, Germany)

Cultures were generated as previously described by Moutin et al.[74]. In brief, mice
were decapitated and neocortices (for condition III) or the frontal third of neocortices
(for condition VII) were isolated in ice-cold Hibernate-A medium (Thermo Fisher
Scientific, Waltham, MA, USA) and cut into 5–10 chunks. Chunks were first digested
by papain (dissolved in Hibernate-A; 10 min at 37 °C) and subsequently by papain +
DNAse (125 µL of 20 mg/mL DNAse stock; 5 min at 37 °C). Digestion was stopped
by CM+ medium (Neurobasal A 86.5 %, B27 2 %, Glutamax 0.25 %, L-Glutamine
0.25%, penicillin G/streptomycin 1 %, heat-inactivated FBS 10 %) and chunks were
mechanically dissociated in two steps by 1250 µL pipette tips with gravity decantation
and collection of the supernatant between dissociation steps and after dissociation.
Supernatant was passed through 2 mL of frothed bovine serum albumin, centrifuged
(7 min, 300 x g), the resulting pellet was resuspended in 1–2 mL of CM+ medium and
25.000–30.000 resuspended cells were plated onto poly-L-lysine-coated coverslips.
After 30 min, 0.5 mL of CM+ medium was added to the wells and after 2 days in vitro
(DIV), cytosine arabinoside (ARA-C, 1 µM final concentration) was added to the wells
for 8–18 h. Subsequently, 0.4 mL medium per well was replaced by fresh CM-
medium (BrainPhys 96.75% by Stemcell Technologies, Vancouver, Canada, B27 2%,
Glutamax 0.25%, penicillin G/streptomycin 1%) and further 0.1 mL of medium were
exchanged every 6–7 days with fresh CM- medium until cells were used.
403

Condition IV (Hallermann Lab, Leipzig, Germany)
Cultures were generated following the protocol by Ritzau-Jost et al.[14]. In short, mice
were decapitated and cerebral cortices were removed, dissected in ice-cold Hank's
balanced saline solution, and cut into 5–10 pieces per hemisphere. Samples were
digested by Trypsin (5 mg, dissolved in digestion solution; for composition see below)
for 5 minutes at 37 °C. Trypsin was stopped by ice-cold MEM growth medium
(composition see below), supernatant was discarded, and samples were
mechanically and enzymatically dissociated using Pasteur pipettes of narrowing tip
diameters in DNAse-supplemented MEM growth medium (10 mg/mL). The
supernatant was centrifuged and the resulting pellet was resuspended in MEM
growth medium for plating onto Matrigel-coated 14 mm coverslips (50.000 vital
cells/coverslip). Forty-eight hours after plating, the medium was fully replaced by
MEM growth medium containing 4 µM ARA-C to limit glial growth. After another 48
hours, the medium was fully replaced by MEM growth medium. Cells were incubated

at 37 °C, 93% humidity, and room air plus 5% CO_2 until use. Digestion solution

contained the following (in mM): 137 NaCl, 5 KCl, 7 Na_2HPO_4, 25 HEPES, pH

adjusted to 7.2 by NaOH. MEM growth medium was prepared from 1l MEM (Earle's

salts + L-Glutamine) supplemented with 5 g Glucose, 0.2 g $NaHCO_3$, 0.1 g bovine

holo-Transferrin, 0.025 g bovine insulin, 50 mL FBS and 10 mL B-27.

Conditions V, VI and VIII (Smith Lab, Portland, OR, USA)

Cultures were generated following the protocol by Martiszus *et al.*[75]. Briefly, 1–2-day

old mice were decapitated following general anesthesia with isoflurane, and their

cerebral cortices (for conditions V and VI) or the frontal third of neocortices (for

condition VIII) were dissected out while bathed in ice-cold modified Hank's balanced

saline solution (modified HBSS; Hyclone laboratories, Logan, UT, USA) and cut into

5–10 pieces per hemisphere. Samples were digested by Trypsin (10 mg, dissolved in

digestion solution as mentioned in Condition IV) for 5 minutes at 37 °C. Trypsin was

de-activated by ice-cold HBSS with 20% FBS, supernatant was discarded, and

samples were enzymatically and mechanically dissociated using Pasteur pipettes

with narrowing tip diameters in DNAse-supplemented HBSS. The supernatant was

centrifuged and the resulting pellet was resuspended in 5% MEM growth medium

(see composition above in Condition IV) for plating onto Matrigel-coated 12 mm glass

coverslips. MEM growth medium containing 4 µM ARA-C was added 48 hours later

to limit glial proliferation, and subsequently replaced with MEM growth medium

following another 48 hours. Cells were incubated at 37 °C, 93% humidity, and room

air plus 5% CO_2 until use between days 11–14 *in vitro.*

Preparation of dissociated hippocampal cultures

Condition IX and X (Heine Lab, Mainz, Germany)

In short, mice pups were decapitated and brains were removed. Hippocampi were

dissected in ice-cold Hank's balanced saline solution and digested by trypsin for 15

minutes at 37 °C. Next, the tissue was washed with cold HBSS and mechanically

dissociated using Pasteur pipettes in DNAse-, glutamate- and FBS-supplemented

DMEM growth medium. 100 µL of cell suspension containing approximately 70.000

cells was placed on a glass coverslip and incubated at 37 °C, 5% CO_2 for 1 hour.

Coverslips were then transferred to 12-well plates with glutamate-, B27- and sodium

pyruvate-supplemented Neurobasal A medium (1 mL per well) and cultivated until

use. Double distilled water was added in between the wells of the 12-well plate to prevent evaporation of the medium during cultivation.

Condition XI (Hallermann Lab, Leipzig, Germany)

Cultures were generated as described for Condition III, except that hippocampi from postnatal day 0–1 Sprague-Dawley rats were used for preparation.

Recordings from dissociated cultures

Within conditions, recordings were performed blinded (except conditions IV and VI) and interleaved from randomized sister coverslips of the same culture, at the same recording setup, and by the same experimenter.

Conditions I – IV, VII and XI (Hallermann Lab, Leipzig, Germany)

Voltage- and current-clamp recordings were performed with borosilicate pipettes using a HEKA EPC10/2 amplifier (HEKA Elektronik, Lambrecht/Pfalz, Germany). Borosilicate glass tubes (Science Products, Hofheim, Germany) were pulled by a DMZ Universal Electrode Puller with filament heating (Zeitz Instruments, Martinsried, Germany) to 3–6.5 MΩ pipette resistance. Pipettes were filled with internal solution (see *Solutions and reagents* section) and mounted on a micromanipulator (Kleindiek Nanotechnik, Reutlingen, Germany) using a custom-built pipette holder. Recordings were performed at room temperature (22–25 °C). All current-clamp recordings were filtered with the internal 10 kHz 8-pole Bessel filter of the HEKA EPC10/2 amplifier and subsequently digitized (200 kHz) with the HEKA EPC10/2 using Patchmaster software (HEKA Elektronik, Lambrecht/Pfalz, Germany). Built-in compensation was used for voltage offset and pipette capacitance. Membrane potentials were not corrected for liquid junction potential and bridge compensation was adjusted to minimize the current injection artifact. APs were evoked by brief 1 ms depolarizing current injections of 500–1500 pA or 500 ms hyperpolarizing pulses (300–1000 pA) leading to APs upon return to resting membrane potential ('rebound APs'). Both, depolarization-evoked and rebound APs were first recorded at resting membrane potential without holding current. While depolarization readily evoked APs in all conditions, rebound AP did not occur in all conditions at their resting membrane potential. In these cells, holding currents were used to depolarize cells until rebound APs were observed upon long hyperpolarizing pulses. Spontaneous excitatory

postsynaptic currents were recorded in voltage-clamp mode at -70 mV membrane potential in neurons grown in conditions I and II. Currents were filtered with the internal 3 kHz 8-pole Bessel filter of the HEKA EPC10/2 amplifier, digitized (200 kHz) and detected using the template matching algorithm [76] implemented in NeuroMatic (Version 2.00, 15.09.2008; ref. [77]).

Conditions V, VI and VIII (Smith Lab, Portland, OR, USA)

Current-clamp recordings were performed with borosilicate pipettes using a HEKA EPC10 amplifier (HEKA Elektronik, Lambrecht/Pfalz, Germany). Patch pipettes (5–10 MΩ resistance) were pulled from thin-walled borosilicate glass tubes (Warner Instruments, Holliston, MA, USA) using a P-87 electrode puller (Sutter Instruments, Novato, CA, USA). Pipettes were filled with an internal solution (see *Solutions and reagents* section) and mounted on a micromanipulator (ROE-200, Sutter Instruments, Novato, CA, USA). Recordings were performed at room temperature (22–25 °C). Voltage signals were filtered at 2.9 kHz and digitized at 20 kHz. Membrane potentials were corrected for liquid junction potential and bridge balance was performed manually prior to data acquisition. All APs were evoked by a sequence of transient injections relative to the resting membrane potential and measurements taken from the APs elicited with the minimal stimulus using either depolarizing current injections (500–1000 pA in 100 pA increments for 1–3 ms in 0.5 ms increments in conditions V and VI or 20–140 pA in 20 pA increments for 0.5 s in condition VIII) or following rebound from longer hyperpolarizing pulses (300–1200 pA in 100 pA increments for 500 ms).

Conditions IX & X (Heine Lab, Mainz, Germany)

Voltage- and current-clamp recordings were performed with borosilicate pipettes using a HEKA EPC9 amplifier (HEKA Elektronik, Lambrecht/Pfalz, Germany). Filamented glass tubes (Science Products, Hofheim, Germany) were pulled by a Narishige PC-100 electrode puller. Resistance of the pipettes was 2–6 MΩ. Pipettes were filled with internal solution (see *Solutions and reagents* section) and mounted on a micromanipulator (Luigs & Neumann, Ratingen, Germany). Recordings were performed at room temperature (21–24 °C). Signals were filtered at 2.9 kHz and digitized at 20 kHz using Patchmaster software (HEKA Elektronik, Lambrecht/Pfalz, Germany). Built-in compensation was used for voltage offset and pipette

520 capacitance. Membrane potentials were not corrected for liquid junction potential.

521 APs were evoked by brief 1 ms depolarizing current injection of 50–300 pA or 500 ms

522 hyperpolarizing pulses (300–400 pA) leading to APs upon return to resting

523 membrane potential ('rebound APs').

Preparation of organotypic hippocampal cultures

Cultures were prepared as described previously[78]. In brief, young Wistar rats (P5–

P7) were anesthetized with isoflurane and killed by decapitation, the brain was

removed, and each hippocampus was dissected. Hippocampal slices (350 µm) were

obtained using a Vibratome (Leica, VT1200S, Wetzlar, Germany). They were placed

on 20 mm latex membranes (Millicell; Merck, Darmstadt, Germany) inserted into 35

mm Petri dishes containing 1 mL of culture medium and maintained for up to 8 days

in an incubator at 34 °C, 95% O_2, 5% CO_2. The culture medium contained 25 mL

MEM, 12.5 mL HBSS, 12.5 mL horse serum, 0.5 mL penicillin/streptomycin, 0.8 mL

glucose (1 M), 0.1 mL ascorbic acid (1 mg/mL), 0.4 mL HEPES (1 M), 0.5 mL B27,

and 8.95 mL sterile H_2O. To limit glial proliferation, 5 µM ARA-C was added to the

culture medium starting at DIV 4.

AP recordings from organotypic hippocampal cultures

Whole-cell recordings were obtained from CA3 neurons in organotypic cultures at

DIV 7 to DIV 10. All recordings were made at 34 °C in a temperature-controlled

recording chamber (Luigs & Neumann, Ratingen, Germany) using 7–10 MΩ patch

pipettes. CA3 pyramidal neurons were recorded in current-clamp with a Multiclamp

700B Amplifier (Molecular Devices, San Jose, CA, USA). APs were elicited by

delivering DC step pulses (duration 1.0 s) at an interstimulus interval of 10–15 s.

During the whole recording, ionotropic glutamate and $GABA_A$ receptors were blocked

with 2–4 mM kynurenic acid and 100 µM picrotoxin, respectively.

Analysis of APs

Data were analyzed using IGOR PRO software (WaveMetrics, Lake Oswego, OR,

USA; Version 6.32A or 6.37), the Patcher's Power Tool (Version 2.19 Carbon,

15.03.2011) and NeuroMatic (Version 2.00, 15.09.2008; ref. [77]) extensions, as well as

self-written IGOR PRO scripts. AP durations were quantified as the duration at half-

maximal AP amplitude, amplitude being the difference between AP peak and the

membrane potential preceding stimulation. When AP durations were recorded at half-
maximal amplitude between AP threshold (defined as the membrane potential at
which the voltage rate of change first reaches 10 V/s) and AP peak, AP durations
were still unaffected by TTX treatment (data not shown). To exclude the effect of mis-
balanced bridge compensation on AP shape, only APs that occurred after the end of
the second current injection artifact (elicited by the end of current-injection) were
analyzed. In some recordings, current injection artifacts were blanked over 50 μs at
the beginning and the end of the current injection and only experiments in which
blanking did not affect AP amplitude and duration were subsequently analyzed.

Hippocampal dissociated cultures and transfection for GEVI imaging
Culture preparation and GEVI transfection were performed as previously described
by Sabater et al.[28]. In short, hippocampi were dissected from embryonic day 17.5
Wistar rat pups of either sex, treated with trypsin (Worthington Biochemical
Corporation, Lakewood, NJ, USA) at 0.5 mg/mL, and mechanically dissociated using
fire polished Pasteur pipettes. Neurons were plated on 18 mm glass coverslips pre-
treated with 100 μg/mL poly-L-lysine and coated with 10 μg/mL laminin (Thermo
Fisher Scientific, Waltham, MA, USA). Cultures were maintained in Neurobasal
medium with B27 (1x) and GlutaMAX (1x, all obtained from Thermo Fisher Scientific,
Waltham, MA, USA), supplemented with FBS (2%, Biosera, Nuaille, France) and
Penicillin/Streptomycin (1%, Sigma-Aldrich, St. Louis, MO, USA), at 37°C in a
humidified incubator with 5% CO_2. For transfections with Ace2N-mNeon-4AA (Ace-
mNeon, ref. [79]; distributed by Biolife Solutions Inc., Bothell, WA, USA), the Effectene
transfection reagent (Qiagen, Venlo, Netherlands) was used. The medium was
changed at DIV 3 to culture medium without antibiotic or FBS, and transfections with
Effectene were performed at DIV 7 following the manufacturer's protocol. After
transfection neurons were maintained in serum-free media without antibiotics. For all
experiments 30% of the medium was changed weekly and neurons were imaged 7–
14 days after transfection (14–21 DIV).

GEVI Imaging conditions
GEVI imaging was performed as previously described by Sabater et al.[28]. In short,
neurons were imaged using an inverted Olympus IX71 epifluorescence microscope
with a 60x 1.42 NA oil-immersion objective (Olympus, Tokyo, Japan). Coverslips

were mounted in a heated chamber (Warner instruments, Holliston, MA, USA; total

volume ~500 µL) and placed on an IMTP microscope stage (Scientifica, Uckfield,

United Kingdom). Cells were maintained in external HEPES-buffered saline solution

(HBS: 2 mM $CaCl_2$, 1.6 mM $MgCl_2$, 1.45 mM NaCl, 2.5 mM KCl, 10 mM Glucose, 10

mM HEPES, pH 7.4, Osmolarity 290 mOsm) and experiments were carried out at

near physiological temperatures of 32–35 ºC.

In Ace-mNeon imaging experiments, ~505 nm excitation illumination was provided

using a 525 nm LED (Solis LED, Thorlabs, Newton, NJ, USA), a 500/20 nm excitation

filter, and a 510 nm long-pass dichroic mirror. Ace-mNeon emission transmitted

through the dichroic was filtered using a 520 nm long-pass filters (Chroma, Bellows

Falls, VT, USA). A power density of 10 mW/mm^2 was obtained at the specimen

plane. Images were acquired at 3.2 kHz with an ORCA-Flash4.0 V2 C11440-22CU

scientific CMOS camera (Hamamatsu Photonics, Hamamatsu, Japan) cooled to ~ -

20 ºC with the Exos2 water cooling system (Koolance, Auburn, WA, USA). Images

were acquired with HCImage software (Hamamatsu, Hamamatsu, Japan), binned to

4 × 4 and cropped to a 16 × 512-pixel region of interest, necessary to achieve the

high image acquisition rates. Images were saved in CXD format.

Stimulation of APs was achieved with an extracellular tungsten parallel bipolar

electrode (FHC, Bowdoin, ME, USA) mounted on a PatchStar motorized micro-

manipulator (Scientifica, Uckfield, United Kingdom). To reliably induce an AP, we

delivered a 1 ms 10 mV pulse at the soma. The timing of the LED illumination, the

acquisition and stimulation were triggered externally through Clampex software

(pClamp 10, Molecular Devices, San Jose, CA, USA).

All GEVI Imaging recordings were performed in presence of NBQX (10 µM),

Gabazine (10 µM) and D-2-amino-5-phosphonovalerate (APV, 25 µM; all from Bio-

Techne, Wiesbaden-Nordenstadt, Germany) in order to block synaptic transmission

and ensure that the observed events were due to the stimulation only. IbTx was

delivered using a custom gravity-fed perfusion system and bath volume was

maintained by a peristaltic pump (Watson-Marlow 120s; Watson-Marlow,

Rommerskirchen, Germany).

Image analysis

Image analysis was performed as previously described by Sabater *et al.*[28]. In short, voltage imaging recordings were analyzed in ImageJ and custom scripts within MATLAB software as follows. Prior to analysis, if drift had occurred in the x and y axis during the recording, the images were aligned using the TurboReg ImageJ plugin[80]. Images were then imported into MATLAB and fluorescence intensity profiles were obtained from regions of interest (ROIs) manually drawn around neuronal structures identified visually on a maximum projection image of the time series by averaging across the ROI pixels in each frame. The fluorescence profile was separated into single trials and corrected for background camera noise by subtracting the average intensity value of the timepoints where no illumination light was applied. Then, recordings were reconstructed at a sampling rate of 100 kHz using cubic spline interpolation. Bleaching was estimated by fitting a single exponential function to each individual trace smoothed using an averaging filter with a window of 3 and interpolated. The resulting curve was used to correct an unfiltered interpolated version of the recording. An accurate representation of the AP waveform with an enhanced SNR was obtained by averaging over the repeats and extracting the AP parameters from the resulting trace. The fluorescence profile during the 20 ms preceding a response was used as baseline.

Solutions and reagents

Reagents were purchased from Sigma-Aldrich (St. Louis, MO, USA) if not indicated otherwise.

Conditions I, II, VII and XI

The extracellular bath solution contained (in mM): 122 NaCl, 3 KCl, 10 D-Glucose, 26 $NaHCO_3$, 1.25 Na_2HPO_4, 1.3 $MgCl_2$, and 2.0 $CaCl_2$. After equilibration with 95% O_2–5% CO_2, pH was adjusted to 7.35 if needed by NaOH or HCl. The intracellular pipette solution contained (in mM): 130 K gluconate, 1 $MgCl_2$, 10 HEPES, 0.3 EGTA, 10 Tris-Phosphocreatine, 4 Mg-ATP, 0.3 Na-GTP. When testing the contribution of BK channels, 300 nM Iberiotoxin (Abcam, Cambridge, United Kingdom) was included in the bath solution.

Conditions III and IV

The extracellular bath solution contained (in mM): 125 NaCl, 3 KCl, 25 Glucose, 25

$NaHCO_3$, 1.25 Na_2HPO_4, 1.1 $MgCl_2$, and 1.1 $CaCl_2$. After equilibration with 95% O_2–

5% CO_2, pH was adjusted to 7.35 if needed by NaOH or HCl. The intracellular pipette

solution contained (in mM): 150 K Gluconate, 10 K-HEPES, 3 Mg-ATP, 0.3 Na-GTP,

0.05 EGTA, 10 NaCl.

Conditions V and VI

The extracellular bath solution contained (in mM): 150 NaCl, 4 KCl, 10 HEPES, 10

glucose, 1.0 $MgCl_2$, and 2.0 $CaCl_2$, and was equilibrated with 95% O_2–5% CO_2. The

intracellular pipette solution contained (in mM): 135 K gluconate, 4 NaCl, 0.05 EGTA,

10 HEPES, 3 $MgCl_2$, 0.3 NaGTP, 3 Na_2ATP, 10 Na_2-Creatine phosphate with pH

normalized to 7.25 using KOH.

Condition VIII

The extracellular bath solution contained (in mM): 122 NaCl, 3 KCl, 1.25 NaH_2PO_4,

26 $NaHCO_3$, 10 glucose, 1.3 $MgCl_2$, and 2.0 $CaCl_2$, and was bubbled with 95% O_2–

5% CO_2. The intracellular pipette solution contained (in mM): 130 K gluconate, 0.3

EGTA, 10 HEPES, 2 $MgCl_2$, 0.3 NaGTP, 4 Na_2ATP (or Mg-ATP), 10 Na_2-Creatine

phosphate (or Tris-Creatinine phosphate).

Conditions IX and X

The extracellular bath solution contained (in mM): 145 NaCl, 2.5 KCl, 10 HEPES, 24

glucose, 1.3 $MgCl_2$, and 2.0 $CaCl_2$, and was equilibrated with 95% O_2–5% CO_2. The

intracellular pipette solution contained (in mM): 140 K gluconate, 2 NaCl, 0.5 EGTA,

20 HEPES, 3 $MgCl_2$, 0.3 NaGTP, 4 Na_2ATP, with pH adjusted to 7.25 using KOH.

Organotypic hippocampal neuron recordings

The external bath solution contained (in mM): 125 NaCl, 26 $NaHCO_3$, 3 $CaCl_2$, 2.5

KCl, 2 $MgCl_2$, 0.8 NaH_2PO_4, and 10 D-glucose, and was equilibrated with 95% O_2–

5% CO_2. The intracellular pipette solution contained (mM): 120 K gluconate, 20 KCl,

10 HEPES, 0.5 EGTA, 2 $MgCl_2$, 2 Na_2ATP, and 0.3 NaGTP (pH 7.4).

Statistics

If not otherwise noted, data are expressed as box plots with boxes reflecting 25th–

75th percentile. The median is indicated by a line within the box, whiskers indicate

10th and 90th percentile. Sample sizes (n) provide the number of individual somatic

recordings. All data sets were obtained from ≥ 2 independent cell cultures. Data in

Fig. 3A were tested for statistical difference using Fisher's exact test. Confidence

limits for 95% confidence intervals were calculated as Wilson interval[81]. Other

experimental groups were independent and tested by the Mann-Whitney *U* test.

Tests were performed in IGOR PRO or GraphPad Prism 8. Results were considered

significant at $p < 0.05$.

Acknowledgment

This work was supported by a European Research Council Consolidator Grant (ERC CoG 865634) the German Research Foundation (HA6386/10-2) to S.H. and by grants awarded by the U.S. Department of Veterans Affairs (BX002547) and NIGMS (GM134110) to S.M.S.

Declaration of interests

The authors declare no competing interests.

References

1. Borst, J.G.G., and Sakmann, B. (1999). Effect of changes in action potential shape on
calcium currents and transmitter release in a calyx-type synapse of the rat auditory
brainstem. Philosophical Transactions of the Royal Society B: Biological Sciences *354*,
347–355. https://doi.org/10.1098/RSTB.1999.0386.
2. Christie, B.R., Eliot, L.S., Ito, K.I., Miyakawa, H., and Johnston, D. (1995). Different
Ca2+ channels in soma and dendrites of hippocampal pyramidal neurons mediate
spike-induced Ca2+ influx. J Neurophysiol *73*, 2553–2557.
https://doi.org/10.1152/jn.1995.73.6.2553.
3. Watanabe, S., Hoffman, D.A., Migliore, M., and Johnston, D. (2002). Dendritic K+
channels contribute to spike-timing dependent long-term potentiation in hippocampal
pyramidal neurons. Proc Natl Acad Sci U S A *99*, 8366–8371.
https://doi.org/10.1073/PNAS.122210599.
4. Bean, B.P. (2007). The action potential in mammalian central neurons. Nat Rev
Neurosci *8*, 451–465. https://doi.org/10.1038/nrn2148.
5. Debanne, D., Campanac, E., Bialowas, A., Carlier, E., and Alcaraz, G. (2011). Axon
physiology. Physiol Rev *91*, 555–602. https://doi.org/10.1152/PHYSREV.00048.2009.
6. Geiger, J.R.P., and Jonas, P. (2000). Dynamic control of presynaptic Ca2+ inflow by
fast-inactivating K+ channels in hippocampal mossy fiber boutons. Neuron *28*, 927–
939. https://doi.org/10.1016/S0896-6273(00)00164-1.
7. Jackson, M.B., Konnerth, A., and Augustine, G.J. (1991). Action potential broadening
and frequency-dependent facilitation of calcium signals in pituitary nerve terminals.
Proc Natl Acad Sci U S A *88*, 380–384. https://doi.org/10.1073/PNAS.88.2.380.
8. Keck, T., Toyoizumi, T., Chen, L., Doiron, B., Feldman, D.E., Fox, K., Gerstner, W.,
Haydon, P.G., Hübener, M., Lee, H.K., et al. (2017). Integrating Hebbian and
homeostatic plasticity: The current state of the field and future research directions.
Philosophical Transactions of the Royal Society B: Biological Sciences *372*.
https://doi.org/10.1098/rstb.2016.0158.
9. Carta, M., Lanore, F., Rebola, N., Szabo, Z., Da Silva, S.V., Lourenço, J., Verraes, A.,
Nadler, A., Schultz, C., Blanchet, C., et al. (2014). Membrane lipids tune synaptic
transmission by direct modulation of presynaptic potassium channels. Neuron *81*, 787–
799. https://doi.org/10.1016/J.NEURON.2013.12.028.
10. Senn, V., Wolff, S.B.E., Herry, C., Grenier, F.O., Ehrlich, I., Grü, J., Fadok, J.P.,
Müller, C., Letzkus, J.J., and Lüthi, A. (2014). Long-range connectivity defines
behavioral specificity of amygdala neurons. Neuron *81*, 428–437.
https://doi.org/10.1016/j.neuron.2013.11.006.
11. Turrigiano, G.G., and Nelson, S.B. (2004). Homeostatic plasticity in the developing
nervous system. Nat Rev Neurosci *5*, 97–107. https://doi.org/10.1038/nrn1327.
12. Pozo, K., and Goda, Y. (2010). Unraveling mechanisms of homeostatic synaptic
plasticity. Neuron *66*, 337–351. https://doi.org/10.1016/J.NEURON.2010.04.028.
13. Li, B., Suutari, B.S., Sun, S.D., Luo, Z., Wei, C., Chenouard, N., Mandelberg, N.J.,
Zhang, G., Wamsley, B., Tian, G., et al. (2020). Neuronal inactivity co-opts LTP
machinery to drive potassium channel splicing and homeostatic spike widening. Cell
181, 1547-1565.e15. https://doi.org/10.1016/j.cell.2020.05.013.
14. Ritzau-Jost, A., Tsintsadze, T., Krueger, M., Ader, J., Bechmann, I., Eilers, J., Barbour,
B., Smith, S.M., and Hallermann, S. (2021). Large, stable spikes exhibit differential
broadening in excitatory and inhibitory neocortical boutons. Cell Rep *34*, 108612.
https://doi.org/10.1016/j.celrep.2020.108612.
15. Zbili, M., Rama, S., Benitez, M.J., Fronzaroli-Molinieres, L., Bialowas, A.,
Boumedine-Guignon, N., Garrido, J.J., and Debanne, D. (2021). Homeostatic
regulation of axonal Kv1.1 channels accounts for both synaptic and intrinsic

760		modifications in the hippocampal CA3 circuit. Proc Natl Acad Sci U S A *118*.
761		https://doi.org/10.1073/pnas.2110601118.
762	16.	Kim, J., and Tsien, R.W. (2008). Synapse-specific adaptations to inactivity in
763		hippocampal circuits achieve homeostatic gain control while dampening network
764		reverberation. Neuron *58*, 925–937. https://doi.org/10.1016/j.neuron.2008.05.009.
765	17.	Mitra, A., Mitra, S.S., and Tsien, R.W. (2012). Heterogeneous reallocation of
766		presynaptic efficacy in recurrent excitatory circuits adapting to inactivity. Nat Neurosci
767		*15*, 250–257. https://doi.org/10.1038/nn.3004.
768	18.	Castro-Alamancos, M.A., and Connors, B.W. (1997). Distinct forms of short-term
769		plasticity at excitatory synapses of hippocampus and neocortex. Proc Natl Acad Sci U
770		S A *94*, 4161–4166. https://doi.org/10.1073/pnas.94.8.4161.
771	19.	Manahan-Vaughan, D. (2018). Special considerations when using mice for In vivo
772		electrophysiology and long-term studies of hippocampal synaptic plasticity during
773		behavior. Handb Behav Neurosci *28*, 63–84. https://doi.org/10.1016/B978-0-12-
774		812028-6.00003-3.
775	20.	Nguyen, P. V., Duffy, S.N., and Young, J.Z. (2000). Differential maintenance and
776		frequency-dependent tuning of LTP at hippocampal synapses of specific strains of
777		inbred mice. J Neurophysiol *84*, 2484–2493. https://doi.org/10.1152/jn.2000.84.5.2484.
778	21.	Prestigio, C., Ferrante, D., Valente, P., Casagrande, S., Albanesi, E., Yanagawa, Y.,
779		Benfenati, F., and Baldelli, P. (2019). Spike-Related Electrophysiological Identification
780		of Cultured Hippocampal Excitatory and Inhibitory Neurons. Mol Neurobiol *56*, 6276–
781		6292. https://doi.org/10.1007/s12035-019-1506-5.
782	22.	Maffei, A., and Turrigiano, G.G. (2008). Multiple modes of network homeostasis in
783		visual cortical layer 2/3. Journal of Neuroscience *28*, 4377–4384.
784		https://doi.org/10.1523/JNEUROSCI.5298-07.2008.
785	23.	Desai, N.S., Rutherford, L.C., and Turrigiano, G.G. (1999). Plasticity in the intrinsic
786		excitability of cortical pyramidal neurons. Nat Neurosci *2*, 515–520.
787		https://doi.org/10.1038/9165.
788	24.	Turrigiano, G.G., Leslie, K.R., Desai, N.S., Rutherford, L.C., and Nelson, S.B. (1998).
789		Activity-dependent scaling of quantal amplitude in neocortical neurons. Nature *391*,
790		892–896. https://doi.org/10.1038/36103.
791	25.	Hanes, A.L., Koesters, A.G., Fong, M.-F., Altimimi, H.F., Stellwagen, D., Wenner, P.,
792		and Engisch, K.L. (2020). Cellular/molecular divergent synaptic scaling of miniature
793		EPSCs following activity blockade in dissociated neuronal cultures. *40*, 4090–4102.
794		https://doi.org/10.1523/JNEUROSCI.1393-19.2020.
795	26.	Frank, C.A., Kennedy, M.J., Goold, C.P.P., Marek, K.W., and Davis, G.W.W. (2006).
796		Mechanisms underlying the rapid induction and sustained expression of synaptic
797		homeostasis. Neuron *52*, 663–677. https://doi.org/10.1016/J.NEURON.2006.09.029.
798	27.	Aoto, J., Nam, C.I., Poon, M.M., Ting, P., and Chen, L. (2008). Synaptic Signaling by
799		All-Trans Retinoic Acid in Homeostatic Synaptic Plasticity. Neuron *60*, 308–320.
800		https://doi.org/10.1016/j.neuron.2008.08.012.
801	28.	Gonzalez Sabater, V., Rigby, M., and Burrone, J. (2021). Voltage-gated potassium
802		channels ensure action potential shape fidelity in distal axons. The Journal of
803		Neuroscience *41*, 5372–5385. https://doi.org/10.1523/jneurosci.2765-20.2021.
804	29.	Hu, H., Shao, L.-R., Chavoshy, S., Gu, N., Trieb, M., Behrens, R., Laake, P., Pongs,
805		O., Knaus, H.G., Ottersen, O.P., et al. (2001). Presynaptic Ca2+-activated K+ channels
806		in glutamatergic hippocampal terminals and their role in spike repolarization and
807		regulation of transmitter release. The Journal of Neuroscience *21*, 9585–9597.
808		https://doi.org/10.1523/JNEUROSCI.21-24-09585.2001.

30. Karmarkar, U.R., and Buonomano, D. V. (2006). Different forms of homeostatic
plasticity are engaged with distinct temporal profiles. European Journal of
Neuroscience *23*, 1575–1584. https://doi.org/10.1111/j.1460-9568.2006.04692.x.

31. Lee, K.Y., and Chung, H.J. (2014). NMDA receptors and L-type voltage-gated Ca2+
channels mediate the expression of bidirectional homeostatic intrinsic plasticity in
cultured hippocampal neurons. Neuroscience *277*, 610–623.
https://doi.org/10.1016/j.neuroscience.2014.07.038.

32. Lee, K.Y., Royston, S.E., Vest, M.O., Ley, D.J., Lee, S., Bolton, E.C., and Chung, H.J.
(2015). N-methyl-D-aspartate receptors mediate activity-dependent down-regulation of
potassium channel genes during the expression of homeostatic intrinsic plasticity. Mol
Brain *8*, 1–16. https://doi.org/10.1186/s13041-015-0094-1.

33. Zhao, C.J., Dreosti, E., and Lagnado, L. (2011). Homeostatic synaptic plasticity
through changes in presynaptic calcium influx. Journal of Neuroscience *31*, 7492–
7496. https://doi.org/10.1523/JNEUROSCI.6636-10.2011.

34. Desai, N.S., Rutherford, L.C., and Turrigiano, G.G. (1999). BDNF regulates the
intrinsic excitability of cortical neurons. Learning and Memory *6*, 284–291.
https://doi.org/10.1101/lm.6.3.284.

35. Rutherford, L.C., DeWan, A., Lauer, H.M., and Turrigiano, G.G. (1997). Brain-derived
neurotrophic factor mediates the activity-dependent regulation of inhibition in
neocortical cultures. Journal of Neuroscience *17*, 4527–4535.
https://doi.org/10.1523/jneurosci.17-12-04527.1997.

36. Ritzau-Jost, A., Nerlich, J., Kaas, T., Krueger, M., Tsintsadze, T., Eilers, J., Barbour,
B., Smith, S.M., and Hallermann, S. (2023). Direct whole-cell patch-clamp recordings
from small boutons in rodent primary neocortical neuron cultures. STAR Protoc *4*,
102168. https://doi.org/10.1016/j.xpro.2023.102168.

37. Chipman, P.H., Fetter, R.D., Panzera, L.C., Bergerson, S.J., Karmelic, D., Yokoyama,
S., Hoppa, M.B., and Davis, G.W. (2022). NMDAR-dependent presynaptic
homeostasis in adult hippocampus: Synapse growth and cross- modal inhibitory
plasticity. Neuron *110*, ki. https://doi.org/10.1016/j.neuron.2022.08.014.

38. Niday, Z., and Bean, B.P. (2021). BK channel regulation of afterpotentials and burst
firing in cerebellar purkinje neurons. Journal of Neuroscience *41*, 2854–2869.
https://doi.org/10.1523/JNEUROSCI.0192-20.2021.

39. Gittis, A.H., Moghadam, S.H., and Du Lac, S. (2010). Mechanisms of sustained high
firing rates in two classes of vestibular nucleus neurons: Differential contributions of
resurgent Na, Kv3, and BK currents. J Neurophysiol *104*, 1625–1634.
https://doi.org/10.1152/jn.00378.2010.

40. Kimm, T., Khaliq, Z.M., and Bean, B.P. (2015). Differential regulation of action
potential shape and burst-frequency firing by BK and Kv2 channels in substantia nigra
dopaminergic neurons. Journal of Neuroscience *35*, 16404–16417.
https://doi.org/10.1523/JNEUROSCI.5291-14.2015.

41. Müller, A., Kukley, M., Uebachs, M., Beck, H., and Dietrich, D. (2007). Nanodomains
of single Ca2+ channels contribute to action potential repolarization in cortical
neurons. Journal of Neuroscience *27*, 483–495.
https://doi.org/10.1523/JNEUROSCI.3816-06.2007.

42. Bock, T., and Stuart, G.J. (2016). The impact of BK channels on cellular excitability
depends on their subcellular location. Front Cell Neurosci *10*, 1–8.
https://doi.org/10.3389/fncel.2016.00206.

43. Desai, N.S., Cudmore, R.H., Nelson, S.B., and Turrigiano, G.G. (2002). Critical
periods for experience-dependent synaptic scaling in visual cortex. Nat Neurosci *5*,
783–789. https://doi.org/10.1038/nn878.

44. Wiesel, T.N., and Hubel, D.H. (1963). Single-cell responses in striate cortex of kittens
deprived of vision in one eye. J Neurophysiol *26*, 1003–1017.

45. Wen, W., and Turrigiano, G.G. (2021). Developmental regulation of homeostatic
plasticity in mouse primary visual cortex. The Journal of Neuroscience *41*, JN-RM-
1200-21. https://doi.org/10.1523/jneurosci.1200-21.2021.

46. Trojanowski, N.F., and Turrigiano, G.G. (2021). CaMKIV signaling is not essential for
the maintenance of intrinsic or synaptic properties in mouse visual cortex. eNeuro *8*, 1–
12. https://doi.org/10.1523/ENEURO.0135-21.2021.

47. Griffen, T.C., Haley, M.S., Fontanini, A., and Maffei, A. (2017). Rapid plasticity of
visually evoked responses in rat monocular visual cortex. PLoS One *12*, 1–12.
https://doi.org/10.1371/journal.pone.0184618.

48. Gainey, M.A., Aman, J.W., and Feldman, D.E. (2018). Rapid disinhibition by
adjustment of PV intrinsic excitability during whisker map plasticity in mouse S1.
Journal of Neuroscience *38*, 4749–4761. https://doi.org/10.1523/JNEUROSCI.3628-
17.2018.

49. Maravall, M., Stern, E.A., and Svoboda, K. (2004). Development of intrinsic properties
and excitability of layer 2/3 pyramidal neurons during a critical period for sensory
maps in rat barrel cortex. J Neurophysiol *92*, 144–156.
https://doi.org/10.1152/jn.00598.2003.

50. Breton, J.D., and Stuart, G.J. (2009). Loss of sensory input increases the intrinsic
excitability of layer 5 pyramidal neurons in rat barrel cortex. J Physiol *587*, 5107–5119.
https://doi.org/10.1113/JPHYSIOL.2009.180943.

51. Sun, Q.Q. (2009). Experience-dependent intrinsic plasticity in interneurons of barrel
cortex layer IV. J Neurophysiol *102*, 2955–2973.
https://doi.org/10.1152/jn.00562.2009.

52. Li, L., Gainey, M.A., Goldbeck, J.E., and Feldman, D.E. (2014). Rapid homeostasis by
disinhibition during whisker map plasticity. Proc Natl Acad Sci U S A *111*, 1616–
1621. https://doi.org/10.1073/pnas.1312455111.

53. Jamann, N., Dannehl, D., Lehmann, N., Wagener, R., Thielemann, C., Schultz, C.,
Staiger, J., Kole, M.H.P., and Engelhardt, M. (2021). Sensory input drives rapid
homeostatic scaling of the axon initial segment in mouse barrel cortex. Nat Commun
12, 1–14. https://doi.org/10.1038/s41467-020-20232-x.

54. Hickmott, P.W. (2005). Changes in intrinsic properties of pyramidal neurons in adult
rat S1 during cortical reorganization. J Neurophysiol *94*, 501–511.
https://doi.org/10.1152/jn.00924.2004.

55. Francis, H.W., and Manis, P.B. (2000). Effects of deafferentation on the
electrophysiology of ventral cochlear nucleus neurons. Hear Res *149*, 91–105.
https://doi.org/10.1016/S0378-5955(00)00165-9.

56. Lu, Y., Monsivais, P., Tempel, B.L., and Rubel, E.W. (2004). Activity-dependent
regulation of the potassium channel subunits Kv1.1 and Kv3.1. Journal of Comparative
Neurology *470*, 93–106. https://doi.org/10.1002/cne.11037.

57. Wang, Y., and Manis, P.B. (2006). Temporal coding by cochlear nucleus bushy cells in
DBA/2J mice with early onset hearing loss. JARO - Journal of the Association for
Research in Otolaryngology *7*, 412–424. https://doi.org/10.1007/s10162-006-0052-9.

58. Leao, R.N., Berntson, A., Forsythe, I.D., and Walmsley, B. (2004). Reduced low-
voltage activated K+ conductances and enhanced central excitability in a congenitally
deaf (dn/dn) mouse. J Physiol *559*, 25–33.
https://doi.org/10.1113/JPHYSIOL.2004.067421.

59. Him, A., and Dutia, M.B. (2001). Intrinsic excitability changes in vestibular nucleus
neurons after unilateral deafferentation. Brain Res *908*, 58–66.
https://doi.org/10.1016/S0006-8993(01)02600-2.

60. George, N.M., Gentile Polese, A., Merle, L., Macklin, W.B., and Restrepo, D. (2022). Excitable axonal domains adapt to sensory deprivation in the olfactory system. The Journal of Neuroscience 42, 1491–1509. https://doi.org/10.1523/JNEUROSCI.0305-21.2021.

61. Barnes, B.Y.C.A., and McNaughton, B.L. (1980). Physiological compensation for loss of afferent synapses in rat hippocampal granule cells during senescence. J. Physiol 309, 473–485. https://doi.org/10.1113/jphysiol.1980.sp013521ch.

62. Midorikawa, M., and Miyata, M. (2021). Distinct functional developments of surviving and eliminated presynaptic terminals. Proc Natl Acad Sci U S A 118. https://doi.org/10.1073/pnas.2022423118.

63. O'Leary, T., van Rossum, M.C.W., and Wyllie, D.J.A. (2010). Homeostasis of intrinsic excitability in hippocampal neurones: Dynamics and mechanism of the response to chronic depolarization. Journal of Physiology 588, 157–170. https://doi.org/10.1113/jphysiol.2009.181024.

64. Pozzi, D., Lignani, G., Ferrea, E., Contestabile, A., Paonessa, F., D'Alessandro, R., Lippiello, P., Boido, D., Fassio, A., Meldolesi, J., et al. (2013). REST/NRSF-mediated intrinsic homeostasis protects neuronal networks from hyperexcitability. EMBO Journal 32, 2994–3007. https://doi.org/10.1038/emboj.2013.231.

65. Wefelmeyer, W., Cattaert, D., and Burrone, J. (2015). Activity-dependent mismatch between axo-axonic synapses and the axon initial segment controls neuronal output. Proc Natl Acad Sci U S A 112, 9757–9762. https://doi.org/10.1073/pnas.1502902112.

66. Malik, R., and Chattarji, S. (2012). Enhanced intrinsic excitability and EPSP-spike coupling accompany enriched environment-induced facilitation of LTP in hippocampal CA1 pyramidal neurons. J Neurophysiol 107, 1366–1378. https://doi.org/10.1152/jn.01009.2011.

67. Valero-Aracama, M.J., Sauvage, M.M., and Yoshida, M. (2015). Environmental enrichment modulates intrinsic cellular excitability of hippocampal CA1 pyramidal cells in a housing duration and anatomical location-dependent manner. Behavioural Brain Research 292, 209–218. https://doi.org/10.1016/j.bbr.2015.05.032.

68. Kole, M.H.P., Czéh, B., and Fuchs, E. (2004). Homeostatic maintenance in excitability of tree shrew hippocampal CA3 pyramidal neurons after chronic stress. Hippocampus 14, 742–751. https://doi.org/10.1002/hipo.10212.

69. Kamal, A., Notenboom, R.G.E., De Graan, P.N.E., and Ramakers, G.M.J. (2006). Persistent changes in action potential broadening and the slow afterhyperpolarization in rat CA1 pyramidal cells after febrile seizures. European Journal of Neuroscience 23, 2230–2234. https://doi.org/10.1111/j.1460-9568.2006.04732.x.

70. Eshra, A., Hirrlinger, P., and Hallermann, S. (2019). Enriched environment shortens the duration of action potentials in cerebellar granule cells. Front Cell Neurosci 13, 1–9. https://doi.org/10.3389/fncel.2019.00289.

71. Ngodup, T., Goetza, J.A., McGuire, B.C., Sun, W., Lauer, A.M., and Xu-Friedman, M.A. (2015). Activity-dependent, homeostatic regulation of neurotransmitter release from auditory nerve fibers. Proc Natl Acad Sci U S A 112, 6479–6484. https://doi.org/10.1073/PNAS.1420885112/-/DCSUPPLEMENTAL/PNAS.201420885SI.PDF.

72. Simkin, D., Hattori, S., Ybarra, N., Musia, T.F., Buss, E.W., Richter, H., Matthew Oh, M., Nicholson, D.A., and Disterhoft, J.F. (2015). Aging-related hyperexcitability in CA3 pyramidal neurons is mediated by enhanced A-type K+ channel function and expression. Journal of Neuroscience 35, 13206–13218. https://doi.org/10.1523/JNEUROSCI.0193-15.2015.

73. Mehranfard, N., Gholamipour-Badie, H., Motamedi, F., Janahmadi, M., and Naderi, N. (2015). Long-term increases in BK potassium channel underlie increased action

potential firing in dentate granule neurons following pilocarpine-induced status
epilepticus in rats. Neurosci Lett *585*, 88–91.
https://doi.org/10.1016/j.neulet.2014.11.041.
74.   Moutin, E., Hemonnot, A.L., Seube, V., Linck, N., Rassendren, F., Perroy, J., and
Compan, V. (2020). Procedures for culturing and genetically manipulating murine
hippocampal postnatal neurons. Front Synaptic Neurosci *12*, 1–16.
https://doi.org/10.3389/fnsyn.2020.00019.
75.   Martiszus, B.J., Tsintsadze, T., Chang, W., and Smith, S.M. (2021). Enhanced
excitability of cortical neurons in low-divalent solutions is primarily mediated by
altered voltage-dependence of voltage-gated sodium channels. Elife *10*, 1–26.
https://doi.org/10.7554/eLife.67914.
76.   Clements, J.D., and Bekkers, J.M. (1997). Detection of spontaneous synaptic events
with an optimally scaled template. Biophys J *73*, 220–229.
https://doi.org/10.1016/S0006-3495(97)78062-7.
77.   Rothman, J.S., and Silver, R.A. (2018). Neuromatic: An integrated open-source
software toolkit for acquisition, analysis and simulation of electrophysiological data.
Front Neuroinform *12*, 1–21. https://doi.org/10.3389/fninf.2018.00014.
78.   Debanne, D., Boudkkazi, S., Campanac, E., Cudmore, R.H., Giraud, P., Fronzaroli-
Molinieres, L., Carlier, E., and Caillard, O. (2008). Paired-recordings from synaptically
coupled cortical and hippocampal neurons in acute and cultured brain slices. Nat Protoc
3, 1559–1568. https://doi.org/10.1038/nprot.2008.147.
79.   Gong, Y., Huang, C., Li, J.Z., Grewe, B.F., Zhang, Y., Eismann, S., and Schnitzer,
M.J. (2015). High-speed recording of neural spikes in awake mice and flies with a
fluorescent voltage sensor. Science (1979) *350*, 1361–1366.
https://doi.org/10.1126/science.aab0810.
80.   Thevenaz, P., Ruttimann, U.E., and Unser, M. (1998). A pyramid approach to subpixel
registration based on intensity. IEEE Transactions on Image Processing *7*, 27–41.
https://doi.org/10.1109/83.650848.
81.   Brown, L.D., Cai, T.T., and DasGupta, A. (2001). Interval estimation for a binomial
proportion. Statistical Science *16*, 101–133.

996   **Figures and legends**

1000

Figure 1 Cell type- and model-specific AP broadening in hippocampal neurons

(A) Example evoked APs recorded from CA3 pyramidal neurons in organotypic

hippocampal cultures under control condition (black) and following 48 hours of

TTX-treatment (blue).

(B) AP durations recorded under conditions in A.

(C) Example depolarization-evoked APs (500 pA for 1 ms) recorded from neurons in

primary dissociated hippocampal cultures under control condition (black) and

following 48 hours of TTX- (blue), Kyn- (purple), or NBQX-treatment (turquoise).

(D) AP durations recorded under conditions in C.

(E) Example spontaneous APs recorded under the conditions in C.

(F) Spontaneous AP frequency under the conditions in C.

Numbers in brackets reflect number of recorded neurons. Box plots as median ±

interquartile range, whiskers indicate 10–90 percentile. P-values above graphs,

significance tested by Mann-Whitney-U test.

1015

1016

Figure 2 Stable AP duration in primary dissociated neurons during TTX-induced homeostatic plasticity

(A) Example current injection-evoked APs (500 pA for 1 ms) under control condition (black) and following 48 hours of TTX-treatment (blue). AP amplitudes are recorded between the membrane potential preceding the AP and AP peak (arrows), AP durations are recorded at half-maximal amplitude (arrow heads).

(B) AP duration in control and TTX-treated cells (color code as in A) for eleven different experimental conditions (I–XI). Specification of individual conditions is provided in Table S1. Condition XI adopted from Fig. 1D.

(C) Merge of AP durations in control and TTX-treated cells across conditions I–IX normalized to the corresponding condition's control median duration.

(D) AP amplitudes in control and TTX-treated cells as in B.

(E) Cross-conditional merge of AP amplitudes in control and TTX-treated cells as in C.

(F) Example hyperpolarization-evoked "rebound APs" (-150 pA or -60 pA for 500ms) in control and TTX-treated cells (color code as in A).

1035   (G) Rebound AP duration in control and TTX-treated cells as in B.

1036   (H) Cross-conditional merge of normalized durations as in C for rebound APs.

1037   Numbers in brackets reflect number of recorded neurons. Box plots as median ±

1038   interquartile range, whiskers indicate 10–90 percentile. P-values provided above

1039   graphs; significance tested using Mann-Whitney-*U* tests.

1040

1041

1042

1043

1044

Figure 3 Homeostatic plasticity increases network activity and excitatory synaptic transmission

(A) *Left:* Spontaneously active control (black) and TTX-treated cells (blue) in condition I–IV and VII. *Right:* Active neurons merged across conditions I–IV and VII.

(B) Example spontaneous AP bursts in control and TTX-treated cells. Asterisks mark individual APs.

(C) AP number per burst in control and TTX-treated cells across conditions I–IV and VII.

(D) Example spontaneous excitatory postsynaptic currents (sEPSCs) in control and TTX-treated cells.

(E) *Left:* sEPSC frequency and *Right:* sEPSC amplitude in control recordings and after TTX-treatment (merged data from conditions I and II).

1060   Numbers in brackets reflect number of recorded neurons. Bars in A reflect fraction

1061   of spontaneously active neurons. Whiskers in A reflect confidence limits for 95%

1062   confidence interval based on Wilson interval. Box plots as median ± interquartile

1063   range, whiskers indicate 10–90 percentile. P-values above graphs, significance

1064   tested by Fisher's exact test in A or Mann-Whitney-U test elsewhere.

1065

1066

1067

1068

1069

Figure 4 Lack of BK channel-dependent AP broadening in dissociated cultured neurons

(A) *Left:* Example current injection-evoked APs (500 pA for 1 ms) in control solution (black) and solution containing 300 nM Iberiotoxin (IbTx, green). *Right:* AP duration in control and TTX-treated cells recorded IbTx-free (black and blue, respectively) or IbTx-containing solution (green). Data recorded under condition I.

(B) Merged AP duration across control and TTX-treated cells for recordings in IbTx-free and IbTx-containing solution (data in A normalized to the respective group's median duration in control solution).

(C) APs evoked by a 200 ms current injection (top) and overlay of the forst three evoked APs (bottom).

(D) Broadening of AP duration of the 3 first APs during 200 ms current injections in control (black, *left*) and TTX-treated cells (blue, *right*). Bars as mean ± SEM, lines with markers depict recordings from individual neurons.

(E) *Left:* Example Ace-mNeon-recorded somatic, proximal axonal, and distal axonal APs in a primary hippocampal cultured neuron in IbTx-free solution (black) and

1088   following BK channel block by IbTx (100 nM, green). *Right:* Ratio of somatic and

1089   axonal AP durations in IbTx-free and IbTx-containing solution.

1090 (F) *Left:* Example AP train (5 x 20 Hz) and overlay of 1^{st} and 5^{th} train AP at the soma,

1091   proximal, and distal axon in control condition (grey and black) and following BK

1092   channel block by IbTx (yellow and green). *Right:* AP broadening (ratio of 5^{th} over

1093   1^{st} train AP duration) in the three compartments in control and IbTx-containing

1094   solution.

1095 (G) Somatic APs under control condition (grey), following treatment with 4-AP (black),

1096   and following treatment with 30μM 4-AP + 100 nM IbTx (green).

1097 Numbers in brackets reflect number of recorded neurons. Box plots as median ±

1098 interquartile range, whiskers indicate 10–90 percentile. Plots in D–F as mean ± SEM.

1099 P-values above graphs, significance tested by Mann-Whitney-*U* test.

1100

1101

1102   **Figure supplements**

Group	Species	Strain	Brain area	Culture age (days *in vitro*)	Lab which recordings were performed at	External recording solution	External [Ca^{2+}] (mM)	Internal recording solution	Growth medium	Protocol Reference
I	Rat	Sprague-Dawley	Cortex	10–14	Hallermann Lab, Leipzig (Germany)	1.)	2.0	5.)	NbActiv4	Li *et al.*, Cell 2020
II	Rat	Sprague-Dawley	Cortex	16–21	Hallermann Lab, Leipzig (Germany)	1.)	2.0	5.)	NbActiv4	Li *et al.*, Cell 2020
III	Mouse	C57BL/6N	Cortex	15–20	Hallermann Lab, Leipzig (Germany)	2.)	1.1	6.)	BrainPhys	Moutin *et al.*, Front. Synaptic Neurosci 2020
IV	Mouse	C57BL/6N	Cortex	16	Hallermann Lab, Leipzig (Germany)	2.)	1.1	6.)	MEM, 5% serum + supplements	Ritzau-Jost *et al.*, Cell Reports 2021
V	Mouse	C57BL/6J	Cortex	11–13	Smith Lab, Portland (OR, USA)	3.)	2.0	7.)	MEM, 5% serum + supplements	Martiszus BJ *et al.*, eLife 2021
VI	Mouse	CD1	Cortex	11–13	Smith Lab, Portland (OR, USA)	3.)	2.0	7.)	MEM, 5% serum + supplements	Martiszus BJ *et al.*, eLife 2021
VII	Rat	Sprague-Dawley	Pre-frontal cortex	10–14	Hallermann Lab, Leipzig (Germany)	1.)	2.0	5.)	BrainPhys	Moutin *et al.*, Front. Synaptic Neurosci 2020
VIII	Mouse	C57BL/6J	Pre-frontal cortex	10–14	Smith Lab, Portland (OR, USA)	1.)	2.0	8.)	MEM, 5% serum + supplements	Martiszus BJ *et al.*, eLife 2021
IX	Mouse	C57BL/6N	Hippo-campus	32	Heine Lab, Mainz (Germany)	4.)	2.0	9.)	Neurobasal A+ supplements	Li *et al.*, Cell 2020
X	Mouse	C57BL/6J	Hippo-campus	15	Heine Lab, Mainz (Germany)	4.)	2.0	9.)	Neurobasal A + supplements	Li *et al.*, Cell 2020
XI	Rat	Sprague-Dawley	Hippo-campus	16–20	Hallermann Lab, Leipzig (Germany)	1.)	2.0	5.)	BrainPhys	Moutin *et al.*, Front. Synaptic Neurosci 2020

1.)   122 NaCl, 3 KCl, 10 Glucose, 26 NaHCO$_3$, 2 CaCl$_2$, 1.3 MgCl$_2$, 1.25 NaH$_2$PO$_4$
2.)   125 NaCl, 3 KCl, 25 Glucose, 25 NaHCO$_3$, 1.1 CaCl$_2$, 1.1 MgCl$_2$, 1.25 Na$_2$HPO$_4$
3.)   150 NaCl, 4 KCl, 10 Glucose, 10 HEPES, 2 CaCl$_2$, 1 MgCl$_2$
4.)   145 NaCl, 2.5 KCl, 24 Glucose, 10 HEPES, 2 CaCl$_2$, 1.3 MgCl$_2$
5.)   130 K gluconate, 10 HEPES, 1 MgCl$_2$, 0.3 EGTA, 4 Mg-ATP, 0.3 Na-GTP, 10 Tris-Phosphocreatine
6.)   150 K gluconate, 10 K-HEPES, 10 NaCl, 0.05 EGTA, 3 Mg-ATP, 0.3 Na-GTP
7.)   135 K gluconate, 10 HEPES, 4 NaCl, 0.05 EGTA, 3 Na$_2$ATP, 0.3 NaGTP, 3 MgCl$_2$, 10 Na$_2$-Creatine phosphate
8.)   130 K gluconate, 10 HEPES, 0.3 EGTA, 4 Na$_2$ATP (or Mg-ATP), 0.3 NaGTP, 2 MgCl$_2$, 10 Na$_2$-Creatine phosphate (or Tris-Creatinine phosphate)
9.)   140 K gluconate, 20 HEPES, 2 NaCl, 0.5 EGTA, 4 Na$_2$ATP, 0.3 NaGTP, 3 MgCl$_2$

1103

1104

1105   **Figure 2 – figure supplement 1**

1106   **Additional information on recording conditions**

1107   The table lists additional relevant information on the experimental conditions I – XI.

1108

1109

1110

1111

1112

1113

Figure 2 – figure supplement 2

Absence of TTX-induced AP broadening in further selected conditions

(A) Box plots of AP durations under control condition (black) or upon TTX-treatment

(blue) recorded in condition I either unselected or selected for parameters

reflecting unimpaired neuronal function.

(B) AP durations as in A for recordings in condition VII.

(C) AP durations as in A for recordings in condition VIII.

Numbers of recorded neurons provided in brackets. Box plots depict median ±

interquartile range, whiskers reflect 10–90 percentile. P-values calculated by Mann-

Whitney-U test.

1124

1125

1126

1127

1128

Figure 4 – figure supplement 1

BK channel block does not affect rebound AP duration

(A) Example rebound APs in control solution (black) and solution containing 300 nM

Iberiotoxin (IbTx, green).

(B) AP duration in control and TTX-treated cells in IbTx-free (black and blue,

respectively) or IbTx-containing solution (green). Data recorded in condition I.

(C) Merged AP duration across control and TTX-treated cells for recordings in IbTx-

free and IbTx-containing solution (data in A, normalized to the respective group's

median duration in IbTx-free solution).

Numbers of recorded neurons provided in brackets. Box plots depict median ±

interquartile range, whiskers reflect 10–90 percentile. P-values calculated by Mann-

Whitney-*U* test.

1141

1142

1143
1144

1145