Decision Making is Contextually Flexible

Behavior choices amongst grooming, feeding, and courting in *Drosophila* show

contextual flexibility, not an absolute hierarchy of needs

Carla E Ladd[1] and Julie H Simpson[1,2]

[1]Department of Molecular, Cellular, and Developmental Biology, University of California,

Santa Barbara, Santa Barbara, CA, United States of America

[2]Corresponding author: jhsimpson@ucsb.edu

Key Words: action selection, decision-making, behavior choice

Abstract

To determine the algorithmic rules and neural circuits controlling selection amongst

competing behaviors, we established assays where adult *Drosophila melanogaster* choose

between grooming and feeding, grooming and courting, or feeding and courting. We find

that there is not an absolute hierarchy: while flies typically perform grooming first, they can

choose to feed if sufficiently starved, or court if an appropriate female is available. Flies

alternate between competing behaviors, performing short bouts of each action rather than

completely satisfying one drive before transitioning to another. And we do not find evidence

for a common genetic or neuronal locus that affects all decisions, suggesting that the fly

may select the best course of action by discrete circuits regulating each pairwise

comparison. Our results add to a growing body of work on decision-making in *Drosophila*

and provide a foundation for future investigation of the exact neural circuits required to

achieve appropriate choices.

Summary Statement

Flies alternate between competing behaviors. The choice amongst positive actions

depends of context and relative drive levels. No central decision-making locus was

identified.

Introduction

Flexibility in decision-making is a hallmark of a successful species (Devineni and Scaplen,

2021). The best choice can change with circumstances, and an animal that can adjust its

actions according to risks and rewards, strengths of competing drives, and available

resources thrives in a range of environments. Decision-making is impaired in human

neurological conditions including Parkinson's Disease (Brand et al., 2004; Ryterska et al.,

2014), Alcohol Use Disorder (Galandra et al., 2020), and schizophrenia (Barry et al.,

2020). The behavioral rules, and genetic and neural bases of decision-making, have been

studied in humans and in various other species (Bechara et al., 1994; Lakshminarayanan et

al., 2010; Alabi et al., 2019; Rodríguez et al., 2001), but a full understanding of the control

mechanisms and an assessment of common principles has been elusive.

Drosophila melanogaster has been an advantageous animal in which to study the genes,

neurons, and circuits that contribute to perceptual decisions (DasGupta et al., 2014;

Groschner et al., 2018), food selection (Münch et al., 2022), mate choice (Hindmarsh Sten

et al., 2025; Dickson, 2008), mating duration (Gautham et al., 2024), oviposition site

preference (Joseph et al., 2009; Yang et al., 2008), and escape strategies (von Reyn et al.,

2014; von Reyn et al., 2017; Asinof and Card, 2024). Often these assays examine the trade-

off between benefit and cost: will a male fly continue to mate in the presence of a threat

(Cazale-Debat et al., 2024)? Do larvae risk hypoxia for food (Kim et al., 2017), or accept

cold or bitter food, if they are very hungry (Wu et al., 2005)? Will they stay on a mediocre

but available food source or seek richer resources (LeDue et al., 2016; de Belle et al.,

1989)? Switching amongst individual and social behaviors driven by internal mechanisms

(or at least uncontrolled stimuli) has also been assayed (Berman et al., 2016).

Flies make decisions about feeding based on how hungry they are, what specific nutrients
they lack, what food is available, and prior experience (Nhuchhen et al., 2025; Marella et
al., 2012). They can adjust meal size, select sugar or protein-rich foods, and will accept
bitter or cold but nutritious substances if sufficiently motivated (LeDue et al., 2016; de
Belle et al., 1989). And they can avoid food that previously made them sick (Stensmyr et al.,
2012). Genes and neurons that affect feeding decisions include leucokinin (Yurgel et al.,
2019), dopamine (Jin et al., 2023), diuretic hormone 31 (Lin et al., 2022), and tyramine
(Cheriyamkunnel et al., 2021).
66
Flies also make decisions about courtship and mating actions. A male fly will evaluate
whether a potential partner belongs to the same species, is male or female, and if she has
previously mated (von Philipsborn et al., 2023). He will adjust his actions based on current
sensory information and his past mating experience (Simon et al., 2011). Once he initiates
courtship, he can escalate from orientation and chasing to singing with his wings
depending on the cues from the female, and may proceed to tapping, licking, and
attempted copulation (villella and Hall, 2008). The commitment to copulation and its
duration can be influenced by perceived environmental threats: males adjust their choice
between courtship and escape depending on external sensory cues, internal drives, and
past experiences (Simon et al., 2011). Neural and genetic bases include leucokinin (Chen
et al., 2025), tyramine (Cheriyamkunnel et al., 2021), and fruitless (Yamamoto and
Koganezawa, 2013).
79
Our analysis of grooming behavior indicated that choice of which body part to clean is
driven both by the acute distribution of debris and an intrinsic set of priorities: the eyes,
antenna, and head are cleaned first (Seeds et al., 2014) – unless there is much more
stimulation to mechanosensory bristles in posterior regions of the body (Zhang et al.,
2020). An innate hierarchy organizes the grooming sequence from anterior to posterior
body parts, with contextual flexibility to respond to the current sensory load. Neural circuits

that contribute to the grooming sequence include dopamine (Pitmon et al., 2016) and

descending neurons (Guo et al., 2022).

Feeding, courting, and grooming are complex motor sequences made up of choices among

smaller actions. We noticed, anecdotally, that grooming behavior superseded escape,

courtship, and feeding in our standard grooming assays and we wondered if there was an

absolute hierarchy amongst distinct behaviors, analogous to Maslow's hierarchy of needs

(Maslow, 1943). How would a fly prioritize when given multiple options? Would this choice

be fixed or flexible?

Much popular discussion on the disadvantages of multitasking and rapid task switching led

us to examine whether flies finish one task before starting another or alternate between

activities that satisfy competing drives.

We wondered if different decisions share common neural circuits – if there is a brain region

involved in multiple decisions: a control center. We reasoned that if decision-making itself,

rather than performance of a specific behavior, were impaired, then we would see

differences in choices amongst several different behaviors. Altering neural activity in either

a central, integration, or modulatory circuit might show normal sensory response

thresholds for flies in individual behavioral assays but changes in the ratio or relative timing

of behaviors when flies were given a choice. Previous research examined decisions

between two behaviors such as courting and feeding showed roles for specific tyraminergic

neurons (Cheriyamkunnel et al., 2021) and gut hormones (Lin et al., 2022), but whether

these same neurons have a more universal effect across multiple decisions was not

known.

Therefore, we established a set of behavioral assays where the fruit fly *Drosophila*

melanogaster chooses amongst grooming, feeding, and courting (**Fig. 1**). All three of these

behaviors are natural, productive actions, and we can adjust the drive to perform each one

by manipulating internal and external cues. Flies can clean themselves and have access to

food or mates; we measured behavior onset, bout durations, and relative time spent

grooming, feeding, or courting in pairwise assays.

Methods and Materials

Fly strains

To study the behavior of wild type flies during decision making tasks, I used the CantonS

Drosophila melanogaster strain (**Table 1**). The flies were raised on common cornmeal in a

25°C incubator on a 12-hour light/dark cycle. Within 24 hours of becoming adults, the

males and females were separated into sex specific housing. Ice was used to anesthetize

flies during sorting. All lines crossed to *UAS-CsChrimson* were kept in a dark incubator and

raised on food containing 0.4 mM all-trans-retinal and starved on agar containing 0.4 mM

all-trans-retinal. All lines crossed to *UAS-Kir2.1; TubP-GAL80^{ts}* were kept at 30°C for 24

hours and then rested at room temperature (23ºC) for 1 hour prior to filming.

Table 1: Fly Strains			
Strain	**Nickname**	**Bloomington Stock ID**	**Obtained From**
CantonS	CantonS		Simpson Lab
pBDPGal4U in attP2	Empty-GAL4	68384	Simpson Lab
w[*] TI{RFP[3xP3.cUa]=2A-GAL4}Tbh[2A-GAL4.KI]/FM7a	Oct-GAL4	86145	Bloomington
w[*]; P{w[+mC] = Tdc2-GAL4.C}2	Tdc2-GAL4	9313	Bloomington
w[1118]; TI{w[+mE.hs]=GAL4}TyrR[Gal4])	TyrR-GAL4	67129	Montell Lab
Th-GAL4	Th-GAL4		Simpson Lab
Trh-GAL4 (III)	Trh-GAL4		Simpson Lab
w[1118]; P{w[+mC]=Lk-GAL4.TH}2M	LK-GAL4	51993	Bloomington

w[1118]; P{y[+t7.7] w[+mC]=GMR65C07-GAL4}attP2	LKR-GAL4	39344	Bloomington
UAS-Kir2.1; Tub-GAL80ts	UAS-Kir2.1		Louis Lab
w[1118]; P{y[+t7.7] w[+mC]=20XUAS-IVS-CsChrimson.mVenus}attP40	UAS-CsChrimson	55135	Simpson Lab
w[*]; P{w[+mW.hs]=GawB}OK107 ey[OK107]/In(4)ci[D], ci[D] pan[ciD] sv[spa-pol]	OK107-GAL4	854	Bloomington
201Y-GAL4	201Y-GAL4		Simpson Lab
w[1118]; Pbac{WH}FoxPf03746	FoxP	85667	Bloomington

Behavioral experiments

For the competition assays, all filming was done at room temperature. Flies were filmed at
30 fps for 15 minutes. Each fly was only used once in one assay. Flies were anesthetized
using ice for transferring to either dusting chambers or recording chambers, described in
(Seeds et al., 2014). All flies recovered for 15 minutes prior to filming to ensure the flies
were awake. The same recording chambers were used for every experiment; chambers
were cleaned with water after each day of experimentation. Experimental and control flies
were shaken in a consistent and reproducible manner before each filming to replicate the
effects of dusting. All males and all virgin females were 3 days old at the onset of filming.
All filming was done with a FLIR Blackfly S USB3 camera and recorded top down with an
infrared backlight.

Grooming vs Feeding
Prior to filming, flies were starved on agar ("wet-starved": deprived of food but not water).
CantonS flies were starved for 24 hours while OK107-GAL4>UAS-Kir2.1; TubPGAL80ts and

201Y>UAS-Kir2.1; TubPGAL80ts cross progeny were starved for 12 hours. Each fly was

tested alone. Each assay chamber has four separate wells, so four flies can be assayed in

each filming run. Four flies at a time were transferred to a transfer chamber to recover from

being anesthetized for 15 minutes. From there they were transferred to a 24-well corning

tissue culture plate filled with 5 mg Reactive Yellow 86 dust in each of the four center wells.

Flies were shaken in a consistent and reproducible manner to evenly coat the flies in yellow

dust. All dusting was performed under a Misonix WS-6 downflow hood. The flies were then

transferred back into the transfer chamber and then finally into the recording chamber. For

the feeding assay, a small plug of cornmeal food was added before the flies were

introduced. The plug was made by inserting the end of a Pasteur pipette into a standard

food vial and then slicing the tube into disks approximately 2 mm thick. Fresh food was

used for each experiment. As soon as the flies were added, the chamber was placed under

the camera and filming began.

Grooming vs Courting

Male flies were dusted according to the protocol outlined above and then transferred into

recording chambers loaded with virgin females, resulting in four chambers each containing

one dusted male and one clean virgin female. The virgin females were Canton S, aged 3

days, and allowed to recover for 15 minutes from anesthesia after transfer. The female flies

were not dusted because we found that the male flies did not initiate courtship to dusted

females. The flies were placed under the camera and filming began.

Courting vs Feeding

Only male flies were starved prior to filming. After being anesthetized, male flies rested in

an empty recording chamber while females rested in a recording chamber containing a

food plug as described above. Male flies were then shaken to simulate the shaking done in

the grooming experiments. The chamber containing the males was then placed back-to-

back with the chamber containing the females and the food to transfer the males into the

female and food chamber. As soon as the males were added, the chamber was placed

under the camera and filming began.

Triple

The protocol in this experiment was the same as described in the Grooming vs Feeding

protocol, except the dust-covered males were transferred to a recording chamber

containing a both a virgin female and food.

Screen

To study what types of neurons were involved in decision-making, a targeted screen was

run using subsets found during a literature hunt (Pooryasin and Fiala, 2015;

Cheriyamkunnel et al., 2021; Huser et al., 2012; Certel et al., 2010; Münch et al., 2022;

Meschi et al., 2024). The GAL4/UAS system was used for both inactivation and activation of

subsets of neurons. GAL4 males (Oct, Tdc2, TyrR, Th, Trh, LK, LKR, and Empty {Control})

were crossed with *UAS-Kir2.1; TubPGAL80ts* (inactivation) and *UAS-CsChrimson*

(activation) virgin females. The resulting flies were then tested in all three decision-making

assays.

Optogenetic activation

Prior to the optogenetic activation experiments with UAS-Chrimson, flies were raised on

0.4 mM all-trans-retinal. Within 24 hours of becoming adults, males were transferred to

fresh tubes containing their regular cornmeal food mixed with retinol while virgin females

were transferred to a plain cornmeal food tube. During starvation experiments, male flies

were transferred to agar tubes containing 0.4 mM all-trans-retinal mixed into the agar.

During the experiment, an overhead FLDR-i132LA3 ring light (626 nm) producing

approximately 0.85 mW/cm^2 of red light was used for optogenetics activation.

Data analysis

Video annotation and ethograms

All videos were annotated for behavior manually using the VCode system

(http://social.cs.uiuc.edu/projects/vcode.html) for the first five minutes. Courting, feeding,

and grooming behaviors were annotated separately (i.e., the video was watched three

times). Courting behavior was annotated for orienting, chasing, wing singing, attempted

copulation, and copulation. Feeding behavior was annotated for feeding and not feeding.

Grooming behavior was annotated for anterior and posterior grooming; walking and

standing behavior was also noted.

Ethograms showing the temporal distribution of behavior were made using MATLAB.

Grooming, feeding, and courting are displayed on separate rows grouped by the individual

fly to show instances of simultaneous behavior and to highlight the alternation pattern.

Further analysis was done to look at both comparative amounts of each behavior as well as

total behaviors. This analysis and the creation of dot plots, box plots, and line graphs were

made using Excel. Significance was calculated using a T test with two-tailed distribution

and equal variance.

Initiation was calculated as the first instance of a particular behavior. Wing singing was

used as the marker for courtship initiation because it is more specific to courtship than

chasing or orienting. Grooming initiation was not analyzed because eye cleaning always

began before the recording chambers could be placed under the camera.

Bout lengths were calculated as time a fly continuously performed an action. Any standing

or walking breaks in behavior for less than a second were disregarded. Copulation was not

included in bout length analysis.

Results

Characterization of pairwise behavioral choices

We examined how flies choose between grooming and feeding when both drives are

present (**Fig. 2**). When flies are covered in dust or debris, they groom. The mechanosensory

stimulus provides strong external drive and the behavioral response is robust. Most flies

spend 60-70% of their time grooming in a 5-minute assay. When flies are deprived of food

for at least 8 and up to 36 hours and then presented with a food source, they spend up to

80% of their time feeding (**Fig. S1**). Hunger is a strong internal drive and most flies spend

time feeding to satisfy it. There is a large amount of variability in percent time feeding: we

hypothesize that this is caused by differences in fat storage or exact time of last meal

before experimental food deprivation began. When flies are starved for 24 hours, covered in

dust, and presented with food, they alternate between feeding and grooming. The total

time spent grooming and the total time spent feeding are both reduced when compared to

performance without competition. We also noted that starvation without food does not

reduce time spent grooming, and that access to food does not reduce grooming when the

flies are not starved.

We started analysis of the decision-making behavior during competition between grooming

and feeding by first creating *ethograms*, records of the actions the flies perform over the

assayed time (**Fig. 2A**). We observe that the bout lengths of grooming shorten as feeding

competition increases (**Fig. S2**). We also observe that the fly alternates between bouts of

grooming and feeding, supporting the Alternation Model proposed in **Figure 1E**.

Quantifying total grooming time and total feeding time for each fly for the first five minutes

of observation shows individual variability but clear trends (**Fig. 2B**). Flies perform less of

each behavior when they have a choice and there is a trade-off: more feeding means less

grooming and vice versa. This is shown with data points for each individual fly in **Figure 2B**

and quantified as a population average for 10 flies in each competition condition in **Figure**

2C.

We conclude that the presence of two competing drives causes the flies to alternate,

dividing their time between behaviors to reduce both motivating drives.

Similar results occur when grooming and courting drives compete (**Fig. 3**). Clean male flies

spend on average 90% of their time courting when they share a small chamber with a

conspecific virgin female, but when the male is covered with dust, courting is reduced, and

grooming is increased. In the competitive condition, males alternate between grooming

and courting.

Starved male flies will alternate between <u>courtship and feeding</u> when given access to both

virgin females and food (**Fig. 4**). Starvation itself reduces courtship, but the total amount of

time spent courting is more significantly reduced when food is available. Most males

alternate between courting and feeding with bouts of shorter duration, rather than feeding

for a sustained period before switching to courtship.

We observe rare attempts to multitask: occasionally a fly will perform posterior grooming or

courtship singing during feeding and occasionally grooming a wing while pursuing a female.

These combinations are physically possible (**Fig. S3**) but infrequent. We also note that

when both male and female flies are covered in dust, courtship is reduced to near zero;

apparently a dusty female does not elicit male mating response. Whether this is due to

masked olfactory or pheromonal cues, altered visual appearance, or other factors is

unclear.

Behavior initiation is delayed by competing drives

When flies are presented with three competing behavior choices, they initiate grooming

first, followed by feeding, and then courting. This order could reflect priority or opportunity:

the flies are covered in dust, so they receive the mechanosensory stimulation that drives

the grooming response immediately. They usually encounter the virgin female fly moving

around in the assay chamber before they locate the food patch, but while the female is free

roaming and requires pursuit, the food is stationary and readily available. Therefore, the

order of initiation could be a consequence of which opportunity to satisfy a drive they reach

first, rather than relative drive strength or an intrinsic hierarchy of importance. The

chambers are small (15.6 mm in diameter), and the subject male fly usually encounters the

food and/or female quickly, so we believe the allocation of time spent is primarily due to

drive strength rather than just proximity.

The average time of courtship initiation is delayed by the presence of food and/or dust,

while the initiation time of feeding is delayed by dust and/or virgin females (**Fig. 5**). Thus,

measurements of change in behavior initiation time support the analysis of total time spent

in each behavior to indicate that flies faced with competing drives alternate between the

actions required to satisfy them.

As the fly's initiation of grooming, feeding, and courting actions are delayed, we also see a

reduction in bout length (**Fig. S2**). As the competing drive strength increases, the duration

of feeding and courting bouts become shorter compared to the bout lengths when no

competition is present. The feeding bout length reduced from an average of 13.56 seconds

(s) when no competition was present, to 10.36 s when a female was present and 6.30 s

when food was present. The courtship bout length (not including actual mating) reduces

from an average of 4.82 s with no competition to 1.19 s when dust was present and 1.74 s

when food was present. The grooming behavior bouts are also reduced when competition

is present, but less so than the feeding and courting bouts: when no competition is

present, the fly averages 12.45 s grooming bouts, but the presence of a female drops the

duration to 9.05 s and the presence of food drops it to 7.27 s. Together this data supports

the Alternation Model proposed in **Figure 1E** as the way flies organize decision-making

when faced with competing drives.

During the triple competition assay, feeding bout lengths dropped to 3.55 s and courting

bouts dropped to 0.59 s, but the average grooming bout length of 9.54 s had no significant

change. This suggests that grooming is less affected by competition than feeding or

courtship and may support a soft hierarchy of grooming, then feeding, then courting as

proposed in **Figure 1C**.

Candidate genes, neurons and brain regions may affect aspects of decision-making

Previous experiments implicated mutations in the FoxP transcription factor in the time

course of perceptual decisions when sensory stimuli were ambiguous (Palazzo et al.,

2020). We tested whether available *FoxP* mutations behaved differently in our pairwise

competition assays. The genetic background of the mutants affected the starvation

conditions required to balance the choice between feeding and grooming or courting, but

the primary effect we observed was an increase in the average bout duration of anterior

grooming, perhaps indicating that transitions between anterior and posterior grooming

actions was impaired (**Fig. S4**). This observation encouraged us to investigate bout duration

as another manifestation of decision-making, potentially acting over a shorter timescale

and within an overarching behavioral context. *FoxP* mutants do not show differences in

action selection within courtship. Our initial grooming bout duration results were observed

with a hypomorphic *FoxP* allele, f03746 (Mendoza et al., 2014), and when we were unable

to confirm them with the additional knock-out allele 5-SZ-3955 (DasGupta et al., 2014), we

did not pursue this gene candidate further.

A fortuitous observation in our undergraduate lab class showed that activation of the

neurons targeted by *leucokinin-GAL4* caused reduced grooming in response to dust – and

an increase in proboscis extension. This led us to test whether these neurons affected the

choice between feeding and grooming. Activation of *leucokinin-GAL4* neurons in single

behavior control assays showed a small increase in time spent feeding and a large

decrease in grooming. In our two-choice competition assays, grooming time was reduced

but neither feeding nor courtship increased (**Fig. 6**). Silencing the leucokinin-GAL4 neurons

reduced feeding in both single-behavior control assays and in competition without

impairing locomotion, grooming, or courtship. *Leucokinin-GAL4* drives expression in more

than 20 neurons in the brain and ventral nerve cord, and the behavioral effects are more

consistent with roles in individual behavior drives (Nässel, 2021) than defects in decision-

making generally.

Aminergic neurons were previously implicated in various decision tasks (Gautham et al.,

2024; Cheriyamkunnel et al., 2021), but in our competition assays, evaluated by the ratio of

time spent in each behavior, we did not observe strong or specific decision-making

phenotypes (**Fig. S5**). Further higher-resolution behavioral quantification will be needed to

determine if they have more subtle contributions to choice and alternation timing.

Finally, the mushroom bodies are a brain area involved in associating positive and negative

valence to olfactory cues in associative learning paradigms and metabolic contexts (Zhang

et al., 2007; Parnas et al., 2024; Suárez-Grimalt et al., 2024). We tested whether silencing

mushroom body neurons (using broad drivers *201Y-* and *OK107-GAL4*) would alter

grooming, feeding, or courtship in individual behavior assays or competitive choice. We

saw a strong reduction in percent time spent courting in both the individual and choice

assays (**Fig. 7**). There was an increase in time spent grooming in the grooming vs courting

choice assay, but not quite to the level of grooming that occurs when there is no competing

drive. These phenotypes are more consistent with a defect in courtship than decision-

making more generally.

Discussion

We observe variability in behavior choices among individuals. The representative

ethograms shown in **Figures 2, 3,** and **4** illustrate that onset timing, bout durations, and

alternation patterns differ, even in genetically homogeneous flies subjected to the same

experimental conditions. The trends in organization of behavior choice are statistically

significant when comparing groups of flies (histograms), but differences in drive strength

and flexibility in action selection are apparent. Difficulty achieving uniform starvation

conditions, dust distributions, and mate access may explain some of the variability, but

stochasticity in the neural mechanisms of action selection is another possible explanation

for the observed difference in individual behavioral responses. This variability in the choice

outcomes makes screening for neurons or genes with subtle effects challenging – we

would be measuring a change in the relative distribution of time spent in competing actions rather than a change from a baseline/set point. This would require higher throughput analysis, potentially possible with automatic behavior recognition methods.

Future experiments could extend our multiple-choice assay to include aggression or escape, in order to search for small numbers of neurons or genes affecting multiple behavioral choices and thus perhaps decision-making in general. While we have not yet found a locus that influenced multiple decisions, we have yet to test the central complex, another plausible candidate brain region with roles in learning and memory (Solanki et al., 2015). We did see differences in behavioral priorities when comparing CantonS and transgenic lines in the w1118 background. The w1118 flies were less robust to starvation, with a higher fraction dying at 24 hrs starved, and they showed a higher frequency of multitasking behaviors. w1118 also prioritized grooming more strongly in all competitive choice assays, indicating that genetic background can indeed influence aspects of decision-making.

Acknowledgements

We would like to acknowledge our lab mates Neil Zhang, Sunanda Marella, Li Guo, Varun Sane, Shingo Yoshikawa, Primoz Ravbar, Durafshan Sayed, and Diego De Alba for their support and continuous feedback. We thank the UCSB undergraduates who helped score behavior video: Omar Alkhairi, Dipakshi Dhinarkaran, Armaan Lebaschi, Taylor Pettway, Rose Rezaee, Rachel Tajiri, Kylie Viray, Jiaqi Wu, Sierra Gordon, Shii Lin, Meier Mai, Ryan Nishioka, Giulia Pellegrini, and Yifan Xu. We would also like to thank the Montell Lab and the Louis Lab for sharing their strains.

Competing interests

No competing interests declared.

Funding

404 This research received no specific grant from any funding agency in the public, commercial
405 or not-for-profit sectors. JHS is supported by a Brain Initiative award NS132900.
406

Data and resource availability

Data and resource availability: All relevant data and details of resources can be found

within the article and its supplementary information. Reagents available upon request.

References

Alabi, O. O., Fortunato, M. P., & Fuccillo, M. V. (2019). Behavioral Paradigms to Probe

Individual Mouse Differences in Value-Based Decision Making. *Frontiers in*

neuroscience, **13**, 50.

Asinof, S.K., and Card, G.M. (2024). Neural Control of Naturalistic Behavior Choices. *Annu*

Rev Neurosci **47**, 369-388. 10.1146/annurev-neuro-111020-094019.

Barry, A. B., Koeppel, J. A., & Ho, B. C. (2020). Impulsive decision making, brain cortical

thickness and familial schizophrenia risk. *Schizophrenia research*, **220,** 54–60.

Bechara, A., Damasio, A. R., Damasio, H., & Anderson, S. W. (1994). Insensitivity to

future consequences following damage to human prefrontal cortex. *Cognition*, **50**(1-3), 7–

15.

Berman, G.J., Bialek, W., and Shaevitz, J.W. (2016). Predictability and hierarchy in

Drosophila behavior. *Proc Natl Acad Sci U S A* **113,** 11943-11948.

10.1073/pnas.1607601113.

Brand, M., Labudda, K., Kalbe, E., Hilker, R., Emmans, D., Fuchs, G., Kessler, J., &

Markowitsch, H. J. (2004). Decision-making impairments in patients with Parkinson's

disease. *Behavioural neurology*, **15**(3-4), 77–85.

Cazale-Debat, L., Scheunemann, L., Day, M., Fernandez, D.V.A.T., Dimtsi, A., Zhang, Y.,

Blackburn, L.A., Ballardini, C., Greenin-Whitehead, K., Reynolds, E., et al. (2024).

Mating proximity blinds threat perception. *Nature* **634**, 635-643. 10.1038/s41586-024-

07890-3.

Certel, S. J., Leung, A., Lin, C. Y., Perez, P., Chiang, A. S., & Kravitz, E. A. (2010).

Octopamine neuromodulatory effects on a social behavior decision-making network in

Drosophila males. *PloS one*, *5*(10), e13248.

Chen, J., Zhu, P., Jin, S., Zhang, Z., Jiang, S., Li, S., Liu, S., Peng, Q., & Pan, Y. (2025). A

hormone-to-neuropeptide pathway inhibits sexual receptivity in

immature *Drosophila* females. *Proceedings of the National Academy of Sciences of the*

United States of America, *122*(8), e2418481122.

Cheriyamkunnel, S. J., Rose, S., Jacob, P. F., Blackburn, L. A., Glasgow, S., Moorse, J.,

Winstanley, M., Moynihan, P. J., Waddell, S., & Rezaval, C. (2021). A neuronal

mechanism controlling the choice between feeding and sexual behaviors in

Drosophila. *Current biology : CB*, *31*(19), 4231–4245.e4.

DasGupta, S., Ferreira, C.H., and Miesenbock, G. (2014). FoxP influences the speed and

accuracy of a perceptual decision in Drosophila. *Science* *344*, 901-904.

10.1126/science.1252114.

de Belle, J.S., Hilliker, A.J., and Sokolowski, M.B. (1989). Genetic localization of foraging

(for): a major gene for larval behavior in Drosophila melanogaster. *Genetics* *123*, 157-163.

10.1093/genetics/123.1.157.

Devineni, A.V., and Scaplen, K.M. (2021). Neural Circuits Underlying Behavioral Flexibility:

Insights From Drosophila. *Front Behav Neurosci* *15*, 821680. 10.3389/fnbeh.2021.821680.

Dickson, B.J. (2008). Wired for sex: the neurobiology of Drosophila mating decisions.

Science *322*, 904-909. 10.1126/science.1159276.

Galandra, C., Crespi, C., Basso, G., & Canessa, N. (2020). Impaired learning from regret

and disappointment in alcohol use disorder. *Scientific reports*, *10*(1), 12104.

Gautham, A.K., Miner, L.E., Franco, M.N., Thornquist, S.C., and Crickmore, M.A. (2024).

Dopamine biases decisions by limiting temporal integration. *Nature* *632*, 850-857.

10.1038/s41586-024-07749-7.

Groschner, L.N., Chan Wah Hak, L., Bogacz, R., DasGupta, S., and Miesenbock, G.

(2018). Dendritic Integration of Sensory Evidence in Perceptual Decision-Making. *Cell* *173*,

894-905 e813. 10.1016/j.cell.2018.03.075.

Guo, L., Zhang, N., & Simpson, J. H. (2022). Descending neurons coordinate anterior

grooming behavior in Drosophila. *Current biology : CB*, *32*(4), 823–833.e4.

Hindmarsh Sten, T., Li, R., Hollunder, F., Eleazer, S., and Ruta, V. (2025). Male-male

interactions shape mate selection in Drosophila. *Cell 188*, 1486-1503.e1425.

10.1016/j.cell.2025.01.008.

Huser, A., Rohwedder, A., Apostolopoulou, A. A., Widmann, A., Pfitzenmaier, J. E.,

Maiolo, E. M., Selcho, M., Pauls, D., von Essen, A., Gupta, T., Sprecher, S. G., Birman,

S., Riemensperger, T., Stocker, R. F., & Thum, A. S. (2012). The serotonergic central

nervous system of the Drosophila larva: anatomy and behavioral function. *PloS one*, *7*(10),

e47518.

Jin, R., Sun, S., Hu, Y., Zhang, H., & Sun, X. (2023). Neuropeptides Modulate Feeding via

the Dopamine Reward Pathway. *Neurochemical research*, *48*(9), 2622–2643.

Joseph, R.M., Devineni, A.V., King, I.F., and Heberlein, U. (2009). Oviposition preference

for and positional avoidance of acetic acid provide a model for competing behavioral drives

in Drosophila. *Proc Natl Acad Sci U S A 106*, 11352-11357.

Kim, D., Alvarez, M., Lechuga, L.M., and Louis, M. (2017). Species-specific modulation of

food-search behavior by respiration and chemosensation in Drosophila larvae. *Elife 6*.

10.7554/eLife.27057.

Lakshminarayanan VR, Chen MK, Santos LR. (2010). The evolution of decision-making

under risk: Framing effects in monkey risk preferences. *Journal of Experimental Social*

Psychology. 47:689-693.

LeDue, E. E., Mann, K., Koch, E., Chu, B., Dakin, R., & Gordon, M. D. (2016). Starvation-

Induced Depotentiation of Bitter Taste in Drosophila. *Current biology: CB*, *26*(21), 2854–

2861.

Lin, H. H., Kuang, M. C., Hossain, I., Xuan, Y., Beebe, L., Shepherd, A. K., Rolandi, M., &

Wang, J. W. (2022). A nutrient-specific gut hormone arbitrates between courtship and

feeding. *Nature*, *602*(7898), 632–638.

Marella, S., Mann, K., & Scott, K. (2012). Dopaminergic modulation of sucrose

acceptance behavior in Drosophila. *Neuron*, *73*(5), 941–950.

Maslow, A. (1943). A Theory of Human Motivation. *Psychological Review 50*, 370-396.

10.1037/h0054346.

Mendoza, E., Colomb, J., Rybak, J., Pflüger, H. J., Zars, T., Scharff, C., & Brembs, B.

(2014). Drosophila FoxP mutants are deficient in operant self-learning. *PloS one, 9*(6),

e100648.

Meschi, E., Duquenoy, L., Otto, N., Dempsey, G., & Waddell, S. (2024). Compensatory

enhancement of input maintains aversive dopaminergic reinforcement in hungry

Drosophila. *Neuron*, *112*(14), 2315–2332.

Münch, D., Goldschmidt, D., and Ribeiro, C. (2022). The neuronal logic of how internal

states control food choice. *Nature 607*, 747-755. 10.1038/s41586-022-04909-5.

Nässel, D.R. (2021). Leucokinin and Associated Neuropeptides Regulate Multiple Aspects

of Physiology and Behavior in Drosophila. *Int J Mol Sci 22*. 10.3390/ijms22041940.

Nhuchhen Pradhan, R., Montell, C., & Lee, Y. (2025). Cholesterol taste avoidance

in *Drosophila melanogaster*. *eLife*, *14*, RP106256.

Palazzo, O., Rass, M., and Brembs, B. (2020). Identification of FoxP circuits involved in

locomotion and object fixation in Drosophila. *Open Biol 10*, 200295. 10.1098/rsob.200295.

Parnas, M., Manoim, J.E., and Lin, A.C. (2024). Sensory encoding and memory in the

mushroom body: signals, noise, and variability. *Learn Mem 31*. 10.1101/lm.053825.123.

Pitmon, E., Stephens, G., Parkhurst, S. J., Wolf, F. W., Kehne, G., Taylor, M., & Lebestky,

T. (2016). The D1 family dopamine receptor, DopR, potentiates hind leg grooming behavior

in Drosophila. *Genes, brain, and behavior*, *15*(3), 327–334.

Pooryasin, A., & Fiala, A. (2015). Identified Serotonin-Releasing Neurons Induce

Behavioral Quiescence and Suppress Mating in Drosophila. *The Journal of neuroscience :*

the official journal of the Society for Neuroscience, *35*(37), 12792–12812.

Rodríguez A, Andrén H, Jansson G. (2001). Habitat-mediated predation risk and decision

making of small birds at forest edges. *Oikos*. *95*(3):383-396.

Ryterska, A., Jahanshahi, M., & Osman, M. (2014). Decision-making impairments in

Parkinson's disease as a by-product of defective cost-benefit analysis and feedback

processing. *Neurodegenerative disease management*, *4*(4), 317–327.

Seeds, A.M., Ravbar, P., Chung, P., Hampel, S., Midgley, F.M., Jr., Mensh, B.D., and

Simpson, J.H. (2014). A suppression hierarchy among competing motor programs drives

sequential grooming in Drosophila. *Elife 3*, e02951. 10.7554/eLife.02951.

Simon, J. C., Dickson, W. B., & Dickinson, M. H. (2011). Prior mating experience

modulates the dispersal of Drosophila in males more than in females. *Behavior*

genetics, *41*(5), 754–767.

Solanki, N., Wolf, R., & Heisenberg, M. (2015). Central complex and mushroom bodies

mediate novelty choice behavior in Drosophila. *Journal of neurogenetics*, *29*(1), 30–37.

Stensmyr, M.C., Dweck, H.K., Farhan, A., Ibba, I., Strutz, A., Mukunda, L., Linz, J.,

Grabe, V., Steck, K., Lavista-Llanos, S., et al. (2012). A conserved dedicated olfactory

circuit for detecting harmful microbes in Drosophila. *Cell 151*, 1345-1357.

10.1016/j.cell.2012.09.046.

Suárez-Grimalt, R., Grunwald Kadow, I.C., and Scheunemann, L. (2024). An integrative

sensor of body states: how the mushroom body modulates behavior depending on

physiological context. *Learn Mem 31*. 10.1101/lm.053918.124.

Villella, A., & Hall, J. C. (2008). Neurogenetics of courtship and mating in

Drosophila. *Advances in genetics*, *62*, 67–184.

von Philipsborn, A. C., Shohat-Ophir, G., & Rezaval, C. (2023). Courtship

Conditioning/Suppression Assays in *Drosophila*. *Cold Spring Harbor protocols*, *2023*(7),

.pdb.prot108106.

von Reyn, C.R., Breads, P., Peek, M.Y., Zheng, G.Z., Williamson, W.R., Yee, A.L.,

Leonardo, A., and Card, G.M. (2014). A spike-timing mechanism for action selection. *Nat*

Neurosci 17, 962-970. 10.1038/nn.3741.

von Reyn, C.R., Nern, A., Williamson, W.R., Breads, P., Wu, M., Namiki, S., and Card,

G.M. (2017). Feature Integration Drives Probabilistic Behavior in the Drosophila Escape

Response. *Neuron 94*, 1190-1204 e1196. 10.1016/j.neuron.2017.05.036.

Wu, Q., Zhao, Z., and Shen, P. (2005). Regulation of aversion to noxious food by Drosophila

neuropeptide Y- and insulin-like systems. *Nat Neurosci 8*, 1350-1355.

Yamamoto, D., & Koganezawa, M. (2013). Genes and circuits of courtship behaviour in

Drosophila males. *Nature reviews. Neuroscience*, *14*(10), 681–692.

Yang, C.H., Belawat, P., Hafen, E., Jan, L.Y., and Jan, Y.N. (2008). Drosophila egg-laying

site selection as a system to study simple decision-making processes. *Science 319*, 1679-

1683.

Yurgel, M. E., Kakad, P., Zandawala, M., Nässel, D. R., Godenschwege, T. A., & Keene,

A. C. (2019). A single pair of leucokinin neurons are modulated by feeding state and

regulate sleep-metabolism interactions. *PLoS biology*, *17*(2), e2006409.

Zhang, K., Guo, J. Z., Peng, Y., Xi, W., & Guo, A. (2007). Dopamine-mushroom body circuit

regulates saliency-based decision-making in Drosophila. *Science (New York,*

N.Y.), 316(5833), 1901–1904.

Zhang, N., Guo, L., and Simpson, J.H. (2020). Spatial Comparisons of Mechanosensory

Information Govern the Grooming Sequence in Drosophila. *Curr Biol 30*, 988-1001 e1004.

10.1016/j.cub.2020.01.045.

Figure Legends

Figure 1: Conceptual overview and choice assay measuring how flies select among

competing positive actions.

A) A graphical abstract of the decision-making assay. B) An outline of the assay and

analysis pipeline. C) Two models hypothesizing the presence or absence of a behavioral

hierarchy. D) Two models hypothesizing the decision-making process at an anatomical

level. E) Two models hypothesizing temporal distribution of behaviors resultant of

competing drives.

Figure 2: Grooming and feeding behaviors alternate when drives compete.

A) Ethograms depicting the first five minutes of the assay. B) A dot plot comparing the total

percent time spent grooming to the total percent time spent feeding for each fly. C) A box

plot comparing total percent time spent on each behavior.

Figure 3: Grooming and courting behaviors alternate when drives compete.

A) Ethograms depicting the first five minutes of the assay. B) A dot plot comparing the total percent time spent grooming to the total percent time spent courting for each fly. C) A box plot comparing total percent time spent on each behavior.

Figure 2: Courting and feeding behaviors alternate when drives compete.

A) Ethograms depicting the first five minutes of the assay. B) A dot plot comparing the total percent time spent courting to the total percent time spent feeding for each fly. C) A box plot comparing total percent time spent on each behavior.

Figure 5: Competing drives result in delayed behavior initiation.

A) A line graph tracking initiation of wing singing courtship behavior. The dotted line represents the 50% mark. B) A line graph tracking initiation of feeding behavior. The dotted line represents the 50% mark.

Figure 6: Inactivation and activation of leukokinin producing neurons in the three choice assays.

A) Single drive analysis of inactivated leukokinin producing neurons. B) Single drive analysis of activated leukokinin producing neurons. C) Grooming vs Feeding inactivation ethogram. D) Grooming vs Feeding activation ethogram. E) Box plot corresponding to C. F) Box plot corresponding to D. G) Grooming vs Courting inactivation ethogram. H) Grooming vs Courting activation ethogram. I) Box plot corresponding to G. J) Box plot corresponding to H. K) Courting vs Feeding inactivation ethogram. L) Courting vs Feeding activation ethogram. M) Box plot corresponding to K. N) Box plot corresponding to L.

Figure 7: Inactivation of mushroom bodies in the three choice assays.

A) Single drive analysis of inactivated mushroom bodies. B) Grooming vs Feeding inactivation ethogram. C) Grooming vs Courting inactivation ethogram. D) Courting vs

604   Feeding inactivation ethogram. E) Box plot corresponding to B. F) Box plot corresponding to

605   C. G) Box plot corresponding to D.

A

Grooming

Courting

Feeding

B

Competing Drives

Film

Score Videos

Behavior Quantification

Individual Behavior Observation (ethograms)

C

Behavioral Hierarchy

Equal Likelihood Behaviors

D

Central Control (Hub-and-spoke Model)

Distributed Choice (Net Model)

E

Completion Model

Bout

Behavior 1

Behavior 2

Bout

Time

Alternation Model

Bout

Behavior 1

Behavior 2

Bout

Time

A

Female and food

Fly 1
Fly 2
Fly 3

Starved with female and food

Fly 1
Fly 2
Fly 3

0   Minutes   5

■ Courting   ■ Feeding

B

% Total Time Feeding

% Total Time Courting

C

Percent Time %

*

Courting   Feeding

Female	+	+
Food	+	+
Starved	0 hrs	24 hrs

n = 10
* p < 0.05