Decision-Making is Contextually Flexible

Behavior choices amongst grooming, feeding, and courting in *Drosophila* show

contextual flexibility, not an absolute hierarchy of needs

Carla E Ladd[1] and Julie H Simpson[1, 2]

[1]Department of Molecular, Cellular, and Developmental Biology, University of California,

Santa Barbara, Santa Barbara, CA, United States of America

[2]Corresponding author: jhsimpson@ucsb.edu

Key Words: action selection, decision-making, behavior choice

Abstract

To determine the algorithmic rules and neural circuits controlling selection amongst

competing behaviors, we established assays where adult *Drosophila melanogaster*

choose between grooming and feeding, grooming and courting, or feeding and courting.

We find that there is not an absolute hierarchy: while flies typically perform grooming

first, they can choose to feed if sufficiently starved, or court if an appropriate female is

available. Flies alternate between competing behaviors, performing short bouts of each

action rather than completely satisfying one drive before transitioning to another. While

we did not do an exhaustive screen, from the candidates we examined, we did not find

evidence for a specific genetic or neuronal locus that affects all decisions. We did

identify genetic background effects, suggesting that multiple genes may contribute to

decision-making priorities. Our results add to a growing body of work on decision-

making in *Drosophila* and provide a foundation for future investigation of the exact

neural circuits required to achieve appropriate choices.

Summary Statement

Flies alternate between competing behaviors. The choice amongst positive actions

depends on context and relative drive levels. No central decision-making locus was

identified.

Introduction

In 1943, American psychologist Abraham Maslow published *A Theory of Human*

Motivation in which he proposed what is commonly referred to as "Maslow's Hierarchy

of Needs" (Maslow, 1943). Later depicted as a pyramid, this theory states that there are

five universal human needs. They rank, from strongest, highest priority to weakest,

lowest priority: physiological, safety, love, esteem and self-actualization. The implication

is that when needs compete, behaviors that satisfy them will be chosen in priority order.

This hierarchy serves as our framework for studying *Drosophila* decision-making in

these new choice-based behavior assays.

Decision-making impairment can be a symptom of human neurological conditions

including Parkinson's Disease (Brand et al., 2004; Ryterska et al., 2014), Alcohol Use

Disorder (Galandra et al., 2020), Major Depressive Disorder (Rinaldi et al., 2019),

Conduct Disorder (Salvatore & Dick, 2018), schizophrenia (Barry et al., 2020), and drug,

alcohol, and gambling addictions (Mizoguchi & Yamada, 2014; Lawrence et al., 2009).

Human decision-making has been studied using behavioral assays such as Gambling

Tests (Bachara et al., 1994; Rogers et al., 1999), the Dictator Game (Knafo et al., 2008),

and the Information Sampling Test (Clark et al., 2006). A better understanding of

decision-making mechanisms might lead to improved therapies.

Animal models have contributed to decision-making research. In primates, the decision-

making process has been studied using mock market economies (Lakshminarayanan et

al., 2010). In rodents, decision-making is studied using techniques such as cost-benefit

analysis (Alabi et al., 2019; Floresco et al., 2008) and judgment of visual stimuli

(Odoemene et al., 2018). Choices between feeding and parenting (Alcantara et al.,

2025), feeding and safety (de Araujo Salgado et al., 2023), and prioritization of feeding

and courtship (Burnett et al., 2019) have been investigated behaviorally and neuronally

in mice. In birds, decision-making has been studied through field observations

(Rodríguez et al., 2001). Relevant neural circuits have been investigated in pond snails

(Pirger et al., 2014), sea slugs (Gillette & Brown, 2015) and leeches (Briggman et al.,

2005; Gaudry & Kristan, 2009). The comparison of neural circuits and activity dynamics

that underlie decision-making in different animals and contexts has revealed the

complexity of this process; many open questions remain.

We were interested in whether there is a common neural locus that contributes to

decision-making more generally, and what algorithms provide flexibility when all of the

choices are beneficial. *Drosophila melanogaster* has been an advantageous model in

which to study the genes, neurons, and circuits that contribute to perceptual decisions

(DasGupta et al., 2014; Groschner et al., 2018), food selection (Münch et al., 2022),

mate choice (Hindmarsh Sten et al., 2025; Dickson, 2008), mating duration (Gautham et

al., 2024), oviposition site preference (Joseph et al., 2009; Yang et al., 2008), feeding

and courtship (Cheriyamkunnel et al., 2021), feeding and locomotion (Mann et al.,

2013), and among escape strategies (von Reyn et al., 2014; von Reyn et al., 2017;

Asinof and Card, 2024). Often these assays examine the trade-off between benefit and

cost: will a male fly continue to mate in the presence of a threat (Cazale-Debat et al.,

2024)? Do larvae risk hypoxia for food (Kim et al., 2017), or accept cold or bitter food, if

they are very hungry (Wu et al., 2005)? Will they stay on a mediocre but available food

source or seek richer resources (LeDue et al., 2016; de Belle et al., 1989)? Switching

amongst individual and social behaviors driven by internal mechanisms has also been

assayed (Berman et al., 2016).

While Maslow's theory suggests a more rigid hierarchy of priorities, flexibility in decision-

making is a hallmark of successful species (Devineni and Scaplen, 2021). The best

choice can change with circumstances, and an animal that can adjust its actions

according to risks and rewards, strengths of competing drives, and available resources,

thrives in a range of environments. Flies use contextual cues to make behavior choices:

flies make decisions about feeding based on how hungry they are, what specific

nutrients they lack, what food is available, and prior experience (Nhuchhen et al., 2025;

Marella et al., 2012). *Drosophila* can adjust meal size, select sugar or protein-rich foods,

and will accept bitter or cold but nutritious substances if sufficiently motivated (LeDue et

al., 2016; de Belle et al., 1989). Fruit flies can also avoid food that previously made

them sick (Stensmyr et al., 2012). Genes, neurons, and chemical cues that affect

feeding decisions include *leucokinin* (Yurgel et al., 2019), *dopamine* (Jin et al., 2023),

diuretic hormone 31 (Lin et al., 2022), and *tyramine* (Cheriyamkunnel et al., 2021).

Flies also make decisions about courtship and mating actions. A male fly will evaluate

whether a potential partner belongs to the same species, is male or female, and is

mated or virgin (von Philipsborn et al., 2023). Once courtship is initiated, male flies can

escalate from orientation and chasing to wing singing depending on the cues from the

female fly, and may then proceed to tapping, licking, and attempted copulation (Villella

and Hall, 2008). The commitment to copulation and its duration can be influenced by

environmental threats: males adjust their choice between courtship and escape

depending on external sensory cues, internal drives, and past experiences (Simon et

al., 2011). Neural and genetic bases include *leucokinin* (Chen et al., 2025), *tyramine*

(Cheriyamkunnel et al., 2021), and *fruitless* (Yamamoto and Koganezawa, 2013).

Our lab's analysis of grooming behavior indicated that choice of which body part to

clean is driven both by the acute distribution of debris and an intrinsic set of priorities:

the eyes, antenna, and head are cleaned first (Seeds et al., 2014) – unless there is

much more stimulation to mechanosensory bristles in posterior regions of the body

(Zhang et al., 2020). An innate hierarchy organizes the grooming sequence from

anterior to posterior body parts, with contextual flexibility to respond to the current

sensory load. Neural circuits that contribute to the grooming sequence include

dopaminergic (Pitmon et al., 2016) and descending neurons (Guo et al., 2022).

Feeding, courting, and grooming are complex motor sequences made up of choices

among smaller actions. We noticed, anecdotally, that grooming behavior superseded

escape, courtship, and feeding in our standard grooming assays and we wondered if

there was an absolute hierarchy amongst distinct behaviors, analogous to Maslow's

Hierarchy of Needs (Maslow, 1943). How would a fly prioritize when given multiple

positive options? Would this choice be fixed or flexible?

Much popular discussion on the disadvantages of multitasking and rapid task switching

led us to examine whether flies finish one task before starting another or alternate

between activities that satisfy competing drives. We also sought to determine if different

decisions share common neural circuits or are controlled independently. We

hypothesized that if decision-making itself, rather than performance of a specific

behavior, were impaired, flies could show altered choices in multiple behavioral assays.

Previous research examined decisions between two behaviors such as courting and

feeding showed roles for specific tyraminergic neurons (Cheriyamkunnel et al., 2021)

and gut hormones (Lin et al., 2022), but whether these same neurons have a more

universal effect across multiple decisions was not known.

We designed behavioral choice assays to test a fly's choices amongst grooming,

feeding, and courting (**Fig. 1A-B**). All three of these behaviors are natural, productive

actions, and the drive can be adjusted to perform each one by manipulating internal and

external cues. We measured behavior onset, bout durations, and relative time spent

grooming, feeding, or courting in pairwise assays. We proposed competing models to

describe aspects of the decision-making process in fruit flies. These models present

alternative possible answers to three questions: 1) Do fruit flies prioritize certain

behaviors (**Fig. 1C**)? 2) Do different decisions share the same mechanisms (**Fig. 1D**)?

3) How do flies allocate time between two productive actions (**Fig. 1E**)?

Methods and Materials

Fly strains

To study the behavior of wild type flies during decision making tasks, we used the

CantonS Drosophila melanogaster strain (**Table 1**). The flies were raised on a mixture of

cornmeal, molasses, yeast, and agar in a 25°C incubator on a 12-hour light/dark cycle.

Within 24 hours of becoming adults, the males and females were separated into sex

specific housing. Ice was used to anesthetize flies during sorting. All lines crossed to

UAS-CsChrimson were kept in a dark 25°C incubator and raised on food containing 0.4

mM all-trans-retinal and starved on agar containing 0.4 mM all-trans-retinal. All lines

crossed to *UAS-Kir2.1; TubP-GAL80^{ts}* were kept at 30°C for 24 hours and then rested at

room temperature (23⁰C) for 1 hour prior to filming.

Table 1: Fly Strains			
Strain	**Nickname**	**Bloomington Stock ID**	**Obtained From**
CantonS	CantonS		Simpson Lab
pBDPGal4U in attP2	Empty-GAL4	68384	Simpson Lab
w[*] TI{RFP[3xP3.cUa]=2A-GAL4}Tbh[2A-GAL4.KI]/FM7a	Oct-GAL4	86145	Bloomington
w[*]; P{w[+mC] = Tdc2-GAL4.C}2	Tdc2-GAL4	9313	Bloomington
w[1118]; TI{w[+mE.hs]=GAL4}TyrR[Gal4])	TyrR-GAL4	67129	Montell Lab
Th-GAL4	Th-GAL4		Simpson Lab
Trh-GAL4 (III)	Trh-GAL4		Simpson Lab
w[1118]; P{w[+mC]=Lk-GAL4.TH}2M	LK-GAL4	51993	Bloomington
w[1118]; P{y[+t7.7] w[+mC]=GMR65C07-GAL4}attP2	LKR-GAL4	39344	Bloomington
UAS-Kir2.1; Tub-GAL80ts	UAS-Kir2.1		Louis Lab
w[1118]; P{y[+t7.7] w[+mC]=20XUAS-IVS-CsChrimson.mVenus}attP40	UAS-CsChrimson	55135	Simpson Lab
w[*]; P{w[+mW.hs]=GawB}OK107 ey[OK107]/In(4)ci[D], ci[D]	OK107-GAL4	854	Bloomington

pan[ciD] sv[spa-pol]			
201Y-GAL4	201Y-GAL4		Simpson Lab
w[1118]; Pbac{WH}FoxPf03746	FoxP	85667	Bloomington

Behavioral experiments

Filming conditions

For all assays, filming was done at room temperature (23°C). All filming was done with a

FLIR Blackfly S USB3 camera and recorded top down with an infrared backlight at 30

fps.

Each fly was only used once in one assay. Flies were anesthetized by placing vials of

flies on ice for transferring to either dusting chambers (grooming experiments only) or

recording chambers, described in Seeds et al., 2014. All flies recovered from

anesthetization for 15 minutes prior to filming to ensure maximum alertness. The same

recording chambers were used for every experiment. All flies, male and virgin female,

were 3 days old at the onset of filming and at the same subjective time of day to control

for circadian effects.

Dusting

As described in the Seeds et al., 2014 paper, the dusting chambers consist of two 3D

printed plates with 24 cylindrical wells each with a 15.6 mm diameter. Between the two

plates lies a Nitex, 630 µm mesh sheet allowing for dust, light, and air to travel through

the wells, but does contain the fly. The wells are sealed using aluminum sliders slid

through grooves on the top plate. Anesthetized flies were placed in the four center wells,

one fly per well. To dust the flies, 5 mg of Reactive Yellow 86 dust were added to the

chambers containing alert flies. The chamber was then shaken in a reproducible

manner to allow for even coating of dust on the flies. Once shaken, excess dust was

removed by tapping the dusting chamber to allow dust to fall through the mesh. A

corresponding filming chamber was then aligned with the dusting chamber to allow for

transfer of the dusted flies. All dusting was performed under a Misonix WS-6 downflow

hood.

The filming chambers used in this experiment are made of clear fabricated plates with

four wells with a 15.6 mm diameter aligning with the four center wells of the dusting

chamber. The chambers consist of three plates with a Nitex mesh sheet between the

bottom two layers. The middle plate served as the chamber the fly resided in while the

top plate and a clear slider acted as a lid which contained the fly but could be filmed

through.

As soon as flies were transferred from the dusting chamber to the filming chamber, the

filming chamber was then placed under the camera and filming commenced.

Grooming vs feeding

Prior to filming, flies were starved on agar ("wet-starved": deprived of food but not

water). *OK107-GAL4>UAS-Kir2.1; TubPGAL80ts,201Y>UAS-Kir2.1; TubPGAL80ts*, and

their control strain cross progeny were starved for 12 hours, all other strains were

starved for 24 hours. This time length was chosen based on a starvation titration

performed on *CantonS* flies (**Fig. S1**).

Male flies were anesthetized and then placed in the dusting chambers and dusted using

the dusting protocol described above. The dusted males were transferred to a filming

chamber. In each well of the filming chamber, a small plug of cornmeal food was placed

on the center of the mesh before the flies were introduced. The plug was made by

inserting the wide end of a glass Pasteur pipette (7 mm diameter) into a standard food

vial and then slicing the tube into disks approximately 2 mm thick. Fresh food was used

for each experiment. Filming commenced as soon as flies were transferred.

Grooming vs courting

Both male and virgin females were anesthetized using ice. Males were transferred into

a dusting chamber while females were transferred into a filming chamber. After allowing

for 15 minutes recovery, the male flies were dusted using the above-described protocol.

The female flies were not dusted because we found that the male flies did not initiate

courtship to dusted females. Whether this is due to masked olfactory or pheromonal

cues, altered visual appearance, or other factors is unclear. The male flies were then

transferred directly into the chambers containing the virgin females; one male and one

virgin female per well. Once flies were transferred, the filming chambers were placed

under the camera and filming began.

Courting vs feeding

Only male flies were starved prior to filming. After being anesthetized, male flies rested

in an empty filming chamber while females rested in a filming chamber containing a

food plug as described above. Male flies were then shaken to simulate the shaking

done in the grooming experiments. The male flies were then directly transferred into the

chambers containing the female and the food. After transfer, the chamber was placed

under the camera and filming began.

Triple competition assay

The protocol in this experiment was the same as described in the courting vs feeding

protocol, except the males were dusted as in the dusting protocol prior to being

transferred to the filming chamber consisting of a virgin female and a food plug. Filming

began once flies were transferred.

Screen

To study what types of neurons were involved in decision-making, a targeted screen

was run using subsets found during a literature hunt (Pooryasin and Fiala, 2015;

Cheriyamkunnel et al., 2021; Huser et al., 2012; Certel et al., 2010; Münch et al., 2022;

Meschi et al., 2024). The GAL4/UAS system was used for both inactivation and

activation of subsets of neurons. GAL4 males (Oct, Tdc2, TyrR, Th, Trh, LK, LKR, and

Empty {Control}) were crossed with *UAS-Kir2.1; TubPGAL80ts* (inactivation) and *UAS-*

CsChrimson (activation) virgin females. The resulting flies were then tested in all three

decision-making assays. All strains in the screen had a *w1118* background.

Optogenetic activation

Prior to the optogenetic activation experiments with *UAS-Chrimson*, flies were raised on

0.4 mM all-trans-retinal. Within 24 hours of becoming adults, males were transferred to

fresh tubes containing their regular cornmeal food mixed with all-trans-retinal while

virgin females were transferred to a plain cornmeal food tube as described previously.

During starvation experiments, male flies were transferred to agar tubes containing 0.4

mM all-trans-retinal mixed into the agar. During the experiment, an overhead FLDR-

i132LA3 ring light (626 nm) producing approximately 0.85 mW/cm^2 of red light was used

for optogenetics activation. Prior to filming, flies were raised in the dark. During filming,

the red light was on consistently throughout process.

Data analysis

Video annotation

All videos were annotated for behavior manually using the VCode system

(http://social.cs.uiuc.edu/projects/vcode.html) for the first five minutes. Courting,

feeding, and grooming behaviors were annotated separately (i.e., the video was

watched three times). Courting behavior was annotated for orienting (when a male

stood facing a female within two body lengths), chasing (when a male walked towards a

female within two body lengths), wing singing (when the male extends its wing to the

side while facing the female), attempted copulation, and copulation. Feeding behavior

was annotated for feeding (proboscis extension, contact, and retraction) and not

feeding. Grooming behavior was annotated for anterior grooming (head sweeps and

front leg rubbing) and posterior grooming (back leg rubbing, abdominal sweeps, wing

cleaning, and thoracic sweeps); walking and standing behavior was also noted. Manual

annotation was performed at a frame-by-frame resolution.

Ethograms

Ethograms showing the temporal distribution of behavior were made using MATLAB.

Grooming, feeding, and courting are displayed on separate rows grouped by the

individual fly to show instances of simultaneous behavior and to highlight the alternation

pattern. Ethograms omitted walking behavior and showed all behavior subsets as their

composite behavior courting, feeding, or grooming. Further analysis was done to look at

both comparative amounts of each behavior as well as total behaviors. This statistical

analysis and the creation of dot plots, box plots, Kaplan-Meier curves, preference

indexes, and ternary plots were made using Excel.

Behavior initiation

Initiation was calculated as the first instance of a particular behavior. Wing singing, a

courtship specific behavior, was used as the marker for courtship initiation because

chasing and orienting behaviors can both be incidental of two free moving flies in a

small space. Feeding initiation was marked as the first time a male fly's proboscis

contacted food. Grooming initiation was not analyzed because eye cleaning

commences instantly and could not be captured before filming began. All flies used in

Figure 5 were *CantonS* flies. Behavior initiation was shown using a Kaplan-Meier curve.

This graph, though most used in the medical field, was selected because it tracks time

to incidence while also accounting for lack of incidence, i.e. when the fly never fed or

wing sang.

Bout length analysis

Bout lengths were calculated as time a fly continuously performed an action. Any

standing or walking breaks in behavior for less than a second were disregarded.

Copulation was not included in courtship bout length analysis. Outliers were excluded

and how many and when are noted in each figure legend. Outliers were defined as

bouts that were above the upper quartile. No more than two bouts per experiment were

excluded. All flies used in **Supplemental Figure 2** were *CantonS* flies.

Preference index

The preference index tracks which of two behaviors in the pairwise competition assays

was preferred. This calculation was done by (Behavior A - Behavior B)/(Behavior A +

Behavior B). A preference index of 1 indicates the fly performed Behavior A but never

Behavior B, an index of -1 indicates the fly performed Behavior B but never Behavior A,

and an index of 0 indicates the fly spent equal time on each behavior.

Ternary plot

Though most often used in physical sciences, we chose this plot in **Figure 8B** because

it portrays a three-dimensional preference index for the triple competition assay. Each

data point has three coordinates adding up to 100. Each coordinate indicates the

percentage of time spent on a behavior of the time spent on any of the three behaviors.

The coordinates are determined by calculating [Behavior A – (Behavior B + Behavior

C)]/(Behavior A + Behavior B + Behavior C) x 100 for each of the three behaviors for

each fly.

Statistical analysis

P values were calculated for two groups with the Mann-Whitney test. For more than two

groups, a Kruskal-Wallis test followed by a Dunn's multiple comparison test and

Bonferroni correction were performed. The p value for the Kaplan-Meier curve was

found using a log-rank analysis followed by a Bonferroni correction. The dot plots in

Figures 2B, **3B**, and **4B** serve to demonstrate variation in the population, and because

each dot conveyed time spent for two different behaviors, p values were not found for

these plots.

Results

Characterization of pairwise behavioral choices

We first established what flies do in our assay chambers when only a single behavioral

option is presented (**Fig. 2A**). When flies are covered in dust or debris, and not starved,

they groom. There is individual variability in timing and amounts (**Fig. 2B**), but the

overall grooming behavioral response is robust: most flies spend 60-80% (**Fig. 2C**) of

their time grooming, often in long bouts in a 5-minute assay. We hypothesize that the

mechanosensory stimulus from the dust provides strong external drive.

When flies are starved but not dusted, hunger is a strong internal drive and flies spend

time feeding to satisfy it (**Fig. 2A**). There is individual variability (**Fig. 2B**) but most flies

spend 20-80% of their time feeding (**Fig. 2C**), often in long bouts, in a 5-minute assay.

We hypothesize that variability is caused by differences in fat storage or exact time of

last meal before experimental food deprivation began.

We then examined how flies choose between grooming and feeding when both drives

are present. When both dusted and starved, the ethograms, records of the actions the

flies perform over the assayed time (**Fig. 2A**), show that the flies alternate between

bouts of grooming and feeding, supporting the Alternation Model proposed in **Figure**

1E. The ratio of behaviors for each individual fly is variable, as shown in **Fig 2B**, but on

average, most flies spend 40-70% of their time grooming and 5-45% of their time

feeding (**Fig. 2C**), often in shorter alternating bouts between feeding and grooming (**Fig.**

S2). Therefore, the total time spent grooming and the total time spent feeding are both

reduced when compared to performance without competition. We conclude that the

presence of two competing drives causes the flies to alternate, dividing their time

between behaviors to reduce both motivating drives.

Similar results occur when grooming and courting drives compete (**Fig. 3**). Males

alternate between grooming and courting (**Fig. 3A**). Individual flies demonstrate

variability in the courting behavior (**Fig. 3B**), which we hypothesize is due in part to the

response of free roaming virgin females. On average, clean male flies spend 90% of

their time courting (**Fig. 3C**) when they share a small chamber with a conspecific virgin

female, but when the male is covered with dust, courting is reduced, and grooming is

increased, suggesting a trade-off in time spent on each behavior.

We also see alternation in the courting vs feeding assay. Starved male flies will alternate

between courtship and feeding when given access to both virgin females and food (**Fig.**

4A). Most males alternate between courting and feeding with bouts of shorter duration

(**Fig. S2**), rather than feeding for a sustained period before switching to courtship. As in

the grooming vs feeding assay, the amount of time spent feeding is variable (**Fig. 4B**).

The total amount of time spent courting is more significantly reduced when food is

available (**Fig. 4C**).

During the three pairwise competition assays, we observe rare attempts to multitask:

occasionally a fly will perform posterior grooming during feeding or courtship singing

and occasionally feed while courtship singing. These combinations are physically

possible (**Fig. S3**) but infrequent.

Behavior initiation is delayed by competing drives

We next evaluated flies' response to competition using behavior initiation time. We

measured when they initiate courtship (**Fig. 5A**) or feeding (**Fig. 5B**) with Kaplan-Meier

curves to depict percentage of flies that have initiated the behavior at each timepoint.

(Note that we used wing-song to mark the onset of courtship because it is more specific

to courtship than orienting or following, and we could not measure grooming onset time

accurately because it always began as soon as the flies were exposed to dust and

before we could begin filming).

In all of our assays, flies predominantly initiated grooming first, followed by feeding, and

then courting. This order could reflect priority or opportunity: the flies are covered in

dust, so they receive the mechanosensory stimulation that drives the grooming

response immediately. Since the chambers are small (15.6mm in diameter), males

typically encounter both the moving female and the stationary food patch quickly.

Therefore, we believe the initiation and allocation of time spent is primarily due to

relative drive strength or priority rather than just proximity.

The average time of courtship initiation is delayed by the presence of food or dust (**Fig.**

5A), while the initiation time of feeding is delayed when more than one alternative is

present: dust and virgin females (**Fig. 5B**). In a triple competition assay (where the flies

were starved, dusted, and presented with females and food), the flies never initiated

wing singing (**Fig. 5A**) and delayed feeding (**Fig. 5B**): only 60% of flies fed at all. Thus,

measurements of change in behavior initiation time support the analysis of total time

spent in each behavior to indicate that flies faced with competing drives alternate

between the actions required to satisfy them as proposed in **Fig. 1E**.

Bout length is another temporal feature that shows the effects of competition. The
amount of time a fly spends performing each behavior is reduced as competition
increases (**Fig. S2**). Feeding and courting bouts become shorter. For example, feeding
bouts average 13.6 s without competition, 10.4 s when a female is present, 6.30 s when
food was present, and 3.55 s in triple competition. Average courtship bout length is
reduced from 4.82 s to 1.19 s when dust was present or 1.74 s when food was present
and 0.59 s when both dust and food are available. Grooming bouts are also reduced by
competition, but less so (from 12.45 s to 9.05 s with female, to 7.27 s with food, and to
9.54 s with both). Both onset time and bout length stability support a soft hierarchy of
grooming, then feeding, then courting as proposed in **Figure 1C**.
413
Candidate genes, neurons, and brain regions may affect aspects of decision-
making
Candidate genes
To further study the decision-making behavior of *Drosophila melanogaster*, we
considered candidate genes, neurons, and brain regions. Previous experiments
implicated mutations in the *FoxP* transcription factor in the time course of perceptual
decisions when sensory stimuli were ambiguous (Palazzo et al., 2020). We tested
whether available *FoxP* mutations behaved differently in our pairwise competition
assays. The genetic background of the mutants affected the starvation conditions
required to balance the choice between feeding and grooming or courting, but the
primary effect we observed was an increase in the average bout duration of anterior
grooming, perhaps indicating that transitions between anterior and posterior grooming
actions was impaired (**Fig. S4**). This observation encouraged us to investigate bout
duration as another manifestation of decision-making, potentially acting over a shorter
timescale and within an overarching behavioral context (**Fig. S4**). *FoxP* mutants do not
show differences in action selection within courtship, indicating the grooming effect is
not due to a wider switching impairment (*data not shown*). Our initial grooming bout
duration results were observed with a hypomorphic *FoxP* allele, f03746 (Mendoza et al.,
2014), and when we were unable to confirm them with the additional knock-out allele 5-
SZ-3955 (DasGupta et al., 2014), we did not pursue this gene candidate further.

Candidate neurons

A fortuitous observation in an undergraduate lab class showed that activation of the
neurons targeted by *leucokinin-GAL4* caused reduced grooming in response to dust –
and an increase in proboscis extension. This led us to test whether these neurons
affected the choice between feeding and grooming. We conducted a small screen of
candidate neuron subsets including neurons expressing leucokinin and leucokinin
receptor, tyramine and tyramine receptor (Cheriyamkunnel et al., 2021), and other
aminergic neurons (Gautham et al., 2024) (**Fig. S5**). We silenced and activated these
neuron populations in the three pairwise competition assays, corroborating the
published findings on tyraminergic neurons in a new assay and quantifying the
anecdotal observations on the neurons expressing *leucokinin-GAL4.*

Leucokinin-GAL4 drives expression in more than 20 neurons in the brain and ventral
nerve cord (Nässel, 2021). Leucokinins are a group of neuropeptides associated with
feeding – satiation and termination, sleep, nociception, and memory formation.
Leucokinin has been implicated in the interaction of hunger and noxious heat (Ohashi &
Sakai, 2018) and in sexual receptivity in juvenile females (Chen et al., 2025).

In single behavior assays, silencing Leucokinin neurons (*leucokinin-GAL4>UAS-
Kir2.1;tubGAL80ts)* results in a decrease in feeding behavior (**Fig. 6A**) while acute
activation (*leucokinin-GAL4>UAS-CsChrimson)* increased feeding and reduced
grooming (**Fig. 6B**). Activation of the neurons expressing *leucokinin-GAL4* produced
flies that had no noticeable motor deficits in grooming, courting, or feeding assays, but
continuously extended their proboscis parallel to the surface.

In pairwise competitions, we observed trends consistent with the single behavior
assays: silencing caused decreases in feeding (**Fig. 6C, E, K, M**) while activation
decreased grooming (**Fig. 6D, F, H, J**). While these neural manipulations could affect
either behavior prioritization or execution, we favor execution because we did not
observe reciprocal increases in competing behaviors. That the reduction of grooming is

not accompanied by an increase in feeding (**Fig. 6F)** or courtship (**Fig. 6J**) could be due

to a general failure to switch between behaviors or a motor deficit, we do not believe

this explanation is likely because alternation was normal in the courting vs feeding

assay (**Fig. 6L, N**).

Though there was no significant change observed in the percentage of total time spent

on courtship in either the individual or competition assays, we observed a more subtle

change in the courtship micro-behaviors (**Fig. S6**): more late-stage behaviors such as

attempted copulation and copulation were observed when leucokinin neurons were

silenced (**Fig. 6K, M**) and fewer late-stage behaviors were observed when leucokinin

neurons were activated (**Fig. 6L, N**).

Candidate brain regions

The mushroom bodies are a brain area involved in associating positive and negative

valence to olfactory cues in associative learning and memory paradigms, as well as a

wealth of other higher order adaptive behaviors (Zhang et al., 2007; Parnas et al., 2024;

Suárez-Grimalt et al., 2024; Saito et al., 2022; Kasoch et al., 2015; Chan et al., 2024).

We chose to test this brain region as a candidate for a central control region for

decision-making.

We tested whether silencing mushroom body neurons (using broad drivers *201Y-* and

OK107-GAL4) would alter grooming, feeding, or courtship in individual behavior assays

or competitive choice. We saw a strong reduction in percentage of total time spent

courting in the individual assay (**Fig. 7A**) but not the grooming vs courting or courting vs

feeding choice assays (**Fig. 7C, F, D, G**). The increase in feeding behavior (**Fig. 7A**)

observed in the single behavior assay was not observed in the competition assays (**Fig.**

7B, E, D, G). These results suggest that the changes caused by mushroom body

inactivation may be due to reduced locomotor activity rather than defects in decision-

making or changes in priority.

The effect of genetic background on the triple competition assay

The strains used in the experiments in this paper have two different genetic

backgrounds *CantonS* (**Fig. 2-5**) and *w1118* (**Fig. 6-7**). The *CantonS* flies were used to

initially characterize the decision-making behavior in the pairwise competitions. The

GAL4/UAS strains with a *w1118* background were used to target the effects of specific

genes, neuron subsets, and brain regions on decision-making. Comparing the two sets

of data we observed the *w1118* background flies were overwhelmingly grooming when

dust was introduced (**Fig. 6-7**). We know this is not due to the flies not being starved

enough or interested in the females because they performed well in the courting vs

feeding assay as well as the individual behavior assays. We hypothesize that the *w1118*

flies have a higher priority on grooming than the other behaviors, which would also

support our conclusion for the *leucokinin-GAL4* results (**Fig. 6**).

To test the effect of the genetic background, we conducted a triple competition assay

comparing *CantonS* flies and *w1118* flies (**Fig. 8A**). We observed that the *w1118* flies

had a greater preference for grooming than the other two behaviors. To plot this, we

used a ternary plot to show the ratio of time spent between the three behaviors (**Fig.**

8B). To read, follow the grey line headed towards 100 for each behavior. Resulting three

coordinates will total 100. Ex. The *w1118* outlier sits at 45 courting, 28 feeding, and 27

grooming. We also observed that the percent of total time spent grooming increased for

w1118 and feeding decreased (**Fig. 8C**). Courting remained unchanged and the flies

never performed any courting specific behaviors (orienting and chasing can be

incidental). We also looked at bout lengths for each of the three behaviors. There was

an increase in grooming bout durations in the *w1118* flies (**Fig. 8D**) and no change in

the other two behaviors (**Fig. 8E-F**). The *w1118* grooming bout length is twice as long as

the *CantonS* grooming bout length and longer than the *CantonS* grooming only behavior

as well (**Fig. S2**). This coincides with the previous remarked upon data (**Fig. 2A, 3A,**

4A, S2) where we observed that as priority lowered (competition introduced) bout length

lowered. We hypothesize, that in *w1118* we observe an increase in bout length because

there is an increase in priority.

Discussion

Characterization of pairwise decision-making

We characterized decision-making behavior in *Drosophila melanogaster* males during

pairwise behavior competition assays. We described how the internal states of hunger

and mating drive interact with external stimulation from dust in the presence of different

environmental resources such as food and virgin females to influence decision-making.

We show that flies alternate between behaviors (**Fig. 2A, 3A, 4A**): we propose that they

do this as their actions satisfy competing drives. Other explanations are possible: the

drives may decay or habituate with different time-courses, or arousal thresholds may

affect each behavior differently. Competition resulted in changes in time allocation

across behaviors (**Fig. 2C, 3C, 4C**) and delayed behavior initiation (**Fig. 5**). The

behavior alternation indicates that while flies do have a hierarchy of needs, they can

adapt their responses based on context (**Fig 1C, E**). This contextual flexibility resembles

need-based prioritization of feeding, social behaviors, and defensive responses in mice

(Burnett et al., 2013), indicating that this may be a general, adaptive mechanism across

species. We find individual variability in behavior choices, even in genetically

homogenous flies subjected to the same experimental conditions (**Fig. 2B, 3B, 4B**).

Feeding is the most variable behavior, which we propose is due differences in hunger

drive from timing of last meal before starvation period or fat storage capacity. We also

report that flies can perform two behaviors simultaneously (**Fig. S3**), suggesting that

these control circuits are not exclusive.

Effects of genes and neurons on decision-making

We tested some candidate genetic mutations (*FoxP*) and neuronal populations

(leucokinin, aminergic, and Mushroom Body neurons) for defects in decision-making. In

the *FoxP* mutants, we saw no change in decision-making overall but observed longer

bout durations of anterior grooming (**Fig. S4**), suggesting an avenue for further

investigation. Manipulating activity in the neurons expressing the leucokinin

neuropeptide affects performance of individual behaviors: silencing reduces feeding

while activating reduces grooming (**Fig. 6A, B**); these effects on individual behavior

performance likely account for the changes in the pairwise choice assays (**Fig 6E, F**

and **J**). We do not yet know whether this reflect altered drive or ability to execute. We

also observed that silencing leucokinin neurons accelerated onset of late-stage

courtship behaviors (**Fig. S6**). These results suggest that leucokinin neurons warrant

further investigation. We also tested whether activity of aminergic neurons affects

decision-making in our assays (**Fig. S5**). Our results align with previously published

work implicating tyraminergic neurons in choices between feeding and courtship.

Because the mushroom body neurons have established roles in olfactory perceptual

decision-making and learned association with valence, we tested inactivation of these

neurons in our assays (**Fig. 7**). Single behavior analysis revealed a decrease in courting

and an increase in feeding behaviors, but the competition assay between courting and

feeding showed no significant changes. We hypothesize that mushroom body

inactivation affects behavior performance rather than drive strength or decision-making,

but confirmation requires further experiments.

We expected that disruption of a central decision-making center would affect choices in

multiple assays and none of the neurons we tested produced this outcome. Therefore,

we did not uncover any evidence identifying such a hub. But we did not rule out its

existence, either. We tested a limited set of neuronal populations, so perhaps a broader

screen would uncover neurons with a more global effect. Alternatively, the expectation

of a global effect may be flawed: if decision-making is distributed or shows functional

redundancy, neurons that affect only one of our assays may still contribute to decision-

making more broadly. Our current results neither confirm nor exclude a central

mechanism for decision-making (**Fig. 1D**).

Genetic background affects decision-making

Our mutations and transgenic lines were in a different genetic background (*w1118*) from

the *CantonS* wild-type flies we used to establish the initial choice assay conditions. We

observed increased grooming in all of the assays with *w1118* flies relative to *CantonS*.

Although flies with the *w1118* background were capable of all the individual behaviors, in

the competition assays, grooming tended to dominate. To further elucidate effects of

genetic background, we tested *w1118* and *CantonS* flies in a triple competition assay:

grooming vs feeding vs courting (**Fig. 8**). We found that the *w1118* flies groomed more,

fed less, and had longer bouts of grooming, indicating that genetic variants can
influence behavioral priorities or performance. While we did not map genetic causes
contributing to these differences, our observations underscore the importance of
considering genetic background and comparing to appropriate controls.
593

Conclusion

The neural basis of decision-making remains an open and challenging area of
investigation, with new experiments representing different levels of analysis and
employing the respective strengths of different animal models (International Brain
Laboratory et al., 2025). Our experiments here further characterize decision-making
behavior in *Drosophila melanogaster*. We show that fruit flies do have a hierarchy
among the three behaviors tested but show flexibility in response to drive strengths and
available resources. We show that flies alternate among competing behaviors. And we
do not identify or exclude a central locus that affects decision-making. These assays
and results guide future exploration into both behavioral algorithms and neural circuit
mechanisms controlling how decisions are made.
605

Acknowledgements

We would like to acknowledge our lab mates Neil Zhang, Sunanda Marella, Li Guo,
Varun Sane, Shingo Yoshikawa, Primoz Ravbar, Durafshan Sayed, and Diego De Alba
for their support and continuous feedback. We thank the UCSB undergraduates who
helped score behavior video: Omar Alkhairi, Dipakshi Dhinakaran, Armaan Lebaschi,
Taylor Pettway, Rose Rezaee, Rachel Tajiri, Kylie Viray, Jiaqi Wu, Sierra Gordon,
Shuwei Lin, Meier Mai, Ryan Nishioka, Giulia Pellegrini, and Yifan Xu. We would also
like to thank the Montell Lab and the Louis Lab for sharing their strains and Dr. Sung
Soo Kim for his advice on statistical tests.
615

616 **Competing interests**

617 No competing interests declared.
618

Funding

This research received no specific grant from any funding agency in the public,

commercial or not-for-profit sectors. JHS is supported by an NIH Brain Initiative award

NS132900.

Data and resource availability

Data and resource availability: All relevant data and details of resources can be found

within the article and its supplementary information. Reagents available upon request.

References

Agrawal, A., Gans, J., & Goldfarb, A. (2019). Prediction, Judgment, and Complexity: A

Theory of Decision-Making and Artificial Intelligence. In *The Economics of Artificial*

Intelligence: An Agenda, pp. 89-110. University of Chicago Press.

Alabi, O. O., Fortunato, M. P., & Fuccillo, M. V. (2019). Behavioral Paradigms to

Probe Individual Mouse Differences in Value-Based Decision Making. *Frontiers in*

neuroscience, *13*, 50.

Alcantara, I. C., Li, C., Gao, C., Rodriguez González, S., Mickelsen, L. E., Papas, B.

N., Goldschmidt, A. I., Cohen, I. M., Mazzone, C. M., de Araujo Salgado, I., Piñol, R.

A., Xiao, C., Karolczak, E. O., Li, J. L., Cui, G., Reitman, M. L., & Krashes, M. J.

(2025). A hypothalamic circuit that modulates feeding and parenting behaviours. *Nature*,

645(8082), 981–990.

Asinof, S.K., and Card, G.M. (2024). Neural Control of Naturalistic Behavior Choices.

Annu Rev Neurosci 47, 369-388. 10.1146/annurev-neuro-111020-094019.

Barry, A. B., Koeppel, J. A., & Ho, B. C. (2020). Impulsive decision making, brain

cortical thickness and familial schizophrenia risk. *Schizophrenia research*, *220,* 54–60.

Bechara, A., Damasio, A. R., Damasio, H., & Anderson, S. W. (1994). Insensitivity to

future consequences following damage to human prefrontal cortex. *Cognition*, *50*(1-3),

7–15.

Berman, G.J., Bialek, W., and Shaevitz, J.W. (2016). Predictability and hierarchy in

Drosophila behavior. *Proc Natl Acad Sci U S A 113*, 11943-11948.

10.1073/pnas.1607601113.

Brand, M., Labudda, K., Kalbe, E., Hilker, R., Emmans, D., Fuchs, G., Kessler, J., &
Markowitsch, H. J. (2004). Decision-making impairments in patients with Parkinson's
disease. *Behavioural neurology*, **15**(3-4), 77–85.

Briggman, K. L., Abarbanel, H. D., & Kristan, W. B., Jr (2005). Optical imaging of
neuronal populations during decision-making. *Science (New York, N.Y.)*, **307**(5711),
896–901.

Burnett, C. J., Funderburk, S. C., Navarrete, J., Sabol, A., Liang-Guallpa, J.,
Desrochers, T. M., & Krashes, M. J. (2019). Need-based prioritization of behavior.
eLife, **8**, e44527.

Castells-Nobau, A., Eidhof, I., Fenckova, M., Brenman-Suttner, D. B., Scheffer-de
Gooyert, J. M., Christine, S., Schellevis, R. L., van der Laan, K., Quentin, C., van
Ninhuijs, L., Hofmann, F., Ejsmont, R., Fisher, S. E., Kramer, J. M., Sigrist, S. J.,
Simon, A. F., & Schenck, A. (2019). Conserved regulation of neurodevelopmental
processes and behavior by FoxP in Drosophila. *PloS one*, **14**(2), e0211652.

Cazale-Debat, L., Scheunemann, L., Day, M., Fernandez, D.V.A.T., Dimtsi, A.,
Zhang, Y., Blackburn, L.A., Ballardini, C., Greenin-Whitehead, K., Reynolds, E., et
al. (2024). Mating proximity blinds threat perception. *Nature* **634**, 635-643.
10.1038/s41586-024-07890-3.

Certel, S. J., Leung, A., Lin, C. Y., Perez, P., Chiang, A. S., & Kravitz, E. A. (2010).
Octopamine neuromodulatory effects on a social behavior decision-making network in
Drosophila males. *PloS one*, **5**(10), e13248.

Chan, I. C. W., Chen, N., Hernandez, J., Meltzer, H., Park, A., & Stahl, A. (2024).
Future avenues in *Drosophila* mushroom body research. *Learning & memory (Cold*
Spring Harbor, N.Y.), **31**(5), a053863.

Chen, J., Zhu, P., Jin, S., Zhang, Z., Jiang, S., Li, S., Liu, S., Peng, Q., & Pan, Y.
(2025). A hormone-to-neuropeptide pathway inhibits sexual receptivity in
immature *Drosophila* females. *Proceedings of the National Academy of Sciences of the*
United States of America, **122**(8), e2418481122.

Cheriyamkunnel, S. J., Rose, S., Jacob, P. F., Blackburn, L. A., Glasgow, S.,
Moorse, J., Winstanley, M., Moynihan, P. J., Waddell, S., & Rezaval, C. (2021). A

neuronal mechanism controlling the choice between feeding and sexual behaviors in Drosophila. *Current biology: CB*, *31*(19), 4231–4245.e4.

Clark, L., Robbins, T. W., Ersche, K. D., & Sahakian, B. J. (2006). Reflection impulsivity in current and former substance users. *Biological psychiatry*, *60*(5), 515–522.

DasGupta, S., Ferreira, C.H., and Miesenbock, G. (2014). FoxP influences the speed and accuracy of a perceptual decision in Drosophila. *Science 344*, 901-904. 10.1126/science.1252114.

de Araujo Salgado, I., Li, C., Burnett, C. J., Rodriguez Gonzalez, S., Becker, J. J., Horvath, A., Earnest, T., Kravitz, A. V., & Krashes, M. J. (2023). Toggling between food-seeking and self-preservation behaviors via hypothalamic response networks. *Neuron*, *111*(18), 2899–2917.e6.

de Belle, J.S., Hilliker, A.J., and Sokolowski, M.B. (1989). Genetic localization of foraging (for): a major gene for larval behavior in Drosophila melanogaster. *Genetics 123*, 157-163. 10.1093/genetics/123.1.157.

Devineni, A.V., and Scaplen, K.M. (2021). Neural Circuits Underlying Behavioral Flexibility: Insights From Drosophila. *Front Behav Neurosci 15*, 821680. 10.3389/fnbeh.2021.821680.

Dickson, B.J. (2008). Wired for sex: the neurobiology of Drosophila mating decisions. *Science 322*, 904-909. 10.1126/science.1159276.

Floresco, S. B., St Onge, J. R., Ghods-Sharifi, S., & Winstanley, C. A. (2008). Cortico-limbic-striatal circuits subserving different forms of cost-benefit decision making. *Cognitive, affective & behavioral neuroscience*, *8*(4), 375–389.

Galandra, C., Crespi, C., Basso, G., & Canessa, N. (2020). Impaired learning from regret and disappointment in alcohol use disorder. *Scientific reports*, *10*(1), 12104.

Gaudry, Q., & Kristan, W. B., Jr (2009). Behavioral choice by presynaptic inhibition of tactile sensory terminals. *Nature neuroscience*, *12*(11), 1450–1457.

Gautham, A.K., Miner, L.E., Franco, M.N., Thornquist, S.C., and Crickmore, M.A. (2024). Dopamine biases decisions by limiting temporal integration. *Nature 632*, 850-857. 10.1038/s41586-024-07749-7.

Gillette, R., & Brown, J. W. (2015). The Sea Slug, Pleurobranchaea californica: A
Signpost Species in the Evolution of Complex Nervous Systems and
Behavior. *Integrative and comparative biology*, *55*(6), 1058–1069.

Groschner, L.N., Chan Wah Hak, L., Bogacz, R., DasGupta, S., and Miesenbock, G.
(2018). Dendritic Integration of Sensory Evidence in Perceptual Decision-Making. *Cell*
173, 894-905 e813. 10.1016/j.cell.2018.03.075.

Guo, L., Zhang, N., & Simpson, J. H. (2022). Descending neurons coordinate anterior
grooming behavior in Drosophila. *Current biology : CB*, *32*(4), 823–833.e4.

Hindmarsh Sten, T., Li, R., Hollunder, F., Eleazer, S., and Ruta, V. (2025). Male-male
interactions shape mate selection in Drosophila. *Cell 188*, 1486-1503.e1425.
10.1016/j.cell.2025.01.008.

Huser, A., Rohwedder, A., Apostolopoulou, A. A., Widmann, A., Pfitzenmaier, J. E.,
Maiolo, E. M., Selcho, M., Pauls, D., von Essen, A., Gupta, T., Sprecher, S. G.,
Birman, S., Riemensperger, T., Stocker, R. F., & Thum, A. S. (2012). The
serotonergic central nervous system of the Drosophila larva: anatomy and behavioral
function. *PloS one*, *7*(10), e47518.

International Brain Laboratory, Angelaki, D., Benson, B., Benson, J., Birman, D.,
Bonacchi, N., Bougrova, K., Bruijns, S. A., Carandini, M., Catarino, J. A., Chapuis, G.
A., Churchland, A. K., Dan, Y., Davatolhagh, F., Dayan, P., DeWitt, E. E., Engel, T. A.,
Fabbri, M., Faulkner, M., Fiete, I. R., ... Witten, I. B. (2025). A brain-wide map of neural
activity during complex behaviour. *Nature, 645*(8079), 177–191.

Jin, R., Sun, S., Hu, Y., Zhang, H., & Sun, X. (2023). Neuropeptides Modulate Feeding
via the Dopamine Reward Pathway. *Neurochemical research*, *48*(9), 2622–2643.

Joseph, R.M., Devineni, A.V., King, I.F., and Heberlein, U. (2009). Oviposition
preference for and positional avoidance of acetic acid provide a model for competing
behavioral drives in Drosophila. *Proc Natl Acad Sci U S A 106*, 11352-11357.

Kacsoh, B. Z., Bozler, J., Hodge, S., Ramaswami, M., & Bosco, G. (2015). A novel
paradigm for nonassociative long-term memory in Drosophila: predator-induced
changes in oviposition behavior. *Genetics*, *199*(4), 1143–1157.

Kenrick, D. T., Griskevicius, V., Neuberg, S. L., & Schaller, M. (2010). Renovating the

Pyramid of Needs: Contemporary Extensions Built Upon Ancient

Foundations. *Perspectives on psychological science : a journal of the Association for*

Psychological Science, *5*(3), 292–314.

Kim, D., Alvarez, M., Lechuga, L.M., and Louis, M. (2017). Species-specific

modulation of food-search behavior by respiration and chemosensation in Drosophila

larvae. *Elife 6*. 10.7554/eLife.27057.

Knafo, A., Israel, S., Darvasi, A., Bachner-Melman, R., Uzefovsky, F., Cohen, L.,

Feldman, E., Lerer, E., Laiba, E., Raz, Y., Nemanov, L., Gritsenko, I., Dina, C.,

Agam, G., Dean, B., Bornstein, G., & Ebstein, R. P. (2008). Individual differences in

allocation of funds in the dictator game associated with length of the arginine

vasopressin 1a receptor RS3 promoter region and correlation between RS3 length and

hippocampal mRNA. *Genes, brain, and behavior*, *7*(3), 266–275.

Lakshminarayanan VR, Chen MK, Santos LR. (2010). The evolution of decision-

making under risk: Framing effects in monkey risk preferences. *Journal of Experimental*

Social Psychology. 47:689-693.

LeDue, E. E., Mann, K., Koch, E., Chu, B., Dakin, R., & Gordon, M. D. (2016).

Starvation-Induced Depotentiation of Bitter Taste in Drosophila. *Current biology:*

CB, *26*(21), 2854–2861.

Lin, H. H., Kuang, M. C., Hossain, I., Xuan, Y., Beebe, L., Shepherd, A. K., Rolandi,

M., & Wang, J. W. (2022). A nutrient-specific gut hormone arbitrates between courtship

and feeding. *Nature*, *602*(7898), 632–638.

Mann, K., Gordon, M. D., & Scott, K. (2013). A pair of interneurons influences the

choice between feeding and locomotion in Drosophila. *Neuron*, *79*(4), 754–765.

Marella, S., Mann, K., & Scott, K. (2012). Dopaminergic modulation of sucrose

acceptance behavior in Drosophila. *Neuron*, *73*(5), 941–950.

Maslow, A. (1943). A Theory of Human Motivation. *Psychological Review 50*, 370-396.

10.1037/h0054346.

Mendoza, E., Colomb, J., Rybak, J., Pflüger, H. J., Zars, T., Scharff, C., & Brembs,

B. (2014). Drosophila FoxP mutants are deficient in operant self-learning. *PloS*

one, *9*(6), e100648.

Meschi, E., Duquenoy, L., Otto, N., Dempsey, G., & Waddell, S. (2024). Compensatory enhancement of input maintains aversive dopaminergic reinforcement in hungry Drosophila. *Neuron*, *112*(14), 2315–2332.

Mizoguchi, H., & Yamada, K. (2019). Methamphetamine use causes cognitive impairment and altered decision-making. Neurochemistry international, **124**, 106–113.

Münch, D., Goldschmidt, D., and Ribeiro, C. (2022). The neuronal logic of how internal states control food choice. *Nature* **607**, 747-755. 10.1038/s41586-022-04909-5.

Nässel, D.R. (2021). Leucokinin and Associated Neuropeptides Regulate Multiple Aspects of Physiology and Behavior in Drosophila. *Int J Mol Sci* **22**. 10.3390/ijms22041940.

Nhuchhen Pradhan, R., Montell, C., & Lee, Y. (2025). Cholesterol taste avoidance in *Drosophila melanogaster*. *eLife*, *14*, RP106256.

Odoemene, O., Pisupati, S., Nguyen, H., & Churchland, A. K. (2018). Visual Evidence Accumulation Guides Decision-Making in Unrestrained Mice. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, *38*(47), 10143–10155.

Ohashi, H., & Sakai, T. (2018). Leucokinin signaling regulates hunger-driven reduction of behavioral responses to noxious heat in Drosophila. *Biochemical and biophysical research communications*, *499*(2), 221–226.

Palazzo, O., Rass, M., and Brembs, B. (2020). Identification of FoxP circuits involved in locomotion and object fixation in Drosophila. *Open Biol 10*, 200295. 10.1098/rsob.200295.

Parnas, M., Manoim, J.E., and Lin, A.C. (2024). Sensory encoding and memory in the mushroom body: signals, noise, and variability. *Learn Mem 31*. 10.1101/lm.053825.123.

Pirger, Z., Crossley, M., László, Z., Naskar, S., Kemenes, G., O'Shea, M., Benjamin, P. R., & Kemenes, I. (2014). Interneuronal mechanism for Tinbergen's hierarchical model of behavioral choice. *Current biology : CB*, *24*(17), 2018–2024.

Pitmon, E., Stephens, G., Parkhurst, S. J., Wolf, F. W., Kehne, G., Taylor, M., & Lebestky, T. (2016). The D1 family dopamine receptor, DopR, potentiates hind leg grooming behavior in Drosophila. *Genes, brain, and behavior*, *15*(3), 327–334.

Pooryasin, A., & Fiala, A. (2015). Identified Serotonin-Releasing Neurons Induce Behavioral Quiescence and Suppress Mating in Drosophila. *The Journal of*

neuroscience : the official journal of the Society for Neuroscience, *35*(37), 12792–

12812.

Rinaldi, R., Lefebvre, L., Joachim, A., & Rossignol, M. (2020). Decision-making of

patients with major depressive disorder in the framework of action control. Cognitive

neuropsychiatry, **25**(1), 71–83.

Rodríguez A, Andrén H, Jansson G. (2001). Habitat-mediated predation risk and

decision making of small birds at forest edges. *Oikos.* *95*(3):383-396.

Rogers, R. D., Everitt, B. J., Baldacchino, A., Blackshaw, A. J., Swainson, R.,

Wynne, K., Baker, N. B., Hunter, J., Carthy, T., Booker, E., London, M., Deakin, J. F.,

Sahakian, B. J., & Robbins, T. W. (1999). Dissociable deficits in the decision-making

cognition of chronic amphetamine abusers, opiate abusers, patients with focal damage

to prefrontal cortex, and tryptophan-depleted normal volunteers: evidence for

monoaminergic mechanisms. *Neuropsychopharmacology : official publication of the*

American College of Neuropsychopharmacology, *20*(4), 322–339.

Ryterska, A., Jahanshahi, M., & Osman, M. (2014). Decision-making impairments in

Parkinson's disease as a by-product of defective cost-benefit analysis and feedback

processing. *Neurodegenerative disease management*, *4*(4), 317–327.

Saitoe, M., Naganos, S., Miyashita, T., Matsuno, M., & Ueno, K. (2022). A non-

canonical on-demand dopaminergic transmission underlying olfactory aversive

learning. *Neuroscience research*, *178*, 1–9.

Salvatore, J. E., & Dick, D. M. (2018). Genetic influences on conduct disorder.

Neuroscience and biobehavioral reviews, **91**, 91–101.

Seeds, A.M., Ravbar, P., Chung, P., Hampel, S., Midgley, F.M., Jr., Mensh, B.D., and

Simpson, J.H. (2014). A suppression hierarchy among competing motor programs

drives sequential grooming in Drosophila. *Elife* *3*, e02951. 10.7554/eLife.02951.

Simon, J. C., Dickson, W. B., & Dickinson, M. H. (2011). Prior mating experience

modulates the dispersal of Drosophila in males more than in females. *Behavior*

genetics, *41*(5), 754–767.

Solanki, N., Wolf, R., & Heisenberg, M. (2015). Central complex and mushroom

bodies mediate novelty choice behavior in Drosophila. *Journal of neurogenetics*, *29*(1),

30–37.

Stensmyr, M.C., Dweck, H.K., Farhan, A., Ibba, I., Strutz, A., Mukunda, L., Linz, J.,

Grabe, V., Steck, K., Lavista-Llanos, S., et al. (2012). A conserved dedicated olfactory

circuit for detecting harmful microbes in Drosophila. *Cell 151*, 1345-1357.

10.1016/j.cell.2012.09.046.

Suárez-Grimalt, R., Grunwald Kadow, I.C., and Scheunemann, L. (2024). An

integrative sensor of body states: how the mushroom body modulates behavior

depending on physiological context. *Learn Mem 31*. 10.1101/lm.053918.124.

van Alphen, B., Semenza, E. R., Yap, M., van Swinderen, B., & Allada, R. (2021). A

deep sleep stage in *Drosophila* with a functional role in waste clearance. *Science*

advances, *7*(4), eabc2999.

Villella, A., & Hall, J. C. (2008). Neurogenetics of courtship and mating in

Drosophila. *Advances in genetics*, *62*, 67–184.

von Philipsborn, A. C., Shohat-Ophir, G., & Rezaval, C. (2023). Courtship

Conditioning/Suppression Assays in *Drosophila. Cold Spring Harbor protocols*, *2023*(7),

.pdb.prot108106.

von Reyn, C.R., Breads, P., Peek, M.Y., Zheng, G.Z., Williamson, W.R., Yee, A.L.,

Leonardo, A., and Card, G.M. (2014). A spike-timing mechanism for action selection.

Nat Neurosci 17, 962-970. 10.1038/nn.3741.

von Reyn, C.R., Nern, A., Williamson, W.R., Breads, P., Wu, M., Namiki, S., and

Card, G.M. (2017). Feature Integration Drives Probabilistic Behavior in the Drosophila

Escape Response. *Neuron 94*, 1190-1204 e1196. 10.1016/j.neuron.2017.05.036.

Wu, Q., Zhao, Z., and Shen, P. (2005). Regulation of aversion to noxious food by

Drosophila neuropeptide Y- and insulin-like systems. *Nat Neurosci 8*, 1350-1355.

Yamamoto, D., & Koganezawa, M. (2013). Genes and circuits of courtship behaviour

in Drosophila males. *Nature reviews. Neuroscience*, *14*(10), 681–692.

Yang, C.H., Belawat, P., Hafen, E., Jan, L.Y., and Jan, Y.N. (2008). Drosophila egg-

laying site selection as a system to study simple decision-making processes. *Science*

319, 1679-1683.

Yurgel, M. E., Kakad, P., Zandawala, M., Nässel, D. R., Godenschwege, T. A., &

Keene, A. C. (2019). A single pair of leucokinin neurons are modulated by feeding state

and regulate sleep-metabolism interactions. *PLoS biology*, *17*(2), e2006409.

Zhang, K., Guo, J. Z., Peng, Y., Xi, W., & Guo, A. (2007). Dopamine-mushroom body

circuit regulates saliency-based decision-making in Drosophila. *Science (New York,*

*N.Y.), **316**(5833), 1901–1904.

Zhang, N., Guo, L., and Simpson, J.H. (2020). Spatial Comparisons of

Mechanosensory Information Govern the Grooming Sequence in Drosophila. *Curr Biol*

***30**, 988-1001 e1004. 10.1016/j.cub.2020.01.045.

Figure Legends

Figure 1: Conceptual overview and choice assay measuring how flies select

among competing positive actions.

A) A graphical abstract of the decision-making assay. B) An outline of the assay and

analysis pipeline. C) Two models proposing the presence or absence of a behavioral

hierarchy. D) Two models representing the decision-making process at an anatomical

level: central vs distributed decision-making circuits. E) Two models hypothesizing about

temporal distribution of behaviors resulting from competing drives: alternation vs

sequential completion.

Figure 2: Grooming and feeding behaviors alternate when drives compete.

A) Ethograms depicting the first five minutes of the grooming vs feeding assay. Male

flies are provided with food and are starved or dusted as indicated. B) Dot plot

comparing the percent of total time spent grooming to the percent of total time spent

feeding for each individual fly. C) Box plot comparing the average time spend for all

flies. X denotes mean, bar denotes median, and whiskers denote the lower and upper

quartile. Statistical significance was assessed using a Kruskal-Wallis test followed by a

Dunn's multiple comparison test and Bonferroni correction.

Figure 3: Grooming and courting behaviors alternate when drives compete.

A) Ethograms depicting the first five minutes of the grooming vs courting assay. Male

flies were raised in sex isolation and presented with a virgin female. Dust is present

when indicated. B) Dot plot comparing percent of total time spent grooming to the

percent of total time spent courting for each fly. C) Box plot comparing percent of total

time spent on each behavior. X denotes mean, bar denotes median, and whiskers

denote the lower and upper quartile. Statistical significance was assessed using a

Mann-Whitney test.

Figure 4: Courting and feeding behaviors alternate when drives compete.

A) Ethograms depicting the first five minutes of the courting vs feeding assay. Male flies

have been starved and raised in sex isolation and have been presented with a virgin

female and food. B) Dot plot comparing the percent of total time spent courting to the

percent of total time spent feeding for each fly. C) Box plot comparing percent of total

time spent on each behavior. X denotes mean, bar denotes median, and whiskers

denote the lower and upper quartile. Statistical significance was assessed using a

Mann-Whitney test.

Figure 5: Competing drives result in delayed behavior initiation.

A) A line graph depicting a Kaplan-Meier curve tracking initiation of wing singing

courtship behavior. The dotted line represents the 50% mark. B) A line graph depicting a

Kaplan-Meier curve tracking initiation of feeding behavior. The dotted line represents the

50% mark. Statistical significance was assessed using a log rank analysis and a

Bonferroni correction.

Figure 6: Inactivation and activation of leucokinin producing neurons in the three

choice assays.

A) Single drive analysis of inactivated leucokinin producing neurons. B) Single drive

analysis of activated leucokinin producing neurons. C) Grooming vs feeding inactivation

ethograms. D) Grooming vs feeding activation ethograms. E) Box plot corresponding to

C. F) Box plot corresponding to D. G) Grooming vs courting inactivation ethograms. H)

Grooming vs courting activation ethograms. I) Box plot corresponding to G. J) Box plot

corresponding to H. K) Courting vs feeding inactivation ethograms. L) Courting vs

feeding activation ethograms. M) Box plot corresponding to K. N) Box plot

corresponding to L. In the box plots, the X denotes mean, bar denotes median, and

whiskers denote the lower and upper quartile. Statistical significance was assessed

using a Mann-Whitney test. All figures are n = 10.

Figure 7: Inactivation of mushroom bodies in the three choice assays.

A) Single drive analysis of inactivated mushroom bodies. B) Grooming vs feeding

inactivation ethograms. C) Grooming vs courting inactivation ethograms. D) Courting vs

feeding inactivation ethograms. E) Box plot corresponding to B. F) Box plot

corresponding to C. G) Box plot corresponding to D. In the box plots, the X denotes

mean, bar denotes median, and whiskers denote the lower and upper quartile.

Statistical significance was assessed using a Mann-Whitney test. All figures are n = 10.

Figure 8: Effects of genetic background on the triple competition assay.

A) Ethograms depicting *CantonS* and *w1118* flies in the grooming vs feeding vs courting

triple competition assay. B) Ternary plot of individual *CantonS* and *w1118* flies depicting

percent of time spent among three competing behaviors totaling 100%. C) Box plots

comparing percent of total time spent on grooming, feeding, and courting during the first

five minutes of the assay. D) Box plot with a dot plot overlay depicting the length of

individual bouts of grooming during the triple competition assay. E) Box plot with a dot

plot overlay depicting the length of individual bouts of feeding during the triple

competition assay. F) Box plot with a dot plot overlay depicting the length of individual

bouts of courting during the triple competition assay. [†]2 outlying bouts were removed [‡]1

outlying bout was removed. In the box plots, the X denotes mean, bar denotes median,

and whiskers denote the lower and upper quartile. Statistical significance was assessed

using a Mann-Whitney test.

A

Grooming

Courting

Feeding

B

Competing Drives

Film

Score Videos

Behavior Quantification

Individual Behavior Observation (ethograms)

C

Behavioral Hierarchy

Equal Likelihood Behaviors

D

Central Control (Hub-and-spoke Model)

Distributed Choice (Net Model)

E

Completion Model

Bout

Behavior 1

Behavior 2

Bout

Time

Alternation Model

Bout

Behavior 1

Behavior 2

Bout

Time

A

Dusted with food

Fly 1
Fly 2
Fly 3
Fly 4
Fly 5
Fly 6
Fly 7
Fly 8
Fly 9
Fly 10

Dusted and starved with food

Fly 1
Fly 2
Fly 3
Fly 4
Fly 5
Fly 6
Fly 7
Fly 8
Fly 9
Fly 10

Starved with food

Fly 1
Fly 2
Fly 3
Fly 4
Fly 5
Fly 6
Fly 7
Fly 8
Fly 9
Fly 10

0   Minutes   5

■ Grooming   ■ Feeding

B

% Total Time Feeding vs % Total Time Grooming

C

Percent Time %

Grooming   Feeding

	●	●	○	
Dust	+	+	−	n = 10
Food	+	+	+	* p < 0.05
Starved	0 hrs	24 hrs	24 hrs	** p < 0.01
				*** p < 0.001

A Wing Singing Initiation				
Dust	-	-	+	+
Food	-	+	-	+
Female	+	+	+	+
Starved	24 hrs	24 hrs	24 hrs	24 hrs

n = 10

B Feeding Inititation				
Dust	-	-	+	+
Food	+	+	+	+
Female	-	+	-	+
Starved	24 hrs	24 hrs	24 hrs	24 hrs

n = 10
* p < 0.05
** p < 0.01
*** p < 0.001