Differential Decay of Multiple eNA Components from a Cetacean

Pedro FP Brandão-Dias[1*]; Megan Shaffer[1]; Gledis Guri[1]; Kim M Parsons[2]; Ryan P Kelly[1]; Elizabeth Andruszkiewicz Allan[1]

1: School of Marine and Environmental Affairs, University of Washington, Seattle, 98105, Washington, USA

2: Conservation Biology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, 98112, Washington, USA

*: Corresponding author. E-mail: pedrobdfp@gmail.com

ABSTRACT

Environmental nucleic acids (eNA), such as DNA and RNA, are powerful tools for monitoring biodiversity. Yet, interpreting eNA detections requires understanding of their environmental persistence. We conducted a decay experiment using seawater from an open enclosure to track degradation of six eNA components derived from *Tursiops truncatus*: mitochondrial eDNA of varying lengths, ribosomal eRNA, and messenger eRNA. Targets were quantified over seven days via digital droplet PCR (ddPCR). Decay followed a biphasic exponential model with rapid initial loss (~24 hours at 15 °C), followed by slower degradation. Cytb messenger eRNA was least stable, disappearing within four hours. Ribosomal eRNA persisted longer but degraded slightly faster than its eDNA counterpart ($\lambda_1 = 0.236$ vs. 0.165 hr^{-1}). Longest eDNA fragments decayed more rapidly ($\lambda_1 = 0.190$ hr^{-1}) than shorter ones ($\lambda_1 = 0.114$ hr^{-1}). These findings support the use of eDNA fragment length as a proxy for degradation state and reinforce that combining multiple eNA components with distinct stabilities can potentially provide a molecular clock for inferring eNA age. This approach improves the spatiotemporal resolution of eNA-based monitoring, particularly for rare marine-mammal that act as point sources. We also emphasize the importance of explicitly distinguishing between RNA types (ribosomal vs. messenger) in environmental studies, given their divergent stability and interpretability.

KEYWORDS: Removal, ddPCR, Bayesian, Experimental, Marine Mammal

INTRODUCTION

Environmental DNA (eDNA) and, more recently, environmental RNA (eRNA)—collectively referred to as environmental nucleic acids (eNA)—have emerged as powerful tools for ecological monitoring and biodiversity assessments [1,2]. Unlike traditional direct observation methods, eNA-based surveys rely on indirect detection of species through trace amounts of genetic material that organisms release into their environment [3]. Due to the eNA accumulation in space, eNA surveys are fundamentally integrative, containing genetic information accumulated over time rather than providing an instantaneous snapshot. This integrative property enhances detection sensitivity compared to nets or visual surveys which may integrate over space but reflect a single point in time; that is, they are snapshot-based methods, making eNA detection analogous to a short movie rather than a single photographic frame. However, the extent to which eNA surveys integrate species presence over both time and space depends on environmental and molecular factors that govern its persistence and movement, thus making the integration variable [4].

Accurately interpreting eNA detections requires understanding how much time and space is integrated into each detection event. This, in turn, demands detailed knowledge about the origin, state, transport, and persistence (or decay) of nucleic acids within the environment, collectively referred to as the ecology of eNA [3,5]. Among these processes, the decay rate of eNAs has been among the most frequently estimated parameters [6–8]. The prominence of decay rate studies stems not only from the relative ease of estimating decay rates compared to transport or production, but also from the fundamental role decay plays in determining the temporal resolution of eNA surveys. While transport primarily governs the spatial resolution of detections [9], largely shaped by hydrological, oceanographic, or atmospheric processes, decay dictates how long genetic signals remain detectable, helping set the temporal bounds of the eNA movie, which reflects a steady-state balance among production, transport, and degradation.

Beyond simply establishing an upper temporal limit for eNA detection, certain applications require more precise aging of the biological source of detections within a sample. This is particularly relevant for point sources or rare detections, such as those of marine mammals [10] or invasive species [11]. To address this, previous research has proposed using the ratio of eRNA to eDNA as a "molecular clock" for eNA persistence [12]. Since eRNA generally degrades faster than eDNA, qualitatively, a sample with a high proportion of eRNA to eDNA suggests a recent biological source, whereas a sample containing only eDNA indicates an older signal [8,12]. This is analogous to applications in the forensic literature, where the ratio of presence between messenger RNA and DNA, as well as the degradation state of RNA, can be reliably used to infer time since death [13].

Nonetheless, this framework extends beyond eDNA/eRNA ratios to any eNA
component with distinct degradation rates. Environmental nucleic acids exist in multiple
molecular forms, including intracellular, particle-adsorbed, dissolved, nuclear,
mitochondrial, and other fractions [14–16]. Provided these components decay at different rates,
then their relative proportions over time and space provide a molecular signature for
estimating detection age [17]. Naturally, this approach can be applied not only to molecular
forms of eDNA, but also to experimentally defined fractions such as fragment lengths
assessed through multiple genetic markers [18,19], variations in eNA particle size distributions
captured by sequential filtration [20,21], differences between eNA extracted from distinct
environmental media, such as sediment versus water [22], eDNA to eRNA ratios [12], among
many others.

In theory, the most effective molecular components for estimating detection age are
those with strongly contrasting decay rates, as their relative proportions shift measurably
over short timescales, thus enhancing spatiotemporal resolution. However, in practice, the
choice of eNA components must balance temporal sensitivity with logistical and
methodological feasibility. Numerous studies have measured eNA decay rates, consistently
highlighting that these rates are highly context dependent. Decay dynamics vary
substantially with environmental conditions such as biological activity and temperature [23,24],
depend on target organism [25], and differ significantly due to methodological differences such
as filter pore size and molecular marker length [17,20,26]. Consequently, generalizing decay rates
across systems or conditions is challenging, especially since controlled experimental
setups seldom fully represent the complexities of natural environments. As a result,
accurate molecular time inference requires decay estimates tailored to or closely
approximating the specific environmental conditions and molecular targets of interest.

In freshwater systems, multiple studies have compared eRNA and eDNA decay,
generally finding that eRNA—particularly messenger RNA (emRNA)—decays more rapidly
than eDNA [27,28]. In marine environments, however, relatively few studies have directly
compared eRNA and eDNA decay. For example, Qian et al.[29] observed significantly faster
decay of prawn-derived emRNA compared to eDNA, particularly under colder temperatures.
Conversely, Wood et al.[30] examined decay rates of worm- and tunicate-derived emRNA,
finding only slightly elevated decay rates compared to eDNA. Similarly, Scriver et al. [31] found
no significant difference in decay between emRNA and eDNA from marine worms. To date,
no studies have directly assessed decay rates of ribosomal eRNA (erRNA) in marine
environments, although Miyata et al.[32] noted that erRNA yielded more ecologically relevant
metabarcoding detections than eDNA, suggesting greater transience and thus potentially
faster decay. Crucially, despite this emerging body of work on eRNA, no studies have yet

evaluated the decay rates of eNA components, eDNA or eRNA, specifically derived from marine mammals.

Here, we experimentally assessed the differential decay rates of multiple eNA components from the common bottlenose dolphin, *Tursiops truncatus* (Montagu, 1821). Using seawater sourced from an open environment, netted dolphin enclosure, we conducted a controlled decay experiment, tracking the persistence of eDNA of several molecular lengths, emRNA, and erRNA over time (Fig. 1). We hypothesized that eRNA would degrade faster than eDNA, with emRNA decaying the fastest due to its transient nature. We expected erRNA to persist longer than emRNA due to its structural properties, but still degrade more rapidly than eDNA. Lastly, we anticipated that larger DNA fragments would decay faster than shorter ones, following previous findings [20,33] and known fragmentation patterns. By quantifying these decay rates, we aimed to improve the temporal resolution of marine mammal detections and refine our understanding of eNA persistence in marine environments.

Figure 1: Overview of the experimental design and the four mitochondrial targets quantified. (a) Water-decay experiment. Seawater was scooped from inside an open-net enclosure housing an Atlantic bottlenose dolphin (*Tursiops truncatus*) and poured into three

25 L carboys (biological replicates). A fourth carboy containing de-ionized (DI) water served
as cross-contamination control. Carboys were held in a 15 °C environmental chamber with
continuous aeration and a 16 h light / 8 h dark cycle. Two-liter subsamples were withdrawn
at nine timepoints (0–188 h) for size-fractionated filtration and downstream DNA/RNA
extraction. (b) Locations of the mitochondrial assays. Arrows show the relative positions of
the 79 bp Cytb amplicon (protein-coding), the 390 bp D-loop amplicon (non-coding control
region), and the 146 S rRNA amplicon (ribosomal) on the circular dolphin mtDNA. The
2746 bp segment spanning D-loop to 16 S (black arc labelled "Bridge") represents the
minimum intact bridging fragment between 16S and Dloop loci. Figure created in
https://BioRender.com.

METHODS

Environmental DNA source and decay experiment

A small, managed group of non-native Atlantic bottlenose dolphins (*Tursiops*
truncatus) is housed in a defined area along the eastern bank of Hood Canal, Washington,
as part of the U.S. Navy's Marine Mammal Program. These dolphins frequently inhabit netted
enclosures adjacent to a pier, where seawater freely exchanges with the surrounding marine
environment. In October 2024 we collected water from within one of these netted
enclosures containing a single dolphin at the time of sampling. Using a 2-liter pitcher, we
filled three replicate 25-liter carboys, which served as the three biological replicates for the
decay study (Fig. 1a). To minimize contamination, collectors wore gloves during sampling,
and both the pitcher and carboys had been cleaned overnight with 1% bleach and
extensively rinsed with deionized water before use.

After collection, water within each carboy was mixed, and a 2-liter baseline sample
was filtered from each carboy to assess initial eNA concentrations before transport. These
samples were immediately filtered and preserved (see below for filtration and preservation
details). The carboys were then sealed and transported at ambient temperature to the
laboratory and an environmental chamber for the decay experiment. Temperature loggers
placed inside one of the carboys recorded transport temperatures ranging from 16°C to 12°C
over the course of ~3 hours of transport back to the laboratory.

At the laboratory, the three biological replicate carboys were decontaminated
externally with bleach before being placed inside an environmental chamber, where they
remained for the duration of the experiment. A fourth carboy, filled with deionized (DI) water,
was also placed in the environmental chamber as a cross-contamination control. The
environmental chamber was maintained at a constant 15°C with a 16-hour daylight cycle

throughout the experiment. To ensure water circulation, we inserted air hoses with stones
powered by a Whisper 100 (Tetra) into each carboy, in addition to a U-shaped plastic
sampling hose, both of which remained in place for the entire experiment. Like the carboys,
each of those were cleaned by overnight 1% bleach bath and extensively rinsed with
deionized water before use.

To evaluate the decay of eNA over time, 2-liter water samples were collected from
each carboy at nine timepoints: 0, 4, 8, 12, 24, 48, 96, 140, and 188 hours (Fig. 1a). At each
timepoint, aeration was temporarily turned off to prevent aerosolization, and water was
drawn via the plastic sampling hoses. We used sequential filtration to capture eDNA across
different particle sizes. Each sample was filtered through a tandem system of three filter
housings (Smith-Root, non-self-preserving) equipped with 5 µm, 1.0 µm, and 0.45 µm mixed
cellulose ester (MCE) 47 mm filters (Advantec), each backed by polyester drain disks
(Sterlitech). Filters were connected downstream to an eDNA Citizen Science Sampler
(Smith-Root) for vacuum filtration. Immediately after filtration, filter housings were opened,
and sterilized forceps were used to fold each filter twice, sample side in, before placing it
into 5 mL Lo-Bind tubes (Eppendorf) containing 2 mL of DNA/RNA Shield (Zymo). Samples
were incubated at room temperature for 30 minutes, then frozen at -80°C until DNA
extraction within one month.

To prevent cross-contamination, all forceps, filter holders, adapters, and tubing were
submerged in 5% bleach baths, thoroughly rinsed with deionized (DI) water, and dried before
reuse. Two independent sets of filtration equipment were available, allowing for alternating
use between timepoints. Additionally, at the end of each timepoint, we filtered 2 liters of the
DI water used for post-bleach rinsing through a 0.45 µm filter as a contamination control to
verify the effectiveness of our decontamination procedures.

DNA and RNA extraction

Nucleotides were extracted from samples using a two-step process. First, frozen
filters with buffer were thawed and incubated with agitation at 37 °C for ~15 minutes to
dissolve any precipitation. The samples were then thoroughly vortexed, and the buffer (2 mL)
was transferred to an Amicon Ultra-15 30 kDa centrifugal filter unit (Millipore-Sigma) and
centrifuged at 4,000 rpm for 30 minutes, concentrating the buffer to ~400 µL. Both eDNA and
eRNA were extracted simultaneously using the Quick DNA/RNA Miniprep Kit (Zymo
Research), following the manufacturer's protocol with a 30-minute proteinase K incubation
before adding the binding buffer.

This extraction process results in two sequential extracts (Fig. S2a). The first primarily
contains the DNA (hereafter "eDNA"), whereas the second is expected to contain RNA;
however, empirical results showed that it also contained a substantial amount of DNA,
hereafter referred to as "carryover eDNA". One-third of this second extract —containing both
eRNA and carryover eDNA— was retained for carryover eDNA analysis (see supplement,
Figs. S2, S3), while the remaining two-thirds were treated with ezDNAse (ThermoFisher
Scientific) at 37 °C for 30 minutes to remove the carryover DNA in preparation for cDNA
synthesis.

The DNAse-treated portion of the extract was subjected to a SuperScript IV
(ThermoFisher Scientific) first-strand cDNA synthesis reaction using random hexamers in
the denaturation step following the manufacturer's protocol, to convert RNA into cDNA. This
product is hereafter referred to as "eRNA." Each sample was also subjected to a second
control reaction, identical in components and cycling conditions, but without the reverse
transcriptase enzyme. This reaction product is hereafter referred to as the "No-RT," and it was
used to assess post-DNAse treatment carryover eDNA, ensuring that quantifications from
the eRNA extract were from eRNA-derived cDNA rather than residual carryover eDNA. For
details on carryover eDNA analysis, see the supplementary material.

Target eNA quantification

Target eNA were quantified using three assays, all of which target mitochondrial loci
(Table 1, Table S1). The first was a 79 bp fragment of *Cytochrome b* (Cytb), a protein-coding
gene, from which we quantified three components: eDNA, messenger eRNA (hereafter
emRNA), and the No-RT control. The second was a 146 bp fragment of the *16S* ribosomal
RNA gene, used to quantify eDNA, ribosomal eRNA (erRNA), and No-RT control. The third
was a 390 bp fragment of the *D-loop* control region, from which only eDNA was analyzed as
it is a non-coding region.

In addition to these three markers quantified directly by their respective primers and
assays, a fourth mitochondrial DNA (mtDNA) component—referred to as the "16S-Dloop
bridge sequence", hereafter "Bridge" for short—was quantified within the duplex ddPCR
reaction targeting both *16S* and *D-loop*. This Bridge signal arises from intact (2746+ bps)
mtDNA molecules that physically link the two markers (Fig. 1b, see details below and in the
supplement). Altogether, this yielded six distinct eNA components to be analyzed: Cytb
eDNA, Cytb emRNA, 16S eDNA, 16S erRNA, Dloop eDNA, and Bridge eDNA (Table 1).

All markers were quantified using digital droplet PCR (ddPCR). Cytb eDNA, Cytb
emRNA, Cytb No-RT, 16S erRNA and 16S No-RT were quantified using monoplex reactions.
Each 22 µL reaction contained 11 µL of ddPCR Supermix for Probes (Bio-Rad Inc., Hercules,
CA), 250 nM of probe, 900 nM of each primer (IDT), and 2 µL of template extract. The

thermocycling conditions were: 4°C for 10 min; 95°C for 10 min; 45 cycles of 94°C for 30 s
and 60°C for 60 s (annealing/extension); followed by 98°C for 10 min and a final hold at 4°C.

Table 1: Markers analyzed herein. "Bridge" refers to the 16S-Dloop bridge sequence (Figs.
1b, S1).

Mitochondrial locus	Nucleotide type	Extract	Assay	Target length (base pairs)
Cytb	DNA	eDNA	Cytb monoplex	79
Cytb	mRNA	eRNA	Cytb monoplex	79
16S	DNA	eDNA	16S-Dloop duplex	146
16S	rRNA	eRNA	16S monoplex	146
Dloop	DNA	eDNA	16S-Dloop duplex	390
Bridge	DNA	eDNA	16S-Dloop duplex*	2746-16390

* See supplement for calculation details

For *16S* and *D-loop* eDNA, we used a duplex ddPCR reaction. Each 22 µL reaction
included 11 µL of Supermix, 900 nM of *D-loop* primers, 600 nM of *16S* primers, 250 nM of
each probe, and 2 µL of extract. Given the longer amplicon sizes, we adjusted cycling
conditions: 4°C for 10 min; 95°C for 10 min; 44 cycles of 94°C for 30 s, 56°C for 30 s
(annealing), and 72°C for 120 s (extension); followed by 98°C for 10 min and a 4°C hold. This
duplex design allowed us to simultaneously quantify 16S, D-loop, and the mtDNA bridge
between them (Fig. 1b).

As mentioned above, this duplex reaction also allowed us to quantify a fourth marker,
Bridge, given by ddPCR droplet sorting and linkage between markers (Fig. S1). Droplet PCR
partitions the reaction into ~20,000 droplets, enabling absolute quantification of target
molecules using Poisson statistics. If targets are unlinked, they sort into droplets
independently, and double-positive droplets arise randomly. However, due to the circular
nature of mtDNA, *16S* and *D-loop* are physically linked in intact molecules. The shortest
span between them is 2746 bp, though linkage could span the full ~16,388 bp mitochondrial
genome (Louis et al., 2023; Xiong et al., 2009). Thus, in a hypothetical situation where all
mtDNA molecules are intact in the sample, all droplets containing 16S would automatically
also contain Dloop, and vice versa, making all droplets positive for both markers.

However, because eDNA in environmental samples is partially fragmented, droplets
in the duplex assay include a mixture of single-positive and double-positive events. Double-
positive droplets can arise from two mechanisms: (1) random co-localization of
independently degraded DNA fragments containing 16S and D-loop sequences, or (2) intact

DNA molecules physically bridging the two markers (i.e., ≥2746 bp). Using the counts of
single-positive droplets for each marker, one can calculate the expected number of double-
positive droplets that would result from mechanism (1) alone. This expected value is then
subtracted from the observed number of double-positive droplets, and the remainder is
attributed to mechanism (2), providing an estimate of the number of intact, bridged
molecules. See supplements for full derivation and schematics.

For all ddPCR assays, droplets were generated using the AutoDG Droplet Generator
(Bio-Rad Inc.), thermocycling was performed on C1000 Touch Thermal Cyclers (Bio-Rad
Inc.), and droplet fluorescence was read using a QX200 Droplet Reader (Bio-Rad Inc.). Each
plate included one positive control and three no-template controls (NTCs), used to define
positive thresholds and screen for contamination.

Finally, to confirm the absence of non-target marine mammals in the decay
experiment, we opportunistically performed metabarcoding on eDNA samples filtered at the
first timepoint of each carboy. We detected no other cetacean species in the decay water
(Tables S2, S3). Details of metabarcoding methods and results are provided in the
Supplement.

Statistical analysis and decay rate estimation

We built a Bayesian hierarchical model to jointly estimate the eNA concentrations
and the decay rates for each component and marker combination *i* (Table 1) from the
observed ddPCR droplet counts (W). To estimate the eNA concentrations from ddPCR
droplet reader observations, we followed the same approach used in Guri et al. [36] where the
number of positive droplets from technical replicates (*r*) of the same sample collected at
time (*t*) are modeled as a Binomial distribution:

$$W_{irt} \sim \text{Binomial}\,(U_{irt}, p_{it}) \tag{1}$$

$$cloglog(p_{it}) = ln(\omega_{it}) \tag{2}$$

Where p_{it} is the probability that an individual droplet positively amplifying the target locus *i*
at sampled time *t* from the total number of droplets generated U_{irt} and ω_{it} is the respective
eNA molecules (in copies/µL reaction volume)

For inferring the eNA concentration in the mesocosm (C_i; units copies/L) we
normalize for the volume filtered, eluted, and diluted as follows:

$$ln(\omega_{it}) = ln(C_{it}) - ln(rvol) + ln(tvol) + ln(d_{it}) - \ln(evol) - \ln(filt_{it}) - ln(dvol) \tag{3}$$

where *rvol* is the total ddPCR reaction volume (22 μL), $tvol$ is the template volume added to
the reaction (2 μL), d_i is a sample-specific template dilution (caused by reverse transcription
reaction), *evol* is the extraction eluted volume (50 μL), $filt_i$ is the sample-specific filtered
volume, and *dvol* is the ddPCR droplet volume (~0.85 nL), as specific by Bio-Rad and used
by Guri et al. (2024).

To calculate decay for each of the components (c), we tested six different decay
models and ultimately used the Biphasic Exponential Decay model which yielded the
highest likelihood (see Supplements for model testing). The biphasic exponential decay
model assumes two exponential decay phases with a transition time specific for each
component and marker (t_{ix}):

292
$$C_{cmt} = \begin{cases} C_{i,t=0} \times e^{-\lambda 1_i \times t}, & t < t_{ix}, \\ C_{i,t=t_{ix}} \times e^{-\lambda 2_i \times (t-t_{ix})}, & t \geq t_{ix}. \end{cases} \tag{4}$$

where C_{it} is the eNA concentration of each component (*c*) and marker (*m*) at time (*t*); $C_{i,t=0}$
is the respective initial concentration at t = 0; $\lambda 1_i$ and $\lambda 2_i$ are the decay rate constants for the
first and second phases, respectively; t_{ix} is the time where decay rate changes; and $C_{i,t=t_{ix}}$
is the concentration at t_{ix} determined as $C_{cm,t=t_{ix}} = C_{i,t=0} \times e^{-\lambda 1_i \times t_{ix}}$.

The concentrations C_{cmt} are jointly modeled from two technical replicates (r), and the
decay rates ($\lambda 1_i$ and $\lambda 2_i$) and t_x for each component and marker combination considering
all three biological replicates. The model was implemented in the Stan language with the R
package Rstan [37] running four independent Markov chain Monte Carlo (MCMC) chains with
5000 warmup and 10000 sampling iterations. Model's effective sample size was above 500,
and model's $\hat{R}$ convergence parameter was < 1.005 for all estimated parameters.

RESULTS

Controls

All target eNA assays from the fourth (control) carboy, filtration negative controls,
extraction blanks, and no reverse transcriptase (No-RT) controls yielded zero positive
droplets, with the exception of occasional single positive droplets in No-RT controls. These
rare events indicate minimal residual DNA carryover in RNA extracts despite DNase
treatment. To account for this, we subtracted the droplet count observed in each No-RT
control from the corresponding eRNA droplet count of the same sample before proceeding
with downstream analyses.

Little eDNA found in smaller pore sizes

To assess the particle-size distribution of dolphin eDNA, we filtered each time-point
sample through a 3-stage serial filtration system (5 µm, 1.0 µm, 0.45 µm). Preliminary
sample screening of the first timepoint samples found that >95% of the Cytb signal was
captured on the 5 µm filter at time zero. Therefore, all downstream analyses used this
fraction exclusively (Fig. 2).

Figure 2: Particle size distribution of Bottlenose Dolphin eDNA in the first timepoint.
Concentration of dolphin mitochondrial Cytb eDNA recovered at the first sampling time-
point as a function of pore size. Filters with pore sizes < 5 µm retained negligible eDNA, so
subsequent analyses rely exclusively on extracts from the 5 µm fraction.

Cytb messenger RNA was short-lived, and eNA decay was biphasic

We quantified decay of six different eNA components, including four eDNA markers
of varying length (Fig. 1b), one erRNA marker, and one emRNA marker (Table 1).

We detected dolphin-derived eDNA and 16S ribosomal eRNA across multiple
timepoints, with signals persisting up to 48 hours after the start of the experiment. In
contrast, Cytochrome b (Cytb) messenger eRNA was only detected in the first pre-transport
sample (Figs. 3, S4). This signal was lost during the ~3-hour transport period, and no Cytb
emRNA was detected in subsequent laboratory samples, suggesting extremely rapid
degradation.

Beyond this specific marker behavior, we observed that environmental nucleic acids
(eNA) generally followed a biphasic decay pattern. This was especially clear for short eDNA
fragments, which showed a steep decline within the first 48 hours, followed by a plateau
phase in which low concentrations persisted through the end of the 7-day experiment. To
formalize this observation, we compared six candidate decay models (Supplement), and the
biphasic exponential model provided the best fit to the data based on leave-one-out cross-
validation (Table S3).

Table 2: Posterior decay rates estimated across markers. "λ_1" is the decay rate for the first
phase of the exponential decay; "λ_1 2.5%" and "λ_1 97.5%" represent it's 95% confidence
interval; "t_x" is the time break at which the decay rate was found to change from the first
phase to the second; and "λ_2" is the decay rate for the second phase of the exponential
decay. NA values indicate no detections were made at the relevant timepoint, making values
unidentifiable.

Marker	λ_1	λ_1 2.5%	λ_1 97.5%	t_x	λ_2
Cytb eDNA	0.114 hr^{-1}	0.110 hr^{-1}	0.118 hr^{-1}	41 hrs	0.026 hr^{-1}
Cytb emRNA	1.615 hr^{-1}	0.877 hr^{-1}	2.763 hr^{-1}	NA	NA
16S eDNA	0.165 hr^{-1}	0.161 hr^{-1}	0.169 hr^{-1}	33 hrs	0.028 hr^{-1}
16S erRNA	0.236 hr^{-1}	0.207 hr^{-1}	0.267 hr^{-1}	29 hrs	0.054 hr^{-1}
Dloop eDNA	0.166 hr^{-1}	0.160 hr^{-1}	0.173 hr^{-1}	36 hrs	0.021 hr^{-1}
Bridge eDNA	0.190 hr^{-1}	0.175 hr^{-1}	0.206 hr^{-1}	28 hrs	0.044 hr^{-1}

Decay rate estimates varied by marker (Table 2). Although Cytb emRNA showed the
fastest apparent initial decay (λ_1 = 1.615 hr^{-1}), this estimate is based on a single detection
and subsequent non-detections, and should be interpreted with caution. Among
consistently detected markers, 16S erRNA had the highest initial decay rate (λ_1 = 0.236 hr^{-1};
Fig. 3b).

For eDNA, molecular length was a strong predictor of persistence (Fig. 3a). Shorter
targets, such as Cytb (λ_1 = 0.114 hr^{-1}), decayed more slowly, while longer targets like the
Bridge fragment spanning 16S and D-loop decayed faster (λ_1 = 0.190 hr^{-1}). The transition
between the two decay phases consistently fell between 24 and 48 hours, and all targets
excibitted a slower decay rate on the second phase (λ_2 between 0.054 and 0.021, Table 2).

Figure 3: Biphasic-model decay of dolphin environmental nucleic acids (eNAs). (a) Mitochondrial DNA targets. (b) RNA targets. Solid lines are the posterior mean trajectories from the biphasic Bayesian ddPCR model; filled circles are the model-predicted concentrations at each sampling time for every biological replicate (three carboys). Colors distinguish markers and correspond to amplicon length: Cyt b 79 bp, 16 S 146 bp, DLL1 390 bp, and the 2 746 bp Bridge (DNA); Cyt b emRNA 79 bp and 16 S erRNA 146 bp (RNA). Axes are log-scaled; concentrations are expressed as copies L^{-1}.

DISCUSSION

This study provides the first empirical estimates of decay rates for multiple environmental nucleic acid (eNA) components derived from a marine mammal. Using a controlled experiment with seawater from a dolphin enclosure, we documented the degradation of mitochondrial eDNA fragments of different lengths, ribosomal eRNA, and messenger eRNA from *Tursiops truncatus* over seven days. We found that decay patterns were highly dependent on molecule type and length, with messenger RNA degrading the fastest, ribosomal RNA decaying slightly faster than its DNA counterpart, and longer DNA fragments decaying more rapidly than shorter ones. Across most targets, decay followed a biphasic trajectory, with a steep initial decline and a slower second phase. These results confirm that eNA components differ markedly in environmental stability and underscore the potential of combining multiple markers to improve the temporal resolution of eNA-based detections.

Rethinking molecular strategies for eDNA age estimation

The ratio between eDNA and emRNA has been proposed as a molecular proxy for time since eDNA shedding[12], consistent with the spatiotemporally variable eDNA composition hypothesis [17]. This approach assumes that RNA decays more rapidly than DNA, such that high RNA:DNA ratios indicate recent activity. However, our results suggest that messenger RNA (emRNA) is not well-suited for this application in marine systems. Cytb emRNA was only detectable at the initial timepoint and degraded beyond detection during transport—highlighting the molecule's extreme lability. Similar challenges have been noted in other marine studies, where emRNA is often either undetectable or not measurably different in decay from eDNA [30,38]. Beyond this, emRNA analysis requires transcript-specific assays, rapid stabilization, and higher-cost laboratory workflows [2,39], which limit its practicality for most field applications.

Instead, our findings support the use of eDNA fragment length as a proxy for degradation state. Longer mitochondrial markers decayed consistently faster than shorter ones, as shown here and in prior work [20,33], making them well-suited for relative age estimation [19]. This method is logistically simple—relying on standard eDNA workflows and multiplexed assays—and broadly applicable across taxa and systems. Importantly, quantitative estimation of eDNA age requires both accurate decay rates and knowledge of the initial proportion of each component, so that their relative abundance over time can be meaningfully interpreted. When targeting mitochondrial loci, the physical linkage of gene regions provides a useful simplification: short and long fragments are expected to originate at roughly equal copy number, enabling decay-based inferences without requiring independent normalization of starting ratios. On the other hand, because mRNA expression

levels change by context, the starting proportion between mRNA and DNA in environmental
samples will always vary, and initial proportions cannot be accurately established. Thus,
pairing mitochondrial markers of different lengths potentially offers a scalable, cost-
effective alternative to RNA-based strategies. Rather than relying on a single target, a duplex
assay can be designed to simultaneously amplify two mitochondrial regions with a
deliberate difference in length, maximizing the contrast in decay rates while maintaining
amplification efficiency.

Mechanisms and practical significance of Biphasic decay

Dolphin eDNA in our experiment followed a biphasic exponential decay: a steep first-
phase loss lasting ~48 h, followed by a low-level "tail" that persisted for days. Similar two-
phase kinetics have been reported from rivers, lakes, and coastal waters [25,40,41], but often go
unrecognized when decay experiments end after only one or two days [42,43]. We detected no
compelling biphasic signal for either ribosomal or messenger eRNA, but it is possible RNA
also follows the same pattern.

The processes that generate two-phase decay remain unresolved. Three non-
exclusive explanations have been previously proposed: (i) physical shielding, in which DNA
adsorbed to mineral or organic particles become inaccessible to nucleases [44,45]; (ii)
encapsulation, whereby intact cells and mitochondria only release DNA after membrane
rupture [46,47]; and (iii) particle-size-dependent removal, where large aggregates settle or are
grazed quickly while finer particles or dissolved DNA persist in suspension [47]. Given these
mechanisms are associated with different eDNA states, disentangling them will require
experiments that characterize or separate DNA states.

Nonetheless, from a monitoring perspective, the slow second phase probably
contributes little to routine eDNA surveys: most metabarcoding data will be dominated by
the highly concentrated but rapidly decaying fraction, as the residual tail is easily swamped
by fresh inputs. Nevertheless, understanding the process could improve source-age
estimates, especially when low concentrations are observed.

Distinguishing Ribosomal and Messenger eRNA in Environmental Studies

In our study, we found cytochrome b (Cytb) messenger eRNA (emRNA) at very low
concentrations, detectable only at the initial time point before water samples were
transferred to the environmental chamber. Low levels of Cytb emRNA in environmental
samples has been found in other studies [48], and these low concentrations are the likely
explained by low generation and expression of Cytb from eNA-generating tissue. In contrast,
ribosomal eRNA (erRNA) was present at much higher concentrations in our samples,
consistent with its constitutive expression, and decayed only slightly faster than its

equivalent DNA. This contrast in stability and concentrations of both forms of RNA aligns
with well-documented differences in RNA stability from forensic and molecular biology
research, where ribosomal RNA's secondary structure may protect it from degradation [49,50].
These findings highlight a fundamental distinction between erRNA and emRNA: while erRNA
in found in high concentrations and can be reliably detected, emRNA is highly transient,
making it a good candidate for detection of more recent targets, but potentially unreliable in
yielding consistent taxa detections given its low starting concentrations.

The rapid degradation of mRNA is associated with its biological function. Within cells,
mRNA is a highly labile molecule that allows for rapid and dynamic gene expression
regulation [51]. To maintain precise control over protein synthesis, cells possess multiple
pathways for degrading mRNA efficiently, including exonuclease-mediated decay and other
complex biomolecular pathways [52,53]. Additionally, mRNA's single-stranded structure and
chemical composition (i.e., an extra hydroxyl group) make it inherently less stable than DNA,
leading to its rapid degradation both within cells and in the environment, particularly in high
temperatures and alkaline conditions [27,54]. In contrast, rRNA is constitutively (permanently)
expressed in large quantities, representing over 80% of all RNA within cells in many
estimates, and plays a structural role in ribosome assembly [55–57]. Its highly structured
secondary and tertiary conformations enhance its resistance to enzymatic degradation [50,58],
which likely extends to higher stability in the environment as well.

Despite these well-established molecular differences, the term "eRNA" is often used
interchangeably in the eDNA literature to refer to both erRNA and emRNA, despite their
fundamentally different properties. This conflation can lead to incorrect assumptions about
eRNA persistence, decay rates, and ecological interpretability. Specifically, our results and
previous literature indicate that emRNA degrades much faster than eDNA in environmental
samples, and its initial concentration depends heavily on gene expression in the tissues
shedding eDNA. Because gene expression varies across tissues, developmental stages, and
environmental conditions, interpreting emRNA abundance requires transcriptomic
knowledge of the target species—information that is often unavailable, but can be used to
inform emRNA assay development [39]. Consequently, assays designed for both eDNA and
emRNA detection (e.g., targeting Cytb, as tested here) may not be widely effective, as the
emRNA marker will have to be customized for each application.

In contrast, erRNA is more abundant, making it a more stable biomarker but
unsuitable for most other applications where emRNA would be advantageous over eDNA,
such as metabolic inferences [2]. Supporting this, previous metabarcoding studies using
erRNA and eDNA to detect species with a 12S marker found minimal differences in detection
rates, though erRNA showed faster species accumulation curves [38,59]. Macher et al.[38] also
tested emRNA-based detection using cytochrome oxidase I (COI), and as expected, emRNA

was less effective than eDNA in species detection due to its rapid degradation (see above).
However, its faster decay resulted in stronger spatiotemporal patterns, reflecting its short-
lived nature in the environment.

These findings reinforce the importance of distinguishing rRNA from mRNA in
environmental studies. The ecological patterns observed for rRNA cannot be assumed to
apply to mRNA, and vice versa. Given their fundamental biological and functional
differences, treating both molecules collectively as "eRNA" is misleading and can obscure
key differences in their degradation rates and interpretability in environmental monitoring.
Future research should clarify this distinction and ensure that eRNA studies account for the
vastly different behaviors of these two RNA types in environmental contexts.

Dolphin eDNA Predominantly Retained in Large Particle Fractions

Interestingly, we found that the vast majority of dolphin eDNA was captured on the
largest pore size filter (5 μm), with only residual amounts recovered in smaller size fractions
(Fig. 2). This pattern contrasts with previous studies on the particle size distribution (PSD) of
eDNA from other marine taxa, such as teleost fish (Jo et al., 2019) and elasmobranchs [60],
whose eDNA is more evenly distributed down to small filter pore sizes (< 1 μm). While eDNA
PSD is known to vary with environmental conditions and organism [44,61], the pattern observed
here closely resembles that reported for freshwater fish in the presence of clay or titanium
dioxide, which strongly adsorb to eDNA [62]. However, such particles were not present in our
samples in substantial quantities. There are two possible alternative explanations for the
observed PSD. First, it is possible that filtration failure occurred at the smaller pore sizes,
preventing accurate capture of eDNA in those fractions. Nonetheless, we observed the same
pattern across three replicate carboys, which were filtered separately. Alternatively, if
experimental error can be excluded, the results may indicate that marine mammal eNA is
predominantly associated with larger particles—possibly due to tissue origin, mucus
binding, or aggregation. If this is the case, it has important implications for sampling design:
using larger pore size filters may enhance eDNA yield and improve detection probability for
marine mammals by allowing greater volumes to be processed.

Carryover DNA and components of eDNA

Commercially available extractions columns were originally designed to extract high
molecular yield DNA from tissue samples. However, they have been commonly used for
extracting eDNA from filters over the past two decades and have yielded sufficient DNA for
most applications [1]. To our knowledge, no other studies have captured and quantified eDNA
that passes through the DNA extraction columns to quantify this inefficiency. Here, we found
a substantial amount of eDNA passed through the initial column, largely comprising shorter
DNA molecules as measured by a TapeStation (Supplement Fig. 2). This finding points to a

potential blind spot in many eDNA workflows: studies seeking rare targets or high detection sensitivity should be cautious about relying on column-based extraction methods, despite their convenience and scalability.

In addition to quantifying the amount of carryover eDNA, we also quantified the decay rate of carryover eDNA with one marker (Cytb) and found it to decay significantly slower than the eDNA fraction captured on the first extraction column (Fig. S3). This is likely due to a cascading effect, where the decay of longer eDNA molecules creates shorter eDNA molecules, resulting is a slower apparent decay of the smaller fragment size [17]. Nonetheless, it is unclear if this method of splitting eNA components (i.e., eDNA vs. carryover eDNA) will be useful in the future because the fraction of eNA passing through the first column may not be stable, and may vary substantially under varying total quantities of DNA in the sample and other chemical characteristics of the sample. Thus, while splitting eNA by molecule length (in basepairs) may find its applications, separation of multiple eNA components may be more reliable with other methods such as sequential filtration [20,63], evaluating multiple markers of varying lengths as here, or differential extraction methods [14].

ACKNOWLEDGEMENTS:

The authors thank the U.S. Naval Base Kitsap-Bangor for logistic support, Kate Bertko for administrative and field support, Krista Nichols, Amy Van Cise, Elizabeth Brasseale, and Olivia Scott for field sampling support, and Andrew Olaf Shelton for sampling support and assistance discussing the STAN decay model. This research was funded by the Office of Naval Research under Award Number (N00014-22-1-2719) and supported through collaborative research efforts with the U.S. Navy's Marine Mammals Program. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

AUTHOR CONTRIBUTIONS

All authors worked together to conceive and design the experiment and analyses; MS, KP, RK, EAA planned fieldwork and sampled water; PBD prepared and executed decay experiment; MS designed assays; PBD performed lab work; PBD and GG wrote the hierarchical Bayesian model; PBD, GG and EAA analyzed and visualized data; KP, RK, EAA obtained funding and supervised the project. All authors discussed results, contributed to and approved the final manuscript draft.

DATA AVAILABILITY STATEMENT:

545   Raw dataset and corresponding metadata will be available as supplement files upon
546 publication

COMPETING INTERESTS

548   The authors declare no competing interests.

SUPPORTING INFORMATION:

550   Additional description of ddPCR methods, additional visualizations of the raw data,
551 and analysis of carryover eDNA is provided in a pdf as Supplementary Information.

REFERENCES

(1) Takahashi, M.; Saccò, M.; Kestel, J. H.; Nester, G.; Campbell, M. A.; Van Der Heyde, M.; Heydenrych, M. J.; Juszkiewicz, D. J.; Nevill, P.; Dawkins, K. L.; Bessey, C.; Fernandes, K.; Miller, H.; Power, M.; Mousavi-Derazmahalleh, M.; Newton, J. P.; White, N. E.; Richards, Z. T.; Allentoft, M. E. Aquatic Environmental DNA: A Review of the Macro-Organismal Biomonitoring Revolution. *Sci. Total Environ.* **2023**, *873*, 162322. https://doi.org/10.1016/j.scitotenv.2023.162322.

(2) Yates, M. C.; Derry, A. M.; Cristescu, M. E. Environmental RNA: A Revolution in Ecological Resolution? *Trends Ecol. Evol.* **2021**, *36* (7), 601–609. https://doi.org/10.1016/j.tree.2021.03.001.

(3) Barnes, M. A.; Turner, C. R. The Ecology of Environmental DNA and Implications for Conservation Genetics. *Conserv. Genet.* **2016**, *17* (1), 1–17. https://doi.org/10.1007/s10592-015-0775-4.

(4) Allan, E. A.; DiBenedetto, M. H.; Lavery, A. C.; Govindarajan, A. F.; Zhang, W. G. Modeling Characterization of the Vertical and Temporal Variability of Environmental

DNA in the Mesopelagic Ocean. *Sci. Rep.* **2021**, *11* (1), 21273.

https://doi.org/10.1038/s41598-021-00288-5.

(5) Harrison, J. B.; Sunday, J. M.; Rogers, S. M. Predicting the Fate of eDNA in the

Environment and Implications for Studying Biodiversity. *Proc. R. Soc. B Biol. Sci.* **2019**,

286 (1915), 20191409. https://doi.org/10.1098/rspb.2019.1409.

(6) Barnes, M. A.; Turner, C. R.; Jerde, C. L.; Renshaw, M. A.; Chadderton, W. L.; Lodge, D.

M. Environmental Conditions Influence eDNA Persistence in Aquatic Systems. *Environ.*

Sci. Technol. **2014**, *48* (3), 1819–1827. https://doi.org/10.1021/es404734p.

(7) Jo, T.; Arimoto, M.; Murakami, H.; Masuda, R.; Minamoto, T. Estimating Shedding and

Decay Rates of Environmental Nuclear DNA with Relation to Water Temperature and

Biomass. *Environ. DNA* **2020**, *2* (2), 140–151. https://doi.org/10.1002/edn3.51.

(8) Scriver, M.; Zaiko, A.; Pochon, X.; von Ammon, U. Harnessing Decay Rates for Coastal

Marine Biosecurity Applications: A Review of Environmental DNA and RNA Fate.

Environ. DNA **2023**, *5* (5), 960–972. https://doi.org/10.1002/edn3.405.

(9) Xiong, J.; MacCready, P.; Brasseale, E.; Andruszkiewicz Allan, E.; Ramón-Laca, A.;

Parsons, K. M.; Shaffer, M.; Kelly, R. P. Advective Transport Drives Environmental DNA

Dispersal in an Estuary. *Environ. Sci. Technol.* **2025**, *59* (15), 7506–7516.

https://doi.org/10.1021/acs.est.5c01286.

(10)   Suarez-Bregua, P.; Álvarez-González, M.; Parsons, K. M.; Rotllant, J.; Pierce, G. J.;

Saavedra, C. Environmental DNA (eDNA) for Monitoring Marine Mammals: Challenges

and Opportunities. *Front. Mar. Sci.* **2022**, *9*.

https://doi.org/10.3389/fmars.2022.987774.

(11)   Jerde, C. L.; Mahon, A. R.; Chadderton, W. L.; Lodge, D. M. "Sight-Unseen" Detection

591   of Rare Aquatic Species Using Environmental DNA: eDNA Surveillance of Rare Aquatic

592   Species. *Conserv. Lett.* **2011**, *4* (2), 150–157. https://doi.org/10.1111/j.1755-

593   263X.2010.00158.x.

594   (12)   Marshall, N. T.; Vanderploeg, H. A.; Chaganti, S. R. Environmental (e)RNA Advances

595   the Reliability of eDNA by Predicting Its Age. *Sci. Rep.* **2021**, *11* (1), 2769.

596   https://doi.org/10.1038/s41598-021-82205-4.

597   (13)   Sampaio-Silva, F.; Magalhães, T.; Carvalho, F.; Dinis-Oliveira, R. J.; Silvestre, R.

598   Profiling of RNA Degradation for Estimation of Post Morterm Interval. *PLOS ONE* **2013**, *8*

599   (2), e56507. https://doi.org/10.1371/journal.pone.0056507.

600   (14)   Kirtane, A.; Kleyer, H.; Deiner, K. Sorting States of Environmental DNA: Effects of

601   Isolation Method and Water Matrix on the Recovery of Membrane-Bound, Dissolved,

602   and Adsorbed States of eDNA. *Environ. DNA* **2023**, *5* (3), 582–596.

603   https://doi.org/10.1002/edn3.417.

604   (15)   Mauvisseau, Q.; Harper, L.; Sander, M.; Hanner, R. H.; Kleyer, H.; Deiner, K. *The*

605   *Multiple States of Environmental DNA and What Is Known about Their Persistence in*

606   *Aquatic Environments*; preprint; Preprints, 2021.

607   https://doi.org/10.22541/au.163638394.41572509/v1.

608   (16)   Nagler, M.; Podmirseg, S. M.; Ascher-Jenull, J.; Sint, D.; Traugott, M. Why ᴇDNA

609   Fractions Need Consideration in Biomonitoring. *Mol. Ecol. Resour.* **2022**, *22* (7), 2458–

610   2470. https://doi.org/10.1111/1755-0998.13658.

(17)   Brandão-Dias, P. F. P.; Hallack, D. M. C.; Snyder, E. D.; Tank, J. L.; Bolster, D.; Volponi,

S.; Shogren, A. J.; Lamberti, G. A.; Bibby, K.; Egan, S. P. Particle Size Influences Decay

Rates of Environmental DNA in Aquatic Systems. *Mol. Ecol. Resour.* **2023**, *23* (4), 756–

770. https://doi.org/10.1111/1755-0998.13751.

(18)   Bylemans, J.; Furlan, E. M.; Gleeson, D. M.; Hardy, C. M.; Duncan, R. P. Does Size

Matter? An Experimental Evaluation of the Relative Abundance and Decay Rates of

Aquatic Environmental DNA. *Environ. Sci. Technol.* **2018**, *52* (11), 6408–6416.

https://doi.org/10.1021/acs.est.8b01071.

(19)   Jo, T.; Murakami, H.; Masuda, R.; Minamoto, T. Selective Collection of Long

Fragments of Environmental DNA Using Larger Pore Size Filter. *Sci. Total Environ.* **2020**,

735, 139462. https://doi.org/10.1016/j.scitotenv.2020.139462.

(20)   Brandão-Dias, P. F. P.; Snyder, E.; Tank, J. L.; Mahl, U. H.; Peters, B.; Shogren, A. J.;

Bolster, D.; Lamberti, G. A.; Bibby, K.; Egan, S. P. Comparing the Fate of eDNA by

Particle Sizes and Molecule Lengths in Recirculating Streams. *Environ. DNA* **2025**, *7* (2),

e70066. https://doi.org/10.1002/edn3.70066.

(21)   Jo, T.; Arimoto, M.; Murakami, H.; Masuda, R.; Minamoto, T. Particle Size Distribution

of Environmental DNA from the Nuclei of Marine Fish. *Environ. Sci. Technol.* **2019**, *53*

(16), 9947–9956. https://doi.org/10.1021/acs.est.9b02833.

(22)   Sakata, M. K.; Yamamoto, S.; Gotoh, R. O.; Miya, M.; Yamanaka, H.; Minamoto, T.

Sedimentary eDNA Provides Different Information on Timescale and Fish Species

Composition Compared with Aqueous eDNA. *Environ. DNA* **2020**, *2* (4), 505–518.

https://doi.org/10.1002/edn3.75.

(23)   Joseph, C.; Faiq, M. E.; Li, Z.; Chen, G. Persistence and Degradation Dynamics of

634   eDNA Affected by Environmental Factors in Aquatic Ecosystems. *Hydrobiologia* **2022**,

635   *849* (19), 4119–4133. https://doi.org/10.1007/s10750-022-04959-w.

636   (24)   Lamb, P. D.; Fonseca, V. G.; Maxwell, D. L.; Nnanatu, C. C. Systematic Review and

637   Meta-Analysis: Water Type and Temperature Affect Environmental DNA Decay. *Mol.*

638   *Ecol. Resour.* **2022**, *22* (7), 2494–2505. https://doi.org/10.1111/1755-0998.13627.

639   (25)   Andruszkiewicz Allan, E.; Zhang, W. G.; C. Lavery, A.; F. Govindarajan, A.

640   Environmental DNA Shedding and Decay Rates from Diverse Animal Forms and

641   Thermal Regimes. *Environ. DNA* **2021**, *3* (2), 492–514.

642   https://doi.org/10.1002/edn3.141.

643   (26)   Jo, T.; Murakami, H.; Masuda, R.; Sakata, M. K.; Yamamoto, S.; Minamoto, T. Rapid

644   Degradation of Longer DNA Fragments Enables the Improved Estimation of Distribution

645   and Biomass Using Environmental DNA. *Mol. Ecol. Resour.* **2017**, *17* (6).

646   https://doi.org/10.1111/1755-0998.12685.

647   (27)   Jo, T.; Tsuri, K.; Hirohara, T.; Yamanaka, H. Warm Temperature and Alkaline

648   Conditions Accelerate Environmental RNA Degradation. *Environ. DNA* **2023**, *5* (5), 836–

649   848. https://doi.org/10.1002/edn3.334.

650   (28)   Kagzi, K.; Hechler, R. M.; Fussmann, G. F.; Cristescu, M. E. Environmental RNA

651   Degrades More Rapidly than Environmental DNA across a Broad Range of pH

652   Conditions. *Mol. Ecol. Resour.* **2022**, *22* (7), 2640–2650. https://doi.org/10.1111/1755-

653   0998.13655.

(29) Qian, T.; Shan, X.; Wang, W.; Jin, X. Effects of Temperature on the Timeliness of

eDNA/eRNA: A Case Study of Fenneropenaeus Chinensis. *Water* **2022**, *14* (7), 1155.

https://doi.org/10.3390/w14071155.

(30) Wood, S. A.; Biessy, L.; Latchford, J. L.; Zaiko, A.; von Ammon, U.; Audrezet, F.;

Cristescu, M. E.; Pochon, X. Release and Degradation of Environmental DNA and RNA

in a Marine System. *Sci. Total Environ.* **2020**, *704*, 135314.

https://doi.org/10.1016/j.scitotenv.2019.135314.

(31) Scriver, M.; von Ammon, U.; Pochon, X.; Arranz, V.; Stanton, J.-A. L.; Gemmell, N. J.;

Zaiko, A. Environmental DNA–RNA Dynamics Provide Insights for Effective Monitoring of

Marine Invasive Species. *Environ. DNA* **2024**, *6* (2), e531.

https://doi.org/10.1002/edn3.531.

(32) Miyata, K.; Inoue, Y.; Amano, Y.; Nishioka, T.; Yamane, M.; Kawaguchi, T.; Morita, O.;

Honda, H. Fish Environmental RNA Enables Precise Ecological Surveys with High

Positive Predictivity. *Ecol. Indic.* **2021**, *128*, 107796.

https://doi.org/10.1016/j.ecolind.2021.107796.

(33) Hirohara, T.; Tsuri, K.; Miyagawa, K.; Paine, R. T. R.; Yamanaka, H. The Application of

PMA (Propidium Monoazide) to Different Target Sequence Lengths of Zebrafish eDNA: A

New Approach Aimed Toward Improving Environmental DNA Ecology and Biological

Surveillance. *Front. Ecol. Evol.* **2021**, *9*.

(34) Louis, M.; Korlević, P.; Nykänen, M.; Archer, F.; Berrow, S.; Brownlow, A.; Lorenzen, E.

D.; O'Brien, J.; Post, K.; Racimo, F.; Rogan, E.; Rosel, P. E.; Sinding, M.-H. S.; van der Es,

H.; Wales, N.; Fontaine, M. C.; Gaggiotti, O. E.; Foote, A. D. Ancient Dolphin Genomes

Reveal Rapid Repeated Adaptation to Coastal Waters. *Nat. Commun.* **2023**, *14* (1),

4020. https://doi.org/10.1038/s41467-023-39532-z.

(35)   Xiong, Y.; Brandley, M. C.; Xu, S.; Zhou, K.; Yang, G. Seven New Dolphin

Mitochondrial Genomes and a Time-Calibrated Phylogeny of Whales. *BMC Evol. Biol.*

2009, *9* (1), 20. https://doi.org/10.1186/1471-2148-9-20.

(36)   Guri, G.; Ray, J. L.; Shelton, A. O.; Kelly, R. P.; Præbel, K.; Andruszkiewicz Allan, E.;

Yoccoz, N.; Johansen, T.; Wangensteen, O. S.; Hanebrekke, T.; Westgaard, J.-I.

Quantifying the Detection Sensitivity and Precision of qPCR and ddPCR Mechanisms

for eDNA Samples. *Ecol. Evol.* **2024**, *14* (12), e70678.

https://doi.org/10.1002/ece3.70678.

(37)   Carpenter, B.; Gelman, A.; Hoffman, M. D.; Lee, D.; Goodrich, B.; Betancourt, M.;

Brubaker, M.; Guo, J.; Li, P.; Riddell, A. Stan: A Probabilistic Programming Language. *J.*

Stat. Softw. **2017**, *76*, 1–32. https://doi.org/10.18637/jss.v076.i01.

(38)   Macher, T.-H.; Arle, J.; Beermann, A. J.; Frank, L.; Hupało, K.; Koschorreck, J.; Schütz,

R.; Leese, F. Is It Worth the Extra Mile? Comparing Environmental DNA and RNA

Metabarcoding for Vertebrate and Invertebrate Biodiversity Surveys in a Lowland

Stream. *PeerJ* **2024**, *12*, e18016. https://doi.org/10.7717/peerj.18016.

(39)   Parsley, M. B.; Goldberg, C. S. Environmental RNA Can Distinguish Life Stages in

Amphibian Populations. *Mol. Ecol. Resour.* **2023**.

(40)   Jo, T. S. Factors Affecting Biphasic Degradation of eDNA Released by Japanese Jack

Mackerel (*Trachurus Japonicus*). *J. Exp. Mar. Biol. Ecol.* **2023**, *568*, 151941.

https://doi.org/10.1016/j.jembe.2023.151941.

(41)   Shogren, A. J.; Tank, J. L.; Egan, S. P.; August, O.; Rosi, E. J.; Hanrahan, B. R.;

699   Renshaw, M. A.; Gantz, C. A.; Bolster, D. Water Flow and Biofilm Cover Influence

700   Environmental DNA Detection in Recirculating Streams. *Environ. Sci. Technol.* **2018**, *52*

701   (15), 8530–8537. https://doi.org/10.1021/acs.est.8b01822.

702   (42)   Sassoubre, L. M.; Yamahara, K. M.; Gardner, L. D.; Block, B. A.; Boehm, A. B.

703   Quantification of Environmental DNA (eDNA) Shedding and Decay Rates for Three

704   Marine Fish. *Environ. Sci. Technol.* **2016**, *50* (19), 10456–10464.

705   https://doi.org/10.1021/acs.est.6b03114.

706   (43)   Snyder, E. D.; Tank, J. L.; Brandão-Dias, P. F. P.; Bibby, K.; Shogren, A. J.; Bivins, A. W.;

707   Peters, B.; Curtis, E. M.; Bolster, D.; Egan, S. P.; Lamberti, G. A. Environmental DNA

708   (eDNA) Removal Rates in Streams Differ by Particle Size under Varying Substrate and

709   Light Conditions. *Sci. Total Environ.* **2023**, *903*, 166469.

710   https://doi.org/10.1016/j.scitotenv.2023.166469.

711   (44)   Barnes, M. A.; Chadderton, W. L.; Jerde, C. L.; Mahon, A. R.; Turner, C. R.; Lodge, D.

712   M. Environmental Conditions Influence eDNA Particle Size Distribution in Aquatic

713   Systems. *Environ. DNA* **2021**, *3* (3), 643–653. https://doi.org/10.1002/edn3.160.

714   (45)   Cai, P.; Huang, Q.-Y.; Zhang, X.-W. Interactions of DNA with Clay Minerals and Soil

715   Colloidal Particles and Protection against Degradation by DNase. *Environ. Sci. Technol.*

716   **2006**, *40* (9), 2971–2976. https://doi.org/10.1021/es0522985.

717   (46)   Moushomi, R.; Wilgar, G.; Carvalho, G.; Creer, S.; Seymour, M. Environmental DNA

718   Size Sorting and Degradation Experiment Indicates the State of Daphnia Magna

Mitochondrial and Nuclear eDNA Is Subcellular. *Sci. Rep.* **2019**, *9* (1), 12500.

https://doi.org/10.1038/s41598-019-48984-7.

(47)   Torti, A.; Lever, M. A.; Jørgensen, B. B. Origin, Dynamics, and Implications of

Extracellular DNA Pools in Marine Sediments. *Mar. Genomics* **2015**, *24*, 185–196.

https://doi.org/10.1016/j.margen.2015.08.007.

(48)   Jo, T. S.; Ozaki, Y.; Matsuda, N.; Yamanaka, H. Assessment of Allometry in

Environmental DNA and RNA Production From Ayu (Plecoglossus Altivelis) in an

Experimental Condition Using Mitochondrial and Nuclear Gene Markers. *Environ. DNA*

2024, *6* (6), e70052. https://doi.org/10.1002/edn3.70052.

(49)   Houseley, J.; Tollervey, D. The Many Pathways of RNA Degradation. *Cell* **2009**, *136*

(4), 763–776. https://doi.org/10.1016/j.cell.2009.01.019.

(50)   Sidova, M.; Tomankova, S.; Abaffy, P.; Kubista, M.; Sindelka, R. Effects of *Post-*

Mortem and Physical Degradation on RNA Integrity and Quality. *Biomol. Detect.*

Quantif. **2015**, *5*, 3–9. https://doi.org/10.1016/j.bdq.2015.08.002.

(51)   Sachs, A. B. Messenger RNA Degradation in Eukaryotes. *Cell* **1993**, *74* (3), 413–421.

https://doi.org/10.1016/0092-8674(93)80043-E.

(52)   Belasco, J. G.; Brawerman, G. *Control of Messenger RNA Stability*; Elsevier, 2012.

(53)   Shyu, A.; Wilkinson, M. F.; van Hoof, A. Messenger RNA Regulation: To Translate or to

Degrade. *EMBO J.* **2008**, *27* (3), 471–481. https://doi.org/10.1038/sj.emboj.7601977.

(54)   Li, Y.; Breaker, R. R. Kinetics of RNA Degradation by Specific Base Catalysis of

Transesterification Involving the 2'-Hydroxyl Group. *J. Am. Chem. Soc.* **1999**, *121* (23),

5364–5372. https://doi.org/10.1021/ja990592p.

(55)   Blanco, A.; Blanco, G. Chapter 6 - Nucleic Acids. In *Medical Biochemistry*; Blanco,

A., Blanco, G., Eds.; Academic Press, 2017; pp 121–140. https://doi.org/10.1016/B978-

0-12-803550-4.00006-9.

(56)   Lewis, J. D.; Tollervey, D. Like Attracts Like: Getting RNA Processing Together in the

Nucleus. *Science* **2000**, *288* (5470), 1385–1389.

https://doi.org/10.1126/science.288.5470.1385.

(57)   Warner, J. R. The Economics of Ribosome Biosynthesis in Yeast. *Trends Biochem.*

Sci. **1999**, *24* (11), 437–440. https://doi.org/10.1016/S0968-0004(99)01460-7.

(58)   Zhang, K.; Hodge, J.; Chatterjee, A.; Moon, T. S.; Parker, K. M. Duplex Structure of

Double-Stranded RNA Provides Stability against Hydrolysis Relative to Single-Stranded

RNA. *Environ. Sci. Technol.* **2021**, *55* (12), 8045–8053.

https://doi.org/10.1021/acs.est.1c01255.

(59)   Littlefair, J. E.; Rennie, M. D.; Cristescu, M. E. Environmental Nucleic Acids: A Field-

Based Comparison for Monitoring Freshwater Habitats Using eDNA and eRNA. *Mol.*

Ecol. Resour. **2022**, *22* (8), 2928–2940. https://doi.org/10.1111/1755-0998.13671.

(60)   Cooper, M. K.; Villacorta-Rath, C.; Burrows, D.; Jerry, D. R.; Carr, L.; Barnett, A.;

Huveneers, C.; Simpfendorfer, C. A. Practical eDNA Sampling Methods Inferred from

Particle Size Distribution and Comparison of Capture Techniques for a Critically

Endangered Elasmobranch. *Environ. DNA* **2022**, *4* (5), 1011–1023.

https://doi.org/10.1002/edn3.279.

(61)   Zhao, B.; Van Bodegom, P. M.; Trimbos, K. The Particle Size Distribution of

Environmental DNA Varies with Species and Degradation. *Sci. Total Environ.* **2021**, *797*,

149175. https://doi.org/10.1016/j.scitotenv.2021.149175.

(62)   Brandão-Dias, P. F. . P.; Tank, J. L.; Snyder, E. D.; Mahl, U. H.; Peters, B.; Bolster, D.;

Shogren, A. J.; Lamberti, G. A.; Bibby, K.; Egan, S. P. Suspended Materials Affect Particle

Size Distribution and Removal of Environmental DNA in Flowing Waters. *Environ. Sci.*

Technol. **2023**, *57* (35), 13161–13171. https://doi.org/10.1021/acs.est.3c02638.

(63)   Turner, C. R.; Barnes, M. A.; Xu, C. C. Y.; Jones, S. E.; Jerde, C. L.; Lodge, D. M.

Particle Size Distribution and Optimal Capture of Aqueous Macrobial ᴇDNA. *Methods*

Ecol. Evol. **2014**, *5* (7), 676–684. https://doi.org/10.1111/2041-210X.12206.

771