1

Cancer cells differentially modulate mitochondrial respiration to alter redox state and

enable biomass synthesis in nutrient-limited environments

Sarah M. Chang[1,2,3], Muhammad Bin Munim[1,2], Sonia E. Trojan[1,4], Huel Cox III[1,3,], Anna

Shevzov-Zebrun[1,2], Keene L. Abbott[1,2], Renee Chang[1,2], and Matthew G. Vander Heiden[1,2,5*]

[1]Koch Institute for Integrative Cancer Research, Massachusetts Institute of Technology,

Cambridge, MA, USA

[2]Department of Biology, Massachusetts Institute of Technology, Cambridge, MA, USA

[3]Harvard/MIT MD-PhD Program, Boston, MA, USA

[4]Jagiellonian University Medical College, Faculty of Medicine, Chair of Medical Biochemistry,

Krakow, Poland

[5]Dana-Farber Cancer Institute, Boston, MA, USA

*Correspondence: mvh@mit.edu

Abstract

The cell NAD+/NADH ratio can constrain biomass synthesis and influence proliferation in nutrient-

limited environments. However, which cell processes regulate the NAD+/NADH ratio is not known.

Here, we find that some cancer cells elevate the NAD+/NADH ratio in response to serine

deprivation by increasing mitochondrial respiration. Cancer cells that elevate mitochondrial

respiration have higher serine production and proliferation in serine limiting conditions than cells

with no mitochondrial respiration response, independent of serine synthesis enzyme expression.

Increases in mitochondrial respiration and the NAD+/NADH ratio promote serine synthesis

regardless of whether serine is environmentally limiting. Lipid deprivation can increase the

NAD+/NADH ratio via mitochondrial respiration in some cells, including cells that do not increase

respiration following serine deprivation. Thus, in cancer cells where lipid depletion raises the

NAD+/NADH ratio, proliferation in serine depleted environments improves when lipids are also

depleted. Taken together, these data suggest that changes in mitochondrial respiration in

response to nutrient deprivation can influence the NAD+/NADH ratio in a cell-specific manner to

impact oxidative biomass synthesis and proliferation. Given the complexity of tumor

microenvironments, this work provides a metabolic framework for understanding how levels of

more than one environmental nutrient affect cancer cell proliferation.

Introduction

Rapidly proliferating cells, including cancer cells, must acquire biomass precursors such

as amino acids, lipids, and nucleotides to support cell growth and division (Deberardinis, 2008;

Faubert, 2020; Hanahan, 2011; Pavlova, 2017; Vander Heiden, 2017). While cells can obtain

biomass precursors from the environment, many tissue and tumor environments can be deficient

in necessary nutrients (Cognet, 2024; Gullino, 1964; Lyssiotis, 2017). In these conditions, cells

depend on synthesis pathways to accumulate sufficient biomass for proliferation (Apiz Saab, 2023;

Ferraro, 2021; Ngo, 2020; Sciacovelli, 2022; Sullivan, 2019b; Sullivan, 2019a). Synthesis of many

biomass precursors involve oxidation reactions that require the redox cofactor NAD+ to act as an

electron acceptor. There is accumulating evidence that NAD+ availability and the cell redox state

(NAD+/NADH ratio) can limit the synthesis of biomass precursors, including aspartate (Birsoy,

2015; Sullivan, 2015), asparagine (Krall, 2021), fatty acids (Li, 2022), serine (Baksh, 2020; Bao,

2016; Broeks, 2023; Diehl, 2019), and nucleotides (Bao, 2016; Diehl, 2019; Wu, 2024). Thus, the

NAD+/NADH ratio impacts cancer cell proliferation in nutrient environments that increase

dependence on biosynthetic oxidation reactions. However, what determines the cell NAD+/NADH

ratio and whether the NAD+/NADH ratio is modulated in response to environmental nutrient

availability is not well characterized. Moreover, whether differences in the NAD+/NADH ratio

across cell types affect sensitivity to different nutrient limitations is not known.

Some tumor microenvironments have low levels of serine, an amino acid used to produce

many biomass components, including proteins, nucleotides, glutathione, and lipids. Thus, low

serine environments can limit cancer cell proliferation and tumor growth (Banh, 2020; Maddocks,

2013; 2017; Ngo, 2020; Sullivan, 2019b). In serine depleted environments, cancer cells must

increase serine synthesis to maintain proliferation (DeNicola, 2015; Labuschagne, 2014;

Locasale, 2011; Possemato, 2011; Sullivan, 2019b). Serine synthesis involves the conversion of

the glycolytic intermediate 3-phosphoglycerate (3-PG) into serine. The production of 3-PG from

glucose involves an oxidation reaction catalyzed by glyceraldehyde 3-phosphate dehydrogenase

(GADPH). 3-PG is then oxidized by phosphoglycerate dehydrogenase (PHGDH) in the first step

of the serine synthesis pathway. Both GADPH and PHGDH require NAD+ as an electron acceptor

to enable serine synthesis. Experimentally blunting NAD+ regeneration from NADH reduces

serine synthesis and is rescued by oxidizing the NAD+/NADH ratio via exogenous electron

acceptor supplementation (Bao, 2016; Diehl, 2019; Gravel, 2014). Similarly, de novo fatty acid

synthesis can be constrained by the NAD+/NADH ratio, as the production of fatty acids involves

multiple oxidation reactions to generate citrate from either glucose or glutamine. Thus, decreasing

the NAD+/NADH ratio sensitizes cells to environmental lipid depletion (Li, 2022). Together, these

findings demonstrate that exogenous changes to the NAD+/NADH ratio can alter serine and

citrate production and impact proliferation in serine- or lipid-depleted environments. However,

auxotrophy for a single nutrient is not sufficient to determine whether cancer cells can form tumors

in a tissue, where levels of variable nutrients may be environmentally limiting for proliferation

(Abbott, 2026; Sivanand, 2024; Sullivan, 2019a).Thus, we aimed to uncover how the endogenous

NAD+/NADH ratio of cancer cells is regulated to influence biosynthesis in different nutrient

depleted environments and its role in determining cancer cell proliferation when multiple oxidized

nutrients are limiting.

By culturing cancer cells in serine depleted conditions and examining the relationship

between the NAD+/NADH ratio and serine synthesis, we find that the NAD+/NADH ratio is

elevated in response to serine deprivation in some but not all cancer cells. The elevated

NAD+/NADH ratio is driven by increased mitochondrial respiration and promotes serine synthesis

to confer resistance to environmental serine depletion. A similar relationship between

mitochondrial respiration and the NAD+/NADH ratio was also observed in lipid depleted

environments, where higher mitochondrial respiration in some cells led to elevated NAD+/NADH

ratios and enhanced citrate synthesis. Lastly, we find that any perturbation that increases the

NAD+/NADH ratio, including lipid deprivation, could paradoxically improve the proliferation of cells

in serine depleted conditions. Together, these data reveal that the NAD+/NADH ratio is regulated

by mitochondrial respiration in response to environmental nutrients in a cancer cell-specific

manner and that this influences oxidative biosynthesis and proliferation. These data also provide

insight into how levels of multiple nutrients in tissue microenvironments may cooperate to enable

biomass synthesis and cell proliferation.

Results

Modulating the cell NAD+/NADH ratio proportionally alters serine synthesis

To begin investigating the processes that determine the cell NAD+/NADH ratio and

influence biomass synthesis, we examined the relationship between the NAD+/NADH ratio and

serine synthesis in serine-replete conditions (400 µM serine, the concentration of serine in DMEM)

and serine depleted conditions (DMEM media formulated without serine). Previous studies have

found that decreasing the cell NAD+/NADH ratio (more reduced) can suppress serine production

(Baksh, 2020; Bao, 2016; Broeks, 2023; Diehl, 2019). To confirm this, we varied the cell

NAD+/NADH ratio upon serine withdrawal and measured the serine synthesis rate of A549 non-

small cell lung cancer cells, which transcriptionally upregulate the serine synthesis enzyme

PHGDH (DeNicola, 2015). We increased the NAD+/NADH ratio by treating cells with the

exogenous electron acceptor alpha-ketobutyrate (AKB), which can be reduced to alpha-

hydroxybutyrate to regenerate NAD+ (Sullivan, 2015) (**Figure 1A**). We decreased the

NAD+/NADH ratio by treating cells with rotenone, an inhibitor of mitochondrial complex I, which

regenerates NAD+ in support of respiration. AKB and rotenone dose-dependently increased and

decreased the cell NAD+/NADH ratio, respectively (**Figure 1B**). To measure the serine synthesis

rate in cells, we performed kinetic isotope tracing using uniformly ^{13}C-labeled glucose (U-^{13}C-

glucose) and measured production of M+3 labeled serine (**Figure 1C**). Because intracellular

serine, including newly synthesized serine, rapidly exchanges with the extracellular environment

(Labuschagne, 2014), we collected both media and cells to help improve the detection of newly

synthesized serine produced from glucose. We find that increasing the cell NAD+/NADH ratio

with AKB led to proportionally higher serine synthesis rates whereas lowering the NAD+/NADH

ratio with rotenone led to proportionally lower serine synthesis rates (**Figure 1D**). This was not

explained by changes in PHGDH protein expression, as altering the NAD+/NADH ratio with AKB

or rotenone did not affect PHGDH protein levels despite modulating serine synthesis

(**Supplementary Figure 1A,B**). Additionally, neither rotenone treatment nor serine deprivation

had a large effect on cell viability (**Supplementary Figure 1C**). Of note, we find that serine

synthesis rate is positively correlated with the cell NAD+/NADH ratio (**Figure 1E**). Together, these

data demonstrate that modulating the NAD+/NADH ratio leads to proportional changes in serine

synthesis rates.

Cancer cells with elevated NAD+/NADH ratios following serine withdrawal exhibit greater

serine synthesis

Based on our finding that the NAD+/NADH ratio correlates with serine synthesis rate, we

wondered whether endogenous cell NAD+/NADH ratios, which can differ across cancer cells, are

predictive of proliferation rate following serine deprivation. To test this, we measured how

proliferation rate varied in a panel of cancer cells derived from various tissues-of-origin and with

different genetic driver mutations, and found a wide range of sensitivity to serine deprivation

(**Figure 2A**, **Supplementary Figure 2A, Supplementary Table 1**). Because PHGDH expression

is important for proliferation in serine depleted conditions (DeNicola, 2015; Locasale, 2011;

Maddocks, 2017; Possemato, 2011; Sullivan, 2019b), we first assessed the protein expression of

the serine synthesis enzymes PHGDH, PSAT1, and PSPH across the different cancer cells to

examine how well this explained environmental serine dependence. As expected, cells with low

serine synthesis enzyme protein levels (MCF7, MDA-MB-231) were most sensitive to serine

withdrawal while cells with higher serine synthesis enzyme protein levels were more resistant to

serine withdrawal (PHGDH-overexpressing MDA-MB-231, A375) (**Supplementary Figure 2A-C**).

However, for many cancer cells (Calu6, A549, 8988T, MIA PaCa-2, H1299, HCT116), serine

synthesis enzyme expression alone could not fully explain sensitivity to serine withdrawal

(**Supplementary Figure 2C**). For example, when comparing the non-small cell lung cancer cells

A549 and H1299, H1299 cells express less PHGDH protein and similar PSAT1 and PSPH protein

levels than A549 cells but are more resistant to serine deprivation **(Supplementary Figure 2B-**

E). Consistent with lower PHGDH expression, H1299 cells express lower levels of NRF2, a

transcriptional regulator of serine biosynthesis enzymes (DeNicola, 2015). When examining the

cancer cells where serine synthesis enzyme expression could not fully explain the differences in

proliferation upon serine depletion, there was a surprising negative correlation between

proliferation and PHGDH protein expression, no relationship between proliferation and PSAT1

protein expression, and a positive correlation between proliferation and PSPH protein expression

(**Supplementary Figure 2F**). Given our finding that the NAD+/NADH ratio correlates with serine

synthesis rate, we hypothesized that differences in the NAD+/NADH ratio upon serine deprivation

could contribute to the variability in sensitivity to serine withdrawal across cells with similar serine

synthesis pathway protein levels. To test this, we measured the cell NAD+/NADH ratios in the

presence and absence of exogenous serine. Interestingly, we find that serine starvation elevated

the NAD+/NADH ratio in some cells, which we term "redox responder cells" and label in blue. In

contrast, we also find that certain cancer cells minimally change the NAD+/NADH ratio in

response to serine depletion, which we term "redox non-responder cells" and label in yellow

(**Figure 2B, Supplementary Figure 2G**). Of note, whether the NAD+/NADH ratio of a cell was

more or less oxidized in serine-replete conditions was not predictive of response to serine

withdrawal (**Supplementary Figure 2G**). Strikingly, redox responder cells were more resistant to

serine depletion compared to redox non-responder cells (**Figure 2C**). The NAD+/NADH ratio was

also elevated in cancer cells with higher serine synthesis enzyme levels upon serine depletion

and were the most resistant to serine depletion (A375, PHGDH-overexpressing MDA-MB-231)

(**Supplementary Figure 2B,C,G**). To confirm that an elevated NAD+/NADH ratio in response to

serine depletion was associated with elevated serine synthesis, we compared how serine

synthesis rates changed following serine withdrawal in A549 and H1299 cells. We performed

focused comparisons between A549 and H1299 cells because they exhibit differences in

proliferation upon serine deprivation that are not explained by PHGDH protein expression,

demonstrate differing responses of the cell NAD+/NADH ratio upon serine deprivation, and have

similar basal proliferation rates. We find that after 24 hours in serine depleted conditions, H1299

cells demonstrated a higher rate of serine synthesis, a greater increase in serine synthesis, and

greater intracellular serine levels than A549 cells, consistent with the elevated NAD+/NADH ratio

in H1299 cells (**Figure 2D,E; Supplementary Figure 2H,I**). Importantly, we confirmed kinetic U-

^{13}C-glucose tracing was performed at metabolic steady state by ensuring metabolite levels were

stable at each collected time point (**Supplementary Figure 2I**). Taken together, these data

suggest that in some cells, the cell NAD+/NADH ratio is responsive to serine availability. Moreover,

increases in the NAD+/NADH ratio correlate with increased serine synthesis and proliferation

rates in serine depleted conditions independent of PHGDH protein expression.

Increased mitochondrial respiration is associated with an elevated NAD+/NADH ratio and

increased serine synthesis

The finding that some cancer cells exhibit a more oxidized NAD+/NADH ratio upon serine

starvation is surprising, particularly when increased serine synthesis will result in a higher demand

for NAD+ to support increased flux through GAPDH and PHGDH (Diehl, 2019; Maddocks, 2017).

We hypothesized that a more oxidized NAD+/NADH ratio could support greater serine synthesis

and thus sought to identify the processes that increase the NAD+/NADH ratio in some but not all

cancer cells. Lactate production via lactate dehydrogenase (LDH) is a major NAD+ regenerating

process, particularly in cancer cells with high glucose fermentation (Fantin, 2006; Luengo, 2021;

Wang, 2022) (**Supplementary Figure 3A**); however, lactate secretion relative to glucose

consumption was not elevated following serine deprivation in any cells tested (**Supplementary**

Figure 3B). The NAD+ salvage pathway regenerates NAD+ from nicotinamide via nicotinamide

phosphoribosyl transferase (NAMPT) and nicotinamide mononucleotide adenylyltransferase

(NMNAT) and can be important for supporting PHGDH-driven serine synthesis (**Supplementary**

Figure 3A) (Murphy, 2018). While A549 cells were more sensitive to the NAMPT inhibitor FK866

in serine-replete conditions, there was no difference in the sensitivity of A549 and H1299 cells to

FK866 in serine depleted conditions (**Supplementary Figure 3C**). These data suggest that redox

responder and non-responder cells do not have differential dependency on the NAD+ salvage

pathway when cultured without serine. Moreover, supplementing redox non-responder A549 cells

with nicotinamide mononucleotide (NMN), a precursor to NAD+, did not improve proliferation in

serine depleted conditions (**Supplementary Figure 3D**). Mitochondrial respiration is another

major way cells regenerate NAD+ from NADH (Handy, 2012). Thus, we tested whether changes

in mitochondrial respiration, as measured by cellular oxygen consumption rate (OCR),

accompany the increase in the NAD+/NADH ratio following serine withdrawal. Indeed, cells that

increased their NAD+/NADH ratio following serine deprivation also exhibited increased OCR,

whereas cells with unaltered NAD+/NADH ratios following serine withdrawal did not (**Figure 3A,**

Supplementary Figure 3E). To test whether elevated oxygen consumption was related to serine

deprivation, we cultured H1299 and A549 cells in varying amounts of extracellular serine and

measured OCR. Oxygen consumption of A549 cells was not impacted by varying extracellular

serine levels, while oxygen consumption of H1299 cells increased as extracellular serine

decreased below 100 µM, the serine concentration where proliferation begins to be affected by

serine limitation (**Figure 3B, Supplementary Figure 3F**). The increase in OCR also

corresponded with the NAD+/NADH ratio at different serine levels in H1299 cells, while the

NAD+/NADH ratio in A549 cells was unchanged by extracellular serine availability (**Figure 3C,**

D). Consistently, the NAD+/NADH ratio of H1299 cells cultured in different concentrations of

serine positively correlated with serine synthesis, whereas A549 cells displayed minimal changes

in serine synthesis rate (**Figure 3E, F**). Notably, PHGDH protein levels remained constant as

extracellular serine levels were decreased in both A549 and H1299 cells (**Supplementary Figure**

3G). Together, these findings suggest that elevated NAD+/NADH ratios in response to decreasing

environmental serine are linked to elevated mitochondrial respiration in select cancer cells and

correlate with increased serine synthesis.

Mitochondrial respiration governs the cell NAD+/NADH ratio and influences serine

synthesis rate

To test whether increased mitochondrial respiration following serine depletion causes an

increase in the cell NAD+/NADH ratio, we treated redox non-responder cells with the proton

ionophore FCCP (trifluoromethoxy carbonylcyanide phenylhydrazone), which pharmacologically

increases mitochondrial oxygen consumption by uncoupling respiration from ATP synthesis

(McLaughlin, 1980; Terada, 1990). Increasing respiration with FCCP elevates mitochondrial

NAD+ regeneration and the cell NAD+/NADH ratio (Luengo, 2021) (**Figure 4A**). Indeed, FCCP

treatment improved proliferation of redox non-responder cells in serine depleted conditions

(**Figure 4B, Supplementary Figure 4A**). In contrast, FCCP did not improve the proliferation of

redox responder cells cultured without serine (**Figure 4C, Supplementary Figure 4B**).

Consistent with its impact on proliferation, FCCP treatment led to higher serine synthesis rates

only in redox non-responder A549 cells while having no impact on the serine synthesis rates of

redox responder H1299 cells (**Figure 4D,E, Supplementary Figure 5A-D**). As expected,

uncoupling mitochondrial respiration from ATP synthase with FCCP led to elevated oxygen

consumption in serine depleted A549 cells (**Figure 4F).** FCCP also increased oxygen

consumption in serine depleted H1299 cells (**Supplementary Figure 4C**). Interestingly, FCCP

raised the cell NAD+/NADH ratios of both A549 and H1299 cells following serine withdrawal

despite only improving serine synthesis and proliferation in A549 cells (**Figure 4G,**

Supplementary Figure 4D). The lack of impact on serine synthesis by FCCP in H1299 cells

suggests that NAD+ availability may not further constrain serine synthesis in cells that already

elevate mitochondrial respiration following environmental serine depletion. In this context, we

considered whether another constraint such as PHGDH protein expression was limiting serine

synthesis in redox responder cells treated with FCCP. Indeed, overexpressing PHGDH in serine

depleted redox responder H1299 and HCT116 cells fully restored proliferation to that observed in

serine-replete conditions (**Supplementary Figure 4E,G**). In contrast, proliferation of the redox

non-responder cells A549 and Calu6 following serine withdrawal had the greatest improvement

with simultaneous PHGDH overexpression and FCCP treatment (**Supplementary Figure 4F,G**).

255 This underscores that changes in both the cell NAD+/NADH ratio and relevant enzyme expression

256 impact serine synthesis in serine depleted environments. As an orthogonal approach to FCCP,

257 we constitutively expressed the NADH oxidase from *Lactobacillus brevis* (*Lb*NOX) in both the

258 cytoplasm and the mitochondria of A549 cells (**Supplementary Figure 6A**). We reasoned that if

259 expressing *Lb*NOX in either the cytoplasm or the mitochondrial were to raise the cell NAD+/NADH

260 ratio, proliferation of A549 cells upon serine depletion would be improved. To ensure *Lb*NOX

261 activity was present in cells, we treated cells with rotenone and measured proliferation. As

262 expected, we observed a proliferation improvement when *Lb*NOX was expressed in either the

263 cytoplasm or the mitochondria (**Supplementary Figure 6B**). However, A549 cyto*Lb*NOX and

264 mito*Lb*NOX cells did not exhibit an elevated cell NAD+/NADH ratio in either serine-replete or

265 serine depleted conditions (**Supplementary Figure 6C**), consistent with prior observations (Titov,

266 2016). There was also no change in proliferation upon serine withdrawal when *Lb*NOX was

267 expressed in the cytoplasm or mitochondria (**Supplementary Figure 6D**).

268  Because FCCP treatment led to a robust improvement in proliferation of redox non-

269 responder cells cultured without serine, we aimed to test whether FCCP improved serine

270 synthesis and proliferation of redox non-responder cells through oxidizing the NAD+/NADH ratio.

271 We treated cells with the mitochondrial complex I inhibitor rotenone in conjunction with FCCP.

272 Rotenone treatment ablated the ability of FCCP to improve both proliferation and serine synthesis

273 and blocked the ability of FCCP to increase the NAD+/NADH ratio (**Figure 4H-J, Supplementary**

274 **Figure 5E,F**). In contrast, treating cells with oligomycin, an ATP synthase inhibitor, did not affect

275 the ability of FCCP to improve proliferation (**Supplemental Figure 4H,I**). This supports the

276 conclusion that FCCP increases serine synthesis and proliferation in serine depleted

277 environments by increasing the NAD+/NADH ratio via mitochondrial complex I-mediated NAD+

278 regeneration. Moreover, FCCP treatment neither increased PHGDH protein expression nor

279 glucose consumption, indicating that improved proliferation and serine synthesis is not caused by

280 changes in enzyme expression or carbon availability for serine synthesis (**Supplementary Figure**

4J,K). Together, these data argue that mitochondrial respiration, either endogenously increased

in response to serine deprivation, or as a result of FCCP treatment, can govern the cell

NAD+/NADH ratio and directly influence the serine synthesis rate in response to serine withdrawal.

The malate aspartate shuttle enzymes MDH1 and GOT2 support elevated mitochondrial

respiration in redox responder H1299 cells in serine-depleted environments

The malate aspartate shuttle (MAS) transports reducing equivalents between the

cytoplasm and the mitochondria and can be important for de novo serine synthesis (Broeks, 2023).

Thus, we hypothesized that in serine-deplete H1299 cells, the MAS allows increases in

mitochondrial respiration to support cytosolic serine synthesis. To test this, we used

CRISPR/Cas9 in H1299 cells to generate isogenic cell lines where either GOT1, MDH1, or GOT2

was deleted (**Supplementary Figure 7A**). Using these knock-out cell lines, we measured

proliferation in serine depleted conditions and found that MDH1 and GOT2 knock-out cell lines

were more sensitive to serine deprivation compared to non-targeting controls (NTC)

(**Supplementary Figure 7B**). We then measured the cell NAD+/NADH ratio in knock-out cells

cultured without serine and observed that unlike H1299 NTC and GOT1 KO cells, H1299 MDH1

and GOT2 KO cells did not exhibit elevated NAD+/NADH ratios upon serine depletion

(**Supplementary Figure 7C**). Consistently, unlike H1299 NTC cells, H1299 MDH1 and GOT2 KO

cells did not demonstrate elevated mitochondrial respiration upon serine depletion

(**Supplementary Figure 7D**). This corresponded with H1299 MDH1 and GOT2 KO cells having

blunted serine synthesis rates compared to H1299 NTC cells (**Supplementary Figure 7E**).

Together, this suggests that MDH1 and GOT2 activity support the process by which mitochondrial

NAD+ regeneration is transmitted to the cytoplasm to support serine synthesis.

The NAD+/NADH ratio influences oxidative citrate synthesis in either serine- or lipid-

depleted environments

Oxidative biosynthetic reactions other than serine synthesis can also be constrained by

the NAD+/NADH ratio. For example, cancer cells deprived of environmental lipids increase

oxidative citrate production, and we have previously found that citrate synthesis, either through

glucose oxidation or glutamine oxidation, is limited by NAD+ availability (Li, 2022) (**Figure 5A,**

Supplementary Figure 8A). Thus, we sought to uncover whether the increase in the cell

NAD+/NADH ratio by mitochondrial respiration in response to serine withdrawal specifically

supports greater serine synthesis or also leads to greater oxidative citrate production. To answer

this question, we first assessed labeled citrate production using kinetic U-^{13}C-glucose tracing in

cells that showed increased NAD+/NADH ratios following serine deprivation. We found that the

redox responder cells largely showed elevated M+2 citrate production despite no change in

environmental lipid availability (**Figure 5B; Supplementary Figure 8B-E**). In contrast, cells with

unaltered NAD+/NADH ratios upon serine depletion did not exhibit higher M+2 citrate production

(Figure 5B; Supplementary Figure 8B, F-H). Consistently, redox responder H1299 cells, but

not redox non-responder A549 cells, displayed evidence of elevated citrate production from

glutamine either through oxidative decarboxylation or reductive carboxylation, both of which

require oxidation reactions that can be limited by electron acceptor availability (Hosios and Vander

Heiden, 2018; Li, 2022) (**Supplementary Figure 8I, J**). To confirm that elevated M+2 citrate

synthesis was driven by an increase in the NAD+/NADH ratio, we treated A549 cells depleted of

serine with FCCP and examined glucose-derived citrate synthesis. Indeed, FCCP treatment led

to a four-fold increase in M+2 citrate from U-^{13}C-glucose in A549 cells depleted of serine whereas

M+2 citrate synthesis was less affected by FCCP in H1299 cells (**Figure 5C**). The elevated M+2

citrate production in FCCP-treated A549 cells was dependent on mitochondrial complex I, as

rotenone prevented the elevated citrate production associated with FCCP treatment

(**Supplementary Figure 9A-C**). To confirm that higher M+2 citrate production following FCCP

treatment was related to the NAD+/NADH ratio independent of serine deprivation, we treated

A549 cells with increasing doses of AKB and FCCP in serine-replete conditions and find that

increasing the NAD+/NADH ratio with either AKB or FCCP led to elevated M+2 citrate synthesis

(**Figure 5D**). This argues that changes to the NAD+/NADH ratio regardless of the nutrient

environment can impact citrate synthesis.

We next asked whether increasing mitochondrial respiration in response to low

extracellular serine was specific to serine deprivation or represented a general response to any

nutrient deprivation that increased NAD+ demand, such as lipid deprivation (Li, 2022). To examine

this, we depleted A549 and H1299 cells of extracellular lipids by culturing cells in media with lipid-

depleted serum (Hosios, 2018) and compared oxygen consumption rates to those of cells cultured

in media with lipid-containing serum. Surprisingly, we find that A549 cells, which do not increase

mitochondrial respiration in response to serine depletion, showed a trend towards a mildly

elevated oxygen consumption rate upon lipid depletion (**Figures 5E**). On the other hand, H1299

cells do not increase oxygen consumption in lipid depleted conditions (**Figure 5E**). We next

measured the NAD+/NADH ratio of A549 and H1299 cells depleted of environmental lipids.

Consistent with the changes in oxygen consumption, A549 cells displayed an increased

NAD+/NADH ratio while H1299 cells did not alter the NAD+/NADH ratio upon lipid deprivation

(**Figure 5F**). Interestingly, both serine and citrate produced via glucose oxidation was elevated in

lipid depleted A549 cells in comparison to H1299 cells, which exhibited unchanged serine

synthesis in response to environmental lipid depletion (**Figure 5G-H, Supplementary Figure 10**).

These findings suggest that the NAD+/NADH ratio can be a determinant of both oxidative citrate

and serine synthesis, regardless of environmental demand for the nutrient. Additionally, the

different metabolic responses to lipid and serine deprivation between cancer cell types suggests

that different nutrient environments impact mitochondrial respiration through distinct processes to

influence the NAD+/NADH ratio.

Lipid depletion can enhance proliferation in serine deprived conditions

Our finding that both serine and oxidative citrate synthesis can be impacted by levels of a

different nutrient than that produced by each pathway raised the possibility that the NAD+/NADH

ratio coordinates biomass synthesis in response to the availability of multiple oxidized nutrients.

Specifically, we hypothesized that changes to the NAD+/NADH ratio caused by one oxidized

nutrient depletion could influence the synthesis of, and in turn the dependency on, another

oxidized nutrient for proliferation. To begin testing this, we first examined how mitochondrial

respiration and the NAD+/NADH ratio were impacted by dual lipid and serine depletion compared

to single nutrient depletions in both A549 and H1299 cells. We find that dual depletion of serine

and lipids in A549 cells led to elevated oxygen consumption that is higher than that observed with

serine depletion alone **(Figure 6A)**. Similarly, dual depletion of serine and lipids in H1299 cells

led to elevated oxygen consumption compared to that observed with lipid depletion alone **(Figure**

6A). The resulting oxygen consumption rate corresponded with the cell NAD+/NADH ratio in each

condition, which was collectively higher upon dual serine and lipid starvation than either serine

depletion in A549 cells or lipid depletion in H1299 cells (**Figure 6B**). We then measured the serine

and citrate synthesis rates of cells depleted of both extracellular serine and lipids via kinetic U-

^{13}C-glucose tracing. Consistent with elevation of the NAD+/NADH ratio after lipid deprivation,

A549 cells depleted of both environmental serine and lipids trended towards elevated serine

synthesis rates when compared to serine deprivation alone, though was not statistically significant

(Figure 6C). Moreover, lipid depletion appeared to lead to a greater fraction of total serine derived

from glucose in serine depleted A549 cells **(Supplementary Figure 11A)**. In contrast, lipid

starvation did not alter serine synthesis in serine deprived H1299 cells (**Figure 6D,**

Supplementary Figure 11A,B). Interestingly, in both A549 and H1299 cells, dual serine and lipid

deprivation led to a trend toward increased citrate production compared to lipid deprivation alone

despite only H1299 cells exhibiting a more oxidized NAD+/NADH upon serine depletion (**Figure**

6E,F; Supplementary Figure 11C,D). This underscores the multifactorial regulation of glucose

oxidation to citrate (Holness, 2003; Novelli, 1950; Patel, 2006; Srere, 1974). We next asked how

cancer cell proliferation was impacted by dual serine and lipid depletion compared to single

nutrient depletions. Strikingly, proliferation of serine depleted A549 cells was improved with

depletion of extracellular lipids (**Figure 6G**), consistent with the elevation in mitochondrial

respiration, NAD+/NADH ratio, and a trend towards greater serine synthesis in A549 cells upon

dual serine and lipid depletion. Unlike serine depletion, lipid depletion does not cause a major

proliferation defect in either A549 or H1299 cells, and dual serine and lipid deprivation did not

greatly alter the proliferation of H1299 cells compared to serine depletion alone (**Figure 6H**). This

is consistent with lipid deprivation causing no major changes in oxygen consumption,

NAD+/NADH ratio, and serine synthesis in H1299 cells. These data argue that the cell

NAD+/NADH ratio is modulated in response to multiple nutrient availabilities in a cancer cell-

specific fashion. The changes in NAD+/NADH ratio in turn may influence the capacity for serine

and citrate synthesis, impacting the proliferation rate of cancer cells in nutrient environments with

differing demands for oxidative biomass synthesis. Taken together, we propose a model where

environmental nutrient availability can impact mitochondrial respiration based on the specific

cancer. Because mitochondrial respiration is a major pathway that regenerates NAD^+, changes

to mitochondrial respiration can alter the cell NAD+/NADH ratio, influencing the activity of major

NAD^+-requiring metabolic reactions such as serine synthesis and citrate synthesis that can be

important for proliferation. We further propose that changes to the cell NAD+/NADH ratio can

impact all oxidative biosynthetic reactions if the enzyme machinery is present, but that specificity

for how the cell NAD+/NADH ratio changes is dependent on both cell-intrinsic factors and cell-

extrinsic factors (**Figure 7**).

Discussion

Proliferating cancer cells must acquire sufficient biomass and rely on contributions from

both environmental nutrients and de novo synthesis pathways, but what determines which cells

can adapt to different nutrient environments that impose constraints on salvage and synthesis is

not known. This study highlights that differential cell responses to changes in the NAD+/NADH

ratio enable biomass synthesis and determine whether cells can adapt to proliferate in different

nutrient conditions. Importantly, this provides one explanation for why single nutrient levels in

tissues may not predict the ability of auxotrophy for that nutrient to determine cancer proliferation

in a tissue (Abbott, 2026). It also suggests why cancer proliferation in different nutrient

environments is controlled by a complex relationship between cell-intrinsic and cell-extrinsic

factors. These factors may be coordinated in part through processes that set the NAD+/NADH

ratio in cells.

Most prior work to determine what shapes the metabolic dependencies of cell proliferation

has focused on factors that govern biosynthetic enzyme expression and activity. For example,

genomic and transcriptional analyses of cancer discovered that increased PHGDH expression

promotes tumor growth (DeNicola, 2015; Locasale, 2011; Newman, 2017; Possemato, 2011) and

provides a proliferative advantage in serine depleted tumor microenvironments (Ngo, 2020;

Sullivan, 2019b). Our findings show that enzyme expression alone cannot fully explain nutrient

dependencies, as PHGDH expression can be discordant with sensitivity to serine depletion.

Rather both the NAD+/NADH ratio and enzyme expression play integral roles in determining

cancer proliferation in specific nutrient environments. These data are consistent with biosynthetic

gene expression and single nutrient levels being unable to predict the tumor tissue environments

where cancer cells can form metastases (Abbott, 2026).

Extensive work has highlighted the importance of cell NAD+/NADH ratio in influencing

oxidized biomass production, nutrient utilization, and nutrient dependencies (Baksh, 2020; Bao,

2016; Birsoy, 2015; Broeks, 2023; Christensen, 2014; Diehl, 2019; Garcia-Bermudez, 2018;

Gibson, 1976; Krall, 2021; Li, 2022; Sullivan, 2015; Wu, 2024). Yet, the processes that govern

the endogenous NAD+/NADH ratio upon shifts in nutrient availability have been largely

uncharacterized. We find that regulation of mitochondrial respiration is a major mechanism that

dictates the NAD+/NADH ratio in response to environmental nutrient fluctuations. The

mitochondrial respiration response to withdrawal of specific nutrients appears to vary across

different cancer cells. Some cancer cells increased mitochondrial respiration to serine depletion,

while others had minimal responses. We found that in serine redox responder cancer cells, the

malate aspartate shuttle (MAS) enzymes MDH1 and GOT2 were required for the elevation in

mitochondrial respiration and the NAD+/NADH ratio in response to serine depletion. Thus,

differences in the activity of MAS, which can vary greatly across cancer cells (Wang, 2022), may

influence the capacity for cancer cells to increase mitochondrial respiration and NAD+

regeneration to support oxidative biosynthetic pathways. Interestingly, serine is an allosteric

activator of the glycolytic enzyme pyruvate kinase, which converts phosphoenolpyruvate to

pyruvate and generates ATP (Chaneton, 2012). Thus, decreased environment serine availability

in addition to differences in pyruvate kinase activity may yield lower glycolytic ATP, resulting in

greater mitochondrial respiration in serine redox responder cancer cells. Additionally, it has been

recently shown that differences in mtDNA in non-small cell lung cancer cells can lead to defective

mitochondrial respiration and imbalanced NAD+/NADH ratios, suppressing serine synthesis and

tumor growth (Lopes, 2025). These findings suggest some reasons why respiration may be

differentially responsive to serine withdrawal. However, lipid depletion yielded mitochondrial

respiration changes in cancer cells that did not respond to serine withdrawal, arguing against

respiration defects explaining non-response. No clear common genetic factor or tissue-of-origin

was found to align with those cells that do or do not have mitochondrial and redox responses to

serine and/or lipid depletion. Further work is needed to understand the mechanisms that enable

increased mitochondrial respiration in response to deprivation of different nutrients. One

possibility is that differences in ATP consumption following nutrient deprivation contribute to the

response. Stimulating ATP consumption can increase mitochondrial respiration and the

NAD+/NADH ratio (Bertholet, 2019; Buttgereit, 1995; Fisher-Wellman, 2012; Luengo, 2021;

Nobes, 1989; Vander Heiden, 1999). ATP synthase activity can restrain complex I activity to

different extents across cancers, modulating the endogenous NAD+/NADH ratio and sensitivity

to serine or lipid depletion. While the regulation of complex I, ATP synthesis, and mitochondrial

coupling efficiency is complicated and incompletely understood (Alavian, 2014; Gao, 2018;

Lapuente-Brun, 2013; Luengo, 2021; Qiu, 2014; Wang, 2022; Yao, 2019; Zhao, 2019), our

findings suggest an important connection between the regulation of mitochondrial bioenergetics

and specific metabolic vulnerabilities of cancer growth, underscoring an important relationship

between environmental nutrient levels and mitochondrial respiration to influence the cell

NAD+/NADH ratio.

Efforts to understand the processes that shape cancer cell metabolic dependencies have

largely focused on how cancer cells respond to single nutrient changes. However, findings from

dual serine and lipid depletion experiments suggest that single nutrient availabilities will be unable

to explain cancer metabolic dependencies in some contexts. For example, depleting A549 cells

of both extracellular serine and lipids paradoxically improved proliferation relative to serine

deprivation alone, consistent with an elevated NAD+/NADH ratio in response to lipid deprivation

that can support greater serine synthesis. This suggests that the NAD+/NADH ratio can serve as

a metabolic regulatory node that orchestrates biomass synthesis and integrates changes to

multiple oxidized biomass components in the environment, providing a framework for

understanding why cancer nutrient dependencies vary across tissue microenvironments (Abbott,

2026). Because oxidizing the cell redox state can increase flux through multiple pathways, the

impact of cell redox state on biosynthesis may also explain why some cells lose enzyme

expression in pathways like serine synthesis, as this could redirect limited resources toward

biomass components that can otherwise be scavenged from the environment (Delage, 2010;

Pavlova, 2017; Rabinovich, 2015). Better understanding the mechanisms cells use to alter

respiration and adjust the NAD+/NADH ratio in response to available nutrients could inform the

complex interplay between cell-intrinsic and cell-extrinsic factors that determine cancer metabolic

dependencies. This is particularly important to consider when targeting metabolism for cancer

treatment. Many newer therapies targeting metabolism have not been successful in part because

of metabolic plasticity to nutrient shifts (Amoedo, 2017; Fendt, 2020; Xiao, 2023). Co-targeting

mitochondrial function limits metabolic adaptations and may also help predict the tissue nutrient

conditions that result in pathway dependencies for specific cancers. Thus, better understanding

how the cell NAD+/NADH ratio responds to nutrient levels in different cancers could improve

selection of patients for cancer therapies that impact metabolism.

Material and Methods

Cell Culture Experiments

Cell lines were maintained in Dulbecco's Modified Eagle's Medium (DMEM) without sodium

pyruvate (Corning 50-013-PC) supplemented with sodium bicarbonate (Sigma S6014) and 10%

heat-inactivated Fetal Bovine Serum (FBS). All cells were cultured at 37°C with 5% CO_2, tested

for mycoplasma contamination regularly, and confirmed negative before experimentation. Cell

line information is listed in Supplementary Table 1.

Proliferation Assays

Cells were plated in six-well plates in 2 ml of DMEM with 10% heat-inactivated FBS at an initial

seeding density of 40,000 cells. Cells were permitted to settle overnight and then quantified using

a sulforhodamine B (SRB) (Sigma Aldrich 230162) colorimetric assay to measure the starting cell

absorbance at the start of the experiment as previously described (Vichai, 2006). For treatment

conditions, cells were washed three times with 1 ml of 1X PBS, and 2 ml of treatment media were

added to each well. All treatment media was made with 10% dialyzed FBS. Serine-free medium

was made by adding a mix of amino acids (Sigma Aldrich) at DMEM concentrations lacking serine

and glycine to DMEM with low glucose, without sodium pyruvate, and amino acids (US Biological

D9800-13-50L). Glucose and glycine were supplemented for a final concentration for 25 mM and

0.4 mM, respectively. Three days after the initial treatment, cells were quantified using SRB assay.

All SRB measurements are normalized to a blank to correct for background signal. Proliferation

rates were calculated with the following formula:

Doublings per day = <u>log$_2$(final SRB absorbance / SRB absorbance day 0)</u>

(total # of treatment days)

Cell Number and Viability Measurements

Cells were seeded at an initial density of 2,000 cells per well in a 96-well plate in 200 μl of

treatment media. After allowing cells to settle for 1-2 hours, cell number was monitored over time

using imaging with the Incucyte Live-Cell Analysis System (Sartorius). Doublings per day were

calculated between hour 0 of monitoring and hour 72 of monitoring using the following formula:

Doublings per day = <u>log$_2$(cell number at hour 72 / cell number at hour 0)</u>

(3 days of treatment)

For viability, cells were seeded at an initial density of 2,000 cells per well in a 96-well plate in 200

μl of treatment media with 20 nM Sytox Green (Invitrogen S7020). After 72 hours, the number of

live and dead cells present was monitored using imaging with the Incucyte Live-Cell Analysis

System (Sartorius).

NAD+/NADH Measurements

NAD+/NADH measurements were performed using the NAD/NADH-Glo Assay (Promega G9072)

with a modified version of manufacturer instructions as previously reported (Sullivan, 2015). Cells

were plated at an initial seeding density of 20,000 cells (A549, H1299, HCT116) or 40,000 cells

(Calu6, MDA-MB-231). Cells were permitted to settle overnight. The next day, cells were washed

once with 1X PBS and then incubated with 2 ml of treatment media for the indicated times prior

to preparation of cell extracts. For extraction, cells were washed once in ice cold 1X PBS and

extracted in 100 μl ice-cold 1% dodecyltrimethylammonium bromide (DTAB) in 0.2N NaOH diluted

1:1 with 1X PBS. Each sample was flash-frozen in liquid nitrogen and immediately stored at -

80°C. To measure NADH, 10 µl of sample was moved to PCR tubes, diluted with 10 µl of DTAB,

and incubated at 75 °C for 30 min, where basic conditions selectively degrade NAD+. To measure

NAD+, 10 µl of the samples was moved to PCR tubes containing 30 µl DTAB and 20 µl 0.4 N HCl

and incubated at 60°C for 15 min, where acidic conditions selectively degrade NADH. Samples

were then allowed to equilibrate to room temperature and quenched by neutralizing with 20 µl

0.25 M Tris in 0.2 N HCl (for NADH) or 20 µl 0.5 M Tris base (for NAD+). Manufacturer instructions

were followed thereafter to measure NAD+/NADH using a luminometer (Tecan Infinite M200Pro).

A standard curve with representative samples was done with each assay to confirm that NAD+

and NADH measurements were in the linear range of detection.

Glucose Consumption and Lactate Secretion Measurements

Cells were plated at a density of 40,000 cells per well in a six well plate. The next day, cells were

washed three times with 1X PBS before 2 ml of treatment media were added to each well. In

parallel, 2 ml of treatment media were added to plates without cells. An initial cell number before

beginning treatment was quantified via Cellometer (Nexcelom Bioscience). 72 hours later, media

was collected from exponentially proliferating cells with a parallel collection of media from wells

without cells, which we used to measure initial media concentrations to take into consideration

evaporative changes. Cell number was quantified via Cellometer along with the volume of media

on cells to consider evaporative changes to metabolite concentrations that may have occurred

over 72 hours in culture. Glucose concentrations from media samples were measured on a YSI-

2900 Biochemistry Analyzer (Yellow Spring Instruments). Every assay included a glucose (Sigma

Aldrich G8270) standard curve (0, 2.5, 5, 10, 20, and 40 mM glucose) and lactate (Sigma Aldrich

71716) standard curve (0, 1.25, 2.5, 5, 10, and 20 mM sodium lactate), made in both control and

treatment media. To calculate consumption rate per cell for a given nutrient, fmol of glucose or

lactate over time were plotted relative to the area under the curve (fmol / cells · time) of an

exponential function fit to the number of cells at the start of the assay and after 72 hours of

treatment, as previously described (Hosios, 2016).

Oxygen Consumption Measurements

An Agilent Seahorse Bioscience Extracellular Flux Analyzer (XFe96) was used to measure

oxygen consumption rates (OCR). Cells were plated at 20,000 – 40,000 cells per well in a

Seahorse Bioscience 96-well plate in 80 µl of DMEM without pyruvate supplemented with 10%

heat-inactivated FBS. Cells were not plated on the perimeter of the plate to avoid edge effect.

The following day, cells were washed twice with 180 µl of treatment media supplemented with 10%

dialyzed FBS before incubating cells with 180 µl of treatment media for the indicated time prior to

OCR data acquisition. For 24 hour treatment periods, treatment media was replaced with 180 µl

of fresh treatment media two hours before OCR data acquisition. The day before OCR data

acquisition, the XFe96 cartridge was hydrated by submerging the sensor cartridge in 200 µl of

sterile water per well of the utility plate. Parafilm was wrapped around the perimeter of the

cartridge to avoid evaporation. The XFe96 cartridge was then incubated in a 37°C non-CO_2

incubator overnight. The next day, at least two hours before data acquisition, the water in the

utility plate was replaced with 200 µl of the Agilent XF Calibrant per well and the sensory cartridge

was submerged and returned to the 37°C non-CO_2 incubator until OCR was ready to be measured.

After OCR data acquisition, six to eight wells per treatment condition were collected for cell

number quantification (Cellometer) to normalize OCR values. Basal OCR was calculated by

subtracting residual OCR, the OCR remaining following the addition of rotenone and antimycin A

for a final concentration of 2 µM.

Kinetic U-[13]C-Glucose Isotope Tracing Experiments

Cells were seeded at 150,000 cells per well in a six-well plate in 2ml of DMEM without sodium

pyruvate supplemented with 10% heat-inactivated FBS. The following day, cells were washed

once with 1ml of 1X PBS and then incubated in 2ml of treatment media supplemented with 10%

dialyzed FBS for the indicated treatment time and when metabolic steady state is reached. Two

hours before the start of the kinetic isotope tracing, media on cells was exchanged with fresh

treatment media with 10mM unlabeled glucose. To measure serine labeling over time in serine

depleted conditions, cells and media were extracted together. Before the initiation of the kinetic

U-^{13}C-glucose isotope tracing, serine-deplete cells were washed three times with 1X PBS and

then cultured with 400 µl of tracing medium containing 10 mM of U-^{13}C-glucose (Cambridge

Isotope Laboratories, CLM-481-0) for the rapid time points. Tracing media was equivalent to

treatment media except with 10 mM of U-^{13}C- glucose. Following incubation, 1.6ml of extraction

buffer containing ice cold 100% HPLC-grade methanol (Sigma Aldrich, 646377) with norvaline

(Sigma, N7627) was added onto cells with tracing media for a final sample volume of 2 ml (80%

extraction buffer and 20% tracing media) and a final concentration of 1 µg per 400 µl norvaline.

1.6 ml of the lysate was placed into a fresh Eppendorf tube, vortexed at 4°C at maximum speed

for ten minutes, and then spun down at 4°C at maximum speed for thirty minutes. The supernatant

was collected and dried under nitrogen gas to prepare for metabolic analysis. To measure serine

and citrate production in serine replete conditions, only intracellular metabolites were quantified.

Before the initiation of kinetic U-^{13}C-glucose isotope tracing, serine-replete cells were washed

three times with 1X PBS and then cultured with 1 ml of tracing medium containing 10 mM of U-

^{13}C-glucose (Cambridge Isotope Laboratories, CLM-481-0) for the rapid time points. Tracing

media was equivalent to treatment media except with 10 mM of U-^{13}C- glucose. Following

incubation, cells were rapidly washed twice with ice cold blood bank saline. To extract, 500 µl of

extraction buffer containing 80% ice cold HPLC-grade methanol and 20% LC-grade water spiked

with 1 µg per 400 µl norvaline were added onto cells. Cells were then lysed as previously

described above.

Gas Chromatography-Mass Spectrometry (GC-MS) Polar Metabolite Measurements

Dried samples were derivatized by adding 16 µl of methoxamine reagent (Thermo Fisher, TS-

45950) and incubated for an hour at 37°C followed by the addition of 20 µl of *N-tert-*

butyldimethylsilyl-*N*-methyltrifluoroacetamide with 1% *tert*-butyldimethylchlorosilane (Sigma

375934). Samples were then incubated for two hours at 60°C. Afterwards, samples were

centrifuged at maximum speed for ten minutes, and 20 µl of supernatant was used for analysis.

Following derivatization, samples were analyzed using a DB-35MS column (30 m × 0.25 mm

i.d. × 0.25 µm, Agilent J&W Scientific) in an Agilent 7890 gas chromatograph coupled to an Agilent

5975C mass spectrometer (GC–MS). All ion counts were normalized to internal standard

norvaline and cell number measured via Cellometer.

Immunoblotting

Cells were washed with ice cold 1X PBS and lysed in cold RIPA buffer containing Halt Protease

and Phosphatase Inhibitor Cocktail (Thermo Scientific 78442). Lysates were clarified by rocking

samples for thirty minutes at 4°C and then spun at maximum speed for ten minutes for collection

of supernatants. Protein concentration was calculating using the BCA Protein Assay (Pierce,

23225) with BSA as a standard. Lysates were resolved by SDS-PAGE using NuPAGE 4-12% Bis-

Tris Protein Gels (Thermo Fisher Scientific) and run at 100V. Proteins were transferred onto

nitrocellulose membranes via wet transfer at 100V. Membranes were blocked in 5% BSA in 1x

TBST before incubating membranes with primary antibodies at 4°C overnight. The next day,

membranes were washed three times with 1x TBST at room temperature and then incubated in

secondary antibodies for one hour at room temperature. The primary antibodies used were

PHGDH (Sigma Aldrich, HPA021241 [1:1000]), PSAT1 (Abcam, ab96136 [1:1000]), PSPH

(Abcam, ab96414 [1:1000]), HSP90 (Cell Signaling Technology, 4874 [1:1000]), Vinculin (E1E9V)

XP® (Cell Signaling Technology, 13901 [1:1000]), GOT1 (Cell Signaling Technology, 34423S

[1:1000], MDH1 (MDHC H-6, Santa Cruz Biotechnology, sc-166879 [1:1000]), GOT2 (Cell

Signaling Technology, 71692S [1:1000]), FLAG (Cell Signaling Technology, 14793 [1:1000]). The

secondary antibodies used were anti-rabbit IgG horseradish peroxidase-linked antibody (1:4000

dilution; Cell Signaling Technologies, 7074S) or anti-mouse IgG horseradish peroxidase-linked

antibody (1:4000, 7076S). Protein expression was quantified by using ImageJ analysis.

Genetic Manipulation of Enzyme Expression in Cells

Cell lines overexpressing either EGFP or human PHGDH were generated via lentiviral infection.

pLJM1-EGFP was obtained from Addgene (Addgene 19319). pLJM1-PHGDH was constructed

using pLJM1-Empty (Addgene 91980) as a backbone. pLJM1-Empty was digested using NheI

and EcoRI and gel purified. Human PHGDH cDNA was amplified from pLHCX-PHGDH (PMID:

31598584) with the following oligonucleotide primers, where the capitalized sequence denotes

PHGDH homology:

PHGDH F: agtgaaccgtcagatccggctagcgccaccATGGCTTTTGCAAATCTGCG

PHGDH R: tactgccatttgtctcgaggtcgagaattcTTAGAAGTGGAACTGGAAGGCT

PHGDH insert was gradient PCR amplified using Phusion High-Fidelity DNA polymerase (NEB

M0530) from 60 to 70°C and gel purified. Digested pLJM1-Empty and amplified PHGDH insert

were assembled via Gibson Assembly, and the resulting plasmid was transformed, and

validated by Sanger sequencing. Lentiviral production was done by transfecting constructs into

LentiX293T cells with Mirus Transit293T (Mirus Bio MIR2700) following manufacturer's protocol.

Briefly, 800,000 cells were plated per well in a 6 well plate in DMEM media with 10% heat-

inactivated FBS. The next day, cells were transfected with 1.6 µg vector (either pLJM1-EGFP or

pLJM1-PHGDH), 800 ng of pMDLg (Addgene 12251) packaging plasmid, 400 ng of pMD2.G

(Addgene 12259) envelope plasmid, and 400 ng of pRSV-REV (Addgene 12253) in OptiMEM

(Thermo Fisher Scientific 31985070). DNA transfection mix was incubated with cells for 48

hours before harvesting virus and filtering through a 0.45 µm polyethersulfone membrane. Sub-

confluent cells were infected with 1 ml of virus-containing media in each well of a 6 well plate

with polybrene (8 µg/ml, EMD Millipore TR-1003-G). 24 hours after infection, selection with 1

µg/mL puromycin (Sigma Aldrich P7255) began. All experiments with generated cell lines

expressing PHGDH were conducted on polyclonal cell populations.

cyto*Lb*NOX-FLAG cDNA and mito*Lb*NOX-FLAG cDNA were PCR amplified from

pUC57-*Lb*NOX or pUC57-mito*Lb*NOX, respectively. pUC57-*Lb*NOX (Addgene 75285) and

pUC57-mito*Lb*NOX (Addgene 74448) were a gift from Vamsi Mootha. The PCR products were

subsequently cloned into pLJM1 using Gibson Assembly (NEB E2611S). pLJM1-EGFP was a

gift from David Sabatini (Addgene 19319). Cells were then transduced with lentivirus containing

pLJM1-*Lb*NOX-FLAG, pLJM1-mito*Lb*NOX-FLAG, or pLJM1-EGFP. The infected cells were

selected in 1 µg/mL puromycin (Sigma Aldrich P7255). All experiments with generated cell lines

involving *Lb*NOX were conducted on a polyclonal cell population.

GOT1, GOT2, and MDH1 Knockout H1299 cells were generated by using lentiCRISPRv2

(Addgene 52961). LentiCRISPRv2 was a gift from Feng Zhang. Guide RNAs (gRNAs) for each

gene of interest were designed by CRISPick

(https://portals.broadinstitute.org/gppx/crispick/public) and cloned into lentiCRISPRv2. The

guide sequences used are as follows:

NTC: GTGTAGTTCGACCATTCGTG

GOT1: GGAGGTGTGCAATCTTTGGG

GOT2: CGGACGCGGGTCCACTCCCG

MDH1: CGTCCAGGACACCCATCATG

Cell lines were generated by transducing cells with viral particles containing lentiCRISPRv2

followed by selection with 1 µg/mL puromycin (Sigma Aldrich P7255). Single cells were plated

into individual wells of 96 well plates in RPMI medium to obtain single clone knock out cells.

Following expansion, immunoblotting was used to confirmed complete knockout.

Statistics and Reproducibility

All statistical tests using experimental data were performed with Prism 10 software. No statistical

method was used to predetermine sample sizes. Samples sizes were chosen based on pilot

experiments using three or more technical replicates. The experiments were not randomized and

the investigators were not blinded during experiments nor data analysis.

Acknowledgements

We thank all members of the Vander Heiden lab for helpful discussions, Emma Dawson, Grace

Wolczanski, and Alejandra Rosario-Crespo for technical assistance and Jessica Spinelli for

assistance with using Seahorse equipment during the COVID-19 pandemic. The authors

acknowledge support from F30CA268633 (SMC), MSTP T32GM007753 (SMC), T32GM007287

(SMC, MBM, ASZ, KLA), the Kosciuszko Foundation (SET), T32GM144273 (HC), NSF DGE-

1122374 (KLA), and F31CA271787 (KLA), as well as support to MGVH from the Ludwig Center

at MIT, the MIT Center for Precision Cancer Medicine, and the NCI (R35CA242379,

R01CA259253, P30CA1405141).

Author Contributions

Conceptualization: SMC, MGVH; Intellectual Discussion: SMC, MBM, MGVH; Proliferation rates:

SMC, SET, ASZ, HC; Incucyte: SMC, MBM; Viability: SMC, MBM; Kinetic Isotope Tracing: SMC,

SET, MBM, HC; Immunoblotting: SMC, SET; NAD+/NADH measurements: SMC, SET; Oxygen

consumption: SMC; PHGDH overexpression constructs: KLA; PHGDH overexpression cell line

generation: SMC; GOT1, GOT2, and MDH1 KO generation: RC; Writing: SMC and MGVH; Editing:

SMC, MBM, SET, ASZ, KLA, MGVH; Funding Acquisition: MGVH

Competing Interests

M.G.V.H. discloses that he is or was a scientific advisor for Agios Pharmaceuticals, iTeos

Therapeutics, Sage Therapeutics, Pretzel Therapeutics, Lime Therapeutics, Faeth Therapeutics,

Droia Ventures, MPM Capital and Auron Therapeutics. All remaining authors declare no competing interests.

References

Abbott, K.L., Subudhi, S., Ferreira, R., Gültekin, Y., Steinbuch, S.C., Bin Munim, M., Ramesh, D.L., Honeder, S.E., Kumar, A.S., Wu, M., Hansen, J.A., Shevzov-Zebrun, A., Rashan, E.H., Eghbalian, K.M., Sivanand, S., Barbeau, A.M., Riedmayr, L.M., Duquette, M., Ali, A., Henning, N., Trojan, S.E., Waite, M., Kunchok, T., Nakano, M.A., Gourgue, F, Ferraro, G.B., Do, B.T., Spanoudaki, V., Sánchez-Rivera, F.J., Jin, X., Church, G.M., Jain, R.K., Vander Heiden, M.G. (2026). Nutrient requirements of organ-specific metastasis in breast cancer. Nature *649*, 1292-1301. https://doi.org/10.1038/s41586-025-09898-9.

Alavian, K.N., Beutner, G., Lazrove, E., Sacchetti, S., Park, H.-A., Licznerski, P., Li, H., Nabili, P., Hockensmith, K., Graham, M., Porter, G. A., & Jonas, E. A. (2014). An uncoupling channel within the c-subunit ring of the F1FO ATP synthase is the mitochondrial permeability transition pore. PNAS *111*, 10580–10585. https://doi.org/10.1073/pnas.1401591111.

Amoedo, N.D., Obre, E., Rossignol, R. (2017). Drug discovery strategies in the field of tumor energy metabolism: Limitations by metabolic flexibility and metabolic resistance to chemotherapy. Biochimica et Biophysica Acta - Bioenergetics *1858*, 674-685. https://doi.org/10.1016/j.bbabio.2017.02.005.

Apiz Saab, J.J., Dzierozynski, L.N., Jonker, P.B., AminiTabrizi, R., Shah, H., Elena Menjivar, R., Scott, A.J., Nwosu, Z.C., Zhu, Z., Chen, R.N., Oh, M., Sheehan, C., Wahl, D.R., Rasca di Magliano, M., Lyssiotis, C.A., Macleod, K.F., Weber, C.R., Muir, A. (2023). Pancreatic tumors exhibit myeloid-driven amino acid stress and upregulate arginine biosynthesis. eLife *12*. https://doi.org/10.7554/eLife.81289.

Baksh, S.C., Todorova, P.K., Gur-Cohen, S., Hurwitz, B., Ge, Y., Novak, J.S.S., Tierney, M.T., dela Cruz-Racelis, J., Fuchs, E., Finley, L.W.S. (2020). Extracellular serine controls epidermal stem cell fate and tumour initiation. Nature Cell Biology *22*, 779-790. https://doi.org/10.1038/s41556-020-0525-9.

Banh, R.S., Biancur, D.E., Yamamoto, K., Sohn, A.S.W., Walters, B., Kuljanin, M., Gikandi, A., Wang, H., Mancias, J.D., Schneider, R.J., Pacold, M.E., Kimmelman, A.C. (2020). Neurons Release Serine to Support mRNA Translation in Pancreatic Cancer. Cell *183*, 1202-1218. https://doi.org/10.1016/j.cell.2020.10.016.

Bao, X.R., Ong, S.-E., Goldberger, O., Peng, J., Sharma, R., Thompson, D. A., Vafai, S. B., Cox, A. G., Marutani, E., Ichinose, F., Goessling, W., Regev, A., Carr, S. A., Clish, C. B., & Mootha, V. K. (2016). Mitochondrial dysfunction remodels one-carbon metabolism in human cells. eLife *5*. https://doi.org/10.7554/elife.10575.

Bertholet, A.M., Chouchani, E. T., Kazak, L., Angelin, A., Fedorenko, A., Long, J. Z., Vidoni, S., Garrity, R., Cho, J., Terada, N., Wallace, D. C., Spiegelman, B. M., & Kirichok, Y. (2019). H+ transport is an integral function of the mitochondrial ADP/ATP carrier. Nature *571*, 515–520. https://doi.org/10.1038/s41586-019-1400-3.

Birsoy, K., Wang, T., Chen, W.W., Freinkman, E., Abu-Remaileh, M., Sabatini, D.M. (2015). An
Essential Role of the Mitochondrial Electron Transport Chain in Cell Proliferation Is to Enable
Aspartate Synthesis. Cell *162*, 540-551. https://doi.org/10.1016/j.cell.2015.07.016.

Broeks, M.H., Meijer, N.W.F., Westland, D., Bosma, M., Gerrits, J., German, H.M., Ciapaite, J.,
van Karnebeek, C.D.M., Wanders, R.J.A., Zwartkruis, F.J.T., Verhoeven-Duif, N., Jans, J.J.M.
(2023). The malate-aspartate shuttle is important for de novo serine biosynthesis. Cell Reports
42. https://doi.org/10.1016/j.celrep.2023.113043.

Buttgereit, F., Brand, M.D. (1995). A hierarchy of ATP-consuming processes in mammalian
cells. Biochemical Journal *312*, 163-167. https://doi.org/10.1042/bj3120163.

Chaneton, B., Hillman, P., Zheng, L., Martin, A.C.L., Maddocks, O.D.K., Chokkathukalam, A.,
Coyle, J.E., Jankevics, A., Holding, F.P., Vousden, K.H., Frezza, C., O'Reilly, M., Gottlieb, E.
(2012). Serine is a natural ligand and allosteric activator of pyruvate kinase M2. Nature *491*,
458-462. https://doi.org/10.1038/nature11540.

Christensen, C.E., Karlsson, M., Winther, J.R., Jensen, P.R., Lerche, M.H. (2014). Non-invasive
In-cell Determination of Free Cytosolic [NAD+]/[NADH] Ratios Using Hyperpolarized Glucose
Show Large Variations in Metabolic Phenotypes. Journal of Biological Chemistry *289*, 2344-
2352. https://doi.org/10.1074/jbc.M113.498626.

Cognet, G., Muir, A. (2024). Identifying metabolic limitations in the tumor microenvironment.
Science Advances *10*. https://doi.org/10.1126/sciadv.adq7305.

Deberardinis, R.J., Lum, J.J., Hatzivassiliou, G., Thompson, C.B. (2008). The Biology of
Cancer: Metabolic Reprogramming Fuels Cell Growth and Proliferation. Cell Metabolism *7*, 11-
20. https://doi.org/10.1016/j.cmet.2007.10.002.

Delage, B., Fennell, D.A., Nicholson, L., McNeish, I., Lemoine, N.R., Crook, T., Szlosarek, P.W.
(2010). Arginine deprivation and argininosuccinate synthetase expression in the treatment of
cancer. International Journal of Cancer *126*, 2762-2772. https://doi.org/10.1002/ijc.25202.

DeNicola, G.M., Chen P., Mullarky E., Sudderth, J.A., Hu, Z., Wu, D., Tang, H., Xie, Y., Asara
J.M., Huffman, K.E., Wistuba, I.I., Minna, J.D., DeBerardinis, R.J., Cantley, L.C. (2015). NFR2
regulates serine biosynthesis in non-small cell lung cancer. Nature Genetics *47*, 1475-1481.
https://doi.org/10.1038/ng.3421.

Diehl, F.F., Lewis, C.A., Fiske, B.P., Vander Heiden, M.G. (2019). Cellular redox state
constrains serine synthesis and nucleotide production to impact cell proliferation. Nature
Metabolism *1*, 861-867. https://doi.org/10.1038/s42255-019-0108-x.

Fantin, V.R., St-Pierre, J., Leder, P. (2006). Attenuation of LDH-A expression uncovers a link
between glycolysis, mitochondrial physiology, and tumor maintenance. Cancer Cell *9*, 425-434.
https://doi.org/10.1016/j.ccr.2006.04.023.

Faubert, B., Solmonson, A., Deberardinis, R.J. (2020). Metabolic reprogramming and cancer
progression. Science *368*. https://doi.org/10.1126/science.aaw5473.

Fendt, S., Frezza, C., Erez, A. (2020). Targeting Metabolic Plasticity and Flexibility Dynamics for
Cancer Therapy. Cancer Discovery *10*, 1797-1807. https://doi.org/10.1158/2159-8290.CD-20-
0844.
818
Ferraro, G.B., Ali, A., Luengo, A., Kodack, D.P., Deik, A., Abbott, K.L., Bezwada, D., Blanc, L.,
Prideaux, B., Jin, X., Possada, J.M., Chen, J., Chin, C.R., Amoozgar, Z., Ferreira, R., Chen, I.,
Naxerova, K., Ng, C., Westermark, A.M., Duquette, M., Roberge, S., Lindeman, N.I., Lyssiotis,
C.A., Nielsen, J., Housman, D.E., Duda, D.G., Brachtel, E., Golub, T.R., Cantley, L.C., Asara,
J.M., Davidson, S.M., Fukumua, D., Dartois, V.A., Clish, C.B., Jain, R.K., Vander Heiden, M.G.
(2021). Fatty acid synthesis is required for breast cancer brain metastasis. Nature Cancer *2*,
414-428. https://doi.org/10.1038/s43018-021-00183-y.
826
827 Fisher-Wellman, K.H., Neufer, P.D. (2012). Linking mitochondrial bioenergetics to insulin
828 resistance via redox biology. Trends in Endocrinology & Metabolism *23*, 142-153.
829 https://doi.org/10.1016/j.tem.2011.12.008.
830
831 Gao, X., Lee, K., Reid, M.A., Sanderson, S.M., Qiu, C., Li, S., Liu, J., Locasale, J.W. (2018).
832 Serine Availability Influences Mitochondrial Dynamics and Function through Lipid Metabolism.
833 Cell Reports *22*, 3507-3520. https://doi.org/10.1016/j.celrep.2018.03.017.
834
835 Garcia-Bermudez, J., Baudrier, L., La, K., Ge Zhu, X., Fidelin, J., Sviderskiy, V.O.,
836 Papagiannakopoulos, T., Molina, H., Snuderl, M., Lewis, C.A., Possemato, R.L., Birsoy, K.
837 (2018). Aspartate is a limiting metabolite for cancer cell proliferation under hypoxia and in
838 tumours. Nature Cell Biology *20*, 775-781. https://doi.org/10.1038/s41556-018-0118-z.
839
840 Gibson, G., Blass, J.P. (1976). A Relation between [NAD+]/[NADH] Potentials and Glucose
841 Utilization in Rat Brain Slices. Journal of Biological Chemistry *251* (*13*), 4127-4130.
842
843 Gravel, S., Hulea, L., Toban, N., Birman, E., Blouin, M., Zakikhani, M., Zhao, Y., Topisirovic, I.,
844 St-Pierre, J., Pollak, M. (2014). Serine Deprivation Enhances Antineoplastic Activity of
845 Biguanides. Cancer Research *74*, 7521-7533. https://doi.org/10.1158/0008-5472.CAN-14-2643-
846 T.
847
848 Gullino, P.M., Clark, S.H., Grantham, F.H. (1964). The Interstitial Fluid of Solid Tumors. Cancer
849 Research *24*, 780-797.
850
851 Hanahan, D., Weinberg, R.A. (2011). Hallmarks of Cancer: The Next Generation. Cell *144*,
852 646–674. https://doi.org/10.1016/j.cell.2011.02.013.
853
854 Handy, D.E., Loscalzo, J. (2012). Redox Regulation of Mitochondrial Function. Antioxidants &
855 Redox Signaling *16*, 1323-1367. https://doi.org/10.1089/ars.2011.4123.
856
857 Holness, M.J., Sugden, M.C. (2003). Regulation of pyruvate dehydrogenase complex activity by
858 reversible phosphorylation. Biochemical Society Transactions *31* (*6*), 1143-1151.
859 https://doi.org/10.1042/bst0311143.
860
861 Hosios, A.M., Hecht, V.C., Danai, L.V., Johnson, M.O., Rathmell, J.C., Steinhauser, M.L.,
862 Manalis, S.R., Vander Heiden, M.G. (2016). Amino acids rather than glucose account for the
863 majority of cell mass in proliferating mammalian cells. Developmental Cell *36*, 540-549.
864 https://doi.org/10.1016/j.devcel.2016.02.012.
865

866 Hosios, A.M., Li, Z., Lien, E.C., Vander Heiden, M.G. (2018). Preparation of Lipid-Stripped
867 Serum for the Study of Lipid Metabolism in Cell Culture. Bio-protocol *8*.
868 https://doi.org/10.21769/BioProtoc.2876
869
870 Hosios, A.M., and Vander Heiden, M.G. (2018). The redox requirement of proliferation
871 mammalian cells. Journal of Biological Chemistry *293*, 7490-7498.
872 https://doi.org/10.1074/jbc.TM117.000239.
873
874 Krall, A.S., Mullen, P.J., Surjano, F., Momcilovic, M., Schmid, E.W., Halbrook, C.J., Thambundit,
875 A., Mittelman, S.D., Lyssiotis, C.A., Shackelford, D.B., Knott, S.R.V., Christofk, H.R. (2021).
876 Asparagine couples mitochondrial respiration to ATF4 activity and tumor growth. Cell
877 Metabolism *33*, 1013-1026. https://doi.org/10.1016/j.cmet.2021.02.001.
878
879 Labuschagne, C.F., van den Broek, N.J.F., Mackay, G.M., Vousden, K.H., Maddocks, O.D.K.
880 (2014). Serine, but not glycine, supports one-carbon metabolism and proliferation of cancer
881 cells. Cell Reports *7*. https://doi.org/10.1016/j.celrep.2014.04.045.
882
Lapuente-Brun, E., Moreno-Loshuertos, R., Acin-Perez, R., Latorre-Pellicer, A., Colas, C.,
Balsa, E., Perales-Clemente, E., Quiros, P.M., Calvo, E., Rodriguez-Hernandez, M.A., Navas,
P., Cruz, R., Carracedo, A., Lopez-Otin, C., Perez-Martos, A., Fernandez-Silva, P., Fernandez-
Vizarra, E., Enriquez, J.A. (2013). Supercomplex Assembly Determines Electron Flux in the
Mitochondrial Electron Transport Chain. Science *340*, 1567-1570.
https://doi.org/10.1126/science.1230381.
889
890 Li, Z., Ji, B.W., Dixit, P.D., Lien, E.C., Tchourine, K., Hosios, A.M., Abbott, K.L., Westermark,
891 A.M., Gorodetsky, E.F., Sullivan, L.B., Vander Heiden, M.G., Vitkup, D. (2022). Cancer cells
892 depend on environmental lipids for proliferation when electron acceptors are limited. Nature
893 Metabolism *4*, 711-723. https://doi.org/10.1038/s42255-022-00588-8.
894
Locasale, J.W., Grassian, A.R., Melman, T., Lyssiotis, C.A., Mattaini, K.R., Bass, A.J., Heffron,
G., Metallo, C.M., Muranen, T., Sharfi, H., Sasaki, A.T., Anastasiou, D., Mullarky, E., Vokes,
N.I., Sasaki, M., Beroukhim, R., Stephanopoulos, G., Ligon, A.H., Meyerson, M., Richardson,
A.L., Chin, L., Wagner, G., Asara, J.M., Brugge, J.S., Cantley, L.C., Vander Heiden, M.G.
(2011). Phosphoglycerate dehydrogenase diverts glycolytic flux and contributes to oncogenesis.
Nature Genetics *43*, 869-874. https://doi.org/10.1038/ng.890.
901
902 Lopes, E.C., Shi, F., Sawant, A., Ibrahim, M., Gomez-Jenkins, M., Hu, Z., Manchiraju, P., Bhatt,
903 V., Wang, W., Hinrichs, C.S., Wallace, D.C., Su, X., Rabinowitz, J.D., Chan, C.S., Guo, J.Y.,
904 Ganesan, S., Lattime, E.C., White, E. (2025). Respiration defects limit serine synthesis required
905 for lung cancer growth and survival. Nature Communications *16*.
906 https://doi.org/10.1038/s41467-025-62911-7.
907
908 Luengo, A., Li, Z., Gui, D. Y., Sullivan, L. B., Zagorulya, M., Do, B. T., Ferreira, R., Naamati, A.,
909 Ali, A., Lewis, C. A., Thomas, C. J., Spranger, S., Matheson, N. J., & Vander Heiden, M. G.
910 (2021). Increased demand for NAD+ relative to ATP drives aerobic glycolysis. Molecular Cell
911 *81*, 691–707. https://doi.org/10.1016/j.molcel.2020.12.012.
912
913 Lyssiotis, C.A., Kimmelman, A.C. (2017). Metabolic Interactions in the Tumor Microenvironment.
914 Trends in Cell Biology *27*, 863-875. https://doi.org/10.1016/j.tcb.2017.06.003.
915

Maddocks, O.D.K., Athineos, D., Cheung, E. C., Lee, P., Zhang, T., Van Den Broek, N. J. F., Mackay, G. M., Labuschagne, C. F., Gay, D., Kruiswijk, F., Blagih, J., Vincent, D. F., Campbell, K. J., Ceteci, F., Sansom, O. J., Blyth, K., & Vousden, K. H. (2017). Modulating the therapeutic response of tumours to dietary serine and glycine starvation. Nature *544*, 372–376. https://doi.org/10.1038/nature22056.

Maddocks, O.D.K., Berkers, C. R., Mason, S. M., Zheng, L., Blyth, K., Gottlieb, E., & Vousden, K. H. (2013). Serine starvation induces stress and p53-dependent metabolic remodelling in cancer cells. Nature *493* (*7433*), 542–546. https://doi.org/10.1038/nature11743.

McLaughlin, S.G., Dilger, J.P. (1980). Transport of protons across membranes by weak acids. Physiological Reviews *60* (*3*), 825-863. https://doi.org/10.1152/physrev.1980.60.3.825.

Murphy, J.P., Giacomantonio, M. A., Paulo, J. A., Everley, R. A., Kennedy, B. E., Pathak, G. P., Clements, D. R., Kim, Y., Dai, C., Sharif, T., Gygi, S. P., & Gujar, S. (2018). The NAD+ Salvage Pathway Supports PHGDH-Driven Serine Biosynthesis. Cell Reports *24*, 2381–2391.e2385. https://doi.org/10.1016/j.celrep.2018.07.086.

Newman, A.C., Maddocks, O.D.K. (2017). One-carbon metabolism in cancer. British Journal of Cancer *116*, 1499-1504. https://doi.org/10.1038/bjc.2017.118.

Ngo, B., Kim, E., Osorio-Vasquez, V., Doll, S., Bustraan, S., Liang, R. J., Luengo, A., Davidson, S. M., Ali, A., Ferraro, G. B., Fischer, G. M., Eskandari, R., Kang, D. S., Ni, J., Plasger, A., Rajasekhar, V. K., Kastenhuber, E. R., Bacha, S., Sriram, R. K., … Pacold, M. E. (2020). Limited Environmental Serine and Glycine Confer Brain Metastasis Sensitivity to PHGDH Inhibition. Cancer Discovery *10*, 1352–1373. https://doi.org/10.1158/2159-8290.cd-19-1228.

Nobes, C.D., Lakin-Thomas, P.L., Brand, M.D. (1989). The contribution of ATP turnover by the Na+/K+-ATPase to the rate of respiration of hepatocytes: Effects of thyroid status and fatty acids. Biochimica Et Biophysica Acta (BBA) - Bioenergetics *976*, 241-245. https://doi.org/10.1016/S0005-2728(89)80236-1.

Novelli, G.D., Lipmann, F. (1950). The Catalytic Function of Coenzyme A in Citric Acid Synthesis. Journal of Biological Chemistry *182*, 213-228. https://doi.org/10.1016/S0021-9258(18)56541-8.

Patel, M.S., Korotchkina, L.G. (2006). Regulation of the pyruvate dehydrogenase complex. Biochemical Society Transactions *34* (*2*), 217-222. https://doi.org/10.1042/BST20060217.

Pavlova, N.N., Thompson, C.B. (2017). The Emerging Hallmarks of Cancer Metabolism. Cell Metabolism *23*, 27-47. https://doi.org/10.1016/j.cmet.2015.12.006.

Possemato, R., Marks, K.M., Shaul, Y.D., Pacold, M.E., Kim, D., Birsoy, K., Sethumadhavan, S., Woo, H., Jang, H.G., Jha, A.K., Chen, W.W., Barrett, F.G., Stransky, N., Tsun, Z., Cowley, G.S., Barretina, J., Kalaany, N.Y., Hsu, P.P., Ottina, K., Chan, A.M., Yuan, B., Garrayway, L.A., Root, D.E., Mino-Kenudson, M., Brachtel, E.F., Driggers, E.M., Sabatini, D.M. (2011). Functional genomics reveals serine synthesis is essential in PHGDH-amplified breast cancer. Nature *476*, 346-350. https://doi.org/10.1038/nature10350.

Qiu, F., Chen, Y.-R., Liu, X., Chu, C.-Y., Shen, L.-J., Xu, J., Gaur, S., Forman, H. J., Zhang, H., Zheng, S., Yen, Y., Huang, J., Kung, H.-J., & Ann, D. K. (2014). Arginine Starvation Impairs

Mitochondrial Respiratory Function in ASS1-Deficient Breast Cancer Cells. Science Signaling *7*, ra31. https://doi.org/10.1126/scisignal.2004761.

Rabinovich, S., Adler, L., Yizhak, K., Sarver, A., Silberman, A., Agron, S., Stettner, N., Sun, Q., Brandis, A., Helbling, D., Korman, S., Itzkovitz, S., Dimmock, D., Ulitsky, I., Nagamani, S. C. S., Ruppin, E., & Erez, A. (2015). Diversion of aspartate in ASS1-deficient tumours fosters de novo pyrimidine synthesis. Nature *527*, 379-383. https://doi.org/10.1038/nature15529.

Sciacovelli, M., Dugourd, A., Valcarcel Jimenez, L., Yang, M., Nikitopoulou, E., Costa, A.S.H., Tronci, L., Caraffini, V., Rodrigues, P., Schmidt, C., Gerard Ryan, D., Young, T., Zecchini, V.R., Rossi, S.H., Massie, C., Lohoff, C., Masid, M., Hatzimanikatis, V., Kuppe, C., Kriegsheim, A.V., Kramann, R., Gnanapragasam, V., Warren, A.Y., Stewart, G.D., Erez, A., Vanharanta, S., Saez-Rodriguez, J., Frezza, C. (2022). Dynamic partitioning of branched-chain amino acids-derived nitrogen supports renal cancer progression. Nature Communications *13* (*7830*). https://doi.org/10.1038/s41467-022-35036-4.

Sivanand, S., Gultekin, Y., Winter, P.S., Vermeulen, S.Y., Tchourine, K.M., Abbott, K.L., Danai, L.V., Gourge, F., Do, B.T., Crowder, K., Kunchok, T., Lau, A.N., Darnell, A.M., Jefferson, A., Morita, S., Duda, D.G., Aguirre, A.J., Wolpin, B.M., Henning, N., Spanoudaki, V., Maiorino, L., Irvine, D.J., Yilmaz, O.H., Lewis, C.A., Vitkup, D., Shalek, A.K., Vander Heiden, M.G. (2024). Cancer tissue of origin constrains the growth and metabolism of metastases. Nature Metabolism *6*, 1668-1681. https://doi.org/10.1038/s42255-024-01105-9.

Srere, P.A. (1974). Controls of citrate synthase activity. Life Sciences *15*, 1695-1710. https://doi.org/10.1016/0024-3205(74)90172-6.

Sullivan, L.B., Gui, D.Y., Hosios, A.M., Bush, L.N., Freinkman, E., Vander Heiden M.G. (2015). Supporting Aspartate Biosynthesis Is an Essential Function of Respiration in Proliferation Cells. Cell *162*, 552-563. https://doi.org/10.1016/j.cell.2015.07.017.

Sullivan, M.R., Danai, L. V., Lewis, C. A., Chan, S. H., Gui, D. Y., Kunchok, T., Dennstedt, E. A., Vander Heiden, M. G., Muir, A. (2019a). Quantification of microenvironmental metabolites in murine cancers reveals determinants of tumor nutrient availability. eLife *8*. https://doi.org/10.7554/elife.44235.

Sullivan, M.R., Mattaini, K.R., Dennstedt, E.A., Nguyen, A.A., Reilly, M.F., Meeth, K., Muir, A., Darnell, A.M., Bosenberg, M.W., Lewis, C.A., Vander Heiden, M.G. (2019b). Increased serine synthesis provides an advantage for tumors arising in tissues where serine levels are limiting. Cell Metabolism *29*, 1410-1421,e1414. https://doi.org/10.1016/j.cmet.2019.02.015.

Terada, H. (1990). Uncouplers of Oxidative Phosphorylation. Environmental Health Perspectives *87*, 213-218. https://doi.org/10.1289/ehp.9087213.

Titov, D.V., Cracan, V., Goodman, R. P., Peng, J., Grabarek, Z., & Mootha, V. K. (2016). Complementation of mitochondrial electron transport chain by manipulation of the NAD+/NADH ratio. Science *352*, 231-235. https://doi.org/10.1126/science.aad4017.

Vander Heiden, M.G., Chandel, N.S., Schumacker, P.T., Thompson, C.B. (1999). Bcl-xL Prevents Cell Death following Growth Factor Withdrawal by Facilitating Mitochondrial ATP/ADP Exchange. Molecular Cell *3*, 159-167. https://doi.org/10.1016/S1097-2765(00)80307-X.

Vander Heiden, M.G., Deberardinis R.J. (2017). Understanding the Intersections between
Metabolism and Cancer Biology. Cell *168*, 657-669. https://doi.org/10.1016/j.cell.2016.12.039.

Vichai, V., Kirtikara, K. (2006). Sulforhodamine B colorimetric assay for cytotoxicity screening.
Nature Protocols *1*, 1112-1116. https://doi.org/10.1038/nprot.2006.179.

Wang, Y., Stancliffe, E., Fowle-Grider, R., Wang, R., Wang, C., Schwaiger-Haber, M., Shriver,
L.P., Patti, G.J. (2022). Saturation of the mitochondrial NADH shuttles drives aerobic glycolysis
in proliferating cells. Molecular Cell *82*, 3270-3283. https://doi.org/10.1016/j.molcel.2022.07.007.

Wu, Z., Bezwada, D., Cai, F., Harris, R.C., Ko, B., Sondhi, V., Pan, C., Vu, H.S., Nguyen, P.T.,
Faubert, B., Cai, L., Chen, H., Martin-Sandoval, M., Do, D., Gu, W., Zhang, Y., Zhang, Y.,
Brooks, B., Kelekar, S., Zacharias, L.G., Oaxaca, K.C., Patricio, J.S., Mathews, T.P., Garcia-
Bermudez, J., Ni, M., DeBerardinis, R.J., (2024). Electron transport chain inhibition increases
cellular dependence on purine transport and salvage. Cell Metabolism *36*, 1504-1520.
https://doi.org/10.1016/j.cmet.2024.05.014.

Xiao, Y., Yu, T., Xu, Y., Ding, R., Wang, Y., Jiang, Y., Shao, Z. (2023). Emerging therapies in
cancer metabolism. Cell Metabolism *35*, 1283-1303. https://doi.org/10.1016/j.cmet.2023.07.006.

Yao, C., Wang, R., Wang, Y., Kung, C., Weber, J.D., Patti, G.J. (2019). Mitochondrial fusion
supports increased oxidative phosphorylation during cell proliferation. eLife *8*.
https://doi.org/10.7554/eLife.41351.

Zhao, R., Jiang, S., Zhang, L., Yu, Z. (2019). Mitochondrial electron transport chain, ROS
generation, and uncoupling International Journal of Molecular Medicine *44*, 3-15.
https://doi.org/10.3892/ijmm.2019.4188.

Figure Legends

Figure 1. The NAD+/NADH ratio is proportional to serine synthesis rate. (A) α-ketobutyrate

(AKB) is an exogenous electron acceptor that promotes the oxidation of NADH into NAD+ through

its conversion to α-hydroxybutyrate (AHB). Rotenone inhibits activity of complex I of the

mitochondrial electron transport chain (ETC) to block NADH oxidation into NAD+. **(B)**

NAD+/NADH ratio measured in A549 cells cultured without serine and indicated concentrations

of AKB (left) and rotenone (right) for 24 hours. NAD+/NADH ratios are normalized to untreated

cells cultured without serine, n=3. Pearson correlation coefficients and P-values were calculated

by simple linear regression, ***p<0.005, ****p<0.001. **(C)** Schematic depicting NAD+ requirement

to produce serine labeled (M+3) from U-^{13}C-glucose. Under culture conditions without serine,

intracellular serine rapidly effluxes from cells into culture media. Thus, cells and media were jointly

analyzed at each time point of kinetic tracing to better capture levels of newly synthesized serine.

(D) Serine labeled (M+3) from U-[13]C-glucose over time in A549 cells cultured without serine and

with indicated concentrations of AKB or rotenone for 24 hours prior to U-[13]C-glucose exposure.

Serine levels are normalized to internal norvaline standard and cell number measured in indicated

conditions, n=3. **(E)** Relative NAD+/NADH ratios from A549 cells cultured without serine after

exposure to 250 µM AKB, 1000 µM AKB, 20 nM rotenone, or 40 nM rotenone for 24 hours are

plotted against the serine synthesis rates from corresponding conditions as shown in (D). Serine

synthesis rates are calculated from labeled serine (M+3) after 1 and 15 minutes of U-[13]C-glucose

exposure and normalized to untreated culture conditions without serine, n=3. Pearson correlation

coefficients and P-values were calculated by simple linear regression, ****p<0.001. Data shown

for all panels represent mean ± SD.

1068

Figure 2. The NAD+/NADH ratio differs between cancer cells upon serine withdrawal and

correlates with proliferation in serine depleted conditions. (A) Proliferation rate (doublings

per day) of indicated cancer cells cultured without serine for 72 hours normalized to proliferation

rate of corresponding cancer cells cultured with serine, n=3. **(B)** NAD+/NADH ratios of indicated

cells cultured with or without serine for 24 hours. Yellow indicates cells with unaltered

NAD+/NADH ratios (redox non-responder cells) and blue indicates cells with elevated

NAD+/NADH ratios when cultured without serine (redox responder cells). Values are normalized

to the NAD+/NADH ratios of cells cultured with serine, n=3. P-values were calculated by unpaired

Student's t-test, **p<0.01, ***p<0.005. **(C)** Relative NAD+/NADH ratios of cells cultured without

serine are plotted against proliferation rates of corresponding cells cultured without serine. Data

points are labeled to denote the different cell lines (Calu6, A549, 8988T, MIA PaCa-2, H1299,

and HCT116). Pearson correlation coefficient and P-values were calculated by simple linear

regression, *p<0.05. **(D)** Serine labeled (M+3) from U-[13]C-glucose over time in A549 and H1299

cells cultured with or without serine for 24 hours prior to U-[13]C-glucose exposure. Serine levels

were normalized to internal norvaline standard and cell number measured in indicated conditions,

n=3. **(E)** Absolute fold change of serine labeled (M+3) from U-[13]C-glucose in A549 and H1299

cells cultured without serine relative to serine labeled (M+3) from U-[13]C-glucose in A549 and

H1299 cells cultured with serine for 24 hours after 12 minutes of U-[13]C-glucose exposure, n=3.

P-values were calculated by unpaired Student's t-test, **p<0.01. Data shown for all panels are

means ± SD.

1089

Figure 3. Elevated mitochondrial respiration is associated with increased NAD+/NADH

ratios and serine synthesis rates. (A) Oxygen consumption rate (OCR) of indicated cells

cultured with or without serine for 24 hours. Values are averages of three repeat measurements,

n=14-20. P-values were calculated by unpaired Student's t-test, *p<0.05, ****p<0.001. **(B)** OCR

of cells (H1299 top, A549 bottom) cultured with indicated concentrations of serine for 24 hours.

Values are averages of three repeat measurements, n=3-9. P-values were calculated by unpaired

Student's t-test, ****p<0.001. **(C)** NAD+/NADH ratios of cells (H1299 top, A549 bottom) cultured

with indicated concentrations of serine for 24 hours. Values are normalized to the NAD+/NADH

ratio of cells cultured in 400 μM serine (serine concentration in DMEM), n=3. P-values were

calculated by unpaired Student's t-test, **p<0.01, ***p<0.005. **(D)** Relative OCR plotted against

relative NAD+/NADH ratios in H1299 and A549 cells cultured in 0, 25, 50, and 100 μM serine for

24 hours. OCR and NAD+/NADH ratios are normalized to measurements from the 100 μM serine

condition, n=3. Pearson correlation coefficients and P-values were calculated by simple linear

regression, *p<0.05, ****p<0.001. **(E)** Serine labeled (M+3) from U-[13]C-glucose over time in cells

(H1299 top, A549 bottom) cultured with indicated serine concentrations for 24 hours prior to U-

[13]C-glucose exposure. Serine levels are normalized to internal norvaline standard and cell number

measured in indicated conditions, n=3. **(F)** Relative NAD+/NADH ratios of cells (H1299 top, A549

bottom) cultured with 0, 25, 50, and 100 μM serine for 24 hours plotted against relative serine

synthesis rates of cells cultured in corresponding conditions. Serine synthesis rates are calculated

from labeled serine (M+3) after 1 and 15 minutes of U-^{13}C-glucose exposure and normalized to

untreated culture conditions without serine, n=3. Pearson correlation coefficients and P-values

were calculated by simple linear regression, ***p<0.005. Data shown for all panels are means ±

SD.

1113

Figure 4. Mitochondrial respiration governs the endogenous cell NAD+/NADH ratio and

influences serine synthesis. (A) Schematic depicting the mitochondrial electron transport chain

(ETC) and the protonophore activity of FCCP that uncouples the ETC from ATP synthase activity.

CoQ(H$_2$) indicates the oxidized and reduced (H$_2$) forms of coenzyme Q, cyt$_c$ denotes cytochrome

c. **(B,C)** Proliferation rate (doublings per day) of A549 (B) and H1299 cells (C) cultured with or

without serine with indicated concentrations of FCCP for 72 hours, n=3. Increasing doses of FCCP

statistically increased proliferation of A549 cells cultured without serine (p<0.001). P-values were

calculated by simple linear regression. **(D)** Serine labeled (M+3) from U-^{13}C-glucose over time in

A549 cells without serine and with or without 2 μM FCCP treatment for 24 hours prior to U-^{13}C-

glucose exposure. Serine levels are normalized to internal norvaline standard and cell number

measured in indicated conditions, n=3. **(E)** Serine labeled (M+3) from U-^{13}C-glucose over time in

H1299 cells without serine, and with or without 2 μM FCCP treatment for 24 hours prior to U-^{13}C-

glucose exposure. Serine levels are normalized to internal norvaline standard and cell number

measured in indicated conditions, n=3. **(F)** Oxygen consumption rate (OCR) of A549 cells cultured

with or without serine with indicated FCCP treatment for 24 hours. Values are averages of three

repeat measurements, n=5-9. P-values were calculated by unpaired Student's t-test, ****p<0.001.

(G) Relative NAD+/NADH ratio of A549 cells cultured with or without serine with indicated FCCP

treatment for 24 hours. NAD+/NADH ratios are normalized to the NAD+/NADH ratio in A549 cells

cultured with serine, n=3. P-values were calculated by unpaired Student's t-test, *p<0.05,

***p<0.005. **(H)** Proliferation rate (doublings per day) of A549 cells cultured with or without serine,

2 μM FCCP, and indicated concentrations of rotenone for 72 hours, n=3. P-values were calculated

by unpaired Student's t-test, *p<0.05. **(I)** Serine labeled (M+3) from U-^{13}C-glucose over time in

A549 cells cultured without serine and with or without 2 μM FCCP or 20 nM rotenone as indicated

for 24 hours prior to U-^{13}C-glucose exposure. Serine levels are normalized to internal norvaline

standard and cell number measured in indicated conditions, n=3. **(J)** Relative NAD+/NADH ratio

of A549 cells cultured with or without serine, 2 μM FCCP, or 20 nM rotenone as indicated.

NAD+/NADH ratios are normalized to the NAD+/NADH ratio in A549 cells cultured with serine,

n=3. P-values were calculated by unpaired Student's t-test, **p<0.01, ***p<0.005. Data shown for

all panels are means ± SD.

1143

Figure 5. Lipid depletion enhances cell-specific elevation in mitochondrial respiration

and the NAD+/NADH ratio, influencing oxidative citrate and serine synthesis. (A)

Schematic depicting glucose and glutamine oxidative routes for synthesizing citrate, highlighting

NAD+ requiring oxidation reactions. **(B)** Citrate labeled (M+2) from U-^{13}C-glucose over time in

A549 and H1299 cells cultured with or without serine for 24 hours prior to U-^{13}C-glucose

exposure. Citrate levels are normalized to internal norvaline standard and cell number

measured in indicated conditions, n=3. **(C)** Citrate labeled (M+2) from U-^{13}C-glucose over time

in A549 and H1299 cells cultured without serine and with or without 2 μM FCCP for 24 hours

prior to U-^{13}C-glucose exposure. Citrate levels are normalized to internal norvaline standard and

cell number measured in indicated conditions, n=3. **(D)** Citrate labeled (M+2) from U-^{13}C-

glucose over time in A549 cells cultured with serine and with indicated treatment conditions for

24 hours prior to U-^{13}C-glucose exposure. Citrate levels are normalized to internal norvaline

standard and cell number measured in indicated conditions, n=3. **(E)** Oxygen consumption rate

(OCR) in A549 and H1299 cells cultured with or without lipids for 24 hours. Values are averages

of three repeat measurements, n=10. Green indicates lipid redox responder cells. Purple

indicates lipid redox non-responder cells. **(F)** Relative NAD+/NADH ratios of A549 and H1299

cells cultured with or without lipids for 24 hours. NAD+/NADH ratios are normalized to the

NAD+/NADH ratio of the corresponding cell type cultured with lipids, n=3. P-values were

calculated by unpaired Student's t-test, *p<0.05. **(G)** Citrate labeled (M+2) from U-^{13}C-glucose

over time in A549 and H1299 cells cultured with or without lipids for 24 hours prior to U-^{13}C-

glucose exposure. Citrate levels are normalized to internal norvaline standard and cell number

measured in indicated conditions, n=3. **(H)** Serine labeled (M+3) from U-^{13}C-glucose over time

in A549 and H1299 cells cultured with or without lipids for 24 hours prior to U-^{13}C-glucose

exposure. Serine levels are normalized to internal norvaline standard and cell number

measured in indicated conditions, n=3. Data shown for all panels are means ± SD.

1169

Figure 6. Lipid depletion increases mitochondrial respiration and the NAD+/NADH ratio in

a cell-specific manner to influence serine synthesis and proliferation in serine depleted

conditions. (A) Oxygen consumption rate (OCR) of A549 and H1299 cells cultured with or without

serine or lipids for 24 hours. Values are the average of three repeat measurements, n=14-15. P-

values were calculated by one-way ANOVA followed by post-hoc Tukey HSD test, ****p<0.001.

(B) Relative NAD+/NADH ratios of A549 and H1299 cells cultured with or without serine or lipids

for 24 hours. NAD+/NADH ratio is normalized to the NAD+/NADH ratio in corresponding cells

cultured with serine and lipids, n=3. P-values were calculated by one-way ANOVA followed by

post-hoc Tukey HSD test, *p<0.05, ****p<0.001. **(C, D)** Left: Serine labeled (M+3) from U-^{13}C-

glucose over time in A549 cells (C) and H1299 cells (D) cultured without serine and with or without

lipids for 24 hours prior to U-^{13}C-glucose exposure. Right: Serine synthesis rates of A549 cells (C)

and H1299 cells (D) cultured in indicated conditions. Serine synthesis rates are calculated from

labeled serine (M+3) after 1 and 15 minutes of U-^{13}C-glucose exposure and normalized to the

serine synthesis rates of the corresponding cell line cultured without serine and with lipids. Serine

levels are normalized to internal norvaline standard and cell number measured in indicated

conditions, n=3. Single data point (H1299, -serine +lipid, t=15 min) removed due to detection error

with no signal. (**E,F**) Left: Citrate labeled (M+2) from U-^{13}C-glucose over time in A549 cells (E)

and H1299 cells (F) cultured without lipids and with or without serine for 24 hours prior to U-^{13}C-

glucose exposure. Right: Citrate synthesis rates of A549 cells (E) and H1299 cells (F) cultured in

indicated conditions. Citrate synthesis rates are calculated from labeled citrate (M+2) after 1 and

15 minutes of U-^{13}C-glucose exposure and normalized to the citrate synthesis rates of the

corresponding cell line cultured without lipids and with serine. Citrate levels are normalized to

internal norvaline standard and cell number measured in indicated conditions, n=3. **(G)**

Proliferation rate (doublings per day) of A549 cells cultured with or without serine or lipids for 72

hours, n=3. P-values were calculated by one-way ANOVA followed by post-hoc Tukey HSD test,

p<0.01 **(H) Proliferation rate (doublings per day) of H1299 cells cultured with or without serine

or lipids for 72 hours, n=3. Data shown for all panels are means ± SD.

1197

Figure 7. The cell NAD+/NADH ratio is regulated in a cancer cell-specific manner by

changes to mitochondrial respiration in response to nutrient availability, impacting

oxidative biosynthetic reactions. (A) In serine and lipid-replete conditions, cells can acquire

environmental serine and lipids to support proliferation. Cells can also oxidize glucose to produce

serine, as well as citrate for lipid synthesis. The oxidation reactions involved can be limited by

NAD+ availability, which is impacted by mitochondrial respiration. **(B)** In nutrient depleted

conditions, the ability to support oxidative reactions is dependent on the mitochondrial response

to the nutrient depletion. For redox responders, environmental depletion of serine or lipids

increases the cell NAD+/NADH ratio to support both oxidative citrate and serine synthesis. This

ability to increase respiration enables proliferation in conditions lacking serine. **(C)** In redox non-

responder cells, neither mitochondrial respiration nor the NAD+/NADH ratio increases following

nutrient depletion. This limits oxidative synthesis of serine (and citrate), subsequently limiting

proliferation in environments that lack serine.

1211

1212

1213

Supplementary Material

Supplementary Figure 1

Supplementary Figure 1. Altering NAD+/NADH with α-ketobutyrate (AKB) or rotenone does not change PHGDH protein levels and minimally changes cell viability. (A,B) Immunoblots assessing PHGDH protein levels in A549 cells cultured with or without serine, 1 mM AKB (A), or 20 nM rotenone (B), as indicated for 24 hours. HSP90 expression is shown as a loading control. **(C)** Percent of A549 cells dead, measured using dye exclusion 72 hours after exposure to the indicated concentrations of rotenone in serine-replete or serine depleted conditions, n=3. Serine depletion led to a statistically significant increase in cell death compared to serine-replete conditions at all concentrations of rotenone with the exception of 40 nM rotenone, p<0.05. There was no statistical difference in cell death between 0 and 20 nM rotenone in either serine-replete or serine depleted conditions. P-values were calculated using unpaired Student's t-test. Data shown are means ± SD.

Supplementary Figure 2

Supplementary Figure 2. Relationship between PHGDH protein expression, cell NAD+/NADH ratio, and cell proliferation rate following serine deprivation. (A) Proliferation rates (doublings per day) of cells cultured with or without serine for 72 hours, n=3. **(B)** Above: Schematic of the serine synthesis pathway. Below: Immunoblot assessing protein levels of PHGDH, PSAT1, and PSPH in the indicated cells cultured with or without serine for 24 hours.

HSP90 protein expression is shown as a loading control. These data are representative of three independent experiments. Abbreviations – 3-PG: 3-phosphoglycerate, PHGDH: phosphoglycerate dehydrogenase, 3PHP: 3-phosphohydroxypyruvate, glu: glutamate, αKG: α-ketoglutarate, PSAT1: phosphoserine aminotransferase 1, 3PS: 3-phosphoserine, PSPH: phosphoserine phosphatase. **(C)** Quantification of PHGDH protein expression for the indicated cells under serine depleted conditions (normalized to HSP90 protein expression) from three biological replicates, correlated with the corresponding relative proliferation rate of indicated cells under serine depleted conditions based on data presented in (A). Cell lines enclosed in the light blue shaded box indicate cancer cell lines with serine synthesis enzyme expression that could not fully predict proliferation in the absence of serine that we chose to examine further. Pearson correlation coefficient calculated by simple linear regression. **(D)** Cell number over time for either A549 cells (yellow) or H1299 cells (blue) in either serine-replete or serine depleted conditions, n=6. **(E)** Calculated doublings per day from the data in (D) for the indicated cells and conditions, n=6. P-values were calculated using unpaired Student's t-test, ****p<0.001. **(F)** Correlation between the protein expression of serine synthesis enzymes PHGDH, PSAT1, and PSPH across three biological replicates and proliferation rate in serine depleted conditions for the indicated cell lines, n=3. Pearson correlation coefficient and P-values were calculated by simple linear regression, *p<0.05, ***p<0.005. **(G)** Unnormalized NAD+/NADH ratios measured for the indicated cell lines cultured with or without serine for 24 hours, n=3 per each cell line. P-values were calculated by unpaired Student's t-test, *p<0.05, ***p<0.005. **(H)** Fraction of total serine labeled from U-^{13}C-glucose over time in A549 and H1299 cells that were cultured with or without serine for 24 hours prior to U-^{13}C-glucose exposure as indicated, n=3. **(I)** Left: Total intracellular serine levels over time from the kinetic tracing with A549 and H1299 cells cultured with or without serine for 24 hours prior to U-^{13}C-glucose exposure as described in (H). Right: Average of total intracellular serine levels from all time points of the kinetic U-^{13}C-glucose isotope tracing experiment shown in (H). Serine levels were normalized to internal norvaline standard and cell number. P-values were calculated by unpaired Student's t-test, ***p<0.005. Data shown for all panels are means ± SD.

Supplementary Figure 3

Supplementary Figure 3. Lactate dehydrogenase activity, reliance on the NAD+ salvage pathway, mitochondrial respiration, and PHGDH protein levels following serine deprivation. (A) Schematic depicting the contributions of lactate dehydrogenase (LDH) and the NAD+ salvage

pathway to NAD+ generation. Glucose is consumed by cells and converted to pyruvate via glycolysis. LDH can convert pyruvate to lactate by oxidizing NADH to NAD+. In the NAD+ salvage pathway, nicotinamide can be converted to nicotinamide mononucleotide (NMN) by the enzyme nicotinamide phosphoribosyltransferase (NAMPT), which is inhibited by the small molecule FK866. NMN is subsequently converted to NAD+ via NMN adenylyltransferase. **(B)** Lactate secretion rate normalized to glucose consumption rate as measured in the indicated cells cultured with or without serine for 72 hours, n=3. **(C)** Proliferation rate (doublings per day) of A549 (above) and H1299 cells (below) cultured with or without serine and with or without the indicated concentrations of FK866 for 72 hours, n=3. There was a statistically significant difference in the sensitivity of A549 and H1299 cells to FK866 in the serine-replete condition ($p<0.001$), but not in the serine depleted conditions. P-values were calculated by ANCOVA analysis followed by post-hoc Tukey HSD test. **(D)** Proliferation rate (doublings per day) of A549 (left) and H1299 cells (right) cultured with or without serine and either NMN (100 μM) or FK866 (10 μM) as indicated for 72 hours, n=3. P-values were calculated using unpaired Student's t-test, ***$p<0.005$, ****$p<0.001$. **(E)** Oxygen consumption rate (OCR) of cells cultured with or without serine for 24 hours, n=10-20. Rotenone and antimycin A were injected between the third and fourth OCR measurements to define non-mitochondrial OCR. Data shown are normalized to cell number. **(F)** Proliferation rate (doublings per day) of A549 and H1299 cells cultured in media with the indicated concentrations of serine for 72 hours. Media was replenished every day to avoid serine depletion in lower serine-containing conditions, n=3. **(G)** Immunoblot assessing PHGDH protein levels in A549 or H1299 cells cultured in media with the indicated concentrations of serine for 24 hours. HSP90 expression is shown as a loading control. Data shown for all panels are means ± SD.

Supplementary Figure 4

Supplementary Figure 4. FCCP increases both serine synthesis and proliferation of redox non-responder cells in serine depleted conditions. (A) Proliferation rate (doublings per day) of redox non-responder cells Calu6 and 8988T cultured with or without serine and the indicated concentration of FCCP for 72 hours, n=3. Increasing doses of FCCP statistically increased the proliferation of Calu6 cells cultured without serine, p<0.001. P-values were calculated by simple

linear regression. All doses of FCCP statistically increased the proliferation of 8988T cells cultured without serine but not in a dose-dependent manner, $p < 0.001$. P-values were calculated by unpaired Student's t-test. **(B)** Proliferation rate (doublings per day) of redox responder cells MIA PaCa-2 and HCT116 cultured with or without serine and the indicated concentration of FCCP for 72 hours, n=3. **(C)** Mitochondrial oxygen consumption rate (OCR) of H1299 cells cultured with or without serine and the indicated concentration of FCCP for 24 hours, n=5. Values are averages of three repeat measurements. P-values were calculated by unpaired Student's t-test, *$p < 0.05$, **$p < 0.01$, ****$p < 0.001$. **(D)** NAD+/NADH ratio measured in H1299 cells cultured with or without serine and the indicated concentration of FCCP for 24 hours, n=3. P-values were calculated by unpaired Student's t-test, **$p < 0.01$. **(E)** Proliferation rate (doublings per day) of redox responder cells H1299 and HCT116 with or without PHGDH overexpression and cultured with or without serine and 2 µM FCCP, as indicated, for 72 hours, n=3. P-values were calculated by two-way ANOVA followed by post-hoc Tukey HSD test, *$p < 0.05$, ***$p < 0.005$, ****$p < 0.001$. **(F)** Proliferation rate (doublings per day) of redox non-responder cells A549 and Calu6 with or without PHGDH overexpression and cultured with or without serine and 2 µM FCCP for 72 hours, n=3. P-values were calculated by two-way ANOVA followed by post-hoc Tukey HSD test, ***$p < 0.005$, ****$p < 0.001$. **(G)** Immunoblot assessing PHGDH protein levels in the indicated cell lines without (pLJM1-EGFP) or with (pLJM1-PHGDH) exogenous PHGDH expression. Vinculin expression is shown as a loading control. **(H,I)** Proliferation rate (doublings per day) of A549 cells (H) and H1299 cells (I) cultured with or without serine, with or without 2 µM FCCP, or the indicated concentration of oligomycin for 72 hours, n=3. P-values were calculated by unpaired Student's t-test, *$p < 0.05$, ***$p < 0.005$, ****$p < 0.001$. **(J)** Immunoblot assessing PHGDH protein levels in A549 cells cultured with or without serine and with or without 2 µM FCCP for 24 hours. HSP90 expression is shown as a loading control. **(K)** Glucose consumption rates of A549 cells cultured with or without serine and 2 µM FCCP for 72 hours, n=3. Data shown for all panels are means ± SD.

Supplementary Figure 5

Supplementary Figure 5. Kinetic tracing of labeled carbon from U-^{13}C-glucose into serine following serine withdrawal and FCCP treatment. (A) Fraction of total serine labeled (M+3 serine) from U-^{13}C-glucose over time in A549 cells cultured without serine, and with or without 2 µM FCCP, for 24 hours prior to U-^{13}C-glucose exposure as indicated, n=3. **(B)** Total serine measured in A549 cells cultured without serine and with or without 2 µM FCCP for each time point shown in (A), n=3. **(C)** Fraction of total serine labeled (M+3 serine) from U-^{13}C-glucose over time in H1299 cells cultured without serine and with or without 2 µM FCCP for 24 hours prior to U-^{13}C-glucose exposure as indicated, n=3. **(D)** Total serine measured in H1299 cells cultured without serine and with or without 2 µM FCCP for each time point shown in (C), n=3. **(E)** Fraction of total serine labeled (M+3 serine) from U-^{13}C-glucose over time in A549 cells cultured without serine and with or without 2 µM FCCP and 20 nM rotenone for 24 hours prior to U-^{13}C-glucose exposure as indicated, n=3. **(F)** Total serine measured in A549 cells cultured without serine and with or without 2 µM FCCP and 20 nM rotenone for each time point shown in (E), n=3. All serine measurements were normalized to internal norvaline standard and cell number. Data shown for all panels are means ± SD.

Supplementary Figure 6

Supplementary Figure 6. Cytosolic or mitochondrial *Lb*NOX expression does not change the cell NAD+/NADH ratio or proliferation of A549 cells cultured without serine. (A) Immunoblot for expression of FLAG-tagged cyto*Lb*NOX or mito*Lb*NOX as indicated. GFP-expressing cells were used as a control. Vinculin expression is shown as a loading control. **(B)** Relative proliferation rate (doublings per day) of A549 cells expressing either cyto*Lb*NOX or mito*Lb*NOX and treated with 50 nM rotenone for 72 hours, n=3. P-values were calculated using one-way ANOVA followed by post-hoc Tukey HSD test, ***p<0.005, ****p<0.001. **(C)** Relative NAD+/NADH ratios of cells expressing either GFP, cyto*Lb*NOX, or mito*Lb*NOX and cultured with or without serine for 24 hours. NAD+/NADH ratios were normalized to the GFP control cells cultured with serine, n=3. **(D)** Proliferation rate (doublings per day) of cells expressing either GFP, cyto*Lb*NOX, or mito*Lb*NOX and cultured with or without serine for 72 hours, n=3. Data shown for all panels are means ± SD.

Supplementary Figure 7

Supplementary Figure 7. MDH1 and GOT2 support the increase in serine synthesis observed in H1299 cells following serine withdrawal. (A) Immunoblots for GOT1 (left), MDH1 (middle), and GOT2 (right) in control cells (NTC, non-targeting control) or cells with knockout (KO) of GOT1, MDH1, or GOT2 as indicated. Vinculin expression is shown as a loading control. **(B)** Proliferation rate (doublings per day) of the indicated cells from (A) cultured with or without serine for 72 hours, n=3. P-values were calculated by nested ANOVA to compare sensitivity to serine depletion between cell lines, ****p<0.001. **(C)** Relative NAD+/NADH ratios of the indicated cells cultured with or without serine for 24 hours, n=3. NAD+/NADH ratios were normalized to H1299 NTC cells cultured with serine. P-values were calculated using unpaired Student's t-test, *p<0.05. **(D)** Oxygen consumption rate (OCR) of the indicated cells cultured with or without serine for 24 hours, n=6. P-values were calculated by unpaired Student's t-test, **p<0.01. **(E)** Serine labeled from U-^{13}C-glucose (M+3) measured over time from the indicated cells cultured with or without serine for 24 hours prior to U-^{13}C-glucose exposure, n=3. Serine levels were normalized to internal standard norvaline and cell number. Data shown for all panels are means ± SD.

Supplementary Figure 8

Supplementary Figure 8. Kinetic tracing of labeled carbon from U-^{13}C-glucose and U-^{13}C-glutamine into citrate in cells cultured with or without serine. (A) Schematic depicting isotopically-labeled carbon contribution to citrate from U-^{13}C-glucose and U-^{13}C-glutamine. Abbreviations: PDH – pyruvate dehydrogenase, CS – citrate synthase, IDH – isocitrate dehydrogenase, SDH – succinate dehydrogenase. **(B)** Citrate labeled (M+2) from U-^{13}C-glucose

over time in Calu6, 8988T, MIA PaCa-2, and HCT116 cells cultured with or without serine for 24 hours prior to U-^{13}C-glucose exposure, n=3. **(C-H)** Left of each figure panel: Fraction of citrate labeled (M+2) from U-^{13}C-glucose over time in indicated cells cultured with or without serine for 24 hours, n=3. Right of each figure panel: Total citrate levels in the indicated cells cultured with or without serine for 24 hours for each time point shown in left panel, n=3. **(I)** Citrate labeled (M+4) from U-^{13}C-glutamine over time in A549 and H1299 cells cultured with or without serine for 24 hours, n=3. **(J)** Citrate labeled (M+5) from U-^{13}C-glutamine over time in A549 and H1299 cells cultured with or without serine for 24 hours, n=3. All citrate measurements were normalized to internal norvaline standard and cell number measured in indicated conditions. Data shown for all panels are means ± SD.

Supplementary Figure 9

Supplementary Figure 9. Kinetic tracing of labeled carbon from U-^{13}C-glucose into citrate in cells cultured without serine and in the presence or absence of FCCP and/or rotenone. **(A)** Fraction of citrate labeled (M+2) from U-^{13}C-glucose over time in A549 cells cultured without serine for 24 hours, and with or without the addition of 2 µM FCCP or 20 nM rotenone as indicated, n=3. **(B)** Fraction of citrate labeled (M+2) from U-^{13}C-glucose over time in A549 cells cultured without serine for 24 hours, and with or without the addition of 2 µM FCCP or 20 nM rotenone as indicated, n=3. **(C)** Total citrate levels in A549 cells cultured without serine for 24 hours, and with or without the addition of 2 µM FCCP or 20 nM rotenone as indicated, n=3. All citrate measurements were normalized to internal norvaline standard and cell number. Data shown for all panels are means ± SD.

Supplementary Figure 10

Supplementary Figure 10. Kinetic tracing of labeled carbon from U-^{13}C-glucose into citrate in cells cultured in the presence or absence of lipids. (A) Fraction of citrate labeled (M+2) from U-^{13}C-glucose over time in A549 and H1299 cells cultured with or without lipids for 24 hours as indicated, n=3. **(B)** Total citrate levels in A549 and H1299 cells cultured with or without lipids for 24 hours for each time point shown in (A), n=3. **(C)** Fraction of serine labeled (M+3) from U-^{13}C-glucose over time in A549 and H1299 cells cultured with or without lipids for 24 hours as indicated, n=3. **(D)** Total serine levels in A549 and H1299 cells cultured with or without lipids for

24 hours for each time point shown in (C), n=3. All citrate and serine measurements were normalized to internal norvaline standard and cell number. Data shown for all panels are means ± SD.

Supplementary Figure 11

Supplementary Figure 11. Kinetic tracing of labeled carbon from U-^{13}C-glucose into serine and citrate in cells cultured in the absence of serine, lipids, or both serine and lipids. (A) Fraction of serine labeled (M+3) from U-^{13}C-glucose over time in A549 and H1299 cells cultured without serine, and with or without lipids for 24 hours as indicated, n=3. **(B)** Total serine levels in A549 and H1299 cells cultured without serine, and with or without lipids for 24 hours for each time point shown in (A), n=3. **(C)** Fraction of citrate labeled (M+2) from U-^{13}C-glucose over time in A549 and H1299 cells cultured without lipids, and with or without serine for 24 hours as indicated, n=3. **(D)** Total citrate levels in A549 and H1299 cells cultured without lipids, and with or without serine for 24 hours for each time point shown in (C), n=3. All citrate and serine measurements were normalized to internal norvaline standard and cell number. Data shown for all panels are means ± SD.

Supplementary Table 1

Cell Line	Genetic Mutations	Cancer Type	Source
Calu6	Kras Q61K TP53 R196*	Lung Adenocarcinoma	pleural effusion
A549	Kras G12S Stk11 Q37*	Lung Adenocarcinoma	lung
PaTu 8988T	Kras G12V TP53 R282W Smad4 del EP300 del	Pancreatic ductal adenocarcinoma	liver metastasis
MIA PaCa-2	Kras G12C TP53 R248W CDKN2A del	Pancreatic ductal adenocarcinoma	pancreas
H1299	Nras Q61K TP53 del	Lung Adenocarcinoma	lymph node metastasis
HCT116	Kras G13D BRCA2 I2675Dfs CDKN2A mut EP300 mut PIK3CA H1047R PPMID L450* TGFBR2 K128Sfs	Colon Carcinoma	colon
A375	BRAF V600E CDKNA E61* TERTc.1-146C>T	Melanoma	skin
MDA-MB-231	BRAF G464V Kras G13D TP53 R280K CDKN2A/B del TERTc.1-146C>T	Triple Negative Breast Adenocarcinoma	pleural effusion
MCF7	PIK3CA E545K GATA3 D336Gfs CDKN2A del	ER+ HER2- Breast Carcinoma	pleural effusion

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7