1

Cancer cells differentially modulate mitochondrial respiration to alter redox state and

enable biomass synthesis in nutrient-limited environments

Sarah M. Chang[1,2], Muhammad Bin Munim[1,2], Sonia E. Trojan[1,3], Anna Shevzov-Zebrun[1,2],

Keene L. Abbott[1,2], and Matthew G. Vander Heiden[1,2,4*]

[1]Koch Institute for Integrative Cancer Research, Massachusetts Institute of Technology,

Cambridge, MA, USA

[2]Department of Biology, Massachusetts Institute of Technology, Cambridge, MA, USA

[3]Jagiellonian University Medical College, Faculty of Medicine, Chair of Medical Biochemistry,

Krakow, Poland

[4]Dana-Farber Cancer Institute, Boston, MA, USA

*Correspondence: mvh@mit.edu

Abstract

The cell NAD+/NADH ratio can constrain biomass synthesis and influence proliferation in nutrient-limited environments. However, which cell processes regulate the NAD+/NADH ratio is not known. Here, we find that some cancer cells elevate the NAD+/NADH ratio in response to serine deprivation by increasing mitochondrial respiration. Cancer cells that elevate mitochondrial respiration have higher serine production and proliferation in serine limiting conditions than cells with no mitochondrial respiration response, independent of serine synthesis enzyme expression. Increases in mitochondrial respiration and the NAD+/NADH ratio promote serine synthesis regardless of whether serine is environmentally limiting. Lipid deprivation can also increase the NAD+/NADH ratio via mitochondrial respiration in some cells, including cells that do not increase respiration following serine deprivation. Thus, in cancer cells where lipid depletion raises the NAD+/NADH ratio, proliferation in serine depleted environments improves when lipids are also depleted. Taken together, these data suggest that changes in mitochondrial respiration in response to nutrient deprivation can influence the NAD+/NADH ratio in a cell-specific manner to impact oxidative biomass synthesis and proliferation. Given the complexity of tumor microenvironments, this work provides a metabolic framework for understanding how levels of more than one environmental nutrient affects cancer cell proliferation.

Introduction

Rapidly proliferating cells, including cancer cells, must acquire biomass precursors such as amino acids, lipids, and nucleotides to support cell growth and division (Deberardinis, 2008; Faubert, 2020; Hanahan, 2011; Pavlova, 2017; Vander Heiden, 2017). While cells can obtain biomass precursors from the environment, many tissue and tumor environments can be deficient in necessary nutrients (Cognet, 2024; Gullino, 1964; Lyssiotis, 2017). In these conditions, cells depend on biosynthesis to accumulate sufficient biomass for proliferation (Apiz Saab, 2023; Ferraro, 2021; Ngo, 2020; Sciacovelli, 2022; Sullivan, 2019b; Sullivan, 2019a). Synthesis of many

biomass precursors involve oxidation reactions, which require the redox cofactor NAD+ to act as

an electron acceptor. There is accumulating evidence that NAD+ availability and the cell redox

state (NAD+/NADH ratio) can limit the synthesis of biomass precursors, including aspartate

(Birsoy, 2015; Sullivan, 2015), asparagine (Krall, 2021), fatty acids (Li, 2022), serine (Baksh, 2020;

Bao, 2016; Broeks, 2023; Diehl, 2019), and nucleotides (Bao, 2016; Diehl, 2019; Wu, 2024). Thus,

the NAD+/NADH ratio impacts cancer cell proliferation in nutrient environments that increase the

dependence on biosynthetic oxidation reactions. However, what determines the cellular

NAD+/NADH ratio and whether the NAD+/NADH ratio is modulated in response to environmental

nutrient availability is not well characterized. Moreover, whether differences in the NAD+/NADH

ratio across cell types affect sensitivity to different nutrient limitations is not known.

Some tumor microenvironments have low levels of serine, an amino acid used to produce

many biomass components, including proteins, nucleotides, glutathione, and lipids. Thus, low

serine environments can limit cancer cell proliferation and tumor growth (Banh, 2020; Maddocks,

2013; 2017; Ngo, 2020; Sullivan, 2019b). In serine depleted environments, cancer cells must

increase serine synthesis to maintain proliferation (DeNicola, 2015; Labuschagne, 2014;

Locasale, 2011; Possemato, 2011; Sullivan, 2019b). Serine synthesis involves the conversion of

the glycolytic intermediate 3-phosphoglycerate (3-PG) into serine. The production of 3-PG from

glucose involves an oxidation reaction catalyzed by glyceraldehyde 3-phosphate dehydrogenase

(GADPH). 3-PG is then oxidized by phosphoglycerate dehydrogenase (PHGDH) in the first step

of the serine synthesis pathway. Both GADPH and PHGDH require NAD+ as an electron acceptor

to enable serine synthesis. Experimentally suppressing NAD+ regeneration from NADH

decreases serine synthesis, which can be rescued by oxidizing the NAD+/NADH ratio via

exogenous electron acceptor supplementation (Bao, 2016; Diehl, 2019; Gravel, 2014). Similarly,

de novo fatty acid synthesis can be constrained by the NAD+/NADH ratio, as the production of

fatty acids involves multiple oxidation reactions to generate citrate from either glucose or

glutamine. Thus, decreasing the NAD+/NADH ratio sensitizes cells to environmental lipid

depletion (Li, 2022). Together, these findings demonstrate that exogenous changes to the

NAD+/NADH ratio can alter serine and citrate production and impact proliferation in serine or lipid-

depleted environments. However, auxotrophy for a single nutrient is not sufficient to determine

whether cancer cells can form tumors in a tissue, where levels of multiple nutrients may be

environmentally limiting for proliferation (Abbott, 2024; Sivanand, 2024; Sullivan, 2019a).Thus,

we aimed to uncover how the endogenous NAD+/NADH ratio of cancer cells is regulated to

influence biosynthesis in different nutrient depleted environments and its role in determining

cancer cell proliferation when multiple oxidized nutrients are limiting.

By culturing cancer cells in serine depleted conditions and examining the relationship

between the NAD+/NADH ratio and serine synthesis, we find that the NAD+/NADH ratio is

elevated in response to serine deprivation in some but not all cancer cells. The elevated

NAD+/NADH ratio is driven by increased mitochondrial respiration and promotes serine synthesis

to confer resistance to environmental serine depletion. A similar relationship between

mitochondrial respiration and the NAD+/NADH ratio was observed in lipid-depleted environments,

where higher mitochondrial respiration in some cells led to elevated NAD+/NADH ratios and

enhanced citrate synthesis. Lastly, we find that any perturbation that increases the NAD+/NADH

ratio, including either serine or lipid deprivation, led to both elevated serine and citrate production,

regardless of what nutrient was depleted from the environment. Together, these data reveal that

the NAD+/NADH ratio is regulated by mitochondrial respiration in response to environmental

nutrients in a cancer cell-specific manner and that this influences oxidative biosynthesis and

proliferation. These data also provide insight into how levels of multiple nutrients in tissue

microenvironments may cooperate to enable biomass synthesis and cell proliferation. This study

is based in part on work performed for a PhD dissertation (Chang, 2024).

Results

Modulating the cell NAD+/NADH ratio proportionally alters serine synthesis

To begin investigating the processes that determine the cell NAD+/NADH ratio and

influence biomass synthesis, we examined the relationship between the NAD+/NADH ratio and

serine synthesis in serine replete and serine limited conditions. Previous studies have found that

decreasing the cell NAD+/NADH ratio (more reduced) can suppress serine production (Baksh,

2020; Bao, 2016; Broeks, 2023; Diehl, 2019). To confirm this, we varied the cell NAD+/NADH

ratio upon serine withdrawal and measured the serine synthesis rate of A549 non-small cell lung

cancer cells, which transcriptionally upregulate the serine synthesis enzyme PHGDH (DeNicola,

2015). We increased the NAD+/NADH ratio by treating cells with the exogenous electron acceptor

alpha-ketobutyrate (AKB), which can be reduced to alpha-hydroxybutyrate to regenerate NAD+

(Sullivan, 2015) (**Figure 1A**). We decreased the NAD+/NADH ratio by treating cells with rotenone,

an inhibitor of mitochondrial complex I, which regenerates NAD+ in support of respiration. AKB

and rotenone dose-dependently increased and decreased the cell NAD+/NADH ratio, respectively

(**Figure 1B**). To measure the serine synthesis rate in cells, we performed kinetic isotope tracing

using uniformly ^{13}C-labeled glucose (U-^{13}C-glucose) and measured production of M+3 labeled

serine (**Figure 1C**). Because intracellular serine, including newly synthesized serine, rapidly

exchanges with the serine-free extracellular environment (Labuschagne, 2014), we collected both

media and cells to ensure detection of all newly synthesized serine produced from glucose. We

find that increasing the cell NAD+/NADH ratio with AKB led to proportially higher serine

synthesis rates whereas lowering the cellular NAD+/NADH ratio with rotenone led to

proportionally lower serine synthesis rates (**Figure 1D**). This was not explained by changes in

PHGDH protein expression, as altering the NAD+/NADH ratio with AKB or rotenone did not affect

PHGDH protein levels despite modulating serine synthesis (**Supplementary Figure 1A,B**). Of

note, we find that serine synthesis rate is positively correlated with the cell NAD+/NADH ratio

(**Figure 1E**). Accordingly, the NAD+/NADH ratio is also positively correlated with proliferation rate

following serine depletion (**Figure 1F,G; Supplementary Figure 1C**). Together, these data

demonstrate that modulating the NAD+/NADH ratio leads to proportional changes in serine

synthesis rates and proliferation in serine-depleted environments.

Cancer cells with elevated NAD+/NADH ratios following serine withdrawal exhibit greater

serine synthesis

Based on our finding that the NAD+/NADH ratio correlates with serine synthesis rate, we

wondered whether endogenous cell NAD+/NADH ratios, which differ across cancer cells, are

predictive of proliferation rate following serine deprivation. To test this, we measured how

proliferation rate varied in a panel of cancer cells derived from various tissues-of-origin and with

different genetic driver mutations, and found a wide range of sensitivity to serine deprivation

(**Figure 2A**, **Supplementary Figure 2A, Supplementary Table 1**). Because PHGDH expression

is important for proliferation in serine depleted conditions (DeNicola, 2015; Locasale, 2011;

Maddocks, 2017; Possemato, 2011; Sullivan, 2019b), we first assessed PHGDH protein

expression across the different cancer cells to examine how well this explained environmental

serine dependence. As expected, some cells with low PHGDH protein expression (MCF7, MDA-

MB-231) were most sensitive to serine withdrawal, and some cells with higher PHGDH protein

levels were more resistant to serine withdrawal (PHGDH-overexpressing MDA-MB-231, A375)

(**Supplementary Figure 2A-C**). However, for many cancer cells PHGDH expression alone could

not fully explain sensitivity to serine withdrawal (**Supplementary Figure 2D**). For example, when

comparing the non-small cell lung cancer cells A549 and H1299, H1299 cells express less

PHGDH protein than A549 cells but are more resistant to serine deprivation **(Supplementary**

Figure 2B-D). Consistent with lower PHGDH protein expression, H1299 cells express lower

levels of NRF2, a transcriptional regulator of serine biosynthesis enzymes, including PHGDH

(DeNicola, 2015). Importantly, A549 and H1299 cells proliferate similarly in serine replete

conditions (A549: 1.17 doublings per day; H1299: 1.33 doublings per day), arguing that

differences in basal proliferation cannot explain the different sensitivities to serine depletion

(**Supplementary Figure 2A**). Given the finding that the NAD+/NADH ratio correlates with serine

synthesis rate, we hypothesized that differences in the NAD+/NADH ratio upon serine deprivation

could contribute to the variability in sensitivity to serine withdrawal across cells with similar

PHGDH protein levels (Calu6, A549, PaTu8988T, MIA PaCa-2, H1299, and HCT116)

(**Supplementary Figure 2B-D**). To test this, we measured the cell NAD+/NADH ratios in the

presence and absence of exogenous serine. Interestingly, we find that serine starvation elevated

the NAD+/NADH ratio in some cells, while minimally changing the NAD+/NADH ratio in others

(**Figure 2B, Supplementary Figure 2E**). Strikingly, cells with elevated cell NAD+/NADH ratios

following serine withdrawal were more resistant to serine depletion compared to cells with

unchanged NAD+/NADH ratios (**Figure 2C**). The NAD+/NADH ratio was also elevated in cancer

cells with higher PHGDH protein levels upon serine depletion (A375, PHGDH-overexpressing

MDA-MB-231) (**Supplementary Figure 2B,E**). To confirm that an elevated NAD+/NADH ratio in

response to serine depletion was associated with elevated serine synthesis, we compared how

serine synthesis rates changed following serine withdrawal in A549 and H1299 cells. Indeed, in

serine depleted conditions, H1299 cells demonstrated a higher rate of serine synthesis, a larger

increase in serine synthesis, and had higher serine levels than A549 cells, consistent with the

elevated NAD+/NADH ratio in H1299 cells (**Figure 2D,E; Supplementary Figure 2F,G**). Taken

together, these data suggest that in some cells, the cell NAD+/NADH ratio is responsive to serine

availability and increases in the NAD+/NADH ratio correlate with increased serine synthesis and

proliferation rates in serine-depleted conditions independent of PHGDH protein expression.

Increased mitochondrial respiration is associated with an elevated NAD+/NADH ratio and

increased serine synthesis

The finding that some cancer cells exhibit a more oxidized NAD+/NADH ratio upon serine

starvation is surprising, particularly when increased serine synthesis will result in a higher demand

for NAD+ to support increased flux through GAPDH and PHGDH (Diehl, 2019; Maddocks, 2017).

We hypothesized that the elevated NAD+/NADH ratio represented a cellular response to make

the NAD+/NADH ratio more oxidized to enable serine synthesis and thus sought to identify the

processes that led to the increased NAD+/NADH ratio in some but not all cancer cells. Lactate

production via lactate dehydrogenase (LDH) is a major NAD+ regenerating process, particularly

in cancer cells with high glucose fermentation (Fantin, 2006; Luengo, 2021; Wang, 2022)

(**Supplementary Figure 3A**); however, lactate secretion relative to glucose consumption was not

elevated following serine deprivation in any cells tested (**Supplementary Figure 3B**).

Mitochondrial respiration is another major way cells regenerate NAD+ from NADH (Handy, 2012).

Thus, we tested whether changes in mitochondrial respiration, as measured by cellular oxygen

consumption, accompany the increase in NAD+/NADH ratio following serine withdrawal. Indeed,

cells that increased their NAD+/NADH ratio following serine deprivation also exhibited increased

rates of oxygen consumption, whereas cells with unaltered NAD+/NADH ratios following serine

withdrawal did not (**Figure 3A, Supplementary Figure 3C**). To test whether elevated oxygen

consumption was related to serine deprivation, we cultured H1299 and A549 cells in varying

amounts of extracellular serine and measured oxygen consumption rates. Oxygen consumption

of A549 cells was not impacted by varying extracellular serine levels, while oxygen consumption

of H1299 cells increased as extracellular serine decreased below 100 µM, the serine

concentration where proliferation begins to be affected most by serine limitation (**Figure 3B,**

Supplementary Figure 3D). The increase in oxygen consumption rate also corresponded with

the NAD+/NADH ratio in H1299 cells at different serine levels, while the NAD+/NADH ratio in

A549 cells was unchanged by extracellular serine availability (**Figure 3C, D).** Consistently, the

NAD+/NADH ratio in H1299 cells cultured in different concentrations of serine positively

correlated with serine synthesis, whereas A549 cells displayed minimal changes in serine

synthesis rate (**Figure 3E, F**). Notably, PHGDH protein levels remained constant as extracellular

serine levels were decreased in both A549 and H1299 cells (**Supplementary Figure 3E**).

Together, these findings suggest that increased NAD+/NADH ratios in response to decreasing

environmental serine are linked to elevated mitochondrial respiration in select cancer cells, and

that this enables increased serine synthesis.

Mitochondrial respiration governs the cell NAD+/NADH ratio and influences serine

synthesis rate

To test whether increased mitochondrial respiration following serine depletion causes an

increase in the cell NAD+/NADH ratio, we treated redox unresponsive cells (i.e. cells that do not

elevate oxygen consumption or the NAD+/NADH ratio following serine withdrawal) with the proton

ionophore carbonyl cyanide-4-(trifluoromethoxy)phenylhydrazone (FCCP), which

pharmacologically increases mitochondrial oxygen consumption by uncoupling respiration from

ATP synthesis (McLaughlin, 1980; Terada, 1990). Increasing respiration with FCCP elevates

mitochondrial NAD+ regeneration and the cell NAD+/NADH ratio (Luengo, 2021) (**Figure 4A**).

Indeed, FCCP treatment dose-dependently improved proliferation of redox unresponsive cells in

serine-depleted conditions (**Figure 4B, Supplementary Figure 4A**). In contrast, FCCP did not

improve the proliferation of redox responsive cells grown without serine (**Figure 4C,**

Supplementary Figure 4B). Consistent with its impact on proliferation, FCCP treatment led to

higher serine synthesis rates only in redox unresponsive A549 cells while having no impact on

the serine synthesis rates of redox responsive H1299 cells (**Figure 4D,E, Supplementary Figure**

5A-D). As expected, uncoupling mitochondrial respiration from ATP synthase with FCCP led to

elevated oxygen consumption in serine depleted A549 cells (**Figure 4F).** FCCP also increased

oxygen consumption in serine depleted H1299 cells (**Supplementary Figure 4C**). Interestingly,

FCCP raised the NAD+/NADH ratios of both A549 and H1299 cells following serine withdrawal

despite only improving serine synthesis and proliferation in A549 cells (**Figure 4G,**

Supplementary Figure 4D). The lack of impact on serine synthesis by FCCP in H1299 cells

suggests that NAD+ availability may not further constrain serine synthesis in cells that already

elevate mitochondrial respiration following environmental serine depletion. In this context, we

considered whether another constraint such as PHGDH protein expression was limiting serine

synthesis in redox responsive cells treated with FCCP. Indeed, overexpressing PHGDH in serine-

depleted redox responsive H1299 and HCT116 cells fully restored proliferation to that observed

in serine-replete conditions (**Supplementary Figure 4E,G**). In contrast, proliferation of redox

unresponsive cells following serine withdrawal had the greatest improvement with simultaneous

PHGDH overexpression and FCCP treatment (**Supplementary Figure 4F,G**). This underscores

that changes in both the NAD+/NADH ratio and relevant enzyme expression impact serine

synthesis in serine depleted environments.

To investigate whether FCCP improved serine synthesis and proliferation of redox

unresponsive cells through oxidizing the NAD+/NADH ratio, we treated cells with the

mitochondrial complex I inhibitor rotenone in conjunction with FCCP. Rotenone treatment blocked

the ability of FCCP to improve both proliferation and serine synthesis and suppressed the

increase in the NAD+/NADH ratio (**Figure 4H-J, Supplementary Figure 5E,F**). In contrast,

treating cells with oligomycin, an ATP synthase inhibitor, did not affect the ability of FCCP to

improve proliferation (**Supplementary Figure 4H,I**). This supports the conclusion that FCCP

increases serine synthesis and proliferation in serine depleted environments by increasing the

NAD+/NADH ratio via mitochondrial complex I-mediated NAD+ regeneration. Moreover, FCCP

treatment neither increased PHGDH protein expression nor glucose consumption, indicating that

improved proliferation and serine synthesis is not caused by changes in enzyme expression or

carbon availability for serine synthesis (**Supplementary Figure 4J,K**). Together, these data

argue that mitochondrial respiration, either endogenously increased in response to serine

deprivation or as a result of FCCP treatment, can govern the cell NAD+/NADH ratio and directly

influence the serine synthesis rate in response to serine withdrawal.

The NAD+/NADH ratio influences oxidative citrate synthesis in either serine- or lipid-

depleted environments

Oxidative biosynthetic reactions other than serine synthesis can be constrained by the

NAD+/NADH ratio. For example, cancer cells deprived of environmental lipids increase oxidative

citrate production, which can be limited by NAD+ regeneration (Li, 2022) (**Figure 5A,**

Supplementary Figure 6A). Interestingly, analysis of kinetic U-^{13}C-glucose tracing in cells that

showed increased NAD+/NADH ratios following serine deprivation largely showed elevated M+2

citrate production despite no change in environmental lipid availability (**Figure 5B;**

Supplementary Figure 6B-E). In contrast, cells with unaltered NAD+/NADH ratios upon serine

depletion did not exhibit higher M+2 citrate production **(Figure 5B; Supplementary Figure 6B,**

F-H). Similarly, only cells that increased the NAD+/NADH ratio in response to serine depletion

displayed evidence of elevated citrate production from glutamine either through oxidative

decarboxylation or reductive carboxylation, both of which require oxidation reactions that can be

limited by electron acceptor availability (Hosios and Vander Heiden, 2018; Li, 2022)

(**Supplementary Figure 6A, I, J**). To confirm that elevated M+2 citrate synthesis was driven by

an increase in the NAD+/NADH ratio, we treated A549 cells depleted of serine with FCCP and

examined glucose derived citrate synthesis. Indeed, FCCP treatment led to a four-fold increase

in M+2 citrate from U-^{13}C-glucose in A549 cells depleted of serine whereas M+2 citrate synthesis

was less affected by FCCP in H1299 cells (**Figure 5C**). The elevated M+2 citrate production in

FCCP treated-A549 cells was dependent on mitochondrial complex I, as rotenone prevented the

elevated citrate production associated with FCCP treatment (**Supplementary Figure 7A-C**). To

confirm that higher M+2 citrate production following FCCP treatment was related to the

NAD+/NADH ratio independent of serine deprivation, we treated A549 cells with increasing doses

of AKB and FCCP in serine replete conditions and find that increasing the NAD+/NADH ratio with

either AKB or FCCP led to elevated M+2 citrate synthesis (**Figure 5D**). This emphasizes that

changes to the NAD+/NADH ratio regardless of the nutrient environment can impact citrate

synthesis.

We next asked whether increasing mitochondrial respiration in response to low

extracellular serine was a general response to other nutrient deprivations that increased NAD+

demand. To examine this, we depleted A549 and H1299 cells of extracellular lipids by culturing

them in media with lipid-depleted serum (Hosios, 2018) and compared oxygen consumption rates

to those of cells cultured in media with lipid-containing serum. Surprisingly, we find that A549 cells,

which do not increase mitochondrial respiration in response to serine depletion, mildly increase

oxygen consumption rates upon lipid depletion. On the other hand, H1299 cells do not increase

oxygen consumption in lipid depleted conditions (**Figure 5E**). We next measured the

NAD+/NADH ratio of A549 and H1299 cells depleted of environmental lipids. Consistent with the

changes in oxygen consumption, A549 cells displayed a slightly increased NAD+/NADH ratio

while H1299 cells did not alter the NAD+/NADH ratio upon lipid deprivation (**Figure 5F**).

Interestingly, both serine and citrate produced via glucose oxidation were elevated in lipid-

deprived A549 cells in comparison to H1299 cells, which exhibited unchanged serine synthesis

in response to environmental lipid depletion (**Figure 5G-H, Supplementary Figure 8A-D**). These

findings demonstrate that the NAD+/NADH ratio is a determinant of both oxidative citrate and

serine synthesis, regardless of environmental demand for the nutrient. Additionally, the different

metabolic responses to lipid and serine deprivation between cancer cell types suggests that

different nutrient environments may impact mitochondrial respiration through distinct processes

to influence the cell NAD+/NADH ratio.

Lipid depletion can enhance serine synthesis and proliferation in serine deprived

conditions

Our finding that both serine and oxidative citrate synthesis are impacted by levels of a

different nutrient than that produced by each pathway raised the possibility that the NAD+/NADH

ratio coordinates biomass synthesis in response to the availability of multiple oxidized nutrients.

Specifically, we hypothesized that changes to the NAD+/NADH ratio caused by one oxidized

nutrient depletion could influence the synthesis of, and in turn the dependency on, another

oxidized nutrient for proliferation. To begin testing this, we first examined how mitochondrial

respiration and the NAD+/NADH ratio were impacted by dual lipid and serine depletion compared

to single nutrient depletions in both A549 and H1299 cells. We find that dual depletion of serine

and lipids in A549 cells led to elevated oxygen consumption that is comparable to that observed

with lipid depletion alone, and higher than that observed with serine depletion alone **(Figure 6A)**.

Similarly, dual depletion of serine and lipids in H1299 cells led to elevated oxygen consumption

that is comparable to that observed with serine depletion alone, and higher than that observed

with lipid depletion alone **(Figure 6A)**. The resulting oxygen consumption rates upon dual serine

and lipid starvation corresponded with changes to the cell NAD+/NADH ratio (**Figure 6B**). We

then measured the serine and citrate synthesis rates of cells depleted of both extracellular serine

and lipids via kinetic U-^{13}C-glucose tracing. Consistent with elevation of the NAD+/NADH ratio

after lipid deprivation, A549 cells depleted of both environmental serine and lipids trended towards

exhibiting a slightly elevated serine synthesis rate when compared to serine deprivation alone

(Figure 6C). Moreover, lipid depletion led to a greater fraction of total serine derived from glucose

in serine depleted A549 cells **(Supplementary Figure 9A)**. In contrast, lipid starvation did not

alter serine synthesis in serine-depleted H1299 cells (**Figure 6D, Supplementary Figure 9B**).

Interestingly, in both A549 and H1299 cells, dual serine and lipid deprivation led to elevated citrate

production (to a greater degree in H1299 cells) compared to lipid deprivation alone despite only

H1299 cells exhibiting a more oxidized NAD+/NADH ratio upon serine depletion (**Figure 6E,F;**

Supplementary Figure 9C,D). This underscores the multifactorial regulation of glucose oxidation

to citrate (Holness, 2003; Novelli, 1950; Patel, 2006; Srere, 1974). We next asked how cancer

cell proliferation was impacted by dual serine and lipid depletion compared to single nutrient

depletions. Strikingly, proliferation of serine-depleted A549 cells was improved with depletion of

extracellular lipids (**Figure 6G**), consistent with an elevation in mitochondrial respiration,

NAD+/NADH ratio, and serine synthesis in A549 cells upon dual serine and lipid depletion. Unlike

serine depletion, lipid depletion does not cause a major proliferation defect in either A549 or

H1299 cells, and dual serine and lipid deprivation did not greatly alter the proliferation of H1299

cells compared to serine depletion alone (**Figure 6H**). This is consistent with lipid deprivation

causing no major change in oxygen consumption, NAD+/NADH ratio, and serine synthesis in

H1299 cells. Taken together, these data argue that the cell NAD+/NADH ratio is modulated in

response to multiple nutrient availabilities in a cancer cell-specific fashion. The changes in

NAD+/NADH ratio in turn influence the capacity to synthesize oxidative biomass precursors such

as serine and citrate, impacting the proliferation rate of cancer cells in nutrient environments with

differing demands for oxidative biomass synthesis.

Discussion

Proliferating cancer cells must acquire sufficient biomass and rely on contributions from

both environmental biomass precursors and de novo biomass synthesis, but what determines

which cells can adapt to different nutrient environments that impose constraints on salvage and

de novo synthesis is not known. This study highlights that differential cell responses to change

NAD+/NADH ratio enables biomass synthesis and determines whether cells can adapt to

proliferate in different nutrient conditions. Importantly, this provides one explanation for why single

nutrient levels in tissues may not predict the ability of auxotrophy for a nutrient to determine cancer

proliferation in a tissue (Abbott, 2024). It also suggests why cancer proliferation in different

nutrient environments is controlled by a complex relationship between cell-intrinsic and cell-

extrinsic factors. These factors may be coordinated in part through processes that set the cell

NAD+/NADH ratio.

Most prior work to determine what shapes the metabolic dependencies of cell proliferation

has focused on factors that govern biosynthetic enzyme expression and activity. For example,

genomic and transcriptional analyses of cancer discovered that increased PHGDH expression

promotes tumor growth (DeNicola, 2015; Locasale, 2011; Newman, 2017; Possemato, 2011) and

provides a proliferative advantage in serine depleted tumor microenvironments (Ngo, 2020;

Sullivan, 2019b). Our findings show that enzyme expression alone cannot fully explain nutrient

dependencies, as PHGDH expression can be discordant with sensitivity to serine depletion.

Rather both the NAD+/NADH ratio and enzyme expression play integral roles in determining

cancer proliferation in specific nutrient environments. These data are consistent with biosynthetic

gene expression and single nutrient levels being unable to predict the tumor tissue environments

where cancer cells can form metastases (Abbott, 2024).

Extensive work has highlighted the importance of cell NAD+/NADH ratio in influencing

oxidized biomass production, nutrient utilization, and nutrient dependencies (Baksh, 2020; Bao,

2016; Birsoy, 2015; Broeks, 2023; Christensen, 2014; Diehl, 2019; Garcia-Bermudez, 2018;

Gibson, 1976; Krall, 2021; Li, 2022; Schwörer, 2020; Sullivan, 2015; Wu, 2024). Yet, the

processes that govern the endogenous NAD+/NADH ratio upon shifts in nutrient availability have

been largely uncharacterized. We find that regulation of mitochondrial respiration is a major

mechanism that dictates the NAD+/NADH ratio in response to environmental nutrient fluctuations.

The mitochondrial respiration response to withdrawal of specific nutrients appears to vary across

different cancers. Some cancer cells increased mitochondrial respiration to serine depletion, while

others had minimal responses. On the other hand, lipid depletion yielded mitochondrial respiration

changes in a different set of cells. No clear common genetic factor or tissue-of-origin was found

to align with those cells that do or do not have mitochondrial and redox responses to serine and/or

lipid depletion, and further work is needed to understand the mechanisms that enable increased

mitochondrial respiration to respond to nutrient deprivation. One possibility is that differences in

ATP consumption in response to nutrient deprivation contribute to the response. Stimulating ATP

consumption can increase mitochondrial respiration and the NAD+/NADH ratio by modulating

mitochondrial membrane potential (Bertholet, 2019; Buttgereit, 1995; Fisher-Wellman, 2012;

Luengo, 2021; Nobes, 1989; Vander Heiden, 1999). This suggests that ATP synthase activity can

constrain complex I activity to different extents across cancers, modulating the NAD+/NADH ratio

and sensitivity to serine or lipid depletion. However, the regulation of complex I, ATP synthesis,

and mitochondrial coupling efficiency is complicated and incompletely understood (Alavian, 2014;

Gao, 2018; Lapuente-Brun, 2013; Luengo, 2021; Qiu, 2014; Wang, 2022; Yao, 2019; Zhao, 2019).

Nevertheless, our findings underscore a relationship between environmental nutrient levels and

mitochondrial respiration to influence the NAD+/NADH ratio in cells.

Efforts to understand the processes that shape cancer cell metabolic dependencies have

largely focused on how cancer cells respond to single nutrient changes. However, through dual

serine and lipid depletion experiments, these findings suggest that single nutrient availabilities will

be unable to explain cancer metabolic dependencies in some contexts. For example, depleting

A549 cells of both extracellular serine and lipids paradoxically improved proliferation relative to

serine deprivation alone because the cell NAD+/NADH ratio responds to lipid deprivation to

enable increased serine synthesis. This suggests that the NAD+/NADH ratio can serve as a

metabolic regulatory node that orchestrates biomass synthesis and integrates changes to multiple

oxidized biomass components in the environment, providing a framework for understanding why

cancer nutrient dependencies vary across tissue microenvironments (Abbott, 2024). Because

oxidizing the cell redox state can increase flux through multiple pathways, the impact of cell redox

state on biosynthesis may also explain why some cells lose enzyme expression as this could

redirect limited resources toward biomass components that can otherwise be scavenged from the

environment (Delage, 2010; Pavlova, 2017; Rabinovich, 2015). Better understanding the

mechanisms cells use to alter respiration to adjust the NAD+/NADH ratio in response to nutrient

changes could lead to better understanding of the complex interplay between cell-intrinsic and

cell-extrinsic factors that determine cancer metabolic dependencies and enable better selection

of patients who will respond to therapies targeting metabolism.

Materials and Methods

Cell Culture Experiments

Cell lines were maintained in Dulbecco's Modified Eagle's Medium (DMEM) without sodium

pyruvate (Corning 50-013-PC) supplemented with sodium bicarbonate (Sigma S6014) and 10%

heat-inactivated Fetal Bovine Serum (FBS). All cells were cultured at 37°C with 5% CO_2, tested

for mycoplasma contamination regularly, and confirmed negative before experimentation. Cell

line information is listed in Supplementary Table 1.

Proliferation Assays

Cells were plated in six-well plates in 2 ml of DMEM with 10% heat-inactivated FBS at an initial

seeding density of 40,000 cells. Cells were permitted to settle overnight and then quantified using

a sulforhodamine B (SRB) (Sigma Aldrich 230162) colorimetric assay to measure the cell

absorbance at the start of the experiment as previously described (Vichai, 2006). For treatment

conditions, cells were washed three times with 1 ml of 1X PBS, and 2 ml of treatment media were

added to each well. All treatment media was made with 10% dialyzed FBS. Serine-free medium

was made by adding a mix of amino acids (Sigma Aldrich) at DMEM concentrations lacking serine

and glycine to DMEM with low glucose, without sodium pyruvate, and no amino acids (US

Biological D9800-13-50L). Glucose and glycine were supplemented for a final concentration of 25

mM and 0.4 mM, respectively. 72 hours after the initial treatment, cells were quantified using SRB

assay. All SRB measurements are normalized to a blank to correct for background signal.

Proliferation rates were calculated with the following formula:

Doublings per day = $\underline{log_2(\text{ final SRB absorbance / SRB absorbance day 0})}$

430   *(total # of treatment days)*

NAD+/NADH Measurements

NAD+/NADH measurements were performed using the NAD/NADH-Glo Assay (Promega G9072)

with a modified version of manufacturer instructions as previously reported (Sullivan, 2015). Cells

were plated at an initial seeding density of 20,000 cells (A549, H1299, HCT116) or 40,000 cells

(Calu6, MDA-MB-231). Cells were permitted to settle overnight. The next day, cells were washed

once with 1X PBS and then incubated with 2 ml of treatment media for the indicated times prior

to preparation of cell extracts. For extraction, cells were washed once in ice cold 1X PBS and

extracted in 100 µl ice-cold 1% dodecyltrimethylammonium bromide (DTAB) in 0.2 N NaOH

diluted 1:1 with 1X PBS. Each sample was flash-frozen in liquid nitrogen and immediately stored

at -80°C. To measure NADH, 10 µl of sample was moved to PCR tubes, diluted with 10 µl of

DTAB, and incubated at 75 °C for 30 min, where basic conditions selectively degrade NAD+. To

measure NAD+, 10 µl of the samples was moved to PCR tubes containing 30 µl DTAB and 20 µl

0.4 N HCl and incubated at 60°C for 15 min, where acidic conditions selectively degrade NADH.

Samples were then allowed to equilibrate to room temperature and quenched by neutralizing with

20 µl 0.25 M Tris in 0.2 N HCl (for NADH) or 20 µl 0.5 M Tris base (for NAD+). Manufacturer

instructions were followed thereafter to measure NAD+/NADH using a luminometer (Tecan Infinite

M200Pro). A standard curve with representative samples was done with each assay to confirm

that NAD+ and NADH measurements were in the linear range of detection.

Glucose Consumption and Lactate Secretion Measurements

Cells were plated at a density of 40,000 cells per well in a six well plate. The next day, cells were

washed three times with 1X PBS before 2 ml of treatment media were added to each well. In

parallel, 2 ml of treatment media were added to plates without cells. An initial cell number before

beginning treatment was quantified via Cellometer (Nexcelom Bioscience). 72 hours later, media

was collected from exponentially proliferating cells with a parallel collection of media from wells

without cells, which we used to measure initial media concentrations to take into consideration

evaporative changes. Cell number was quantified via Cellometer along with the volume of media

on cells to consider evaporative changes to metabolite concentrations that may have occurred

over 72 hours in culture. Glucose concentrations from media samples were measured on a YSI-

2900 Biochemistry Analyzer (Yellow Spring Instruments). Every assay included a glucose (Sigma

Aldrich G8270) standard curve (0, 2.5, 5, 10, 20, and 40 mM glucose) and lactate (Sigma Aldrich

71716) standard curve (0, 1.25, 2.5, 5, 10, and 20 mM sodium lactate), made in both control and

treatment media. To calculate consumption rate per cell for a given nutrient, fmol of glucose or

lactate over time were plotted relative to the area under the curve (fmol / cells · time) of an

exponential function fit to the number of cells at the start of the assay and after 72 hours of

treatment, as previously described (Hosios, 2016).

Oxygen Consumption Measurements

An Agilent Seahorse Bioscience Extracellular Flux Analyzer (XFe96) was used to measure

oxygen consumption rates (OCR). Cells were plated at 20,000 – 40,000 cells per well in a

Seahorse Bioscience 96-well plate in 80 µl of DMEM without pyruvate supplemented with 10%

heat-inactivated FBS. Cells were not plated on the perimeter of the plate to avoid edge effect.

The following day, cells were washed twice with 180 µl of treatment media supplemented with 10%

dialyzed FBS before incubating cells with 180 µl of treatment media for the indicated time prior to

OCR data acquisition. For 24 hour treatment periods, treatment media was replaced with 180 µl

of fresh treatment media two hours before OCR data acquisition. The day before OCR data

acquisition, the XFe96 cartridge was hydrated by submerging the sensor cartridge in 200 µl of

sterile water per well of the utility plate. Parafilm was wrapped around the perimeter of the

cartridge to avoid evaporation. The XFe96 cartridge was then incubated in a 37°C non-CO_2

incubator overnight. The next day, at least two hours before data acquisition, the water in the

utility plate was replaced with 200 µl of the Agilent XF Calibrant per well and the sensory cartridge

was submerged and returned to the 37°C non-CO_2 incubator until OCR was ready to be measured.

After OCR data acquisition, six to eight wells per treatment condition were collected for cell

number quantification (Cellometer) to normalize OCR values. Basal OCR was calculated by

subtracting residual OCR, the OCR remaining following the addition of rotenone and antimycin A

for a final concentration of 2 µM each.

Kinetic U-^{13}C-Glucose Isotope Tracing Experiments

Cells were seeded at 150,000 cells per well in a six-well plate in 2 ml of DMEM without sodium

pyruvate supplemented with 10% heat-inactivated FBS. The following day, cells were washed

once with 1 ml of 1X PBS and then incubated in 2 ml of treatment media supplemented with 10%

dialyzed FBS for the indicated treatment time and when metabolic steady state is reached. Two

hours before the start of the kinetic isotope tracing, media on cells was exchanged with fresh

treatment media with 10 mM unlabeled glucose. To measure serine labeling over time in serine-

depleted conditions, cells and media were extracted together. Before the initiation of the kinetic

U-^{13}C-glucose isotope tracing, serine-depleted cells were washed three times with 1X PBS and

then cultured with 400 µl of tracing medium containing 10 mM of U-^{13}C-glucose (Cambridge

Isotope Laboratories, CLM-481-0) for the rapid time points. Tracing media was equivalent to

treatment media except with 10 mM of U-^{13}C- glucose. Following incubation, 1.6 ml of extraction

buffer containing ice cold 100% HPLC-grade methanol (Sigma Aldrich, 646377) with norvaline

(Sigma, N7627) was added onto cells with tracing media for a final sample volume of 2 ml (80%

extraction buffer and 20% tracing media) and a final concentration of 1 µg per 400 µl norvaline.

1.6 ml of the lysate was placed into a fresh Eppendorf tube, vortexed at 4°C at maximum speed

for ten minutes, and then spun down at 4°C at maximum speed for thirty minutes. The supernatant

was collected and dried under nitrogen gas to prepare for metabolic analysis. To measure serine

and citrate production in serine replete conditions, only intracellular metabolites were quantified.

Before the initiation of kinetic U-^{13}C-glucose isotope tracing, serine-replete cells were washed

three times with 1X PBS and then cultured with 1 ml of tracing medium containing 10 mM of U-

^{13}C-glucose (Cambridge Isotope Laboratories, CLM-481-0) for the rapid time points. Tracing

media was equivalent to treatment media except with 10 mM of U-^{13}C- glucose. Following

incubation, cells were rapidly washed twice with ice cold blood bank saline. To extract, 500 µl of

extraction buffer containing 80% ice cold HPLC-grade methanol and 20% LC-grade water spiked

513 with 1 µg per 400 µl norvaline were added onto cells. Cells were then lysed as previously

514 described above.

Gas Chromatography-Mass Spectrometry (GC-MS) Polar Metabolite Measurements

Dried samples were derivatized by adding 16 µl of methoxamine reagent (Thermo Fisher, TS-

45950) and incubated for an hour at 37°C followed by the addition of 20 µl of *N-tert*-

butyldimethylsilyl-*N*-methyltrifluoroacetamide with 1% *tert*-butyldimethylchlorosilane (Sigma

375934). Samples were then incubated for two hours at 60°C. Afterwards, samples were

centrifuged at maximum speed for ten minutes, and 20 µl of supernatant was used for analysis.

Following derivatization, samples were analyzed using a DB-35MS column (30 m × 0.25 mm

i.d. × 0.25 µm, Agilent J&W Scientific) in an Agilent 7890 gas chromatograph coupled to an Agilent

5975C mass spectrometer (GC–MS). All ion counts were normalized to internal standard

norvaline and cell number measured via Cellometer.

Immunoblotting

Cells were washed with ice cold 1X PBS and lysed in cold RIPA buffer containing Halt Protease

and Phosphatase Inhibitor Cocktail (Thermo Scientific 78442). Lysates were clarified by rocking

samples for thirty minutes at 4°C and then spun at maximum speed for ten minutes for collection

of supernatant. Protein concentration was calculating using the BCA Protein Assay (Pierce, 23225)

with BSA as a standard. Lysates were resolved by SDS-PAGE using NuPAGE 4-12% Bis-Tris

Protein Gels and run at 100 V. Proteins were transferred onto nitrocellulose membranes via wet

transfer at 100 V. Membranes were blocked in 5% BSA in 1x TBST before incubating membranes

with primary antibodies at 4°C overnight. The next day, membranes were washed three times

with 1x TBST at room temperature and then incubated in secondary antibodies for one hour at

room temperature. The primary antibodies used were PHGDH (Sigma Aldrich, HPA021241

[1:1000]), HSP90 (Cell Signaling Technology, 4874 [1:1000]), and Vinculin (E1E9V) XP® (Cell

Signaling Technology, 13901 [1:1000]). The secondary antibodies used were anti-rabbit IgG

horseradish peroxidase-linked antibody (1:4000 dilution; Cell Signaling Technologies, 7074S) or

anti-mouse IgG horseradish peroxidase-linked antibody (1:4000, 7076S).

Cell Line Generation

Cell lines overexpressing either EGFP or human PHGDH were generated via lentiviral infection.

pLJM1-EGFP was obtained from Addgene (Addgene plasmid # 19319). pLJM1-PHGDH was

constructed using pLJM1-Empty (Addgene plasmid # 91980) as a backbone. pLJM1-Empty was

digested using NheI and EcoRI and gel purified. Human PHGDH cDNA was amplified from

pLHCX-PHGDH (PMID: 31598584) with the following oligonucleotide primers, where the

capitalized sequence denotes PHGDH homology:

PHGDH F: agtgaaccgtcagatccggctagcgccaccATGGCTTTTGCAAATCTGCG

PHGDH R: tactgccatttgtctcgaggtcgagaattcTTAGAAGTGGAACTGGAAGGCT

PHGDH insert was gradient PCR amplified using Phusion High-Fidelity DNA polymerase (NEB

M0530) from 60 to 70°C and gel purified. Digested pLJM1-Empty and amplified PHGDH insert

were assembled via Gibson Assembly, and the resulting plasmid was transformed, and

validated by Sanger sequencing. Lentiviral production was done by transfecting constructs into

LentiX293T cells with Mirus Transit293T (Mirus Bio MIR2700) following manufacturer's protocol.

Briefly, 800,000 cells were plated per well in a 6 well plate in DMEM media with 10% heat-

inactivated FBS. The next day, cells were transfected with 1.6 µg vector (either pLJM1-EGFP or

pLJM1-PHGDH), 800 ng of pMDLg (Addgene plasmid # 12251) packaging plasmid, 400 ng of

pMD2.G (Addgene plasmid # 12259) envelope plasmid, and 400 ng of pRSV-REV (Addgene

plasmid # 12253) in OptiMEM (Thermo Fisher Scientific 31985070). DNA transfection mix was

incubated with cells for 48 hours before harvesting virus and filtering through a 0.45 µm

polyethersulfone membrane. Sub-confluent cells were infected with 1 ml of virus-containing

media in each well of a 6 well plate with polybrene (8 µg/ml, EMD Millipore TR-1003-G). 24

hours after infection, selection with 1 µg/mL puromycin (Sigma Aldrich P7255) began. All

experiments with generated cell lines were conducted on a polyclonal cell population.

Statistics and Reproducibility

All statistical tests using experimental data were performed with Prism 10 software. No statistical

method was used to predetermine sample sizes. Samples sizes were chosen based on pilot

experiments using three or more technical replicates. Certain data points were excluded from

Seahorse analyzer assays due to premature injection of rotenone and antimycin. The experiments

were not randomized and the investigators were not blinded during experiments nor data analysis.

Acknowledgements

We thank all members of the Vander Heiden lab for helpful discussions, Emma Dawson, Grace

Wolczanski, and Alejandra Rosario-Crespo for technical assistance, and Jessica Spinelli for

assistance with using Seahorse equipment during the COVID-19 pandemic. The authors

acknowledge support from F30CA268633 (SMC), T32GM007287 (SMC, MBM, ASZ, KLA), the

Kosciuszko Foundation (SET), NSF DGE-1122374 (KLA), and F31CA271787 (KLA), as well as

support to MGVH from the Ludwig Center at MIT, the MIT Center for Precision Cancer Medicine,

and the NCI (R35CA242379, R01CA259253, P30CA1405141).

Author Contributions

Conceptualization: SMC, MGVH; Intellectual Discussion: SMC, MBM, MGVH; Proliferation rates:

SMC, SET, ASZ; Kinetic Isotope Tracing: SMC, SET, MBM; Immunoblotting: SMC, SET;

NAD+/NADH measurements: SMC, SET; Oxygen consumption: SMC; Overexpression constructs:

KLA; Cell line generation: SMC; Writing: SMC and MGVH; Editing: SMC, MBM, SET, ASZ, KLA,

MGVH; Funding Acquisition: MGVH

Competing Interests

M.G.V.H. discloses that he is a scientific advisor for Agios Pharmaceuticals, iTeos Therapeutics, Sage Therapeutics, Pretzel Therapeutics, Lime Therapeutics, Faeth Therapeutics, Droia Ventures, MPM Capital and Auron Therapeutics. All remaining authors declare no competing interests.

References

Abbott, K.L., Subudhi, S., Ferreira, R., Gültekin, Y., Steinbuch, S.C., Bin Munim, M., Honeder, S.E., Kumar, A.S., Ramesh, D.L., Wu, M., Hansen, J.A., Sivanand, S., Reidmayr, L.M., Duquette, M., Ali, A., Henning, N., Shevzov-Zebrun, A., Gourgue, F., Barbeau, A.M., Waite, M., Kunchok, T., Ferraro, G.B., Do, B.T., Spanoudaki, V., Sánchez-Rivera, F.J., Jin, X., Church, G.M., Jain, R.K., Vander Heiden, M.G. (2024). Site of breast cancer metastasis is independent of single nutrient levels. bioRxiv. 10.1101/2024.10.24.616714.

Alavian, K.N., Beutner, G., Lazrove, E., Sacchetti, S., Park, H.-A., Licznerski, P., Li, H., Nabili, P., Hockensmith, K., Graham, M., Porter, G. A., & Jonas, E. A. (2014). An uncoupling channel within the c-subunit ring of the F1FO ATP synthase is the mitochondrial permeability transition pore. PNAS *111*, 10580–10585. https://doi.org/10.1073/pnas.1401591111.

Apiz Saab, J.J., Dzierozynski, L.N., Jonker, P.B., AminiTabrizi, R., Shah, H., Elena Menjivar, R., Scott, A.J., Nwosu, Z.C., Zhu, Z., Chen, R.N., Oh, M., Sheehan, C., Wahl, D.R., Rasca di Magliano, M., Lyssiotis, C.A., Macleod, K.F., Weber, C.R., Muir, A. (2023). Pancreatic tumors exhibit myeloid-driven amino acid stress and upregulate arginine biosynthesis. eLife *12*. https://doi.org/10.7554/eLife.81289.

Baksh, S.C., Todorova, P.K., Gur-Cohen, S., Hurwitz, B., Ge, Y., Novak, J.S.S., Tierney, M.T., dela Cruz-Racelis, J., Fuchs, E., Finley, L.W.S. (2020). Extracellular serine controls epidermal stem cell fate and tumour initiation. Nature Cell Biology *22*, 779-790. https://doi.org/10.1038/s41556-020-0525-9.

Banh, R.S., Biancur, D.E., Yamamoto, K., Sohn, A.S.W., Walters, B., Kuljanin, M., Gikandi, A., Wang, H., Mancias, J.D., Schneider, R.J., Pacold, M.E., Kimmelman, A.C. (2020). Neurons Release Serine to Support mRNA Translation in Pancreatic Cancer. Cell *183*, 1202-1218. 10.1016/j.cell.2020.10.016.

Bao, X.R., Ong, S.-E., Goldberger, O., Peng, J., Sharma, R., Thompson, D. A., Vafai, S. B., Cox, A. G., Marutani, E., Ichinose, F., Goessling, W., Regev, A., Carr, S. A., Clish, C. B., & Mootha, V. K. (2016). Mitochondrial dysfunction remodels one-carbon metabolism in human cells. eLife *5*. https://doi.org/10.7554/elife.10575.

Bertholet, A.M., Chouchani, E. T., Kazak, L., Angelin, A., Fedorenko, A., Long, J. Z., Vidoni, S., Garrity, R., Cho, J., Terada, N., Wallace, D. C., Spiegelman, B. M., & Kirichok, Y. (2019). H+ transport is an integral function of the mitochondrial ADP/ATP carrier. Nature *571*, 515–520. https://doi.org/10.1038/s41586-019-1400-3.

635
636 Birsoy, K., Wang, T., Chen, W.W., Freinkman, E., Abu-Remaileh, M., Sabatini, D.M. (2015). An
637 Essential Role of the Mitochondrial Electron Transport Chain in Cell Proliferation Is to Enable
638 Aspartate Synthesis. Cell *162*, 540-551. https://doi.org/10.1016/j.cell.2015.07.016.
639
640 Broeks, M.H., Meijer, N.W.F., Westland, D., Bosma, M., Gerrits, J., German, H.M., Ciapaite, J.,
641 van Karnebeek, C.D.M., Wanders, R.J.A., Zwartkruis, F.J.T., Verhoeven-Duif, N., Jans, J.J.M.
642 (2023). The malate-aspartate shuttle is important for de novo serine biosynthesis. Cell Reports
643 *42*. https://doi.org/10.1016/j.celrep.2023.113043.
644
645 Buttgereit, F., Brand, M.D. (1995). A hierarchy of ATP-consuming processes in mammalian
646 cells. Biochemical Journal *312*, 163-167. A hierarchy of ATP-consuming processes in
647 mammalian cells.
648
649 Chang, S.M. (2024). Control of Cellular Redox State and Biomass Synthesis. PhD Thesis
650 *Massachusetts Institute of Technology* https://dspace.mit.edu/handle/1721.1/155353.
651
652 Christensen, C.E., Karlsson, M., Winther, J.R., Jensen, P.R., Lerche, M.H. (2014). Non-invasive
653 In-cell Determination of Free Cytosolic [NAD+]/[NADH] Ratios Using Hyperpolarized Glucose
654 Show Large Variations in Metabolic Phenotypes. Journal of Biological Chemistry *289*, 2344-
655 2352. 10.1074/jbc.M113.498626.
656
657 Cognet, G., Muir, A. (2024). Identifying metabolic limitations in the tumor microenvironment.
658 Science Advances *10*. 10.1126/sciadv.adq7305.
659
660 Deberardinis, R.J., Lum, J.J., Hatzivassiliou, G., Thompson, C.B. (2008). The Biology of
661 Cancer: Metabolic Reprogramming Fuels Cell Growth and Proliferation. Cell Metabolism *7*, 11-
662 20.
663
664 Delage, B., Fennell, D.A., Nicholson, L., McNeish, I., Lemoine, N.R., Crook, T., Szlosarek, P.W.
665 (2010). Arginine deprivation and argininosuccinate synthetase expression in the treatment of
666 cancer. International Journal of Cancer *126*, 2762-2772. 10.1002/ijc.25202.
667
668 DeNicola, G.M., Chen P., Mullarky E., Sudderth, J.A., Hu, Z., Wu, D., Tang, H., Xie, Y., Asara
669 J.M., Huffman, K.E., Wistuba, I.I., Minna, J.D., DeBerardinis, R.J., Cantley, L.C. (2015). NFR2
670 regulates serine biosynthesis in non-small cell lung cancer. Nature Genetics *47*, 1475-1481.
671 10.1038/ng.3421.
672
673 Diehl, F.F., Lewis, C.A., Fiske, B.P., Vander Heiden, M.G. (2019). Cellular redox state
674 constrains serine synthesis and nucleotide production to impact cell proliferation. Nature
675 Metabolism *1*, 861-867. doi:10.1038/s42255-019-0108-x.
676
677 Fantin, V.R., St-Pierre, J., Leder, P. (2006). Attenuation of LDH-A expression uncovers a link
678 between glycolysis, mitochondrial physiology, and tumor maintenance. Cancer Cell *9*, 425-434.
679 10.1016/j.ccr.2006.04.023.
680
681 Faubert, B., Solmonson, A., Deberardinis, R.J. (2020). Metabolic reprogramming and cancer
682 progression. Science *368*. DOI: 10.1126/science.aaw5473.
683
684 Ferraro, G.B., Ali, A., Luengo, A., Kodack, D.P., Deik, A., Abbott, K.L., Bezwada, D., Blanc, L.,
685 Prideaux, B., Jin, X., Possada, J.M., Chen, J., Chin, C.R., Amoozgar, Z., Ferreira, R., Chen, I.,

Naxerova, K., Ng, C., Westermark, A.M., Duquette, M., Roberge, S., Lindeman, N.I., Lyssiotis, C.A., Nielsen, J., Housman, D.E., Duda, D.G., Brachtel, E., Golub, T.R., Cantley, L.C., Asara, J.M., Davidson, S.M., Fukumua, D., Dartois, V.A., Clish, C.B., Jain, R.K., Vander Heiden, M.G. (2021). Fatty acid synthesis is required for breast cancer brain metastasis. Nature Cancer *2*, 414-428. https://doi.org/10.1038/s43018-021-00183-y.

Fisher-Wellman, K.H., Neufer, P.D. (2012). Linking mitochondrial bioenergetics to insulin resistance via redox biology. Trends in Endocrinology & Metabolism *23*, 142-153. 10.1016/j.tem.2011.12.008.

Gao, X., Lee, K., Reid, M.A., Sanderson, S.M., Qiu, C., Li, S., Liu, J., Locasale, J.W. (2018). Serine Availability Influences Mitochondrial Dynamics and Function through Lipid Metabolism. Cell Reports *22*, 3507-3520. https://doi.org/10.1016/j.celrep.2018.03.017.

Garcia-Bermudez, J., Baudrier, L., La, K., Ge Zhu, X., Fidelin, J., Sviderskiy, V.O., Papagiannakopoulos, T., Molina, H., Snuderl, M., Lewis, C.A., Possemato, R.L., Birsoy, K. (2018). Aspartate is a limiting metabolite for cancer cell proliferation under hypoxia and in tumours. Nature Cell Biology *20*, 775-781. 10.1038/s41556-018-0118-z.

Gibson, G., Blass, J.P. (1976). A Relation between [NAD+]/[NADH] Potentials and Glucose Utilization in Rat Brain Slices. Journal of Biological Chemistry *251* (*13*), 4127-4130.

Gravel, S., Hulea, L., Toban, N., Birman, E., Blouin, M., Zakikhani, M., Zhao, Y., Topisirovic, I., St-Pierre, J., Pollak, M. (2014). Serine Deprivation Enhances Antineoplastic Activity of Biguanides. Cancer Research *74*, 7521-7533. https://doi.org/10.1158/0008-5472.CAN-14-2643-T.

Gullino, P.M., Clark, S.H., Grantham, F.H. (1964). The Interstitial Fluid of Solid Tumors. Cancer Research *24*, 780-797.

Hanahan, D., Weinberg, R.A. (2011). Hallmarks of Cancer: The Next Generation. Cell *144*, 646–674. https://doi.org/10.1016/j.cell.2011.02.013.

Handy, D.E., Loscalzo, J. (2012). Redox Regulation of Mitochondrial Function. Antioxidants & Redox Signaling *16*, 1323-1367. 10.1089/ars.2011.4123.

Holness, M.J., Sugden, M.C. (2003). Regulation of pyruvate dehydrogenase complex activity by reversible phosphorylation. Biochemical Society Transactions *31* (*6*), 1143-1151. 10.1042/bst0311143.

Hosios, A.M., Hecht, V.C., Danai, L.V., Johnson, M.O., Rathmell, J.C., Steinhauser, M.L., Manalis, S.R., Vander Heiden, M.G. (2016). Amino acids rather than glucose account for the majority of cell mass in proliferating mammalian cells. Developmental Cell *36*, 540-549. doi: 10.1016/j.devcel.2016.02.012.

Hosios, A.M., Li, Z., Lien, E.C., Vander Heiden, M.G. (2018). Preparation of Lipid-Stripped Serum for the Study of Lipid Metabolism in Cell Culture. Bio-protocol *8*. 10.21769/BioProtoc.2876

735   Hosios, A.M., and Vander Heiden, M.G. (2018). The redox requirement of proliferation
736   mammalian cells. Journal of Biological Chemistry *293*, 7490-7498.
737   https://doi.org/10.1074/jbc.TM117.000239.
738
739   Krall, A.S., Mullen, P.J., Surjano, F., Momcilovic, M., Schmid, E.W., Halbrook, C.J., Thambundit,
740   A., Mittelman, S.D., Lyssiotis, C.A., Shackelford, D.B., Knott, S.R.V., Christofk, H.R. (2021).
741   Asparagine couples mitochondrial respiration to ATF4 activity and tumor growth. Cell
742   Metabolism *33*, 1013-1026. https://doi.org/10.1016/j.cmet.2021.02.001.
743
744   Labuschagne, C.F., van den Broek, N.J.F., Mackay, G.M., Vousden, K.H., Maddocks, O.D.K.
745   (2014). Serine, but not glycine, supports one-carbon metabolism and proliferation of cancer
746   cells. Cell Reports *7*. doi: 10.1016/j.celrep.2014.04.045.
747
748   Lapuente-Brun, E., Moreno-Loshuertos, R., Acin-Perez, R., Latorre-Pellicer, A., Colas, C.,
749   Balsa, E., Perales-Clemente, E., Quiros, P.M., Calvo, E., Rodriguez-Hernandez, M.A., Navas,
750   P., Cruz, R., Carracedo, A., Lopez-Otin, C., Perez-Martos, A., Fernandez-Silva, P., Fernandez-
751   Vizarra, E., Enriquez, J.A. (2013). Supercomplex Assembly Determines Electron Flux in the
752   Mitochondrial Electron Transport Chain. Science *340*, 1567-1570. 10.1126/science.1230381.
753
754   Li, Z., Ji, B.W., Dixit, P.D., Lien, E.C., Tchourine, K., Hosios, A.M., Abbott, K.L., Westermark,
755   A.M., Gorodetsky, E.F., Sullivan, L.B., Vander Heiden, M.G., Vitkup, D. (2022). Cancer cells
756   depend on environmental lipids for proliferation when electron acceptors are limited. Nature
757   Metabolism *4*, 711-723. https://doi.org/10.1038/s42255-022-00588-8.
758
Locasale, J.W., Grassian, A.R., Melman, T., Lyssiotis, C.A., Mattaini, K.R., Bass, A.J., Heffron,
G., Metallo, C.M., Muranen, T., Sharfi, H., Sasaki, A.T., Anastasiou, D., Mullarky, E., Vokes,
N.I., Sasaki, M., Beroukhim, R., Stephanopoulos, G., Ligon, A.H., Meyerson, M., Richardson,
A.L., Chin, L., Wagner, G., Asara, J.M., Brugge, J.S., Cantley, L.C., Vander Heiden, M.G.
(2011). Phosphoglycerate dehydrogenase diverts glycolytic flux and contributes to oncogenesis.
Nature Genetics *43*, 869-874. 10.1038/ng.890.
765
766   Luengo, A., Li, Z., Gui, D. Y., Sullivan, L. B., Zagorulya, M., Do, B. T., Ferreira, R., Naamati, A.,
767   Ali, A., Lewis, C. A., Thomas, C. J., Spranger, S., Matheson, N. J., & Vander Heiden, M. G.
768   (2021). Increased demand for NAD+ relative to ATP drives aerobic glycolysis. Molecular Cell
769   *81*, 691–707. https://doi.org/10.1016/j.molcel.2020.12.012.
770
771   Lyssiotis, C.A., Kimmelman, A.C. (2017). Metabolic Interactions in the Tumor Microenvironment.
772   Trends in Cell Biology *27*, 863-875. https://doi.org/10.1016/j.tcb.2017.06.003.
773
774   Maddocks, O.D.K., Athineos, D., Cheung, E. C., Lee, P., Zhang, T., Van Den Broek, N. J. F.,
775   Mackay, G. M., Labuschagne, C. F., Gay, D., Kruiswijk, F., Blagih, J., Vincent, D. F., Campbell,
776   K. J., Ceteci, F., Sansom, O. J., Blyth, K., & Vousden, K. H. (2017). Modulating the therapeutic
777   response of tumours to dietary serine and glycine starvation. Nature *544*, 372–376.
778   https://doi.org/10.1038/nature22056.
779
780   Maddocks, O.D.K., Berkers, C. R., Mason, S. M., Zheng, L., Blyth, K., Gottlieb, E., & Vousden,
781   K. H. (2013). Serine starvation induces stress and p53-dependent metabolic remodelling in
782   cancer cells. Nature *493* (*7433*), 542–546. https://doi.org/10.1038/nature11743.
783
McLaughlin, S.G., Dilger, J.P. (1980). Transport of protons across membranes by weak acids.
Physiological Reviews *60* (*3*), 825-863. 10.1152/physrev.1980.60.3.825.

Newman, A.C., Maddocks, O.D.K. (2017). One-carbon metabolism in cancer. British Journal of
Cancer *116*, 1499-1504. https://doi.org/10.1038/bjc.2017.118.

Ngo, B., Kim, E., Osorio-Vasquez, V., Doll, S., Bustraan, S., Liang, R. J., Luengo, A., Davidson,
S. M., Ali, A., Ferraro, G. B., Fischer, G. M., Eskandari, R., Kang, D. S., Ni, J., Plasger, A.,
Rajasekhar, V. K., Kastenhuber, E. R., Bacha, S., Sriram, R. K., … Pacold, M. E. (2020).
Limited Environmental Serine and Glycine Confer Brain Metastasis Sensitivity to PHGDH
Inhibition. Cancer Discovery *10*, 1352–1373. https://doi.org/10.1158/2159-8290.cd-19-1228.

Nobes, C.D., Lakin-Thomas, P.L., Brand, M.D. (1989). The contribution of ATP turnover by the
Na+/K+-ATPase to the rate of respiration of hepatocytes: Effects of thyroid status and fatty
acids. Biochimica Et Biophysica Acta (BBA) - Bioenergetics *976*, 241-245.
https://doi.org/10.1016/S0005-2728(89)80236-1.

Novelli, G.D., Lipmann, F. (1950). The Catalytic Function of Coenzyme A in Citric Acid
Synthesis. Journal of Biological Chemistry *182*, 213-228.

Patel, M.S., Korotchkina, L.G. (2006). Regulation of the pyruvate dehydrogenase complex.
Biochemical Society Transactions *34* (*2*), 217-222. 10.1042/BST20060217.

Pavlova, N.N., Thompson, C.B. (2017). The Emerging Hallmarks of Cancer Metabolism. Cell
Metabolism *23*, 27-47. 10.1016/j.cmet.2015.12.006.

Possemato, R., Marks, K.M., Shaul, Y.D., Pacold, M.E., Kim, D., Birsoy, K., Sethumadhavan,
S., Woo, H., Jang, H.G., Jha, A.K., Chen, W.W., Barrett, F.G., Stransky, N., Tsun, Z., Cowley,
G.S., Barretina, J., Kalaany, N.Y., Hsu, P.P., Ottina, K., Chan, A.M., Yuan, B., Garrayway, L.A.,
Root, D.E., Mino-Kenudson, M., Brachtel, E.F., Driggers, E.M., Sabatini, D.M. (2011).
Functional genomics reveals serine synthesis is essential in PHGDH-amplified breast cancer.
Nature *476*, 346-350. 10.1038/nature10350.

Qiu, F., Chen, Y.-R., Liu, X., Chu, C.-Y., Shen, L.-J., Xu, J., Gaur, S., Forman, H. J., Zhang, H.,
Zheng, S., Yen, Y., Huang, J., Kung, H.-J., & Ann, D. K. (2014). Arginine Starvation Impairs
Mitochondrial Respiratory Function in ASS1-Deficient Breast Cancer Cells. Science Signaling *7*,
ra31.

Rabinovich, S., Adler, L., Yizhak, K., Sarver, A., Silberman, A., Agron, S., Stettner, N., Sun, Q.,
Brandis, A., Helbling, D., Korman, S., Itzkovitz, S., Dimmock, D., Ulitsky, I., Nagamani, S. C. S.,
Ruppin, E., & Erez, A. (2015). Diversion of aspartate in ASS1-deficient tumours fosters de novo
pyrimidine synthesis. Nature *527*, 379-383. https://doi.org/10.1038/nature15529.

Schwörer, S., Berisa, M., Violante, S., Qin, W., Zhu, J., Hendrickson, R.C., Cross, J.R.,
Thompson, C.B. (2020). Proline biosynthesis is a vent for TGFβ-induced mitochondrial redox
stress. The EMBO Journal *39*. https://doi.org/10.15252/embj.2019103334.

Sciacovelli, M., Dugourd, A., Valcarcel Jimenez, L., Yang, M., Nikitopoulou, E., Costa, A.S.H.,
Tronci, L., Caraffini, V., Rodrigues, P., Schmidt, C., Gerard Ryan, D., Young, T., Zecchini, V.R.,
Rossi, S.H., Massie, C., Lohoff, C., Masid, M., Hatzimanikatis, V., Kuppe, C., Kriegsheim, A.V.,
Kramann, R., Gnanapragasam, V., Warren, A.Y., Stewart, G.D., Erez, A., Vanharanta, S., Saez-
Rodriguez, J., Frezza, C. (2022). Dynamic partitioning of branched-chain amino acids-derived
nitrogen supports renal cancer progression. Nature Communications *13* (*7830*).
10.1038/s41467-022-35036-4.

837
Sivanand, S., Gultekin, Y., Winter, P.S., Vermeulen, S.Y., Tchourine, K.M., Abbott, K.L., Danai,
L.V., Gourge, F., Do, B.T., Crowder, K., Kunchok, T., Lau, A.N., Darnell, A.M., Jefferson, A.,
Morita, S., Duda, D.G., Aguirre, A.J., Wolpin, B.M., Henning, N., Spanoudaki, V., Maiorino, L.,
Irvine, D.J., Yilmaz, O.H., Lewis, C.A., Vitkup, D., Shalek, A.K., Vander Heiden, M.G. (2024).
Cancer tissue of origin constrains the growth and metabolism of metastases. Nature Metabolism
6, 1668-1681. 10.1038/s42255-024-01105-9.
844
845 Srere, P.A. (1974). Controls of citrate synthase activity. Life Sciences *15*, 1695-1710.
846
847 Sullivan, L.B., Gui, D.Y., Hosios, A.M., Bush, L.N., Freinkman, E., Vander Heiden M.G. (2015).
848 Supporting Aspartate Biosynthesis Is an Essential Function of Respiration in Proliferation Cells.
849 Cell *162*, 552-563. doi:10.1016/j.cell.2015.07.017.
850
851 Sullivan, M.R., Danai, L. V., Lewis, C. A., Chan, S. H., Gui, D. Y., Kunchok, T., Dennstedt, E. A.,
852 Vander Heiden, M. G., Muir, A. (2019a). Quantification of microenvironmental metabolites in
853 murine cancers reveals determinants of tumor nutrient availability. eLife *8*.
854 https://doi.org/10.7554/elife.44235.
855
856 Sullivan, M.R., Mattaini, K.R., Dennstedt, E.A., Nguyen, A.A., Reilly, M.F., Meeth, K., Muir, A.,
857 Darnell, A.M., Bosenberg, M.W., Lewis, C.A., Vander Heiden, M.G. (2019b). Increased serine
858 synthesis provides an advantage for tumors arising in tissues where serine levels are limiting.
859 Cell Metabolism *29*, 1410-1421,e1414. doi: 10.1016/j.cmet.2019.02.015.
860
861 Terada, H. (1990). Uncouplers of Oxidative Phosphorylation. Environmental Health
862 Perspectives *87*, 213-218. 10.1289/ehp.9087213.
863
864 Vander Heiden, M.G., Chandel, N.S., Schumacker, P.T., Thompson, C.B. (1999). Bcl-xL
865 Prevents Cell Death following Growth Factor Withdrawal by Facilitating Mitochondrial ATP/ADP
866 Exchange. Molecular Cell *3*, 159-167. https://doi.org/10.1016/S1097-2765(00)80307-X.
867
868 Vander Heiden, M.G., Deberardinis R.J. (2017). Understanding the Intersections between
869 Metabolism and Cancer Biology. Cell *168*, 657-669. doi:10.1016/j.cell.2016.12.039.
870
871 Vichai, V., Kirtikara, K. (2006). Sulforhodamine B colorimetric assay for cytotoxicity screening.
872 Nature Protocols *1*, 1112-1116. https://doi-org.ezp-
873 prod1.hul.harvard.edu/10.1038/nprot.2006.179.
874
875 Wang, Y., Stancliffe, E., Fowle-Grider, R., Wang, R., Wang, C., Schwaiger-Haber, M., Shriver,
876 L.P., Patti, G.J. (2022). Saturation of the mitochondrial NADH shuttles drives aerobic glycolysis
877 in proliferating cells. Molecular Cell *82*, 3270-3283. https://doi.org/10.1016/j.molcel.2022.07.007.
878
Wu, Z., Bezwada, D., Cai, F., Harris, R.C., Ko, B., Sondhi, V., Pan, C., Vu, H.S., Nguyen, P.T.,
Faubert, B., Cai, L., Chen, H., Martin-Sandoval, M., Do, D., Gu, W., Zhang, Y., Zhang, Y.,
Brooks, B., Kelekar, S., Zacharias, L.G., Oaxaca, K.C., Patricio, J.S., Mathews, T.P., Garcia-
Bermudez, J., Ni, M., DeBerardinis, R.J., (2024). Electron transport chain inhibition increases
cellular dependence on purine transport and salvage. Cell Metabolism *36*, 1504-1520.
https://doi.org/10.1016/j.cmet.2024.05.014.
885

Yao, C., Wang, R., Wang, Y., Kung, C., Weber, J.D., Patti, G.J. (2019). Mitochondrial fusion
supports increased oxidative phosphorylation during cell proliferation. eLife *8*.
10.7554/eLife.41351.

Zhao, R., Jiang, S., Zhang, L., Yu, Z. (2019). Mitochondrial electron transport chain, ROS
generation, and uncoupling International Journal of Molecular Medicine *44*, 3-15.
10.3892/ijmm.2019.4188.

Figure Legends

Figure 1. The NAD+/NADH ratio is proportional to serine synthesis rate. (A) α-ketobutyrate

(AKB) is an exogenous electron acceptor that promotes the oxidation of NADH into NAD+ through

its conversion to α-hydroxybutyrate (AHB). Rotenone inhibits activity of complex I of the

mitochondrial electron transport chain (ETC) to block NADH oxidation into NAD+. **(B)**

NAD+/NADH ratio measured in A549 cells cultured without serine and indicated concentrations

of AKB (left) and rotenone (right) for 24 hours. NAD+/NADH ratios are normalized to untreated

cells cultured without serine, n=3. **(C)** Schematic depicting NAD+ requirement to produce serine

labeled (M+3) from U-^{13}C-glucose. Under culture conditions without serine, intracellular serine

rapidly effluxes from cells into culture media. Thus, cells and media were jointly analyzed at each

time point of kinetic tracing to better capture serine levels. **(D)** Serine labeled (M+3) from U-^{13}C-

glucose over time in A549 cells cultured without serine and with indicated concentrations of AKB

and rotenone for 24 hours prior to U-^{13}C-glucose exposure. Serine levels are normalized to

internal norvaline standard and cell number measured in indicated conditions, n=3. **(E)** Relative

NAD+/NADH ratios from A549 cells cultured without serine after exposure to 250 µM AKB, 1000

µM AKB, 20 nM rotenone, or 40 nM rotenone for 24 hours are plotted against the serine synthesis

rates from corresponding conditions as shown in D. Serine synthesis rates are calculated from

labeled serine (M+3) after 1 and 15 minutes of U-^{13}C-glucose exposure and normalized to

untreated and serine-free culture conditions, n=3. **(F)** Proliferation rate (doublings per day) of

A549 cells cultured with and without serine and the indicated concentration of AKB for 72 hours,

n=3. **(G)** Proliferation rate (doublings per day) of A549 cells cultured with and without serine and

the indicated concentration of rotenone for 72 hours, n=3. Data shown represent means ± SD.

Figure 2. The NAD+/NADH ratio differs between cancer cells upon serine withdrawal and

correlates with proliferation in serine-depleted conditions. (A) Proliferation rate (doublings

per day) of indicated cancer cells cultured without serine for 72 hours normalized to proliferation

rate of corresponding cancer cells cultured with serine, n=3. **(B)** NAD+/NADH ratios of indicated

cells cultured with and without serine for 24 hours. Yellow indicates cells with unaltered

NAD+/NADH ratios and blue indicates cells with elevated NAD+/NADH ratios when cultured

without serine. Values are normalized to the NAD+/NADH ratios of cells cultured with serine, n=3.

(C) Relative NAD+/NADH ratios of cells cultured without serine as shown in B are plotted against

proliferation rates of corresponding cells cultured without serine. Data points denote the different

cell lines (Calu6, A549, 8988T, MIA PaCa-2, H1299, and HCT116). **(D)** Serine labeled (M+3) from

U-^{13}C-glucose over time in A549 and H1299 cells cultured with and without serine for 24 hours

prior to U-^{13}C-glucose exposure. Serine levels are normalized to internal norvaline standard and

cell number measured in indicated conditions, n=3. **(E)** Absolute fold change of serine labeled

(M+3) from U-^{13}C-glucose in A549 and H1299 cells cultured without serine relative to serine

labeled (M+3) from U-^{13}C-glucose in A549 and H1299 cells cultured with serine for 24 hours after

twelve minutes of U-^{13}C-glucose exposure, n=3. Data shown are means ± SD. P-values were

calculated by unpaired Student's t-test, **p<0.01, ***p<0.001

Figure 3. Elevated mitochondrial respiration is associated with increased NAD+/NADH

ratios and serine synthesis rates. (A) Oxygen consumption rate (OCR) of indicated cells

cultured with and without serine for 24 hours. Values are averages of three repeat measurements,

n=14-20. **(B)** OCR of cells (H1299 top, A549 bottom) cultured with indicated concentrations of

serine for 24 hours. Values are averages of three repeat measurements, n=3-9. **(C)** NAD+/NADH

ratios of cells (H1299 top, A549 bottom) cultured with indicated concentrations of serine for 24

hours. Values are normalized to the NAD+/NADH ratio of cells cultured in 400μM serine (serine

concentration in DMEM), n=3. **(D)** Relative OCR plotted against NAD+/NADH ratios in H1299 and

A549 cells cultured in 0, 25, 50, and 100 μM serine for 24 hours. OCR and NAD+/NADH ratios

are normalized to measurements from the 100 μM serine condition, n=3. **(E)** Serine labeled (M+3)

from U-^{13}C-glucose over time in cells (H1299 top, A549 bottom) cultured with indicated serine

concentrations for 24 hours prior to U-^{13}C-glucose exposure. Serine levels are normalized to

internal norvaline standard and cell number measured in indicated conditions, n=3. **(F)** Relative

NAD+/NADH ratios of cells (H1299 top, A549 bottom) cultured with 0, 25, 50, and 100 μM serine

for 24 hours plotted against relative serine synthesis rates of cells cultured in corresponding

conditions. Serine synthesis rates are calculated from labeled serine (M+3) after 1 and 15 minutes

of U-^{13}C-glucose exposure and normalized to culture conditions with serine, n=3. Data shown are

means ± SD. P-values were calculated by unpaired Student's t-test, *$p<0.05$, **$p<0.01$,

***$p<0.001$ ****$p<0.0001$, "ns" denotes not significant.

Figure 4. Mitochondrial respiration governs the cell NAD+/NADH ratio and influences

serine synthesis. (A) Schematic depicting the mitochondrial electron transport chain (ETC) and

the protonophore activity of FCCP that uncouples the ETC from ATP synthase activity. IMS,

intermembrane space; CoQ(H$_2$) indicates the oxidized and reduced (H$_2$) forms of coenzyme Q;

cyt$_c$, cytochrome c. **(B)** Proliferation rate (doublings per day) of A549 cells cultured with and

without serine with indicated concentrations of FCCP for 72 hours, n=3. **(C)** Proliferation rate

(doublings per day) of H1299 cells cultured with and without serine with indicated concentrations

of FCCP for 72 hours, n=3. **(D)** Serine labeled (M+3) from U-^{13}C-glucose over time in A549 cells

cultured without serine, and with and without 2 μM FCCP treatment for 24 hours prior to U-^{13}C-

glucose exposure. Serine levels are normalized to internal norvaline standard and cell number

measured in indicated conditions, n=3. **(E)** Serine labeled (M+3) from U-^{13}C-glucose over time in

H1299 cells cultured without serine, and with and without 2 µM FCCP treatment for 24 hours prior

to U-^{13}C-glucose exposure. Serine levels are normalized to internal norvaline standard and cell

number measured in indicated conditions, n=3. **(F)** Oxygen consumption rate (OCR) of A549 cells

cultured with and without serine with indicated FCCP treatment for 24 hours. Values are averages

of three repeat measurements, n=5-9. **(G)** Relative NAD+/NADH ratio of A549 cells cultured with

and without serine with indicated FCCP treatment for 24 hours. NAD+/NADH ratios are

normalized to the NAD+/NADH ratio in A549 cells culture with serine, n=3. **(H)** Proliferation rate

(doublings per day) of A549 cells cultured with and without serine, 2 µM FCCP, and indicated

concentrations of rotenone for 72 hours, n=3. **(I)** Serine labeled (M+3) from U-^{13}C-glucose over

time in A549 cells cultured without serine, and with and without 2 µM FCCP and 20 nM rotenone

as indicated, for 24 hours prior to U-^{13}C-glucose exposure. Serine levels are normalized to internal

norvaline standard and cell number measured in indicated conditions, n=3. **(J)** Relative

NAD+/NADH ratio of A549 cells cultured with and without serine, 2 µM FCCP, and 20 nM

rotenone as indicated. NAD+/NADH ratios are normalized to the NAD+/NADH ratio in untreated

A549 cells cultured with serine, n=3. Data shown are means ± SD. P-values were calculated by

unpaired Student's t-test, *p<0.05, **p<0.01, ***p<0.005, ****p<0.001, "ns" denotes not significant.

Figure 5. Lipid depletion leads to a cell-specific elevation in mitochondrial respiration

and the NAD+/NADH ratio, influencing oxidative citrate and serine synthesis. (A)

Schematic depicting glucose and glutamine oxidative routes for synthesizing citrate, highlighting

NAD+ requiring oxidation reactions. **(B)** Citrate labeled (M+2) from U-^{13}C-glucose over time in

H1299 and A549 cells cultured with and without serine for 24 hours prior to U-^{13}C-glucose

exposure. Citrate levels are normalized to internal norvaline standard and cell number

measured in indicated conditions, n=3. **(C)** Citrate labeled (M+2) from U-^{13}C-glucose over time

in A549 and H1299 cells cultured without serine, and with and without 2 µM FCCP for 24 hours

prior to U-^{13}C-glucose exposure. Citrate levels are normalized to internal norvaline standard and

cell number measured in indicated conditions, n=3. **(D)** Citrate labeled (M+2) from U-^{13}C-

glucose over time in A549 cells cultured with serine, and with indicated concentrations of AKB

or FCCP for 24 hours prior to U-^{13}C-glucose exposure. Citrate levels are normalized to internal

norvaline standard and cell number measured in indicated conditions, n=3. **(E)** Oxygen

consumption rate (OCR) in A549 and H1299 cells cultured with and without lipids for 24 hours.

Values are averages of three repeat measurements, n=10. **(F)** Relative NAD+/NADH ratios of

A549 and H1299 cells cultured with and without lipids for 24 hours. NAD+/NADH ratios are

normalized to the NAD+/NADH ratio of the corresponding cell type cultured with lipids, n=3. **(G)**

Citrate labeled (M+2) from U-^{13}C-glucose over time in A549 and H1299 cells cultured with and

without lipids for 24 hours prior to U-^{13}C-glucose exposure. Citrate levels are normalized to

internal norvaline standard and cell number measured in indicated conditions, n=3. **(H)** Serine

labeled (M+3) from U-^{13}C-glucose over time in A549 and H1299 cells cultured with and without

lipids for 24 hours prior to U-^{13}C-glucose exposure. Serine levels are normalized to internal

norvaline standard and cell number measured in indicated conditions, n=3. Data shown are

means ± SD. P values were calculated by unpaired Student's t-test, *p<0.05.

1009

Figure 6. Lipid depletion increases mitochondrial respiration and the NAD+/NADH ratio in

a cell specific manner to influence serine synthesis and proliferation in serine-depleted

conditions. (A) Oxygen consumption rate (OCR) of A549 and H1299 cells cultured with and

without serine or lipids for 24 hours. Values are the average of three repeat measurements, n=14-

15. **(B)** Relative NAD+/NADH ratios of A549 and H1299 cells culture with and without serine or

lipids for 24 hours. NAD+/NADH ratio is normalized to the NAD+/NADH ratio in corresponding

cell type cultured with serine and lipids, n=3. **(C, D)** Left: Serine labeled (M+3) from U-^{13}C-glucose

over time in A549 cells (C) and H1299 cells (D) cultured without serine, and with or without lipids

for 24 hours prior to U-^{13}C-glucose exposure. Right: Serine synthesis rates of A549 cells (C) and

H1299 cells (D) cultured in conditions shown on left. Serine synthesis rates are calculated from

labeled serine (M+3) after 1 and 15 minutes of U-^{13}C-glucose exposure and normalized to the

serine synthesis rates of the corresponding cell type cultured without serine and with lipids. Serine

levels are normalized to internal norvaline standard and cell number measured in indicated

conditions, n=3. Single data point (H1299, -serine +lipid, t=15 min) removed due to error in sample

processing. (**E,F**) Left: Citrate labeled (M+2) from U-^{13}C-glucose over time in A549 cells (E) and

H1299 cells (F) cultured without lipids, and with or without serine for 24 hours prior to U-^{13}C-

glucose exposure. Right: Citrate synthesis rates of A549 cells (E) and H1299 cells (F) cultured in

conditions shown on left. Citrate synthesis rates are calculated from labeled citrate (M+2) after 1

and 15 minutes of U-^{13}C-glucose exposure and normalized to the citrate synthesis rates of the

corresponding cell type cultured without lipids and with serine. Citrate levels are normalized to

internal norvaline standard and cell number measured in indicated conditions, n=3. (**G**)

Proliferation rate (doublings per day) of A549 cells cultured with and without serine or lipids for 72

hours, n=3. (**H**) Proliferation rate (doublings per day) of H1299 cells cultured with and without

serine or lipids for 72 hours, n=3. Data shown are means ± SD. P-values were calculated by

unpaired Student's t-test, *p<0.05 ***p<0.005. ****p<0.001.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6