Phylogenetic relationships of *ATM* polymorphic loci haplotypes in Eurasian and African

populations

Marina V. Olkova[1,*], Sergey M. Koshel[2], Georgiy Yu. Ponomarev[1], Andrei A. Alimov[1]

[1] Research Centre for Medical Genetics, Moscow, 115478, Russia

[2] Department of Cartography and Geoinformatics, Lomonosov Moscow State University,

Moscow, 119991, Russia

Corresponding author email: genetics@inbox.ru

ABSTRACT

The *ATM* gene encodes a key kinase involved in DNA repair and cell cycle control, and

mutations in it lead to the development of ataxia-telangiectasia and increase susceptibility to

cancer. In this context, the study of the distribution patterns of *ATM* haplotypes in

metapopulations living in Eurasia and Africa using population genetic and genogeographic

methods appears to be a socially important scientific endeavor.

Polymorphic markers of the *ATM* gene were studied in 3876 individuals from 46

metapopulations, *ATM* gene haplotypes and genotypes based on *ATM* haplotypes were

determined, and population frequencies of haplotypes and genotypes were calculated. An

analysis of phylogenetic relationships of haplotypes, the Tajima test to assess population

changes, and mapping of population frequencies of *ATM* gene haplotypes and genotypes were

performed. The study showed the presence of a linkage block of 28 polymorphic markers of the

ATM gene in the Eurasian and African populations. Two dominant *ATM* haplotypes in Eurasia,

which are the result of–independent evolution of two branches from the ancestral *ATM*

haplotype, are formed by mutually exclusive alleles and correspond to the "yin-yang" concept.

The greatest diversity of *ATM* haplotypes is observed on the African continent. It is suggested

that the process of haplotype formation occurred on the African continent, with subsequent

migration events to the territory of Eurasia.

Key words: *ATM* haplotypes, LD-block, yin-yang haplotypes, mutually exclusive alleles,

evolutionary separation

INTRODUCTION

The human *ATM* gene (ataxia-telangiectasia mutated), which encodes a serine-threonine

protein kinase, is located on chromosome 11, locus 11q22-23. Activation of the gene's protein

product occurs as a result of double-stranded DNA breaks, leading to the initiation of signaling pathways and the phosphorylation of numerous substrates (several hundred have been identified to date). The final result of cascade reactions is cell cycle arrest and DNA repair. In addition to double-stranded DNA breaks, activation of the *ATM* protein can be caused by oxidative stress, deficiency of a number of metabolites, hypoxia, and some other factors [1]. *ATM* is also involved in the activation of the spindle checkpoint during mitosis. Loss of *ATM* function results in a defective spindle checkpoint and increases the likelihood of aneuploidy [2].

Mutations in the *ATM* gene are associated with the development of autosomal recessive Louis-Barr syndrome (ataxia-telangiectasia) and with the risk of developing cancer, which is one of the reasons for the research interest in studying the distribution of frequencies of polymorphic variants of this gene in populations [3–6]. One of the most important stages of genetic data research is the linkage disequilibrium (LD) analysis, which allows to identify new clinically significant variants of the *ATM* gene associated with changes in the function of the gene's protein product and disease severity. As a result of previous scientific studies, blocks of LD in the *ATM* gene have been identified and described in a region of ~150 kb, covering the entire length of the gene [7–9]. In particular, Tornston et al. identified seven haplotypes of the *ATM* gene, two of which were present in 82% of the 93 DNA samples examined from representatives of seven human populations living on all continents. Two of the seven identified haplotypes were found only in African populations with a frequency of 33%. The presence of blocks of LD of varying length was also demonstrated for several other autosomal regions of the human genome. In particular, Bonnen's article assessed the LD of non-coding polymorphic markers for 5 genes (TP53, RAG51, BRCA1, BRCA2 and *ATM*) associated with the cancer development. It was shown that for all genes, since the study included markers evenly distributed along the gene, there was no tendency for the LD value to decrease with increasing distance. The authors

concluded that the LD value can vary widely regardless of the physical distribution of the

markers along the length of the DNA sequence of the gene. The observed phenomenon testified

to the existence of polymorphic loci in the genome, linked in blocks, with the degree of linkage

independent of distance. A similar conclusion was reached by Gabriel et al., who reported the

presence of SNP blocks with a small number of common haplotypes, evenly distributed across

the human genome. The study was conducted using DNA samples from 275 representatives of

the American population of African, European and Asian origin. A link was found between the

origins of the populations and specific haplotypes. Haplotypes with mutually exclusive alleles

of polymorphic markers, called "yin" and "yang", were also identified [10–12].

Our study aims to perform a population analysis of combined into haplotypes SNP variants of

the *ATM* gene, associated with oncological pathology. In addition, a genogeographic

comparison of the North Eurasian gene pool of *ATM* haplotypes with the gene pools of the

peoples of Eurasia and Africa was conducted. Phylogenetic analysis based on all identified

ATM haplotypes was performed to better understand the structural changes accumulated in the

gene during the evolutionary process.

MATERIALS AND METHODS

Ethical approvals

This study was conducted under the condition that written informed consent was obtained from

all subjects. This consent was approved by the Ethics Committee of the Research Center for

Medical Genetics.

83
Collection of DNA genotype data and populations

1975 DNA samples from individuals representing 27 populations living in Russia and adjacent

countries were provided by the Northern Eurasia Biobank. Population sample sizes ranged

from 41 to 129 individuals. Individual allelic variants of 269 polymorphic markers of the *ATM*

gene were extracted from a genome-wide genotyping dataset using the Infinium Omni5Exome

BeadChip Kit (Illumina).

Genotype information for 1901 individuals from 19 populations in Eurasia and Africa, with

population sample sizes ranging from 85 to 113 individuals, was obtained from the 1000

Genomes Project Phase 3 data (Ensembl).

The general list of populations, their abbreviated names and the size of population samples are

presented in Table 1.

The "ancestral" haplotype was determined based on the "ancestral" alleles of the polymorphic

markers specified in the Ensembl project, identified based on a three-stage multiple alignment

of EPO (Enredo, Pecan, Ortheus).

Statistical data processing

Microsoft Excel (Microsoft Corporation) and RStudio 2024.12.1+563 - R version 4.4.2

(RStudio, PBC) were used for statistical data processing.

DNA sample genotyping and polymorphic ATM loci quality assessment

Polymorphic markers and samples with genotype quality below 95% were excluded from the

study (using the *snpStats* R-package).

Isolation of haplotypes. Calculation of population frequencies of *ATM* haplotypes and

genotypes

The isolation of haplotypes and the calculation of their population frequencies were performed

in the Haploview 4.1 program (Broad Institute).

The "confidence interval method" (Gabriel) was used to isolate blocks of polymorphic variants

forming a haplotype. A haplotype block was defined if 95% of SNP pairs within it had D > 0.8.

Calculation of population frequencies of genotypes based on *ATM* haplotypes was performed

using the *plyr* R-package.

Phylogenetic analysis

117   Phylogenetic networks of *ATM* haplotype representation in populations were constructed using

118   the *pegas* and *geneHapR* R-packages according to the "parsimony principle". The preparatory

119   files were filled with data on population frequencies of haplotypes, multiplied by 1000.

Calculation of Tajima test for populations and individual "yin" and "yang" branches

122   The Tajima test was calculated based on the population haplotype frequency data obtained in

123   Haploview 4.1 software. The population number of haplotypes of each type was obtained by

124   multiplying the haplotype population frequency by 1000. The Tajima test D-value and p-value

125   were calculated using the *pegas* R-package.

Cartographic analysis

Cartographic visualization of the population frequencies of haplotypes and genotypes of the

ATM gene was performed using the original GeneGeo cartographic package [13]. Frequency

calculations for the entire mapped area were performed using the weighted average

interpolation method with a weighting function inversely proportional to the cube of the

distance and an influence radius of 4000 km. To avoid the influence of map projection

distortions calculations, all distances in the interpolation process were calculated on a sphere.

The Winkel triple projection with the central meridian 60^{o} E was used to create the maps.

137 RESULTS

138 Determination of *ATM* gene haplotypes and their population frequencies

The genotype data of the combined population of Northern Eurasia allowed the identification of

a LD block of 28 linked polymorphic loci using Haploview 4.1 software – rs189037, rs228590,

rs623860, rs600931, rs228592, rs672655, rs627418, rs664677, rs609261, rs645485, rs619972,

rs599558, rs595747, rs609655, rs227061, rs227062, rs227064, rs227068, rs227070, rs227074,

rs227075, rs425538, rs419716, rs227040, rs664143, rs227094, rs227092 and rs4585, located

over 145,795 bp from the 5′-untranslated region to the 3′-untranslated region of the *ATM* gene

(Figure S1).

The value of the square correlation coefficient (r^2) between all pairs of loci in the LD block

exceeded 0.8. Of 268,435,456 possible haplotypes for 28 polymorphic loci of the LD block in

the combined population of peoples living in the territory of Northern Eurasia, 4 haplotypes

were found, and 2 of them were predominant: G-A-C-C-C-A-T-C-T-A-C-C-C-G-A-G-T-A-T-

A-T-C-A-T-A-C-G-G and A-G-T-T-A-G-C-T-C-G-T-T-T-A-G-A-C-G-G-G-C-A-C-C-G-A-T-

T with a frequency of 39.2% and 54.5%, respectively. The alleles in the two most frequent

haplotypes were mutually exclusive, which allows us to classify them as "yin"/"yang"

haplotypes. The remaining two detected haplotypes were much less frequent in the population

(with a frequency of about 2%) and were the result of recombination between the "yin" and

"yang" haplotypes.

Similarly, haplotypes were determined, and their population frequencies were calculated for 28

polymorphic markers of the *ATM* haplotype block in each of the 27 populations of Northern

Eurasia and in 19 populations of Eurasia and Africa using data from the 1000 Genomes Project

Phase 3 (Ensembl). From the total set of haplotypes detected in all the above Eurasian and

African populations, rare (with a population frequency of up to 5%) found in single

populations, as well as haplotypes that most likely arose as a result of recombination between

the "yin-yang" haplotypes, were excluded. As a result, a list of 11 haplotypes of the *ATM* gene

(Figure 1) was determined, occurring in 46 populations of Eurasia and Africa. The "ancestral"

haplotype (H0) of the human *ATM* gene was determined using the Ensembl genome browser,

which indicates the "ancestral" alleles for each of the 28 linked loci of this gene.

ATM **haplotypes-based phylogenetic analysis**

Based on the isolated haplotypes, a phylogenetic network was constructed reflecting the

evolutionary variability of the *ATM* gene haplotype block in the human population and the

frequency ratio of *ATM* haplotypes in modern peoples of Eurasia and Africa (Figure 2).

The presence of two independently evolving phylogenetic branches, which we called I and J

branches, was discovered, leading from the "ancestral" haplotype to the most common in

Eurasia H6i and H5j ("yin" and "yang") haplotypes (Figure 3). All haplotypes analyzed were

found in African populations (Figure 4, S2a).

In addition to the main "yin/yang" haplotypes, the H2i and H4i (branch I), H3j (branch J)

haplotypes were also present in the Eurasia area. Of these three haplotypes, only H2i was found

in populations of European origin (Figure S2b), in contrast to populations of Asian origin

(Figure S2c). All three haplotypes are rare (up to 2%) in Eurasian populations. Based on these

data, it can be concluded that the formation of the ancestral haplotype and the subsequent

divergent evolution of two isolated branches probably occurred on the territory of the African

continent.

Tajima test for populations and individual branches of "yin" and "yang" haplotypes

For all populations of Eurasia and Africa, both divided by origin and geographical location, and

combined by these criteria, a positive value of the D parameter of the Tajima test (from 4.7 to

9.3) with a p value << 0.05 was observed, which may be due to a long-term subdivision of the

population into several subgroups with limited gene flow between them, balancing selection, a

sharp reduction in population size, or other genetic and demographic processes.

For each of these populations, a significant deviation from the neutral model of evolution was

observed. At the same time, when considering each of the two evolutionary branches leading

from the "ancestral" haplotype to the H6i and H5j haplotypes separately, this test showed a

statistically insignificant result for the D parameter and p-value ($p \gg 0.05$), which indicating

that both divergence paths (I and J) of the "ancestral" *ATM* haplotype conform to the neutral

model of evolution (Table S1).

Geographical analysis of population frequencies of *ATM* haplotypes and genotypes

The frequencies of *ATM* haplotypes, as well as their homozygous and heterozygous

combinations, are presented in Tables 2-3 and on maps (Fig. S3-S6).

<u>"Yin" and "yang" haplotypes</u>

Cartographic analysis of the prevalence of population frequencies of haplotypes H6i and H5j,

as well as homozygous carriers of these haplotypes showed similar results.

Haplotype H5j was in southern Eurasia with maxima in the southern of Western Europe (~

69%), in Transcaucasia (~72%) and in the north-west of South Asia (~70%). There was also a

significant increase in the frequency of this haplotype in the populations of the indigenous

populations of southern Siberia (with a maximum among Tuvans and Tofalars ~66%) and

Kamchatka (Chukchi, Koryaks, Itelmens ~59%), with a significant decrease in East Asia to a

frequency of ~36%. The H5j haplotype is the second most common in Africa, although its

population frequencies are noticeable lower than in Eurasia - from 18% in most African

populations to 24% in the Mende people living on the coast of West Africa.

The H6i haplotype has two geographic maxima - on the east coast of Asia (~56% in the Dai

people in southwest China) and in eastern Europe (~49% in the Bashkirs). Between the

maxima, there is a marked decrease in population frequencies in central Siberia, extending to

the northern coast of Eurasia and the south of the continent - along the northern border of the

Alpide belt. The lowest population frequency of the H6i haplotype in Eurasia is in Eastern

Transcaucasia (~21%). The H6i haplotype is the fourth most common in African populations –

from 4% in the Mende of Sierra Leone to 12% in the Yoruba of Nigeria.

In summary, it can be said that in the Eurasian area, apart from East Asia, the H5j haplotype

predominates between the two mutually exclusive and most common haplotypes. In the

populations of East Asia, the H6i haplotype predominates. In Africa, the population frequencies

of the "yin" and "yang" haplotypes are significantly lower than in Eurasian populations. On

average, the H5j haplotype is 2.8 times less common in African populations than in Eurasian

populations, and the H6i haplotype is 4.6 times less common.

<u>Heterozygous combination of "yin" and "yang" haplotypes</u>

The heterozygous genotype H6i/H5j turned out was one of the most common in Eurasian

populations (from 16.4% to 52.5%). The exception was 9 out of 42 Eurasian populations (Table

5). The prevalence of homozygotes for the H5j haplotype was found in 2 populations of

northwestern South Asia (41.7% in Gujaratis of India and 51% in Punjabis of Pakistan), in

populations of Transcaucasia (42-44.6%), in the Italian population of southern Europe (51.4%),

as well as in 3 metapopulations of Northern Eurasia - "Chukchi, Koryaks, Itelmens" (34.5%),

"Mari, Chuvash" (35.3%) and "Tuvans, Tofalars" (41.3%). The homozygous genotype for the

H6i haplotype turned out was the most common in only one population - the East Asian Han,

where its frequency was ~ 34%. In Africa, the frequency of H6i/H5j heterozygotes was an order

of magnitude lower than in Eurasian (on average ~3.5%) than in Eurasia (~37%). Meanwhile,

among the Yoruba population, the genotype H6i/H5j was the most frequent (6.5%), despite the

complete absence of homozygotes for haplotype H6i and a very low population frequency of

homozygotes for haplotype H5j (~1%). This may be related to the manifestation of balancing

selection or migration processes.

The peaks of the H6i/H5j genotype in Eurasia are located on the coasts: in the east of the

Indochinese peninsula (~52.5%), in the Spanish (~46.7%) and British (~47.3%) populations of

Western Europe, and in the Arkhangelsk region (~51.4%) of Northern Europe. There is also a

significant increase in heterozygote frequencies in the Eastern European Tatar population

(48.1%).

The Eurasian minimum of population frequencies of "yin-yang" heterozygotes is observed

among the "Chukchi, Koryaks and Itelmens" (~16.4%).

<u>Haplotype H3j</u>

Haplotype H3j differs from H5j at 3 positions - rs189037, rs672655, rs619972. With a

frequency of 1% to 7%, H3j is found in most Eurasian populations of Asian origin (except for

the West Siberian Plain and Vietnam) but has not been found in European populations.

Homozygotes for H3j have not been found, apparently because of the low frequencies of the

haplotype.

In Africa, this haplotype ranks 5th in population frequency ranging - from 4.5% in the Luhya

population of Kenya to 11% in the Yoruba of Nigeria.

<u>Geography of population frequencies of heterozygotes for the haplotype H3j in combination</u>

<u>with H6i, H5j, H4i and H2i</u>

The geographical distribution of the H3j haplotype in combination with H6i and H5j generally

corresponds to that of the H3j haplotype in populations. The H3j/H6i genotypes are more

common along the line running from southern Asia to northern Eurasia, with the highest frequencies among Sri Lankan Tamils (7.8%) and Bashkirs (7.1%). The H3j/H5j genotype is observed with the highest frequency (10.5%) in the Bengal population of South Asia. The H3j/H4i and H3j/H2i genotypes are found only in Africa, the former with a frequency of 1-2% and the latter with a frequency of 2-6%.

Haplotype H2i

 The most common in Africa, H2i (ranging from 19% to 31%), was found at very low levels (~2%) in some populations of East Asia (southern China and Vietnam) and Europe (UK, Western Caucasus, south-eastern central Russia, Mordovia and Tatarstan).

Prevalence of heterozygous carriers of the haplotype H2i in combination with H6i, H5j and H4i

The geographical distribution of heterozygotes for mixed haplotypes of the most common African H2i and the predominant Eurasian H6i and H5j is quite revealing.

The population frequencies of the H2i/H6i genotype are widespread, although at low frequencies, not only in African populations, where its occurrence is 2-5%, but also in some Eurasian populations, where the frequency of this genotype is 1-2%.

Heterozygous carriers of the H2i/H5j haplotypes are more common in Africa than heterozygotes for H2i/H6i, the range of population frequency for H2i/H5j is from 5-7%, and in the Mandinka population of Gambia this genotype is the most common.

Carriers of the H2i/H6i genotype have also been found in all populations of East Asia, Western Europe and some European populations of Northern Eurasia with a population frequency of 1-2%.

The H2i/H4i genotype is found in African populations with a frequency of 3-5%, with only the

Mandinka having a frequency of about 2%. This genotype was also found in the southern East

Asia and in the British and Italian populations of Europe with a frequency of 1-2%.

<u>Population characteristics of ATM haplotypes in Africa</u>

Haplotypes H1j, H1i, H2j, H3i, H4j, and H5i are exclusively characteristic of the African

continent. H4i is third in frequency in Africa (from 9% among Yoruba to 13.5% among

Mandinka and Mende) but has been detected outside Africa only in the Bashkir population at a

frequency of 1%. This observation likely reflects the heterogeneity of this population, which

was formed through the contribution of various ethnic groups.

The haplotype H1j occurs in African populations with frequencies ranging from 3% to 9% (on

the western coast) and differs from the 'ancestral' one only at a single position (G in rs645485).

H1j was found in the homozygous state exclusively among the Mende from Sierra Leone,

where the frequency of homozygous carriers reached ~3.5%. H1i is most prevalent on both the

eastern and western African coasts (7-8%) and almost completely absent in the central part of

the continent (0-1%). Homozygous carriers of H1i were detected with a population frequency

of ~1% among Mandinka from Gambia and Luhya from Kenya.

H2j was found at a frequency of 1% in Luhya and Mende populations. H3i is rarest in the

Mende population (~1%), while its population frequency increases to 5-6% in Luhya and

Mandinka populations on the eastern and western coasts.

H4j is most prevalent among Mandinka (~7%) in western Africa, with a decreased presence of

~1.5% observed in Luhya populations in the east. In contrast, H5i reaches its highest frequency

on the eastern coast (~10%), while in other African populations, its population frequencies

range from ~4-5%.

Overall, it can be concluded that the African continent exhibits maximum diversity among

ATM haplotypes and unique patterns of their population distribution.

DISCUSSION

The study of *ATM* haplotypes in human populations of Eurasia and Africa from the standpoint

of the "yin-yang" concept showed the presence in all populations of a conservative LD-block

consisting of 28 extra-exon loci of the *ATM* gene located throughout the DNA sequence of the

gene, including 5' and 3' untranslated regions of the *ATM* gene. Analysis of the phylogenetic

relationships between 11 identified *ATM* haplotypes revealed the presence of two independent

branches of *ATM* gene evolution from the ancestral haplotype, leading to the formation and

widespread distribution of haplotypes with mutually exclusive alleles according to the "yin-

yang" principle. In the samples analyzed outside the African continent, the dominant

haplotypes in all human populations were the H6i haplotype (from 21 to 56%) and the H5j

haplotype (from 32 to 72%), while the population frequency of the remaining haplotypes did

not exceed 7% in Eurasian populations. In the sample from the African continent, 11 haplotypes

were found, reflecting all stages of gene evolution from the "ancestral" haplotype. The most

common was the H2i haplotype, its median frequency in African populations was 23%. On the

territory of Eurasia, the H2i haplotype was found in only 7 populations, where its occurrence

did not exceed 2%, which may be the result of evolutionary and demographic processes.

Combining the obtained information on the evolutionary variability of *ATM* haplotypes with

data on their frequencies in modern populations of Africa and Eurasia, it can be assumed that

the process of formation of existing *ATM* haplotypes occurred on the African continent.

Phylogenetic analysis of *ATM* haplotypes demonstrates the existence of two isolated ancestral

human populations in the past, followed by their merging into a single mixed population on the

African continent (Figure 3). Similar conclusions were reached in the project [11], which was

conducted using data from the human genome of the 1000 Genomes Project, as well as the genomes of Denisovans and Neanderthals - Eastern and Western. Dutta et al. found an abundance of mutually exclusive pairs of common haplotypes for population-frequent (>25%) polymorphic markers in the human genome. Moreover, these "yin" and "yang" haplotypes had close population frequencies in Eurasia. At the same time, this study identified a category of haplotypes that are mostly observed in Africa. These haplotypes are composed of fragments of "yin" and "yang" haplotypes, hence the term "mosaic" used by the authors.

The results of the Tajima test in our study confirm the fact of the probable division of the original human population into several independent groups, and the statistically insignificant values of D when calculating the Tajima test for the haplotypes of the "yin" and "yang" branches suggest a neutral nature of evolution for each of the branches (Table S1).

The combined mixed population of carriers of *ATM* haplotypes, based on phylogenetic data, continued to exist on the African continent. Only a part of the *ATM* haplotypes from the "African" set were found on the territory of Eurasia. As a result of genetic drift and, possibly, balancing selection, the "yin" and "yang" haplotypes became dominant in Eurasia, while the others either disappeared completely or survived in small quantities in individual populations.

The peoples of Asia differ from those of European origin, firstly, by the widespread occurrence of the H3j haplotype (up to 7%) and, secondly, by the prevalence of the H6i haplotype in East Asia. It is important to note that in South Asia, the dominant haplotype among peoples of European origin, H5j, prevails, which allows us to consider the modal haplotype H5j in the population as a distinctive Indo-European feature. An analysis of literary data on the connection of the analyzed polymorphic loci with oncological diseases revealed one important contradiction. In populations of East Asian origin, the H5j haplotype, containing the rs189037, rs609261 and rs4585 loci, is associated with the manifestation of negative effects in oncopathology. In peoples of European origin, this association is not observed, which may be

due to the presence of an additional factor, yet unknown, which acts as an additive in East Asian populations. It is also worth paying attention to the prevalence of heterozygotes for the "yin" and "yang" haplotypes in most Eurasian populations. There are scientific data on the positive effect of heterozygous carrier references between the GA genotype and longevity, which the authors associate with a moderate level of *ATM* protein expression [14]. A study by Ding et al. showed that heterozygous genotype for rs189037 is associated with a reduced risk of type 2 diabetes in older adults [15].

CONCLUSION

Our study of evolutionary changes in the *ATM* gene haplotypes, represented by high-frequency polymorphic loci throughout the DNA sequence of this gene, indicates a very ancient, significant in time African period of isolation of two branches of the ancestral human population and the subsequent merging of these branches into a single population space on the African continent. Repeated migration events to the territory of Eurasia probably occurred after the unification of ancient isolated subpopulations. Some differences in the composition of *ATM* haplotypes in the east and west of the Eurasian continent, as well as in individual populations, are probably due to the effect of gene drift.

DECLARATION OF INTERESTS

Authors have no conflict of interest to declare.

ACKNOWLEDGMENTS

Authors express the gratitude to the "Biobank of North Eurasia" for providing DNA samples, and to E.V. Balanovskaya for her comments and suggestions that help improve the article. The research was carried out within the state assignment of MSHE of RF for RCMG.

AUTHOR CONTRIBUTIONS

Marina V. Olkova – article idea, bioinformatics analysis, primary genogeographic analysis, writing the article text

Sergey M. Koshel – genogeographic analysis

Georgiy Yu. Ponomarev – statistical and cartographic analysis

Andrei A. Alimov – research supervision, article editing

DATA AND CODE AVAILABILITY

The 1000 Genomes data can be obtained from the 1000 Genomes Project (Ensembl) Phase 3 data. The published article includes all data on Northern Eurasian populations analyzed during this study. R code is available at https://github.com/molkova/ATM-haplotypes/tree/main.

REFERENCES

1. Ueno S, Sudo T, Hirasawa A. *ATM*: Functions of *ATM* Kinase and Its Relevance to Hereditary Tumors. Int J Mol Sci. 2022;23(1):523. https://www.mdpi.com/1422-0067/23/1/523

2. Yang C, Tang X, Guo X, Niikura Y, Kitagawa K, Cui K, et al. Aurora-B Mediated ATM Serine 1403 Phosphorylation Is Required for Mitotic *ATM* Activation and the Spindle Checkpoint. Mol Cell. 2011;44(4):597–608. https://linkinghub.elsevier.com/retrieve/pii/S1097276511008148

3. Zhao Z liang, Xia L, Zhao C, Yao J. *ATM* rs189037 (G > A) polymorphism increased the risk of cancer: an updated meta-analysis. BMC Med Genet. 2019;20(1):28. https://bmcmedgenet.biomedcentral.com/articles/10.1186/s12881-019-0760-8

4. Tao Y, Mei Y, Ying R, Chen S, Wei Z. The *ATM* rs189037 G>A polymorphism is associated with the risk and prognosis of gastric cancer in Chinese individuals: A case–control study. Gene. 2020;741:144578. https://www.sciencedirect.com/science/article/pii/S037811192030247X

5. Zhou K, Bellenguez C, Spencer CCA, Bennett AJ, Coleman RL, Tavendale R, et al. Common variants near *ATM* are associated with glycemic response to metformin in type 2 diabetes. Nat Genet. 2011;43(2):117–20. https://doi.org/10.1038/ng.735

6. Lee SA, Lee KM, Lee SJ, Yoo KY, Park SK, Noh DY, et al. Antioxidant Vitamins Intake, Ataxia Telangiectasia Mutated (*ATM*) Genetic Polymorphisms, and Breast Cancer Risk. Nutr Cancer. 2010;62(8):1087–94. https://doi.org/10.1080/01635581.2010.492088

7. Bonnen PE, Story MD, Ashorn CL, Buchholz TA, Weil MM, Nelson DL. Haplotypes at *ATM* Identify Coding-Sequence Variation and Indicate a Region of Extensive Linkage

Disequilibrium. Am J Hum Genet. 2000;67(6):1437–51.

https://www.sciencedirect.com/science/article/pii/S0002929707632133

8. Bonnen PE, Wang PJ, Kimmel M, Chakraborty R, Nelson DL. Haplotype and Linkage

Disequilibrium Architecture for Human Cancer-Associated Genes. Genome Res.

2002;12(12):1846–53. http://genome.cshlp.org/lookup/doi/10.1101/gr.483802

9. Thorstenson YR, Shen P, Tusher VG, Wayne TL, Davis RW, Chu G, et al. Global

Analysis of *ATM* Polymorphism Reveals Significant Functional Constraint. Am J Hum

Genet. 2001;69(2):396–412.

https://linkinghub.elsevier.com/retrieve/pii/S0002929707610854

10. Zhang J, Rowe WL, Clark AG, Buetow KH. Genomewide Distribution of High-

Frequency, Completely Mismatching SNP Haplotype Pairs Observed To Be Common

across Human Populations. Am J Hum Genet. 2003;73(5):1073–81.

https://linkinghub.elsevier.com/retrieve/pii/S0002929707619703

11. Dutta R, Mainsah J, Yatskiv Y, Chakrabortty S, Brennan P, Khuder B, et al. Intricacies in

arrangement of SNP haplotypes suggest "Great Admixture" that created modern humans.

BMC Genomics. 2017;18(1):433.

http://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-017-3776-5

12. Curtis D, Amos W. The human genome harbours widespread exclusive yin yang

haplotypes. Eur J Hum Genet. 2024;32(6):691–6.

https://www.nature.com/articles/s41431-023-01399-5

13. Koshel SM. Geoinformacionnye tehnologii v genogeografii // Sovremennaya

geograficheskaya kartografiya. 2012. P. 158–166.

14. Chen T, Dong B, Lu Z, Tian B, Zhang J, Zhou J, et al. A functional single nucleotide

polymorphism in promoter of *ATM* is associated with longevity. Mech Ageing Dev.

2010;131(10):636–40.

https://www.sciencedirect.com/science/article/pii/S0047637410001582

15. Ding X, Hao Q, Yang M, Chen T, Chen S, Yue J, et al. Polymorphism rs189037C > T in

the promoter region of the *ATM* gene may associate with reduced risk of T2DM in older

adults in China: a case control study. BMC Med Genet. 2017;18(1):84.

https://bmcmedgenet.biomedcentral.com/articles/10.1186/s12881-017-0446-z

FIGURES

Figure 1 – *ATM* haplotypes

Figure 2 - Phylogenetic network of *ATM* haplotypes in Eurasian and African populations

Figure 3 - Scheme of phylogenetic relationships of *ATM* haplotypes

Figure 4 – Medians of population frequencies of *ATM* haplotypes in populations by origin:

AFR – African,

EAS – East Asian,

EEUR – Eastern European,

NEAS – Asian of Northern Eurasia,

SAS – South Asian,

WEUR – Western European

Figure S1 - LD-block of *ATM* SNPs. The numbers in the cells are the r^2 value multiplied by 100

468
469
470 Figure S2 - Phylogenetic networks based on medians of population frequencies of *ATM*

471 haplotypes for populations of African (a), European (b) and Asian (c) origin

473　Figure S3 - Geographic distribution of the most common *ATM* haplotypes

Figure S4 - Geographic distribution of the most widespread Eurasian *ATM* genotypes

476
477 Figure S5 - Geographic distribution of African-specific *ATM* haplotypes

478
479 Figure S6 - Geographic distribution of *ATM* genotypes defined by African-specific

480 haplotypes

Abbreviation	Population	Population size
ALT	Peoples of Altai	83
ANC	Ancestral (Ensembl)	-
BASH	Bashkirs	44
BEB	Bengali	86
BKHY	Buryats, Khamnigans, Yakuts	59
BRNW	Belarusians and Russians of the North-West of the European Russia	61
CDX	Chinese Dai in Xishuangbanna, China	93
CHB	Han Chinese in Bejing, China	103
CHKI	Chukchi, Koryaks, Itelmen	75
CHS	Southern Han Chinese, China	105
ECAU	Peoples of the North-East Caucasus	123
ESN	Esan in Nigeria	99
ETRC	Peoples of Eastern Transcaucasia	87
FE	Peoples of the Far East	104
FIN	Finnish in Finland	99
GBR	British in England and Scotland	91
GIH	Gujarati Indian	103
GWD	Gambian in Western Division, The Gambia	113
IBS	Iberian populations in Spain	107
ITU	Indian Telugu	102
JPT	Japanese in Tokyo, Japan	104
KHMN	Khanty, Mansi, Nenets	63
KHV	Kinh in Ho Chi Minh City, Vietnam	99
KKUN	Kazakhs, Karakalpaks, Uighurs, Nogays	53
KLMN	Mongols (other groups) and Kalmyks	79
KOUD	Komi and Udmurts	84
KV	Karelians and Vepsians	59
LWK	Luhya in Webuye, Kenya	99
MCH	Mari and Chuvash	53
MKH	Mongols (Khalkha)	49
MOR	Mordovians	41
MSL	Mende in Sierra Leone	85
PJL	Punjabi in Lahore, Pakistan	96
RN	Russians of the North of the European Russia	84
RNAR	Russians from the north of the Arkhangelsk region	75
RSE	Russians of the South-East of Central Russia	62
ST	Siberian Tatars	68
STU	Sri Lankan Tamil	102
TAD	Tajiks, Pamiris, Yagnobis	72
TAT	Tatars	60
TSI	Toscani in Italy	107
TUTO	Tuvans and Tofalars	63
URSW	Ukrainians and Russians of the South-West of Central Russia	129
UZTK	Uzbeks, Turkmens, Kyrgyz	98
WCAU	Peoples of the North-West Caucasus	91
WTRC	Peoples of the Western Transcaucasia	56
YRI	Yoruba in Ibadan, Nigeria	108

483   Table 1 - Studied populations

Population	H1i	H1j	H2i	H2j	H3i	H3j	H4i	H4j	H5i	H5j	H6i
ALT										0.556	0.344
BASH						0.048	0.012			0.440	0.488
BEB						0.070				0.587	0.314
BKHY						0.036				0.464	0.455
BRNW										0.458	0.467
CDX			0.016			0.011				0.360	0.559
CHB						0.015				0.374	0.544
CHKI						0.055				0.591	0.280
CHS						0.014				0.390	0.524
ECAU										0.604	0.346
ESN	0.015	0.071	0.308		0.035	0.066	0.116	0.051	0.035	0.177	0.096
ETRC										0.715	0.209
FE						0.011				0.517	0.422
FIN										0.581	0.399
GBR			0.016							0.588	0.379
GIH						0.053				0.631	0.306
GWD	0.084	0.031	0.186		0.058	0.067	0.133	0.071	0.049	0.177	0.097
IBS										0.659	0.322
ITU						0.034				0.618	0.333
JPT						0.019				0.322	0.514
KHMN						0.012				0.492	0.359
KHV			0.015							0.399	0.545
KKUN										0.442	0.450
KLMN						0.019				0.494	0.430
KOUD										0.499	0.402
KV										0.533	0.440
LWK	0.066	0.061	0.232	0.010	0.051	0.045	0.111	0.015	0.101	0.182	0.076
MCH										0.616	0.353
MKH						0.014				0.443	0.471
MOR			0.012							0.499	0.390
MSL	0.044	0.088	0.250	0.012	0.014	0.059	0.135	0.024	0.053	0.241	0.041
PJL						0.036				0.703	0.250
RN										0.537	0.431
RNAR										0.568	0.412
RSE			0.018							0.500	0.456
ST										0.571	0.375
STU						0.064				0.554	0.353
TAD						0.017				0.517	0.441
TAT			0.019							0.497	0.442
TSI										0.692	0.299
TUTO										0.659	0.254
URSW										0.536	0.411
UZTK						0.026				0.500	0.408
WCAU			0.017							0.565	0.365
WTRC										0.687	0.295
YRI		0.083	0.199		0.032	0.106	0.093	0.051	0.046	0.176	0.120

485 Table 2 - Population frequencies of *ATM* haplotypes

486 See Table 1 for population abbreviations

Population	Population frequency of homozygous genotype									Population frequency of heterozygous genotype					
	H6i/H6i	H5j/H5j	H4i/H4i	H3i/H3i	H2i/H2i	H1j/H1j	H1i/H1i	H5i/H5i	H4j/H4j	H6i/H5j	H2i/H6i	H2i/H5j	H3j/H6i	H3j/H5j	H4i/H2i
ALT	0.088	0.238								0.400			0.013		
BASH	0.238	0.143								0.357			0.071	0.024	
BEB	0.116	0.337								0.349			0.035	0.105	
BKHY	0.232	0.161								0.286	0.018		0.036	0.018	
BRNW	0.217	0.250								0.383					
CDX	0.333	0.151								0.355	0.022	0.011	0.011	0.011	0.022
CHB	0.340	0.175								0.330		0.010	0.019	0.010	
CHKI	0.145	0.345								0.164			0.036	0.055	
CHS	0.267	0.114								0.448	0.010	0.010	0.019	0.010	0.010
ECAU	0.123	0.298								0.360		0.009			
ESN	0.020	0.040	0.020	0.010	0.101					0.010	0.051	0.071	0.010	0.020	0.051
ETRC	0.012	0.420								0.272					
FE	0.178	0.211								0.378			0.011	0.011	
FIN	0.202	0.364								0.394		0.010			
GBR	0.121	0.341								0.473	0.022	0.011			0.022
GIH	0.097	0.417								0.359		0.010	0.058	0.049	
GWD	0.009	0.035	0.044	0.009	0.018		0.009		0.009	0.044	0.018	0.053	0.018	0.027	0.018
IBS	0.084	0.411								0.467		0.009			
ITU	0.108	0.382								0.422			0.020	0.039	
JPT	0.279	0.115								0.279		0.019	0.029	0.010	
KHMN	0.097	0.210								0.339			0.016		
KHV	0.263	0.101								0.525	0.010	0.020	0.010		0.010
KKUN	0.154	0.154								0.404			0.019		
KLMN	0.203	0.215								0.342			0.013	0.025	
KOUD	0.159	0.244								0.341	0.017				
KV	0.224	0.328								0.362					
LWK		0.030	0.020		0.061		0.010	0.010		0.030	0.040	0.051		0.020	0.040
MCH	0.137	0.353								0.294			0.020		
MKH	0.200	0.244								0.356				0.022	
MOR	0.098	0.244								0.317	0.024				
MSL		0.059	0.024		0.082	0.035				0.024	0.047	0.071		0.047	0.047
PJL	0.083	0.510								0.313			0.010	0.063	
RN	0.213	0.338								0.363		0.013			
RNAR	0.135	0.297								0.514					
RSE	0.211	0.228								0.421	0.018	0.018		0.018	
ST	0.107	0.250								0.429					
STU	0.147	0.363								0.314			0.078	0.039	
TAD	0.136	0.220								0.424			0.034		
TAT	0.154	0.173								0.481		0.038			
TSI	0.121	0.514								0.346	0.009	0.009			0.009
TUTO	0.063	0.413								0.254				0.016	
URSW	0.195	0.285								0.350					
UZTK	0.143	0.235								0.398	0.010		0.020	0.031	
WCAU	0.101	0.292								0.360	0.011				
WTRC	0.071	0.446								0.411				0.018	
YRI		0.009	0.009		0.019					0.065	0.046	0.046	0.046	0.037	0.046

489   Table 3 - Population genotype frequencies based on *ATM* haplotypes

490   See Table 1 for population abbreviations

Population	D	p-value	Origin
Combined population of Eurasia and Africa	6.5	$\ll 0.05$	
Africa	6.1	$\ll 0.05$	AFR
Northern Eurasia	8.9	$\ll 0.05$	EEUR and NEAS
EUR, EAS and SAS	8.4	$\ll 0.05$	EUR, EAS and SAS
Set of ATM haplotypes in the I branch	-0.6	> 0.05	ATM branch I
Set of ATM haplotypes in the J branch	-0.9	> 0.05	ATM branch J
EUR	7.0	$\ll 0.05$	EUR
EAS	7.8	$\ll 0.05$	EAS
SAS	6.7	$\ll 0.05$	SAS
FIN	6.5	$\ll 0.05$	EUR
GBR	6.4	$\ll 0.05$	
IBS	5.7	$\ll 0.05$	
TSI	5.3	$\ll 0.05$	
BEB	5.6	$\ll 0.05$	SAS
GIH	5.5	$\ll 0.05$	
ITU	5.8	$\ll 0.05$	
PJL	4.6	$\ll 0.05$	
STU	6.1	$\ll 0.05$	
CDX	6.5	$\ll 0.05$	EAS
CHB	6.5	$\ll 0.05$	
CHS	6.7	$\ll 0.05$	
JPT	6.5	$\ll 0.05$	
KHV	6.6	$\ll 0.05$	
MSL	4.9	$\ll 0.05$	AFR
ESN	4.8	$\ll 0.05$	
GWD	5.2	$\ll 0.05$	
LWK	4.8	$\ll 0.05$	
YRI	5.4	$\ll 0.05$	
ALT	6.3	$\ll 0.05$	NEAS
BASH	6.8	$\ll 0.05$	
BKHY	6.7	$\ll 0.05$	
CHKI	6.0	$\ll 0.05$	
FE	6.7	$\ll 0.05$	
KHMN	6.7	$\ll 0.05$	
KKUN	6.7	$\ll 0.05$	
KLMN	6.7	$\ll 0.05$	
MKH	6.7	$\ll 0.05$	
ST	6.4	$\ll 0.05$	
TAD	6.7	$\ll 0.05$	
TUTO	5.5	$\ll 0.05$	
UZTK	6.6	$\ll 0.05$	
BRNW	6.8	$\ll 0.05$	EEUR
ECAU	6.1	$\ll 0.05$	
ETRC	4.6	$\ll 0.05$	
KOUD	6.7	$\ll 0.05$	
KV	6.6	$\ll 0.05$	
MCH	6.0	$\ll 0.05$	
MOR	6.6	$\ll 0.05$	
RN	6.6	$\ll 0.05$	
RNAR	6.5	$\ll 0.05$	
RSE	6.7	$\ll 0.05$	
TAT	6.7	$\ll 0.05$	
URSW	6.5	$\ll 0.05$	
WCAU	6.3	$\ll 0.05$	
WTRC	5.3	$\ll 0.05$	

492   Table S1 - Tajima's test for *ATM* haplotypes

493   Abbreviations for populations can be found in Table S1. and for populations by origin:

494   AFR – African, EAS – East Asian, EEUR – Eastern European, NEAS – Asian of Northern

495   Eurasia, SAS – South Asian, WEUR – Western European