Phylogenetic relationships of ATM polymorphic loci haplotypes in Eurasian and African populations

Marina V. Olkova[1,*], Sergey M. Koshel[2], Georgiy Yu. Ponomarev[1], Andrei A. Alimov[1]

[1] Research Centre for Medical Genetics, Moscow, Russia

[2] Lomonosov Moscow State University, Moscow, Russia

Corresponding author email: genetics@inbox.ru

ABSTRACT

The ATM gene encodes a key kinase involved in DNA repair and cell cycle control, and mutations in it lead to the development of ataxia-telangiectasia and increase susceptibility to cancer. In this context, the study of the distribution patterns of ATM haplotypes in metapopulations living in Eurasia and Africa using population genetic and genogeographic methods appears to be a socially important scientific endeavor.

Polymorphic markers of the ATM gene were studied in 3876 individuals from 46 metapopulations, ATM gene haplotypes and genotypes based on ATM haplotypes were determined, and population frequencies of haplotypes and genotypes were calculated. An analysis of phylogenetic relationships of haplotypes, the Tajima test to assess population changes, and mapping of population frequencies of ATM gene haplotypes and genotypes were performed. The study showed the presence of a linkage block of 28 polymorphic markers of the ATM gene in the Eurasian and African populations. Two dominant ATM haplotypes in Eurasia, which are the result of independent evolution of two branches from the ancestral ATM haplotype, are formed by mutually exclusive alleles and correspond to the "yin-yang" concept.

The greatest diversity of ATM haplotypes is observed on the African continent. It is suggested that the process of haplotype formation occurred on the African continent, with subsequent migration events to the territory of Eurasia.

Key words: ATM haplotypes, LD-block, "yin-yang" concept, mutually exclusive alleles, evolutionary separation

INTRODUCTION

The human ATM gene (ataxia-telangiectasia mutated), which encodes a serine-threonine protein kinase, is located on chromosome 11, locus 11q22-23. Activation of the gene's protein product occurs as a result of double-stranded DNA breaks, leading to the initiation of signaling pathways and the phosphorylation of numerous substrates (several hundred have been identified to date). The final result of cascade reactions is cell cycle arrest and DNA repair. In addition to double-stranded DNA breaks, activation of the ATM protein can be caused by oxidative stress, deficiency of a number of metabolites, hypoxia, and some other factors [1]. ATM is also involved in the activation of the spindle checkpoint during mitosis. Loss of ATM function results in a defective spindle checkpoint and increases the likelihood of aneuploidy [2].

Mutations in the ATM gene are associated with the development of autosomal recessive Louis-Barr syndrome (ataxia-telangiectasia) and with the risk of developing cancer, which is one of the reasons for the research interest in studying the distribution of frequencies of polymorphic variants of this gene in populations [3–6]. One of the most important stages of genetic data research is the linkage disequilibrium (LD) analysis, which allows to identify new clinically significant variants of the ATM gene associated with changes in the function of the gene's protein product and disease severity. As a result of previous scientific studies, blocks of LD in

the ATM gene have been identified and described in a region of ~150 kb, covering the entire length of the gene [7–9]. In particular, Tornston et al. identified seven haplotypes of the ATM gene, two of which were present in 82% of the 93 DNA samples examined from representatives of seven human populations living on all continents. Two of the seven identified haplotypes were found only in African populations with a frequency of 33%. The presence of blocks of LD of varying length was also demonstrated for a number of other autosomal regions of the human genome. In particular, Bonnen's article assessed the LD of non-coding polymorphic markers for 5 genes (TP53, RAG51, BRCA1, BRCA2 and ATM) associated with the cancer development. It was shown that for all genes, taking into account the fact that the study included markers evenly distributed along the gene, there was no tendency for the LD value to decrease with increasing distance. The authors concluded that the LD value can vary widely regardless of the physical distribution of the markers along the length of the DNA sequence of the gene. The observed phenomenon testified to the existence of polymorphic loci in the genome, linked in blocks, with the degree of linkage independent of distance. A similar conclusion was reached by Gabriel et al., who reported the presence of SNP blocks with a small number of common haplotypes, evenly distributed across the human genome. The study was conducted using DNA samples from 275 representatives of the American population of African, European and Asian origin. A link was found between the origins of the populations and specific haplotypes. Haplotypes with mutually exclusive alleles of polymorphic markers, called "yin" and "yang", were also identified [10–12].

Our study aims to perform a population analysis of combined into haplotypes SNP variants of the ATM gene, associated with oncological pathology. In addition, a genogeographic comparison of the North Eurasian gene pool of ATM haplotypes with the gene pools of the peoples of Eurasia and Africa was conducted. Phylogenetic analysis based on all identified

ATM haplotypes was performed to better understand the structural changes accumulated in the gene during the evolutionary process.

MATERIALS AND METHODS

Ethical approvals

This study was conducted under the condition that written informed consent was obtained from all subjects. This consent was approved by the Ethics Committee of the Research Center for Medical Genetics.

Collection of DNA genotype data and populations

1975 DNA samples from individuals representing 27 populations living in Russia and adjacent countries were provided by the Northern Eurasia Biobank. Population sample sizes ranged from 41 to 129 individuals. Individual allelic variants of 269 polymorphic markers of the ATM gene were extracted from a genome-wide genotyping dataset using the Infinium Omni5Exome BeadChip Kit (Illumina Inc., San Diego, California).

Genotype information for 1901 individuals from 19 populations in Eurasia and Africa, with population sample sizes ranging from 85 to 113 individuals, was obtained from the 1000 Genomes Project Phase 3 data (Ensembl).

The general list of populations, their abbreviated names and the size of population samples are presented in Table 1.

The "ancestral" haplotype was determined based on the "ancestral" alleles of the polymorphic markers specified in the Ensembl project, identified based on a three-stage multiple alignment of EPO (Enredo, Pecan, Ortheus).

Statistical data processing

Microsoft Excel (Microsoft Corporation, Redmond, WA) and RStudio 2024.12.1+563 - R version 4.4.2 (RStudio, PBC) were used for statistical data processing.

DNA sample genotyping and polymorphic ATM loci quality assessment

Calculation of the percentage of genotyping of markers and DNA samples was performed using the snpStats package in the RStudio software. Polymorphic markers and samples with a genotyping quality of less than 95% were excluded from the study.

Isolation of haplotypes. Calculation of population frequencies of ATM haplotypes and genotypes

The isolation of haplotypes and the calculation of their population frequencies were performed in the Haploview 4.1 program (Broad Institute, Cambridge).

The "confidence interval method" (Gabriel et al.) was used to isolate blocks of polymorphic variants forming a haplotype. A haplotype block was defined if 95% of SNP pairs within it had D' > 0.8.

Calculation of population frequencies of genotypes based on ATM haplotypes was performed with RStudio software using the plyr package.

Phylogenetic analysis

Phylogenetic networks of ATM haplotype representation in populations were constructed in RStudio using the pegas and geneHapR packages according to the "parsimony principle" (parsimony network). The preparatory files were filled with data on population frequencies of haplotypes, multiplied by 1000.

Calculation of Tajima test for populations and individual "yin" and "yang" branches

The Tajima test was calculated based on the population haplotype frequency data obtained in Haploview 4.1 software. The population number of haplotypes of each type was obtained by multiplying the population frequency of the haplotype by 1000. The Tajima test D-value and p-value were calculated using the pegas package (RStudio).

Cartographic analysis

Cartographic visualization of the population frequencies of haplotypes and genotypes of the ATM gene was performed using the original GeneGeo cartographic package [13]. Frequency calculations for the entire mapped area were performed using the weighted average interpolation method with a weighting function inversely proportional to the cube of the distance and an influence radius of 4000 km. To avoid the influence of map projection distortions calculations, all distances in the interpolation process were calculated on a sphere. The Winkel triple projection with the central meridian 60^{0} E was used to create the maps.

RESULTS

Determination of ATM gene haplotypes and their population frequencies

The genotype data of the combined population of Northern Eurasia allowed the identification of a linkage disequilibrium block (LD-block) of 28 linked polymorphic loci using Haploview 4.1 software − rs189037, rs228590, rs623860, rs600931, rs228592, rs672655, rs627418, rs664677, rs609261, rs645485, rs619972, rs599558, rs595747, rs609655, rs227061, rs227062, rs227064, rs227068, rs227070, rs227074, rs227075, rs425538, rs419716, rs227040, rs664143, rs227094, rs227092 and rs4585, located over 145,795 bp from the 5′-untranslated region to the 3′-untranslated region of the ATM gene (Figure S1).

The value of the square correlation coefficient (r^2) between all pairs of loci in the LD block exceeded 0.8. Of 268,435,456 possible haplotypes for 28 polymorphic loci of the LD block in the combined population of peoples living in the territory of Northern Eurasia, 4 haplotypes were found, and 2 of them were predominant: G-A-C-C-C-A-T-C-T-A-C-C-C-G-A-G-T-A-T-A-T-C-A-T-A-C-G-G and A-G-T-T-A-G-C-T-C-G-T-T-T-A-G-A-C-G-G-G-C-A-C-C-G-A-T-T with a frequency of 39.2% and 54.5%, respectively. The alleles in the two most frequent haplotypes were mutually exclusive, which allows us to classify them as "yin"/"yang" haplotypes. The remaining two detected haplotypes were much less frequent in the population (with a frequency of about 2%) and were the result of recombination between the "yin" and "yang" haplotypes.

Similarly, haplotypes were determined, and their population frequencies were calculated for 28 polymorphic markers of the ATM haplotype block in each of the 27 populations of Northern Eurasia and in 19 populations of Eurasia and Africa using data from the 1000 Genomes Project Phase 3 (Ensembl). From the total set of haplotypes detected in all of the above Eurasian and African populations, rare (with a population frequency of up to 5%) found in single populations, as well as haplotypes that most likely arose as a result of recombination between the "yin-yang" haplotypes, were excluded. As a result, a list of 11 haplotypes of the ATM gene (Table 2) was determined, occurring in 46 populations of Eurasia and Africa. The "ancestral" haplotype (H0) of the human ATM gene was determined using the Ensembl genome browser, which indicates the "ancestral" alleles for each of the 28 linked loci of this gene.

ATM haplotypes-based phylogenetic analysis

Based on the isolated haplotypes, a phylogenetic network was constructed reflecting the evolutionary variability of the ATM gene haplotype block in the human population and the frequency ratio of ATM haplotypes in modern peoples of Eurasia and Africa (Figure 1).

The presence of two independently evolving phylogenetic branches, which we called I and J branches, was discovered, leading from the "ancestral" haplotype to the most common in Eurasia H6i and H5j ("yin" and "yang") haplotypes (Figure 2). All haplotypes analyzed were found in African populations (Figure 3, S2a).

In addition to the main "yin/yang" haplotypes, the H2i and H4i (branch I), H3j (branch J) haplotypes were also present in the Eurasia area. Of these three haplotypes, only H2i was found in populations of European origin (Figure S2b), in contrast to populations of Asian origin (Figure S2c). All three haplotypes are rare (up to 2%) in Eurasian populations. Based on these data, it can be concluded that the formation of the ancestral haplotype and the subsequent divergent evolution of two isolated branches probably occurred on the territory of the African continent.

Tajima test for populations and individual branches of "yin" and "yang" haplotypes

For all populations of Eurasia and Africa, both divided by origin and geographical location, and combined by these criteria, a positive value of the D parameter of the Tajima test (from 4.7 to 9.3) with a p value $\ll 0.05$ was observed, which may be due to a long-term subdivision of the population into several subgroups with limited gene flow between them, balancing selection, a sharp reduction in population size, or other genetic and demographic processes. For each of these populations, a significant deviation from the neutral model of evolution was observed. At the same time, when considering each of the two evolutionary branches leading from the "ancestral" haplotype to the H6i and H5j haplotypes separately, this test showed a statistically insignificant result for the D parameter and p-value ($p \gg 0.05$), which indicating that both divergence paths (I and J) of the "ancestral" ATM haplotype conform to the neutral model of evolution (Table S1).

Geographical analysis of population frequencies of ATM haplotypes and genotypes

The frequencies of ATM haplotypes, as well as their homozygous and heterozygous combinations, are presented in Tables S2-S3 and on maps (Fig. 3-4, Supplement: Fig. S3-S4).

"Yin" and "yang" haplotypes

Cartographic analysis of the prevalence of population frequencies of haplotypes H6i and H5j, as well as homozygous carriers of these haplotypes showed similar results.

Haplotype H5j was in southern Eurasia with maxima in the southern of Western Europe (~ 69%), in Transcaucasia (~ 72%) and in the north-west of South Asia (~ 70%). There was also a significant increase in the frequency of this haplotype in the populations of the indigenous populations of southern Siberia (with a maximum among Tuvans and Tofalars ~ 66%) and Kamchatka (Chukchi, Koryaks, Itelmens ~ 59%), with a significant decrease in East Asia to a frequency of ~36%. The H5j haplotype is the second most common in Africa, although its population frequencies are noticeable lower than in Eurasia - from 18% in most African populations to 24% in the Mende people living on the coast of West Africa.

The H6i haplotype has two geographic maxima - on the east coast of Asia (~ 56% in the Dai people in southwest China) and in eastern Europe (~ 49% in the Bashkirs). Between the maxima, there is a marked decrease in population frequencies in central Siberia, extending to the northern coast of Eurasia and the south of the continent - along the northern border of the Alpide belt. The lowest population frequency of the H6i haplotype in Eurasia is in Eastern Transcaucasia (~ 21%). The H6i haplotype is the fourth most common in African populations – from 4% in the Mende of Sierra Leone to 12% in the Yoruba of Nigeria.

In summary, it can be said that in the Eurasian area, with the exception of East Asia, the H5j haplotype predominates between the two mutually exclusive and most common haplotypes. In the populations of East Asia, the H6i haplotype predominates. In Africa, the population

frequencies of the "yin" and "yang" haplotypes are significantly lower than in Eurasian populations. On average, the H5j haplotype is 2.8 times less common in African populations than in Eurasian populations, and the H6i haplotype is 4.6 times less common.

Heterozygous combination of "yin" and "yang" haplotypes

The heterozygous genotype H6i/H5j turned out was one of the most common in Eurasian populations (from 16.4% to 52.5%). The exception was 9 out of 42 Eurasian populations (Table 5). The prevalence of homozygotes for the H5j haplotype was found in 2 populations of northwestern South Asia (41.7% in Gujaratis of India and 51% in Punjabis of Pakistan), in populations of Transcaucasia (42% in Eastern Transcaucasia and 44.6% in the west), in the Italian population of southern Europe (51.4%), as well as in 3 metapopulations of Northern Eurasia - "Chukchi, Koryaks, Itelmens" (34.5%), "Mari, Chuvash" (35.3%) and "Tuvans, Tofalars" (41.3%). The homozygous genotype for the H6i haplotype turned out was the most common in only one population - the East Asian Han from Beijing, where its frequency was ~ 34%. In Africa, the frequency of H6i/H5j heterozygotes was an order of magnitude lower than in Eurasian (on average ~ 3.5%) than in Eurasia (~ 37%), although in the Yoruba, the H6i/H5j genotype was the most common population genotype (6.5%), despite the complete absence of homozygotes for the H6i haplotype and a very low population frequency of homozygotes for the H5j haplotype (~ 1%). This may be due to the manifestation of balancing selection or migration processes.

The peaks of the H6i/H5j genotype in Eurasia are located on the coasts: in the east of the Indochinese peninsula (~ 52.5%), in the Spanish (~ 46.7%) and British (~ 47.3%) populations of Western Europe, and in the Arkhangelsk region (~ 51.4%) of Northern Europe. There is also a significant increase in heterozygote frequencies in the Eastern European Tatar population (48.1%).

The Eurasian minimum of population frequencies of "yin-yang" heterozygotes is observed among the Chukchi, Koryaks and Itelmens (~ 16.4%), which is unusual, given the population frequencies of homozygous carriers of these haplotypes, which are at the "average European" levels.

Haplotype H3j

Haplotype H3j differs from H5j at 3 positions - rs189037, rs672655, rs619972. With a frequency of 1% to 7%, H3j is found in most Eurasian populations of Asian origin (with the exception of the West Siberian Plain and Vietnam) but has not been found in European populations. Homozygotes for H3j have not been found, apparently because of the low frequencies of the haplotype.

In Africa, this haplotype ranks 5th in population frequency ranging - from 4.5% in the Luhya population of Kenya to 11% in the Yoruba of Nigeria.

Geography of population frequencies of heterozygotes for the haplotype H3j in combination with H6i, H5j, H4i and H2i

The geographical distribution of the H3j haplotype in combination with H6i and H5j generally corresponds to that of the H3j haplotype in populations. The H3j/H6i genotypes are more common along the line running from southern Asia to northern Eurasia, with the highest frequencies among Sri Lankan Tamils (7.8%) and Bashkirs (7.1%). The H3j/H5j genotype is observed with the highest frequency (10.5%) in the Bengal population of South Asia.

The H3j/H4i and H3j/H2i genotypes are found only in Africa, the former with a frequency of 1-2% and the latter with a frequency of 2-6%.

Haplotype H2i

 The most common in Africa, H2i (ranging from 19% to 31%), was found at very low levels (~2%) in some populations of East Asia (southern China and Vietnam) and Europe (UK, Western Caucasus, south-eastern central Russia, Mordovia and Tatarstan).

Prevalence of heterozygous carriers of the haplotype H2i in combination with H6i, H5j and H4i

The geographical distribution of heterozygotes for mixed haplotypes of the most common African H2i and the predominant Eurasian H6i and H5j is quite revealing.

The population frequencies of the H2i/H6i genotype are widespread, although at low frequencies, not only in African populations, where its occurrence is 2-5%, but also in some Eurasian populations, where the frequency of this genotype is 1-2%.

Heterozygous carriers of the H2i/H5j haplotypes are more common in Africa than heterozygotes for H2i/H6i, the range of population frequency for H2i/H5j is from 5 - 7%, and in the Mandinka population of Gambia this genotype is the most common.

Carriers of the H2i/H6i genotype have also been found in all populations of East Asia, Western Europe and some European populations of Northern Eurasia with a population frequency of 1-2%.

The H2i/H4i genotype is found in African populations with a frequency of 3-5%, with only the Mandinka having a frequency of about 2%. This genotype was also found in the southern East Asia and in the British and Italian populations of Europe with a frequency of 1-2%.

Haplotype H4i

Haplotype H4i was found to be the third most common in Africa. It differs from the "East Asian" H6i in 3 positions (rs623860, rs664677, rs609655) and its population frequencies in

Africa range from 9% in the Yoruba to 13.5% in the Mandinka of the Gambia and the Mende of Sierra Leone.

In Eurasia, this haplotype was found only in the Bashkir population (frequency ~1%), probably due to the heterogeneity of this population, which was formed by different ethnic groups.

Population features of ATM haplotypes in Africa

Haplotypes H1i, H3i, H5i, H1j, H2j and H4j were found only in African populations. The "ancestral" haplotype H0 is located in the phylogenetic network between haplotypes H1j and H1i.

The H1j haplotype, found in African populations with a frequency of 3% to 9% (on the west coast), differs from the "ancestral" haplotype in only one position (the G allele in the polymorphic marker rs645485, characteristic of H5j). The H1j haplotype was only found homozygous in the Mende of Sierra Leone, where the population frequency of its homozygous carriers was ~3.5%. A less common haplotype, H1i, is most common on the East and West African coasts (7-8%) and is almost absent in the central part of the continent (0 - 1%). H1i is genetically distant by 4 nucleotides from the ancestral H0 in branch I due to the alleles of the polymorphic markers rs227062, rs227074, rs227040 and rs4585. Homozygous carriers of the H1i haplotype have been found at a population frequency of about 1% in two African peoples, the Mandinka of Gambia and the Luhya of Kenya.

The H2j haplotype was found in the Luhya and Mende populations with a frequency of 1%.

The H3i haplotype was the rarest in the Mende population (~ 1%), it was slightly more common among the Yoruba and Ishan of Nigeria (3 - 3.5%), and its population frequency increased to 5 - 6% among the Luhya and Mandinka on the east and west coasts.

H4j is most common among the Mandinka (~ 7%) in West Africa, with its presence gradually decreasing in the east among the Luhya (1.5%).

In contrast, H5i has its maximum frequency on the east coast (~ 10%), while in other African populations its population frequencies are ~ 4 - 5%.

In general, the African continent is characterized by the greatest diversity of ATM haplotypes and the originality of their population distribution.

DISCUSSION

The study of ATM haplotypes in human populations of Eurasia and Africa from the standpoint of the "yin-yang" concept showed the presence in all populations of a conservative LD-block consisting of 28 extra-exon loci of the ATM gene located throughout the DNA sequence of the gene, including 5' and 3' untranslated regions of the ATM gene. Analysis of the phylogenetic relationships between 11 identified ATM haplotypes revealed the presence of two independent branches of ATM gene evolution from the ancestral haplotype, leading to the formation and widespread distribution of haplotypes with mutually exclusive alleles according to the "yin-yang" principle. In the samples analyzed outside the African continent, the dominant haplotypes in all human populations were the H6i haplotype (from 21 to 56%) and the H5j haplotype (from 32 to 72%), while the population frequency of the remaining haplotypes did not exceed 7% in Eurasian populations. In the sample from the African continent, 11 haplotypes were found, reflecting all stages of gene evolution from the "ancestral" haplotype. The most common was the H2i haplotype, its median frequency in African populations was 23%. On the territory of Eurasia, the H2i haplotype was found in only 7 populations, where its occurrence did not exceed 2%, which may be the result of evolutionary and demographic processes. Combining the obtained information on the evolutionary variability of ATM haplotypes with data on their frequencies in modern populations of Africa and Eurasia, it can be assumed that the process of formation of existing ATM haplotypes occurred on the African continent. Phylogenetic analysis of ATM haplotypes demonstrates the existence of two isolated ancestral

human populations in the past, followed by their merging into a single mixed population on the African continent (Figure 3). Similar conclusions were reached in the project[11], which was conducted using data from the human genome of the 1000 Genomes Project, as well as the genomes of Denisovans and Neanderthals - Eastern and Western. Dutta et al. found an abundance of mutually exclusive pairs of common haplotypes for population-frequent (>25%) polymorphic markers in the human genome. Moreover, these "yin" and "yang" haplotypes had close population frequencies in Eurasia. At the same time, this study identified a category of haplotypes that are mostly observed in Africa. These haplotypes are composed of fragments of "yin" and "yang" haplotypes, hence the term "mosaic" used by the authors.

The results of the Tajima test (D > 4.7, p << 0.05) for the combined population and individual populations of the world in our study confirm the fact of the probable division of the ancestral human population into several independent groups, and statistically insignificant D-values in the calculation of the Tajima test for haplotypes of the "yin" and "yang" branches ((- 0.6) and (- 0.8), at p >> 0.05) suggest a neutral nature of evolution for each of the branches.

The combined mixed population of carriers of ATM haplotypes, based on phylogenetic data, continued to exist on the African continent. Only a part of the ATM haplotypes from the "African" set were found on the territory of Eurasia. As a result of genetic drift and, possibly, balancing selection, the "yin" and "yang" haplotypes became dominant in Eurasia, while the others either disappeared completely or survived in small quantities in individual populations. The peoples of Asia differ from those of European origin, firstly, by the widespread occurrence of the H3j haplotype (up to 7%) and, secondly, by the prevalence of the H6i haplotype in East Asia. It is important to note that in South Asia, the dominant haplotype among peoples of European origin, H5j, prevails, which allows us to consider the modal haplotype H5j in the population as a distinctive Indo-European feature. An analysis of literary data on the connection of the analyzed polymorphic loci with oncological diseases revealed one important

contradiction. In particular, in populations of East Asian origin, the H5j haplotype, containing the rs189037, rs609261 and rs4585 loci, is associated with the manifestation of negative effects in oncopathology. In peoples of European origin, this association is not observed, which may be due to the presence of an additional factor, as yet unknown, which acts as an additive in East Asian populations. It is also worth paying attention to the prevalence of heterozygotes for the "yin" and "yang" haplotypes in most Eurasian populations. There are scientific data on the positive effect of heterozygous carrier references between the GA genotype and longevity, which the authors associate with a moderate level of ATM protein expression [14]. A study by Ding et al. showed that heterozygous genotype for rs189037 is associated with a reduced risk of type 2 diabetes in older adults [15].

CONCLUSION

Our study of evolutionary changes in the ATM gene haplotypes, represented by high-frequency polymorphic loci throughout the DNA sequence of this gene, indicates a very ancient, significant in time African period of isolation of two branches of the ancestral human population and the subsequent merging of these branches into a single population space on the African continent. Repeated migration events to the territory of Eurasia probably occurred after the unification of ancient isolated subpopulations. Some differences in the composition of ATM haplotypes in the east and west of the Eurasian continent, as well as in individual populations, are probably due to the effect of gene drift.

DATA AND CODE AVAILABILITY

The 1000 Genomes data can be obtained from the 1000 Genomes Project (Ensembl) Phase 3 data. Data on populations of Northern Eurasia and the R code used in this work are available upon reasonable request.

REFERENCES

1. Ueno S, Sudo T, Hirasawa A. ATM: Functions of ATM Kinase and Its Relevance to Hereditary Tumors. Int J Mol Sci. 2022;23(1):523. https://www.mdpi.com/1422-0067/23/1/523

2. Yang C, Tang X, Guo X, Niikura Y, Kitagawa K, Cui K, et al. Aurora-B Mediated ATM Serine 1403 Phosphorylation Is Required for Mitotic ATM Activation and the Spindle Checkpoint. Mol Cell. 2011;44(4):597–608. https://linkinghub.elsevier.com/retrieve/pii/S1097276511008148

3. Zhao Z liang, Xia L, Zhao C, Yao J. ATM rs189037 (G > A) polymorphism increased the risk of cancer: an updated meta-analysis. BMC Med Genet. 2019;20(1):28. https://bmcmedgenet.biomedcentral.com/articles/10.1186/s12881-019-0760-8

4. Tao Y, Mei Y, Ying R, Chen S, Wei Z. The ATM rs189037 G>A polymorphism is associated with the risk and prognosis of gastric cancer in Chinese individuals: A case–control study. Gene. 2020;741:144578. https://www.sciencedirect.com/science/article/pii/S037811192030247X

5. Zhou K, Bellenguez C, Spencer CCA, Bennett AJ, Coleman RL, Tavendale R, et al. Common variants near ATM are associated with glycemic response to metformin in type 2 diabetes. Nat Genet. 2011;43(2):117–20. https://doi.org/10.1038/ng.735

6. Lee SA, Lee KM, Lee SJ, Yoo KY, Park SK, Noh DY, et al. Antioxidant Vitamins Intake, Ataxia Telangiectasia Mutated (ATM) Genetic Polymorphisms, and Breast Cancer Risk. Nutr Cancer. 2010;62(8):1087–94. https://doi.org/10.1080/01635581.2010.492088

7. Bonnen PE, Story MD, Ashorn CL, Buchholz TA, Weil MM, Nelson DL. Haplotypes at ATM Identify Coding-Sequence Variation and Indicate a Region of Extensive Linkage

Disequilibrium. Am J Hum Genet. 2000;67(6):1437–51.

https://www.sciencedirect.com/science/article/pii/S0002929707632133

8. Bonnen PE, Wang PJ, Kimmel M, Chakraborty R, Nelson DL. Haplotype and Linkage
   Disequilibrium Architecture for Human Cancer-Associated Genes. Genome Res.
   2002;12(12):1846–53. http://genome.cshlp.org/lookup/doi/10.1101/gr.483802

9. Thorstenson YR, Shen P, Tusher VG, Wayne TL, Davis RW, Chu G, et al. Global
   Analysis of ATM Polymorphism Reveals Significant Functional Constraint. Am J Hum
   Genet. 2001;69(2):396–412.
   https://linkinghub.elsevier.com/retrieve/pii/S0002929707610854

10. Zhang J, Rowe WL, Clark AG, Buetow KH. Genomewide Distribution of High-
    Frequency, Completely Mismatching SNP Haplotype Pairs Observed To Be Common
    across Human Populations. Am J Hum Genet. 2003;73(5):1073–81.
    https://linkinghub.elsevier.com/retrieve/pii/S0002929707619703

11. Dutta R, Mainsah J, Yatskiv Y, Chakrabortty S, Brennan P, Khuder B, et al. Intricacies in
    arrangement of SNP haplotypes suggest "Great Admixture" that created modern humans.
    BMC Genomics. 2017;18(1):433.
    http://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-017-3776-5

12. Curtis D, Amos W. The human genome harbours widespread exclusive yin yang
    haplotypes. Eur J Hum Genet. 2024;32(6):691–6.
    https://www.nature.com/articles/s41431-023-01399-5

13. Koshel SM. Geoinformacionnye tehnologii v genogeografii // Sovremennaya
    geograficheskaya kartografiya. 2012. P. 158–166.

14. Chen T, Dong B, Lu Z, Tian B, Zhang J, Zhou J, et al. A functional single nucleotide polymorphism in promoter of ATM is associated with longevity. Mech Ageing Dev. 2010;131(10):636–40.

https://www.sciencedirect.com/science/article/pii/S0047637410001582

15. Ding X, Hao Q, Yang M, Chen T, Chen S, Yue J, et al. Polymorphism rs189037C > T in the promoter region of the ATM gene may associate with reduced risk of T2DM in older adults in China: a case control study. BMC Med Genet. 2017;18(1):84.

https://bmcmedgenet.biomedcentral.com/articles/10.1186/s12881-017-0446-z

TABLES

Abbreviation	Population	Population size
ALT	Peoples of Altai	83
ANC	Ancestral (Ensembl)	-
BASH	Bashkirs	44
BEB	Bengali	86
BKHY	Buryats, Khamnigans, Yakuts	59
BRNW	Belarusians and Russians of the North-West of the European Russia	61
CDX	Chinese Dai in Xishuangbanna, China	93
CHB	Han Chinese in Bejing, China	103
CHKI	Chukchi, Koryaks, Itelmen	75
CHS	Southern Han Chinese, China	105
ECAU	Peoples of the North-East Caucasus	123
ESN	Esan in Nigeria	99
ETRC	Peoples of Eastern Transcaucasia	87
FE	Peoples of the Far East	104
FIN	Finnish in Finland	99
GBR	British in England and Scotland	91
GIH	Gujarati Indian	103
GWD	Gambian in Western Division, The Gambia	113
IBS	Iberian populations in Spain	107
ITU	Indian Telugu	102
JPT	Japanese in Tokyo, Japan	104
KHMN	Khanty, Mansi, Nenets	63
KHV	Kinh in Ho Chi Minh City, Vietnam	99
KKUN	Kazakhs, Karakalpaks, Uighurs, Nogays	53
KLMN	Mongols (other groups) and Kalmyks	79
KOUD	Komi and Udmurts	84
KV	Karelians and Vepsians	59
LWK	Luhya in Webuye, Kenya	99
MCH	Mari and Chuvash	53
MKH	Mongols (Khalkha)	49
MOR	Mordovians	41
MSL	Mende in Sierra Leone	85
PJL	Punjabi in Lahore, Pakistan	96
RN	Russians of the North of the European Russia	84
RNAR	Russians from the north of the Arkhangelsk region	75
RSE	Russians of the South-East of Central Russia	62

Abbreviation	Population	Population size
ST	Siberian Tatars	68
STU	Sri Lankan Tamil	102
TAD	Tajiks, Pamiris, Yagnobis	72
TAT	Tatars	60
TSI	Toscani in Italy	107
TUTO	Tuvans and Tofalars	63
URSW	Ukrainians and Russians of the South-West of Central Russia	129
UZTK	Uzbeks, Turkmens, Kyrgyz	98
WCAU	Peoples of the North-West Caucasus	91
WTRC	Peoples of the Western Transcaucasia	56
YRI	Yoruba in Ibadan, Nigeria	108

Table 1 - Studied populations

Table 2 – ATM haplotypes

FIGURES

Figure 1 - Phylogenetic network of ATM haplotypes in Eurasian and African populations

Figure 2 - Scheme of phylogenetic relationships of ATM haplotypes

Figure 3 – Medians of population frequencies of ATM haplotypes in populations by origin:

AFR – African,

EAS – East Asian,

EEUR – Eastern European,

NEAS – Asian of Northern Eurasia,

SAS – South Asian,

WEUR – Western European

Figure 3 - Geographic distribution of the most common ATM haplotypes

Figure 4 - Geographic distribution of the most widespread Eurasian ATM genotypes

SUPPLEMENT

Tables

Population	D	p-value	Origin
Combined population of Eurasia and Africa	6,5	<< 0.05	
Africa	6,1	<< 0.05	AFR
Northern Eurasia	8,9	<< 0.05	EEUR&NEAS
EUR&EAS&SAS	8,4	<< 0.05	EUR&EAS&SAS
Set of ATM haplotypes in the I branch	-0,6	> 0.05	ATM branch I
Set of ATM haplotypes in the J branch	-0,9	> 0.05	ATM branch J
EUR	7,0	<< 0.05	EUR
EAS	7,8	<< 0.05	EAS
SAS	6,7	<< 0.05	SAS
FIN	6,5	<< 0.05	EUR
GBR	6,4	<< 0.05	
IBS	5,7	<< 0.05	
TSI	5,3	<< 0.05	
BEB	5,6	<< 0.05	SAS
GIH	5,5	<< 0.05	
ITU	5,8	<< 0.05	
PJL	4,6	<< 0.05	
STU	6,1	<< 0.05	
CDX	6,5	<< 0.05	EAS
CHB	6,5	<< 0.05	
CHS	6,7	<< 0.05	
JPT	6,5	<< 0.05	
KHV	6,6	<< 0.05	
MSL	4,9	<< 0.05	AFR
ESN	4,8	<< 0.05	
GWD	5,2	<< 0.05	
LWK	4,8	<< 0.05	
YRI	5,4	<< 0.05	
ALT	6,3	<< 0.05	NEAS
BASH	6,8	<< 0.05	
BKHY	6,7	<< 0.05	
CHKI	6,0	<< 0.05	
FE	6,7	<< 0.05	
KHMN	6,7	<< 0.05	
KKUN	6,7	<< 0.05	
KLMN	6,7	<< 0.05	
MKH	6,7	<< 0.05	
ST	6,4	<< 0.05	
TAD	6,7	<< 0.05	
TUTO	5,5	<< 0.05	
UZTK	6,6	<< 0.05	
BRNW	6,8	<< 0.05	EEUR
ECAU	6,1	<< 0.05	
ETRC	4,6	<< 0.05	
KOUD	6,7	<< 0.05	
KV	6,6	<< 0.05	
MCH	6,0	<< 0.05	
MOR	6,6	<< 0.05	
RN	6,6	<< 0.05	
RNAR	6,5	<< 0.05	
RSE	6,7	<< 0.05	
TAT	6,7	<< 0.05	
URSW	6,5	<< 0.05	
WCAU	6,3	<< 0.05	
WTRC	5,3	<< 0.05	

Table S1 - Tajima's test for ATM haplotypes

Population	H1i	H1j	H2i	H2j	H3i	H3j	H4i	H4j	H5i	H5j	H6i
ALT										0,556	0,344
BASH						0,048	0,012			0,44	0,488
BEB						0,07				0,587	0,314
BKHY						0,036				0,464	0,455
BRNW										0,458	0,467
CDX			0,016			0,011				0,36	0,559
CHB						0,015				0,374	0,544
CHKI						0,055				0,591	0,28
CHS						0,014				0,39	0,524
ECAU										0,604	0,346
ESN	0,015	0,071	0,308		0,035	0,066	0,116	0,051	0,035	0,177	0,096
ETRC										0,715	0,209
FE						0,011				0,517	0,422
FIN										0,581	0,399
GBR			0,016							0,588	0,379
GIH						0,053				0,631	0,306
GWD	0,084	0,031	0,186		0,058	0,067	0,133	0,071	0,049	0,177	0,097
IBS										0,659	0,322
ITU						0,034				0,618	0,333
JPT						0,019				0,322	0,514
KHMN						0,012				0,492	0,359
KHV			0,015							0,399	0,545
KKUN										0,442	0,45
KLMN						0,019				0,494	0,43
KOUD										0,499	0,402
KV										0,533	0,44
LWK	0,066	0,061	0,232	0,01	0,051	0,045	0,111	0,015	0,101	0,182	0,076
MCH										0,616	0,353
MKH						0,014				0,443	0,471
MOR			0,012							0,499	0,39
MSL	0,044	0,088	0,25	0,012	0,014	0,059	0,135	0,024	0,053	0,241	0,041
PJL						0,036				0,703	0,25
RN										0,537	0,431
RNAR										0,568	0,412
RSE			0,018							0,5	0,456
ST										0,571	0,375
STU						0,064				0,554	0,353
TAD						0,017				0,517	0,441
TAT			0,019							0,497	0,442
TSI										0,692	0,299
TUTO										0,659	0,254
URSW										0,536	0,411
UZTK						0,026				0,5	0,408
WCAU			0,017							0,565	0,365
WTRC										0,687	0,295
YRI		0,083	0,199		0,032	0,106	0,093	0,051	0,046	0,176	0,12

Table S2 - Population frequencies of ATM haplotypes

Population	Population frequency of homozygous genotype									Population frequency of heterozygous genotype					
	H6i/H6i	H5j/H5j	H4i/H4i	H3i/H3i	H2iH2i	H1j/H1j	H1i/H1i	H5i/H5i	H4j/H4j	H6i/H5j	H2i/H6i	H2i/H5j	H3j/H6i	H3j/H5j	H4i/H2i
ALT	0,088	0,238								0,400			0,013		
BASH	0,238	0,143								0,357			0,071	0,024	
BEB	0,116	0,337								0,349			0,035	0,105	
BKHY	0,232	0,161								0,286	0,018		0,036	0,018	
BRNW	0,217	0,250								0,383					
CDX	0,333	0,151								0,355	0,022	0,011	0,011	0,011	0,022
CHB	0,340	0,175								0,330		0,010	0,019	0,010	
CHKI	0,145	0,345								0,164			0,036	0,055	
CHS	0,267	0,114								0,448	0,010	0,010	0,019	0,010	0,010
ECAU	0,123	0,298								0,360		0,009			
ESN	0,020	0,040	0,020	0,010	0,101					0,010	0,051	0,071	0,010	0,020	0,051
ETRC	0,012	0,420								0,272					
FE	0,178	0,211								0,378			0,011	0,011	
FIN	0,202	0,364								0,394		0,010			
GBR	0,121	0,341								0,473	0,022	0,011			0,022
GIH	0,097	0,417								0,359		0,010	0,058	0,049	
GWD	0,009	0,035	0,044	0,009	0,018		0,009		0,009	0,044	0,018	0,053	0,018	0,027	0,018
IBS	0,084	0,411								0,467		0,009			
ITU	0,108	0,382								0,422			0,020	0,039	
JPT	0,279	0,115								0,279		0,019	0,029	0,010	
KHMN	0,097	0,210								0,339			0,016		
KHV	0,263	0,101								0,525	0,010	0,020	0,010		0,010
KKUN	0,154	0,154								0,404			0,019		
KLMN	0,203	0,215								0,342			0,013	0,025	
KOUD	0,159	0,244								0,341	0,017				
KV	0,224	0,328								0,362					
LWK		0,030	0,020		0,061		0,010	0,010		0,030	0,040	0,051		0,020	0,040
MCH	0,137	0,353								0,294			0,020		
MKH	0,200	0,244								0,356				0,022	
MOR	0,098	0,244								0,317	0,024				
MSL		0,059	0,024		0,082	0,035				0,024	0,047	0,071		0,047	0,047
PJL	0,083	0,510								0,313			0,010	0,063	
RN	0,213	0,338								0,363		0,013			
RNAR	0,135	0,297								0,514					
RSE	0,211	0,228								0,421	0,018	0,018		0,018	
ST	0,107	0,250								0,429					
STU	0,147	0,363								0,314			0,078	0,039	
TAD	0,136	0,220								0,424			0,034		
TAT	0,154	0,173								0,481		0,038			
TSI	0,121	0,514								0,346	0,009	0,009			0,009
TUTO	0,063	0,413								0,254				0,016	
URSW	0,195	0,285								0,350					
UZTK	0,143	0,235								0,398	0,010		0,020	0,031	
WCAU	0,101	0,292								0,360	0,011				
WTRC	0,071	0,446								0,411				0,018	
YRI		0,009	0,009		0,019					0,065	0,046	0,046	0,046	0,037	0,046

Table S3 - Population genotype frequencies based on ATM haplotypes

Figures

Figure S1 - LD-block of ATM SNPs. The numbers in the cells are the r^2 value multiplied by 100

Figure S2 - Phylogenetic networks based on medians of population frequencies of ATM haplotypes for populations of African (a), European (b) and Asian (c) origin

Figure S3 - Geographic distribution of African-specific ATM haplotypes

Figure S4 - Geographic distribution of ATM genotypes defined by African-specific haplotypes