# Locus coeruleus-related insula activation supports implicit learning

**Authors:**

Martin J. Dahl*[1,2,3], Tiantian Li[1,], Mara Mather[3,4,5°], & Markus Werkle-Bergner[1°]

**Affiliations:**

[1]Center for Lifespan Psychology, Max Planck Institute for Human Development, Berlin, Germany

[2]Max Planck UCL Centre for Computational Psychiatry and Ageing Research, Berlin, Germany.

[3]Davis School of Gerontology, University of Southern California, Los Angeles, CA, USA

[4]Department of Psychology, University of Southern California, Los Angeles, CA, USA

[5]Department of Biomedical Engineering, University of Southern California, Los Angeles, CA, USA

**\* Corresponding author**:

MJD (dahl@mpib-berlin.mpg.de)

° These authors contributed equally (shared senior authorship) and are listed alphabetically.

**ORCIDs:**

0000-0003-3525-0539
0009-0004-8578-1092
0000-0003-4331-6112
0000-0002-6399-9996

**Abstract**

The noradrenergic locus coeruleus and its neuromodulatory cortical projections are
critical for adaptive behavior, yet their contributions to implicit learning in novel
environments remain incompletely understood, due to challenges in non-invasive assessment.
Here, we combined multimodal neuroimaging—including locus-coeruleus-sensitive structural
MRI, concurrent pupillometry–fMRI, and PET-derived noradrenergic transporter maps—with
repeated behavioral assessments to investigate noradrenergic contributions to implicit
learning across younger and older adults (n = 77).
Salient expectation-violating stimuli elicited pupil dilation, indicating enhanced
neuromodulation, activated the action-mode network and deactivated the default-mode
network. Pupil-linked BOLD responses suggested a functional coupling between the locus
coeruleus and action-mode network, further supported by spatial overlap of activation patterns
with PET-derived noradrenergic transporter maps. Locus coeruleus MRI-guided functional
connectivity analyses demonstrated that locus coeruleus activity is coupled to anterior insula
activation, suggesting a noradrenergic role in shifting cortical dynamics toward action-
oriented processing.
Behaviorally, participants implicitly learned the statistical task structure over time, as
evidenced by reaction time adjustments based on stimulus probabilities. Critically, stronger
locus coeruleus integrity, greater task-related anterior insula activation, and more pronounced
pupil dilation were associated with enhanced implicit learning, highlighting the behavioral
relevance of noradrenergic neuromodulation. Notably, noradrenergic responses and their link
to learning were preserved across age groups, suggesting a robust noradrenergic role in
supporting adaptive behavior throughout adulthood.
These findings provide novel insights into the neuromodulatory mechanisms
underlying learning and cognitive flexibility, emphasizing the pivotal role of locus coeruleus–
action-mode network interactions in behavioral adaptation.

*Keywords:* locus coeruleus, neuromodulation, noradrenaline, norepinephrine, MRI,
pupillometry, attention, memory, implicit learning, aging

## 1. Introduction

Optimal behavior requires dynamic adaptation to changing environments [1]. On a neurochemical level, unexpected environmental changes transiently increase noradrenaline release from the locus coeruleus, a brainstem nucleus that facilitates attention and learning [2–4]. Specifically, noradrenaline release has long been conceptualized as a neural interrupt signal that promotes rapid behavioral adaptation to new environmental imperatives [5–8]. The momentary processing of salient stimuli is thereby facilitated by a noradrenergic enhancement of neural gain, effectively increasing signal-to-noise ratios in neural circuits [2,9–12]. By contrast, long-term adjustments to salient events are mediated by noradrenergic influences on hippocampal synaptic plasticity [3,13–16].

In a similar vein, more recent accounts propose that the locus coeruleus is activated when predictions about the world are violated. The resulting noradrenaline release helps refine internal model of the world by adjusting the rate of learning in cortical target regions [17–21].

To enable these functions, the locus coeruleus receives critical contextual information about stimulus salience and utility from cortical structures such as the anterior cingulate and insula [2,9]. In primates, these forebrain structures send major direct inputs to the locus coeruleus to recruit noradrenergic neuromodulation and modulate brain-wide processing according to environmental demands [9].

The anterior cingulate and insular cortices have recently been grouped into the action-mode network [22], overlapping with the cingulo-opercular network or ventral attention network [6,23,24]. The action-mode network is closely linked to initiating and maintaining states of heightened arousal and focused attention, as well as the generation and updating of actions based on salient internal or external signals [22,25]. In this context, arousal is defined as a continuum of sensitivity to environmental stimuli [26], tightly regulated by noradrenergic neuromodulation [27–29]. In line with this, it has been proposed that the locus coeruleus modulates or is part of the action-mode network [6,22,30], but direct in vivo evidence remains limited.

Aging dysregulates the noradrenergic system [31–33], possibly due to the age-related accumulation of pathology [34–37]. In accordance with this, converging evidence based on cognitive assessments suggests that age differences in proxies of noradrenergic neuromodulation contribute to late-life impairments in attention [38–40], learning, and memory [41–43]; i.e., in situations when participants were *explicitly* instructed about the cognitive testing. However, the role of the locus coeruleus in how younger and older adults *implicitly* learn to adapt their behavior in new situations remains underexplored.

Here, we investigated the implications of locus coeruleus–cortical network interactions for implicit behavioral adaptation using a multimodal age-comparative approach. To this end, we used a modified conditioned oddball task, previously shown to reliably increase locus coeruleus spiking in non-human primates [44]. In this variant of the oddball task, participants detect infrequent, distinctive stimuli amid a series of repetitive standard stimuli. To further increase recruitment of the locus coeruleus, infrequent oddball stimuli are pre-conditioned by repeatedly pairing them with appetitive or aversive outcomes [40,44–46]. Our participants completed the conditioned oddball task twice on consecutive days. Trial-level behavioral responses from the task allowed us to examine implicit learning within and across sessions. In this context, implicit learning refers to how initially naïve participants pick up the statistical regularities of the originally new but increasingly familiar task environment and optimize their behavior over time (i.e., without explicit instructions).

To overcome challenges in non-invasive assessment posed by the locus coeruleus' small size and location in the brainstem [47–49], we relied on several recent additions to the neuroscience toolkit. In particular, dedicated structural magnetic resonance imaging (MRI) sequences can reveal the locus coeruleus as a cluster of distinct bright (hyperintense) voxels

bordering the fourth ventricle[41,49]. Transgenic animal[50,51] and human post-mortem[52] validation
studies suggest this signal may serve as an in-vivo proxy for locus coeruleus integrity. Here,
we relied on neuromodulation-sensitive MRI to assess individual differences in locus
coeruleus integrity and to reliably localize the small brainstem nucleus in functional analyses.
To additionally track locus coeruleus activation patterns, we combined fMRI with concurrent
pupillometry, as increasing arousal-related neuromodulatory activity, including in the
noradrenergic system, dilates pupils[53–55]. Finally, we compared the observed pupillometry–
fMRI activation maps to the brain-wide distribution of noradrenergic and other
neuromodulatory transporters[56] to gauge potential dependencies. In sum, younger and older
adults completed a conditioned oddball task while we assessed multiple proxies for locus
coeruleus structure and function to explore their role in implicit learning.
In a brief synopsis of our findings, salient stimuli reliably dilated younger and older
participants' pupils and activated the action-mode network. Pupil-indexed neuromodulation
correlated with brainstem and action-mode network activation, and was associated with
noradrenergic transporter distribution, suggesting potential links to the locus coeruleus, which
we confirmed using functional connectivity analyses. Initially naïve participants implicitly
learned the structure of the new task environment and adapted their behavior over time.
Behavioral adaptation (i.e., implicit learning) was positively associated with structural and
functional locus coeruleus and action-mode network measures. Together, these results suggest
that noradrenergic signals support refining internal models of environmental contingencies for
behavioral adaptation by modulating attention and memory following salient events. By
integrating advanced MRI, pupillometry, and functional connectivity analyses, we reveal
critical insights into how neuromodulatory signals dynamically drive behavioral flexibility.
103

## a Conditioned oddball task

## b Experimental design

104
***Figure 1.* Overview of the conditioned oddball task (a) and experimental phases (b)**. As part of a
three-stimulus oddball task, previously conditioned stimuli (CS+) are presented as infrequent oddball
stimuli while non-conditioned perceptually matched control stimuli (CS−) serve as frequent standards.
A second oddball category consists of perceptually matched but previously not shown stimuli,
hypothesized to elicit a novelty response. (**b**) Fear conditioning sessions are repeated over three
consecutive days while collecting eye tracking and EEG data. The conditioned oddball task takes
place on Day 2 (pupillometry-EEG) and Day 3 (pupillometry-fMRI). During the oddball tasks, no
electrical stimulation is applied.

113
**2. Results**

*Conditioning reliably increases pupil-indexed neuromodulation across age groups*
To increase recruitment of the locus coeruleus, experimental stimuli that were
subsequently shown in the main task were pre-conditioned by repeatedly pairing them with an
aversive electrical stimulation (Figure 1,[44]). Robust pupil dilation following presentation of
conditioned stimuli (CS) relative to perceptually matched control stimuli (CS−) suggests this
procedure enhanced neuromodulatory activation, including in the noradrenergic system[55],
across age groups and assessment days (Figure 2a; all $p_{\text{cluster-corr}} \leq 0.002$).
122

123
*Figure 2*. **Stimulus-related pupil responses during fear conditioning (a) and conditioned oddball**
**tasks (b)**. During daily fear conditioning sessions, fear conditioned stimuli (CS+; red) elicit stronger
pupil dilation relative to perceptually matched control stimuli (CS–, black), suggesting
neuromodulatory activation[53,54]. Note that both stimuli initially elicit comparable pupil constriction
due to higher luminance compared to the background (pupil light reflex after stimulus presentation at
time 0). (**b**) During subsequent pupillometry-EEG (day 1) and pupillometry-fMRI (day 2) conditioned
oddball tasks, oddball presentation (red) is linked to larger pupil dilation relative to perceptually
matched control stimuli (standard; black). Differences in pupil light reflexes between oddball
acquisitions result from different background luminance in the two laboratories (EEG, MRI). The
black horizontal lines indicate the temporal extent of the significant group level cluster, controlling for
multiple comparisons, which is comparable across days. Shaded areas represent standard errors of the
mean.

136
*Conditioned oddball stimuli dilate the pupil and activate the action-mode network*
During the subsequent main task, salient stimuli—oddballs that were either pre-
conditioned or novel—reliably dilated participants' pupils ($p_{cluster-corr} \leq 0.002$) and activated
the action-mode network relative to perceptually matched control stimuli (standards [CS–];
Figures 2b and 3).
Action-mode network recruitment encompassed the bilateral anterior insula ($T =$
11.81, $p_{FWE-corr} < 0.001$ MNI: 30, 22, 10), middle cingulate gyrus ($T = 8.95$, $p_{FWE} < 0.001$
MNI: 3.5, 7.5, 31.5), thalamus ($T = 7$, $p_{FWE} < 0.001$ MNI: 12, –20, 12) and cerebellum ($T =$
7.67, $p_{FWE} < 0.001$ MNI: 30, –50, –24;[22,25]). In addition, salient stimuli elicited prominent
dorsal attention-network activation (bilateral superior parietal lobule; $T = 13.8$, $p_{FWE-corr} <$
0.001 MNI: 42, –38, 50;[6,25]) and default-mode network *de*activation (e.g., bilateral frontal
pole, $T = –11.64$, $p_{FWE} < 0.001$ MNI: –4, 68, 4;[22]). The un/thresholded whole-brain contrast
maps are available at osf.io/4t5hj.
Together, this suggests task-related neuromodulatory activation, as indicated by pupil
dilation, as well as activation of the action-mode network, reflecting major cortical in- and
output regions of the noradrenergic system[6,9,57]. This may indicate that in response to salient
expectation-violating stimuli, there is a transition from default-mode to action-mode network
activation, likely facilitated by noradrenergic neuromodulation, which supports coping with
changing task demands[5,17,22].
156

*Figure 3*. **Salient oddball stimuli elicit action-mode network (a) and dorsal attention network**
**activation (b) and deactivate the default mode network (c).** For visualization, group contrast maps
(oddballs vs standards) are thresholded at $T = 3$, main text results report family-wise error corrected
statistics. Contrast maps are available at: osf.io/4t5hj. Numbers below the labeled regions indicate
MNI XYZ-coordinates. **(d-e)** Overlap of group-level activation **(d)** and deactivation **(e)** maps with
three published functional network atlases[23,58,59], quantified as Dice coefficients. Only networks with a
significant overlap based on permutation tests [24] and Dice coefficients > 0.2 are labeled. Note that the
*Action-mode network* has previously been termed *Cingulo-opercular network*[22] and the *Salience*
*network* is also termed *Ventral attention network*[24].

*Pupil-indexed neuromodulation is associated with action-mode network and brainstem activation*

By integrating pupil time series into the fMRI analyses, we directly probe the relationship between the detected cortical activation patterns and markers of neuromodulation. This approach allowed us to test which brain regions' activity correlated with moment-to-moment pupil fluctuations, which are likely driven by release of neuromodulatory neurotransmitters[53–55].

Supporting an association between noradrenergic neuromodulation and action-mode activation, network hubs such as bilateral anterior insula ($T = 7.31$, $p_{FWE-corr} < 0.001$ MNI: –32, 20, –8), middle cingulate cortex ($T = 5.55$, $p_{FWE-corr} < 0.001$ MNI: –4 –0 44), and thalamus ($T = 7.88$, $p_{FWE-corr} < 0.001$ MNI: 14 –12 10) scaled their activity with pupil size (Figure 4). The un/thresholded whole-brain contrast maps are available at osf.io/4t5hj. These findings corroborate recent observations—linking human intracranial insula recordings with pupil dynamics[30]—and extend them to a functional network level.

Importantly, these pupil–fMRI analyses also revealed a bilateral brainstem–midbrain cluster ($T = 7.27$, $p_{FWE-corr} < 0.001$ MNI: –4 –30 –14), overlapping with key noradrenergic, serotonergic, and dopaminergic nuclei, in accordance with recent work in rodents[55,60,61] and primates[62–64].

In sum, we found that neuromodulatory and action-mode network regions activated when pupils dilated, suggesting neuromodulatory drive of network transitions.

**Correlation with pupil size**

anterior insula
(–4 –30 –14)

brainstem–midbrain
(–7, –30, –14)

brainstem–midbrain
(–7, –32, –14)

*Figure 4.* **Moment-to-moment changes in pupil size correlate with action-mode network (a) and neuromodulatory brainstem–midbrain activation (b–c).** For visualization, group contrast maps are thresholded at $T = 3$, main text results report family-wise error corrected statistics. Contrast maps are available at: osf.io/4t5hj. Numbers below the labeled regions indicate MNI XYZ-coordinates.

*Noradrenergic transporter distribution overlaps with pupil-linked BOLD activation patterns*

If indeed the observed activation pattern is supported by noradrenergic release in the cortex, one would expect overlapping spatial distributions with the noradrenergic system. To test this, we leveraged public PET-derived maps of the cortical and subcortical distribution of the noradrenergic transporter[56,65–69]. We then assessed the voxel-wise association of noradrenergic transporter expression and pupil-linked BOLD activation. We detected a robust correlation (Hesse et al. map: $r = 0.189$; $p_{permutation-corr} < 0.001$), that we replicated using an

alternative noradrenergic transporter map (Ding et al. map: $r = 0.161$; $p_{\text{permutation-corr}} < 0.001$; Figure 5). This suggests that those parts of the brain that activated when pupils dilated overlapped with those housing noradrenergic receptors.

To further contextualize these results, we computed two negative control analyses. Specifically, we linked two brain-wide $GABA_A$ receptor distribution maps[70,71] to our pupil-linked BOLD maps, which we assumed to show no or even a negative association[72]. In line with this, we found that $GABA_A$ receptor expression showed a non-significant negative relation with the pupil–fMRI pattern (Nørgaard et al map: $r = -0.028$; $p_{\text{permutation-corr}} = 0.063$; Dukart et al. map: $r = -0.045$; $p_{\text{permutation-corr}} = 0.056$; Figure 5).

Finally, given that large parts of the brainstem–midbrain, overlapping with noradrenergic, but also serotonergic and dopaminergic nuclei, activated when pupils dilated, we ran two additional analyses to determine whether similar effects extended to other neuromodulatory systems. Both serotonergic (Beliveau et al. map: $r = 0.229$; $p_{\text{permutation-corr}} < 0.001$;[73]) and dopaminergic (Sasaki et al. map: $r = 0.241$; $p_{\text{permutation-corr}} < 0.001$;[74]) transporter expression showed a comparable spatial correspondence with the pupil–fMRI activation pattern. While the cortical innervation of dopaminergic, serotonergic and noradrenergic nuclei is partially overlapping, there are also considerable differences between neuromodulatory systems[3]. To test if the expression pattern of noradrenergic transporters explains unique variance in the observed pupil-linked BOLD maps over and above the other neuromodulatory systems, we estimated multiple regression analyses. Specifically, we compared a base model (including serotonergic and dopaminergic transporter expression as predictors) to a full model additionally including a predictor for the noradrenergic transporter (Table 1). Importantly, we found the full model substantially outperformed the base model, suggesting a unique association of noradrenergic transporter expression and pupil-linked BOLD activation ($\Delta\chi^2(df = 1) = 42909.5$; $p_{\text{permutation-corr}} < 0.001$).

**Table 1:** Multiple regression predicting pupil-linked BOLD activation based on neuromodulatory transporter maps

| Predictor | Estimate | Standard error | *t*-value | *p* |
|---|---|---|---|---|
| NAT | 0.18997 | 0.0009112 | 208.48 | <0.001 |
| 5HTT | 0.084247 | 0.00087668 | 96.098 | <0.001 |
| DAT | 0.20462 | 0.00081556 | 250.89 | <0.001 |

*Note*: NAT, 5HTT, DAT refer to PET-derived noradrenergic, serotonergic and dopaminergic transporter expression maps. To obtain a single noradrenergic predictor, the Hesse et al. and Ding et al. maps were averaged.

Taken together, these analyses suggest that key ascending arousal-promoting systems, including noradrenergic neuromodulation, contribute to the observed pupil-linked subcortical and action-mode network activation.

*Figure 5.* **Voxel-wise association of pupil-linked BOLD activation patterns with PET-derived maps of noradrenergic transporter (a–b) and GABAergic receptor (c–d) distribution**. (**a**) Relative distribution of noradrenergic transporters based on[69]. (**b**) Brain regions activating when pupils dilate overlap with noradrenergic transporter distribution based on two alternative PET maps. (**c**) Relative distribution of $GABA_A$ receptors based on[70]. (**d**) Brain regions activating when pupils dilate show no systematic overlap with GABAergic receptor distribution based on two alternative PET maps. Observed correlation coefficients are tested against a reference distribution obtained using phase-randomization permutation.

*Locus coeruleus is functionally connected to action-mode network hubs*

So far, our analyses suggest that pupil dilation, likely reflecting neuromodulatory drive, is associated with brainstem and action-mode-network activations. Moreover, the

activated regions closely overlap with noradrenergic receptor distributions as derived from
PET atlases. These observations prompted us to test if locus coeruleus and action-mode
network activation were directly connected.
Using dedicated neuromodulation-sensitive MRI[49], we estimated person-level proxies
for locus coeruleus integrity as well as a group image that visualizes the spatial location of the
locus coeruleus adjacent to the fourth ventricle (Figure 6 and osf.io/4t5hj). Notably, the locus
coeruleus-related hyperintensity was adequately captured by a previously generated high-
confidence mask of this structure ([75]; available at: https://osf.io/sf2ky/).
After this validation step, we used the derived coordinates overlapping with the
masked region as seed region for trial-level whole-brain functional connectivity analyses,
probing which region's activity co-fluctuated with the locus coeruleus during the conditioned
oddball task[76,77]. A bilateral anterior insula ($T = 8.93$, $p_{\text{FWE-corr}} < 0.001$ MNI: $-38$ $12$ $-10$;
Figure 6) and thalamus ($T = 6.79$, $p_{\text{FWE-corr}} < 0.001$ MNI: $-6.5$ $-22.5$ $-2.5$) cluster survived
correction for multiple comparisons, in addition to a broad cerebellar and pontine activation.
The un/thresholded whole-brain contrast maps are available at osf.io/4t5hj. These results
point to a co-fluctuation of activity in the locus coeruleus and key action-mode network hubs,
in line with known direct anatomical projections[9], as well as our pupil–fMRI and
noradrenergic transporter findings.
However, the functional connectivity analyses do not speak to the directionality of this
association (brainstem→action-mode network vs. action-mode network→brainstem). Thus,
we ran a separate set of analyses, using the TR-level BOLD timeseries of the locus coeruleus
and anterior insula (i.e., on a sample level, averaged over voxels). We submitted these to a
cross-correlation analysis, which measures the temporal relationship between two signals by
quantifying how one variable aligns with shifts in another over time[78], allowing us to assess
the temporal precedence of brainstem vs. action-mode network activation. First, we observed
a strong coinciding activation (mean cross-correlation coefficient at lag zero: 0.363; $T =$
11.892), replicating our functional connectivity findings, that stood out above the general
association of the two signals (mean cross-correlation coefficient across all lags: 0.217). Note
that to better judge the temporal relation of activations, we removed this overall background
association by demeaning before group-level cluster-statistics (Figure 6).
Adding to our connectivity analyses, we found that anterior insula activations
preceded and predicted subsequent locus coeruleus activations by up to four seconds (mean
cross-correlation coefficient at lag $-4$ s: 0.264; $T = 4.69$; at lag $-2$ s: 0.255; $T = 2.896$; cluster
permutation test: $T_{\text{sum}} = 22.103$; $p_{\text{cluster-corr}} \leq 0.002$; cluster including lags: $(-4)–(+2)$ s). We
also observed that preceding locus coeruleus activity was followed by action mode network
activation two seconds later (mean cross-correlation coefficient at lag $+2$ s: 0.25; $T = 2.626$),
suggesting a potentially slightly later recurrent connection (brainstem↔action-mode network;
contrast of cross-correlation coefficient at lags $[-4$ to $-2$; anterior insula leading] vs $[+2$ to $+4$;
locus coeruleus]: $T = 2.421$; $p = 0.018$). While these analyses do not allow for causal
interpretations, they indicate that anterior insula activity may recruit noradrenergic
neuromodulation to facilitate subsequent action-mode network activation[5,6]. Taken together,
we observed that brainstem activation was functionally connected with action-mode network
activity and that this crosstalk may be initiated by the anterior insula, potentially helping
participants to efficiently respond to unexpected stimuli[2].
295

*Figure 6.* **Locus coeruleus MRI (a) guides whole-brain functional connectivity analyses (b–c), revealing brainstem activation is coupled to and precedes action-mode network activity (d–e).** (**a**) Neuromodulation-sensitive MRI reveals locus coeruleus-related hyperintensity bordering the fourth ventricle that is accurately captured by a high-confidence volume of interest (red;[75]). (**b**) Using this region as seed for whole-brain task-based functional connectivity shows locus coeruleus–anterior insula co-fluctuations. (**d–e**) Cross-correlations replicate the locus coeruleus–anterior insula association and suggest this is initiated by the anterior insula, showing a temporal precedence of up to 4 s (contrast of lags [(–4)–(–2)] vs [(+2)–(+4)]).

*Locus coeruleus and action-mode network support implicit learning of task structure*

To investigate the implications of locus coeruleus–action-mode network interactions during the conditioned oddball task, we analyzed behavioral data. As participants showed high overall accuracy (group mean >0.95), our analyses focused on reaction time dynamics. Participants entered the task without prior knowledge of its statistical structure, yet mean reaction times systematically varied with trial likelihood across the fMRI session (Figure 7a;

$F = 173.83$; $p < 0.001$; mean reaction times per condition: stay-standard: –0.437±0.021,
switch-to standard: 0.173±0.031, stay-oddball: 0.442±0.043, switch-to oddball: 0.602±0.040;
note that all reaction times are expressed in $Z$-values after log-transformation and within-
person standardization). This pattern suggests that participants prioritized responding quickly
to frequent trial types at the cost of slower responses to rare events. Supporting this, a strong
negative association of the reaction times for stay-standard and switch-to-oddball conditions
indicates a performance trade-off between these categories (Figure 7b; $r = -0.795$; $p < 0.001$).
We thus assumed that participants implicitly learned the statistical structure of the novel task
environment and adapted their behavior, which implies that this reaction time pattern should
emerge over the course of the experiment. Combining behavioral data from both repetitions of
the conditioned oddball task, we observed that early in the experiment, responses were
generally slow and did not differentiate between frequent and rare trials (Figure 7c). However,
as the experiment progressed, reaction times to common stimuli became faster, while rare
oddballs elicited increasingly slower responses. Accordingly, we found that the difference in
reaction times between switch-to-oddball and stay-standard categories increased with time,
suggesting implicit learning (Figure 7d; correlation of reaction-time difference and time bin
on a group level: $r = 0.509$; $p = 0.003$). This learning pattern detected on a group level was
also evident within participants (dependent samples $t$-test of within-participant time-bin beta
parameters against zero: $T = 3.194$, $p = 0.002$; mean beta parameter: 0.0097±0.00035; Figure
S1). Collectively, these results indicate that participants, initially unfamiliar with the task,
gradually internalized its structure and adjusted their behavior accordingly.
333

334
335    *Figure 7*. **Participant focus their behavior on more likely trial categories (a) at the expense of less**
336    **likely categories (b). This bias emerges across the course of the experiments (c) indicating**
337    **implicit learning of the task structure (d).** Day 1 indicates the pupillometry-EEG assessment, while
338    day 2 indicates the pupillometry-fMRI oddball assessment. (**a–b**) show day 2 data. (**c–d**) include data
339    of both days.

340            Finally, we combined imaging and behavioral data to probe if individual differences in
markers of the noradrenergic system and its cortical target regions explained implicit learning
performance. Specifically, we combined dedicated structural locus coeruleus imaging (Figure
6a and Figure 8a left), task-related pupil dilation (cf. Figure 2b) and locus coeruleus-related
anterior insula activation (cf. Figure 3a) in a multivariate partial least-squares correlation with
condition-wise reaction times (stay-standard; switch-to standard; stay-oddball; switch-to
oddball; Figure 7 and Figure 8a right). We identified one reliable latent component ($r = 0.488$;
$p_{permutation-corrected} = 0.001$). Participants with stronger task-related anterior insula activation
(loading onto latent variable: 0.752; bootstrap ratio: 3.402 [interpreted analogous to $Z$-
values]), higher locus coeruleus integrity (loading: 0.491; bootstrap ratio: 2.306) and task-
related pupil dilation (loading: 0.44; bootstrap ratio: 1.873; trend level association) showed
more pronounced differentiation in reaction times between frequent and rare stimuli (cf.
Figure 8b). In particular, these participants tended to show quicker responses for frequent trial
categories (loadings; stay-standard: –0.468; switch-to-standard: –0.396) at the expense of

slower responses for infrequent trial categories (loadings; stay-oddball: 0.364, switch-to-
oddball: 0.701)—recapitulating the implicit-learning pattern we have identified in behavioral
analyses (Figure 7c). Post-hoc age group comparisons revealed that younger and older adults
showed comparable latent behavioral ($T = -1.36$; $p = 0.179$) and neural scores ($T = -0.209$; $p$
$= 0.835$) and that the latent association did not differ across groups ($Z = 1.111$; $p = 0.133$).
As a sensitivity analysis, we replicated partial least squares-correlation findings using
a simpler analytical strategy ($r = 0.466$; $p < 0.001$; by averaging over standardized neural
indicators (i.e., insula BOLD activity, locus coeruleus integrity, and pupil dilation, see PLSC
analyses) and correlating this composite score with the difference in reaction times to oddball
and standard stimuli).
In summary, participants adapted their responses over time to reflect the statistical
regularities of the experimental task, demonstrating implicit learning. This implicit learning
pattern was stronger expressed in those participants with higher insula activation, higher
structural locus coeruleus integrity, and a stronger pupil response to salient stimuli. Together
those findings collectively support a role of the locus coeruleus-noradrenergic system in
guiding attention and learning to adapt behavior in novel environments[3,6,41].
370

371
*Figure 8.* **Individual differences in noradrenergic markers (insula BOLD activity, locus
coeruleus integrity, pupil dilation) are associated with implicit learning.** (**a**) Multivariate partial
least squares correlation links noradrenergic markers to reaction times for frequent and infrequent trial
categories (standards, oddballs, respectively), which reflect that participants implicitly learned the
statistical regularities of the task. (**b**) Noradrenergic markers positively contribute to the latent brain
variable (e.g., the more activation, the higher scores). On a behavioral level, quicker reaction times for
frequent trial categories at the expense of slower reaction times for rare categories are linked to higher
latent behavioral scores. (**c**) Participants with higher values in markers of the noradrenergic system
demonstrate more implicit learning, a pattern that is comparable across age groups.

381
## 3.    Discussion
This study investigated the role of the locus coeruleus and its cortical targets in how
younger and older participants adapt their behavior in dynamic environments with novel
observations. We took advantage of repeated assessments of an experimental task shown to
modulate noradrenergic activity in primates[44] and a dedicated multimodal imaging protocol to
identify neuromodulatory contributions to implicit learning.
We found that salient expectation-violating stimuli dilated participants' pupils,
indicating heightened neuromodulatory activity, and triggered a shift from default-mode to
action-mode network activation. Combining pupil and BOLD data, we observed that
neuromodulatory brainstem–midbrain as well as action-mode network regions scaled their

activity with pupil size. This suggests an interplay between the noradrenergic system and the
action-mode network, which we confirmed using PET-derived transporter maps and locus
coeruleus-centered functional connectivity analyses. Behavioral data showed that participants
implicitly learned the statistical structure of the novel task environment and focused their
behavior on more prevalent trial categories, resulting in quicker responses at the expense of
slower responses to rare oddballs. Linking brain and behavior, we found that this implicit
learning pattern was more pronounced in participants with higher levels of markers associated
with the noradrenergic system and its cortical targets. Taken together, this study suggests
noradrenergic neuromodulation may upregulate the action-mode network to facilitate
(implicit) learning for behavioral adaptation.
402
403    *Fear conditioned stimuli evoke robust neuromodulatory responses*
404            To increase recruitment of the locus coeruleus, a subset of our experimental stimuli
was pre-conditioned before the main task by repeatedly pairing them with an aversive
outcome[40,44–46]. Conditioned stimuli (CS+) elicited greater pupil dilation than perceptually
matched control stimuli (CS–), suggesting increased neuromodulatory drive. Moreover, these
effects remained stable across age groups and assessment days, highlighting the robustness of
the conditioned response. Our findings are in line with earlier work in rodents demonstrating
increased locus coeruleus spiking[79] and noradrenergic axonal activation[45] during conditioning,
assumed to support fear learning.
412
413    *Noradrenergic neuromodulatory activation is coupled with the action-mode network*
414            Building on this, during the conditioned oddball task, salient stimuli dilated
participants' pupils and activated the action-mode network, including the anterior insula,
middle cingulate cortex, thalamus, and cerebellum. This aligns with prior findings implicating
the brain's action-mode of function with states of externally focused attention, heightened
arousal, and the processing of action-relevant bottom up signals[22,25]. Importantly, these
activations were accompanied by default-mode network deactivation, suggesting a functional
transition from introspective to externally oriented processing[6]. This is in line with the
proposed antagonism of the default-mode and action-mode networks[22] and the more general
noradrenergic role in facilitating dynamic reorganizations of target neural networks (also
termed network reset[5]).
424            Combined analyses of pupil and BOLD time series revealed that moment-to-moment
fluctuations in pupil size systematically co-varied with activation in the action-mode network
(anterior insula, middle cingulate cortex, thalamus). This supports previous intracranial work
in humans linking anterior insula activations to variations in pupil size[30] and extends them to
a network level. In addition, a brainstem–midbrain cluster encompassing the locus coeruleus
and other neuromodulatory nuclei activated when pupils dilated, in line with recent animal
research[55,60–62], suggesting that pupil dilation reflects broad neuromodulatory engagement
across cortical and subcortical regions[63,64].
432            An analysis of the spatial overlap between publicly available PET-derived
noradrenergic transporter maps and pupil-linked BOLD activation further underscores a role
for noradrenergic neuromodulation in regulating these activation patterns[56,65–69]. We found a
robust positive correlation, replicating across independent transporter maps, suggesting that
brain regions with increased activity during pupil dilation overlap with areas rich in
noradrenergic transporters. Negative control analyses with GABAergic receptor maps[70,71]
showed no such association, further strengthening the specificity of the association between
pupil size, locus coeruleus and action-mode network activation. In line with the broad
brainstem–midbrain activation, also serotonergic[73] and dopaminergic[74] transporter maps were
positively coupled with the pupil-linked BOLD pattern[63,64]. Crucially, however, noradrenergic

transporter expression explained unique variance in pupil-linked BOLD activation over and
above the other neuromodulatory systems.
To more precisely map locus coeruleus activation to its cortical targets, we employed
functional connectivity analyses guided by dedicated neuromodulation-sensitive MRI[75].
These analyses revealed that locus coeruleus activity co-fluctuated with action-mode network
hubs, particularly the anterior insula and thalamus, consistent with established direct
projections in primates[9]. Cross-correlation analyses further elucidated the temporal dynamics
of this relationship, suggesting that the anterior insula may recruit locus coeruleus activation
up to four seconds later. This finding together with evidence of a recurrent connection support
the notion that the insula-initiated recruitment of noradrenergic signaling may facilitate a
cortical shift toward an action-oriented state, optimizing adaptive responses to environmental
changes[5,6,22]. In this circuit, the anterior insula likely provides crucial contextual signals
without which the locus coeruleus would be blind to salient events that require gain-
modulation[2]. Our findings are in broad agreement with a recent rodent study in which direct
stimulation of the cingulate cortex, another action-mode network hub region, elicited pupil
dilation through locus coeruleus activation, suggesting top-down recruitment of brainstem
neuromodulation[80]. However, we cannot yet rule out the possibility that differences in the
shape of subcortical and cortical hemodynamic responses may contribute to the observed
effects, though previous findings do not point in this direction[81].
Notably, beyond the action-mode network, also the salience network includes hub
regions in the anterior insula and cingulate cortex[82], and in some atlases networks with these
names are overlapping[24]. Similar to the action-mode network, salience network activity has
been linked to ascending neuromodulatory systems, including noradrenaline[83]. While our
study cannot conclusively disentangle the spatially adjacent and functionally linked networks,
the more posterior cingulate and more dorsal anterior insula activation favors an action-mode
network involvement ([22,82], see Figure 3).
468
*Locus coeruleus and action-mode network together support implicit learning*
Finally, we examined behavioral adaptations over the course of the experiments.
Participants, initially unfamiliar with the task structure, gradually optimized their reaction
times to reflect stimulus probabilities, indicating implicit learning. Optimized reaction times
may arise from preparatory movement initiation, resulting in quicker responses for expected
stimuli, but slower responses for unexpected stimuli. Crucially, individuals with greater MRI-
indexed locus coeruleus integrity, stronger task-related anterior insula activation, and more
pronounced pupil dilation showed greater differentiation in reaction times between frequent
and infrequent stimuli. This suggests that the locus coeruleus and its cortical targets play a
key role in learning and adapting to novel environments, matching previous findings from
rodents demonstrating that noradrenergic activity is causal for task execution *and*
optimization[84]. In addition, our observations are in line with a recent study in humans relating
low MRI-indexed locus coeruleus integrity to diminished practice effects[85]. These result
moreover support and extend earlier work using *explicit* cognitive assessments to link
noradrenergic markers to attention[38,40], learning, and memory[41,43].
Post-mortem research identified the locus coeruleus as one of the starting points of
Alzheimer's-related tau accumulation[36], which may dysregulate noradrenergic activity with
increasing age[31,34] and impair late-life cognition[13,33]. Here we found that a latent measure
reflecting locus coeruleus integrity as well as pupil dilation and insula activation to salient
stimuli did not differ between younger and older adults. This may indicate that externally-
driven noradrenergic responses may be comparatively less impaired in aging[38,86]. In addition,
we found that this noradrenergic latent variable similarly explained implicit learning across
age groups, highlighting its behavioral relevance over the adult lifespan.
492

*Conclusions*
By combining neuromodulation-sensitive MRI, concurrent pupillometry–fMRI, and
PET-derived neuromodulatory transporter maps, our study suggests the noradrenergic locus
coeruleus may contribute to a transition from default-mode to action-mode network
activation. This mechanism may support adapting behavior to novel environments,
reinforcing the importance of neuromodulatory systems in attention, learning and memory.
499

**4.        Material and methods**

*Study design*
Data were collected as part of a larger project investigating neural and cognitive
correlates of age-related differences in the noradrenergic system (for details, see[38]).
Participants attended three consecutive days of lab-based testing, with the conditioned oddball
task administered on the second and third days (Figure 1). On the second day, concurrent
pupillometry and electroencephalography (EEG) data were recorded during task performance,
while third-day assessments incorporated MRI and pupillometry measurements. To enhance
the salience of oddball stimuli, each assessment day included a preceding fear conditioning
session, during which the oddball stimuli were repeatedly paired with aversive electrical
stimulation. The institutional review board of the German Psychological Association
approved the study protocol.

*Participants*
The final sample included 39 younger adults (YAs; mean age ± standard deviation:
25.23 ± 3.23 years; range: 20.17–31 years) and 38 older adults (OAs; mean age: 70.61 ± 2.71
years; range: 65.50–75.92 years), all of whom were male. Two younger adults were excluded
from the study after the first assessment day due to low-quality pupil data. Six participants
(three per age group) did not participate in the fMRI part of the study (n = 4) or had
incomplete eye tracking data (n = 2), leaving a sample of n = 71 for fMRI–pupillometry
analyses. All participants were healthy, right-handed, MRI-compatible, fluent German
speakers with normal or corrected-to-normal vision, for further characteristics, see[38]. All
participants provided written informed consent and were reimbursed for their participation.
Exclusion criteria included the use of centrally active drugs, particularly medications affecting
the noradrenergic system (e.g., beta-blockers).

*Experimental procedures and stimuli*
*Fear conditioning*
Participants underwent fear conditioning on each testing day to increase the salience of
experimental stimuli and their likelihood of eliciting locus coeruleus activation [40,44–46]. During
this phase, a sinusoidal luminance pattern (Gabor patch; CS+) with either horizontal or
vertical orientation (0° or 90°) was paired with an aversive electrical shock (US) on 80% of
trials. A perceptually matched control stimulus (CS–) with the alternate orientation (vertical
or horizontal) was never paired with the shock. Stimulus orientation–reinforcement pairings
remained consistent for each participant across days and were counterbalanced within age
groups (younger adults [YA]: 21:18; older adults [OA]: 20:18).
Each conditioning session comprised 40 trials, with 20 presentations each of CS+ and
CS– in pseudorandomized order across assessment days. Trials began with a one-second
fixation cross, followed by a two-second visual stimulus (CS+ or CS–). Shocks were
delivered for 0.5 seconds immediately after CS+ offset via a bipolar current stimulator (DS5;
Digitimer) connected to a ring electrode affixed to participant's left or right index finger.
Hand assignment was counterbalanced within age groups (YA: 19:20; OA: 19:19). Each trial
concluded with a six-second inter-trial interval, allowing pupil responses to return to baseline.
Participants selected an individually calibrated stimulation intensity deemed
unpleasant but not painful before each session[38]. Throughout all conditioning sessions, pupil
dilation and gaze position were recorded, and external distractions were minimized.
A procedural error occurred for one older participant, resulting in a reversal of
stimulus orientation–shock assignment between the first and second assessment days. This
error reduced the likelihood of detecting differences between conditions (CS+/CS–; i.e.,

worked against our hypothesis), but the participant's data were retained for analysis to
maintain a larger sample size.
552
*Conditioned oddball task*
On the second and third assessment days (EEG–pupillometry and fMRI–pupillometry,
respectively), participants completed a modified version of a conditioned oddball task
previously shown to increase locus coeruleus activity[44].
Specifically, on each trial participants viewed a Gabor patch in one of three possible
orientations (horizontal, diagonal, vertical; [0°, 45°, 90°]). Two of these orientations (0°, 90°)
were extensively familiarized during preceding fear conditioning (see above) and other tasks
within the larger project (see[38]), whereas the third orientation (45°) benefited from relative
novelty, hypothesized to enhance neuromodulatory activity[87].
During the conditioned oddball task, one of the stimulus orientations was presented
frequently (standard trials; 0° or 90°), while the two other orientations were rarely presented
(oddballs). In total, the task consisted of 160 trials presented in pseudorandomized order that
differed over assessment days, with the standard, non-conditioned stimulus (CS–) appearing
on 70% of trials (112 trials), while the two oddball stimuli (CS+ and the novel 45°
orientation) appeared less frequently, each comprising 15% of trials (24 each).
Gabor patches (0°, 45°, and 90° orientation) were displayed for two seconds followed
by an inter-trial interval ranging from 2–12 seconds (mean 3.625 seconds; skewed distribution
generated with OptSeq2, http://surfer.nmr.mgh.harvard.edu/optseq/).
During Gabor patch presentation, participants were instructed to respond as quickly
and accurately as possible by pressing a button corresponding to the orientation of each
stimulus using the hand not used for fear conditioning (see above). Accuracy and reaction
times were recorded for each trial.
575
*Physiological data recording and preprocessing*
*Eye tracking*
During all experiments, participants' pupil dilation was measured as an indirect
marker of central neuromodulatory activity[53–55] alongside gaze position, using a video-based
infrared eye tracker (SR Research EyeLink 1000). On the second day of assessments (during
EEG), the desktop mount configuration was used, while on the third day (during fMRI), the
long-range mount was employed. The systems operated in a monocular configuration with a
spatial resolution of up to 0.25° and a sampling rate of 1000 Hz.
To minimize head movements, participants were seated with their forehead and chin
stabilized at a fixed distance of 53.5 cm from the display during the second day. For the MRI
session (third day), the eye tracker was positioned within the scanner bore, and data were
collected via a mirror affixed to the head coil, which redirected participants' view to the
stimulus display.
Participants were instructed to maintain central fixation throughout the experiments.
Before each experiment, the eye tracker was (re)calibrated using a standard 5-point grid.
Calibration was considered successful if fixation errors were under 0.5°.
Eye tracking data were preprocessed and missing samples were imputed using
published standardized routines[88]. The percentage of imputed samples was logged as an
objective quality measure for potential subsequent participant exclusion (proportion of valid
eye tracking samples during the fMRI–pupillometry assessment (mean ± standard error):
$0.803 \pm 0.024$).
597
*Magnetic resonance imaging*
MRI data were collected using a 3T Magnetom TIM Trio (Siemens Healthcare) with a
12-channel head coil.

An axial, T1-weighted neuromodulation-sensitive Fast Spin Echo (FSE) sequence was
collected to assess markers for locus coeruleus integrity. The following parameters were
used[41,89]: acquisition matrix: $440 \times 512$, 10 slices, voxel size: $0.5 \times 0.5 \times 2.5$ mm, 20 % gap
between slices, repetition time (TR): 600 ms, echo time (TE): 11 ms, flip angle (FA): 120 °,
acquisition time: 11:48 min. To increase signal to noise ratios in brainstem imaging, the
sequence included four online averages and yielded two images, extracted locus coeruleus
parameters were averaged to boost stability of estimates[48].

In addition, a sagittal T1-weighted Magnetization-Prepared Gradient Echo
(MPRAGE) sequence with the following parameters was collected to facilitate co-registration
to standard space: acquisition matrix: $256 \times 256 \times 192$, voxel size: 1 mm isotropic, repetition
time (TR): 2500 ms, echo time (TE): 4.77 ms, inversion time (TI): 1100 ms, flip angle (FA):
10 °, acquisition time: 9:20 min.

Finally, a T2*-weighted whole-brain Echo Planar Imaging (EPI) sequence was used to
track blood oxygenation level dependent (BOLD)-changes during the conditioned oddball
task. The following imaging parameters were used: acquisition matrix: $72 \times 72$, 36 slices, 450
images, voxel size: 3 mm isotropic, repetition time (TR): 2000 ms, echo time (TE): 30 ms,
flip angle (FA): 80 °, acquisition time: 15 min.

MPRAGE and EPI data were preprocessed using standardized routines as
implemented in HeuDiConv[90], MRIQC[91,92], and fMRIprep[93] and as detailed in the
supplementary information. Data were transformed to 2 mm isotropic MNI 152 non-linear
2009c asymmetric space and smoothed with an isotropic 8 mm full-width half-maximum
(FWHM) kernel for further analyses, following recommendations for smoothing kernels to be
at least twice the voxel resolution[94]. A time series of confounds was derived from head-
motion estimates, and global signals (within the white matter, the cerebrospinal fluid (CSF)
and whole brain mask). These confounds were expanded with the inclusion of temporal
derivatives and quadratic terms for each, yielding a total of 36 confounds that were included
in within-participant analyses[95,96].

*Physiological data analyses*
*Pupillometry analysis*
Pupil time series were analyzed using the EEGlab[97], eye-EEG[98], and fieldtrip[99]
toolboxes. The continuous pupil time series were segmented into trials surrounding
presentation of the Gabor patches (for conditioning and oddball experiments). Within-
participant trial-level analyses then contrasted pupil responses across experimental conditions
after baseline correction (–500–0 ms relative to stimulus onset). That is, for conditioning
experiments, CS+ and CS– trials were contrasted, while for the oddball experiments standard
and oddball trials were compared (i.e., collapsing the two oddball categories) using sample-
wise independent samples $t$-tests. Within-participant analyses were repeated over $n_{Bootstraps} =$
50 and the resulting $t$-value time series were averaged[38]. Subsequent across-participant
analyses tested for the consistency of within-participant effects on a group level (dependent
samples $t$-test against zero), while controlling for multiple comparisons using cluster-based
random permutation tests[100]. For more details, see[38].

*Functional magnetic resonance imaging analysis*
fMRI data were analyzed using the Statistical Parametric Mapping (SPM) 12
toolbox[101] to identify brain responses to oddball stimuli. Within-participant event-related
analyses included stick functions for each stimulus onset (standards, oddballs) convolved with
a canonical hemodynamic response function as well as its temporal and dispersion
derivatives[102]. General linear models contrasted BOLD responses to standard and oddball
stimuli (aggregating over the two oddball categories to increase power). Subsequent across-

participant analyses evaluated the consistency of within-participants effects on a group level
with family-wise error correction for multiple comparisons.
Overlap of observed group-level effects with several published functional network
atlases (see Figure 3,[23,58,59]) was evaluated using Dice coefficients and permutation tests, as
implemented in the Network Correspondence Toolbox[24]. For overlap estimations, statistical
maps were thresholded at $T = 3$.
Follow-up within-participant functional connectivity analyses first calculated trial-by-
trial activation patterns for each voxel[76] and next assessed which brain regions' time series
correlated with the locus coeruleus[77]. For this, a locus coeruleus consensus mask was applied
as seed region[75] that reliably segmented the locus coeruleus-related hyperintensity in this
study, as shown in the results. That is, our functional connectivity analyses were centered on
the region in which we localized the locus coeruleus using dedicated high-resolution
structural scans.
We detected a reliable association of locus coeruleus and anterior insula time series. To
further characterize the temporal properties of this association, we estimated cross-
correlations between TR-level locus coeruleus and insula time series on a participant level
(using *crosscorr* in *Matlab*;[30]). To this end, we averaged BOLD signals across voxels within
the locus coeruleus consensus mask[75], as well as anterior insula functional-connectivity
cluster. Cluster-based random permutation tests probed the consistency of the participant-level
cross-correlation estimates on a group level after de-meaning[100].
671
*Structural magnetic resonance imaging analysis*
Neuromodulation-sensitive FSE images were analyzed using a previously established
semi-automatic procedure to extract proxies for locus coeruleus integrity[41,75,89] using
Advanced Normalization Tools (version 2.3.3.; ANTs[103,104]).
In brief, first MPRAGE scans were linearly resampled to 0.5 mm isotropic resolution
before estimating a whole-brain group template (six iterations;
*antsMultivariateTemplateConstruction,* including *N4BiasFieldCorrection*). Next, native-
space FSE scans were non-linearly aligned to within-person template-space MPRAGE images
using *antsRegistrationSyNQuick* to facilitate subsequent brainstem template construction
based on the outputs of this transformation (six iterations;
*antsMultivariateTemplateConstruction,* including *N4BiasFieldCorrection*). Finally, the
brainstem template was moved to 0.5 mm linear MNI space using the whole-brain template as
an intermediate target (*antsRegistration,* facilitated by a custom coregistration mask). All
transformation matrices were then concatenated and applied to individual neuromodulation-
sensitive FSE scans, bringing them directly from native to MNI space in a single step
(*antsApplyTransforms*).
To extract proxies for locus coeruleus integrity, standard space FSE images were
masked using a high-confidence locus coeruleus consensus mask[75] that captured the observed
locus coeruleus hyperintensity well (see Figure 6). To normalize intensity values for across-
participant analyses, scans were also masked with a pontine reference area mask[75]. The peak
voxel intensity for each slice of each region of interest was then automatically identified.
Finally, intensity ratios for each slice of the locus coeruleus were computed as a measure of
structural integrity (see Equation 1). For each slice, these ratios were derived by dividing the
difference between the peak intensity in the locus coeruleus and the corresponding reference
region by the peak intensity of the reference region, following previously established
methods[41,75,89].
698
699 **Equation 1:** $$LocusCoeruleusRatio = \frac{\max(LocusCoeruleus) - \max(Reference)}{\max(Reference)}$$
700

For subsequent analyses, extracted slice-wise values were averaged over hemispheres and then the peak value across slices was selected, while excluding the two most rostral slices to avoid unreliable intensity values at the edge of the acquisition[41].

*Behavioral data analysis*

In the conditioned oddball task, participants were instructed to press a specific button corresponding to the orientation of the presented Gabor stimulus. The majority of trials (70%) featured the standard stimulus, creating a high probability that a standard stimulus would follow another standard stimulus. We assumed that due to this consistency participants would build up a tendency (bias) to respond with the corresponding key. We termed these *stay-standard* trials. Occasionally, this expectation was violated by the presentation of one of the oddball stimuli, requiring participants to switch response keys (*switch-to-oddball trials*). Following an oddball stimulus, most trials returned to the standard stimulus (*switch-to-standard*), while less frequently, another oddball appeared (*stay-oddball*). Following this reasoning, we estimated mean reaction times for each of these four trial categories for each participant (2 (stay/switch) × 2 (standard/oddball)) after log-transformation and Z-standardization of reaction times across trials.

Crucially, participants began the task without prior knowledge of the statistical structure or trial distribution. The extent to which participants adjusted their behavior over time in response to the predictability of the stimuli is thus interpreted as an implicit measure of (task structure) learning. To quantify if participant indeed learned the statistical structure of the task, we down-sampled the trial-level reaction-time time series of the repeated oddball experiments (320 trials in total, two repetitions of 160 trials each; cf. Figure 7). That is, we averaged reaction times over 10 adjacent trials within the four trial categories, yielding 32 reaction-time time bins, which we contrasted across trial categories. Finally, we evaluated if participants over the course of the experiment developed a tendency to preferentially focus on one trial category at the expense of another. To this end, we probed if the contrast of reaction times (*switch-to-oddball – stay-standard*) increased over time. We repeated these analyses on a group level (correlation of group-averaged reaction time contrast and time bin) and on a single-participant level (linear regression predicting reaction time contrasts by time bin). Finally, participant-level beta estimates (quantifying implicit learning) were contrasted against zero using dependent-samples *t*-test to evaluate the consistency of implicit learning on a group level.

*Multimodal data analyses*
*Combined pupillometry–functional magnetic resonance imaging analyses*

Presentation of oddball stimuli elicited pupil dilation and distributed cortical activation, including the action-mode network. To test which brain regions' activation scaled with pupil-indexed neuromodulation, we estimated a separate set of general linear models. First, we down-sampled each participant's pupil time series to the fMRI temporal resolution (TR = 2 s) and added it as regressor to the participant-level models[105]. The same confounds as in unimodal fMRI analyses were applied (see above), while task regressors were omitted. Subsequent analyses across participants assessed the consistency of within-participant effects at the group level, applying family-wise error correction to account for multiple comparisons. These analyses were restricted to participants with high-quality pupil data (>50% valid non-imputed samples; 64/71 participants[88]).

*Association of pupillometry–functional magnetic resonance imaging patterns and PET-derived transporter maps*

Combined pupillometry–fMRI analyses showed that brainstem–midbrain and insula activation correlated with pupil size variations, suggesting neuromodulatory influences. To

more directly test this, we leveraged publicly available maps of neuromodulatory transporter
distributions[56]. PET-derived maps were resampled to the spatial resolution of our pupil–fMRI
activation patterns (2 mm MNI-ICBM 152 non-linear 2009 version c space). Next, we probed
the voxel-wise association of noradrenergic transporter expression and pupil-linked BOLD
activation using two alternative transporter maps using Pearson's correlations[56,65–69]. To judge
the reliability of these effects, we ran two negative control analyses, using two maps of
GABAergic receptor distribution acquired from two different sources[70,71] as well as two
comparative analyses, using maps of the dopaminergic [74] and serotonergic transporter[73]
distribution. For each of these, observed effects were contrasted to reference distributions of
correlation coefficients estimated with permuted pupillometry-fMRI data while preserving the
spatial autocorrelation of the observed data. Specifically, we applied a phase-randomization
approach, which retains the power spectrum of the original data while randomizing its phase
in the Fourier domain[106]. This method generates surrogate datasets that maintain the
autocorrelation properties of the original data (i.e., the pupil–fMRI activation map) while
breaking voxel-wise correspondences with the PET-derived transporter/receptor maps. We
computed correlation coefficients between the shuffled pupil-linked BOLD activation pattern
and the transporter/receptor maps across 10000 iterations, forming a null distribution, to
which we compared the observed correlation coefficients.
770            Finally, to determine if the voxel-wise noradrenergic transporter expression explained
variance in pupil-linked BOLD patterns (*PupilBOLD*), we used multiple regression analyses
(using *fitglm* in Matlab). Specifically, we compared a base model (*PupilBOLD ~ 1 + 5HTT +*
*DAT*) incorporating the serotonergic (*5HTT*) and dopaminergic (*DAT*) transporter expression
to a full model including all three neuromodulatory systems (*PupilBOLD ~ 1 + 5HTT + DAT*
*+ NAT*). To obtain a single noradrenergic transporter predictor (*NAT*), the two available maps
were averaged[65,69]. Differences in model fit between base and full models were evaluated
using a likelihood-ratio test and compared to a null distribution obtained from adding shuffled
noradrenergic transporter predictors (10000 iterations).
779
780    *Combined brain–behavior analyses*
781            Our physiological analyses yielded multiple locus coeruleus-related measures (MRI-
782    indexed locus coeruleus integrity; pupil-indexed neuromodulation; locus coeruleus-linked
783    action-mode network activation). Given the prominent involvement of the noradrenergic
784    system in learning and memory [3], we next tested if these explained individual differences in
785    implicit learning during the conditioned oddball task, using partial least squares correlations
786    (PLSC[107,108]).
787            In short, partial least squares correlation is a multivariate statistical technique used to
identify patterns of covariance between two sets of variables. In this study, behavioral partial
least squares correlation was applied to explore the relationship between locus coeruleus-
related measures and task-derived measures of implicit learning. By decomposing the
covariance structure with Singular Value Decomposition, partial least squares correlation
identifies latent variables (LVs) that maximize the shared variance between the datasets (i.e.,
locus coeruleus-related measures and implicit learning). Each latent variable is characterized
by a pair of weighted combinations of variables—one from each dataset—allowing for the
assessment of how strongly these patterns are expressed across participants. Statistical
significance of the latent variables was determined using permutation tests ($n_{permutations}$ =
10,000), and the reliability of individual variable contributions was evaluated with
bootstrapping ($n_{bootstraps}$ = 10,000). A ratio of a variable's weight and its bootstrapped standard
error, termed bootstrap ratio, (BSR), allows identifying indicators reliably contributing to the
latent variable ($|BSR| \geq 1.96$, interpreted analogous to Z-scores). Taken together, partial least
squares correlation provides a robust method for examining complex, high-dimensional

relationships, integrating locus coeruleus-related and learning-related measures. For further
details, see[75].
The following neural and behavioral variables were used as input to the partial least-
squares analyses: (1) individual task-related left anterior insula activations, extracted using
MarsBar[109], averaged over voxels of the cluster that showed reliable associations with pupil-
linked neuromodulation (Figure 4); (2) individual task-related pupil dilation, extracted using
the cluster shown in Figure 2 and averaged over samples; (3) peak locus coeruleus contrast
ratios, averaged over hemispheres; and (4–8) individual reaction times for each trial category,
2 (stay/switch) × 2 (standard/oddball), averaged over trials after log-transformation and $Z$-
standardization, as shown in Figure 7. All input variables were standardized across
individuals, and participants with absolute values >3 were dropped before analyses (n = 3).
Post-hoc independent samples $t$-tests probed age differences in the latent brain and behavioral
variables. As a sensitivity analysis, we replicated partial least squares-correlation findings
using a simpler analytical strategy that averaged over standardized neural indicators (i.e.,
calculated composite scores based on (1)-(3), see PLSC analyses), estimated the difference in
response times for oddball and standard trials, and then correlated these measures.
818
819

### Acknowledgements

MW-B received support from the German Research Foundation (DFG, WE 4269/5-1) and the Jacobs Foundation (Early Career Research Fellowship 2017–2019).

MM's work was supported by an Alexander von Humboldt fellowship and by National Institutes of Health grant R01AG025340 and R01AG080652.

During the work on this article, MJD was a member of the International Max Planck Research School on Computational Methods in Psychiatry and Ageing Research (IMPRS COMP2PSYCH, https://www.mps-uclcentre.mpg.de/comp2psych. Participating institutions: Max Planck Institute for Human Development, University College London). At the time of writing, TL was a PhD student in the Max Planck School of Cognition.

Acknowledgment is made to the donors of the ADR A2024006F, a program of the BrightFocus Foundation, for support of this research (MJD).

This work was in part conducted at the Max Planck Dahlem Campus of Cognition (MPDCC) of the Max Planck Institute for Human Development, Berlin, Germany.

### Author contributions

MJD, MM and MWB designed the study. MJD performed the experiments together with a team of research assistants. MJD and TL analyzed the data. MJD wrote the paper. All authors gave conceptual advice and revised the paper.

### Competing interests

The authors declare no competing interests.

### Data availability

Participants in this study did not consent to the public sharing of their data. To obtain access to participant-level data, please contact the corresponding author (MJD) with a signed data sharing agreement (template available at: https://osf.io/4t5hj/). Un/thresholded group-level data are available at https://osf.io/4t5hj/.

### Material and correspondence

Correspondence and requests for materials should be addressed to Martin J. Dahl.

**References**

1.  Uddin, L. Q. Cognitive and behavioural flexibility: neural mechanisms and clinical considerations. *Nat. Rev. Neurosci.* 1–13 (2021) doi:10.1038/s41583-021-00428-w.

2.  Mather, M., Clewett, D., Sakaki, M. & Harley, C. W. Norepinephrine ignites local hotspots of neuronal excitation: How arousal amplifies selectivity in perception and memory. *Behav. Brain Sci.* **39**, e200 (2016).

3.  Sara, S. J. The locus coeruleus and noradrenergic modulation of cognition. *Nat. Rev. Neurosci.* **10**, 211–223 (2009).

4.  Dahl, M. J., Mather, M. & Werkle-Bergner, M. Noradrenergic modulation of rhythmic neural activity shapes selective attention. *Trends Cogn. Sci.* **26**, 38–52 (2022).

5.  Bouret, S. & Sara, S. J. Network reset: A simplified overarching theory of locus coeruleus noradrenaline function. *Trends Neurosci.* **28**, 574–582 (2005).

6.  Corbetta, M., Patel, G. & Shulman, G. L. The reorienting system of the human brain: From environment to theory of mind. *Neuron* **58**, 306–324 (2008).

7.  Dayan, P. & Yu, A. J. Phasic norepinephrine: A neural interrupt signal for unexpected events. *Netw. Comput. Neural Syst.* **17**, 335–350 (2006).

8.  Nassar, M. R. Toward a computational role for locus coeruleus/norepinephrine arousal systems. *Curr. Opin. Behav. Sci.* **59**, 101407 (2024).

9.  Aston-Jones, G. & Cohen, J. D. An integrative theory of locus coeruleus-norepinephrine function: Adaptive gain and optimal performance. *Annu. Rev. Neurosci.* **28**, 403–450 (2005).

10. Thiele, A. & Bellgrove, M. A. Neuromodulation of attention. *Neuron* **97**, 769–785 (2018).

11. Servan-Schreiber, D., Printz, H. & Cohen, J. D. A network model of catecholamine effects: Gain, signal-to-noise ratio, and behavior. *Science (80-. ).* **249**, 892–5 (1990).

12. Ghosh, S. & Maunsell, J. H. R. Locus coeruleus norepinephrine contributes to visual-spatial attention by selectively enhancing perceptual sensitivity. *Neuron* **112**, 2231-2240.e5 (2024).

13. Dahl, M. J., Kulesza, A., Werkle-Bergner, M. & Mather, M. Declining locus coeruleus-dopaminergic and noradrenergic modulation of long-term memory in aging and Alzheimer's disease. *Neurosci. Biobehav. Rev.* 105358 (2023) doi:10.1016/J.NEUBIOREV.2023.105358.

14. Hagena, H., Hansen, N. & Manahan-Vaughan, D. β-adrenergic control of hippocampal function: Subserving the choreography of synaptic information storage and memory. *Cereb. Cortex* **26**, 1349–1364 (2016).

15. Eschenko, O. The role of the locus coeruleus in cellular and systems memory consolidation. in *Handbook of behavioral neuroscience: Handbook of in vivo neural plasticity techniques* (ed. Manahan-Vaughan, D.) vol. 28 327–347 (Elsevier, 2019).

16. Hagena, H. & Manahan-Vaughan, D. Oppositional and competitive instigation of hippocampal synaptic plasticity by the VTA and locus coeruleus. *Proc. Natl. Acad. Sci.* **122**, e2402356122 (2024).

17. Jordan, R. The locus coeruleus as a global model failure system. *Trends Neurosci.* **47**, 92–105 (2024).

18. Jepma, M. *et al.* Catecholaminergic Regulation of Learning Rate in a Dynamic Environment. *PLOS Comput. Biol.* **12**, e1005171 (2016).

19. Silvetti, M., Vassena, E., Abrahamse, E. & Verguts, T. Dorsal anterior cingulate-brainstem ensemble as a reinforcement meta-learner. *PLoS Comput. Biol.* **14**, e1006370 (2018).

20. Li, T., Marble, H., Chen, T., Razmi, N. & Nassar, M. R. Fluctuations in arousal reflect latent state transitions that facilitate behavioral optimization. *bioRxiv* 2025.02.06.636887 (2025) doi:10.1101/2025.02.06.636887.

21. Nassar, M. R., Wilson, R. C., Heasly, B. & Gold, J. I. An Approximately Bayesian Delta-Rule Model Explains the Dynamics of Belief Updating in a Changing Environment. *J. Neurosci.* **30**, 12366–12378 (2010).

22. Dosenbach, N. U. F., Raichle, M. E. & Gordon, E. M. The brain's action-mode network. *Nat. Rev. Neurosci. 2025* 1–11 (2025) doi:10.1038/s41583-024-00895-x.

23. Yeo, B. T. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J. Neurophysiol.* **106**, 1125–1165 (2011).

24. Kong, R. *et al.* A network correspondence toolbox for quantitative evaluation of novel neuroimaging results. *Nat. Commun. 2025 161* **16**, 1–16 (2025).

25.  Dosenbach, N. U. F., Fair, D. A., Cohen, A. L., Schlaggar, B. L. & Petersen, S. E. A dual-
networks architecture of top-down control. *Trends Cogn. Sci.* **12**, 99–105 (2008).

26.  Berridge, C. W. Noradrenergic modulation of arousal. *Brain Res. Rev.* **58**, 1–17 (2008).

27.  Carter, M. E. *et al.* Tuning arousal with optogenetic modulation of locus coeruleus neurons.
*Nat. Neurosci.* **13**, 1526–1535 (2010).

28.  Hayat, H. *et al.* Locus coeruleus norepinephrine activity mediates sensory-evoked awakenings
from sleep. *Sci. Adv.* **6**, eaaz4232 (2020).

29.  Osorio-Forero, A. *et al.* Infraslow noradrenergic locus coeruleus activity fluctuations are
gatekeepers of the NREM–REM sleep cycle. *Nat. Neurosci.* (2024) doi:10.1038/s41593-024-
01822-0.

30.  Kucyi, A. & Parvizi, J. Pupillary dynamics link spontaneous and task-evoked activations
recorded directly from human insula. *J. Neurosci.* **40**, 6207–6218 (2020).

31.  Kelberman, M. A. *et al.* Diversity of ancestral brainstem noradrenergic neurons across species
and multiple biological factors. *bioRxiv Prepr. Serv. Biol.* (2024)
doi:10.1101/2024.10.14.618224.

32.  Theofilas, P. *et al.* Locus coeruleus volume and cell population changes during Alzheimer's
disease progression: A stereological study in human postmortem brains with potential
implication for early-stage biomarker discovery. *Alzheimer's Dement.* **13**, 236–246 (2017).

33.  Mooraj, Z. *et al.* Toward a functional future for the cognitive neuroscience of human aging.
*Neuron* **113**, 154–183 (2025).

34.  Weinshenker, D. Long road to ruin: Noradrenergic dysfunction in neurodegenerative disease.
*Trends Neurosci.* **41**, 211–223 (2018).

35.  Mather, M. & Harley, C. W. The locus coeruleus: Essential for maintaining cognitive function
and the aging brain. *Trends Cogn. Sci.* **20**, 214–226 (2016).

36.  Braak, H., Thal, D. R., Ghebremedhin, E. & Del Tredici, K. Stages of the pathologic process in
Alzheimer disease: Age categories from 1 to 100 years. *J. Neuropathol. Exp. Neurol.* **70**, 960–
969 (2011).

37.  Ehrenberg, A. J. *et al.* Priorities for research on neuromodulatory subcortical systems in
Alzheimer's disease: Position paper from the NSS PIA of ISTAART. *Alzheimer's Dement.* **8**,
21 (2023).

38.  Dahl, M. J., Mather, M., Sander, M. C. & Werkle-Bergner, M. Noradrenergic responsiveness
supports selective attention across the adult lifespan. *J. Neurosci.* **40**, 4372–4390 (2020).

39.  Kosciessa, J. Q., Mayr, U., Lindenberger, U. & Garrett, D. D. Broadscale dampening of
uncertainty adjustment in the aging brain. *Nat. Commun. 2024 151* **15**, 1–18 (2024).

40.  Lee, T.-H. *et al.* Arousal increases neural gain via the locus coeruleus–noradrenaline system in
younger adults but not in older adults. *Nat. Hum. Behav.* **2**, 356–366 (2018).

41.  Dahl, M. J. *et al.* The integrity of dopaminergic and noradrenergic brain regions is associated
with different aspects of late-life memory performance. *Nat. Aging 2023* 1–16 (2023)
doi:10.1038/s43587-023-00469-z.

42.  Bueichekú, E. *et al.* Spatiotemporal patterns of locus coeruleus integrity predict cortical tau and
cognition. *Nat. Aging 2024* 1–13 (2024) doi:10.1038/s43587-024-00626-y.

43.  Jacobs, H. I. L. *et al.* In vivo and neuropathology data support locus coeruleus integrity as
indicator of Alzheimer's disease pathology and cognitive decline. *Sci. Transl. Med.* **13**,
eabj2511 (2021).

44.  Aston-Jones, G., Rajkowski, J., Kubiak, P. & Alexinsky, T. Locus coeruleus neurons in
monkey are selectively activated by attended cues in a vigilance task. *J. Neurosci.* **14**, 4467–
4480 (1994).

45.  Deitcher, Y., Leibner, Y., Kutzkel, S., Zylbermann, N. & London, M. Nonlinear relationship
between multimodal adrenergic responses and local dendritic activity in primary sensory
cortices. *bioRxiv* 814657 (2019) doi:10.1101/814657.

46.  Wilmot, J. H. *et al.* Phasic locus coeruleus activity enhances trace fear conditioning by
increasing dopamine release in the hippocampus. *Elife* **12**, (2023).

47.  Astafiev, S. V., Snyder, A. Z., Shulman, G. L. & Corbetta, M. Comment on 'Modafinil shifts
human locus coeruleus to low-tonic, high-phasic activity during functional MRI' and
'Homeostatic sleep pressure and responses to sustained attention in the suprachiasmatic area'.

*Science (80-. ).* **328**, 309 (2010).

48.  Forstmann, B. U., De Hollander, G., Van Maanen, L., Alkemade, A. & Keuken, M. C.
Towards a mechanistic understanding of the human subcortex. *Nat. Rev. Neurosci.* **18**, 57–65
(2016).

49.  Betts, M. J. *et al.* Locus coeruleus imaging as a biomarker for noradrenergic dysfunction in
neurodegenerative diseases. *Brain* **142**, 2558–2571 (2019).

50.  Pérot, J.-B. *et al.* Longitudinal neuromelanin changes in prodromal and early Parkinson's
disease in humans and rat model. *bioRxiv* 2024.10.22.619619 (2024)
doi:10.1101/2024.10.22.619619.

51.  Watanabe, T., Tan, Z., Wang, X., Martinez-Hernandez, A. & Frahm, J. Magnetic resonance
imaging of noradrenergic neurons. *Brain Struct. Funct.* **224**, 1609–1625 (2019).

52.  Keren, N. I. *et al.* Histologic validation of locus coeruleus MRI contrast in post-mortem tissue.
*Neuroimage* **113**, 235–245 (2015).

53.  Joshi, S. & Gold, J. I. Pupil size as a window on neural substrates of cognition. *Trends Cogn.*
*Sci.* **24**, 466–480 (2020).

54.  Grujic, N., Polania, R. & Burdakov, D. Neurobehavioral meaning of pupil size. *Neuron* **0**,
(2024).

55.  Privitera, M. *et al.* A complete pupillometry toolbox for real-time monitoring of locus
coeruleus activity in rodents. *Nat. Protoc.* **15**, 2301–2320 (2020).

56.  Hansen, J. Y. *et al.* Mapping neurotransmitter systems to the structural and functional
organization of the human neocortex. *Nat. Neurosci. 2022* 1–13 (2022) doi:10.1038/s41593-
022-01186-3.

57.  George, M. S. & Aston-Jones, G. Noninvasive techniques for probing neurocircuitry and
treating illness: vagus nerve stimulation (VNS), transcranial magnetic stimulation (TMS) and
transcranial direct current stimulation (tDCS). *Neuropsychopharmacol. 2010 351* **35**, 301–316
(2009).

58.  Gordon, E. M. *et al.* Precision Functional Mapping of Individual Human Brains. *Neuron* **95**,
791-807.e7 (2017).

59.  Ji, J. L. *et al.* Mapping the human brain's cortical-subcortical functional network organization.
*Neuroimage* **185**, 35–57 (2019).

60.  Cazettes, F., Reato, D., Morais, J. P., Renart, A. & Mainen, Z. F. Phasic activation of dorsal
raphe serotonergic neurons increases pupil size. *Curr. Biol.* **31**, 192–197.e4 (2020).

61.  Reimer, J. *et al.* Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in
cortex. *Nat. Commun.* **7**, 13289 (2016).

62.  Joshi, S., Li, Y., Kalwani, R. M. & Gold, J. I. Relationships between pupil diameter and
neuronal activity in the locus coeruleus, colliculi, and cingulate cortex. *Neuron* **89**, 221–234
(2016).

63.  Meissner, S. N. *et al.* Self-regulating arousal via pupil-based biofeedback. *Nat. Hum. Behav.*
*2023* 1–20 (2023) doi:10.1038/s41562-023-01729-z.

64.  de Gee, J. W. *et al.* Dynamic modulation of decision biases by brainstem arousal systems. *Elife*
**6**, 1–36 (2017).

65.  Hesse, S. *et al.* Central noradrenaline transporter availability in highly obese, non-depressed
individuals. *Eur. J. Nucl. Med. Mol. Imaging* **44**, 1056–1064 (2017).

66.  Belfort-Deaguiar, R. *et al.* Noradrenergic Activity in the Human Brain: A Mechanism
Supporting the Defense Against Hypoglycemia. *J. Clin. Endocrinol. Metab.* **103**, 2244–2252
(2018).

67.  Sanchez-Rangel, E. *et al.* Norepinephrine transporter availability in brown fat is reduced in
obesity: a human PET study with [11C] MRB. *Int. J. Obes. 2019 444* **44**, 964–967 (2019).

68.  Li, C. R. *et al.* Decreased norepinephrine transporter availability in obesity: Positron Emission
Tomography imaging with (S,S)-[11C]O-methylreboxetine. *Neuroimage* **86**, 306–310 (2014).

69.  Ding, Y. S. *et al.* PET imaging of the effects of age and cocaine on the norepinephrine
transporter in the human brain using (S,S)-[(11)C]O-methylreboxetine and HRRT. *Synapse* **64**,
30–38 (2010).

70.  Dukart, J. *et al.* Cerebral blood flow predicts differential neurotransmitter activity. *Sci. Reports*
*2018 81* **8**, 1–11 (2018).

71. Nørgaard, M. *et al.* A high-resolution in vivo atlas of the human brain's benzodiazepine binding site of GABAA receptors. *Neuroimage* **232**, (2021).

72. Breton-Provencher, V. & Sur, M. Active control of arousal by a locus coeruleus GABAergic circuit. *Nat. Neurosci.* **22**, 218–228 (2019).

73. Beliveau, V. *et al.* A High-Resolution In Vivo Atlas of the Human Brain's Serotonin System. *J. Neurosci.* **37**, 120 (2017).

74. Sasaki, T. *et al.* Quantification of dopamine transporter in human brain using PET with 18F-FE-PE2I. *J. Nucl. Med.* **53**, 1065–1073 (2012).

75. Dahl, M. J. *et al.* Locus coeruleus integrity is related to tau burden and memory loss in autosomal-dominant Alzheimer's disease. *Neurobiol. Aging* **112**, 39–54 (2022).

76. Mumford, J. A., Turner, B. O., Ashby, F. G. & Poldrack, R. A. Deconvolving BOLD activation in event-related designs for multivoxel pattern classification analyses. *Neuroimage* **59**, 2636–2643 (2012).

77. Rissman, J., Gazzaley, A. & D'Esposito, M. Measuring functional connectivity during distinct stages of a cognitive task. *Neuroimage* **23**, 752–763 (2004).

78. Box, G. E. P., Jenkins, G. M. & Reinsel, G. C. *Time series analysis: forecasting and control* . (Prentice Hall, 1994).

79. Uematsu, A. *et al.* Modular organization of the brainstem noradrenaline system coordinates opposing learning states. *Nat. Neurosci.* **20**, 1602–1611 (2017).

80. Salvi, V. *et al.* Cingulate cortex stimulation drives distinct pupillary responses in rat via recruitment of noradrenergic neurons in the locus coeruleus. *Cereb. Cortex* **35**, (2025).

81. Prokopiou, P. C. *et al.* Lower novelty-related locus coeruleus function is associated with Aβ-related cognitive decline in clinically healthy individuals. *Nat. Commun. 2022 131* **13**, 1–14 (2022).

82. Seeley, W. W. The Salience Network: A Neural System for Perceiving and Responding to Homeostatic Demands. *J. Neurosci.* **39**, 9878–9882 (2019).

83. Hermans, E. J. *et al.* Stress-related noradrenergic activity prompty large-scale neural network reconfiguration. *Science (80-. ).* **334**, 1151–1153 (2011).

84. Breton-Provencher, V., Drummond, G. T., Feng, J., Li, Y. & Sur, M. Spatiotemporal dynamics of noradrenaline during learned behaviour. *Nat. 2022* 1–7 (2022) doi:10.1038/s41586-022-04782-2.

85. Smegal, L. F. *et al.* Lower locus coeruleus integrity is associated with diminished practice effects in clinically unimpaired older individuals. *Neurobiol. Aging* **152**, 13–24 (2025).

86. Lindenberger, U. & Mayr, U. Cognitive aging: Is there a dark side to environmental support? *Trends Cogn. Sci.* **18**, 7–15 (2014).

87. Duszkiewicz, A. J., McNamara, C. G., Takeuchi, T. & Genzel, L. Novelty and dopaminergic modulation of memory persistence: A tale of two systems. *Trends Neurosci.* **42**, 102–114 (2019).

88. Kret, M. E. & Sjak-Shie, E. E. Preprocessing pupil size data: Guidelines and code. *Behav. Res. Methods* **51**, 1336–1342 (2019).

89. Dahl, M. J. *et al.* Rostral locus coeruleus integrity is associated with better memory performance in older adults. *Nat. Hum. Behav.* **3**, 1203–1214 (2019).

90. Halchenko, Y. O. *et al.* HeuDiConv — flexible DICOM conversion into structured directory layouts. *J. Open Source Softw.* **9**, 5839 (2024).

91. Esteban, O. *et al.* Crowdsourced MRI quality metrics and expert quality annotations for training of humans and machines. *Sci. data* **6**, 30 (2019).

92. Esteban, O. *et al.* MRIQC: Advancing the automatic prediction of image quality in MRI from unseen sites. *PLoS One* **12**, e0184661 (2017).

93. Esteban, O. *et al.* fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat. Methods* **16**, 111–116 (2019).

94. Worsley, K. J. & Friston, K. J. Analysis of fMRI Time-Series Revisited—Again. *Neuroimage* **2**, 173–181 (1995).

95. Satterthwaite, T. D. *et al.* An improved framework for confound regression and filtering for control of motion artifact in the preprocessing of resting-state functional connectivity data. *Neuroimage* **64**, 240–256 (2013).

96. Friston, K. J., Williams, S., Howard, R., Frackowiak, R. S. J. & Turner, R. Movement-Related
effects in fMRI time-series. *Magn. Reson. Med.* **35**, 346–355 (1996).
97. Delorme, A. & Makeig, S. EEGLAB: An open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. *J. Neurosci. Methods* **134**, 9–21 (2004).
98. Dimigen, O., Sommer, W., Hohlfeld, A., Jacobs, A. M. & Kliegl, R. Coregistration of eye
movements and EEG in natural reading: Analyses and review. *J. Exp. Psychol. Gen.* **140**, 552–
572 (2011).
99. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J. M. FieldTrip: Open source software for
advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput. Intell.*
*Neurosci.* **2011**, 1–9 (2011).
100. Maris, E. & Oostenveld, R. Nonparametric statistical testing of EEG- and MEG-data. *J.*
*Neurosci. Methods* **164**, 177–190 (2007).
101. Friston, K. J., Ashburner, J., Kiebel, S., Nichols, T. & Penny, W. *Statistical Parametric*
*Mapping: The Analysis of Functional Brain Images. Statistical Parametric Mapping: The*
*Analysis of Functional Brain Images* (Academic Press, 2007). doi:10.1016/B978-0-12-372560-
8.X5000-1.
102. Friston, K. J. *et al.* Event-Related fMRI: Characterizing Differential Responses. *Neuroimage* **7**,
30–40 (1998).
103. Avants, B. B., Tustison, N. & Song, G. Advanced Normalization Tools: V1.0. *Insight J.* **2**,
(2009).
104. Tustison, N. J. *et al.* The ANTsX ecosystem for quantitative biological and medical imaging.
*Sci. Reports 2021 111* **11**, 1–13 (2021).
105. Murphy, P. R., O'Connell, R. G., O'Sullivan, M., Robertson, I. H. & Balsters, J. H. Pupil
diameter covaries with BOLD activity in human locus coeruleus. *Hum. Brain Mapp.* **35**, 4140–
4154 (2014).
106. Theiler, J., Eubank, S., Longtin, A., Galdrikian, B. & Doyne Farmer, J. Testing for nonlinearity
in time series: the method of surrogate data. *Phys. D Nonlinear Phenom.* **58**, 77–94 (1992).
107. Krishnan, A., Williams, L. J., McIntosh, A. R. & Abdi, H. Partial Least Squares (PLS) methods
for neuroimaging: A tutorial and review. *Neuroimage* **56**, 455–475 (2011).
108. McIntosh, A. R. & Lobaugh, N. J. Partial least squares analysis of neuroimaging data:
Applications and advances. in *NeuroImage* vol. 23 (Neuroimage, 2004).
109. Brett, M., Anton, J.-L., Valabregue, R. & Poline, J.-B. Region of interest analysis using an
SPM toolbox. in *Human Brain Mapping conference* (2002).
1104