Learned statistical regularity drives anticipatory micro-saccades

toward suppressed distractor locations

Sirui Chen[1,2], Xin Zhang[1,2,3], Xinyu Li[3], Ole Jensen[4], Jan Theeuwes[5], and Benchi Wang[1,2*]

[1]Key Laboratory of Mental Health of the Ministry of Education, Guangdong-Hong Kong-Macao Greater Bay Area

Center for Brain Science and Brain-Inspired Intelligence, Guangdong-Hong Kong Joint Laboratory for Psychiatric

Disorders, Guangdong Province Key Laboratory of Psychiatric Disorders, Guangdong Basic Research Center of

Excellence for Integrated Traditional and Western Medicine for Qingzhi Diseases, Department of Neurobiology,

School of Basic Medical Sciences, Southern Medical University, Guangzhou, China

[2]Institute for Brain Research and Rehabilitation, South China Normal University, Guangzhou, China

[3]Department of Psychology, Zhejiang Normal University, Jinhua, China

[4]Department of Psychiatry and Experimental Psychology, University of Oxford, Oxford, United Kingdom

[5]Department of Experimental and Applied Psychology, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

Author notes

BW designed the experiment, XZ and XL collected the data, SC and BW analyzed the data. SC, OJ, JT and BW wrote the paper. All authors approved the final version of the manuscript for submission. All authors declared no competing interests. This research was supported by the Brain Science and Brain-like Intelligence Technology−National Science and Technology Major Project (2022ZD0204800) and the Natural Science Foundation of Guangdong grant (2023A1515012789) to BW; the Wellcome Trust Discovery Award (227420) and the NIHR Oxford Health Biomedical Research Centre (203316) to OJ. Correspondence should be addressed to Benchi Wang, wangbenchi.swift@gmail.com. The code and the data used for reproducing the results shown in the figures can be accessed through https://doi.org/10.17605/OSF.IO/DFRPQ.

Abstract

Statistical learning enables individuals to suppress locations associated with salient distractors, yet the mechanisms underlying this suppression remain unclear. Proactive accounts propose that suppression operates without prior attentional allocation to distractor locations, whereas reactive accounts suggest that suppression requires covert attention to the distractor location before it can be engaged. To address this ongoing debate, the current study recorded micro-saccades—an index of covert attention—during the pre-stimulus interval, alongside electroencephalogram (EEG) data collected during a visual search task. Participants were instructed to ignore a salient distractor that occurred more frequently at a specific location. Statistical learning reduced attentional capture for distractors presented at this high-probability location, and this effect was accompanied by oculomotor markers: micro-saccade rates decreased before stimulus onset relative to a control condition. Strikingly, these anticipatory micro-saccades were more often directed toward high-probability distractor locations than away from them, consistent with a reactive suppression mechanism. In parallel, alpha-band activity (8–14 Hz) carried decodable representations of high-probability distractor locations, indicating preparatory neural tuning. Overall, these findings provide evidence that the oculomotor system is closely involved in encoding and responding to learned spatial regularities.

Keywords: Statistical learning, distractor suppression, micro-saccades, alpha suppression

Introduction

The ability to extract statistical regularities (i.e., statistical learning) from the environment is fundamental to human survival. Recent studies have highlighted the role of statistical learning on attentional selection[1–3], as well as the suppression of distracting salient stimuli[4,5]. For instance, distractors appearing frequently at specific locations cause less interference than those appearing at other locations, indicating that implicitly learned regularities facilitate distractor suppression[6–11]. This learned suppression likely enhances perceptual efficiency and supports action planning when navigating in complex environments.

Despite its importance, the neural mechanism underlying learned suppression remains poorly understood, with ongoing debate regarding the nature of attentional suppression[12]. The suppression can, in principle, be implemented in two ways: proactively (before the display appears) or reactively (following attentional engagement). Studies on statistical learning propose that the spatial priority map is proactively altered, reducing competition for attention at distractor locations[1,13–18]. This proactive adjustment diminishes the distractor saliency and minimizes their attentional capture over sustained periods. Conversely, other studies provide evidence for reactive distractor suppression, where suppression occurs only after attention has been engaged[19–26], resembling the classic phenomenon of inhibition of return[27,28].

Notably, the human oculomotor system might offer a compelling avenue for investigating this issue, given its close link to covert spatial attention, as evidenced by micro-saccades. It is well-established that micro-saccades tend to occur in the direction of covert attention[29–32]. Furthermore, micro-saccades occur not only when directing attention to external locations[29,33], but also when directing attention to internal memory representations by retro-cues[31,34]. This highlights the role of micro-saccades in reflecting the allocation of covert attention to specific locations or memory-based representations, even in the absence of visual stimuli. These findings provide a unique method, by measuring micro-saccades, for examining whether

covert attention precedes attentional suppression (supporting reactive mechanism) or

suppression can occur independently of prior attention allocation (supporting

proactive mechanism).

Thus, in the current study, we measured micro-saccades during the pre-stimulus

interval, assessing the involvement of covert attention prior to display onset, and

examined its relationship with neural activity. We simultaneously recorded

electroencephalogram (EEG) and eye movements while participants performed the

additional singleton task[9,35]. In this task, participants were instructed to search for a

unique shape (the target) while ignoring a salient distractor colored differently (see

Fig. 1a). The salient distractor was presented more frequently at one specific location

(high-probability location) compared to others (low-probability location; see Fig. 1b

left panel and Methods for details), introducing a statistical learning component,

resulting in learned suppression of the salient distractor. Notably, we also ran a control

experiment where the locations of the distractors were not predictable. In this setup,

the distractor appeared equally often at all possible locations, so participants couldn't

learn or anticipate where it would be (see Fig. 1b right panel).

Figure 1. Task, behavior, and micro-saccade dynamics. a) Search task: participants reported line
orientation inside the unique shape (e.g., circle) while ignoring a salient distractor (e.g., red diamond).
Distractor-present and -absent trials were intermixed. b) Distractor location probabilities. In the main
experiment, one side (left/right, counterbalanced) had 65% probability; each of five other locations had
7% probability. In the control experiment, each location had 16.67% probability. c) Main experiment:
Left panel shows response times (RTs) for different distractor conditions, comparing high- and
low-probability distractor locations and no-distractor trials. Right panel shows RTs for distractor-absent
trials, comparing target appearances at high- versus low-probability distractor locations. d) Mean RTs
in the control experiment, comparing distractor present and absent trials. Individual RTs are represented
by scatters. e) Micro-saccade rates in the main and control experiments, with data variability
represented by ±1 standard error of the mean (SEM). Time window, showing that micro-saccade rates
were significantly different between the control (blue) and main (red) experiments, is highlighted by
the solid gray line above (cluster-based permutation test, $p < .05$). f) The left panel shows
polar-histograms of anticipatory micro-saccades directions during the pre-stimulus period (-2500 ms to
0 ms), the middle panel shows ratios of micro-saccades directed toward or away from high-probability
distractor locations, and the right panel shows time-binned ratios, with significant bins marked (*$p <$

0.05, FDR corrected). Ratios represent the proportion of micro-saccades directed toward or away from

the learned distractor location (count in each direction divided by total micro-saccade count), which is

equivalent to directional micro-saccade rate because the total micro-saccade count does not vary

between conditions. g) Pre-stimulus micro-saccades in control experiment showing no directional bias.

h) Scatter plots illustrating correlations between micro-saccade onset times and RTs for micro-saccades

directed towards or away from high-probability locations (linear fits in red/pink). *$p < .05$, n. s. denotes

non-significant. i) Dynamic time warping (DTW) between learned suppression and micro-saccade bias

(see Methods for details). Left panel shows DTW path, illustrating co-variation between learned

suppression and micro-saccade bias. Right panel shows permutation results of the DTW distance, with

red line indicating the true value and the gray histogram showing the permutation distribution.

Results

Mean response times (RTs) were slower when distractors were present compared

to absent (757 ms), regardless of whether they appeared at high-probability locations

(827 ms), $t(27) = 8.59$, $p < .001$, or low-probability locations (872 ms), $t(27) = 9.55$, p

$< .001$ (Fig. 1c left panel), reflecting capture by salient distractors[35,36]. Notably, faster

responses were observed when the salient distractor was presented at the high-

compared to low-probability locations, $t(27) = 6.64$, $p < .001$ (Fig. 1c), suggesting

suppression of the high-probability distractor location[9]. In addition, we investigated

the learned suppression effect over time by performing a 2 (condition: high- vs.

low-probability locations) $\times$ 10 (blocks) repeated-measures ANOVA. This revealed

significant main effects of condition, $F(1,27) = 40.38$, $p < .001$, and block, $F(9,243) =$

5.21, $p < .001$, but no significant interaction, $F(9,243) = 0.93$, $p = .495$ (see Fig. S1 in

Supplementary Information). These results indicate that although overall performance

systematically changed across blocks, learned suppression consistently persisted over

time, consistent with previous studies (e.g., refs[37]).

In the distractor absent trials, responses were slower when the target appeared at

high-probability distractor locations (792 ms) compared to low-probability locations

(751 ms), $t(27) = 4.2$, $p < .001$ (Fig. 1c right panel). This suggests that suppression is

spatially based, rather than driven by the suppression of the salient feature itself.

Moreover, the behavioral results in the control experiment showed that participants

responded more slowly when a distractor was present (985 ms) compared to when it

was absent (862 ms), $t(19) = 8.55$, $p < .001$, purely reflecting attentional capture by

salient distractors.

Because participants were not explicitly asked whether they noticed the spatial

regularity, we cannot rule out that some participants acquired explicit knowledge of

the high-probability distractor locations. However, previous work suggests that

explicit knowledge alone is insufficient to produce the characteristic distractor

suppression observed in such learning paradigms[10].

Anticipatory micro-saccades during pre-stimulus gazes

As hypothesized, learned suppression may engage the oculomotor system before

stimulus onset. To test this, we examined the micro-saccade rate and direction during

the fixation period preceding the search array onset. As shown in Fig. 1e, overall

micro-saccade rates exhibited a significant decrease after fixation presentation in the

main experiment compared to the control condition, during the interval from -2500 to

113 ms relative to stimulus onset (cluster-based permutation test based on two-sample

t-tests, $p < .05$). This finding is consistent with previous studies[29,38,39], which

demonstrated that covert attention can suppress micro-saccade rate. Such suppression

is generally interpreted as an indicator of covert attentional engagement in preparation

for the coming task.

Critically, a key finding was that a greater proportion of pre-stimulus

micro-saccades during the pre-stimulus period were directed toward the

high-probability distractor location (37.2%) compared to those directed away (29.5%),

$t(27) = 3.05$, $p = .005$ (see left and middle panels in Fig. 1f). This indicates that covert

attention was proactively biased toward high-probability locations[29,40], even though

the behavioral data showed that this location was subsequently suppressed. To further

examine the temporal evolution of micro-saccade dynamics, we performed a

sliding-window analysis using a 500 ms window with a 250 ms overlap across the

entire trial. Within each time-bin, paired t-tests compared the proportion of

micro-saccades directed toward versus away from the high-probability location. A

significant bias toward the high-probability location emerged from -2000 ms

(time-bin 2) to -250 ms (time-bin 8) before stimulus onset (all $ps < .05$; Fig. 1f, right

panel), indicating sustained proactive attentional deployment. Notably, in the control

experiment, we found no pre-stimulus micro-saccade bias. Specifically, there was no

difference in micro-saccade direction toward the distractor side (contralateral) versus

the opposite side (ipsilateral), for the leftmost distractor (0.235 vs. 0.241 respectively),

$t(19) = 0.57$, $p = .58$, $BF_{10} = 0.74$, the rightmost distractor (0.235 vs. 0.229

respectively), $t(19) = 0.76$, $p = .55$, $BF_{10} = 0.84$ (see Fig. 1g), and the other four

locations, all $ps > .54$. Notably, the micro-saccade directional bias in the main

experiment ($ratio_{toward} - ratio_{away}$) was significantly stronger than the corresponding

directional bias in the control experiment ($ratio_{contralateral} - ratio_{ipsilateral}$) for both the

leftmost distractor, $t(46) = 2.31$, $p = .026$, and the rightmost distractor, $t(46) = 2.69$, p

$= .02$.

To directly link behavior with micro-saccades, we tested whether the timing of

micro-saccades toward the high-probability distractor location was related to

behavioral RTs. We predicted that earlier micro-saccades—indicating earlier

attentional orienting—would lead to faster responses. As expected, a significant

positive correlation was found when micro-saccades were directed toward the

high-probability location, $r = 0.41$, $p = .03$, but not when they were directed away, $r =$

0.10, $p = .61$ (see Fig. 1h). Because these micro-saccades occurred well before

stimulus onset, this relationship likely reflects variability in preparatory attentional

states rather than a direct influence of individual eye movements on behavior. In

contrast, the control experiment showed no significant correlation between

micro-saccade direction and performance (toward left: $r = 0.04$, $p = .85$; toward right:

$r = -0.01$, $p = .97$; see Fig. S2), supporting the idea that this effect reflects learned

spatial suppression, not general oculomotor dynamics toward distractor salience.

To explore how micro-saccade bias and behavioral suppression developed

together over time, we employed a Dynamic Time Warping (DTW) analysis to

compare their learning-related temporal trajectories (see Methods for details). We

divided the main experiment into 20 time-bins and created two normalized time series:

one tracking micro-saccade bias (the ratio of micro-saccades toward vs. away from

the high-probability location), and the other tracking suppression (the RT difference

between distractors at the high- and low-probability locations). The DTW analysis

showed strong temporal alignment between the two signals (DTW distance = 0.31;

Fig. 1i), confirmed by a nonparametric permutation test (p = .01). These findings

reveal that micro-saccade bias and behavioral suppression develop in sync over time,

rather than emerging independently. This provides evidence that predictive

micro-saccades both reflect and potentially contribute to the gradual development of

learned attentional suppression.

Tracking learned high-probability locations from pre-stimulus alpha oscillation

Previous studies have identified contralateral pre-stimulus alpha oscillations over

the parietal and occipital cortex associated with statistically learned suppression at

high-probability locations[18]. Yet, in the present study, we did not observe such

lateralized modulation (see Fig. 2a). Instead, pre-stimulus alpha power showed a

sustained increase over both contralateral and ipsilateral electrodes relative to the

high-probability location. This increase spanned from –2324 ms to –514 ms over

contralateral electrodes and from –2328 ms to –532 ms over ipsilateral electrodes

before stimulus onset. This was followed by a reduction in alpha power from 252 ms

to 1000 ms and from 260 ms to 1000 ms after the onset of the search, respectively

(cluster-based permutation test, p < .05; Fig. 2a). To determine whether the observed

alpha oscillation reflected attention being directed toward or away from the

high-probability location, we employed a multi-variate inverted encoding model[41],

which is capable of decoding neuronal representations associated with direction of

spatial attention from the brain activity distributed across parietal and occipital

electrodes. We successfully tracked spatial representation of the high-probability

location from alpha power during the -400 to -20 ms interval relative to stimulus onset

(cluster-based permutation test, p < .05; Fig. 2b), while no such representation was

observed for low-probability locations. Moreover, a significant difference between

high- and low-probability location emerged during the -480 to 0 ms interval relative

to the stimulus onset (cluster-based permutation test, $p < .05$; Fig. 2b). These findings

suggest that the spatial distribution of pre-stimulus alpha oscillation may reflect

enhanced feedforward control mechanisms over high-probability locations,

establishing a preparatory state that anticipates and encodes the spatial representation

of the high-probability location.

Notably, the decodable representation of the high-probability location emerged

only after the peaks of both alpha power and micro-saccade activity, suggesting a

temporally ordered cascade in which early anticipatory micro-saccades prime

subsequent alpha-based spatial tuning, thereby facilitating later suppression processes.

Both alpha-band desynchronization and micro-saccade suppression have been linked

to heightened attentional engagement and cortical excitability[29,42], consistent with the

idea that robust spatial representations arise when attention is stably anchored to the

expected distractor location—explaining peak decoding despite reduced alpha power

and micro-saccade rate. Supporting this view, the ratio of early micro-saccades

directed toward the high-probability location significantly predicted later alpha-based

decoding strength, $r = 0.50$, $p = .006$, whereas no such relationship was found for

micro-saccades directed away from it, $r = 0.14$, $p = .48$ (see Fig. 2c).

Figure 2. a) Pre-stimulus alpha oscillation over the contralateral and ipsilateral electrodes relative to the high-probability distractor location (cluster-based permutation test, $p < .05$). b) Spatial representations of high- and low-probability distractor locations tracked within alpha oscillations (8-14 Hz) using an inverted encoding model (see Methods for details). Significant CTF slopes for high-probability locations are marked in red horizontal line (cluster-based permutation test, $p < .05$). The difference between different types of locations are marked in black horizontal line (cluster-based permutation test, $p < .05$). c) Scatter plots illustrate the correlation between the alpha CTF slopes and the ratio of micro-saccades directed towards (left panel) and away from (right panel) high-probability locations (linear fits in red/pink). $**p < .01$, n. s. denotes non-significant. d) Saccade-related potential (SRP) waveform from posterior regions. e) SRP topography averaged from -500 to 1000 ms relative to micro-saccade onset. Red dots present 17 posterior electrodes used in all analyses. f) Z-scored SRP power spectrum from -500 to 1000 ms. Asterisk indicates significant activations in frequencies > 4 Hz ($p < .05$). Shaded areas denote ±1 SEM in sub-figures a, b, d, and f. g) Left: time–frequency power spectrum of micro-saccade-locked activity. The red dashed box highlights the 8–14 Hz band. Right: classification accuracy for anticipatory micro-saccades directed toward vs. away from the high-probability location based on alpha activity.

Micro-saccade-locked activity

To further validate the role of micro-saccades in driving neural activity associated with learned suppression, we investigated whether micro-saccade-locked brain

activity across multiple frequency bands encoded information about learned statistical

regularities. Previous studies have demonstrated that micro-saccades elicit posterior

EEG activity[43–45], closely linking them to visual processing. This raises the possibility

that micro-saccade-locked brain activity is also sensitive to learned suppression, as

suppression acts by modulating visual processing through statistical learning. To test

this hypothesis, we analyzed the power of micro-saccade-locked neural oscillations

over occipital-parietal cortices across multiple frequency bands (3-40 Hz; see

Methods for details).

We computed saccade-related potential (SRP) time-locked to micro-saccade

onsets and observed a prominent transient spike at 0 ms (Fig. 2d). This short-lived

broadband response peaked precisely at micro-saccade onset and corresponds to the

well-documented saccadic spike potential[46]. Fig. 2e presents the SRP topography

averaged from -500 ms to 1000 ms relative to micro-saccade onset. Beyond this

transient effect, we also observed oscillatory activity surrounding the micro-saccade

onset, reflecting corresponding rhythmic fluctuations in specific frequency bands. A

spectral analysis of the -500 to 1000 ms window around micro-saccade onset revealed

a clear alpha-band peak against zero (10-11 Hz, $p < .05$) superimposed on the typical

1/f distribution (Fig. 2f), indicating alpha as the dominant oscillatory component.

Importantly, no sustained higher-frequency activity was present, as only oscillatory

rhythms such as alpha remain after accounting for the 1/f background. Figure 2g (left

panel) presents a temporal increase in power across a wide range of frequencies (3-40

Hz) following micro-saccade onset.

When isolating alpha activity (8-14 Hz) from raw EEG signals, we found that

micro-saccade-locked alpha activity encoded learned statistical regularities, tracking

the direction of anticipatory micro-saccades toward high-probability locations.

Specifically, using a multivariate decoding approach (see Methods for details), we

successfully discriminated micro-saccades directed toward and away from the

high-probability location based on micro-saccade-locked alpha activity during interval

of 40 - 440 ms after the anticipatory micro-saccade onset (cluster-based permutation

test, $p < .05$; Fig. 2g, right panel). Notably, micro-saccades toward the

high-probability location were evenly split between left and right directions, and the

same was true for micro-saccades directed away. This balanced distribution confirms

that the decoding results reflect information about the expected location, rather than

simply the movement direction of the micro-saccades.

No significant alpha activity was observed in the micro-saccade-locked signals

recorded from electro-oculogram (EOG) electrodes, ruling out the possibility that the

observed findings were driven by artifacts associated with micro-saccades[43]. These

results suggest that, although micro-saccades may not be obligatory for task

completion, they indeed play a role in triggering neural activity that encodes and

processes learned information.

Discussion

The current study shows that learned suppression proactively recruits the

oculomotor system. Consistent with previous studies, we observed the classic

attentional capture effect[35,36] and reduced capture for distractors presented at the

high-probability location, indicating learned suppression[1,5,7,9–11]. A novel finding of

our study is the decreased micro-saccade rate, compared to the control condition,

observed before the search display onset (Fig. 1e). This indicates the involvement of

covert attention in anticipating the arrival of the search display. Critically, and perhaps

unexpectedly, these anticipatory micro-saccades were more frequently directed

toward high-probability distractor locations than away from them (Fig. 1f). Consistent

with this finding, we successfully tracked high-probability locations in pre-stimulus

alpha power (8-14 Hz). Further analysis revealed that micro-saccades could trigger

neural activity sensitive to learned regularities, with micro-saccade-locked alpha

activity encoding and tracking the direction of anticipatory micro-saccades (Fig. 2g).

Collectively, these findings suggest that learned suppression engages the oculomotor

system, potentially functioning as a monitoring mechanism for locations likely to

contain distractors.

Micro-saccades are deeply integrated into various cognitive processes, including

attentional control[29,33], temporal expectation[47–49], working memory[31,34], and

perceptual learning[50]. Our study extends these findings by showing that anticipatory

micro-saccades are actively involved in learned suppression associated with statistical

regularities (i.e., high-probability distractor locations). While previous studies have

claimed that learned suppression can occur proactively without covert attention first

being directed toward the suppressed location[1,13,15–18,51,52], the current findings suggest

otherwise. We speculate that attending to the distractor location proactively might

provide an advantage, as it allows the observer to eliminate it as a potential target

location. This process triggers the fast disengagement of covert attention from the

distractor location when the search array is presented, ultimately resulting in learned

suppression. Our findings, indicating that anticipatory micro-saccades actively

contribute to learned suppression, align with the "search and destroy" hypothesis[24],

which claims that suppression is only possible after first attending to the location that

needs to be ignored.

Furthermore, from a theoretical perspective, the involvement of the oculomotor

system in learned suppression could be explained by its role in processing spatial

information. As the brain learns to identify and track locations likely to contain

distractors, it may rely on the oculomotor system's precise spatial representation

capabilities. This idea aligns with previous research showing that remembering spatial

locations activates the oculomotor system, leading to eye movements that curve away

from locations marked for later recall[53,54]. Notably, prior research has shown that

suppression effects of the kind observed in the present study do not arise when

participants are explicitly instructed to ignore a salient distractor[10]. Instead, a growing

body of work supports a clear dissociation between suppression driven by implicit

statistical learning and that arising from explicit, strategic control[4,55]. Furthermore,

recent findings indicate that micro-saccade biases are not reliably elicited by

probabilistic cues alone[56], underscoring the importance of distinguishing effects due

to implicit learning mechanisms from those attributable to explicit cueing.

Previous studies have identified pre-stimulus alpha oscillations over the parietal

and occipital cortices associated with statistically learned suppression[18]. Specifically,

increased parieto-occipital alpha power contralateral to high-probability distractor

locations has been observed up to one second before stimulus onset, consistent with

inhibitory processes linked to distractor suppression[42]. However, in the present study,

we did not observe such lateralized modulation. This discrepancy raises an open

question for future research on the specific role of pre-stimulus (lateralized) alpha in

learned suppression.

Notably, we were able to decode the spatial representation of the high-probability

location from pre-stimulus alpha oscillations. Furthermore, we successfully

differentiated micro-saccades directed toward and away from the high-probability

location based on micro-saccade-locked alpha activity. These findings highlight a

potential connection between the human oculomotor system and neural oscillations in

the attention network[57]. Indeed, our results reveal a temporal cascade in which early

micro-saccades bias the oculomotor system and subsequently prime alpha-based

spatial tuning: decodable representations of the high-probability location emerged

only after the peak of micro-saccade activity, and the ratio of early micro-saccades

directed toward this location significantly predicted later decoding strength, whereas

no such relationship was observed for away-directed micro-saccades (Fig. 2c). One

possibility is that micro-saccades toward high-probability locations actively reinforce

neural spatial representations by transiently enhancing posterior cortical activity.

Mechanistically, this may occur through periodic "refreshing" of neural activity via

phase resetting, which sharpens alpha spatial specificity and stabilizes attentional

representations[58]. Thus, micro-saccades are not passive by-products of attention but

active contributors that maintain and refine spatial representations. Future work

should investigate how oculomotor and neural dynamics are jointly orchestrated to

optimize attentional control in complex environments.

Another open question is whether the anticipatory micro-saccade bias observed

here is specific to distractor suppression or reflects a more general expectation of

salient events. In the present study, spatial probabilities were manipulated only for

distractors, allowing us to examine how learning about distractor locations shapes

anticipatory oculomotor behavior. Within this framework, the observed micro-saccade

bias toward high-probability distractor locations is consistent with the idea that covert

attention is proactively allocated to locations that are expected to contain distractors,

thereby facilitating their suppression once the display appears. Future studies that

independently manipulate both target and distractor spatial probabilities could further

clarify whether anticipatory micro-saccades reflect suppression-specific mechanisms

or a broader predictive orienting toward expected salient stimuli.

Overall, our findings expand the traditional understanding of micro-saccades by

demonstrating their involvement in high-level cognitive processes, particularly their

role in statistically learned suppression. Within the context of attentional selection, we

uncovered potential neural mechanisms reflecting learned suppression, contributing to

the ongoing debate on whether attentional processing precedes suppression [12]. Our

results suggest that learned suppression is reactive and only occurs after attention is

directed to the location that needs to be suppressed.

Methods

Participants

Twenty-eight volunteers (13 males, mean age = 20.1 years old) participated in the

experiment for monetary compensation (¥80 per hour). The sample size was

predetermined based on previous studies[9,18]. All participants had normal color vision

and normal or corrected-to-normal visual acuity, and provided written informed

consent before testing. The study was approved by the ethics review committee of

South China Normal University (2020-3-013).

Apparatus and stimuli

414   Each stimulus contained a vertical or horizontal white line ($0.2° \times 1.4°$,

415 Red-green-blue [RGB]: 255,255,255) inside, presented against a black background

416 (RGB: 0, 0, 0). The enclosing shapes included a circle with a radius of $1°$ or a

417 diamond subtending $2° \times 2°$, outlined in either red (RGB: 255, 0, 0) or green (RGB:0,

418 255, 0) outline. These stimuli were presented on an imaginary circle with a radius of

419 $4°$, centered around a white fixation cross (RGB: 255, 255, 255) measuring $1° \times 1°$.

420 Stimulus presentation and response registration were controlled by custom Python

421 scripts. Participants were seated in a dimly lit laboratory, positioned in front of a 27-in.

422 LCD monitor at a viewing distance of 65 cm.

Procedure and design

424   We employed the classic additional singleton task[9,35], which required participants

425 to search for a shape singleton while ignoring a color distractor singleton. At the

426 beginning of each trial, a self-paced drift check was performed, followed by a central

427 fixation cross remaining visible throughout a trial. During the drift check, participants

428 fixated on a central cross to initiate the trial, while the eye-tracking system ensured

429 that their gaze remained within $3°$ of visual angle from the center. After 2500 ms, the

430 search array was presented for 3000 ms or until a response was registered.

431 Participants were instructed to identify the unique shape (target) in the search array,

432 ignoring the salient color singleton (salient distractor), and indicate the orientation

433 (horizontal or vertical) of the line inside the target by pressing the "left" or "up" key,

434 respectively. The experiment contained 10 blocks, each comprising 120 trials,

435 consisting of 40 distractor-absent trials and 80 distractor-present trials. This design

436 provided enough trials to analyze the distractor-present condition. While it may have

437 introduced some potential anticipation of distractor presence, our main comparison

438 was within the distractor-present condition itself—that is, between different types of

439 distractor locations (high- vs. low-probability distractor locations) within distractor

440 present trials. Participants were required to maintain fixation at the display center

441 throughout each trial. Feedback messages were presented for trials where participants

442 blinked, failed to respond, pressed a wrong key, or deviated their gaze more than $3°$

from the fixation cross. Such trials were later retested in a random order after

completing the remaining trials.

Due to the predominance of horizontal micro-saccades[29], the salient distractor

appeared on either the left (180°) or right (0°) side (counterbalanced across

participants) with a probability of 65%, while it appeared at other five locations with

an even probability of 7% (see Fig. 1b). The target was presented at each location

with equal probability. A brief 10-trial practice session was conducted before the main

experiment. Trials on which the response times (RTs) were faster than 200 ms and

slower than 2000 ms were excluded from analyses (1.31% of trials).

Notably, some participants from the same group were successfully invited back

approximately one year later to perform a control experiment, in which no distractor

location probability was manipulated while all other parameters remained identical to

those of the main experiment. That is, the salient distractor appeared equally often at

all possible locations. In total, twenty participants (9 males, mean age = 21.3 years)

took part in the control experiment. It comprised 5 blocks of 120 trials each, with 40

distractor-absent trials and 80 distractor-present trials per block. Trials with response

times (RTs) faster than 200 ms or slower than 2000 ms were excluded from analysis

(1.6% of total trials).

Eye-tracking acquisition

The eye tracker (EyeLink Portable Duo, SR Research) was positioned

approximately 10 cm in front of the monitor, with participants seated ~55 cm away

from the screen. Gaze positions (horizontal and vertical) were continuously recorded

at a sampling rate of 1000 Hz. Before the beginning of each block, the eye-tracking

system was calibrated and validated according to standard 9-point protocols provided

by the EyeLink software, with an acceptance threshold of $\leq 1°$ of visual angle.

Micro-saccade detection

Micro-saccades were detected using an improved version[59] of the algorithm

initially proposed by Engbert and Kliegl (2003). Horizontal and vertical eye positions

were mapped onto a velocity space, with a relatively low velocity threshold of 3

standard deviations used to detect micro-saccades[59]. For a detected saccade to be

considered as a micro-saccade, it needed to exhibit temporal overlap between both

eyes, a minimum duration of 12 ms, and an amplitude below 1°. This gave average

micro-saccade rates of 0.324 Hz in the main experiment and 0.598 Hz in the control

experiment. The reliability of this method was confirmed by two checks: (1) a strong

positive correlation between log-transformed peak velocity and amplitude (main

experiment: $r = 0.91$; control experiment: $r = 0.89$; both ps $< .001$; Fig. S3a), and (2)

the fact that most saccades were smaller than 1° of visual angle (Fig. S3b), consistent

with known micro-saccade dynamics[29]. A 100 ms sliding window (stepped in 1-ms

intervals) was employed to assess the micro-saccade rate following the fixation

onset[29,59,60].

We categorized the micro-saccades based on their directional vectors into 12

distinct bins, with each bin defined by angular ranges of 30° (specifically: 345°-15°,

15°-45°, 45°-75°, 75°-105°, 105°-135°, 135°-165°, 165°-195°, 195°-225°, 225°-255°,

255°-285°, 285°-315°, and 315°-345°). This binning approach allowed for a detailed

analysis of the directional distribution of micro-saccades, facilitating the assessment

of their role in learned suppression.

Since half participants in the learning condition had high-probability locations on

the left side of the visual field while others had them on the right side, we aligned the

data by mirroring the directional bins for those participants with high-probability

locations on the right side. This mirroring ensured that horizontal micro-saccades

within the 345°-15° bin were treated as micro-saccades directed away from the

high-probability location, while those within the 165°-195° bin were treated as

directed toward the high-probability location. We quantified micro-saccades across

the entire 2500 ms pre-stimulus window to examine proactive attention.

Dynamic time warping

To quantify how closely micro-saccade bias and behavioral learned suppression

aligned over time, we used Dynamic Time Warping (DTW)—an algorithm designed

to measure the similarity between two time series that may differ in timing or speed.

DTW identified the optimal alignment between sequences by non-linearly stretching

or compressing the time axis, minimizing the cumulative distance between

corresponding points.

Let $X = \{x_1, x_2, \ldots, x_n\}$ and $Y = \{y_1, y_2, \ldots, y_m\}$ represent the two time

series being compared, where x_i and y_i are the values at time points i and j,

respectively. The DTW distance $D(i,j)$ is computed recursively using the following

equation:

$$D(i,j) = \text{dist}(x_i, y_i) + \min \begin{cases} D(i-1, j) \\ D(i, j-1) \\ D(i-1, j-1) \end{cases}$$

508   The optimal warping path, $P = \{(i_1, j_1), (i_2, j_2), \ldots, (i_k, j_k)\}$, is determined by

509   tracing back from $D(n, m)$ to $D(1,1)$, selecting the path that minimizes the total

510   cumulative cost. The DTW distance between the two time series is defined as the total

511   cost along this optimal path:

$$DTW(X, Y) = D(n, m)$$

For each participant, we constructed two time series: Micro-saccade bias time

series – calculated as the difference between the number of micro-saccades directed

toward the high-probability distractor location and those directed away from it.

Learning suppression time series – measured as the reaction time (RT) difference

between trials where the distractor appeared at the high- and low-probability locations.

To allow direct comparison, both time series were z-scored, standardizing them to the

same scale. DTW was then applied to the average of these normalized series to

evaluate their temporal alignment.

We assessed significance using a non-parametric permutation test by randomly

shuffling group labels 1000 times, recalculating the DTW distances between shuffled

time-series averages, and comparing the observed distance to this null distribution.

The p-value was calculated as the proportion of permutations in which the shuffled

distance was smaller than the observed distance.

EEG recording and preprocessing

EEG data were recorded using 64 Ag/AgCl active electrodes connected to

BrainAmp amplifiers (Brain Products, Munich, Germany), positioned based on the

extended 10-20 system, and sampled at 500 Hz. Notably, horizontal EOG was

recorded from electrodes placed approximately 1 cm lateral to the outer canthi of the

left and right eyes, using electrodes originally assigned to T7 and FT9. Vertical EOG

was recorded from electrodes placed above and below one eye, using electrodes

originally assigned to T8 and FT10. Electrodes originally assigned to TP9 and TP10

were used to record signals from the mastoids, and the electrode originally assigned to

FCz served as the online reference.

The data underwent re-referencing to the mean of the left and right mastoids and

were subjected to high-pass filtering with a cutoff frequency of 1.5 Hz (for

independent component analysis [ICA] only) and 0.1 Hz (for subsequent analyses).

Continuous EEG recordings were epoched from -3500 to 2500 ms relative to the

search array onset. Malfunctioning electrodes were visually detected and temporally

removed from the data; and a 110–140Hz bandpass filter was used to capture muscle

activity[61] and allowed for variable z-score cutoffs per participant based on the

within-subject variance of z-scores. After trial rejection, ICA was performed only on

the clean electrodes. Components associated with eye blinks, eye movements, or other

non-neural artifacts were visually inspected and removed. Specifically, we excluded

components that showed: (1) high-amplitude deflections coinciding with blink events

in the EOG channel, (2) a frontal scalp topography with polarity reversals

characteristic of ocular activity, and (3) temporal waveform dominated by sporadic,

high-amplitude spikes. Subsequently, we interpolated the malfunctioning electrodes

identified earlier.

Time-frequency analysis

551   Preprocessed EEG data were segmented into event-related epochs (-3500 ms to

552   2500 ms relative to search array onset; avoiding edge artifacts), and were bandpass

553   filtered using a two-way least-squares finite-impulse-response filter[62]. A Hilbert

554   transform (MATLAB Signal Processing Toolbox) was applied to the bandpass-filtered

555   data to produce the complex analytic signal, $z(t)$, of the filtered EEG signals, $f(t)$:

$$z(t) = f(t) + i\tilde{f}(t)$$

where $\tilde{f}(t)$ is the Hilbert transform of $f(t)$, and $i = \sqrt{-1}$. The complex

analytic signal was extracted for each electrode.

We applied the Hilbert transform on the bandpass filtered data from 8 Hz to 14

Hz in steps of 1 Hz. Total alpha power was computed by squaring the complex

magnitude of complex analytic signal, followed by averaging it across trials. The

power was then z-scored by subtracting the average value and dividing by the

standard deviation across all trials within the focused epoch (-2500 ms to 1000 ms).

We selected 17 posterior parietal and occipital electrodes (Pz, P3, P7, O1, Oz, O2, P4,

P8, P1, P5, PO7, PO3, POz, PO4, PO8, P6, P2) for all further analysis (see Fig. 2e).

Inverted encoding model

To reconstruct spatial representation of distractor locations, we applied an

inverted encoding model (IEM) to estimate spatial channel-tuning functions (CTFs)

from alpha-band activity over time[41]. The assumption is that the oscillatory activity

recorded from each electrode reflects the weighted sum of six spatially selective

channels, each tuned to a specific angular position (corresponding to the possible

locations of the distractors). The response profile of each spatial channel across

distractor locations was modeled as a half sinusoid raised to the seventh power:

$$R = \sin(0.5\theta)^7$$

where θ is the angular location (ranging from 0° to 359°), and R is the response

of the spatial channel in arbitrary units. To ensure that the peak response of each

spatial channel was centered over one of the six distractor locations, the response

profile was circularly shifted.

We partitioned the data randomly into independent sets of training (2/3 trials) and

test data (1/3 trials) following a cross-validation routine. The training data (B; m

electrodes × n locations) were used to estimate the weights that approximated the

possible contributions of the six spatial channels to the observed oscillatory activity

measured at each electrode and distractor location. We defined C (k channels × n

locations) as a matrix of the predicted response of each spatial channel (estimated by

the basic function for that channel) for each location; and W (m electrodes × k

channels) as a weight matrix characterizing a linear mapping from "channel space" to

"electrode space". We further described the relationships between B, C, and W in a

linear model as follows:

$$B = WC$$

Using the weight matrix (W) obtained via least-squares estimation, we inverted

the model to transform the observed test data into estimated channel responses (i.e.,

reconstructed CTFs). This IEM routine was iterated 10 times to minimize the

influence of idiosyncrasies specific to any trial assignment and obtain averaged

channel-response profiles. We then calculated the slope of the CTFs via linear

regression to quantify attentional processing for different distractor locations over

time. Higher slope values indicate greater spatial selectivity while lower values

indicate less spatial selectivity. Distractor-tuning CTFs were reconstructed according

to the distractor location across participants. The data were down-sampled to 50 Hz to

reduce computational time. The time-resolved CTF slopes were smoothed using a

Gaussian filter with a standard deviation (σ) of 5.

Micro-saccade-locked activity

To analyze micro-saccade-locked activity within the EEG signals, we first aligned

all EEG data to the onset of micro-saccades within the pre-stimulus period (-2500 ms

to 0 ms relative to search array onset). The EEG data were then epoched from -1000

ms to 2000 ms relative to the onset of each micro-saccade. Saccade-related potential

(SRP) waveforms were extracted from -500 to 1000 ms relative to micro-saccade

onset. For each participant, data from the selected electrodes were averaged across

trials and transformed into the frequency domain using fast Fourier transform (FFT).

Power spectral density (PSD) was then computed within 1–40 Hz. To normalize for

inter-individual variability, PSD values were z-scored across frequencies for each

participant. Statistical significance was assessed using one-sample right-tailed t-tests

at each frequency (1–40 Hz) across participants, testing whether normalized spectral

power exceeded zero.

We also applied the same Hilbert transform methods as before, but in this case,

we focused on calculating evoked (phase-locked) power, instead of total power. This

allowed us to examine neural responses that were phase-locked to the micro-saccade

events and to capture the specific neural dynamics associated with these tiny eye

movements.

Classification analysis

We employed a Support Vector Machine with error-correcting output codes

(SVM-ECOC) to create two classifiers[63], aimed at classifying whether anticipatory

micro-saccades were directed toward or away from the high-probability location,

based on micro-saccade-locked activities. Notably, micro-saccades directed toward

the high-probability location were evenly distributed, with half occurring to the left

and half to the right; the same pattern was set for micro-saccades directed away from

that location. This balanced distribution ensures that the classification analysis

provides a reliable test of whether micro-saccade–locked activity carried information

about the expected location. This was implemented using the MATLAB function

'fitcecoc'. Due to significant variability in the number of micro-saccade-locked EEG

epochs across participants, subject-level analysis was not feasible. However,

assuming that responses from the same electrodes across different participants should

be consistent under the same cognitive process, we concatenated all epochs across all

participants. Specifically, with a threefold cross-validation procedure, 2/3 of epochs

were randomly selected for model training, and the remaining 1/3 were used for

testing. To reduce biases from specific grouping, 100 iterations of this procedure were

conducted with distinct randomizations. Additionally, the data were down-sampled to

50 Hz to reduce computational time. The time-resolved classification accuracies were

smoothed using a Gaussian filter with a standard deviation (σ) of 5.

Statistical analysis

To correct for multiple comparisons, we used a cluster-based permutation test[64]

against a null distribution shuffled from 1000 iterations (following the Monte Carlo

randomization procedure) for each temporal-level analysis. Specifically, one-sample

t-tests were performed across participants for conditional difference (or

single-condition activity) against zeros to identify above-chance activity. Time

windows with t values larger than a threshold ($p = .05$) were combined into

contiguous clusters based on adjacency. The cluster statistics was defined as the sum

of the t values within each cluster. The null distribution was formed by randomly

permuting condition labels for 1000 times to get the largest clusters per iteration.

Clusters were determined to be significant if the cluster statistics were larger than the

95th percentile of the null distribution.

For the classification analysis, since subject-level t-tests could not be performed,

we employed a non-parametric approach by repeating the SVM-ECOC procedure

1000 times while randomizing the labels of micro-saccade directions. For each time

point, a null distribution of classification accuracy was generated. Classification

accuracies were determined to be significant if they exceeded the 95th percentile of

the null distribution. To correct for multiple comparisons across time, we used a

cluster-based permutation procedure. For each of 1000 permutations, supra-threshold

time points (exceeding the 95th percentile of the null distribution) were identified, and

the length of the longest contiguous cluster was recorded. This yielded a null

distribution of maximum cluster lengths. In the original (unshuffled) data, clusters

were considered significant if their length exceeded the 95th percentile of this null

distribution.

Moreover, for multiple t-tests, we controlled for false positives using the False

Discovery Rate (FDR) correction. For key non-significant comparisons, we also

calculated Bayes Factors to assess evidence for the null hypothesis. BF_{10} values < 1

indicate support for the null, whereas values > 1 indicate support for the alternative.

In our analyses, BF_{10} values of 0.74 and 0.84 provided only anecdotal (i.e., weak and

inconclusive) evidence in favor of the null hypothesis.

References

1. Ferrante, O. *et al.* Altering spatial priority maps via statistical learning of target selection and distractor filtering. *Cortex* **102**, 67–95 (2018).

2. Jiang, Y. V., Swallow, K. M., Rosenbaum, G. M. & Herzig, C. Rapid acquisition but slow extinction of an attentional bias in space. *J. Exp. Psychol. Hum. Percept. Perform.* **39**, 87–99 (2013).

3. Sha, L. Z., Remington, R. W. & Jiang, Y. V. Short-term and long-term attentional biases to frequently encountered target features. *Atten. Percept. Psychophys.* **79**, 1311–1322 (2017).

4. Theeuwes, J. Attentional Capture and Control. *Annu. Rev. Psychol.* **76**, 251–273 (2025).

5. Theeuwes, J., Bogaerts, L. & Moorselaar, D. van. What to expect where and when: how statistical learning drives visual selection. *Trends Cogn. Sci.* **26**, 860–872 (2022).

6. Goschy, H., Bakos, S., Müller, H. J. & Zehetleitner, M. Probability cueing of distractor locations: both intertrial facilitation and statistical learning mediate interference reduction. *Front. Psychol.* **5**, (2014).

7. Leber, A. B., Gwinn, R. E., Hong, Y. & O'Toole, R. J. Implicitly learned suppression of irrelevant spatial locations. *Psychon. Bull. Rev.* **23**, 1873–1881 (2016).

8. Sauter, M., Liesefeld, H. R., Zehetleitner, M. & Müller, H. J. Region-based shielding of visual search from salient distractors: Target detection is impaired with same- but not different-dimension distractors. *Atten. Percept. Psychophys.* **80**, 622–642 (2018).

9. Wang, B. & Theeuwes, J. Statistical regularities modulate attentional capture. *J. Exp. Psychol. Hum. Percept. Perform.* **44**, 13–17 (2018).

10. Wang, B. & Theeuwes, J. How to inhibit a distractor location? Statistical learning versus active, top-down suppression. *Atten. Percept. Psychophys.* **80**, 860–870 (2018).

11. Wang, B. & Theeuwes, J. Statistical regularities modulate attentional capture independent of search strategy. *Atten. Percept. Psychophys.* **80**, 1763–1774 (2018).

12. Luck, S. J., Gaspelin, N., Folk, C. L., Remington, R. W. & Theeuwes, J. Progress toward resolving the attentional capture debate. *Vis. Cogn.* **29**, 1–21 (2021).

13. Huang, C., Vilotijević, A., Theeuwes, J. & Donk, M. Proactive distractor suppression elicited by statistical regularities in visual search. *Psychon. Bull. Rev.* **28**, 918–927 (2021).

14. Huang, C., Donk, M. & Theeuwes, J. Attentional suppression is in place before display onset. *Atten. Percept. Psychophys.* **85**, 1012–1020 (2023).

15. Kong, S., Li, X., Wang, B. & Theeuwes, J. Proactively location-based suppression elicited by statistical learning. *PLOS ONE* **15**, e0233544 (2020).

16. Munneke, J., Heslenfeld, D. J., Usrey, W. M., Theeuwes, J. & Mangun, G. R. Preparatory Effects of Distractor Suppression: Evidence from Visual Cortex.

PLOS ONE **6**, e27700 (2011).

17. Richter, D., Moorselaar, D. van & Theeuwes, J. Proactive distractor suppression
in early visual cortex. 2024.04.03.587747 Preprint at
https://doi.org/10.1101/2024.04.03.587747 (2024).

18. Wang, B., van Driel, J., Ort, E. & Theeuwes, J. Anticipatory Distractor
Suppression Elicited by Statistical Regularities in Visual Search. *J. Cogn.*
Neurosci. **31**, 1535–1548 (2019).

19. Beck, V. M., Luck, S. J. & Hollingworth, A. Whatever you do, don't look at
the . . .: Evaluating guidance by an exclusionary attentional template. *J. Exp.*
Psychol. Hum. Percept. Perform. **44**, 645–662 (2018).

20. Humphreys, G. W., Stalmann, B. J. & Olivers, C. An analysis of the time course
of attention in preview search. *Percept. Psychophys.* **66**, 713–730 (2004).

21. Lahav, A., Makovski, T. & Tsal, Y. White bear everywhere: Exploring the
boundaries of the attentional white bear phenomenon. *Atten. Percept. Psychophys.*
74, 661–673 (2012).

22. Lahav, A. & Tsal, Y. Allocating Attention to Distractor Locations is Based on
Top-Down Expectations. *Q. J. Exp. Psychol.* **66**, 1873–1880 (2013).

23. Makovski, T. Preparing for distraction: Attention is enhanced prior to the

presentation of distractors. *J. Exp. Psychol. Gen.* **148**, 221–236 (2019).

24. Moher, J. & Egeth, H. E. The ignoring paradox: Cueing distractor features leads
first to selection, then to inhibition of to-be-ignored items. *Atten. Percept.*
Psychophys. **74**, 1590–1605 (2012).

25. Tsal, Y. & Makovski, T. The attentional white bear phenomenon: The mandatory
allocation of attention to expected distractor locations. *J. Exp. Psychol. Hum.*
Percept. Perform. **32**, 351–363 (2006).

26. Won, B.-Y., Kosoyan, M. & Geng, J. J. Evidence for second-order singleton
suppression based on probabilistic expectations. *J. Exp. Psychol. Hum. Percept.*
Perform. **45**, 125–138 (2019).

27. Klein, R. M. Inhibition of return. *Trends Cogn. Sci.* **4**, 138–147 (2000).

28. Wang, B. & Theeuwes, J. Salient distractor processing: inhibition following
attentional capture. *Trends Cogn. Sci.* **28**, 593–594 (2024).

29. Engbert, R. & Kliegl, R. Microsaccades uncover the orientation of covert
attention. *Vision Res.* **43**, 1035–1045 (2003).

30. Jiang, Y. V., Won, B.-Y. & Swallow, K. M. First saccadic eye movement reveals
persistent attentional guidance by implicit learning. *J. Exp. Psychol. Hum. Percept.*
Perform. **40**, 1161–1173 (2014).

31. van Ede, F., Chekroud, S. R. & Nobre, A. C. Human gaze tracks attentional
focusing in memorized visual space. *Nat. Hum. Behav.* **3**, 462–470 (2019).

32. Yuval-Greenberg, S., Merriam, E. P. & Heeger, D. J. Spontaneous Microsaccades
Reflect Shifts in Covert Attention. *J. Neurosci.* **34**, 13693–13700 (2014).

33. Denison, R. N., Yuval-Greenberg, S. & Carrasco, M. Directing Voluntary
Temporal Attention Increases Fixational Stability. *J. Neurosci.* **39**, 353–363
(2019).

34. De Vries, E. & Van Ede, F. Microsaccades Track Location-Based Object Rehearsal in Visual Working Memory. *eneuro* **11**, ENEURO.0276-23.2023 (2024).

35. Theeuwes, J. Perceptual selectivity for color and form. *Percept. Psychophys.* **51**, 599–606 (1992).

36. Lin, R. *et al.* Neural evidence for attentional capture by salient distractors. *Nat. Hum. Behav.* **8**, 932–944 (2024).

37. Lin, R., Li, X., Wang, B. & Theeuwes, J. Spatial suppression due to statistical learning tracks the estimated spatial probability. *Atten. Percept. Psychophys.* **83**, 283–291 (2021).

38. Contadini-Wright, C., Magami, K., Mehta, N. & Chait, M. Pupil Dilation and Microsaccades Provide Complementary Insights into the Dynamics of Arousal and Instantaneous Attention during Effortful Listening. *J. Neurosci.* **43**, 4856–4866 (2023).

39. Rolfs, M., Engbert, R. & Kliegl, R. Crossmodal coupling of oculomotor control and spatial attention in vision and audition. *Exp. Brain Res.* **166**, 427–439 (2005).

40. Hafed, Z. M. & Clark, J. J. Microsaccades as an overt measure of covert attention shifts. *Vision Res.* **42**, 2533–2545 (2002).

41. Foster, J. J., Sutterer, D. W., Serences, J. T., Vogel, E. K. & Awh, E. The topography of alpha-band activity tracks the content of spatial working memory. *J. Neurophysiol.* **115**, 168–177 (2016).

42. Jensen, O. & Mazaheri, A. Shaping Functional Architecture by Oscillatory Alpha Activity: Gating by Inhibition. *Front. Hum. Neurosci.* **4**, (2010).

43. Dimigen, O., Valsecchi, M., Sommer, W. & Kliegl, R. Human Microsaccade-Related Visual Brain Responses. *J. Neurosci.* **29**, 12321–12331 (2009).

44. Liu, B., Nobre, A. C. & van Ede, F. Microsaccades transiently lateralise EEG alpha activity. *Prog. Neurobiol.* **224**, 102433 (2023).

45. Meyberg, S., Werkle-Bergner, M., Sommer, W. & Dimigen, O. Microsaccade-related brain potentials signal the focus of visuospatial attention. *NeuroImage* **104**, 79–88 (2015).

46. Keren, A. S., Yuval-Greenberg, S. & Deouell, L. Y. Saccadic spike potentials in gamma-band EEG: Characterization, detection and suppression. *NeuroImage* **49**, 2248–2263 (2010).

47. Abeles, D., Amit, R., Tal-Perry, N., Carrasco, M. & Yuval-Greenberg, S. Oculomotor inhibition precedes temporally expected auditory targets. *Nat. Commun.* **11**, 3524 (2020).

48. Badde, S., Myers, C. F., Yuval-Greenberg, S. & Carrasco, M. Oculomotor freezing reflects tactile temporal expectation and aids tactile perception. *Nat. Commun.* **11**, 3341 (2020).

49. Dankner, Y., Shalev, L., Carrasco, M. & Yuval-Greenberg, S. Prestimulus Inhibition of Saccades in Adults With and Without Attention-Deficit/Hyperactivity Disorder as an Index of Temporal Expectations. *Psychol. Sci.* **28**, 835–850 (2017).

50. Hung, S.-C. & Carrasco, M. Microsaccades as a long-term oculomotor correlate in visual perceptual learning. *Psychon. Bull. Rev.* **30**, 235–249 (2023).

51. Ruff, C. C. & Driver, J. Attentional Preparation for a Lateralized Visual Distractor: Behavioral and fMRI Evidence. *J. Cogn. Neurosci.* **18**, 522–538 (2006).

52. Serences, J. T., Schwarzbach, J., Courtney, S. M., Golay, X. & Yantis, S. Control of Object-based Attention in Human Cortex. *Cereb. Cortex* **14**, 1346–1357 (2004).

53. Theeuwes, J., Olivers, C. N. L. & Chizk, C. L. Remembering a Location Makes the Eyes Curve Away. *Psychol. Sci.* **16**, 196–199 (2005).

54. Van der Stigchel, S., Meeter, M. & Theeuwes, J. The spatial coding of the inhibition evoked by distractors. *Vision Res.* **47**, 210–218 (2007).

55. Theeuwes, J. The Attentional Capture Debate: When Can We Avoid Salient Distractors and When Not? *J. Cogn.* **6**, 35.

56. Willett, S. M. & Mayo, J. P. Microsaccades are directed toward the midpoint between targets in a variably cued attention task. *Proc. Natl. Acad. Sci.* **120**, (2023).

57. Popov, T., Gips, B., Kastner, S. & Jensen, O. Spatial specificity of alpha oscillations in the human visual system. *Hum. Brain Mapp.* **40**, 4432–4440 (2019).

58. Barczak, A. *et al.* Dynamic Modulation of Cortical Excitability during Visual Active Sensing. *Cell Rep.* **27**, 3447-3459.e3 (2019).

59. Engbert, R. & Mergenthaler, K. Microsaccades are triggered by low retinal image slip. *Proc. Natl. Acad. Sci.* **103**, 7192–7197 (2006).

60. Laubrock, J., Engbert, R. & Kliegl, R. Microsaccade dynamics during covert attention. *Vision Res.* **45**, 721–730 (2005).

61. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Comput. Intell. Neurosci.* **2011**, 1–9 (2011).

62. Delorme, A. & Makeig, S. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J. Neurosci. Methods* **134**, 9–21 (2004).

63. Bae, G.-Y. & Luck, S. J. Dissociable Decoding of Spatial Attention and Working Memory from EEG Oscillations and Sustained Potentials. *J. Neurosci.* **38**, 409–422 (2018).

64. Maris, E. & Oostenveld, R. Nonparametric statistical testing of EEG- and MEG-data. *J. Neurosci. Methods* **164**, 177–190 (2007).