Functional Characterization of Paillotin: An Immune Peptide Regulated by the

Imd Pathway with Pathogen-Specific Roles in *Drosophila* Immunity

Yao Tian[1], Xiaojing Yue[3], Renjie Jiao[4], Mark A. Hanson[1,2]*, Bruno Lemaitre[1]*

1: Global Health Institute, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne,

Switzerland

2: Centre for Ecology and Conservation, University of Exeter, Penryn, United Kingdom

3: Laboratory of Biology and Modelling of the Cell, Ecole Normale Superieure of Lyon, CNRS,

Lyon, France.

4: Sino-French Hoffmann Institute, Guangzhou Medical University, Guangzhou, China

*: Equal last author: M.A. Hanson (M.Hanson@exeter.ac.uk), B.L. Lemaitre

(bruno.lemaitre@epfl.ch)

Abstract

Insects, such as *Drosophila* melanogaster, rely on innate immune defenses to combat

microbial threats. Antimicrobial peptides (AMPs) play an important role in limiting pathogen entry

and colonization. Despite intensive research into the regulation and biochemical properties of

AMPs, their exact significance in vivo has remained uncertain due to the challenges of mutating

small genes. Fortunately, recent technologies have enabled the mutation of individual AMP

genes, overcome previous obstacles, and opened new avenues for research. In this study, we

characterized one novel host-defense peptide, Paillotin (IM18, *CG33706*), using loss-of-

function mutants. Paillotin is an ancient host defense peptide of Diptera, regulated by the Imd

pathway. Loss of Paillotin does not impact the activity of either the Imd or Toll pathways.

Importantly, we found that *Paillotin* mutants are viable but exhibit increased susceptibility to

specific infections, particularly *Providencia burhodogranariea*. Paillotin was further found to

contribute synergistically to defense against *P. burhodogranariea* when combined with other

AMPs. However, we did not detect direct microbicidal activity of Paillotin *in vitro* in our hands.

Taken together, our findings identify Paillotin as a novel host defense peptide acting

downstream of Imd signaling, advancing our understanding of the *Drosophila* antimicrobial

response.

Keywords: antimicrobial peptide, host–pathogen interactions, *Drosophila*, immunity

Introduction

Insects inhabit environments rich in microbes, necessitating a robust immune system to

defend against opportunistic pathogens [1]. Unlike vertebrates, insects rely solely on innate

immune defense mechanisms. The conservation of innate immunity across *Metazoans* and the

powerful genetics of *Drosophila melanogaster* make this insect a powerful model for studying

immune responses [2–4].

Flies exhibit behavioral immunity to prevent pathogens from entering their body cavity,

such as grooming and avoidance of pathogens [5,6]. Once pathogens breach this barrier,

immune responses are activated in different body regions to limit growth. Local immunity

occurs at initial pathogen entry sites, like the gut or trachea. Systemic immunity involves

cellular and humoral responses that take place in the body cavity upon systemic infection.

Cellular responses involve hemocytes (blood cells), while humoral responses involve the

production of host defense peptides as well as other effectors secreted by hemocytes and the fat

body, an organ analogous to the mammalian liver [1,2]. Two NF-κB signaling pathways, the

Toll and Imd pathways, regulate the humoral response. The principal effectors downstream of

the Toll and Imd signaling pathways consist of AMPs, which are synthesized by both the fat

body and hemocytes and then secreted into the hemolymph. AMPs are positively charged and

can bind to the negatively charged membranes of microorganisms, disrupting membrane

integrity through pore formation [1,7]. Upon infection, their expression is markedly increased

to high levels in *Drosophila*, notably *Drosomycin* and *Diptericin,* which are often used as

readouts for Toll and Imd activity, respectively [8]. Besides microbe killing, AMPs have been

implicated in multiple processes such as tumor suppression [9,10], microbiota control [11], and

neurodegeneration [12,13]. Currently, *D. melanogaster* is known to possess eight families of

inducible AMPs, including antifungal peptides like Drosomycin, Baramicin, and Metchnikowin;

Cecropins (four isoforms) exhibiting both antibacterial and antifungal activities *in vitro*; and

Drosocin, Attacins (four isoforms), Defensin, and Diptericins (two isoforms), primarily

displaying antibacterial activity [14–21]. Additionally, the *Drosophila* genome encodes

numerous other host defense peptides, such as the Toll-regulated *Daisho* genes (two genes),

Bomanins (twelve genes), peptides encoded by *Baramicin A*, and the Imd-regulated peptide Buletin (product of the *Drosocin* gene), whose antimicrobial activity *in vitro* remains to be fully demonstrated, but functional studies have indicated their importance *in vivo* for surviving microbial infections [22–26].

Functional studies using single or combined mutations have demonstrated that these host defense peptides largely account for the critical roles of the Imd and Toll pathways in host survival following infection with Gram-negative bacteria, Gram-positive bacteria, or fungi. The antibacterial peptides Diptericin, Cecropin, Attacin, Drosocin, and Buletin account for most of the Imd pathway–mediated resistance to Gram-negative bacterial infection [25,27–29]. Drosomycin, Metchnikowin, Daisho (two genes), Baramicin A mediate much of the Toll-mediated resistance to fungal infection [23,24,26,27,30] , and Bomanins (twelve genes) appear to be the most important family of Toll-regulated host defense peptides contributing to resistance against Gram-positive bacteria and fungi [22,27,31]. These host defense peptides display both additive and synergistic effects, along with a high degree of specificity, whereby individual peptides greatly contribute to survival against a defined pathogen [27,29,32]. Some immune-induced peptides regulated by Toll and Imd have regulatory functions. This includes GNBP-like3 and Ibin A/B which have been reported to modulate Toll pathway activity, as well as Edin, which has been implicated in regulating hemocyte sessility [33–35].

Despite significant advances in understanding *Drosophila* host defense peptides, many peptides remain underexplored. The *Drosophila* genome encodes several other immune-induced peptides that remain poorly characterized, including notably Listericin, CG4269, and IM18 [1,36,37].

In this article, we have characterized the function of IM18, which we renamed Paillotin. We showed that Paillotin is an ancient host defense peptide regulated by the Imd pathway, which does not affect the regulation of the Imd or Toll pathways. Importantly, we found that *Paillotin* mutants exhibit increased susceptibility to specific infections, particularly *Providencia burhodogranariea*. Paillotin was also found to contribute synergistically to defense against *P. burhodogranariea* when combined with other AMPs, specifically Drosocin, Diptericins, and Attacins. However, we did not detect direct microbicidal activity of Paillotin *in vitro*. Taken together, our findings identify Paillotin as a novel host defense peptide

88   downstream of Imd signaling, advancing our understanding of the *Drosophila* antimicrobial

89   response.

Results

Paillotin is a conserved host defense peptide in *Diptera*

A previous proteomic analysis identified an immune inducible peptide called Immune-

induced Molecule 18 (IM18) based on its mass by MALDI-TOF [38]. We rename the *IM18*

gene (*CG33706*) as *Paillotin* (*Pai*) from André Paillot, who in 1920 discovered the humoral

immune response of insects [39,40]. The gene encoding Paillotin was identified as *CG33706* in

a region annotated as polycistronic with another gene, *CG10332*, making it challenging to

distinguish their expression patterns (Figure 1*a*); the *CG10332-RA* transcript is identical to the

Pai-RA transcript and contains the full coding sequence of both *CG10332* and *Pai*. Indeed, in

FlyAtlas data, *IM18* and *CG10332* are shown with identical expression levels [41]. However,

the characterization of IM18 as an immune-induced peptide pointed to the existence of two

distinct gene products, perhaps regulated by the distinct transcripts *Pai-RB* and *CG10332-RA*.

To confirm this, we monitored the expression of either *CG10332-RA* alone or the cumulative

products of *CG10332-RA* and *Pai-RB* upon septic injury with either the Gram-negative

bacterium *Escherichia coli* (*E. coli*) or the Gram-positive bacterium *Micrococcus luteus* (*M.*

luteus). The results showed that the mRNA encoding *Pai-RB* (Figure 2*a*, *b*) but not *CG10332-*

RA was strongly induced upon septic injury (Figure 2*c*, *d*), confirming that only *Pai-RB* is

immune-induced at the transcriptional level.

The *Paillotin* gene encodes a 71-residue precursor peptide composed of a signal peptide,

dipeptidylpeptidase motif, and mature peptide (Figure 1*b*). To investigate the evolutionary

conservation of Paillotin, we used a recursive tBLASTn approach to screen outgroup fly species

[42]. Orthologues of Paillotin were identified in *Diptera* families such as *Tephritidae* and

Diopsidae, which share a common ancestor with *D. melanogaster* approximately 126 million

years ago [43] (Figure 1*b*). The C-terminal domain displayed the highest level of conservation

across species, suggesting a critical role in Paillotin function. The presence of conserved

features, such as the two-residue dipeptidyl peptidase cleavage site (XA/XP) and a positive

charge (+3.89 in *D. melanogaster*) commonly found in AMPs [24,44], supports the hypothesis

that Paillotin plays a key role in host defense and could have microbicidal activities. This

evolutionary conservation underscores Paillotin as an ancient host defense peptide in *Diptera*.

Paillotin is regulated by the Imd pathway

To explore the expression pattern and regulation of *Paillotin* during systemic immune

responses, we analyzed transcript levels in wild-type, Toll, and Imd pathway-deficient flies

(Figure 2). Specifically, we challenged w^{1118} (wild-type), spz^{rm7} (deficient for the Toll pathway),

$Relish^{E20}$ (deficient for Imd pathway) and $Relish^{E20}$, spz^{rm7} double mutant flies with either *E. coli,*

an inducer of the Imd pathway (Figure 2*a*), or *M. luteus,* an inducer of the Toll pathway (Figure

2*b*). Expression of *Paillotin* was monitored at 8h, 16h, and 24h post-infection using qRT-PCR.

We observed that *Paillotin* gene expression was strongly induced after both *M. luteus* and *E.*

coli infections in w^{1118} flies at all time points. However, this induction was completely abolished

in $Relish^{E20}$ and $Relish^{E20}$, spz^{rm7} double mutants, but not in spz^{rm7} flies, indicating *Paillotin* is

regulated by the Imd pathway (Figure 2*a*, *b*). Consistent with this expression profile, we found

multiple putative NF-κB responsive motifs within the first 200bp of promoter sequence

upstream of gene transcription start sites of *Paillotin* [24,45] including five motifs uniquely

matching the *Relish* consensus motif sequence (GGRDNNHHBS), one motif matching a

Dif/Dorsal (GGGHHNNDVH) binding site, and two κB motifs with putative matches to either

Relish and/or *Dif/Dorsal* binding sites. In contrast, no NF-κB binding motifs were identified in

the 250bp upstream region of *IM18-RA/CG10332-RA*, consistent with its non-immune

inducibility.

A recent proteomic analysis of *Drosophila* hemolymph following bacterial infection [46]

indicated that Paillotin peptide was strongly induced in the hemolymph of wild-type flies

infected with *M. luteus* or the Gram-negative bacterium *Pectobacterium carotovorum* (*Ecc15*),

and this induction was abolished in $Relish^{E20}$ flies (Figure 2*e*). Consistent with its regulation by

the Imd pathway, *Paillotin* was more induced upon infection with *Ecc15* than *M. luteus*. We

conclude that Paillotin is a novel host defense peptide regulated at the transcriptional level by

the Imd pathway and secreted into the hemolymph.

Identification and characterization of *Paillotin* mutants

The *Drosophila* genome nexus represents a global sampling of the genetic diversity of *D. melanogaster* wild types subjected to genome sequencing [47]. Using a combination of the genome nexus and more detailed variant information in the *Drosophila* Genetic Reference Panel (DGRP) [48], we identified two strains from Kenya (KR7, KR4N) encoding a premature stop codon in *Paillotin* ("Gly37∗", numbering per start codon). Globally, we could assess the *Paillotin* region in 789 genome-sequenced strains, finding 39 that encode putative loss-of-function mutations (Table S1). The majority of these mutant strains were collected from Winters California and Raleigh North Carolina, which have a high frequency of putative loss-of-function mutations (~26% and ~8% respectively within these populations). We characterized two of the loss-of-function mutations segregating in the DGRP further (Figure 3*a*). The first mutation (2R_19488505_DEL) is found in 17 genotypes from the DGRP, including DGRP-41, which deletes 61 nucleotides within the Paillotin mature peptide coding region, leading to a frameshift and protein truncation (referred to as $Pai^{\triangle 41}$). The second mutation (2R_19488690_SNP), found in DGRP-370 ($Pai^{\triangle 370}$), involves a C-T transition that disrupts the *Paillotin* start codon. We utilized these naturally-occurring mutations to study the role of *Paillotin* in host defense by backcrossing $Pai^{\triangle 41}$ and $Pai^{\triangle 370}$ into the w^{1118} iso DrosDel background for seven generations.

In addition, we used a third mutant referred to as $Pai^{\triangle SFH}$, a gift from the Guangzhou Drosophila Resource Center (Figure 3*a*), which was generated by CRISPR-Cas9 and involves a 20-nucleotide deletion that causes a complex frameshift, resulting in a nonsense peptide. The precise nucleotide alterations associated with each *Paillotin* mutant allele used in this study are presented in Supplementary Figure S1. For $Pai^{\triangle SFHI}$, only the *X* chromosome was replaced by the w^{1118} iso DrosDel *X* chromosome.

RT-qPCR confirmed reduced *Paillotin* gene expression in $Pai^{\triangle 41}$ and $Pai^{\triangle 370}$, while *Paillotin* expression was completely absent in $Pai^{\triangle SFHI}$ (Figure S2). We next performed a MALDI-TOF analysis of hemolymph from adult flies infected with *P. c. carotovara Ecc15*. The results revealed the absence of the 4625Da Paillotin peaks in $Pai^{\triangle 41}$, $Pai^{\triangle 370}$, and $Pai^{\triangle SFHI}$ flies (Figure 3*b*). All the mutants were viable without morphological defects.

Paillotin does not regulate the Imd and Toll pathways

Some host defense peptides, such as mammalian LL37 or β-Defensins, have been shown to function similarly to cytokines, modulating immune system activity [49,50]. This prompted us to assess whether Paillotin could influence Toll and Imd pathway activity. To address this, we monitored the expression of *Diptericin A* (a readout of the Imd pathway) in wild-type (w^{1118}) and *Paillotin* -deficient flies ($Pai^{\triangle 41}$ and $Pai^{\triangle 370}$) at 6- and 12-hours post-infection with *Ecc15* and *Drosomycin* (a readout of the Toll pathway) at 8- and 24-hours post-infection with *M. luteus* (Figure 4*a*, *b*). The expression levels of *Diptericin A* in $Pai^{\triangle 41}$ and $Pai^{\triangle 370}$ mutants were comparable to those in wild-type flies at both time points, indicating no effect of *Paillotin* on Imd pathway activity. Similarly, the expression of *Drosomycin* in $Pai^{\triangle 41}$ and $Pai^{\triangle 370}$ mutants showed no significant difference compared to w^{1118}, suggesting that *Paillotin* is not involved in regulating the Toll pathway. Together, these results demonstrate that *Paillotin* is not a regulator of the Imd or Toll pathways.

Paillotin mutant flies are susceptible to *P. burhodogranariea* infection

Recent studies have demonstrated that the deletion of a single AMP gene can significantly increase susceptibility to specific pathogens. For instance, the *Diptericin A* mutation leads to marked susceptibility to *Providencia rettgeri*, while the *Diptericin B* mutation greatly reduces resistance to *Acetobacter sp.* [29,32]. To investigate the role of Paillotin in resistance to infections, we examined its contribution against a range of Gram-negative, Gram-positive bacteria, and fungi. In these experiments, we compared the survival of $Pai^{\triangle 41}$, $Pai^{\triangle 370}$, and $Pai^{\triangle SFHI}$ flies to that of the wild-type (w^{1118}), $Relish^{E20}$, $Bom^{\triangle 55}$, and spz^{rm7} immune-deficient mutants. Flies were systemically infected by bacteria using a needle dipped in a bacterial pellet, a mode of infection that causes cuticle injury. To exclude a role of *Paillotin* in resistance to injury, we first monitored the susceptibility of *Paillotin* mutants to clean injury, and did not detect any susceptibility after clean injury alone (Figure S3). Regarding Gram-negative bacteria, there was no statistically significant difference in survival between wild-type and *Paillotin* mutants following infections with *P. rettgeri*, *Ecc15*, or *Enterobacter cloacae* (Figure S4*a-c*). However, all three *Paillotin* mutant flies ($Pai^{\triangle 41}$, $Pai^{\triangle 370}$, $Pai^{\triangle SFHI}$) exhibited a modest but significant increase in susceptibility to septic injury with *Providencia burhodogranariea* (*P.*

burhodogranariea) (Figure 5*a*). This susceptibility was less marked than *Relish*[E20] flies,

indicating that other factors downstream of the Imd pathway contribute to survival upon *P.*

burhodogranariea infection. In addition, we observed higher *P. burhodogranariea* bacterial

load in *Paillotin* mutants compared to wild-type, indicating that Paillotin suppresses bacterial

growth (Figure 5*b*). For Gram-positive bacteria, *Paillotin* mutants showed no consistent

increased susceptibility to *Enterococcus faecalis* (Figure 5*c*), *Bacillus subtilis*, *Listeria*

monocytogenes, or *Staphylococcus aureus* infections (Figure S4*d-f*). Similarly, fungal

infections with *Beauveria bassiana* (Figure 5*d*), *Metarhizium anisopliae*, or *Metarhizium*

robertsii did not reveal any increased susceptibility of *Paillotin* mutants compared to wild-type

(Figure S4*g-h*).

No *in vitro* activity of Paillotin peptide

Cecropins were the first inducible AMPs identified in moths [14]. *In vitro* studies have

shown that Cecropins are highly effective against a wide range of Gram-negative bacteria at

concentrations lower than those induced in insects during infection (25–50 μM) [51]. To

investigate the antimicrobial potential of Paillotin *in vitro*, we produced the peptide and

evaluated its activity against *P. burhodogranariea* at 100μM alone or in combination with

sublethal concentrations of Cecropin A. However, Paillotin exhibited no detectable

microbicidal activity against *P. burhodogranariea in vitro* (Figure S5*a*). It also failed to inhibit

the growth of other Gram-negative bacteria, including *Ecc15* and *E. coli* (Figure S5*b, c*). Thus,

we found no activity of Paillotin *in vitro* under these conditions.

Paillotin synergistically contributes to the defense against *P. burhodogranariea*

with other AMPs.

Recent studies have demonstrated synergistic or additive cooperation among AMPs,

notably against *P. burhodogranaria*. While single mutations in *Drosocin*, *Attacins*, or

Diptericins have minimal overall effect, a compound mutant referred to as the "*Group B*" AMP

mutant lacking *Drosocin*, *Attacins A, B, C* and *D*, and *Diptericins A* and *B* [full genotype: *w*[1118]*;*

AttC[Mi]*, Dro-AttAB*[SK2]*, Dpt*[SK1]*; AttD*[SK1]], displays marked susceptibility to *P. burhodogranariea*

infection [27]. To investigate potential interactions between *Paillotin* and these AMPs, we used

genetic crosses to combine the *Paillotin* mutation with *Group B* mutants. We initially infected

Group B mutants, *Paillotin* mutants, and *Group B, Paillotin* compound mutants [full genotype:

w^{1118}; *AttC*^{Mi}, *Dro-AttAB*^{SK2}, *Dpt*^{SK1}, *Pai*] with *P. burhodogranariea*. However, we observed no

synergistic or additive effects, as the introduction of *Paillotin* mutations into *Group B* mutants

did not further increase susceptibility. This lack of an observable effect may be due to the

already high susceptibility of *Group B* AMP mutants to this bacterium, even at a low inoculum

(OD_{600}=0.1), potentially masking any additional contribution from the *Paillotin* mutation

(Figure 6, dashed curves). To circumvent this, we assessed the impact of *Paillotin* homozygous

deficiency in flies heterozygous for *Group B* AMP mutations (*Group B/+*), which depresses the

inducibility of AMPs compared to homozygotes [27]. Notably, in this sensitized genetic

background, the absence of *Paillotin*, regardless of the allele used, significantly increased

susceptibility of *Group B/+* flies to *P. burhodogranariea* (Figure 6*a-c*, solid curves). This co-

occurring loss of resistance reflects a synergistic effect on the Hazard Ratio (HR), evidenced

by increased mortality beyond additive effects (*Paillotin* alone, HR =-0.17, *p* =0.01; *Group B*

alone, HR =-1.06, *p*<0.001; *Paillotin*Group B* interaction, HR =-0.15, *p*=0.02). In addition, a

marginally significant trend consistent with a synergistic effect between *Paillotin* and *Group B*

mutants was also seen for infections with the Gram-negative bacteria *Providencia rustigianii*

and *Providencia vermicola*, but not *E. coli* (Figure S6). This supports the notion that Paillotin

confers a fairly specific resistance to *P. burhodogranariea*, perhaps also extending to other

Providencia species.

Overexpression of Paillotin restores resistance in Paillotin-deficient flies against *P.*

burhodogranariea

The observation that three distinct null mutations in *Paillotin* are associated with a higher

susceptibility and the synergistic effect between Paillotin and other AMPs against infection

provide strong evidence for a role of this host defense peptide in resisting *P. burhodogranariea*.

To reinforce this finding, we performed a genetic rescue experiment by combining the *Paillotin*

mutation with a *UAS-Pai* transgene. We initially observed that overexpression of *Paillotin* in a

wild-type background was not sufficient to enhance resistance to *P. burhodogranariea* (Figure

S7). In contrast, overexpression of *Paillotin* using either the *C564-Gal4* (a Gal4 driver with

strong expression in the fat body) or *Actin-Gal4* (ubiquitous) driver restored pathogen

resistance to near wild-type levels in the *Pai*$^{\triangle 41}$ (Figure 7*a*) and *Pai*$^{\triangle 370}$ (Figure 7*b*) mutant

backgrounds. The observation that three distinct *Pai* alleles confer susceptibility, together with

the results of the rescue experiment, collectively demonstrates that Paillotin acts as a pathogen-

specific host defense peptide essential for restricting *P. burhodogranariea* proliferation.

The Imd and Toll signaling pathways are essential for insect immunity, with double

pathway-deficient flies exhibiting extreme susceptibility to infection due to their inability to

upregulate hundreds of immune effector genes, particularly AMPs [52]. Previous work has

demonstrated that transgenic overexpression of individual AMPs in some cases is sufficient to

restore resistance in *Imd, Toll-deficient* flies [53]. Building on this paradigm, we generated flies

overexpressing *Paillotin* using *Actin-Gal4* or *C564-Gal4* drivers in *Relish*^{E20} mutant flies

(Figure 7*c*). However, *Paillotin* overexpression failed to confer protection against *P.*

burhodogranariea infection. While we cannot exclude that Paillotin was not expressed at a

sufficient level to overcome the extreme susceptibility of *Relish* mutants, this suggests that

Paillotin alone is insufficient and the host requires the presence of other AMPs for the defense

response to function effectively, consistent with the synergistic interactions observed with

Group B/+ heterozygous mutants.

Discussion

In this study, we characterized Paillotin, an evolutionarily conserved host-defense peptide

regulated by the Imd pathway. Use of three *Paillotin* mutants reveals a role of *Paillotin* in the

resistance to *P. burhodogranariea* infection, and perhaps other moderately virulent *Providencia*

species, without altering survival to other tested bacterial or fungal species. Furthermore, the

Paillotin mutation enhances the susceptibility to *P. burhodogranariea* in flies heterozygous for

mutations in *Drosocin*, *Diptericins A and B*, and *Attacins A, B, and C.* Taken together, the

consistent susceptibility of three independent *Paillotin* alleles to *P. burhodogranariea*,

accompanied by elevated bacterial loads, the resistance of *Paillotin*-deficient flies to other

tested pathogens, and the successful rescue of the phenotype by overexpression, all provide

strong evidence that *Paillotin* functions as a host defense peptide specifically effective against

this bacterium. A high degree of specificity has been observed for other host defense peptides,

whose inactivation frequently causes pathogen-specific immune deficiency. This has been

illustrated not only for Diptericin A and B as described previously [29,32], but also for Drosocin,

which provides significant protection against *E. cloacae* [25,27]. The existence of such

specificity between a defined host defense peptide and particular pathogens suggests that these

genes have been evolutionarily maintained under selective pressure imposed by specific

microbes encountered in natural environments [54].

Previous studies have shown that resistance to *P. burhodogranariea* involves multiple

antimicrobial peptides, including Buletin (encoded by the *Drosocin* gene), Diptericins A and B,

and Attacins A, B, C, and D. While single AMP mutations in many genes do not greatly impact

survival following infection with *P. burhodogranariea*, compound mutants lacking *Drosocin*,

Diptericins and *Attacins* exhibit increased susceptibility, suggesting the presence of a

synergistic effect among these peptides [27]. More recently, the absence of Buletin—a peptide

generated by furin cleavage of the Drosocin precursor—results in notable survival differences

following *P. burhodogranariea* infection [25]. Thus, host resistance to *P. burhodogranariea* is

mediated by the collective action of multiple host defense peptides, and our study now adds

Paillotin to this list. Future studies may further reveal additional roles of Paillotin, either alone

or in combination with other host defense peptides, in resistance to other pathogens.

We were unable to detect any direct *in vitro* microbicidal activity of Paillotin against *P.*

burhodogranariea and other bacteria, either alone or in combination with Cecropin. This

reflects a common mismatch between in vitro and in vivo activities. Despite demonstrating

effective antimicrobial activity in vivo in relevant animal models, many AMPs show minimal

or no activity in standard minimal inhibitory concentration (MIC) or minimal microbicidal

concentration (MMC) assays when tested under physiological salt concentrations or in serum-

containing environments [55–58]. Several factors could explain this lack of detectable activity:

i) Paillotin may depend on host-specific factors for its function, such as the requirement for a

cofactor or modulation by environmental cues (e.g., pH or ionic strength), or ii) synergistic

interactions with other immune molecules that are only present *in vivo*. Similar synergistic

mechanisms have been observed in other systems. For instance, in *Galleria mellonella* (the

greater wax moth), an anionic antimicrobial peptide (anionic peptide 2) enhances lysozyme-mediated membrane perforation, facilitating the elimination of *E. coli* [59]. Furthermore, AMPs often enhance each other's efficacy through complementary mechanisms of action [21,60]. For instance, the bumblebee linear peptides hymenoptaecin and abaecin exhibit a synergistic interaction against *E. coli*. Although abaecin alone shows no detectable activity at concentrations up to 200 μM, hymenoptaecin disrupts the bacterial membrane, allowing abaecin to enhance bactericidal effects at doses as low as 1.25 μM [61]. In *Tenebrio molitor*, RNAi-mediated knock-down of the defensin Tenecin-1 did not affect host survival but reduced bacterial load following *S. aureus* infection. However, simultaneous knock-down of Tenecin-1 with other AMPs, such as the coleoptericin Tenecin-2 or the attacin Tenecin-4, resulted in increased mortality and higher bacterial loads, highlighting functional complementarity among AMPs during infection [62]. In vitro experiments showed that Attacin, an antibacterial protein that accumulates in the hemolymph of the giant silk moth *Hyalophora cecropia*, compromises the permeability barrier function of the *E. coli* outer membrane. Pretreatment with Attacin significantly increases the sensitivity of *E. coli* to Cecropin B, another antimicrobial protein present in the hemolymph of *H. cecropia* [63]. In addition, recent studies have pointed to the role of host defense peptides in resilience mechanisms, including tolerance to host immune molecules [46] and pathogen toxins [26,31]. However, we did not detect a role for Paillotin in resilience to the fungal toxins Destruxin A or Verruculogen (Figure S8). Interestingly, in contrast to recent studies [26,31], we did not detect a susceptibility of Toll-deficient *spz^{rm7}* flies to these two toxins (Figure S8). Of note, the description of DIMs has been done largely by collecting peptides in Trifluoroacetic acid, which is known to modify peptides by, for example, deamidating NG motifs [24,64]. It is possible that the endogenous peptide products of the host immune response differ from the peptides observed in proteomic analysis and synthetic peptides used in recent studies, in unknown ways. Early studies on AMPs typically found striking activity *in vitro* using peptide extracts purified from endogenous hemolymph or cell line supernatant [14,17,19,21,65]. Future studies may benefit from a return to such approaches, including gel overlay experiments that can simplify the isolation of the activity of single peptides from endogenous samples [66] or collecting crude peptide extracts from overexpressing cells or larval hemolymph. However, future studies might also explore whether

Paillotin protects against virulence factors specific to *P. burhodogranariea*. For now, despite displaying many features of AMPs, we cannot conclusively classify Paillotin as a bona fide AMP. It should be classified as a host defense peptide until its molecular mode of action is fully determined.

The use of compound module-deficient mutants has revealed the major role of the Imd pathway, and to a lesser extent, melanization, but not the cellular response to survive *P. burhodogranariea* infection [67]. Since Paillotin does not affect Toll and Imd pathway activities, we cannot exclude a role of this peptide in the melanization reaction. However, this role would have to be subtle as *Paillotin* mutants display a wild-type blackening reaction upon clean injury (Figure S9*a*). Previous studies have shown that flies lacking 14 AMPs exhibit increased microbiota abundance and altered composition in old flies, suggesting that AMPs are involved not only in pathogen resistance but also in shaping the environmental microbiota [11]. However, we did not detect any significant differences in gut microbiota load between wild-type flies and *Paillotin* mutants (Figure S9*b*).

Interestingly, there is a relatively high frequency of loss-of-function mutations segregating in wild *Drosophila* populations (Table S1), suggesting Paillotin is not integral to the generic host defense. This observation is consistent with our results showing that the *Paillotin* mutation affects defense against only a few bacteria out of the pathogens tested, suggesting it is dispensable in defense against many microbes, but useful against some microbes like *Providencia sp.* whose presence is stochastic in wild-caught flies [68].

To conclude, our study characterizes a new *Drosophila* host defense peptide, Paillotin, which plays a specific role against *P. burhodogranariea*. IM18/Paillotin is the last gene encoding an immune molecule identified by MALDI-TOF in the 1998 study [38] to be characterized, following the successful characterization of Bomanins [30], Daisho [23], Buletin [25], and Baramicin [24,26]. Together, these studies and the present article provide important insights into the *Drosophila* humoral immune response, revealing a more complex picture than initially anticipated. Using the power of *Drosophila* genetics to dissect how immune effectors individually or collectively contribute to host defense provides a general principle on the architecture of innate immune systems.

Materials and Methods

Sequence Comparisons

Genomic sequences were obtained from GenBank default reference assemblies [69], from the DPGP via the PopFly website (www.popfly.uab.cat)[47], and DGRP sequence data from the DGRP database [48] (http://dgrp2.gnets.ncsu.edu/). Sequence comparisons and alignment figures were prepared using Geneious R10 [70]. Putative NF-κB binding sites were annotated using the *Relish* motif "GGRDNNHHBS" as described in Copley et al [45], along with a manually curated amalgam motif "GGGHHNNDVH" described in Hanson et al [24].

Paillotin annotation across species was performed as described in Hanson et al [42]. In brief, a recursive tBLASTn approach was used with an artificially high E-value (E < 1000) to compensate for the short length of the AMP query sequence. All the best BLAST hits were assessed manually. If no Paillotin was found in a genome or lineage, only the conserved C-terminus was used as a query in a repeat attempt (improving sensitivity). No *Paillotin* candidates were discovered outside *Brachycera*, which included screens of the sequenced nematoceran genomes of various mosquitoes, *Chironomus riparius, Clogmia albipunctata, Tipula unca, Lutzomyia longipalpis, Phlebotomus papatasi, Sylvicola cinctus,* and *Bibio marci*.

Fly Strains

Flies were maintained at 25°C on standard cornmeal-agar media. w^{1118} flies were used as wild-type controls. The positive controls for infection assays for Gram-positive and Gram-negative infections were spz^{rm7} and $Relish^{E20}$. The $Pai^{\triangle 41}$ and $Pai^{\triangle 370}$ mutations are loss-of-function variants in the *IM18* gene within the DGRP, and $Pai^{\triangle SFHI}$ was generated via CRISPR/Cas9. The guide RNAs that targeted *Pai* were cloned into the pUAST-attB plasmids before injecting into the *y [1] M {RFP [3xP3.PB] GFP[E.3xP3] =vas-int.Dm}ZH-2A w [*]; M{3xP3-RFP.attP} ZH-86Fb* (BL24749) to get gRNA transgenic flies. The gRNA flies were then crossed with flies containing nos-Cas9 to get mutants. The *UAS-Pai* constructs were prepared using the TOPO pENTR entry vector and cloned into the pTW system. Genetic variants were backcrossed into the DrosDel isogenic background over seven generations as previously described [71] using the following primers that are specific to an indel in the *Pai*

411   locus common to both the *Pai^{Δ41}* and *Pai^{Δ370}*, but which is absent in the DrosDel background:

412   Pai_41F   (ACGTACGGTCAAAAGCAACTACTTCG)   and   Pai_R3mut1

413   (CCAATCCCAAGGGCACTCGTATATG), amplified with a 66°C annealing temperature. All

414   the fly strains used in the paper are listed in Table S2.

MALDI-TOF Proteomic Analysis

Hemolymph samples were collected from *Ecc15*-infected flies (OD600 = 200, 24 hours

post-infection, 50 flies/sample) using a nanoinjector (Nanoject, Drummond Scientific,

Broomall, PA) for MALDI-TOF analysis. Trifluoroacetic acid (0.1% TFA) was added to the

hemolymph before mixing with the acetonitrile matrix. Samples are processed by SPE (Zip-Tip

C18) before MALDI. Peaks were identified based on corresponding m/z values from previous

studies [24,72] in ISIC's Mass Spectrometry facility of EPFL. Spectra were visualized using

Origin software, with figures prepared in Prism 10.

Gene Expression Analysis

Total RNA was extracted from pooled samples of ten flies using TRIzol. cDNA was

synthesized using Takara Reverse Transcriptase. Quantitative PCR (qPCR) was performed

using PowerUP SYBR Green Master Mix with the primers. Results represent the average of

three independent experiments. Primers used in this study were:

Drosomycin-F 5'- CGTGAGAACCTTTTCCAATATGAT-3'

Drosomycin-R 5'- TCCCAGGACCACCAGCAT-3'

Diptericin A-F: 5'-GCTGCGCAATCGCTTCTACT-3'

Diptericin A-R:5'-TGGTGGAGTGGGCTTCATG-3'

Rpl32-F: 5'-GACGCTTCAAGGGACAGTATCTG-3'

Rpl32-R: 5'-AAACGCGGTTCTGCATGAG-3'

Pai-F: 5'-GATCCGTTTCCTGGTCCAGTGTG-3'

Pai-R: 5'-CCATGAAGCTGATCGCATTGTGC-3'

CG10332-F: 5'-TCAGAGGCTGCACCATCAGAAGT-3'

CG10332-R: 5'-CCGCTTCAGAGGCTGCACC-3'

Microbial Culturing Conditions

Micrococcus luteus, *Erwinia carotovora carotovora (Ecc15)*, and *Providencia*

burhodogranariea strain B were cultured overnight in LB broth at 29°C. *Enterobacter cloacae*,

Escherichia coli strain 1106, *Providencia rustigianii*, *Providencia vermicola*, *Providencia*

rettgeri, and *Staphylococcus aureus* were cultured overnight in LB at 37°C. *Enterococcus*

faecalis, *Bacillus subtilis,* and *Listeria monocytogenes* were cultured overnight in BHI at 37°C.

Metarhizium anisopliae and *Metarhizium robertisii* were cultured on malt agar plates at 25°C

for around two weeks until sporulation was observed. *Beauveria bassiana* strain R444

commercial spores were produced by Andermatt Biocontrol, products: BB-PROTEC. Bacteria

were pelleted at 4000 rpm for 15 minutes at 4°C, resuspended in PBS, and diluted to the desired

OD600 value indicated in the figure legends. The fungal material was filtered through glass

wool using 0.05% Tween-80 to collect spores and diluted to the desired concentration (in

spores/ml). All microbial strains used in this study are listed in Table S3.

Microbiota Load Analysis

Flies were collected 30 hours post-infection and surface-sterilized with 70% ethanol.

Whole-fly homogenization was performed in LB broth using a bead beater at 7200 rpm for 30

seconds. Serial dilutions of the homogenates were plated (100 μL per plate) on LB agar and

incubated at 29 °C for 24 hours. Colony-forming units (CFUs) were then manually counted.

For gut microbiota load analysis, 18-day-old conventionally reared flies were collected

and processed using the same procedure, except that MRS medium was used for culturing.

Plates were incubated at 29 °C for 48 hours prior to CFU quantification.

***In Vitro* Antibacterial Activity**

The Paillotin peptide (45 residues) was synthesized by GenicBio with >95% purity, and

Cecropin A (from silk moth) was obtained from Sigma-Aldrich (≥97% purity). Peptides were

dissolved in water. Bacteria were cultured overnight and diluted to OD600 = 0.005 in LB broth,

then incubated for 2 hours. Paillotin peptide and Cecropin A were diluted in LB broth to the desired concentration. Bacterial cultures (1 µL) were added to peptide solutions in a 96-well plate, and OD600 was recorded every 10 minutes using a TECAN plate reader.

Survival Assays

Male flies (3-5 days old) were used for survival assays. The thorax of each fly was pricked with a needle (diameter ~0.2mm) dipped in a bacterial suspension. For the fungi infection, flies were rolled in the plates of *A. fumigatus* or the commercial spores of *B. bassiana,* or were dipped into a spore solution of *M. anisopliae* or *M. robertisii*. Mortality was recorded daily. Infected flies were maintained at 29°C or 25°C and transferred to fresh vials three times per week.

Toxin Injection

Destruxin A (Sigma) was dissolved in high-purity dimethyl sulfoxide (DMSO) and subsequently diluted with PBS to a final concentration of 8 mM. A volume of 9.2 nL of this solution was administered to flies using the Nanoject III microinjector (Drummond) [26]. Verruculogen (Abcam) was prepared in DMSO at a concentration of 1 mg/mL, and 4.6 nL of this solution was injected into the flies [31].

Wounding experiment

Clean injury was performed with a needle sterilized (diameter ~0.1mm) with an EtOH and PBS wash. Cuticle blackening was categorized into normal, weak, or no melanization 24h post-pricking at the injury site.

MAH.   Funding acquisition: BL, MAH. Investigation: YT, XJY, RJJ, FS. Methodology: YT, XJY, RJJ, MAH. Project administration: BL. Validation: YT, MAH. Resources: YT, XJY, MAH. Supervision: MAH, BL. Writing – original draft: YT, BL. Writing – review & editing: MAH, BL.

Conflict of interest declaration. We declare we have no competing interests.

Funding. This project was supported by the SNSF grants 310030_215073 and CRSII5_186397 awarded to B.L. and Wellcome Trust grant 227559/Z/23/Z awarded to M.H.

Acknowledgements. We thank Jean-Philippe Boquette, Guiqing Liu, Samuel Rommelaere, Fanny Schupfer, Faustine Ryckebusch, and Yijie Li for experimental help and Hannah Westlake for editing. We thank Ana Marija Jakšić for providing the initial DGRP flies used to source the natural *Paillotin* mutations. We thank the Guangzhou Drosophila Resource Center (GDRC) for generating the CRISPR–Cas9 *Paillotin* knockout line. We thank Laure Menin from the EPFL ISIC Mass Spectrometry Facility for her technical assistance.

Reference

1. Westlake H, Hanson MA, Lemaitre B. 2024 *The Drosophila immunity handbook*. First. EPFL Press. (doi:10.55430/6304TDIHVA01)

2. Ferrandon D, Imler J-L, Hetru C, Hoffmann JA. 2007 The Drosophila systemic immune response: sensing and signalling during bacterial and fungal infections. *Nat Rev Immunol* **7**, 862–874. (doi:10.1038/nri2194)

3. Buchon N, Silverman N, Cherry S. 2014 Immunity in Drosophila melanogaster — from microbial recognition to whole-organism physiology. *Nat Rev Immunol* **14**, 796–810. (doi:10.1038/nri3763)

4. Pradeu T, Thomma BPHJ, Girardin SE, Lemaitre B. 2024 The conceptual foundations of innate immunity: Taking stock 30 years later. *Immunity* **57**, 613–631. (doi:10.1016/j.immuni.2024.03.007)

5. De Roode JC, Lefèvre T. 2012 Behavioral Immunity in Insects. *Insects* **3**, 789–820. (doi:10.3390/insects3030789)

6. Montanari M, Royet J. 2021 Impact of Microorganisms and Parasites on Neuronally Controlled Drosophila Behaviours. *Cells* **10**, 2350. (doi:10.3390/cells10092350)

7. Brown KL, Hancock RE. 2006 Cationic host defense (antimicrobial) peptides.
Current Opinion in Immunology **18**, 24–30. (doi:10.1016/j.coi.2005.11.004)

8. Hanson MA, Lemaitre B. 2020 New insights on Drosophila antimicrobial peptide
function in host defense and beyond. *Current Opinion in Immunology* **62**, 22–30.
(doi:10.1016/j.coi.2019.11.008)

9. Parvy J-P, Yu Y, Dostalova A, Kondo S, Kurjan A, Bulet P, Lemaître B, Vidal M,
Cordero JB. 2019 The antimicrobial peptide defensin cooperates with tumour
necrosis factor to drive tumour cell death in Drosophila. *eLife* **8**, e45061.
(doi:10.7554/eLife.45061)

10. Hirata M, Nomura T, Inoue YH. 2025 Anti-Tumor Effects of Cecropin A and
Drosocin Incorporated into Macrophage-like Cells Against Hematopoietic Tumors
in Drosophila mxc Mutants. *Cells* **14**, 389. (doi:10.3390/cells14060389)

11. Marra A, Hanson MA, Kondo S, Erkosar B, Lemaitre B. 2021 *Drosophila*
Antimicrobial Peptides and Lysozymes Regulate Gut Microbiota Composition and
Abundance. *mBio* **12**, e00824-21. (doi:10.1128/mBio.00824-21)

12. Kounatidis I, Chtarbanova S, Cao Y, Hayne M, Jayanth D, Ganetzky B,
Ligoxygakis P. 2017 NF-κB Immunity in the Brain Determines Fly Lifespan in
Healthy Aging and Age-Related Neurodegeneration. *Cell Reports* **19**, 836–848.
(doi:10.1016/j.celrep.2017.04.007)

13. E L *et al.* 2018 An Antimicrobial Peptide and Its Neuronal Receptor Regulate
Dendrite Degeneration in Aging and Infection. *Neuron* **97**, 125-138.e5.
(doi:10.1016/j.neuron.2017.12.001)

14. Hultmark D, Steiner H, Rasmuson T, Boman HG. 1980 Insect Immunity.
Purification and Properties of Three Inducible Bactericidal Proteins from
Hemolymph of Immunized Pupae of *Hyalophora cecropia*. *European Journal of*
Biochemistry **106**, 7–16. (doi:10.1111/j.1432-1033.1980.tb05991.x)

15. Steiner H, Hultmark D, Engström Å, Bennich H, Boman HG. 1981 Sequence and
specificity of two antibacterial proteins involved in insect immunity. *Nature* **292**,
246–248. (doi:10.1038/292246a0)

16. Pardi A, Hare DR, Selsted ME, Morrison RD, Bassolino DA, Bach AC. 1988
Solution structures of the rabbit neutrophil defensin NP-5. *Journal of Molecular*
Biology **201**, 625–636. (doi:10.1016/0022-2836(88)90643-2)

17. Fehlbaum P, Bulet P, Michaut L, Lagueux M, Broekaert WF, Hetru C, Hoffmann
JA. 1994 Insect immunity. Septic injury of Drosophila induces the synthesis of a
potent antifungal peptide with sequence homology to plant antifungal peptides.
Journal of Biological Chemistry **269**, 33159–33163. (doi:10.1016/S0021-
9258(20)30111-3)

18. Levashina EA, Ohresser S, Bulet P, Reichhart J, Hetru C, Hoffmann JA. 1995 Metchnikowin, a Novel Immune-Inducible Proline-Rich Peptide from *Drosophila* with Antibacterial and Antifungal Properties. *European Journal of Biochemistry* **233**, 694–700. (doi:10.1111/j.1432-1033.1995.694_2.x)

19. Bulet P, Urge L, Ohresser S, Hetru C, Otvos L. 1996 Enlarged Scale Chemical Synthesis and Range of Activity of Drosocin, an O-Glycosylated Antibacterial Peptide of *Drosophila*. *European Journal of Biochemistry* **238**, 64–69. (doi:10.1111/j.1432-1033.1996.0064q.x)

20. Ekengren S. 1999 Drosophila cecropin as an antifungal agent. *Insect Biochemistry and Molecular Biology* **29**, 965–972. (doi:10.1016/S0965-1748(99)00071-5)

21. Rabel D, Charlet M, Ehret-Sabatier L, Cavicchioli L, Cudic M, Otvos L, Bulet P. 2004 Primary Structure and in Vitro Antibacterial Properties of the Drosophila melanogaster Attacin C Pro-domain. *Journal of Biological Chemistry* **279**, 14853–14859. (doi:10.1074/jbc.M313608200)

22. Clemmons AW, Lindsay SA, Wasserman SA. 2015 An Effector Peptide Family Required for Drosophila Toll-Mediated Immunity. *PLoS Pathog* **11**, e1004876. (doi:10.1371/journal.ppat.1004876)

23. Cohen LB, Lindsay SA, Xu Y, Lin SJH, Wasserman SA. 2020 The Daisho Peptides Mediate Drosophila Defense Against a Subset of Filamentous Fungi. *Front. Immunol.* **11**, 9. (doi:10.3389/fimmu.2020.00009)

24. Hanson MA, Cohen LB, Marra A, Iatsenko I, Wasserman SA, Lemaitre B. 2021 The Drosophila Baramicin polypeptide gene protects against fungal infection. *PLoS Pathog* **17**, e1009846. (doi:10.1371/journal.ppat.1009846)

25. Hanson MA, Kondo S, Lemaitre B. 2022 Drosophila immunity: the Drosocin gene encodes two host defence peptides with pathogen-specific roles.*Proc Biol Sci*. **289**, 20220773.(doi.org/10.1098/rspb.2022.0773)

26. Huang J *et al.* 2023 A Toll pathway effector protects *Drosophila* specifically from distinct toxins secreted by a fungus or a bacterium. *Proc. Natl. Acad. Sci. U.S.A.* **120**, e2205140120. (doi:10.1073/pnas.2205140120)

27. Hanson MA, Dostálová A, Ceroni C, Poidevin M, Kondo S, Lemaitre B. 2019 Synergy and remarkable specificity of antimicrobial peptides in vivo using a systematic knockout approach. *eLife* **8**, e44341. (doi:10.7554/eLife.44341)

28. Carboni AL, Hanson MA, Lindsay SA, Wasserman SA, Lemaitre B. 2022 Cecropins contribute to *Drosophila* host defense against a subset of fungal and Gram-negative bacterial infection. *Genetics* **220**, iyab188. (doi:10.1093/genetics/iyab188)

29. Hanson MA, Grollmus L, Lemaitre B. 2023 Ecology-relevant bacteria drive the evolution of host antimicrobial peptides in *Drosophila*. *Science* **381**, eadg5725. (doi:10.1126/science.adg5725)

30. Lindsay SA, Lin SJH, Wasserman SA. 2018 Short-Form Bomanins Mediate Humoral Immunity in ***Drosophila***. *J Innate Immun* **10**, 306–314. (doi:10.1159/000489831)

31. Xu R *et al.* 2023 The Toll pathway mediates *Drosophila* resilience to *Aspergillus* mycotoxins through specific Bomanins. *EMBO Reports* **24**, e56036. (doi:10.15252/embr.202256036)

32. Unckless RL, Howick VM, Lazzaro BP. 2016 Convergent Balancing Selection on an Antimicrobial Peptide in Drosophila. *Current Biology* **26**, 257–262. (doi:https://doi.org/10.1016/j.cub.2015.11.063)

33. Vanha-aho L-M, Kleino A, Kaustio M, Ulvila J, Wilke B, Hultmark D, Valanne S, Rämet M. 2012 Functional Characterization of the Infection-Inducible Peptide Edin in Drosophila melanogaster. *PLoS ONE* **7**, e37153. (doi:10.1371/journal.pone.0037153)

34. Lu M, Wei D, Shang J, Li S, Song S, Luo Y, Tang G, Wang C. 2024 Suppression of Drosophila antifungal immunity by a parasite effector via blocking GNBP3 and GNBP-like 3, the dual receptors for β-glucans. *Cell Reports* **43**, 113642. (doi:10.1016/j.celrep.2023.113642)

35. K. Maasdorp M, Valanne S, Vesala L, Vornanen P, Haukkavaara E, Tuomela T. 2025 IbinA and IbinB regulate the Toll pathway-mediated immune response in Drosophila melanogaster. *bioRxiv*. (doi.org/10.1101/2025.04.10.648149)

36. Goto A, Yano T, Terashima J, Iwashita S, Oshima Y, Kurata S. 2010 Cooperative Regulation of the Induction of the Novel Antibacterial Listericin by Peptidoglycan Recognition Protein LE and the JAK-STAT Pathway. *Journal of Biological Chemistry* **285**, 15731–15738. (doi:10.1074/jbc.M109.082115)

37. Westlake H, David F, Tian Y, Krakovic K, Dolgikh A, Juravlev L, Esmangart de Bournonville T, Carboni A, Melcarne C, Shan T, Wang Y, Mu Y, Kotwal A, Pirko N, Boquete JP, Schüpfer F, Rommelaere S, Poidevin M, Liu Z, Kondo S, Ratnaparkhi GS, Chakrabarti S, Liu G, Masson F, Li X, Hanson MA, Jiang H, Di Cara F, Kurant E, Lemaitre B. 2025 Reproducibility of Scientific Claims in Drosophila Immunity: A Retrospective Analysis of 400 Publications. *bioRxiv*. (doi.org/10.1101/2025.07.07.663442)

38. Uttenweiler-Joseph S, Moniatte M, Lagueux M, Van Dorsselaer A, Hoffmann JA, Bulet P. 1998 Differential display of peptides induced during the immune response of *Drosophila* : A matrix-assisted laser desorption ionization time-of-flight mass

spectrometry study. *Proc. Natl. Acad. Sci. U.S.A.* **95**, 11342–11347. (doi:10.1073/pnas.95.19.11342)

39. Jehle JA. 2009 André Paillot (1885–1944): His work lives on. *Journal of Invertebrate Pathology* **101**, 162–168. (doi:10.1016/j.jip.2009.03.009)

40. Carton Y. 2019 *Innate Immunity: From Louis Pasteur to Jules Hoffmann*. San Diego: ISTE Press Limited - Elsevier Incorporated.

41. Robinson SW, Herzyk P, Dow JAT, Leader DP. 2013 FlyAtlas: database of gene expression in the tissues of Drosophila melanogaster. *Nucleic Acids Research* **41**, D744–D750. (doi:10.1093/nar/gks1141)

42. Hanson MA, Lemaitre B, Unckless RL. 2019 Dynamic Evolution of Antimicrobial Peptides Underscores Trade-Offs Between Immunity and Ecological Fitness. *Front. Immunol.* **10**, 2620. (doi:10.3389/fimmu.2019.02620)

41. Sanjay Kumar, Sanjay Kumar, Arvind Kumar Singh. 2017 Population genetics of Drosophila: Genetic Variation and Differentiation among Indian Natural Populations of Drosophila ananassae. *Zoological Studies* **56**: e. (doi:10.6620/ZS.2017.56-01)

44. Waumans Y, Baerts L, Kehoe K, Lambeir A-M, De Meester I. 2015 The Dipeptidyl Peptidase Family, Prolyl Oligopeptidase, and Prolyl Carboxypeptidase in the Immune System and Inflammatory Disease, Including Atherosclerosis. *Front. Immunol.* **6**. (doi:10.3389/fimmu.2015.00387)

45. Copley RR, Totrov M, Linnell J, Field S, Ragoussis J, Udalova IA. 2007 Functional conservation of Rel binding sites in drosophilid genomes. *Genome Res.* **17**, 1327–1335. (doi:10.1101/gr.6490707)

46. Rommelaere S, Schüpfer F, Armand F, Hamelin R, Lemaitre B. 2025 An updated proteomic analysis of *Drosophila* haemolymph after bacterial infection. *Fly* **19**, 2485685. (doi:10.1080/19336934.2025.2485685)

47. Lack JB, Lange JD, Tang AD, Corbett-Detig RB, Pool JE. 2016 A Thousand Fly Genomes: An Expanded*Drosophila*Genome Nexus. *Mol Biol Evol* **33**, 3308–3313. (doi:10.1093/molbev/msw195)

48. Mackay TFC *et al.* 2012 The Drosophila melanogaster Genetic Reference Panel. *Nature* **482**, 173–178. (doi:10.1038/nature10811)

49. Scott MG, Davidson DJ, Gold MR, Bowdish D, Hancock REW. 2002 The Human Antimicrobial Peptide LL-37 Is a Multifunctional Modulator of Innate Immune Responses. *The Journal of Immunology* **169**, 3883–3891. (doi:10.4049/jimmunol.169.7.3883)

50. Dong X *et al.* 2022 Keratinocyte-derived defensins activate neutrophil-specific receptors Mrgpra2a/b to prevent skin dysbiosis and bacterial infection. *Immunity* **55**, 1645-1662.e7. (doi:10.1016/j.immuni.2022.06.021)

51. Samakovlis C, Kimbrell DA, Kylsten P, Engström A, Hultmark D. 1990 The immune response in Drosophila: pattern of cecropin expression and biological activity. *The EMBO Journal* **9**, 2969–2976. (doi:10.1002/j.1460-2075.1990.tb07489.x)

52. De Gregorio E. 2002 The Toll and Imd pathways are the major regulators of the immune response in Drosophila. *The EMBO Journal* **21**, 2568–2579. (doi:10.1093/emboj/21.11.2568)

53. Tzou P, Reichhart J-M, Lemaitre B. 2002 Constitutive expression of a single antimicrobial peptide can restore wild-type resistance to infection in immunodeficient *Drosophila* mutants. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 2152–2157. (doi:10.1073/pnas.042411999)

54. Hanson MA. 2024 When the microbiome shapes the host: immune evolution implications for infectious disease. *Phil. Trans. R. Soc. B* **379**, 20230061. (doi:10.1098/rstb.2023.0061)

55. Maiti S, Patro S, Purohit S, Jain S, Senapati S, Dey N. 2014 Effective Control of Salmonella Infections by Employing Combinations of Recombinant Antimicrobial Human β-Defensins hBD-1 and hBD-2. *Antimicrob Agents Chemother* **58**, 6896–6903. (doi:10.1128/aac.03628-14)

56. Myhrman E, Håkansson J, Lindgren K, Björn C, Sjöstrand V, Mahlapuu M. 2013 The novel antimicrobial peptide PXL150 in the local treatment of skin and soft tissue infections. *Appl Microbiol Biotechnol* **97**, 3085–3096. (doi:10.1007/s00253-012-4439-8)

57. Rivas-Santiago B, Rivas Santiago CE, Castañeda-Delgado JE, León–Contreras JC, Hancock REW, Hernandez-Pando R. 2013 Activity of LL-37, CRAMP and antimicrobial peptide-derived compounds E2, E6 and CP26 against Mycobacterium tuberculosis. *International Journal of Antimicrobial Agents* **41**, 143–148. (doi:10.1016/j.ijantimicag.2012.09.015)

58. Björn C, Håkansson J, Myhrman E, Sjöstrand V, Haug T, Lindgren K, Blencke H-M, Stensvåg K, Mahlapuu M. 2012 Anti-infectious and anti-inflammatory effects of peptide fragments sequentially derived from the antimicrobial peptide centrocin 1 isolated from the green sea urchin, Strongylocentrotus droebachiensis. *AMB Expr* **2**. (doi:10.1186/2191-0855-2-67)

59. Zdybicka-Barabas A, Mak P, Klys A, Skrzypiec K, Mendyk E, Fiołka MJ, Cytryńska M. 2012 Synergistic action of Galleria mellonella anionic peptide 2 and

lysozyme against Gram-negative bacteria. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1818**, 2623–2635. (doi:10.1016/j.bbamem.2012.06.008)

60. Rahnamaeian M, Cytryńska M, Zdybicka-Barabas A, Vilcinskas A. 2016 The functional interaction between abaecin and pore-forming peptides indicates a general mechanism of antibacterial potentiation. *Peptides* **78**, 17–23. (doi:10.1016/j.peptides.2016.01.016)

61. Rahnamaeian M *et al.* 2015 Insect antimicrobial peptides show potentiating functional interactions against Gram-negative bacteria. *Proc. R. Soc. B.* **282**, 20150293. (doi:10.1098/rspb.2015.0293)

62. Zanchi C, Johnston PR, Rolff J. 2017 Evolution of defence cocktails: Antimicrobial peptide combinations reduce mortality and persistent infection. *Molecular Ecology* **26**, 5334–5343. (doi:10.1111/mec.14267)

63. Engström P, Carlsson A, Engström A, Tao ZJ, Bennich H. 1984 The antibacterial effect of attacins from the silk moth Hyalophora cecropia is directed against the outer membrane of Escherichia coli. *EMBO J* **3**, 3347–3351. (doi:10.1002/j.1460-2075.1984.tb02302.x)

64. Tyler-Cross R, Schirch V. 1991 Effects of amino acid sequence, buffers, and ionic strength on the rate and mechanism of deamidation of asparagine residues in small peptides. *Journal of Biological Chemistry* **266**, 22549–22556. (doi:10.1016/s0021-9258(18)54607-x)

65. Alkotaini B, Anuar N, Kadhum AAH, Sani AAA. 2013 Detection of secreted antimicrobial peptides isolated from cell-free culture supernatant of *Paenibacillus alvei* AN5. *Journal of Industrial Microbiology and Biotechnology* **40**, 571–579. (doi:10.1007/s10295-013-1259-5)

66. Samakovlis C, Kylsten P, Kimbrell DA, Engström A, Hultmark D. 1991 The andropin gene and its product, a male-specific antibacterial peptide in Drosophila melanogaster. *The EMBO Journal* **10**, 163–169. (doi:10.1002/j.1460-2075.1991.tb07932.x)

63. Ryckebusch F, Tian Y, Rapin M, Schupfer F, Hanson M, Lemeitre B. 2025 Layers of immunity: Deconstructing the Drosophila effector response. *bioRxiv*. (doi: 10.1101/2025.03.28.645959)

68. Chen J-S, Tsaur S-C, Ting C-T, Fang S. 2022 Dietary Utilization Drives the Differentiation of Gut Bacterial Communities between Specialist and Generalist Drosophilid Flies. *Microbiol Spectr* **10**, e01418-22. (doi:10.1128/spectrum.01418-22)

69. Kim BY *et al.* 2021 Highly contiguous assemblies of 101 drosophilid genomes. *eLife* **10**, e66405. (doi:10.7554/eLife.66405)

70. Kearse M *et al.* 2012 Geneious Basic: An integrated and extendable desktop
software platform for the organization and analysis of sequence data.
Bioinformatics **28**, 1647–1649. (doi:10.1093/bioinformatics/bts199)

71. Ferreira ÁG, Naylor H, Esteves SS, Pais IS, Martins NE, Teixeira L. 2014 The Toll-
Dorsal Pathway Is Required for Resistance to Viral Oral Infection in Drosophila.
PLoS Pathog **10**, e1004507. (doi:10.1371/journal.ppat.1004507)

72. Levy F, Rabel D, Charlet M, Bulet P, Hoffmann JA, Ehret-Sabatier L. 2004
Peptidomic and proteomic analyses of the systemic immune response of Drosophila.
Biochimie **86**, 607–616. (doi:10.1016/j.biochi.2004.07.007)

73. Tian Y, Yue XJ, Jiao RJ, Hanson M, Lemaitre B. 2025 Data from: Functional
characterization of Paillotin: an immune peptide regulated by the Imd pathway with
pathogen-specific roles in Drosophila immunity. Dryad Digital
Repository. (doi.org/10.5061/dryad.f4qrfj77k)

Figure 1: The *Paillotin* gene structure and conservation. (*a*) Schematic representation of the *Paillotin* (*Pai*) locus. The *Pai-RB* transcript is annotated as polycistronic, sharing a promoter and transcript with *CG10332* (*CG10332-RA*), indicating both gene CDS regions are co-transcribed. Promoter regions (250 bp upstream) for *Pai-RB* and *CG10332-RA* are shown in cyan. Colored triangles indicate putative NF-κB binding motifs: *Relish*-responsive motifs (GGRDNNHHBS) in teal, *Dif/Dorsal*-responsive motifs (GGGHNNNDVH) in purple. Red triangles represent sites containing overlapping matches to both motifs. (*b*) Sequence alignment of Paillotin orthologues across diverse *Brachyceran* flies revealed a conserved protein structure, including a signal peptide, a dipeptidyl peptidase cleavage site (XA/XP), and the mature Paillotin peptide. Highlighting corresponds to residues with >50% identity across the alignment.

Figure 2: *Paillotin* **is regulated by the Imd pathway**. Transcript levels of *Paillotin*/*CG10332* in w^{1118}, *Relish*^{E20}, *spz*^{rm7}, and *Relish*^{E20}, *spz*^{rm7} flies collected at 8 h, 16h, and 24h post-*E. coli* infection (*a*, *c*) and *M. luteus* infection (*b*, *d*). (*e*) Quantified Paillotin peptide levels of w^{1118}, *Relish*^{E20}, and *spz*^{rm7} hemolymph upon either *Ecc15* or *M. luteus* infection (data extracted from (Rommelaere *et al.*, 2025)). Gene Expression was measured by qRT-PCR and normalized with w^{1118} UC (unchallenged) flies set as a value of 1. The data shown in the figure are based on three independent experiments, with at least 30 flies per genotype at each time point in each experiment. Error bars represent SEM. Statistical significance was determined using two-way ANOVA (*$p<0.05$, ** $p<0.01$, *** $p<0.001$, n.s. = not significant, $p>0.05$).

Figure 3: Description and validation of *Pai△41*, *Pai△370*, and *Pai△SFHI* mutations. (*a*) Multiple sequence alignment between the Dmel_R6 reference genome Paillotin and Paillotin peptide variants. Conserved residues are highlighted by color coding according to sequence similarity. (*b*) MALDI-TOF proteomic analysis of hemolymph from *w1118*, *Pai△41*, *Pai△370*, and *Pai△SFHI* male flies collected 24 hours after septic injury with *Ecc15*.

Figure 4: Paillotin does not impact the Imd or Toll pathways. The expression of *Diptericin A* (*a*) at 6 or 12h post-infection with *Ecc15* and *Drosomycin* (*b*) at 8h or 24h post-infection with *M. luteus* were monitored in w^{1118}, $Pai^{\triangle41}$, and $Pai^{\triangle370}$ flies. $Relish^{E20}$ and spz^{rm7} were used as the positive control. Expression was normalized to w^{1118} UC (unchallenged) flies set as a value of 1. The data shown in the figure are based on three independent experiments, with at least 30 flies per genotype at each time point in each experiment. Error bars represent SEM. Statistical analysis was performed using two-way ANOVA. For *Diptericin A*, significant effects were detected for genotype ($F(3,24) = 148.1$, $p < 0.0001$), time ($F(2,24) = 332.8$, $p < 0.0001$), and their interaction ($F(6,24) = 38.98$, $p < 0.0001$). For *Drosomycin*, significant effects were also observed for genotype ($F(3,24) = 72.82$, $p < 0.0001$), time ($F(2,24) = 191.5$, $p < 0.0001$), and interaction ($F(6,24) = 22.84$, $p < 0.0001$). However, Tukey's post hoc comparisons revealed no significant difference between w^{1118} and *Paillotin* mutants at any time point (n.s. = not significant).

Figure 5: *Paillotin* mutants are susceptible to *P. burhodogranariea* infection. (*a*) The three *Paillotin* mutants exhibit increased susceptibility following septic injury with *P. burhodogranariea* at the indicated bacterial inoculum. (*b*) Increased bacterial load in *Paillotin* mutants at 30 hpi. Bacterial load was $\log_{10}$-transformed prior to statistical analysis. One-way ANOVA revealed a significant genotype effect ($F(3,30) = 18.95$, $p < 0.0001$). Post hoc Dunnett's test confirmed that all three mutant lines carried significantly higher bacterial loads than w^{1118} ($p < 0.0001$ for all comparisons). Data points represent the mean bacterial load per fly, derived from individual experiments using pools of four flies per genotype following *P. burhodogranariea* infection. No consistent effect of the *Paillotin* mutation on resistance to the Gram-positive bacterium *Enterococcus faecalis* (*c*) or the fungus *Beauveria bassiana* (*d*). Survival data were analyzed using the Kaplan–Meier method. Statistical significance is indicated relative to the w^{1118} (*** $p<0.001$, n.s. = not significant, $p>0.05$). N=total number of flies in experiments.

Figure 6: Paillotin synergistically contributes to defense against *P. burhodogranariea* with other Imd-regulated AMPs. (*a-c*) Survival curves showing the synergistic effects between different *Paillotin* mutants (*Pai*$^{\Delta 41}$, *Pai*$^{\Delta 370}$, *Pai*$^{\Delta SFHI}$) and *Group B* mutations (*Drosocin, Attacin A, B, C,* and *Diptericin A, B*) in the heterozygous background (*Group B/+*). Survival curves were analyzed using the Cox proportional hazards (CoxPH) model in R version 3.6.3. (** *p*<0.01, *** *p*<0.001, n.s. = not significant, *p*>0.05). N=total number of flies in experiments.

Figure 7: Over-expression of *Paillotin* rescues the resistance of *Paillotin*-deficient flies against *P. burhodogranariea*. (*a-b*) Overexpression of *Paillotin* via a combination of the *C564-Gal4* or *Actin-Gal4* and *UAS-Pai* constructs enhances the resistance of the Paillotin mutant to *P. burhodogranariea* infection. (*c*) Overexpression of *Paillotin* in the *Relish^E20* background failed to restore resistance. Survival curves were analyzed using the Kaplan-Meier method. (*** $p<0.001$, n.s. = not significant, $p>0.05$). N=total number of flies in experiments.

Supplementary material for

Functional Characterization of Paillotin: An Immune Peptide Regulated by the Imd Pathway with Pathogen-Specific Roles in *Drosophila* Immunity

Yao Tian[1], Xiaojing Yue[3], Renjie Jiao[4], Mark A. Hanson[1,2]*, Bruno Lemaitre[1]*

1: Global Health Institute, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

2: Centre for Ecology and Conservation, University of Exeter, Penryn, United Kingdom

3: Laboratory of Biology and Modelling of the Cell, Ecole Normale Superieure of Lyon, CNRS, Lyon, France.

4: Sino-French Hoffmann Institute, Guangzhou Medical University, Guangzhou, China

*: Equal last author: M.A. Hanson (M.Hanson@exeter.ac.uk), B.L. Lemaitre (bruno.lemaitre@epfl.ch)

Table S1: Collectively, 39 of 789 genome-sequenced fly strains, including 18 of the 205 *Drosophila* Genetic Reference Panel lines encode a loss-of-function mutation in *Paillotin*. Lines encoding *Pai* putative loss-of-function mutations are annotated, alongside the genetic description of each allele. *Pai^A41* "uncertain" and "alternate" alleles per the DGRP variant call dataset are both represented by a string of ~61 uncertain nucleotide calls (Ns) matching the pattern of allele 2R_19488505_DEL in the DPGP3 dataset (downloaded from PopFly). This suggests DGRP strains with "uncertain" allele annotations at base Dmel.R5_2R_19488505 also bear the 2R_19488505_DEL indel. Bolstered by this finding, we scanned the *Drosophila* genome nexus for strings of uncertain variant calls (Ns) that affect at least 3 contiguous codons, treating these as putative indels. This may yield an undercount by ignoring smaller indels, or an overcount by treating regions bearing multiple SNPs in linkage as indels. Nevertheless, this annotation effort draws attention to strains bearing potential loss-of-function mutations in *Paillotin* that may be relevant to future studies. DPGP indel strain summaries are grouped by the location of the first disrupted codon (within the signal or mature peptide).

Mutation	Allele	Reference allele	Alternate allele	Effect	Present in
Pai^{Δ41}	*2R_19488505 _DEL*	(C)CAAT CGGAG CGTCG TAGTA CAGTG GTCGT TGCTG TGAGC TTGGA GACCT GAACG GATTT CC	(C)-	61bp deletion	**DGRP:** 28, 31, 41, 176, 227, 321, 362, 390, 406, 409, 491, 707, 843, 852, 879, 892, 911 **DPGP:** CHB381
Pai^{Δ370}	*2R_19488690 _SNP*	C	T	loss of start codon	**DGRP:** 370 **DPGP:** USW_59 possibly heterozygous
Pai^{G37*}	*Pai*^{G37*}	GGA codon	TGA codon	codon 37 premature stop	**DPGP:** KR7, KR4N
Pai^{ΔM1}	*Pai*^{ΔM1}	ATG codon	ANG codon	putative start codon disruption	**DPGP:** USI06
Signal peptide + DPase indel	*multiple*	ATGAA GCTGA TCGCA TTGTG CTGCC TGCTC CTTTT GGGCC TCCTG GGCTT CCTAG CTGCT CCCGG CGTCG CCTCG CCA	multiple	indel beginning within signal peptide	**DGRP:** 385, 887 **DPGP:** NTH_07, NTH_15, USW_26, USW_37, USW_62, USW_68, USW_49, USW_76, AUS_04, EA_70N, MW_6-2
Mature peptide indel not including Pai^{Δ41}	*multiple*	TCTCG CCACA CTGGA CCAGG AAACG GATCG GGATC TGGAG CTGGG TCCGG AAATC CGTTC AGGTC TCCAA GCTCA CAGCA ACGAC CACTG TACTA CGACG CTCCG ATTGG GAAAC CATCG AAGAC TATGT ACGCC TGA	multiple	indel beginning within mature peptide not including *Pai*^{Δ41}	**DPGP:** USW_74, USW_87, FR_2N

Figure S1. Nucleotide alignment of wild-type and mutant *Paillotin* alleles. Nucleotide sequences of wild-type *Paillotin* (*Pai^Dmel R6*) and the three mutant alleles (*Pai^△41*, *Pai^△370*, and *Pai^△SFHI*) used in this study are aligned.

Figure S2: Expression of *Paillotin* in $Pai^{\Delta 41}$, $Pai^{\Delta 370}$, and $Pai^{\Delta SFHI}$ mutations. Expression was normalized to w^{1118} UC (unchallenged) flies set as a value of 1. The data shown in the figure are based on three independent experiments, with at least 30 flies per genotype at each time point in each experiment. Error bars represent SEM. Statistical analysis was performed using two-way ANOVA (** $p<0.01$, *** $p<0.001$).

Figure S3: No susceptibility of *Paillotin* mutants to clean injury. The thorax of wild-type and mutants was pricked with a clean needle. A total of 80 flies were used in this experiment.

Figure S4: No consistent effect of *Paillotin* mutation on resistance to a broad spectrum of microbes. w^{1118} flies were utilized as wild-type controls, while spz^{rm7} flies lack the Toll pathway, and $Relish^{E20}$ flies lack the Imd pathway, were used as susceptible models for all survival experiments against infections. Tm and OD600 are indicated. Each survival assay was performed with a minimum of 80 flies.

Figure S5: Paillotin did not show microbicidal activity *in vitro*. (*a*) The growth of *P. burhodogranariea* was monitored in the presence of the Paillotin peptide (100 µM) and/or the Cecropin A peptide (2.5 µM). The growth of *Ecc15* (*b*) and *E. coli* (*c*) was monitored in the presence of the Paillotin peptide (100 µM). Optical density (OD) values were measured every 10 minutes over 20 hours to generate bacterial growth curves. Different colored curves represent the conditions tested, as indicated in the figure legend.

Figure S6: *Paillotin* shows a minor synergistic effect with *Group B* mutants against other *Providencia* species at the doses tested. (*a*) Both *Group B/+, Pai* compound mutants suffer a minor increase in mortality compared to *Group B/+* alone upon *Providencia rustigianii* infection. Although the synergistic effect between Paillotin and Group B was only marginally significant (Mutant, Cox Hazard Ratio (HR), *p*-value: *Paillotin*, HR =-0.01, *p*=0.904; *Group B*, HR =-2.8, *p*<0.001; *Paillotin*Group B*, HR = -0.41, *p*=0.09). (*b*) *Group B/+, Pai*$^{\triangle 41}$ mutants suffer a mortality increase compared to *Group B/+*. In addition, *Group B/+ Pai*$^{\triangle 370}$ shows no difference from *Group B/+*, but also *Group B, Pai*$^{\triangle 370}$ compound mutants survive better than *Group B* alone, indicating a complex phenotype after recombination for this infection that may mask the contribution of *Pai*$^{\triangle 370}$ at the chosen low-mortality dose (Mutant, Cox Hazard Ratio (HR), *p*-value: *Paillotin*, HR =0.15, *p*=0.14; *Group B*, HR =-2.7, *p*<0.001; *Paillotin*Group B*, HR = -1.13, *p*=0.03). Differences between *Pai*$^{\triangle 41}$ (truncated) and *Pai*$^{\triangle 370}$ (loss of start) may explain this difference. (*c*) No survival difference was observed for *Group B, Pai* compound mutants against *E. coli*. N=total number of flies in experiments.

Figure S7: Overexpression of *Paillotin* does not confer protection against *P. burhodogranariea* infection in wild-type flies. Survival curves of flies overexpressing *Paillotin* by using *C564-Gal4* and *Actin-Gal4* drivers following septic infection with *P. burhodogranariea*. Survival assay was performed with a minimum of 120 flies.

Figure S8: Paillotin does not contribute to protection against fungal toxins. (*a*) Survival of wild-type and *Paillotin* mutant flies following injection with 9.2 nl of 8 mM Destruxin A, a mycotoxin secreted by *M. robertsii*. (*b*) Survival of wild-type and *Paillotin* mutant flies injected with 4.6 nl of 1 mg/ml Verruculogen, a neurotoxin produced by *A. fumigatus*. (*c*) Survival of all tested fly strains following injection with DMSO as vehicle control. *spz^rm7* flies are used as positive controls. N=total number of flies in experiments.

Figure S9: Paillotin is not involved in melanization response and microbiota regulation.
(*a*) Cuticle blackening after clean injury is not impaired except in melanization-deficient flies
(*PPO1$^\Delta$, PPO2$^\Delta$*). (*b*) CFUs per fly were measured for wild-type, *Paillotin* mutants, and the 14
AMP-deficient mutant (ΔAMP14). Data were analyzed using one-way ANOVA followed by
Dunnett's multiple comparisons test with w^{1118} as the control. A significant increase in bacterial
load was observed only in ΔAMP14, while no significant differences were detected between
wild-type and *Paillotin* mutants. Data points represent the mean bacterial load per fly, derived
from individual experiments using pools of five flies per genotype (*** $p<0.001$, n.s. = not
significant, $p>0.05$).

Table S2: Drosophila strains used in this study.

Drosophila lines	Reference
iso; iso; Rel^{E20}	Hedengrenetal.,1999
iso; iso; spz^{rm7}	Lemaitre et al.,1996
iso; Bom$^{\triangle55}$; iso	Hanson et al., 2019
iso; iso; Rel^{E20}, spz^{rm7}	Ryckebusch et al., 2025
iso; Pai$^{\triangle41}$; iso	DGRP
iso; Pai$^{\triangle370}$; iso	DGRP
; Pai$^{\triangle SFHI}$;	Guangzhou Drosophila Resource Center
; ; UAS-Pai	This paper
iso ;Group B; iso	Hanson et al.,2019
iso ; PPO1$^{\triangle}$,PPO2$^{\triangle}$; iso	Bingelli et al., 2014
$\triangle$AMP 14	Carboni et al., 2022
C564-Gal4	BDSC
Actin-Gal4	BDSC

Table S3: Microbial strains used in this study.

Type	Species	Strain,Source, DOI of previous study
Gram-negative	*Escherichia coli*	1106
	Pectobacterium carotovorum	Ecc15
	Providencia rettgeri	Dmel
	Enterobacter cloacae	B12
	Providencia burhodogranariea	B
	Providencia rustigianii	DSM 4541
	Providencia vermicola	DSM 17385
Gram-negative	*Micrococcus luteus*	doi:10.7554/eLife.44341
	Bacillus subtilis	168
	Enterococcus faecalis	254
	Listeria monocytogenes	BUG 1600
	Staphylococcus aureus	doi:10.7554/eLife.44341
Fungi	*Beauveria bassiana*	R444
	Metarhizium anisopliae	ARSEF-2575
	Metarhizium robertsii	ARSEF-2575