p53-induced RNA-binding protein ZMAT3 inhibits transcription of a hexokinase to suppress mitochondrial respiration

Ravi Kumar[1], Simon Couly[2], Bruna R. Muys[1], Xiao Ling Li[1], Ioannis Grammatikakis[1], Ragini Singh[1], Mary Guest[3], Xinyu Wen[4], Wei Tang[5], Stefan Ambs[5], Lisa M. Jenkins[6], Erica C. Pehrsson[7], Raj Chari[3], Tsung-Ping Su[2] & Ashish Lal[1*]

[1]Regulatory RNAs and Cancer Section, Genetics Branch, Center for Cancer Research (CCR), National Cancer Institute (NCI), National Institutes of Health (NIH), Bethesda, MD, USA
[2]Cellular Pathobiology Section, Integrative Neuroscience Branch, National Institute on Drug Abuse (NIDA), NIH, Baltimore, MD, USA
[3]Genome Modification Core, Frederick National Lab for Cancer Research, NCI, NIH, Frederick, MD, USA.
[4]Oncogenomics Section, Genetics Branch, CCR, NCI, NIH, Bethesda, MD, USA
[5]Laboratory of Human Carcinogenesis, CCR, NCI, NIH, Bethesda, MD, USA
[6]Mass Spectrometry Section, Laboratory of Cell Biology, CCR, NCI, NIH, Bethesda, MD, USA
[7]Advanced Biomedical Computational Science, Frederick National Laboratory for Cancer Research, Frederick, MD, USA

*Corresponding author: ashish.lal@nih.gov

Key words: p53, ZMAT3, HKDC1, JUN, RNA-binding protein, hexokinase, mitochondrial respiration, proliferation, proteomics, transcription, tumor suppressor, cancer.

Abstract

The tumor suppressor p53 is a transcription factor that controls the expression of hundreds of

genes. Emerging evidence suggests that the p53-induced RNA-binding protein ZMAT3 is a key

splicing regulator that functions in p53-dependent tumor suppression *in vitro* and *in vivo*.

However, the mechanism by which ZMAT3 functions in the p53 pathway is largely unclear.

Here, we discovered a function of ZMAT3 in inhibiting transcription of *HKDC1*, a hexokinase

that regulates glucose metabolism and mitochondrial respiration. Using quantitative proteomics,

we identified HKDC1 as the most significantly upregulated protein in *ZMAT3*-depleted

colorectal cancer cells. *ZMAT3* depletion results in increased mitochondrial respiration that was

rescued upon depletion of *HKDC1*, suggesting that HKDC1 is a critical downstream effector of

ZMAT3. Unexpectedly, ZMAT3 did not bind to the *HKDC1* RNA or DNA but the identification

of the ZMAT3-interactome uncovered its interaction with the oncogenic transcription factor

JUN. ZMAT3 depletion resulted in increased JUN binding at the *HKDC1* promoter and increased

HKDC1 transcription that was rescued upon JUN depletion, suggesting that JUN activates

HKDC1 transcription in ZMAT3-depleted cells. Collectively, these data reveal a mechanism by

which ZMAT3 regulates transcription and demonstrates that *HKDC1* is a key component of the

ZMAT3-regulated transcriptome in the context of mitochondrial respiration regulation.

Introduction

TP53 is the most frequently mutated gene in human cancer and functions as a major tumor suppressor[1]. *TP53* mutations in the germline of Li Fraumeni patients and in sporadic cancer, are mostly missense mutations that occur in the DNA-binding domain resulting in loss of tumor suppressor function and in some cases, gain of oncogenic functions[2-4]. Deletion of the *Trp53* gene in mice results in spontaneous tumor development within 6 months of age at 100% penetrance, underscoring the importance of p53 as a tumor suppressor[5,6]. Mechanistically, p53 functions as a sequence-specific transcription factor activating the expression of hundreds of genes that control diverse cellular processes including but not limited to, cell cycle arrest, apoptosis, senescence and DNA repair[7,8]. Despite the undisputed role of p53 in tumor suppression, our understanding of how p53 target genes mediate the effects of p53 is not fully understood.

Among the p53 target genes, some such as *p21* (*CDKN1A*) control p53-dependent cell cycle arrest, whereas *PUMA* and *NOXA* are critical in inducing apoptosis downstream of p53[9-11]. *In vivo* studies in mice have demonstrated that *Abca1*, *Gls2*, *Mlh1*, *Padi4* and *Zmat3* play key roles in mediating the tumor suppressor effects of p53[12-15]. However, triple knockout mice lacking *p21*, *Puma* and *Noxa*, which are regulators of cell cycle arrest and apoptosis in the p53 pathway, do not develop tumors spontaneously, unlike p53 null mice[16]. This has led to search for new mechanisms and effectors of p53-mediated tumor suppression.

An emerging potent mediator of p53 is the p53 target gene *ZMAT3*, which functions as an RNA-binding protein and studies in mice strongly implicate *Zmat3* as a tumor suppressor[17-19]. Mechanistically, using transcriptome-wide approaches from crosslinked cells, we and others recently reported that ZMAT3 directly binds to intronic sequences in thousands of pre-mRNAs and regulates alternative splicing[17,20]. ZMAT3 has also been shown to interact with AU-rich elements in the 3′ untranslated regions (UTR) of target mRNAs either stabilizing its targets or promoting their decay[21-24]. A deeper understanding of the molecular mechanisms by which ZMAT3 functions is necessary to better understand how ZMAT3 functions in p53-mediated tumor suppression.

Here, to better understand the function of ZMAT3, we identified the ZMAT3-interactome and the proteins regulated by ZMAT3. Unexpectedly, this approach revealed that ZMAT3

inhibits mitochondrial respiration by interacting with the transcription factor JUN (c-Jun) to
inhibit transcription of the hexokinase *HKDC1* (hexokinase domain containing 1). We chose to
focus on *HKDC1* because by quantitative proteomics, HKDC1 was the most strongly up-
regulated protein in ZMAT3-depleted colorectal cancer (CRC) cells. Hexokinases are the first
rate-limiting enzymes in the glucose metabolic pathway that phosphorylate glucose to glucose-6-
phosphate and thereby modulate glycolysis, oxidative phosphorylation, and the pentose
phosphate pathway. There are four classic hexokinases (HK1-4), and recently HKDC1 was
identified as the fifth hexokinase[25]. Besides its hexokinase function, HKDC1 also interacts with
the mitochondrial membrane to maintain mitochondrial homeostasis and plays an important role
in preventing cellular senescence[26-28]. Although, HKDC1 is reported to be overexpressed in
many cancers and high HKDC1 expression is associated with poor clinical outcome[27-30], how
HKDC1 expression is regulated remains largely unclear. Our findings provide novel insights on
the regulation and function of HKDC1 in the p53 pathway via transcriptional inhibition by a
ZMAT3/JUN axis.

Results

The hexokinase HKDC1 is the most strongly upregulated protein in ZMAT3-knockout cells

To investigate the mechanism by which ZMAT3 promotes growth suppression, we utilized
CRISPR/Cas9 to knockout *ZMAT3* in HCT116 cells (CRC) using two sgRNAs flanking the p53
response element in the second intron of the *ZMAT3*. Although this approach did not completely
delete the region spanning the two sgRNAs, a ~57 bp region near sgRNA#1 was deleted
resulting in >75% decrease in *ZMAT3* mRNA levels (Figure 1A and B). At the protein level,
ZMAT3 was strongly induced only in *ZMAT3*-WT (wild-type) cells upon treatment with Nutlin,
a small molecule that upregulates p53 and its target genes (Figure S1A). As expected, upon
Nutlin treatment, p53 and its known target p21, were induced in both *ZMAT3*-WT and isogenic
ZMAT3-KO (knockout) cells (Figure S1A). Depletion of ZMAT3 resulted in increased
proliferation and clonogenicity, consistent with previous reports (Figure 1C and S1B)[18-20].

To identify the genes regulated by ZMAT3, we next performed RNA-seq from biological
triplicates of *ZMAT3*-WT and *ZMAT3*-KO HCT116 cells. As expected, *ZMAT3* mRNA levels
markedly decreased (~7.5-fold) in *ZMAT3*-KO cells (Table S1); the recently identified direct
ZMAT3 target gene *MDM4*[17] was modestly but significantly upregulated in *ZMAT3*-KO cells

(Table S1). Transcriptome-wide, upon loss of ZMAT3, 606 genes were significantly up-regulated

(adj. p<0.05 and 1.5-fold change) and 552 were down-regulated with a median fold change of

1.76 and 0.55 for the up- and downregulated genes, respectively (Figure 1D and Table S1).

Because we and others recently reported that ZMAT3 directly regulates alternative splicing [17,20],

we reasoned that ZMAT3-dependent changes in splicing could lead to altered protein levels

without altering mRNA levels. We therefore performed quantitative proteomics from *ZMAT3*-

WT and *ZMAT3*-KO HCT116 cells. At the protein level 228 proteins were significantly (p<0.05)

up-regulated and 108 were down-regulated upon loss of ZMAT3 (Figure 1E) (Table S2).

ZMAT3 directly binds to intronic sequences in pre-mRNAs and can inhibit inclusion of

the neighboring exon[17,20]. Depletion of ZMAT3 can therefore result in increased target gene

expression. Gene set enrichment analysis (GSEA) for the proteins upregulated in *ZMAT3*-KO

cells revealed glycolysis as the most significantly overrepresented biological process (Figure 1F).

GSEA analysis from our RNA-seq data showed that glycolysis was among the top 10

overrepresented biological processes for the mRNAs upregulated in *ZMAT3*-KO cells (Figure

S1D). Interestingly, the protein that was most strongly upregulated (~3.4-fold, p<0.05) in

ZMAT3-KO cells was the hexokinase HKDC1 (Figure 1E, G and Table S2). The increase in

HKDC1 expression was also observed at the mRNA level (Figure 1H, S1C and Table S1); as

expected, *ZMAT3* mRNA levels were decreased in the *ZMAT3*-KO cells. We therefore chose to

focus on *HKDC1* because it was the most strongly upregulated protein upon loss of ZMAT3 and

is directly involved in regulating glucose metabolism and mitochondrial respiration that are

cellular processes not been previously associated with ZMAT3.

ZMAT3 inhibition of *HKDC1* expression is conserved and observed in diverse cell types

To validate inhibition of *HKDC1* expression by ZMAT3, we next performed RT-qPCR from

ZMAT3-WT and *ZMAT3*-KO HCT116 cells. We observed ~4-fold upregulation of *HKDC1*

mRNA upon ZMAT3 depletion (Figure 2A). The increase in HKDC1 expression was further

confirmed at the protein level by immunoblotting (Figure 2B). Because these experiments were

conducted from a single *ZMAT3*-KO clone, we next analyzed our recently published RNA-seq

data[20] conducted in biological triplicates from HCT116 cells transfected with a control siRNA

(siCTRL) or ZMAT3 siRNAs (SMARTpool of 4 siRNAs). *ZMAT3* mRNA was strongly down-

regulated, and *HKDC1* was modestly but significantly up-regulated (Figure S2A and Table S3), a

result that was validated by RT-qPCR (Figure 2C). GSEA analysis for the mRNAs up-regulated upon ZMAT3 knockdown showed that glycolysis was among the top 10 overrepresented biological processes (Figure S2B). Additionally, comparison of the RNA-seq data from the *ZMAT3*-WT vs *ZMAT3*-KO and CTRL siRNA vs ZMAT3 siRNA transfected HCT116 cells, indicated that 1023 genes were commonly up-regulated, and 1042 genes were commonly down-regulated upon loss of ZMAT3 (Figure S2C and D), suggesting that ZMAT3 depletion results in altered expression of thousands of genes. GSEA analysis for the top 500 mRNAs up-regulated upon ZMAT3 knockdown showed that glycolysis was among the top 10 overrepresented biological processes (Figure S2E). Collectively, these data indicate that ZMAT3 inhibits the expression of genes that play important roles in glycolysis.

To determine if ZMAT3 inhibits *HKDC1* expression in multiple cell lines, we performed RT-qPCR after ZMAT3 knockdown in SW1222 (CRC cells), HCEC-1CT (immortalized human colonic epithelial cells), and HepG2 (liver cancer cells). We observed significant increase in *HKDC1* mRNA levels upon ZMAT3 knockdown in these cell lines (Figure 2D). At the protein level, we observed modest increase in HKDC1 levels upon knockdown of ZMAT3 in HCT116 and HepG2 cells (Figure 2E). Moreover, we explored recently published RNA-seq data from *Zmat3* knockout mouse embryonic fibroblasts (MEFs)[31]. *Hkdc1* mRNA was significantly up-regulated (~6-fold) in *Zmat3*-KO MEFs (Figure 2F). And we also observed the expected upregulation of its target gene *Mdm4* while no changes in *Trp53* mRNA expression were observed.

In the context of human CRC, in the TCGA colorectal adenocarcinoma (COAD) cohort, *HKDC1* mRNA levels were significantly higher in tumors as compared to normal tissues (Figure 2G). Further analysis of the TCGA COAD data showed that as compared to p53 wild-type tumors, mutant p53 tumors exhibited significantly higher *HKDC1* mRNA levels (Figure 2G). As expected for the p53 target ZMAT3 and as shown previously in other cancer types[17], ZMAT3 mRNA levels were significantly lower in the mutant p53 tumors as compared to p53 wild-type tumors (Figure S2F and G). Furthermore, RNA-seq from *Trp53* knockout MEFs from the same study mentioned above[31] showed significant upregulation (~8.6-fold) of *Hkdc1* mRNA, and down-regulation of its target genes *Zmat3* and *Cdkn1a* (Figure 2H). These data suggest that

ZMAT3 and *HKDC1* mRNA expressions are negatively correlated in the p53 pathway and the

inhibition of HKDC1 expression by ZMAT3 and p53 is conserved between human and mouse.

ZMAT3 inhibits mitochondrial respiration by downregulating HKDC1

Previous studies suggest that HKDC1 plays a crucial role in regulating glucose metabolism and

cell proliferation in various cell types[27,28]. To determine whether ZMAT3 regulates glucose

metabolism and/or proliferation by regulating HKDC1 expression, we performed glucose

metabolic assays. Since HKDC1 is a hexokinase and can regulate phosphorylation of glucose to

glucose 6-phosphate when it enters the cells, we began with measuring changes in hexokinase

activity upon ZMAT3 loss. We incubated the cells with the non-catabolic glucose analog 2-deoxy

glucose (2-DG) for a short period of time and quantified the conversion of 2-DG to 2-DG6P

using a luminescence-based assay. We observed a significant increase in relative 2-DG6P levels

in *ZMAT3*-KO cells compared to WT cells. Interestingly, this increase was reversed when we

knocked down HKDC1 (Figure 3A). Furthermore, siRNA-mediated knockdown of ZMAT3 in

SW1222 and HEPG2 also showed a similar increase in hexokinase activity and the effect was

reversed upon concurrent knockdown of ZMAT3 and HKDC1, suggesting that ZMAT3 inhibits

glucose uptake, and this effect is HKDC1-dependent (Figure 3A). We then investigated whether

increased hexokinase activity upon ZMAT3 knockdown leads to increased glycolysis in cells. To

measure this, we performed Seahorse assays, measured extracellular acidification rate and

calculated the glycolysis proton efflux. It should be noted that the proton efflux rate measured

with Seahorse is reflecting the lactate production from glycolysis more than the pyruvate end

point of glycolysis that fueled the mitochondria. Thus, we also measured mitochondria activity.

Knockdown of ZMAT3 or HKDC1 by siRNAs resulted in modest increase in basal glycolysis

but it was not significant (Figure 3B).

Recent studies suggest that HKDC1 plays a significant role in regulating mitochondrial

respiration, and depletion of HKDC1 results in mitochondrial dysfunction and senescence[26,27].

We therefore determined a potential function of ZMAT3 in regulation of basal mitochondrial

respiration by regulating HKDC1 expression. We conducted Seahorse assays, and to measure

OCR (oxygen consumption rate) upon knockdown of HKDC1 or ZMAT3 or both. Interestingly,

we found that ZMAT3 knockdown results in significantly increased basal mitochondrial

respiration, and simultaneous ZMAT3 and HKDC1 knockdown rescues this effect (Figure 3C)

and this, without affecting the oxygen consumption that could occur along cellular pathways that

are not involving mitochondria. We did not observe any changes in non-mitochondrial

respiration(Figure S3). Because these phenotypes can be associated with proliferation, we next

asked whether HKDC1 regulates proliferation and whether it is an effector of ZMAT3. Live cell

proliferation assays and CCK8 cell viability assays suggested that knocking down HKDC1 in

both ZMAT3-WT and KO cells resulted in decreased cell proliferation. However, the effect of

HKDC1 knockdown was more significant in *ZMAT3*-KO cells suggesting that HKDC1 plays a

key role in promoting proliferation downstream of ZMAT3 (Figure 3D and E).

The p53/ZMAT3 axis inhibits HKDC1 expression

Because *ZMAT3* transcription is activated by p53, we next examined if the p53 pathway inhibits

HKDC1 expression. We therefore performed RNA-seq from HCT116 cells transfected with

siCTRL or sip53 (SMARTpool of 4 siRNAs). Interestingly, *HKDC1* mRNA was significantly

up-regulated (~2.25-fold) upon p53 knockdown (Figure 4A, 4B and Table S4). As expected,

upon p53 knockdown, the mRNAs encoding p53 and its target genes *p21* and *ZMAT3* were

strongly down-regulated (Figure 4A and 4B). We validated the observed up-regulation of

HKDC1 upon p53 knockdown by RT-qPCR and immunoblotting (Figure 4C and 4D). In these

experiments, mRNA and/or protein levels of p53, p21 and ZMAT3 were markedly decreased

upon p53 knockdown (Figure 4C and 4D). At a transcriptome-wide level, ~2850 genes were

differentially expressed (p<0.05) upon p53 knockdown (Table S4 and Figure S4A). Intersection

of the list of differentially expressed genes upon p53 knockdown with the ZMAT3-regulated

genes identified 351 genes that were commonly upregulated and 425 genes that were commonly

downregulated upon knockdown of p53 and ZMAT3 (Figure S4B and C). GSEA showed

enrichment of glycolysis and the p53 pathway, for the genes upregulated and downregulated

upon knockdown of p53 and ZMAT3, respectively (Figure S4D and E).

Since p53 knockdown resulted in increased HKDC1 expression, we next asked if

inducing p53 would result in down-regulation of HKDC1. Indeed, when we looked at RNA-seq

data, we observed ~40% reduction in *HKDC1* mRNA levels upon Nutlin treatment of siCTRL

transfected HCT116 cells (Figure 4E and Table S3). As expected for a direct p53 target gene,

upon p53 activation by Nutlin, *ZMAT3* and *p21* mRNAs were upregulated ~2.5- and ~6-fold,

respectively; p53 mRNA levels didn't change because Nutlin is known to specifically induce p53

protein levels (Figure 4E). At the protein level, p53 activation by Nutlin resulted in decreased HKDC1 levels and increased ZMAT3 and p21 levels in *ZMAT3*-WT cells (Figure 4F). In *ZMAT3*-KO cells, there was a modest increase in basal HKDC1 levels and a modest decrease upon Nutlin treatment (Figure 4F). Moreover, we generated doxycycline-inducible ZMAT3-FLAG-HA expressing HCT116 cells to overexpress ZMAT3. After 48 hours of doxycycline treatment ZMAT3-HA induction was confirmed by RT-qPCR and immunoblotting (Figure 4G and 4H). Furthermore, knockdown of p53 in these cells resulted in significant increase in both mRNA and protein levels of HKDC1 in untreated cells. Importantly, upon inducing ZMAT3 levels using doxycycline, the upregulation of HKDC1 mRNA and protein associated with p53 knockdown was not observed (Figure 4I and 4J). Collectively these data suggest that the p53/ZMAT3 signaling axis plays a critical role in the regulation of HKDC1 expression, where ZMAT3 operates downstream of p53 to inhibit HKDC1 expression.

ZMAT3 inhibits HKDC1 transcription by interacting with the transcription activator JUN

ZMAT3 is an RNA-binding protein that regulates alternative splicing. We therefore hypothesized that ZMAT3 directly binds to the *HKDC1* pre-mRNA and regulates its splicing. However, when we looked at our ZMAT3 PAR-CLIP data[20] we did not identify binding for ZMAT3 to *HKDC1* pre-mRNA (data not shown). Since the ZMAT3 protein has three C2H2-zinc-finger motifs it has the potential to bind to DNA (Supplementary Figure S5A)[32-34]. A recent report suggested ZMAT3 and other zinc finger proteins as DRBPs (DNA-RNA binding proteins)[33]. To examine if ZMAT3 binds to DNA and regulates HKDC1 expression directly at the transcriptional level we conducted ZMAT3-HA Cut&Run-seq and ChIP-seq in three biological replicates using anti-HA or anti-FLAG or anti-ZMAT3 antibody. However, we did not observe reproducible binding of ZMAT3 to the *HKDC1* locus or other regions of the genome in HCT116 cells (data not shown).

We therefore hypothesized that ZMAT3 inhibits HKDC1 expression by interacting with a specific transcription factor. To identify the proteins that interact with ZMAT3 we performed IPs using an anti-FLAG antibody followed by mass spectrometry from whole cell lysates of untreated or doxycycline-treated ZMAT3-FLAG-HA HCT116 cells (Figure 5A and S5B. After removing common contaminants (<10% in Crapome), this unbiased approach identified 21 ZMAT3-interacting proteins highlighted in red (Figure 5B and Supplemental Table S5). As expected, ZMAT3 was the most strongly enriched protein (~36,000-fold). Of note, due to low

ZMAT3 levels in one of the controls, the p-value for ZMAT3 enrichment was not very significant
(p=0.09) (Table S5).

Interestingly, the transcription factor JUN (c-Jun) was strongly enriched in the ZMAT3-
FLAG pulldowns (~8,500-fold). JUN is a proto-oncogene and has previously been implicated in
glucose metabolism[35,36] and p53 function[37,38] . We next utilized publicly available JUN ChIP-seq
data from various cell lines from ENCODE, with a focus on JUN binding at the HKDC1
promoter. Notably, in 3 cell lines, we found a JUN ChIP-seq peak having consensus JUN binding
motif in the *HKDC1* intron 1. This peak coincided with ChIP-seq peaks for POLR2A, H3K27Ac
and H3K4Me3 in HCT116 cells (Figure 5C). We validated the interaction between ZMAT3 and
JUN proteins by ZMAT3-FLAG IP and immunoblotting (Figure 5D). Importantly, knockdown of
JUN resulted in decreased *HKDC1* mRNA levels in ZMAT3- WT cells and rescued the elevated
HKDC1 mRNA and protein levels in ZMAT3-KO cells (Figure 5E and F). To determine if
ZMAT3 inhibits JUN binding to the *HKDC1* locus we conducted ChIP-qPCR for JUN from
ZMAT3-WT and KO cells. JUN exhibited significantly enhanced binding at the *HKDC1* intron 1
in *ZMAT3*-WT cells compared to the IgG control, and the enrichment of JUN to the *HKDC1*
promoter was significantly increased in *ZMAT3*-KO cells, indicating that ZMAT3 inhibits the
ability of JUN to bind to the *HKDC1* promoter. We next cloned a ~700 bp DNA fragment that
encompassed the JUN and POLR2A binding peaks within the *HKDC1* promoter into the pGL4
basic luciferase reporter vector, designated as pGL4 HKDC1-promoter. Remarkably, knockdown
of JUN in HCT116 cells resulted in a significant decrease in luciferase activity of the pGL4
HKDC1-promoter (Figure 5H). Conversely, ZMAT3 knockdown resulted in a marked increase in
luciferase activity relative to control and simultaneous knockdown of ZMAT3 and JUN rescued
the increase in reporter activity (Figure 5H).

Collectively, these data suggest that ZMAT3 plays a crucial role in controlling
mitochondrial respiration and cell proliferation by suppressing *HKDC1* transcription. We
propose a model according to which in *ZMAT3*-WT cells p53 drives ZMAT3 transcription,
ZMAT3 protein binds to the HKDC1 transcription activator JUN thereby inhibiting its binding
with *HKDC1* promoter which leads to transcriptional repression of HKDC1 thereby controlled
mitochondrial respiration and controlled cell proliferation. In the absence of ZMAT3, JUN binds

to the HKDC1 promoter and upregulates its expression, resulting in increased mitochondrial
respiration and increased cell proliferation (Figure 6).

Discussion

Recent studies suggest that ZMAT3 significantly contributes to the tumor suppressive effects of
p53[17,18]. At the molecular level, ZMAT3 functions as an RNA-binding protein that acts as a key
splicing factor and regulates mRNA stability[17,20]. Here, we unexpectedly found that the
transcription of *HKDC1*, the gene that is most strongly upregulated at the protein level in
ZMAT3-deficient cells, is indirectly repressed by ZMAT3 via interaction with the key
transcription factor, JUN, thereby inhibiting JUN's binding to the *HKDC1* promoter. Consistent
with the well-established function of HKDC1 in glucose metabolism and mitochondrial
respiration, we observed increased mitochondrial respiration upon upregulation of HKDC1 in
ZMAT3-depleted cells and this phenotype was rescued upon concurrent knockdown of ZMAT3
and HKDC1, suggesting that HKDC1 is a key downstream effector of ZMAT3.

ZMAT3 has been known as a p53 target gene for more than two decades[39,40], but its
physiological function and role in tumor suppression are only beginning to be understood.
ZMAT3 belongs to zinc finger family of proteins that play crucial role in regulating gene
expression through specific recognition of DNA sequences[41]. Although they are primarily known
for their involvement in transcription regulation, these proteins have also been found to interact
with RNA and proteins[42,43]. Among them, ZMAT3 is a member of the ZMAT domain-containing
family that has three zinc fingers of C2H2-type zinc fingers motif [44,45]. These domains play
crucial roles transcriptional regulation by specifically binding to target molecules such as DNA
and RNA. A recent study[33] revealed ZMAT3 as a DRBP (DNA- and RNA-binding protein), but
in our hands using Cut&Run-seq and ChIP-seq, we did not observe specific binding of ZMAT3-
FLAG or endogenous ZMAT3 to DNA (data not shown). In these experiments Cut&Run-seq and
ChIP-seq for p53 worked exceedingly well (data not shown), demonstrating that there was no
technical flaw. It may be that ZMAT3 binding to DNA is cell-type specific, or upon DNA
damage but this needs further investigation.

Our findings that ZMAT3 interacts with and inhibits binding of JUN to the HKDC1 locus
provides mechanistic insights on how ZMAT3, without directly interacting with DNA regulates
HKDC1 transcription. JUN is a protooncogene that plays a crucial role in both normal

physiological processes and tumorigenesis by regulating cell proliferation, differentiation,
senescence and metastasis[46-49]. It is a transcription factor function as a key component of AP-1
complex and promote RNA polymerase II mediated transcription of target genes[50]. It would be
interesting to determine which ZMAT3 regulated genes are regulated via JUN. Future
investigations are also needed to determine the domains of JUN that interact with ZMAT3 and
what ZMAT3 does to JUN.

Because ZMAT3 regulates alternative splicing which can lead to changes in protein levels
without altering mRNA levels, in this study we integrated quantitative proteomics data with
RNA-seq data from ZMAT3-WT and isogenic ZMAT3-KO CRC cells. Furthermore, we
integrated these data with RNA-seq from HCT116 cells upon ZMAT3 or p53 knockdown using
siRNAs, to make sure that the findings from the isogenic cell lines were not restricted to a single
KO clone and to determine the role of the p53/ZMAT3 axis in regulating these genes. This
approach identified HKDC1 as the most strongly protein upon ZMAT3 depletion. HKDC1 is
emerging as an important regulator of tumor progression and is frequently upregulated in several
cancers including CRC[51-53]. Besides its hexokinase activity, HKDC1 interacts with the
mitochondrial membrane and plays an essential role in mitochondrial function[26,27]. We further
demonstrated that ZMAT3 suppresses glucose uptake and basal mitochondrial respiration by
inhibiting HKDC1 expression leading to suppression of cell proliferation in CRC cells.

Our data also demonstrates that p53 negatively regulates *HKDC1* expression and this
effect is ZMAT3-dependent, suggesting that the p53/ZMAT3/HKDC1 axis may be an important
component of the p53 network, in the context of mitochondrial respiration and proliferation. The
ability of p53 to induce cell cycle arrest and programmed cell death is important for tumor
suppressor[54]. However, p53 has several other functions that recent data strongly implicate in
tumor suppression, particularly regarding the control of metabolism such as glycolysis,
mitochondrial respiration, and ferroptosis[55,56]. Metabolic reprogramming is a hallmark of cancer
cells, which plays a pivotal role in cancer progression by providing energy and a wide variety of
substrates for biosynthesis to support the rapid proliferation and survival of cancer cells[57-59]. p53
has been reported to play an important role in suppressing tumor development by regulating the
expression and function of metabolic genes, directly *(GLUT1*[60]*, GLUT4*[60]*, PFKFB3*[61] and
PFKFB4[62]) or indirectly (*HK2*[63]*, HIF1α*[64] and *G6PD*[65]). Our data uncovers HKDC1 as an

indirect p53 target gene that is negatively regulated via ZMAT3. Collectively, our findings

provide key insights into the diverse functions of ZMAT3 and their involvement in gene

regulation in the p53 pathway.

Materials and Methods

Cell lines

The cell lines HCEC-1CT, HepG2, HCT116 and SW1222 were purchased from the American

Type Culture Collection (ATCC). Cells were grown in DMEM medium (ThermoFisher

scientific, Catalog no. 11995065), containing 10% Fetal Bovine Serum (FBS) (ThermoFisher

scientific, Catalog no. 10082147) and 100 U/ml of penicillin and 0.1 mg/ml of streptomycin

(ThermoFisher scientific, Catalog no. 15070063), at 37°C and 5% CO2. All cell lines were

regularly screened for mycoplasma using the Venor™ GeM Mycoplasma Detection Kit

(Millipore Sigma, Catalog no. MP0025-1KT).

Targeted deletion of ZMAT3 using CRISPR/Cas9

To generate *ZMAT3*-KO clones, we used the PiggyBac CRISPR/Cas9 system adopted from the

Zhang lab (Shalem et al., Science, 2014). Two sgRNAs flanking the p53RE in the *ZMAT3* intron

2 were designed and separately cloned in pENTR221 vector. These constructs were

electroporated into $1x10^6$ parental HCT116 cells using Amaxa Cell Line Nucleofector Kit

(Lonza, Catalog no. VCA-1005) along with pT3.5-Flag-Cas9, pCDNA-pB7, and pBSB-CG-

LUC-GFP-(puro)(cre+) vectors. After two days, the cells were treated with 2 µg/ml puromycin

(ThermoFisher scientific, Catalog no. A1113803) for three days. Single cells were then seeded in

96-well plates after puromycin selection to select *ZMAT3*-WT and *ZMAT3*-KO clones. The

clones were harvested three weeks later and split into 24-well plates. Total RNA was isolated

from each well of the 24-well plate, and ZMAT3 expression was measured by RT-qPCR

normalized to *GAPDH*. Genomic DNA was extracted from individual clones in which ZMAT3

expression was profoundly decreased, and the DNA flanking the p53RE of ZMAT3 was PCR

amplified and subjected to Sanger sequencing.

Plasmids construction and lentivirus production

The pLVX-Puro-Tet-One vector from TaKaRa (631849) was used as a backbone to construct an

expression vector containing ZMAT3-3xFLAG-2xHA. The vectors were transformed into DH5α

cells (ThermoFisher scientific, Catalog no. 18265017), and the plasmids were purified using the

Monarch plasmid miniprep kit (NEB T1010L). Lentiviruses were produced in 3×10^5 293T cells

after co-transfection of 1 µg DNA with a third-generation lentivirus packaging system using

Lipofectamine 2000 (Thermo Fisher Scientific Catalog no. 11668027). HCT116 cells were

transduced at an MOI~1, and after 2 days, the cells were treated with 2 µg/mL puromycin

(ThermoFisher scientific, Catalog no. A1113803) for 1 week.

Wild-type (pGL4-Basic-HKDC1-WT-promoter) POLR2A binding site present at the HKDC1

promoter construct were cloned into the pGL4-Basic vector (Promega, Catalog no. E6651) for

luciferase assays. We used gene fragments (Twist Bioscience) at region chr10:69,222,339-

69,223,043 (hg38) as HKDC1 JUN binding site, containing restriction sites from KpnI (NEB,

Catalog no. R3142) and XhoI (NEB, Catalog no. R0146S) at 5′ and 3′, respectively. We digested

the pGL4-Basic vector and DNA fragments with these enzymes, purified the products using the

Monarch DNA gel extraction kit (NEB, Catalog no. T1020S) or QIAquick PCR purification kit

(Qiagen, Catalog no. 28106), and ligated the inserts with the vector using T4 DNA ligase (NEB,

Catalog no. M0202S)

siRNA transfections

We used reverse transfection to deliver siRNAs with the use of Lipofectamine RNAiMAX

Transfection Reagent (ThermoFisher Scientific Catalog no.13778075) and optiMEM

(ThermoFisher Scientific Catalog no. 31985062) in HCT116, HCEC-1CT, HepG2 and SW1222

cells according to the manufacturer's protocol. The final concentration of siRNA was 20 nM. For

RT-qPCR and immunoblotting, we conducted two rounds of transfection. The second round was

conducted 48 hours after the first transfection conducted for 72 hours. The HKDC1 transfection

was conducted for only one round after the first transfection with siCTRL or siZMAT3. We used

Negative Control siRNA (Qiagen, Catalog no. 1027281) as control. We used SMARTPool

siRNAs against ZMAT3 (Horizon Discovery, Catalog no. L-017382-00-0005), siJUN (Horizon

Discovery, Catalog no. L-003268-00-0005) and p53 (Horizon Discovery, Catalog no. L-003329-

00-0005). For metabolic assays, we transfected cells using siRNAs against more than one target

(e.g. siHKDC1 and siZMAT3) at 20nM each. HCT116 expressing the ZMAT3-FLAG-HA were

transfected for 48 h and reseeded, the cells were treated with 2μg/ml doxycycline to induce the

ZMAT3 expression.

Luciferase assays

Cells were transfected with siRNAs for 48 hours, and then $1x10^5$ cells were reseeded for

luciferase assay in 24-well plates. The next day, the cells were co-transfected with 250 ng of

pGL4-ZMAT3-WT-promoter and 25 ng of pRL-TK (Promega, Catalog no. E2231) vectors, along

with 20 nM of either Allstars negative control siRNA, siJUN or siZMAT3. Lipofectamine 2000

transfection reagent (ThermoFisher Scientific, Catalog no. 11668027) was used to carry out co-

transfection, as per manufacturer's protocol. After 2 days, firefly and Renilla luminescence from

pGL4 and pRL-TK vectors, respectively, were measured using the dual-luciferase reporter assay

system (Promega, Catalog no. E1910) according to the manufacturer's protocol on EnSight

Multimode plate reader (PerkinElmer). Firefly luminescence was normalized with Renilla

luminescence for transfection efficiency.

RNA extraction and RT-qPCR

Cells that were used to analyze the expression via RT-qPCR were washed with DPBS 1X

(ThermoFisher Scientific, Catalog no. 14190250) after 48 hours of transfection. Then, they were

lysed using 500 μl of TRIzol Reagent (ThermoFisher Scientific, Catalog no. 15596018) 48 hours

after transfection, and RNA was extracted based on the manufacturer's protocol. To prepare

cDNA, 500 ng of RNA was reverse transcribed using iScript™ Reverse Transcription Supermix

(Biorad, Catalog no. 1708841). In the qPCR reaction, 2.5 μl of diluted cDNA was combined with

5 μl of 2x FastStart Universal SYBR Green Master (Rox) (Millipore Sigma, Catlog no.

4913914001), 0.5 μM (final concentration) of each primer in a reaction with a final volume of 10

μl. The reactions were executed on StepOnePlus Real-Time PCR machine (Applied Biosystems),

and *GAPDH* was used as a loading control. Finally, the relative expression was calculated using

the 2-ΔΔCt method.

Immunoblotting

For immunoblotting, cells were lysed using RIPA buffer (ThermoFisher Scientific, Catlog no.

89901). The lysates were sonicated three times for five seconds each at a power set of 50% using

a VirTis VIRSONIC 100. Then, the lysates were centrifuged at 13000g for 10 minutes at 4°C,

and the supernatant was collected. Pierce BCA Protein Assay Kit (ThermoFisher Scientific,
Catalog no. 23225) was used for protein quantitation. For SDS-PAGE the gel was loaded with 20
to 50 µg of protein, transferred to a PVDF membrane using a BioRad semi-dry transfer apparatus
and the membrane was blocked for 1 hour with TBST (Tris-Buffered Saline - 19.98 mM Tris,
136 mM NaCl and Tween 0.05%, pH 7.4) containing 5% milk. Anti-GAPDH antibody (1:6000
dilution; Cell Signaling, 5174S) was used for loading control. The following primary antibodies
were used: anti-FLAG (1:1000 dilution; Sigma F1804), anti-p53 (DO-I) (1:1000 dilution; Santa
Cruz Biotechnology sc-126), anti-ZMAT3 (1:500 dilution; Santa Cruz Biotechnology sc-
398712), anti JUN (1:1000 dilution; Cell signalling, 9165S) and anti-HKDC1 (Proteintech,
Catlog no: 25874-1-AP). After 1 hour of secondary antibody incubation at 1:5000 dilution, the
membranes were developed using ECL™ Prime Western Blotting Detection Reagent (Fisher
Scientific, Catalog no. RPN2232).

Colony formation assays

One thousand ZMAT3-WT and isogenic ZMAT3-KO HCT116 cells were seeded in 6-well
plates. After 12 days, the cells were fixed with ice-cold methanol for 15 minutes and stained with
0.5% crystal violet (prepared in 10% methanol) for 15 minutes. We used the ImageJ software
(version 2.0.0-rc-43/1.52n) to analyze images of the colony coverage area.

Incucyte Proliferation assays

To conduct the proliferation assays, 1000 cells were seeded per well in 96-well plates. The cells
were then incubated on an Incucyte® S3 Live-Cell Analysis Instrument and photographed every
6 hours for 4 days. The images were analyzed using the manufacturer's software to measure %
confluence over time.

Cell viability assays

To determine cell viability, cells were incubated with cell counting kit-8 (Dojindo, Kumamoto,
Japan) for 4 hour and absorbance at 450 nm was measured using a microplate reader Envision
(PerkinElmer).

Glucose uptake assays

siCTRL, siZMAT3 and siHKDC1 were transfected into HCT116, SW1222, and HepG2 cells in

poly-l-lysine–coated white 96-well plates with opaque bottoms (Costar) and incubated at 37°C

for 24 hours. After the incubation period, the growth medium was removed, and cells were

washed twice with PBS to eliminate residual glucose. The Glucose Uptake-Glo kit (Promega,

Catalog no. J1341) was used to measure cellular 2-deoxyglucose uptake, according to the

manufacturer's instructions. Glucose uptake was initiated by the addition of 1 mM 2-

deoxyglucose for 10 minutes at 37°C following luminescence measurement in Envision

Instrument.

Metabolic flux assays

First, HCT116 cells were reverse transfected with siCTRL or siZMAT3 for 48 hours. After 48

hours, a second round of transfection was conducted in 24well plates Seahorse (Agilent

Technologies Inc, Catalog no. 100777-004,) at a concentration of $3x10^4$ per 200µL per well.

After 48 hours cells were washed twice with 500mL of XF DMEM Medium, pH 7.4 (Agilent

Technologies Inc, Catlog no. 103575-10) containing 1mM pyruvate, 2mM of glutamine and 10

mM of glucose (Agilent Technologies Inc, Catalog no. 103578-100, 103579-100, 103577-100).

Eventually cells were incubated 45 minutes in a non-CO2 incubator prior to the assays.

Meanwhile drugs from Mito Stress Test Kit (Agilent Technologies Inc, Catalog no. 103015-10)

and Glycolytic Rate Assay Kit (Agilent Technologies Inc, Catalog no. 103344-100) were

prepared in XF DMEM Medium, pH 7.4 containing 1mM pyruvate, 2mM of glutamine and

10mM of glucose. For the Mito Stress Test Kit the working drug solutions concentration were

Oligomycin 1.5µM, FCCP 1.0µM and Rot/AA 0.5µM. For the glycolytic Rate Assay Kit the

working drug solutions concentration were Rot/AA 0.5µM and 2-DG 50mM. After the 45min of

incubation, the plates were loaded into the Seahorse Analyzer and the commercial protocols for

the drugs distribution were used. Immediately after the assays, media was removed from wells

and cells frozen for later protein concentration measurement. Protein concentrations were

measured with PierceTM BCA Protein Assay kit (ThermoFisher Scientific, Catalog no. 23225)

and used to normalize Seahorse results**.**

Gene set enrichment analysis

GSEA using MSigDB Hallmark gene sets was performed on the website of the Broad Institute

(http://www.gsea-msigdb.org/gsea/msigdb/annotate.jsp). The proteins that showed significant

(p<0.05) upregulation upon ZMAT3-KO were used as input.

Co-immunoprecipitation

ZMAT3-FLAG coimmunoprecipitations were performed using FLAG M2 antibody-coated

magnetic beads (Sigma-Aldrich Catalog no. M8823). 5 x 107 doxycycline treated/untreated

HCT116 ZMAT3-FLAG-HA cells lysed in IP lysis buffer (10 mM Tris/Cl pH 7.5, 150 mM NaCl,

0.5 mM EDTA, 0.5 % NP40) supplemented with 1 mM PMSF, complete protease inhibitor

cocktail (Roche). Lysates were incubated for 30 min at 4°C with periodic mixing and clarified by

centrifugation at maximum speed for 10 min at 4°C. Equal amounts of protein lysates were

incubated with washed M2 beads overnight at 4°C with constant rotating. Beads were

magnetically separated from supernatant and washed four times with IP Wash buffer (10 mM

Tris/Cl pH 7.5, 150 mM NaCl, 0.05 % NP40, 0.5 mM EDTA). Bound proteins were eluted with

FLAG elution buffer supplemented with 125 µg/ml 3xFLAG peptide [F4799; Sigma-Aldrich]).

Equal volumes of eluates were boiled at 100°C for 5 min in Laemmli sample buffer and then

centrifuged at maximum speed for 5 min at RT. Five percent of the total cell lysate was used as

input, and proteins were detected by immunoprecipitation. For IP mass spectrometry, samples

were processed after washing without elution step.

ChIP-qPCR

ChIP-qPCR was carried out using an Active Motif ChIP-IT express kit (catalog no. 53008)

following the manufacturer's instructions. In brief, 5×10^7 ZMAT3-WT and ZMAT3 KO

HCT116 cells grown in 15-cm plates were cross-linked with 1% formaldehyde, scraped, lysed,

and then sheared. The size of the chromatin was verified on a 1% agarose gel. Chromatin was

immunoprecipitated at 4°C overnight with 1µg of anti-JUN and IgG antibody. The IP material

was washed, eluted, and reverse cross-linked overnight at 65°C. ChIP DNA was column purified

(Qiagen PCR purification, catalog no. 28104) and analyzed by qPCR. ChIP-qPCR primers were

designed based on the genomic sequence of the JUN peaks found in the HKDC1 intron-1.

RNA-seq and analysis

RNA-seq was performed in biological triplicates from HCT116 ZMAT3-WT and ZMAT3-KO
cells. Total RNA was isolated using the RNeasy Plus mini kit (Qiagen, Catalog no. 74134)
following the manufacturer's instructions. We used the Illumina Stranded mRNA Ligation library
Kit. We used 450ng of total RNA as the input to an mRNA capture with oligo-dT coated
magnetic beads. The mRNA is fragmented, and then a random-primed cDNA synthesis is
performed. The resulting double-strand cDNA is used as the input to a standard Illumina library
prep with end-repair, adapter ligation and PCR amplification being performed to give you a
sequencing ready library. The final purified product is then quantitated by qPCR before cluster
generation and paired end sequencing on NovaSeq 6000 SP 200 cycles run.

The Illumina bcl2fastq2.20 was used to demultiplex and convert binary base calls and quality
scores to Fastq format. The sequencing reads were trimmed for adapters and low-quality bases
using Cutadapt (v1.18). The trimmed reads were mapped to human reference genome (hg38)
and Gencode annotation v30 using STAR aligner (v2.7.0f) with two-pass alignment option.
RSEM (v1.3.1) was used for gene and transcript quantification based on GENCODE annotation.

Mass Spectrometry Sample Preparation

For total protein identification, cell pellets were suspended in 8M urea buffer supplemented with
protease and phosphatase inhibitors (Roche). All samples were transferred to 2 ml TissueLyser
tubes containing 5 mm steel balls on ice. Samples were lysed in a TissueLyser (Qiagen) for 2 X 2
mins with chilling in-between at -20°C for 2-3 mins. Lysates were centrifuged in a
microcentrifuge at 12,500 rpm for 15 min (4°C). Supernatants were transferred to new tubes and
proteins concentrations were measured using BCA method (Thermo). For downstream
processing, 200 ug of each sample were reduced with 10 mM DTT at 56°C for 1 hour and
alkylated with 20 mM iodoacetamide at room temperature for 30 minutes in the dark. Following
alkylation, samples were diluted 4-fold with 50 mM triethylammonium bicarbonate (TEAB) to
reduce the urea concentration to 2 M and digested with trypsin (40:1) at 37°C overnight.
Digested peptides were desalted using C18 columns and lyophilized. Peptide concentrations
were measured using colorimetric BCA peptide assay (Thermo). For TMT labeling, 100 ug of
digested peptides were labeled with the TMTpro 16-plex reagent at room temperature (dark) for
1 hour. Reactions were stopped by addition of 5% hydroxyl amine and incubation at room
temperature (dark) for 15 mins. Following labeling, peptide samples were pooled and

lyophilized. Lyophilized pooled samples were suspended in 0.1% TFA and fractionated using
high pH Reverse-Phase Peptide Fractionation Kit (Thermo) with nine elution buffers containing
0.1% Triethylamine and 10%, 12.5%, 15%, 17.5%, 20%, 22.5%, 25%, 50%, and 75%
acetonitrile, respectively. Fractions were lyophilized separately.

For ZMAT3-FLAG interacting protein identification, IP samples were solution digested with
trypsin using S traps (Protifi), following the manufacturer's instructions. Briefly, proteins were
denatured in 5% SDS, 50 mM triethylammonium bicarbonate (TEAB) pH 8.5. They were next
reduced with 5 mM Tris(2-carboxyethyl)phosphine (TCEP) and alkylated with 20 mM
iodoacetamide. The proteins were acidified to a final concentration of 2.5% phosphoric acid and
diluted into 100 mM TEAB pH 7.55 in 90% methanol. They were loaded onto the S-traps,
washed four times with 100 mM TEAB pH 7.55 in 90% methanol, and digested with trypsin
overnight at 37 °C. Peptides were eluted from the S-trap using 50 mM TEAB pH 8.5; 0.2%
formic acid in water; and 50% acetonitrile in water. These elutions were pooled and dried by
lyophilization.

Mass Spectrometry Analysis

Dried peptides were resuspended in 5% acetonitrile, 0.05% TFA in water for mass spectrometry
analysis on an Orbitrap Exploris 480 (Thermo) mass spectrometer. The peptides were separated
on a 75 µm x 15 cm, 3 µm Acclaim PepMap reverse phase column (Thermo) at 300 nL/min
using an UltiMate 3000 RSLCnano HPLC (Thermo) and eluted directly into the mass
spectrometer. For analysis, parent full-scan mass spectra acquired at 120,000 FWHM resolution
and product ion spectra at 45,000 resolutin with a 0.7 m/z isolation window. Proteome
Discoverer 3.0 (Thermo) was used to search the data against the human database from Uniprot
using SequestHT and with INFERYS rescoring. The search was limited to tryptic peptides, with
maximally two missed cleavages allowed. Cysteine carbamidomethylation and TMT pro
modification of lysine and peptide N-termini were set as a fixed modification, with methionine
oxidation as a variable modification. The precursor mass tolerance was 10 ppm, and the fragment
mass tolerance was 0.02 Da. The Percolator node was used to score and rank peptide matches
using a 1% false discovery rate. TMT quantitation was performed using the Reporter Ions
Quantifier nodes with correction of the values for lot-specific TMT reagent isotopic impurities.

TCGA COAD gene expression

Gene expression and clinical data of COAD were obtained using the TCGA biolinks (v2.16.0)
package in R (v4.0.0). The GDC query function was used to query samples by setting the data
category to "gene expression," data type to "gene expression quantification," platform to
"Illumina HiSeq," file type to "normalized results," and experimental strategy to "RNA-seq."
Data were then downloaded using GDC download followed by GDC prepare to obtain
normalized gene expression values.

Statistical analysis

The statistical analysis for all data was performed using at least three replicates. The significance
of statistical analyses was tested using two-tailed Student's t-test when comparing two groups or
two-way ANOVA. Pearson coefficient correlation was used for correlation analysis with the
TCGA COAD study.

Acknowledgements
We thank the CCR Genomics Core, CCR, NCI, Bethesda, MD for valuable assistance with
Sanger sequencing and Agilent TapeStation. We also thank the CCR Sequencing Facility, NCI,
Frederick, MD for performing the RNA-seq. Finally, we thank the members of the Lal lab for
discussion and suggestions.
589

590 **Conflict of interest statement**
591 The authors declare no Conflict of interest
592

593 **Author contribution statement**
594 R.K and A.L designed experiments and interpreted the results. R.K., S.C., B.M., X.L and M.G
595 performed experiments and analyzed data. R.C designed the gRNA constructs to KO ZMAT3.
596 E.P., I.G., and X.W analyzed the RNA-seq data. L.J conducted the quantitative mass
597 spectrometry. R.K., and A. L. wrote the manuscript.
598

599 **Funding**
600 This research was supported by the Intramural Research Program (A.L.) of the National Cancer
601 Institute (NCI), Center for Cancer Research (CCR), NIH (ZIA BC011646 to A.L.).

Data Availability

The RNA-seq data from ZMAT3-WT vs ZMAT3-KO HCT116 cells and from siCTRL vs sip53 from HCT116 have been deposited to the GEO (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE280756).

References

Kandoth, C. *et al.* Mutational landscape and significance across 12 major cancer types. *Nature* **502**, 333-339 (2013). https://doi.org/10.1038/nature12634
Olivier, M. *et al.* The IARC TP53 database: new online mutation analysis and recommendations to users. *Hum Mutat* **19**, 607-614 (2002). https://doi.org/10.1002/humu.10081
Malkin, D. *et al.* Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms. *Science* **250**, 1233-1238 (1990). https://doi.org/10.1126/science.1978757
Oren, M. & Prives, C. p53: A tale of complexity and context. *Cell* **187**, 1569-1573 (2024). https://doi.org/10.1016/j.cell.2024.02.043
Donehower, L. A. *et al.* Mice deficient for p53 are developmentally normal but susceptible to spontaneous tumours. *Nature* **356**, 215-221 (1992). https://doi.org/10.1038/356215a0
Donehower, L. A. The p53-deficient mouse: a model for basic and applied cancer studies. *Semin Cancer Biol* **7**, 269-278 (1996). https://doi.org/10.1006/scbi.1996.0035
Boutelle, A. M. & Attardi, L. D. p53 and Tumor Suppression: It Takes a Network. *Trends in cell biology* **31**, 298-310 (2021). https://doi.org/10.1016/j.tcb.2020.12.011
Indeglia, A. & Murphy, M. E. Elucidating the chain of command: our current understanding of critical target genes for p53-mediated tumor suppression. *Crit Rev Biochem Mol Biol* **59**, 128-138 (2024). https://doi.org/10.1080/10409238.2024.2344465
Waldman, T., Kinzler, K. W. & Vogelstein, B. p21 is necessary for the p53-mediated G1 arrest in human cancer cells. *Cancer Res* **55**, 5187-5190 (1995).
Yu, J., Wang, Z., Kinzler, K. W., Vogelstein, B. & Zhang, L. PUMA mediates the apoptotic response to p53 in colorectal cancer cells. *Proc Natl Acad Sci U S A* **100**, 1931-1936 (2003). https://doi.org/10.1073/pnas.2627984100
Shibue, T. *et al.* Integral role of Noxa in p53-mediated apoptotic response. *Genes Dev* **17**, 2233-2238 (2003). https://doi.org/10.1101/gad.1103603
Indeglia, A. *et al.* An African-Specific Variant of TP53 Reveals PADI4 as a Regulator of p53-Mediated Tumor Suppression. *Cancer Discov* **13**, 1696-1719 (2023). https://doi.org/10.1158/2159-8290.CD-22-1315
Janic, A. *et al.* DNA repair processes are critical mediators of p53-dependent tumor suppression. *Nat Med* **24**, 947-953 (2018). https://doi.org/10.1038/s41591-018-0043-5

14 Moon, S. H. *et al.* p53 Represses the Mevalonate Pathway to Mediate Tumor
Suppression. *Cell* **176**, 564-580 e519 (2019).
https://doi.org/10.1016/j.cell.2018.11.011
15 Suzuki, S. *et al.* GLS2 Is a Tumor Suppressor and a Regulator of Ferroptosis in
Hepatocellular Carcinoma. *Cancer Res* **82**, 3209-3222 (2022).
https://doi.org/10.1158/0008-5472.CAN-21-3914
16 Valente, L. J. *et al.* p53 efficiently suppresses tumor development in the complete
absence of its cell-cycle inhibitory and proapoptotic effectors p21, Puma, and
Noxa. *Cell Rep* **3**, 1339-1345 (2013). https://doi.org/10.1016/j.celrep.2013.04.012
17 Bieging-Rolett, K. T. *et al.* Zmat3 Is a Key Splicing Regulator in the p53 Tumor
Suppression Program. *Mol Cell* **80**, 452-469 e459 (2020).
https://doi.org/10.1016/j.molcel.2020.10.022
18 Bieging-Rolett, K. T. & Attardi, L. D. Zmat3 splices together p53-dependent tumor
suppression. *Mol Cell Oncol* **8**, 1898523 (2021).
https://doi.org/10.1080/23723556.2021.1898523
19 Brennan, M. S. *et al.* Combined absence of TRP53 target genes ZMAT3, PUMA
and p21 cause a high incidence of cancer in mice. *Cell Death Differ* **31**, 159-169
(2024). https://doi.org/10.1038/s41418-023-01250-w
20 Muys, B. R. *et al.* The p53-induced RNA-binding protein ZMAT3 is a splicing
regulator that inhibits the splicing of oncogenic CD44 variants in colorectal
carcinoma. *Genes Dev* **35**, 102-116 (2021).
https://doi.org/10.1101/gad.342634.120
21 Vilborg, A. *et al.* The p53 target Wig-1 regulates p53 mRNA stability through an
AU-rich element. *Proc Natl Acad Sci U S A* **106**, 15756-15761 (2009).
https://doi.org/10.1073/pnas.0900862106
22 Bersani, C. *et al.* Genome-wide identification of Wig-1 mRNA targets by RIP-Seq
analysis. *Oncotarget* **7**, 1895-1911 (2016).
https://doi.org/10.18632/oncotarget.6557
23 Kim, B. C. *et al.* Wig1 prevents cellular senescence by regulating p21 mRNA
decay through control of RISC recruitment. *EMBO J* **31**, 4289-4303 (2012).
https://doi.org/10.1038/emboj.2012.286
24 Bersani, C., Xu, L. D., Vilborg, A., Lui, W. O. & Wiman, K. G. Wig-1 regulates cell
cycle arrest and cell death through the p53 targets FAS and 14-3-3sigma.
Oncogene **33**, 4407-4417 (2014). https://doi.org/10.1038/onc.2013.594
25 Khan, M. W., Ding, X., Cotler, S. J., Clarke, M. & Layden, B. T. Studies on the
Tissue Localization of HKDC1, a Putative Novel Fifth Hexokinase, in Humans. *J*
Histochem Cytochem **66**, 385-392 (2018).
https://doi.org/10.1369/0022155418756849
26 Cui, M. *et al.* HKDC1, a target of TFEB, is essential to maintain both
mitochondrial and lysosomal homeostasis, preventing cellular senescence. *Proc*
Natl Acad Sci U S A **121**, e2306454120 (2024).
https://doi.org/10.1073/pnas.2306454120
27 Khan, M. W. *et al.* The hexokinase "HKDC1" interaction with the mitochondria is
essential for liver cancer progression. *Cell Death Dis* **13**, 660 (2022).
https://doi.org/10.1038/s41419-022-04999-z

28   Zapater, J. L., Wicksteed, B. & Layden, B. T. Enterocyte HKDC1 Modulates
Intestinal Glucose Absorption in Male Mice Fed a High-fat Diet. *Endocrinology*
163 (2022). https://doi.org/10.1210/endocr/bqac050
29   Wang, X. *et al.* HKDC1 promotes the tumorigenesis and glycolysis in lung
adenocarcinoma via regulating AMPK/mTOR signaling pathway. *Cancer Cell Int*
20, 450 (2020). https://doi.org/10.1186/s12935-020-01539-7
30   Wang, M. Q. *et al.* HKDC1 upregulation promotes glycolysis and disease
progression, and confers chemoresistance onto gastric cancer. *Cancer Sci* **114**,
1365-1377 (2023). https://doi.org/10.1111/cas.15692
31   Boutelle, A. M. *et al.* Integrative multiomic approaches reveal ZMAT3 and p21 as
conserved hubs in the p53 tumor suppression network. *Cell Death Differ* (2025).
https://doi.org/10.1038/s41418-025-01513-8
32   Brayer, K. J. & Segal, D. J. Keep your fingers off my DNA: protein-protein
interactions mediated by C2H2 zinc finger domains. *Cell Biochem Biophys* **50**,
111-131 (2008). https://doi.org/10.1007/s12013-008-9008-5
33   Gosztyla, M. L. *et al.* Integrated multi-omics analysis of zinc-finger proteins
uncovers roles in RNA regulation. *Mol Cell* (2024).
https://doi.org/10.1016/j.molcel.2024.08.010
34   Nabeel-Shah, S. *et al.* C2H2-zinc-finger transcription factors bind RNA and
function in diverse post-transcriptional regulatory processes. *Mol Cell* **84**, 3810-
3825 e3810 (2024). https://doi.org/10.1016/j.molcel.2024.08.037
35   Belgardt, B. F. *et al.* Hypothalamic and pituitary c-Jun N-terminal kinase 1
signaling coordinately regulates glucose metabolism. *Proc Natl Acad Sci U S A*
107, 6028-6033 (2010). https://doi.org/10.1073/pnas.1001796107
36   Xiao, F. *et al.* Hepatic c-Jun regulates glucose metabolism via FGF21 and
modulates body temperature through the neural signals. *Mol Metab* **20**, 138-148
(2019). https://doi.org/10.1016/j.molmet.2018.12.003
37   Scherer, S. J. *et al.* p53 and c-Jun functionally synergize in the regulation of the
DNA repair gene hMSH2 in response to UV. *J Biol Chem* **275**, 37469-37473
(2000). https://doi.org/10.1074/jbc.M006990200
38   Eferl, R. *et al.* Liver tumor development. c-Jun antagonizes the proapoptotic
activity of p53. *Cell* **112**, 181-192 (2003). https://doi.org/10.1016/s0092-
8674(03)00042-4
39   Hellborg, F. *et al.* Human wig-1, a p53 target gene that encodes a growth
inhibitory zinc finger protein. *Oncogene* **20**, 5466-5474 (2001).
https://doi.org/10.1038/sj.onc.1204722
40   Varmeh-Ziaie, S. *et al.* Wig-1, a new p53-induced gene encoding a zinc finger
protein. *Oncogene* **15**, 2699-2704 (1997). https://doi.org/10.1038/sj.onc.1201454
41   Cassandri, M. *et al.* Zinc-finger proteins in health and disease. *Cell Death Discov*
3, 17071 (2017). https://doi.org/10.1038/cddiscovery.2017.71
42   Kamaliyan, Z. & Clarke, T. L. Zinc finger proteins: guardians of genome stability.
Front Cell Dev Biol **12**, 1448789 (2024).
https://doi.org/10.3389/fcell.2024.1448789
43   Orth, B. *et al.* Identification of an atypical interaction site in the BTB domain of the
MYC-interacting zinc-finger protein 1. *Structure* **29**, 1230-1240 e1235 (2021).
https://doi.org/10.1016/j.str.2021.06.005

44   Font, J. & Mackay, J. P. Beyond DNA: zinc finger domains as RNA-binding
modules. *Methods Mol Biol* **649**, 479-491 (2010). https://doi.org/10.1007/978-1-
60761-753-2_29
45   Krishna, S. S., Majumdar, I. & Grishin, N. V. Structural classification of zinc
fingers: survey and summary. *Nucleic Acids Res* **31**, 532-550 (2003).
https://doi.org/10.1093/nar/gkg161
46   Eferl, R. & Wagner, E. F. AP-1: a double-edged sword in tumorigenesis. *Nat Rev*
Cancer **3**, 859-868 (2003). https://doi.org/10.1038/nrc1209
47   Zhou, C. *et al.* JUN is a key transcriptional regulator of the unfolded protein
response in acute myeloid leukemia. *Leukemia* **31**, 1196-1205 (2017).
https://doi.org/10.1038/leu.2016.329
48   Weiss, C. & Bohmann, D. Deregulated repression of c-Jun provides a potential
link to its role in tumorigenesis. *Cell Cycle* **3**, 111-113 (2004).
49   Spangler, B., Vardimon, L., Bosserhoff, A. K. & Kuphal, S. Post-transcriptional
regulation controlled by E-cadherin is important for c-Jun activity in melanoma.
Pigment Cell Melanoma Res **24**, 148-164 (2011). https://doi.org/10.1111/j.1755-
148X.2010.00787.x
50   Lively, T. N., Ferguson, H. A., Galasinski, S. K., Seto, A. G. & Goodrich, J. A. c-
Jun binds the N terminus of human TAF(II)250 to derepress RNA polymerase II
transcription in vitro. *J Biol Chem* **276**, 25582-25588 (2001).
https://doi.org/10.1074/jbc.M100278200
51   Guo, C. *et al.* Coordinated regulatory variation associated with gestational
hyperglycaemia regulates expression of the novel hexokinase HKDC1. *Nat*
Commun **6**, 6069 (2015). https://doi.org/10.1038/ncomms7069
52   Chen, X. *et al.* PGC1beta Regulates Breast Tumor Growth and Metastasis by
SREBP1-Mediated HKDC1 Expression. *Front Oncol* **9**, 290 (2019).
https://doi.org/10.3389/fonc.2019.00290
53   Xu, Z., Zhang, S., Nian, F. & Xu, S. Identification of a glycolysis-related gene
signature associated with clinical outcome for patients with lung squamous cell
carcinoma. *Cancer Med* **10**, 4017-4029 (2021).
https://doi.org/10.1002/cam4.3945
54   Bieging, K. T., Mello, S. S. & Attardi, L. D. Unravelling mechanisms of p53-
mediated tumour suppression. *Nat Rev Cancer* **14**, 359-370 (2014).
https://doi.org/10.1038/nrc3711
55   Vousden, K. H. & Ryan, K. M. p53 and metabolism. *Nat Rev Cancer* **9**, 691-700
(2009). https://doi.org/10.1038/nrc2715
56   Tarangelo, A. *et al.* p53 Suppresses Metabolic Stress-Induced Ferroptosis in
Cancer Cells. *Cell Rep* **22**, 569-575 (2018).
https://doi.org/10.1016/j.celrep.2017.12.077
57   Cairns, R. A., Harris, I. S. & Mak, T. W. Regulation of cancer cell metabolism. *Nat*
Rev Cancer **11**, 85-95 (2011). https://doi.org/10.1038/nrc2981
58   Pavlova, N. N. & Thompson, C. B. The Emerging Hallmarks of Cancer
Metabolism. *Cell Metab* **23**, 27-47 (2016).
https://doi.org/10.1016/j.cmet.2015.12.006

59 Wolpaw, A. J. & Dang, C. V. Exploiting Metabolic Vulnerabilities of Cancer with
Precision and Accuracy. *Trends in cell biology* **28**, 201-212 (2018).
https://doi.org/10.1016/j.tcb.2017.11.006
60 Schwartzenberg-Bar-Yoseph, F., Armoni, M. & Karnieli, E. The tumor suppressor
p53 down-regulates glucose transporters GLUT1 and GLUT4 gene expression.
Cancer Res **64**, 2627-2633 (2004). https://doi.org/10.1158/0008-5472.can-03-
0846
61 Franklin, D. A. *et al.* p53 coordinates DNA repair with nucleotide synthesis by
suppressing PFKFB3 expression and promoting the pentose phosphate pathway.
Sci Rep **6**, 38067 (2016). https://doi.org/10.1038/srep38067
62 Ros, S. *et al.* 6-Phosphofructo-2-kinase/fructose-2,6-biphosphatase 4 is essential
for p53-null cancer cells. *Oncogene* **36**, 3287-3299 (2017).
https://doi.org/10.1038/onc.2016.477
63 Wang, L. *et al.* Hexokinase 2-mediated Warburg effect is required for PTEN- and
p53-deficiency-driven prostate cancer growth. *Cell Rep* **8**, 1461-1474 (2014).
https://doi.org/10.1016/j.celrep.2014.07.053
64 Ravi, R. *et al.* Regulation of tumor angiogenesis by p53-induced degradation of
hypoxia-inducible factor 1alpha. *Genes Dev* **14**, 34-44 (2000).
65 Jiang, P. *et al.* p53 regulates biosynthesis through direct inactivation of glucose-
6-phosphate dehydrogenase. *Nat Cell Biol* **13**, 310-316 (2011).
https://doi.org/10.1038/ncb2172
66 Andrysik, Z. *et al.* Identification of a core TP53 transcriptional program with highly
distributed tumor suppressive activity. *Genome Res* **27**, 1645-1657 (2017).
https://doi.org/10.1101/gr.220533.117

Figure legends

Figure 1. ZMAT3 depletion results in increased expression of genes related to glucose
metabolism in colorectal cancer cells.

(**A**) IGV snapshot shows location of the two sgRNAs used to generate *ZMAT3*-KO HCT116
cells, the observed 57 bp deletion near sgRNA#2 and the p53 ChIP-seq peak in the *ZMAT3* locus
in response to p53 activation upon Nutlin treatment. The p53 ChIP-seq data was previously
published[66]. (**B**) RT-qPCR was performed from 3 biological replicates of *ZMAT3*-WT and
ZMAT3-KO HCT116 cells. *GAPDH* served as a housekeeping gene control. (**C**) Colony
formation assays were performed from 3 biological replicates of *ZMAT3*-WT and *ZMAT3*-KO
HCT116 cells. (**D**) Notched box plot of the $\log_2$FC in RNA abundance of differentially expressed
genes after RNA-Seq analysis of *ZMAT3*-KO versus *ZMAT3*-WT HCT116 cells. Median values
for each group are indicated at the top, and the number of RNAs for which data were obtained
for each group is indicated at the bottom. (**E**) Volcano plots showing differentially expressed
proteins (shown in red) identified by performing quantitative proteomics from *ZMAT3*-WT and

ZMAT3-KO HCT116 cells. (**F**) Most significantly enriched pathways in the gene set enrichment

analysis of significantly upregulated genes (p<0.05) from the *ZMAT3*-KO/*ZMAT3*-WT

quantitative proteomics analysis. (**G**) TMT mass spectrometry peptide abundance of HKDC1

protein in *ZMAT3*-WT and *ZMAT3*-KO HCT116 cells. The values are the average of five

biological replicates for ZMAT3-WT and four biological replicates for *ZMAT3*-KO cells. (**H**)

IGV snapshot for ZMAT3 and HKDC1 from RNA-seq from *ZMAT3*-WT and *ZMAT3*-KO

HCT116 cells. $*p < 0.05$, $****p < 0.0001$.

824 Figure 2. ZMAT3 negatively regulates HKDC1 expression in diverse cell types.

(**A, B**) RT-qPCR and immunoblotting for HKDC1 from *ZMAT3*-WT and *ZMAT3*-KO HCT116

cells. GAPDH served as a housekeeping gene control. RT-qPCR are values are the average of

three biological replicates. (**C, D**) RT-qPCR was performed in biological triplicates for *ZMAT3*

and *HKDC1* mRNAs from HCT116, SW1222, HCEC-1CT and HEPG2 cells after transfection

with a control (CTRL) siRNA or ZMAT3 siRNAs for 72 hr. GAPDH served as a housekeeping

gene control. (**E**) Immunoblotting was performed for endogenous ZMAT3 and HKDC1 from

HCT116 and HepG2 whole cell lysates after siRNA-mediated knockdown of ZMAT3 or HKDC1

for 72 hr. *GAPDH* served as housekeeping gene control. (**F**) Fold change for *Zmat3, Trp53,*

Mdm4 and *Hkdc1* mRNAs is shown from the RNA-seq from *Zmat3* knock-out and wild-type

MEFs. (**G**) Analysis of *HKDC1* mRNA levels from normal colon tissues or CRC patient samples

from the TCGA COAD cohort. N refers to the number of samples in each group. (**H**) Fold

change *for Trp53, Zmat3, Cdkn1a* and *Hkdc1* mRNAs is shown from the RNA-seq from *Trp53*

knock-out and wild-type MEFs. N refers to the number of samples in each group. $*p < 0.05$,

$***p < 0.001$, $****p < 0.0001$

839 Figure 3. ZMAT3 inhibits mitochondrial respiration and proliferation via HKDC1.

(**A**) 2-Deoxyglucose analog of glucose together with luminescence-based enzymatic assay was

used to assess relative glucose uptake in *ZMAT3*-WT and *ZMAT3*-KO HCT116 cells in presence

and absence of HKDC1. For SW122 and HEPG2 cells relative glucose was measured in the

presence and absence of siRNA-mediated knockdown of HKDC1 and ZMAT3 alone or in

combination. (**B, C**) Metabolic flux assays were performed for basal glycolysis rate and basal

mitochondrial respiration rate in ZMAT3 and/or HKDC1 knockdown in HCT116 cells. (**D, E**)

Incucyte live cell proliferation assays and CCK8-based cell proliferation in HCT116 *ZMAT3*-WT

and KO cells in the presence and absence of siRNA-mediated knockdown HKDC1. ∗p < 0.05,

(∗∗) p < 0.01, (∗∗∗) P<0.001. The results are the average of three independent experiments.

Figure 4. p53 negatively regulates HKDC1 expression in a ZMAT3-dependent manner.

(A) IGV snapshots from the RNA-seq data following knockdown of p53 with p53 siRNAs. Data

shows increased *HKDC1* mRNA levels and decreased *ZMAT3* mRNA levels upon p53

knockdown in HCT116 cells. **(B)** Fold change is shown for *p53, p21, ZMAT3*, and *HKDC1*

mRNAs from the RNA-seq performed from siCTRL and sip53 transfected HCT116 cells. **(C, D)**

HCT116 cells were transfected with CTRL siRNA or p53 siRNAs for 48 hr. The levels of

ZMAT3, p53, and *HKDC1* mRNA or protein were measured by RT-qPCR (C) or immunoblotting

from whole cell lysates (D). *GAPDH* was used as housekeeping gene control. **(E)** Fold change

for ZMAT3, p21, HKDC1 and p53 mRNAs is shown from the RNA-seq from HCT116 cells

treated with DMSO or Nutlin for 6 hr. **(F)** Immunoblotting was performed for HKDC1, ZMAT3

and p21 from *ZMAT3*-WT and *ZMAT3*-KO HCT116 cells with or without Nutlin treatment for 24

hr. GAPDH served as the loading control. ns refers to not significant. **(G,H)** Doxycycline

inducible ZMAT3-FLAG-HA HCT116 cells treated with 2ug/ml doxycycline for 48 hr. The

levels of ZMAT3 mRNA or protein induction were measured by RT-qPCR (G) or

immunoblotting from whole cell lysates against HA antibody(H). *GAPDH* was used as

housekeeping gene control. **(I, J)** Doxycycline inducible ZMAT3-FLAG-HA HCT116 cells were

transfected with CTRL siRNA or p53 siRNAs for 48 hr followed by 48hr doxycycline treatment.

The levels of *ZMAT3, p53*, and *HKDC1* mRNA or protein were measured by RT-qPCR (I) or

immunoblotting from whole cell lysates (J). *GAPDH* was used as housekeeping gene control. ∗p

< 0.05, ∗∗p < 0.01, ∗∗∗p < 0.001, ∗∗∗∗p < 0.0001

Figure 5. ZMAT3 inhibits HKDC1 transcription by interacting with the transcription

factor JUN.

(A) Schematic for identification of ZMAT3-FLAG interacting proteins by IP mass spectrometry

from HCT116 cells expressing doxycycline-induced ZMAT3-FLAG-HA. **(B)** Volcano plot

showing significantly enriched proteins (shown in red) identified by ZMAT3-FLAG IP followed

by mass spectrometry, in presence or absence of doxycycline from ZMAT3-FLAG-HA HCT116

cells. JUN was strongly enriched in the ZMAT3-FLAG IPs. **(C**) IGV snapshot showing JUN,

POLR2A, H3K27Ac and H3K3Me3 peaks at the *HKDC1* locus from the ENCODE project

(accession from top to bottom: ENCSR000FAH, ENCSR000EDG, ENCSR000EEK,
ENCSR000EUU, ENCSR661KMA and ENCSR333OPW). JUN binding motif (TGASTCA) is
shown in blue (positive strand) and in red (negative strand). (**D**) Immunoblotting was performed
using anti-FLAG beads and whole cell lysates from no doxy or doxycycline treated ZMAT3-
FLAG-HA HCT116 cells. 10% of total cell lysate was used for input. GAPDH used as loading
control. (**E, F**) *ZMAT3*- WT and KO HCT116 cells were transfected with CTRL siRNA or JUN
siRNAs for 48 hr. The levels of *ZMAT3*, *JUN*, and *HKDC1* mRNAs or the corresponding
proteins were measured by RT-qPCR (E) or immunoblotting from whole cell lysates (F).
GAPDH was used as housekeeping gene control. (**G**) JUN ChIP-qPCR was performed in
biological triplicates from HCT116 *ZMAT3*-WT and KO cells to determine enrichment of JUN at
the *HKDC1* promoter. (H) Luciferase assays were performed in biological triplicates upon JUN
or ZMAT3 knockdown alone or in combination using the *HKDC1* promoter reporter constructs.
∗p < 0.05, ∗∗p < 0.01

Figure 6

Schematic for ZMAT3-mediated regulation of HKDC1 expression and inhibition of
mitochondrial respiration. In *ZMAT3*-WT cells p53 activates *ZMAT3* transcription, resulting in
ZMAT3 protein binding to the transcription factor JUN. This inhibits the ability of JUN to bind
to the *HKDC1* promoter, low *HKDC1* expression leading to controlled mitochondrial respiration
and controlled cell proliferation. In *ZMAT3*-knockout cells, JUN actively binds to the *HKDC1*
promoter and upregulates its expression resulting in increased mitochondrial respiration and
increased cell proliferation.

Figure S1. Extended results for Figure 1.

(**A**) Immunoblotting for ZMAT3, p53, p21 and GAPDH from *ZMAT3*-WT and *ZMAT3*-KO
HCT116 cells with or without Nutlin treatment (24 hr). GAPDH served as the loading control.
(**B**) Incucyte live cell proliferation assays were performed from *ZMAT3*-WT and *ZMAT3*-KO
HCT116 cells. (**C**) Volcano plot for the differentially expressed genes identified by RNA-Seq
from *ZMAT3*-WT and *ZMAT3*-KO HCT116 cells. Significantly expressed genes are indicated in
red (p<0.05). (**D**) Most significantly enriched pathways identified by GSEA from the top 500
significantly upregulated genes (p<0.05) from *ZMAT3*-KO vs *ZMAT3*-WT RNA-seq.

Figure S2. Extended results for Figure 2.

(A) Volcano plot for the differentially expressed genes identified by RNA-Seq performed after transfection of HCT116 cells with siCTRL or siZMAT3 for 72 hr. Significantly expressed genes are indicated in red ($p<0.05$). **(B)** GSEA analysis was performed for the top 500 significantly upregulated genes ($p<0.05$) in ZMAT3 knockdown HCT116 cells identified by RNA-seq. **(C, D)** Venn diagram of showing comparisons of the indicated RNA-Seq data sets. 1,023 significant upregulated (C) and 1,042 downregulated (D) differentially expressed genes were shared between the *ZMAT3*-KO/*ZMAT3*-WT and siZMAT3/siCTRL comparisons. **(E)** Most significantly enriched pathways in the GSEA for the top 500 genes commonly upregulated genes ($p<0.05$) in ZMAT3-KO vs *ZMAT3*-WT and siZMAT3 vs siCtrl comparisons from the RNA-seq data. **(F, G)** ZMAT3 and HKDC1 mRNA levels were determined in CRC patient samples in the TCGA COAD cohort from p53-WT (wild-type) and p53-Mutant CRC patient samples.

Figure S3. Extended results for Figure 3.

Non-mitochondrial oxygen consumption in ZMAT3 and/or HKDC1 knockdown in HCT116 cells. Values are the average of four independents experiments.

Figure S4. Extended results for Figure 4.

(A) Volcano plot shows the differentially expressed genes from the RNA-Seq from sip53 and siCTRL transfected groups in HCT116 cells. Significantly expressed genes are shown in red ($p<0.05$). **(B, C)** Venn diagram and GSEA analysis for the genes commonly upregulated in RNA-Seq upon p53 knockdown and ZMAT-KO HCT116. **(D, E)** Venn diagram and GSEA analysis for the genes commonly downregulated in RNA-Seq upon p53 knockdown and *ZMAT*-KO HCT116.

Figure S5. Extended results for Figure 5.

(A) Schematic of full-length ZMAT3 protein showing three zinc finger motifs. **(B)** Immunoblot from 10% input and FLAG immunoprecipitation from doxycycline inducible ZMAT3-FLAG-HA HCT116 cell lysates treated with/without doxycycline for 48 hr.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

HCT116

Nutlin: — + — +
ZMAT3-WT ZMAT3-KO

ZMAT3
p53
p21
GAPDH

B   **Figure S1**

HCT116

C

RNA-Seq
ZMAT3-KO vs ZMAT3-WT

HKDC1

$p < 0.05$

$-\log_{10}$ (p value)

$\log_2$ FC (KO/WT)

D

GSEA for mRNAs up in KO

KRAS_SIGNALING_UP
ESTROGEN_RESPONSE_LATE
EPITHELIAL_MESENCHYMAL_TRANSITION
ESTROGEN_RESPONSE_EARLY
P53_PATHWAY
GLYCOLYSIS
HYPOXIA
COAGULATION
IL2_STAT5_SIGNALING
MYOGENESIS

$-\log_{10}$ (p value)

A RNA-Seq siZMAT3 vs siCTRL

B GSEA (Up in siZMAT3 vs siCTRL)

C KO vs WT / siZMAT3 vs siCTRL

p<0.05, log FC>0
(Up in ZMAT3-KO and upon ZMAT3 knockdown)

D KO vs WT / siZMAT3 vs siCTRL

p<0.05, log FC<0
(Down in ZMAT3-KO and upon ZMAT3 knockdown)

E GSEA (Up in ZMAT3-KO and upon ZMAT3 knockdown)

F TCGA COAD tumors

G TCGA COAD tumors

Non mitochondrial oxygen consumption

A

B

D

C

E

A

1 → 289 (Amino acids)

B

Input | IP-FLAG

Doxy: - + | - - + +

ZMAT3 FLAG