Alpha power indexes working memory load for durations

Sophie K. Herbst[1], Izem[1] Mangione, Charbel-Raphaël Segerie[2], Richard Höchenberger[2],
Tadeusz Kononowicz[1,3,4], Alexandre Gramfort[2] & Virginie van Wassenhove[1]

[1] Cognitive Neuroimaging Unit, INSERM, CEA, Université Paris-Saclay, NeuroSpin, 91191 Gif/Yvette, France

[2] Inria, CEA, Université Paris-Saclay, Palaiseau, France

[3] Institute of Psychology, The Polish Academy of Sciences, ul. Jaracza 1, 00-378 Warsaw, Poland

[4] Institut NeuroPSI - UMR9197 CNRS Université Paris-Saclay

Keywords: time perception, duration, working memory, alpha oscillations, magnetoencephalography

Corresponding Authors:
Sophie Herbst, ksherbstgmail.com
Virginie van Wassenhove, virginie.van.wassenhove@gmail.com
CEA/DRF/Joliot, NeuroSpin - INSERM Cognitive Neuroimaging Unit,
Bat 145 PC 156, F-91191, Gif-sur-Yvette, France

Author contributions

VvW, SKH, and TWK developed the research concept and study design, to which IM also contributed. Coding was performed by SKH, IM, and TWK. Testing and data acquisition was performed by IM. The analyses were performed by IM, TWK, and SKH (behavior, MEG preprocessing) and finalized by CRS and RH, supervised by AG, and SKH (decoding, source reconstruction), with regular inputs and guidance by VvW. SKH wrote the first draft, and SKH and VvW revised the manuscript. All authors reviewed and approved the final version of the manuscript.

Acknowledgments

This project has received funding from Agence Nationale de la Recherche 'AutoTime' (ANR-16-CE37-0004-04) granted to Virginie van Wassenhove, Agence Nationale de la Recherche meegBIDS.fr (ANR-19-DATA-0023), granted to Alexandre Gramfort (Coordinator), Sophie Herbst, Maximilien Chaumon, and Virginie van Wassenhove (collaborators), and from the European Union's Horizon 2020 research and innovation program under grant agreement No. 101017727, FET Experience granted to Virginie van Wassenhove.

NeuroSpin, NeuroPSI, UNICOG, Cognition & Brain Dynamics, and the Mind Lab are part of the DIM C-BRAINS, funded by the Conseil Régional d'Ile-de-France. We would also like to thank the NeuroSpin MEG platform staff, and in particular Leila Azizi, for their support during data acquisition. We think Yunyun Shen for enriching discussions about the study, both theoretically and empirically.

Declaration of interests

The authors declare no competing interests.

Supplemental information

Figure S1. Reproduction performance for single items.

Table T1: Relationship between oscillatory power and reproduction precision, per region of interest.

Abstract

Timing, that is estimating, comparing, or remembering how long events last, requires the temporary storage of durations. How durations are stored in working memory is unknown, despite the widely held view of memory systems' central role in timing. Here, we investigated the neural signatures of a sequence of durations (n-item sequence) held in working memory. We recorded human participants using magnetoencephalography (MEG) while they performed an n-item delayed reproduction task, which required to encode a sequence of durations, maintain it, and then reproduce it. The number of items in a sequence (one or three) and the duration of the sequence were orthogonalized. Our results show that during working memory maintenance, the number of durations, not the duration of the sequence, affected recall precision and could be decoded from alpha and beta oscillatory activity. Parieto-occipital alpha power showed a direct link with the precision of temporal reproduction. Our results extend earlier behavioral findings suggesting that durations are itemized in working memory and that their number, not their duration, modulates recall precision (Herbst et al., 2025). Crucially, we establish that alpha power reflects a universal signature of working memory load and mediates recall precision, even for abstract information such as duration.

Introduction

Humans and animals can readily discriminate the durations of sensory events as well as the silent intervals that may separate them. Sensory timing tasks typically quantify this ability by asking individuals to compare or to (re)produce time intervals (Grondin, 2010). In most sensory timing tasks, the encoded temporal information is stored until a decision is made: for instance, in the simplest two-alternative-forced choice design, two time intervals are presented in succession, and the first of the two is held in working memory until the second is presented and a comparison can be made (Coull et al., 2008; Harrington et al., 2010; Rao et al., 2001). The encoding of duration in such tasks and the decision stages have been well explored (Díaz et al., 2025; Ofir & Landau, 2022; Paton & Buonomano, 2018). To the contrary, how durations are maintained in working memory is unknown. Classical information-theoretic timing models postulated a working memory component (Buhusi & Meck, 2005; Gibbon, 1977; Gibbon et al., 1984; Treisman, 1963), but its mechanisms and neural dynamics remain largely speculative (Gu et al., 2015; Teki et al., 2017; van Wassenhove, 2016).

Recent behavioral work suggests that the storage of durations in working memory is comparable to that of any items (Herbst et al., 2025; Manohar & Husain, 2016; Teki & Griffiths, 2014), in that recall precision is indicative of working memory load (Bays, 2015; Bays et al., 2024; Ma et al., 2014). At the cortical level, the maintenance of encoded events in working memory has been linked to oscillatory dynamics (Miller et al., 2018), which provide a self-sustained neural code in the absence of sensory stimulation. Working memory function likely relies on different oscillatory regimes, ranging from theta to alpha, beta, and gamma (Axmacher et al., 2010; Jensen & Lisman, 2005; Kopell et al., 2011; Lisman & Idiart, 1995; Palva & Palva, 2011; Roux & Uhlhaas, 2014). A consistent finding is that alpha power (8 – 12 Hz) increases with working memory load (Jensen et al., 2002), but studies assessing working memory for duration do not clearly align with this seminal observation as discussed below (Y. G. Chen et al., 2015; Y. G. Chen & Huang, 2016).

Herein, we thus investigated whether duration storage follows the known principles of working memory. We focused on recall precision as a measure of working memory load, and neural oscillations as its neural signature. We recorded participants' brain activity with magnetoencephalography (MEG) while they performed an n-items delayed reproduction task, in which they were asked to encode, maintain, and reproduce temporal sequences of varying length. Critically, we orthogonalized the number of items (one or three) in a sequence and the duration of the sequence. Using this task, we previously showed that the number of items, but not the duration of the sequence affected recall precision (Herbst et al., 2025). These results provided evidence that durations can be itemized and stored in working memory as mental abstract magnitudes (Bueti & Walsh, 2009; Gallistel, 2011; Gallistel & Gelman, 2000; Walsh, 2003). Accordingly, in the present study, we expected an increase of alpha power with working memory load depending on the number of durations in a sequence, but not on the duration of the sequences itself.

The modulation of alpha power is a well-established marker of working memory load, resulting from the number of items (Jensen et al., 2002; Klimesch et al., 1999; Scheeringa et al., 2009). Jensen et al. (2002) demonstrated a parametric alpha increase over posterior and parietal regions during a visual Sternberg task (2, 4, and 6 items, electroencephalography (EEG)), a finding that has since been replicated in visual (Ede et al., 2018; Kornblith et al., 2016; Tuladhar et al., 2007) and auditory contexts (Kaufman et al., 1992; Krause et al., 1996; Leiberg et al., 2006; Obleser et al., 2012; see also the review by Wilsch & Obleser, 2016). Working memory load is classically operationalized through the number of items that have to be maintained, but similar results have been obtained when modulating the complexity of the stimuli (X. Chen et al., 2023; Krause et al., 1995; Obleser et al., 2012)

The increase in alpha power is thought to reflect inhibition of task-irrelevant information during retention (Bonnefond & Jensen, 2012; see also Klimesch et al., 2007; Schneider et al., 2021), and can be modulated by temporal expectations about the retention interval (F.-W. Chen et al., 2023; Wilsch et al., 2014). Two studies have directly targeted the neural dynamics of working memory maintenance for duration: one study showed that alpha power decreased with longer visual item durations (Y. G. Chen et al., 2015), whereas the second study, using a visual n-back task, reported a decrease in posterior alpha power with increasing working memory load – together with beta power increase over temporal areas (Y. G. Chen & Huang, 2016). The divergent alpha power trends in timing studies suggest the possibility of distinct mechanisms for durations held in working memory. To investigate this further, we employed alpha power as an index of working memory load, aiming to determine which factor, the number of items (n-items) or the duration of the sequence contributes most to working memory load.

We show that the number of items, but not the duration of the sequence, can be decoded from both alpha and beta (15 – 25 Hz) band activities, with distinct cortical sources. Interestingly, the alpha-band power showed a direct link to the precision of temporal reproduction. Thus, our results suggest that durations are itemized in working memory like any other mental events.

A) Trial structure

B) Durations in 1- and 3-item sequences

C) Behavioural results

Figure 1. N-item delayed reproduction task and behavioural results. **A: Trial structure.** Each trial consisted of four phases: First, participants listened to a sequence of pure tones (1 kHz, 50 ms) demarcating empty time intervals of varying duration (encoding phase, red fixation dot). Following a retention interval (orange fixation dot), participants were asked to reproduce the full temporal sequence as precisely as possible (green fixation dot). In the example depicted here, the sequence was composed of three durations. Following their reproduction, participants received feedback about the relative temporal reproduction error for each duration in the sequence. ITI = inter-trial-interval, RP = reproduction. **B: Items and sequences.** Dots represent tones. Horizontal lines and the numbers above indicate the duration of the empty intervals in seconds. A sequence was composed of 1 or 3 items (blue and green, respectively). The durations of sequences were fixed to 1.6 s (short), 2.4 s (medium), or 3.6 s (long), irrespective of the number of items that composed them, such that the number of items and the duration of the full sequence were orthogonalized. **C: Effects of the number of items and the duration of sequences on relative reproduction (relRP) and inverse precision (coefficient of variation, CV).** Left: relRP significantly increased with the number of items and decreased with the duration of the sequence (** = p < 0.01). Right: CVs were not affected by the duration of the sequence but significantly increased with the number of items, indicating a decrease in reproduction precision with higher working memory load.

Results

Behavioral Results

In an n-item-delayed-reproduction task, participants reproduced sequences of either one or three duration-items following a retention interval (see Figure 1A). The sequences varied orthogonally in the number of items (one or three), and the duration of the sequence (1.6 s / short, 2.4 s / medium, 3.6 s / long, Figure 1B). The behavioral results replicate our previous findings (Herbst et al., 2025). Specifically, the relative reproduction (relRP, defined as the ratio between each item's reproduction and duration) was modulated by the number of items and the sequence duration (Table 1, Figure 1C, Supplementary Figure 1). Consistent with context effects in magnitude estimation (Petzschner et al., 2015), durations in sequences containing more items were reproduced as longer compared to durations that occurred alone. Furthermore, short sequences were overproduced, long sequences were underproduced, an effect typically interpreted as a regression to the mean (Lejeune & Wearden, 2009; Petzschner et al., 2015; Vierordt, 1868).

More importantly, the results confirm that the number of items significantly decrease the precision of temporal reproduction (quantified as the coefficient of variation, CV: mean (item reproduction) divided by standard deviation (item reproduction); $F_{(1, 97)} = 119.06$, $p < 0.001$, $BF_{10} > 1000$). Conversely, the duration of the sequence did not significantly affect the precision of temporal reproduction ($F_{(1, 97)} = 0.63$, $p = 0.43$, $BF_{10} = 0.125$). Thus, the number of durations, but not the duration of the sequence itself constitute the load when it comes to storing multiple durations in working memory.

DV	IV	STAT	P	BF
relRP	**n-item**	**$F_{(1, 97)} = 74.71$**	**< 0.001**	**> 1000**
	sequence duration	**$F_{(1, 97)} = 221.38$**	**< 0.001**	**> 1000**
	IA	$F_{(1, 97)} = 0.01$	0.95	0.092
CV	**n-item**	**$F_{(1, 97)} = 119.06$**	**< 0.001**	**> 1000**
	sequence duration	$F_{(1, 97)} = 0.63$	0.43	0.125
	IA	$F_{(1, 97)} = 4.27$	0.04	0.786

Table 1. Effects of the number of items (n-item) and sequence duration on the relative reproduction (relRP) and precision (CV) of temporal reproductions. Bayes Factors (BF) reflect the Bayesian evidence for the significance of the respective predictor. Bold values indicate a significant effect. DV = dependent variable, IV = independent variable, STAT = statistical parameters, P = p-value, IA = interaction.

Decoding results

Next, we applied common spatial pattern decoding (CSP, Grosse-Wentrup & Buss, 2008; Koles, 1991) on the MEG data to test whether oscillatory dynamics during the stimulus-free retention interval contained information about the number of items and/ or the duration of the sequence. Decoding was applied to induced oscillatory power (after subtraction of the evoked response), in time-frequency bins (2 – 25 Hz, 0.5 – 4 s). Significant decoding performance at the group level was observed only for the number of items (Figure 2A, left), but not for the duration of the sequence (Figure 2E). A cluster permutation test revealed a statistically significant cluster spanning all time-frequency bins (Figure 2A, left). The peak frequency distributions (histograms in Figure 2A, middle) were of bimodal shape, with one peak in the alpha and one in the beta band. The source projection of the decoding patterns for the whole frequency and time ranges showed sources in occipital, parietal, and motor areas (Figure 2A, right). To investigate the cortical dynamics underlying the significant decoding performance, we separated the decoding patterns in the alpha and beta bands before projecting the weights in source space (Figure 2B). This revealed parieto-occipital sources in the alpha band, versus a more pronounced spread into sensorimotor areas for the beta band. Both were relatively lateralized to the left hemisphere. Lastly, we reconstructed the sources directly from the

oscillatory power, rather than from the decoding patterns, which allowed assessing the directionality of the effect: for the alpha band, we observed higher power for 3-item versus 1-item sequences, with a bilateral occipital distribution. In the beta band, power was higher for 1-item compared to 3-item sequences in the motor regions, and higher for 3-items in occipital regions (Figure 2C), both bilateral. While the effect of beta power in the occipital regions likely reflects harmonics of alpha, the opposite directionality and the different topographical distributions of the modulation of beta power in the sensorimotor areas suggests that it reflects an independent effect. These findings thus argue for the separability of alpha and beta dynamics during working memory maintenance.

Alpha power mediates the effect of n-items on precision

We addressed the relationship between the precision of reproduction (quantified by the CV, computed for single trials through a jackknifing approach, see Methods) and oscillatory power in the alpha and beta bands, obtained from six source labels covering the significant decoding patterns: lateral occipital cortex, inferior parietal cortex, superior parietal cortex, supramarginal gyrus, precentral and postcentral gyri (all bilateral). Alpha power in bilateral occipital cortices showed a significant relationship with CV ($F_{(1/4901)} = 4.70$, $p = 0.03$; Fig. 2D, Table 2), so that the precision of temporal reproduction (here: CV inverted through the jackknifing approach) *decreases* with stronger alpha power. No significant effects were found in other regions, nor in the beta band (Supplementary Table 1). We then tested for a direct relationship between the behavioral precision effects and the differences in oscillatory dynamics for 3-vs. 1- items through mediation (Baron & Kenny, 1986; Benwell et al., 2018; Muller et al., 2005). Alpha power in the lateral occipital cortex mediated the effect of n-items on precision: (1) The independent variable n-items had a significant effect on the mediator (here: power; $F(1, 4906) = 38.54$, $p < 0.001$). (2) n-items significantly affected the CV ($F(1, 4903) = 2742.05$, $p < 0.001$). (3) The regression weight of the effect of the n-items on precision was reduced when the oscillatory power was added to the equation (-0.0202 vs. -0.0201; partial mediation). The Sobel test (Sobel, 1982) result further confirmed a significant mediation (t = -3.05, $p < 0.001$). In sum, alpha power in the lateral occipital cortex mediated the effect of the number of items on the precision of temporal reproduction, suggesting that increased working memory load with the number of durations, not with the duration of the sequence.

Caption of Figure 3 (next page)**. MEG decoding. A) N-item decoding in sensor space.** The number of items was significantly decoded from induced power in the alpha and beta bands during the retention interval using common spatial patterns (CSP). Left: decoding performance during the retention interval, computed in time-frequency bins. The coloured pixels reflect significant decoding performance (decoding accuracy > 0.5; cluster permutation test), and the grey ones non-significant decoding performance. The marginal means reflect average decoding performance across frequency (left, red), and time points (below, orange). Middle: The red and orange histograms show the frequencies and latencies with the highest decoding performance for each participant. Right: topography and source projection of the decoding patterns, averaged across participants. AU = absolute units. (Caption continued on next page **Figure 2** (continued from previous page). **B) Sensor topographies and source projections of decoding patterns**, separately for the alpha (8 – 12 Hz) and beta (15 – 25 Hz) bands. Decoding patterns in the alpha and beta band overlapped, but were relatively more shifted to occipital/parietal areas for alpha and central areas for beta, and left-lateralized.
(Caption continued below the Figure, next page)

Figure 2. MEG decoding. (Caption for Panels A, B on previous page) **C) Source contrasts for 3- versus 1-item sequences:** power differences projected in source space and thresholded at 80%. While panels A and B show the decoding patterns (red = decoding better than chance), which do not allow to infer the directionality of the effect, the contrast of relative power between 3- and 1-item sequences allows to appreciate the different directions of the effects in the alpha (red = more power for 3 items) and beta bands (blue = more power for 1 item). **D) Brain-behavior correlations.** We tested whether the power modulations were directly related to the precision of temporal reproductions in each trial. A significant relationship between precision (CV_{JN}, computed from a jack-knifing (JN) approach) and alpha power was found in bilateral lateral occipital cortex. **E) Decoding of sequence duration (short vs. long).** Time-frequency matrix of decoding performance with marginal means reflecting average decoding performance across frequency (left, red), and timepoints (below, orange). No significant decoding performance was observed for sequence duration.

Discussion

Sustained representations of durations are required for tasks like duration comparison or reproduction (Van Wassenhove, 2009), but little is known about duration storage in memory. Here, we investigated whether working memory for duration follows similar principles as working memory for sensory items (Bays et al., 2009, 2024; Herbst et al., 2025; Joseph et al., 2016; Ma et al., 2014; Manohar & Husain, 2016). Specifically, we assessed established behavioral and neural markers of working memory maintenance from the sensory domain, namely recall precision and neural oscillatory power during retention, to test whether working memory load for durations is driven by the *number* of durations (n-items) or their duration (i.e. temporal extent) itself. Manipulating n-items and sequence duration orthogonally in a previously published n-item delayed reproduction task (Herbst et al., 2025), we replicated our previous behavioral findings that recall precision decreased with the number of items, but not with sequence duration (Fig. 1).

During working memory maintenance, a decoding approach applied to MEG data revealed that induced oscillatory dynamics (parieto-occipital alpha and sensorimotor beta) reflected the number of durations (3- vs. 1-item), but not the duration of the sequence (Fig. 2). Furthermore, alpha power statistically mediated the effect of n-items on behavioral precision. A key contribution of this work is the demonstration that time intervals are represented as discrete items in working memory, which extends the principles of sensory working memory to abstract temporal information (Bueti & Walsh, 2009; Gallistel & Gelman, 2000; Herbst et al., 2025; Walsh, 2003).

Alpha power indexes working memory load

Maintaining three versus one duration in working memory increased alpha power over parieto-occipital areas. Previous studies have established that maintaining multiple distinct visual or auditory items (e.g. letters, faces, or spoken syllables) in working memory is associated with an increase of alpha power measured over parieto-occipital regions, proportional to working memory load (Jensen et al., 2002; Kaufman et al., 1992; Klimesch et al., 1999; Obleser et al., 2012; Scheeringa et al., 2009; Tuladhar et al., 2007). The increase of alpha activity is typically interpreted as the disengagement of the dorsal visual stream, to shield the currently retained information from interference by distracting inputs (Bonnefond & Jensen, 2012; X. Chen et al., 2023; Cooper et al., 2003; Jensen & Mazaheri, 2010; Klimesch et al., 2007; Zhou et al., 2023). Our observation extends the signature of alpha power to the maintenance of duration information in working memory. The parietal sources are in line with a previous study reporting load-depended activity modulations in inferior parietal cortex (Teki & Griffiths, 2016), and, more generally, the representation of mental magnitudes in parietal cortices (Bueti & Walsh, 2009), and the representation of durations in chronotopic maps in parietal and motor areas (Harvey et al., 2020; Protopapa et al., 2019).

Critically, alpha power in the lateral occipital cortex statistically mediated the effect of the number of items on recall precision, providing a direct link between the behavioral precision effect and alpha power as an index for working memory load. Furthermore, the absence of a similar effect on alpha power with longer sequence duration, and the inability to decode sequence duration during retention, corroborate the behavioral result that working memory precision depends solely on the number of duration items.

The role of beta power during working memory retention

Beta power in sensorimotor cortices during retention *decreased* with the number of durations maintained in working memory, which suggests a clear functional distinction from the neural dynamics observed in alpha power over parieto-occipital areas. Beta oscillations are a known neural correlate of working memory, (Bastos et al., 2018; Lundqvist et al., 2016; Miller et al., 2018; Spitzer &

Haegens, 2017), with several putative functional roles: top-down inhibition (Bastos et al., 2018; Liljefors et al., 2024), with commonalities and overlaps to alpha dynamics, but also more specifically the maintenance of sequence order (Liljefors et al., 2024; Miller et al., 2018; Salazar et al., 2012). Furthermore, beta power modulations have been reported as neural signatures of interval timing (Kononowicz, 2015; Kononowicz et al., 2018; Kulashekhar et al., 2016) and timing precision (modulated by alpha phase, Grabot et al., 2019; Kononowicz et al., 2020), as well as working memory for duration of a single tactile stimulus (Spitzer et al., 2014).

In line with the inhibitory role of beta, our observation of higher beta power for 1-items here might indicate stronger movement inhibition (Fujioka et al., 2012; Jha et al., 2015; Khanna & Carmena, 2017; Zhang et al., 2008), or enhanced preparation for target encoding (Liljefors et al., 2024), as the maintenance of single items binds less resources than for multi-item sequences. Furthermore, increased complexity in the motor demands (as in the reproduction of several items) could decrease beta power contralateral to the effector side in a motor imagery task (Brinkman et al., 2014), in line with the left lateralization observed in the source reconstructed decoding patterns (Fig. 2B).

With respect to the specific relevance of beta dynamics to timing, our findings point towards a sustained representation of duration in sensorimotor areas used for working memory maintenance. Seminal investigations revealed that working memory maintenance relies on persistent firing of distributed neural populations, both, in areas that encode the respective stimuli (i.e. visual or motor cortices), but also parietal and frontal areas (Bays, 2015; Goldman-Rakic, 1995). Critically, the amplitude of the activity specifically related to the memorized items decreases with load (Sprague et al., 2014), as does recall precision. This decrease can be linked to a more distributed representation of multiple items, suggested to be mediated by beta power (Lundqvist et al., 2016, 2024). Here, the successful decoding of the number of durations in working memory from beta band activity points towards a distributed representation of item durations mediated by beta dynamics.

Conclusions

Here, we addressed how sequences of durations are stored in working memory, through behavioral and neural indices of working memory load: recall precision and neural oscillatory dynamics. The number of duration items but not the duration of the sequence had an effect on recall precision, and was decodable from oscillatory power in the alpha and beta bands, pointing towards an abstract representation of 'duration items' in working memory. Alpha power increased with load, bridging between durations and other kinds of items. Beta power dynamics differed from alpha, suggesting a specialized neural code for duration.

Methods

Data and Code Availability

The analysis code and anonymized behavioral and MEG data (BIDS format) are available in a private repository on EBRAINS (for review only). Upon acceptance, the repository will be made public with a link included in the final version of the paper. The study was not formally pre-registered.

Participants

23 participants (14 women, mean age = 26.2 years, SD = 5.2 years) were recruited at Neurospin. The sample size was chosen in accordance to previous behavioral experiments (Herbst et al., 2025). All participants were right-handed, and had normal hearing and vision, and no self-reported history of audiological or neurological disorders. Participants were naive as to the purpose of the study and received monetary compensation for their participation. Prior to the study, all participants signed a written informed consent in accordance with the Ethics Committee on Human Research at Neurospin: CPP n° 100049 (Gif-sur-Yvette, France), and the Declaration of Helsinki (2013). Three participants were excluded, one because of a problem with the triggers, one had incomplete data, and for the third, no anatomical MRI scan was available. This left 20 participants for the analyses.

General Procedure and Task

Participants performed an *n-item delayed temporal reproduction* task (Herbst et al., 2025). They were presented with a sequence of one or three "empty" intervals (Fig. 1A), delimited by short pure tones (*encoding*; for details about the stimuli see below). Empty time intervals were used to prevent a maintenance strategy based on auditory features, as opposed to duration. They had to maintain the sequence in memory (*retention*), and, upon a prompt, reproduce the whole sequence by pressing a button for each tone (*reproduction*). Participants received feedback after each reproduction of a sequence.

Before the main experiment, the researcher explained the task to the participants, and provided them with written instructions. Participants then started a training block composed of eight trials. The training was repeated if a participant failed on more than half of the trials (average relative temporal reproduction error over all items in the sequence below - 0.5 or above + 0.5). Data from the training session were not included in the analyses. The actual experiment consisted of 8 blocks of 36 trials each; each block lasted about 10 minutes. After each block, the MEG recording was stopped to save the data, resulting in a break for the participant. The total MEG session lasted 90 minutes.

Stimuli

The experimental paradigm was adapted from Experiment 2 in Herbst et al. (2025). All stimuli were created digitally using Psychtoolbox (Brainard, 1997) under MATLAB (2017a, The Mathworks), at a sampling rate of 44.1 kHz. Stimuli consisted of a sequence of pure tones (1 kHz, 50 ms duration including 10 ms onset and offset ramping to avoid onset artifacts), which demarcated the time intervals (hereafter referred to as "items").

Three sequence durations were tested (1.6 s, 2.4 s and 3.6 s) and the number of duration items was one or three for each sequence. Two-item sequences as used in the previous behavioral study were removed to obtain more trials per condition in the MEG experiment. The order of items within each n-item sequence was randomized. We orthogonalized n-items and the duration of the sequences, resulting in a fixed sequence duration regardless of n-items, but shorter individual items in sequences with more items (Figure 1B). Each block contained all possible sequence types in random order: for the 3-item sequences, six permutations of the order exist for each of the three sequence

durations (18 trials), and the three different 1-item sequences were each repeated six times throughout the block, to balance the number of 1- and 3-item sequences.

During encoding, a red fixation dot was displayed on the center of the screen. 0.5 s after the last tone in the sequence, an orange fixation dot signaled the onset of the retention period. The retention period lasted for 4.2 s. This relatively long interval was chosen with the aim to assess the neural dynamics of working memory with MEG during a period free from sensory stimuli. Following the retention period, a green fixation dot prompted participants to reproduce the full sequence. To respond, participants pressed one button on a Fiber Optic Response Pad (FORP, Science Plus Group, DE) with the index finger of their right hand. Participants had unlimited time to initiate their reproduction, and no sounds were played when they pressed the button.

To reproduce the sequence, participants pressed the button as many times as there were tones in the sequence (*e.g.* 4 key presses for 3 items demarcated by 4 tones), trying to reproduce each duration as precisely as possible. If the correct number of presses was registered, visual feedback was displayed on the screen 0.5 s after the last button press. Visual feedback was displayed for 1 s, and followed by a variable inter-trial interval sampled from a uniform distribution ranging from 2 s to 3 s. Feedback was given in the form of a visual bar plot, which depicted the relative signed reproduction error for each item in the sequence: (item reproduction – item duration) / item duration. Thus, bars of zero length indicated perfectly accurate reproduction, while bars above and below zero reflected over- and under-reproduction, respectively. Trials in which the participant pressed too early (during retention), or an incorrect number of times were labelled as error trials and the participant was prompted with 'too early' / 'wrong number of clicks' instead of the feedback described above. These trials were removed from the analyses.

MEG recording

Prior to the arrival of the participant, an empty room recording was performed for one minute to assess the noise level of the MEG sensors. Before undergoing the MEG recording, participants were equipped with external electrodes, positioned to record the electro-occulogram (EOG, horizontal and vertical eye movements) and -cardiogram (ECG). The positions of the EEG electrodes, four head-position indicator coils, and three fiducial points (nasion, left and right pre-auricular areas) were digitized using a 3D digitizer (Polhemus, US/Canada) for subsequent co-registration with the individual's anatomical MRI. The MEG recordings took place in a magnetically shielded chamber, where the participant was seated in an armchair under the MEG helmet. The electromagnetic brain activity was recorded using a whole-head Elekta Neuromag Vector View 306 MEG system (Neuromag Elekta LTD, Helsinki) with 102 triple-sensors elements (two orthogonal planar gradiometers, and one magnetometer per sensor location). Participants were instructed to fixate their gaze on a screen positioned in front of them, at about one meter distance. The chamber was dimly lit. Their head position was measured before each recording run (8 in total) using the head-position indicator coils. MEG recordings were sampled online at 1 kHz, high-pass filtered at 0.03 Hz, and low-pass filtered at 330 Hz. A two-minute-long resting state recording (eyes open) was performed after the task, used to compute the noise covariance matrix for source reconstruction.

Anatomical MRI recordings

To improve the spatial resolution of the source reconstruction, individual high-resolution structural Magnetic Resonance Imaging (MRI) recordings were used. These were recorded on another day, using a Siemens 3 T Magnetom Prisma Fit MRI scanner. Parameters of the sequence were: slice thickness: 1 mm, repetition time TR = 2300 ms, echo time TE = 2.98 ms, and flip angle = 9 degrees.

Analyses

Software

The analyses of the behavioral data were conducted using R (version 4.3.3 R Core Team, 2024). The MEG data were analyzed using MNE Python (version 1.8.0, Gramfort et al., 2013, 2014; Larson et al., 2024), transformed to the standardized brain imaging data structure (BIDS, Niso et al., 2018) using MNE BIDS (version 1.6.0, Appelhoff et al., 2019), and analyzed with the MNE BIDS pipeline (version 1.8, https://mne.tools/mne-bids-pipeline/), plus additional custom-written code. FreeSurfer was used for the reconstruction of the MRI surfaces (Dale et al., 1999).

Behavioral data

The reproduced duration per item was measured as the time between the onsets of the two key presses. We removed reproduction outliers defined as reproductions that exceeded by 3 standard deviations the participant's mean reproduction for that duration. Recall precision is commonly used in working memory experiments as a continuous measure of working memory load (Ma et al., 2014). To ensure that our measure of precision was not confounded by established biases in reproducing durations with varying magnitudes, we employed the **coefficient of variation (CV)**, a widely accepted index of precision in timing research (Gibbon et al., 1984). We computed CV for bins of items of the same duration, and from the same n-item sequence for each participant as follows:

$$CV = \frac{SD(item\ reproduction)}{mean(item\ reproduction)}$$

To be in accordance with previous studies, we also computed the **relative reproduction** per item (**relRP**), in the same bins as CV, defined as follows:

$$relRP = \frac{item\ reproduction}{item\ duration}$$

Parallel statistical analyses were performed on **relRP** and **CV**, testing for effects of the number of items (n-items), sequence duration, and their interactions. To assess statistical significance, we computed linear mixed effect models using R's *lme4* package (Bates et al., 2015), with the Satterthwaite approximation of degrees of freedom, implemented in the *lmerTest* package (Kuznetsova et al., 2016). Additionally, to obtain effect sizes and assess null-effects, we computed Bayes factors (BF) for each predictor in the model using the *bayestestR* package (Makowski et al., 2019) via the Bayesian Information Criterion approximation (Wagenmakers, 2007). The alpha level for assuming statistical significance was set to $p < 0.05$. In addition, we use Bayes factors to support an unequivocal interpretation of the results. BF represent the relative posterior probability of observing the data under the alternative (i.e., full) and null (i.e., reduced) model, which is informative about whether the inclusion of a given predictor into the model improves the explained variance. For BF we chose the criterion of > 3 and < 0.3 when comparing to models (Lee & Wagenmakers, 2014; Rouder et al., 2009). BF between 0.3 and 3 reflect inconclusive outcomes. In Table 1, we report F- and p-values, as well as BF.

MEG preprocessing

To remove artifacts from the recorded data, we used a standardized preprocessing routine, implemented in the MNE-BIDS-pipeline, with the following steps: noisy or flat sensors were identified visually and marked for exclusion. Next, environmental artifacts were removed from the raw data, using signal source separation (SSS or Maxwell-filter, Taulu & Kajola, 2005), which also interpolates the bad sensors. Head position coordinates were read from the first of the eight recording runs, and

all other runs were spatially aligned to these coordinates. The data were then filtered with a low-pass filter of 160 Hz, and a notch-filter at 50, 100, and 150 Hz to remove line noise. No additional high-pass filter was applied (0.03 Hz used during recording). Furthermore, we resampled the data to a sampling frequency of 250 Hz to speed up the subsequent computations.

Next, the data were cut into epochs of -5 s to 10 s around the onset of the retention interval, and epochs with peak-to-peak amplitudes exceeding 50000 fT in the magnetometers were rejected. Independent component analysis (ICA) was performed on the remaining epochs, with an additional 1 Hz high-pass filter applied only for that purpose. For the detection of artefacts related to the electro-occulogram (EOG) and -cardiogram (ECG), we used an inbuilt routine in MNE python, which finds a participant's typical EOG and ECG activity recorded with the external electrodes, and returns the ICA components that correlate with these typical events. Thresholds for returning the ICA components were set to: 0.1 for the ECG (cross-trial phase statistics, Dammers et al., 2008), and 4.0 for the EOG (z-score).

After ICA cleaning, we rejected epochs with peak-to-peak amplitudes exceeding 10,000 fT (magnetometers) and 20,000 fT/cm (gradiometers). In the following, we report only the analyses of the 102 magnetometers, to reduce the dimensionality of the data and simplify the interpretation of the results and topographies. On average, we obtained 271.7 clean epochs per participant (SD = 13.45, minimum: 246, maximum: 288).

Anatomical MRI preprocessing and forward model computation

FreeSurfer's "recon-all" function was used to reconstruct the cortical surfaces of each participant's brain. Individual surfaces were mapped to FreesSurfer's 'fsaverage' template brain, for later averaging. Single-layer individual head models were computed using the boundary element method (watershed BEM), from each participant's aMRI. The coregistration of the MEG data with the individual's structural MRI was carried out by manually realigning the digitized fiducial points with the anatomical landmarks of the head surface reconstructed from the MRI (using MNE Python's command line tools). Second, an iterative refinement procedure was used to realign all digitized points with the individuals' scalp. We then computed a surface-based source space with 4098 candidate dipoles per hemisphere, spaced approximately 4.9 mm apart (24 mm^2 per source area; recursively subdivided octahedrons 'oct6'), a minimal distance of 5 mm from the inner skull. The final forward models were derived from this source space, the co-registration matrices, and the boundary element models.

Common spatial pattern decoding

To investigate how duration sequences are represented in working memory and identify the factors contributing to memory load, we focused our MEG data analysis on the retention interval. During this period devoid of sensory inputs, we examined how sequences varying in length and duration modulate *induced oscillatory activity* (Tallon-Baudry et al., 1996), in line with established neural dynamics of working memory (Miller et al., 2018). We employed a decoding approach using Common Spatial Patterns (CSP), implemented as part of the MNE BIDS pipeline. CSP is a multivariate decoding algorithm suited for the analysis of induced oscillatory activity (Grosse-Wentrup & Buss, 2008; Koles, 1991). It identifies spatial filters that maximize the variance of one class of signals (e.g., sequences with a three versus one item), while minimizing the variance of other classes. This allows for effective classification of brain activity patterns associated with varying memory load.

We performed the CSP decoding in linearly spaced time-frequency bins, ranging from 2 – 30 Hz (bin width of 2 Hz, 14 bins) and 0.5 – 4 s following retention onset (bin width of 0.25 s, 14 bins). Specifically, the decoding was run on each trials' covariance matrix computed on single trial data, band-pass filtered in the range of the desired frequency bin, after subtracting the evoked response to

focus on induced activity. We used a 5-fold cross validation and quantified decoding performance as the area under the receiver-operator curve averaged across the folds.

To assess in which frequency bands and at which time points we could significantly decode the number of items or the sequence duration at the group level, we performed cluster-permutation tests on the decoding accuracy values across time-frequency bins, using an initial cluster forming threshold of $p < 0.01$, a final cluster selection threshold of $p < 0.05$, and 5000 permutations.

Source reconstruction

To further investigate which brain areas contribute to the significant decoding accuracy, we reconstructed the sources of the decoding patterns (Westner & King, 2023). Here, we used the decoding weights (recomputed now for all epochs jointly without cross-validation), and transformed them to the corresponding patterns across sensors (group average depicted in Figure 2). Topographical patterns are closer to an interpretation as neurophysiological activity compared to the classifier weights (Haufe et al., 2014). We then computed linearly constrained minimum variance spatial filters (lcmv-beamformers, Van Veen et al., 1997) from the epoched and band-pass filtered data, and a noise covariance matrix from the resting state recording, and projected the decoding patterns to the source level. Finally, we morphed the individual source activity to a template brain for averaging (fsaverage). We performed the source reconstruction for the complete range of frequencies identified as statistically significant, and second, split into the canonical alpha ($8 - 12$ Hz) and beta ($15 - 25$ Hz) frequency bands. Finally, to investigate the directions of the effects, we also projected the frequency-band power difference (covariance matrix computed from band-pass filtered epochs) for 1-item and 3-item sequences to source space using minimum-norm estimates (dSPM, Dale et al., 2000).

Brain-behavior relationships

To further assess whether the neural dynamics identified as correlates of working memory load for durations directly relate to behavioral reproduction precision, we extracted single trial power in the alpha ($8 - 12$ Hz) and beta ($15 - 25$ Hz) bands from the occipital, parietal, and central regions, in which we found significant decoding (labels: lateraloccipital, inferiorparietal, superiorparietal, supramarginal, precentral, postcentral from the 'aparc' atlas (Destrieux et al., 2010)). We created functional labels for each participant per hemisphere, by sub-selecting the 15 % of voxels with the highest activity in each label (across all conditions). Selecting only the most activated voxels balanced the need for robust labels, despite individual differences, with the goal of maximizing the signal-to-noise ratio by excluding inactive voxels. We then computed the average power for the alpha and beta band for each label, by applying the Hilbert transform to bandpass-filtered data from each epoch to obtain complex time series, and squaring the absolute values averaging across voxels and time points ($0.5 - 4$ s) to obtain oscillatory power.

To obtain precision *for each trial*, a measure which is per se defined *across* trials, we applied a jackknifing approach (Benwell et al., 2018): we iteratively removed one trial and computed the CV from the remaining trials. This allows to compute an 'inverse trace' of each trial's contribution to the CV, and use this single trial measure for correlation with single trial oscillatory power. Outlier data points were excluded as z-values < -2.5 and > 2.5 for each individual. Finally, we computed linear mixed effect models for each label, to examine the modulation of CV by oscillatory power and the number of items, as well as their interaction. CV was z-scored across all participants and trials for model stability, and power values were averaged for the left and right hemispheres and z-scored per individual. All models contained a random intercept across participants.

Mediation analyses

We also assessed whether oscillatory power in the retention window *mediates* the effect of n-items on CV (Baron & Kenny, 1986; Benwell et al., 2018; Muller et al., 2005), i.e. assessed whether statistically the neural dynamics play the hypothesized role of reflecting the effect of working memory load caused by the number of items. The presence of statistical mediation demands that the following conditions be fulfilled: (1) The independent variable (here: n-items) has a significant effect on the mediator (here: power; estimated from a separate model testing the effect of n-items as the only fixed effect and participant as random intercept on power). (2) The independent variable significantly affects the dependent variable (here: CV). (3) The regression weight of the effect of the independent variable on the dependent variable is reduced when the mediator is added to the equation. Additionally, we performed the Sobel test[1] (Sobel, 1982) which tests the indirect effect of the independent variable on the dependent variable via the mediator.

---

[1] https://www.danielsoper.com/statcalc/calculator.aspx?id=31

Supplementary Materials

Supplementary Figure 1. Reproduction of single items. A: Reproduction of single items (in seconds), as a function of item duration. Shorter items were over-produced, and longer items under-produced, relative to their objective value. This reflects a regression of the mean effect, typically found in timing studies. **B:** Relative reproduction (reproduced duration divided by item duration), as a function of item duration. **C:** Coefficient of Variation (CV) of single item reproduction. There was no significant difference between the CV of different item durations, indicating that scalar variability was observed in this task.

frequency band	label	IV	STAT	P
ALPHA	**lateral occipital**	**n-items**	**F(1/ 4901) = 2647.33**	**< 0.001**
		power	**F (1/ 4901) = 4.70**	**0.03**
		interaction	F(1/ 4901) = 0.58	0.45
	inferior parietal	**n-items**	**F(1/ 4901) = 2661.36**	**< 0.001**
		power	F(1/ 4901) = 0.01	0.93
		interaction	F(1/ 4901) = 1.99	0.16
	superior parietal	**n-items**	**F(1/ 4901) = 2642.06**	**< 0.001**
		power	F(1/ 4901)= 0.95	0.33
		interaction	F(1/ 4901) = 0.45	0.50
	supramarginal	**n-items**	**F(1/ 4901) = 2626.51**	**< 0.001**
		power	F(1/ 4901) = 1.93	0.17
		interaction	F(1/ 4901) = 1.07	0.30
	precentral	**n-items**	**F(1/ 4901) = 2668.38**	**< 0.001**
		power	F(1/ 4901) = 0.12	0.73
		interaction	F(1/ 4901) = 1.47	0.23
	postcentral	**n-items**	**F(1/ 4901) = 2640.93**	**< 0.001**
		power	F(1/ 4901) = 0.28	0.59
		interaction	F(1/ 4901) = 0.62	0.43
BETA	lateral occipital	**n-items**	**F(4931) = 2651.04**	**< 0.001**
		power	F(4931) = 2.18	0.14
		interaction	F(4931) = 0.14	0.71
	inferior parietal	**n-items**	**F(4931) = 2658.40**	**< 0.001**
		power	F(4931) = 0.52	0.47
		interaction	F(4931) = 0.59	0.44
	superior parietal	**n-items**	**F(4931) = 2648.58**	**< 0.001**
		power	F(4931) = 0.24	0.62
		interaction	F(4931) = 1.08	0.30
	supramarginal	**n-items**	**F(4931) = 2604.62**	**< 0.001**
		power	F(4931) = 0.07	0.80
		interaction	F(4931) = 0.34	0.56
	precentral	**n-items**	**F(4931) = 2635.56**	**< 0.001**
		power	F(4931) = 2.07	0.15

	interaction	$F_{(4931)} = 0.72$	0.40
postcentral	**n-items**	**$F_{(4931)} = 2613.87$**	**< 0.001**
	power	$F_{(4931)} = 0.73$	0.40
	interaction	$F_{(4931)} = 0.05$	0.05

Supplementary Table 1. Linear mixed effect models computed to address the relationship between power (in ROIs derived from the significant decoding results) and behavior (CV), separately for the alpha and beta frequency bands. Bold values indicate significant results.

References

Appelhoff, S., Sanderson, M., Brooks, T. L., Vliet, M. van, Quentin, R., Holdgraf, C., Chaumon, M., Mikulan, E., Tavabi, K., Höchenberger, R., Welke, D., Brunner, C., Rockhill, A. P., Larson, E., Gramfort, A., & Jas, M. (2019). MNE-BIDS: Organizing electrophysiological data into the BIDS format and facilitating their analysis. *Journal of Open Source Software*, *4*(44), 1896. https://doi.org/10.21105/joss.01896

Axmacher, N., Henseler, M. M., Jensen, O., Weinreich, I., Elger, C. E., & Fell, J. (2010). Cross-frequency coupling supports multi-item working memory in the human hippocampus. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(7), 3228–3233. https://doi.org/10.1073/pnas.0911531107

Baron, R. M., & Kenny, D. A. (1986). The Moderator-Mediator Variable Distinction in Social Psychological Research. Conceptual, Strategic, and Statistical Considerations. *Journal of Personality and Social Psychology*, *51*(6), 1173–1182. Scopus. https://doi.org/10.1037/0022-3514.51.6.1173

Bastos, A. M., Loonis, R., Kornblith, S., Lundqvist, M., & Miller, E. K. (2018). Laminar recordings in frontal cortex suggest distinct layers for maintenance and control of working memory. *Proceedings of the National Academy of Sciences*, *115*(5), 1117–1122. https://doi.org/10.1073/pnas.1710323115

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/jss.v067.i01

Bays, P. M. (2015). Spikes not slots: Noise in neural populations limits working memory. *Trends in Cognitive Sciences*, *19*(8), 431–438. https://doi.org/10.1016/j.tics.2015.06.004

Bays, P. M., Catalao, R. F. G., & Husain, M. (2009). The precision of visual working memory is set by allocation of a shared resource. *Journal of Vision*, *9*(10), 7. https://doi.org/10.1167/9.10.7

Bays, P. M., Schneegans, S., Ma, W. J., & Brady, T. F. (2024). Representation and computation in visual working memory. *Nature Human Behaviour*, *8*(6), 1016–1034. https://doi.org/10.1038/s41562-024-01871-2

Benwell, C. S. Y., Keitel, C., Harvey, M., Gross, J., & Thut, G. (2018). Trial-by-trial co-variation of pre-stimulus EEG alpha power and visuospatial bias reflects a mixture of stochastic and deterministic effects. *The European Journal of Neuroscience*, *48*(7), 2566–2584. https://doi.org/10.1111/ejn.13688

Bonnefond, M., & Jensen, O. (2012). Alpha oscillations serve to protect working memory maintenance against anticipated distracters. *Current Biology: CB*, *22*(20), 1969–1974. https://doi.org/10.1016/j.cub.2012.08.029

Brinkman, L., Stolk, A., Dijkerman, H. C., Lange, F. P. de, & Toni, I. (2014). Distinct Roles for Alpha- and Beta-Band Oscillations during Mental Simulation of Goal-Directed Actions. *Journal of Neuroscience*, *34*(44), 14783–14792. https://doi.org/10.1523/JNEUROSCI.2039-14.2014

Bueti, D., & Walsh, V. (2009). The parietal cortex and the representation of time, space, number and other magnitudes. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *364*(1525), 1831–1840. https://doi.org/10.1098/rstb.2009.0028

Buhusi, C. V., & Meck, W. H. (2005). What makes us tick? Functional and neural mechanisms of interval timing. *Nature Reviews Neuroscience*, *6*(10), 755–765.

Chen, F.-W., Li, C.-H., & Kuo, B.-C. (2023). Temporal expectation based on the duration variability modulates alpha oscillations during working memory retention. *NeuroImage*, *265*, 119789. https://doi.org/10.1016/j.neuroimage.2022.119789

Chen, X., Ma, R., Zhang, W., Zeng, G. Q., Wu, Q., Yimiti, A., Xia, X., Cui, J., Liu, Q., Meng, X., Bu, J., Chen, Q., Pan, Y., Yu, N. X., Wang, S., Deng, Z.-D., Sack, A. T., Laughlin, M. M., & Zhang, X. (2023). Alpha oscillatory activity is causally linked to working memory retention. *PLoS Biology*, *21*(2), e3001999. https://doi.org/10.1371/journal.pbio.3001999

Chen, Y. G., Chen, X., Kuang, C. W., & Huang, X. T. (2015). Neural oscillatory correlates of duration maintenance in working memory. *Neuroscience*, *290*, 389–397. https://doi.org/10.1016/j.neuroscience.2015.01.036

Chen, Y. G., & Huang, X. (2016). Modulation of Alpha and Beta Oscillations during an n-back Task with Varying Temporal Memory Load. *Frontiers in Psychology*, *6*. https://doi.org/10.3389/fpsyg.2015.02031

Cooper, N. R., Croft, R. J., Dominey, S. J. J., Burgess, A. P., & Gruzelier, J. H. (2003). Paradox lost? Exploring the role of alpha oscillations during externally vs. internally directed attention and the implications for idling and inhibition hypotheses. *International Journal of Psychophysiology*, *47*(1), 65–74. https://doi.org/10.1016/S0167-8760(02)00107-1

Coull, J. T., Nazarian, B., & Vidal, F. (2008). Timing, Storage, and Comparison of Stimulus Duration Engage Discrete Anatomical Components of a Perceptual Timing Network. *Journal of Cognitive Neuroscience*, *20*(12), 2185–2197. https://doi.org/10.1162/jocn.2008.20153

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis I. Segmentation and surface reconstruction. *Neuroimage*, *9*(2), 179–194. https://doi.org/10.1006/nimg.1998.0395

Dale, A. M., Liu, A. K., Fischl, B. R., Buckner, R. L., Belliveau, J. W., Lewine, J. D., & Halgren, E. (2000). Dynamic statistical parametric mapping: Combining fMRI and MEG for high-resolution imaging of cortical activity. *Neuron*, *26*(1), 55–67. https://doi.org/10.1016/s0896-6273(00)81138-1

Dammers, J., Schiek, M., Boers, F., Silex, C., Zvyagintsev, M., Pietrzyk, U., & Mathiak, K. (2008). Integration of amplitude and phase statistics for complete artifact removal in independent components of neuromagnetic recordings. *IEEE Transactions on Biomedical Engineering*, *55*(10), 2353–2362. https://doi.org/10.1109/TBME.2008.926677

Destrieux, C., Fischl, B., Dale, A., & Halgren, E. (2010). Automatic parcellation of human cortical gyri and sulci using standard anatomical nomenclature. *NeuroImage*, *53*(1), 1–15. https://doi.org/10.1016/j.neuroimage.2010.06.010

Díaz, H., Bayones, L., Alvarez, M., Andrade-Ortega, B., Valero, S., Zainos, A., Romo, R., & Rossi-Pool, R. (2025). Contextual neural dynamics during time perception in the primate ventral premotor cortex. *Proceedings of the National Academy of Sciences*, *122*(6), e2420356122. https://doi.org/10.1073/pnas.2420356122

Ede, F. van, Chekroud, S. R., Stokes, M. G., & Nobre, A. C. (2018). Decoding the influence of anticipatory states on visual perception in the presence of temporal distractors. *Nature Communications*, *9*(1), 1449. https://doi.org/10.1038/s41467-018-03960-z

Fujioka, T., Trainor, L. J., Large, E. W., & Ross, B. (2012). Internalized Timing of Isochronous Sounds Is Represented in Neuromagnetic Beta Oscillations. *The Journal of Neuroscience*, *32*(5), 1791–1802. https://doi.org/10.1523/JNEUROSCI.4107-11.2012

Gallistel, C. R. (2011). Chapter 1—Mental Magnitudes. In S. Dehaene & E. M. Brannon (Eds.), *Space, Time and Number in the Brain* (pp. 3–12). Academic Press. https://doi.org/10.1016/B978-0-12-385948-8.00001-3

Gallistel, C. R., & Gelman, R. (2000). Non-verbal numerical cognition: From reals to integers. *Trends in Cognitive Sciences*, *4*(2), 59–65. https://doi.org/10.1016/S1364-6613(99)01424-2

Gibbon, J. (1977). Scalar expectancy theory and Weber's law in animal timing. *Psychological Review*, *84*(3), 279–325.

Gibbon, J., Church, R. M., & Meck, W. H. (1984). Scalar timing in memory. *Annals of the New York Academy of Sciences*, *423*(1 Timing and Time Perception), 52–77.

Goldman-Rakic, P. S. (1995). Cellular basis of working memory. *Neuron*, *14*(3), 477–485. https://doi.org/10.1016/0896-6273(95)90304-6

Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., Goj, R., Jas, M., Brooks, T., Parkkonen, L., & Hämäläinen, M. (2013). MEG and EEG data analysis with MNE-Python. *Frontiers in Neuroscience*, *7*. https://doi.org/10.3389/fnins.2013.00267

Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., Parkkonen, L., & Hämäläinen, M. S. (2014). MNE software for processing MEG and EEG data. *NeuroImage*, *86*, 446–460. https://doi.org/10.1016/j.neuroimage.2013.10.027

Grondin, S. (2010). Timing and time perception: A review of recent behavioral and neuroscience findings and theoretical directions. *Attention, Perception, & Psychophysics*, *72*(3), 561–582. https://doi.org/10.3758/APP.72.3.561

Grosse-Wentrup, M., & Buss, M. (2008). Multiclass common spatial patterns and information theoretic feature extraction. *IEEE Transactions on Biomedical Engineering*, *55*(8), 1991–2000.

Gu, B.-M., van Rijn, H., & Meck, W. H. (2015). Oscillatory multiplexing of neural population codes for interval timing and working memory. *Neuroscience & Biobehavioral Reviews*, *48*, 160–185. https://doi.org/10.1016/j.neubiorev.2014.10.008

Harrington, D. L., Zimbelman, J. L., Hinton, S. C., & Rao, S. M. (2010). Neural modulation of temporal encoding, maintenance, and decision processes. *Cerebral Cortex*, *20*(6), 1274–1285.

Harvey, B. M., Dumoulin, S. O., Fracasso, A., & Paul, J. M. (2020). A Network of Topographic Maps in Human Association Cortex Hierarchically Transforms Visual Timing-Selective Responses. *Current Biology*, *0*(0). https://doi.org/10.1016/j.cub.2020.01.090

Haufe, S., Meinecke, F., Görgen, K., Dähne, S., Haynes, J.-D., Blankertz, B., & Bießmann, F. (2014). On the interpretation of weight vectors of linear models in multivariate neuroimaging. *NeuroImage*, *87*, 96–110. https://doi.org/10.1016/j.neuroimage.2013.10.067

Herbst, S. K., Mangione, I., Kononowicz, T., Shen, Y., & Wassenhove, V. van. (2025). Abstracting time in memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition, epub ahead of print*. https://doi.org/10.31234/osf.io/478cx

Jensen, O., Gelfand, J., Kounios, J., & Lisman, J. E. (2002). Oscillations in the Alpha Band (9–12 Hz) Increase with Memory Load during Retention in a Short-term Memory Task. *Cerebral Cortex*, *12*(8), 877–882. https://doi.org/10.1093/cercor/12.8.877

Jensen, O., & Lisman, J. E. (2005). Hippocampal sequence-encoding driven by a cortical multi-item working memory buffer. *Trends in Neurosciences*, *28*(2), 67–72. https://doi.org/10.1016/j.tins.2004.12.001

Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha activity: Gating by inhibition. *Frontiers in Human Neuroscience*, *4*, 186. https://doi.org/10.3389/fnhum.2010.00186

Jha, A., Nachev, P., Barnes, G., Husain, M., Brown, P., & Litvak, V. (2015). The Frontal Control of Stopping. *Cerebral Cortex*, *25*(11), 4392–4406. https://doi.org/10.1093/cercor/bhv027

Joseph, S., Teki, S., Kumar, S., Husain, M., & Griffiths, T. D. (2016). Resource allocation models of auditory working memory. *Brain Research*, *1640*(Pt B), 183–192. https://doi.org/10.1016/j.brainres.2016.01.044

Kaufman, L., Curtis, S., Wang, J.-Z., & Williamson, S. J. (1992). Changes in cortical activity when subjects scan memory for tones. *Electroencephalography and Clinical Neurophysiology*, *82*(4), 266–284. https://doi.org/10.1016/0013-4694(92)90107-S

Khanna, P., & Carmena, J. M. (2017). Beta band oscillations in motor cortex reflect neural population signals that delay movement onset. *eLife*, *6*, e24573. https://doi.org/10.7554/eLife.24573

Klimesch, W., Doppelmayr, M., Schwaiger, J., Auinger, P., & Winkler, T. (1999). "Paradoxical" alpha synchronization in a memory task. *Brain Research. Cognitive Brain Research*, *7*(4), 493–501. https://doi.org/10.1016/s0926-6410(98)00056-1

Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition-timing hypothesis. *Brain Research Reviews*, *53*(1), 63–88. https://doi.org/10.1016/j.brainresrev.2006.06.003

Koles, Z. J. (1991). The quantitative extraction and topographic mapping of the abnormal components in the clinical EEG. *Electroencephalography and Clinical Neurophysiology*, *79*(6), 440–447. https://doi.org/10.1016/0013-4694(91)90163-X

Kononowicz, T. W. (2015). Dopamine-dependent oscillations in frontal cortex index "start-gun" signal in interval timing. *Frontiers in Human Neuroscience*, *9*, 331. https://doi.org/10.3389/fnhum.2015.00331

Kononowicz, T. W., Roger, C., & van Wassenhove, V. (2018). Temporal Metacognition as the Decoding of Self-Generated Brain Dynamics. *Cerebral Cortex (New York, N.Y.: 1991)*. https://doi.org/10.1093/cercor/bhy318

Kopell, N., Whittington, M. A., & Kramer, M. A. (2011). Neuronal assembly dynamics in the beta1 frequency range permits short-term memory. *Proceedings of the National Academy of Sciences*, *108*(9), 3779–3784. https://doi.org/10.1073/pnas.1019676108

Kornblith, S., Buschman, T. J., & Miller, E. K. (2016). Stimulus Load and Oscillatory Activity in Higher Cortex. *Cerebral Cortex (New York, N.Y.: 1991)*, *26*(9), 3772–3784. https://doi.org/10.1093/cercor/bhv182

Krause, C. M., Heikki Lang, A., Laine, M., Kuusisto, M., & Pörn, B. (1996). Event-related. EEG desynchronization and synchronization during an auditory memory task. *Electroencephalography and Clinical Neurophysiology*, *98*(4), 319–326. https://doi.org/10.1016/0013-4694(96)00283-0

Krause, C. M., Lang, H., Laine, M., Kuusisto, M., & Pörn, B. (1995). Cortical processing of vowels and tones as measured by event-related desynchronization. *Brain Topography*, *8*(1), 47–56. https://doi.org/10.1007/BF01187669

Kulashekhar, S., Pekkola, J., Palva, J. M., & Palva, S. (2016). The role of cortical beta oscillations in time estimation. *Human Brain Mapping*, *37*(9), 3262–3281. https://doi.org/10.1002/hbm.23239

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2016). *lmerTest: Tests in Linear Mixed Effects Models*. https://CRAN.R-project.org/package=lmerTest

Larson, E., Gramfort, A., Engemann, D. A., Leppakangas, J., Brodbeck, C., Jas, M., Brooks, T. L., Sassenhagen, J., McCloy, D., Luessi, M., King, J.-R., Höchenberger, R., Goj, R., Favelier, G., Brunner, C., van Vliet, M., Wronkiewicz, M., Rockhill, A., Holdgraf, C., … luzpaz. (2024). *MNE-Python* (Version v1.8.0) [Computer software]. Zenodo. https://doi.org/10.5281/zenodo.13340330

Lee, M. D., & Wagenmakers, E.-J. (2014). *Bayesian Cognitive Modeling: A Practical Course*. Cambridge University Press.

Leiberg, S., Lutzenberger, W., & Kaiser, J. (2006). Effects of memory load on cortical oscillatory activity during auditory pattern working memory. *Brain Research*, *1120*(1), 131–140. https://doi.org/10.1016/j.brainres.2006.08.066

Lejeune, H., & Wearden, J. H. (2009). Vierordt's The Experimental Study of the Time Sense (1868) and its legacy. *European Journal of Cognitive Psychology*, *21*(6), 941–960.

Liljefors, J., Almeida, R., Rane, G., Lundström, J. N., Herman, P., & Lundqvist, M. (2024). Distinct functions for beta and alpha bursts in gating of human working memory. *Nature Communications*, *15*(1), 8950. https://doi.org/10.1038/s41467-024-53257-7

Lisman, J. E., & Idiart, M. A. P. (1995). Storage of 7 ± 2 Short-Term Memories in Oscillatory Subcycles. *Science*, *267*(5203), 1512–1515. https://doi.org/10.1126/science.7878473

Lundqvist, M., Miller, E. K., Nordmark, J., Liljefors, J., & Herman, P. (2024). Beta: Bursts of cognition. *Trends in Cognitive Sciences*, *28*(7), 662–676. https://doi.org/10.1016/j.tics.2024.03.010

Lundqvist, M., Rose, J., Herman, P., Brincat, S. L., Buschman, T. J., & Miller, E. K. (2016). Gamma and Beta Bursts Underlie Working Memory. *Neuron*, *90*(1), 152–164. https://doi.org/10.1016/j.neuron.2016.02.028

Ma, W. J., Husain, M., & Bays, P. M. (2014). Changing concepts of working memory. *Nature Neuroscience*, *17*(3), Article 3. https://doi.org/10.1038/nn.3655

Makowski, D., Ben-Shachar, M. S., & Lüdecke, D. (2019). bayestestR: Describing Effects and their Uncertainty, Existence and Significance within the Bayesian Framework. *Journal of Open Source Software*, *4*(40), 1541. https://doi.org/10.21105/joss.01541

Manohar, S. G., & Husain, M. (2016). Working Memory for Sequences of Temporal Durations Reveals a Volatile Single-Item Store. *Frontiers in Psychology*, *7*. https://doi.org/10.3389/fpsyg.2016.01655

Miller, E. K., Lundqvist, M., & Bastos, A. M. (2018). Working Memory 2.0. *Neuron*, *100*(2), 463–475. https://doi.org/10.1016/j.neuron.2018.09.023

Muller, D., Judd, C. M., & Yzerbyt, V. Y. (2005). When moderation is mediated and mediation is moderated. *Journal of Personality and Social Psychology*, *89*(6), 852–863. https://doi.org/10.1037/0022-3514.89.6.852

Niso, G., Gorgolewski, K. J., Bock, E., Brooks, T. L., Flandin, G., Gramfort, A., Henson, R. N., Jas, M., Litvak, V., T. Moreau, J., Oostenveld, R., Schoffelen, J.-M., Tadel, F., Wexler, J., & Baillet, S. (2018). MEG-BIDS, the brain imaging data structure extended to magnetoencephalography. *Scientific Data*, *5*(1), Article 1. https://doi.org/10.1038/sdata.2018.110

Obleser, J., Wöstmann, M., Hellbernd, N., Wilsch, A., & Maess, B. (2012). Adverse Listening Conditions and Memory Load Drive a Common Alpha Oscillatory Network. *Journal of Neuroscience*, *32*(36), 12376–12383. https://doi.org/10.1523/JNEUROSCI.4908-11.2012

Ofir, N., & Landau, A. N. (2022). Neural signatures of evidence accumulation in temporal decisions. *Current Biology*, *32*(18), 4093-4100.e6. https://doi.org/10.1016/j.cub.2022.08.006

Palva, S., & Palva, J. M. (2011). Functional roles of alpha-band phase synchronization in local and large-scale cortical networks. *Frontiers in Psychology*, *2*, 204. https://doi.org/10.3389/fpsyg.2011.00204

Paton, J. J., & Buonomano, D. V. (2018). The Neural Basis of Timing: Distributed Mechanisms for Diverse Functions. *Neuron*, *98*(4), 687–705. https://doi.org/10.1016/j.neuron.2018.03.045

Petzschner, F. H., Glasauer, S., & Stephan, K. E. (2015). A Bayesian perspective on magnitude estimation. *Trends in Cognitive Sciences*, *19*(5), 285–293. https://doi.org/10.1016/j.tics.2015.03.002

Protopapa, F., Hayashi, M. J., Kulashekhar, S., Zwaag, W. van der, Battistella, G., Murray, M. M., Kanai, R., & Bueti, D. (2019). Chronotopic maps in human supplementary motor area. *PLOS Biology*, *17*(3), e3000026. https://doi.org/10.1371/journal.pbio.3000026

R Core Team. (2024). *R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.* https://www.R-project.org

Rao, S. M., Mayer, A. R., & Harrington, D. L. (2001). The evolution of brain activation during temporal processing. *Nat Neurosci*, *4*(3), 317–323. https://doi.org/10.1038/85191

Rouder, J. N., Speckman, P. L., Sun, D., Morey, R. D., & Iverson, G. (2009). Bayesian t tests for accepting and rejecting the null hypothesis. *Psychonomic Bulletin & Review*, *16*(2), 225–237. https://doi.org/10.3758/PBR.16.2.225

Roux, F., & Uhlhaas, P. J. (2014). Working memory and neural oscillations: Alpha–gamma versus theta–gamma codes for distinct WM information? *Trends in Cognitive Sciences*, *18*(1), 16–25. https://doi.org/10.1016/j.tics.2013.10.010

Scheeringa, R., Petersson, K. M., Oostenveld, R., Norris, D. G., Hagoort, P., & Bastiaansen, M. C. M. (2009). Trial-by-trial coupling between EEG and BOLD identifies networks related to alpha and theta EEG power increases during working memory maintenance. *NeuroImage*, *44*(3), 1224–1238. https://doi.org/10.1016/j.neuroimage.2008.08.041

Schneider, D., Herbst, S. K., Klatt, L.-I., & Wöstmann, M. (2021). Target enhancement or distractor suppression? Functionally distinct alpha oscillations form the basis of attention. *The European Journal of Neuroscience*. https://doi.org/10.1111/ejn.15309

Sobel, M. E. (1982). Asymptotic confidence intervals for indirect effects in structural equation models. *Sociological Methodology*. https://www.jstor.org/stable/270723

Spitzer, B., Gloel, M., Schmidt, T. T., & Blankenburg, F. (2014). Working memory coding of analog stimulus properties in the human prefrontal cortex. *Cerebral Cortex (New York, N.Y.: 1991)*, *24*(8), 2229–2236. https://doi.org/10.1093/cercor/bht084

Spitzer, B., & Haegens, S. (2017). Beyond the Status Quo: A Role for Beta Oscillations in Endogenous Content (Re)Activation. *eNeuro*, *4*(4), ENEURO.0170-17.2017. https://doi.org/10.1523/ENEURO.0170-17.2017

Sprague, T. C., Ester, E. F., & Serences, J. T. (2014). Reconstructions of Information in Visual Spatial Working Memory Degrade with Memory Load. *Current Biology*, *24*(18), 2174–2180. https://doi.org/10.1016/j.cub.2014.07.066

Tallon-Baudry, C., Bertrand, O., Delpuech, C., & Pernier, J. (1996). Stimulus specificity of phase-locked and non-phase-locked 40 Hz visual responses in human. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *16*(13), 4240–4249. https://doi.org/10.1523/JNEUROSCI.16-13-04240.1996

Taulu, S., & Kajola, M. (2005). Presentation of electromagnetic multichannel data: The signal space separation method. *Journal of Applied Physics*, *97*(12), 124905. https://doi.org/10.1063/1.1935742

Teki, S., & Griffiths, T. D. (2014). Working memory for time intervals in auditory rhythmic sequences. *Frontiers in Psychology*, *5*. https://doi.org/10.3389/fpsyg.2014.01329

Teki, S., & Griffiths, T. D. (2016). Brain Bases of Working Memory for Time Intervals in Rhythmic Sequences. *Frontiers in Neuroscience*, *10*. https://doi.org/10.3389/fnins.2016.00239

Teki, S., Gu, B.-M., & Meck, W. H. (2017). The Persistence of Memory: How the Brain Encodes Time in Memory. *Current Opinion in Behavioral Sciences*, *17*, 178–185. https://doi.org/10.1016/j.cobeha.2017.09.003

Treisman, M. (1963). Temporal discrimination and the indifference interval: Implications for a model of the "internal clock." *Psychological Monographs: General and Applied*, *77*(13), 1–31. https://doi.org/10.1037/h0093864

Tuladhar, A. M., Huurne, N. ter, Schoffelen, J.-M., Maris, E., Oostenveld, R., & Jensen, O. (2007). Parieto-occipital sources account for the increase in alpha activity with working memory load. *Human Brain Mapping*, *28*(8), 785–792. https://doi.org/10.1002/hbm.20306

Van Veen, B. D., Van Drongelen, W., Yuchtman, M., & Suzuki, A. (1997). Localization of brain electrical activity via linearly constrained minimum variance spatial filtering. *IEEE Transactions on Biomedical Engineering*, *44*(9), 867–880.

Van Wassenhove, V. (2009). Minding time in an amodal representational space. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *364*(1525), 1815–1830. https://doi.org/10.1098/rstb.2009.0023

van Wassenhove, V. (2016). Temporal cognition and neural oscillations. *Current Opinion in Behavioral Sciences*, *8*, 124–130. https://doi.org/10.1016/j.cobeha.2016.02.012

Vierordt, K. (1868). *Der zeitsinn nach versuchen*. H. Laupp.

Wagenmakers, E.-J. (2007). A practical solution to the pervasive problems ofp values. *Psychonomic Bulletin & Review*, *14*(5), 779–804. https://doi.org/10.3758/BF03194105

Walsh, V. (2003). A theory of magnitude: Common cortical metrics of time, space and quantity. *Trends in Cognitive Sciences*, *7*(11), 483–488. https://doi.org/10.1016/j.tics.2003.09.002

Westner, B. U., & King, J.-R. (2023). *The best of two worlds: Decoding and source-reconstructing M/EEG oscillatory activity with a unique model* (p. 2023.03.24.534080). bioRxiv. https://doi.org/10.1101/2023.03.24.534080

Wilsch, A., Henry, M. J., Herrmann, B., Maess, B., & Obleser, J. (2014). Alpha Oscillatory Dynamics Index Temporal Expectation Benefits in Working Memory. *Cerebral Cortex*, bhu004. https://doi.org/10.1093/cercor/bhu004

Wilsch, A., & Obleser, J. (2016). What works in auditory working memory? A neural oscillations perspective. *Brain Research*, *1640, Part B*, 193–207. https://doi.org/10.1016/j.brainres.2015.10.054

Zhang, Y., Chen, Y., Bressler, S. L., & Ding, M. (2008). Response preparation and inhibition: The role of the cortical sensorimotor beta rhythm. *Neuroscience*, *156*(1), 238–246. https://doi.org/10.1016/j.neuroscience.2008.06.061

Zhou, Y. J., Ramchandran, A., & Haegens, S. (2023). Alpha oscillations protect working memory against distracters in a modality-specific way. *NeuroImage*, *278*, 120290. https://doi.org/10.1016/j.neuroimage.2023.120290