Identification and functional characterization of toluene degradation genes in *Acinetobacter* sp. Tol 5

Short title: Toluene degradation genes in *Acinetobacter* sp. Tol 5

Shogo Yoshimoto[1], Maiko Hattori[1], Shori Inoue[1], Sakura Mori[1], Yuki Ohara[2], Katsutoshi Hori[1,*]

[1]Department of Biomolecular Engineering, Graduate School of Engineering, Nagoya University, Chikusa-ku, Nagoya, Aichi 464-8603, Japan.

[2]Friend Microbe Inc. Chikusa-ku, Nagoya, Aichi 464-0858, Japan.

*Corresponding author: Katsutoshi Hori (khori@chembio.nagoya-u.ac.jp)

Phone: +81-52-789-3339

Email: khori@chembio.nagoya-u.ac.jp

Author contributions

Shogo Yoshimoto: Conceptualization, Validation, Writing - Original Draft. **Maiko Hattori**: Investigation. **Shori Inoue**: Investigation. **Sakura Mori**: Investigation. **Yuki Ohara**: Investigation. **Katsutoshi Hori**: Writing - Review & Editing, Supervision, Funding acquisition

Keywords

Toluene, dioxygenase, degradation, aromatic compound, *Acinetobacter*

Abstract

Microbial degradation of aromatic compounds provides sustainable solutions for environmental remediation and bioconversion. *Acinetobacter* sp. Tol 5 is notable for its strong adhesiveness and potential as a biocatalyst for toluene degradation; however, its toluene metabolic pathway has not been fully elucidated. In this study, genomic analysis identified a cluster of genes in Tol 5 highly similar to the well-known *tod* operon of *Pseudomonas putida*, encoding enzymes responsible for toluene metabolism. Phylogenetic analyses indicated that these *tod* genes, unusual among *Acinetobacter* species, were likely acquired through horizontal gene transfer. Transcriptomic analyses revealed that *todF* and *todC1* are co-transcribed, while the adjacent *fadL2* gene, encoding a putative outer membrane transporter corresponding to *P. putida todX*, is independently transcribed. Functional characterization using gene-knockout mutants revealed that TodC1, the large subunit of dioxygenase, is essential for growth on toluene, whereas FadL2 is not essential. Growth experiments further showed that the *todC1* knockout mutant could metabolize benzoate, but not toluene or benzene, confirming that the TOD pathway is the primary route for toluene and benzene degradation in Tol 5. The identification of the functional TOD pathway, which is unique within *Acinetobacter*, provides genetic and biochemical insights for the development of Tol 5 as an efficient immobilized biocatalyst for the bioremediation and bioconversion of aromatic compounds.

Introduction

Toluene consumption has been increasing due to its widespread use in industrial applications, including solvents, coatings, adhesives, and fuel additives (1). As a major component of BTEX compounds (benzene, toluene, ethylbenzene, and xylene), toluene is released into air and water, resulting in environmental pollution and raising concerns about toxicity and persistence (2). Given the growing demand for green and cost-effective remediation methods, microbial biodegradation and bioconversion have emerged as sustainable and energy-efficient treatments (3, 4). Unlike chemical treatments, microbial processes can achieve complete degradation of toluene under mild conditions.

Bacteria have evolved diverse metabolic pathways to degrade toluene (5, 6). The well-known pathways involve monooxygenases, which oxidize the aromatic ring or the methyl side chain of toluene, leading to the formation of cresols or benzyl alcohol, which are subsequently converted into catechol-type intermediates by other downstream enzymes. Another well-characterized pathway is the toluene dioxygenase (TOD) pathway. It employs a multicomponent enzyme system that comprises a reductase (*todA*), a ferredoxin (*todB*), and a dioxygenase (*todC1* and *todC2*) that oxidize the aromatic ring of toluene, along with other enzymes (*todD*, *E*, *F*, *G*, *H*, and *I*) responsible for downstream metabolism (7). This TOD pathway, originally characterized in *Pseudomonas putida* (8), converts toluene to 3-methylcatechol and finally to pyruvate and acetyl-CoA.

While *Acinetobacter* species are not typically recognized as toluene degraders, some strains have been reported to possess toluene-degrading capabilities. For example, *A. junii* isolated from petroleum-contaminated soil degrades 70% of toluene at 0.15 mg/ml within 72 h (9). *A. baylyi*, previously known as *A. calcoaceticus*, completely degrades toluene at 0.3 mg/ml within 14 days (10). *Acinetobacter* sp. Tol 5 isolated from a biofiltration system completely degrades toluene at 0.43 mg/ml within 37 h (11). Among *Acinetobacter*, Tol 5 is unique in terms of its high adhesiveness to various surfaces through its adhesive nanofiber

protein AtaA in addition to its high toluene-degradation ability (12, 13). Due to its

adhesiveness, Tol 5 cells can be rapidly immobilized onto support and efficiently used for

degradation of toluene (14). However, despite the practical potential for biodegradation and

bioconversion of toluene, the specific metabolic pathway and genes involved in toluene

degradation of Tol 5 remain unclear.

In this study, we elucidated the toluene-degradation pathway of Tol 5. Genomic

analysis identified candidate toluene-degrading genes, and gene-knockout mutants verified

their function. These findings reveal how Tol 5 metabolizes toluene and provide new insights

into toluene metabolism in *Acinetobacter*.

Materials and Methods

Strains, plasmids, and culture conditions

The strains and plasmids used in this study are shown in Table S1. *Escherichia coli* and its

transformants were cultured in Luria-Bertani (LB) medium (20066-95; Nacalai Tesque) at

37 °C. Tol 5 and its mutants were cultured in LB medium or basal salt (BS) medium (11) at

28 °C.

Gene knockout

The gene knockout mutants were generated using the cytidine base editing system for

Acinetobacter according to the previous report (15, 16). The DNA dimer fragment encoding

the sequence of the single guide RNA was prepared by mixing oligo DNAs listed in Table S2

in 50 mM NaCl solution and gradually cooling from 95 °C to 18 °C at 0.1 °C/s. This dimer

was introduced into the BsaI site of pBECAb-apr. The constructed plasmid was

electroporated into Tol 5$_{REK}$, a restriction-modification system and *ataA*-deficient strain (17).

After overnight culture of Tol 5$_{REK}$ harboring the plasmid in BS medium to promote mutation,

the cells were spread on a BS agar plate containing 5 % (w/v) sucrose to remove the plasmid.

Mutation of the target gene was confirmed by sequencing the DNA amplified from the

genome by PCR using KOD FX Neo (TOYOBO, Osaka, Japan).

Bacterial cell growth

The cells cultured in BS medium supplemented with 0.2 % sodium lactate were collected by

centrifugation, washed two times with BS medium, and resuspended in BS medium. A 0.25

mL of the suspension was inoculated into 5 mL of BS medium in a glass test tube. The

following carbon sources were separately added into each tube: 2.4 μL of toluene or 48 μL of

10 % sodium lactate. The test tubes were sealed with rubber stoppers and incubated at 28 °C

with shaking at 200 rpm. After an overnight incubation, photographs of the test tubes were

taken to visually assess the bacterial growth.

To make growth curves, the cells cultured in BS medium supplemented with each

carbon source were collected by centrifugation, washed two times with BS medium, and

resuspended in BS medium. A 0.25 mL of the suspension was inoculated into 5 mL of BS

medium in a glass test tube. The following carbon sources were separately added into each

tube: 1.2 μL of toluene, 1.0 μL of benzene, or 16 μL of 10 % sodium benzoate, and the optical

density was monitored every 60 min using OD-monitorC&T system (TAITEC, Saitama,

Japan) during the culture.

Bioinformatics

To obtain amino acid sequences with similarity to the TodC1 (BCX75806.1) and FadL2

(BCX75810.1), PSI-BLAST searches were performed at an E-value cutoff of 1e-30 against

the nr30 or nr90 databases on the MPI Bioinformatics Toolkit (18). The collected sequences

were clustered using CLANS (19).

Results

Toluene degradation genes of Tol 5

BLAST searches were conducted on the genome sequence of Tol 5 (20) using the toluene

degradation genes, *tod,* from *Pseudomonas putida*. The 12 genes with high sequence

similarity to the *tod* genes*,* annotated as *todF, C1, C2, B, A, D, E, G, I, H, S, and T,* were

identified. Figure 1 illustrates the genetic organization of the *tod* genes in Tol 5, highlighting

its similarity to that of *P. putida*. Surrounding these genes, transposon-related sequences such

as a reverse transcriptase, an integrase, and IS elements were observed. When the amino acid

sequences of TodC1, the toluene dioxygenase large subunit responsible for the enzyme

activity (21), were compared between the two species, they showed a high sequence similarity

(identity = 92 %). Other *tod* enzyme genes and the two-component regulatory system TodS

and TodT controlling the expression of the *tod* genes (22) also showed high sequence

similarities (Table S3). In addition, a gene encoding a putative outer membrane transporter

was identified in Tol 5 at the genomic locus corresponding to *todX*, which encodes an outer

membrane toluene transporter in *P. putida*. Although this gene was predicted to belong to the

outer membrane fatty acid transporter (FadL) family as in *P. putida todX* (23), the sequence

similarity with TodX was relatively low (identity = 39 %). Therefore, the gene was designated

fadL2.

To investigate the evolutionary and phylogenetic background of these *tod* genes

identified in Tol 5, a BLAST analysis was performed using the amino acid sequence of TodC1

and FadL2 of Tol 5 as a query. Many hits were found in bacteria belonging to the genus

Pseudomonas, whereas no homologous genes were detected in the genomes of *A. bereziniae*,

the closely related species of Tol 5, or *Acinetobacter baumannii*, the most extensively studied

species in the genus *Acinetobacter*. Within the current NCBI database, *todC1* and *fadL2* were

identified in only three strains of *Acinetobacter*: *A. johnsonii*, *A. pittii*, and *Acinetobacter* sp.

ANC 4635. Although this BLAST analysis highlighted the rarity of the *tod* genes in

Acinetobacter, it was insufficient to elucidate the evolutionary relationships of *todC1* and

fadL2 from Tol 5 with other aromatic-ring-hydroxylating genes. Therefore, we next conducted

a cluster analysis. Amino acid sequences of aromatic ring hydroxylating dioxygenase alpha

subunits were collected from the nr30 database using PSI-BLAST and subsequently classified

based on their sequence similarity using CLANS, resulting in five distinct clusters (Fig. 1C).

While TodC1 of Tol 5 grouped with TodC1 of *P. putida*, biphenyl 2,3-dioxygenase BphA (24),

and 3-phenylpropionate dioxygenase HcaE (25), some dioxygenases for toluene were

included also in a separate cluster. This seems to reflect the classification of aromatic ring-

hydroxylating oxygenases reported in the previous study (26). A similar clustering analysis

was performed for FadL2 of Tol 5 (Fig. 1D). FadL2 of Tol 5 was grouped with transporters

reported to facilitate the uptake of aromatic compounds, such as TbuX, encoded in the toluene

m-monooxygenase operon (27), CymD, encoded in the *p*-cumate-2,3-dioxygenase operon

(28), and XylN, encoded in the toluene and xylenes monooxygenase operon (29). However,

TodX of *P. putida* formed a distinct cluster from these transporters. While the exact genetic

origin of *fadL2* remains unclear, it appears to have been derived from a transporter associated

with an aromatic oxygenase other than *tod*.

Gene expression of the *tod* genes in Tol 5

The promoter region of *todX* in *P. putida* has been studied in detail (30). It was

shown that the *tod* genes are regulated by the three regions with sequences called TodT-boxes.

Comparing this region with the upstream region of *tod* genes of Tol 5, we found that the

transcription initiation site is located approximately 200 bases upstream of *todF* of Tol 5,

showing high sequence similarity to the *todX* promoter region of *P. putida* (30) (Fig. 2A).

Furthermore, the toluene-binding site of TodS predicted in *P. putida* (31) was also conserved

in that of Tol 5 (Fig. S1). These results suggest that the expression of the *tod* genes of Tol 5 is

regulated similarly by the TodST two-component system, as in *P. putida*. In *P. putida*, the

transcription initiation gene is the *todX*, which encodes an outer membrane protein involved

in toluene uptake. However, in Tol 5, the presumed transcription initiation gene is *todF*,

located downstream of *fadL2* which corresponds to *todX*.

178   To investigate whether *todF* in Tol 5 is transcribed as part of an operon with its

neighboring genes, *fadL2* (upstream) and *todC1* (downstream), we performed reverse

transcription polymerase chain reaction (RT-PCR) using primer sets targeting regions within

and between the three genes. All primer sets successfully amplified DNA fragments from Tol

5 genomic DNA (Fig. 2C; Genomic DNA), confirming the accuracy of the primer design.

When using cDNA synthesized from Tol 5 RNA (Fig. 2C; RNA/RTase(+)), we obtained an

amplification product for the region between *todF* and *todC1*, suggesting that these genes

belong to the same operon. In contrast, no amplification was observed for the region between

todF and *fadL2*, indicating that these genes are transcribed separately. Furthermore, no PCR

amplification was detected in control samples without reverse transcriptase (Fig. 2C;

RNA/RTase(-)), demonstrating that the amplification was derived from RNA and not from

contaminated genomic DNA. These results suggest that, unlike *P. putida*, the *tod* operon in

Tol 5 does not start transcription from *fadL2,* corresponding to *todX,* but instead begins from

todF.

Growth abilities of mutant strains on toluene and other carbon sources

To investigate whether the *tod* genes, which are rarely found in other *Acinetobacter* species,

are involved in toluene degradation in Tol 5, we constructed knockout mutants of *todC1* and

fadL2 using the Tol 5_{REK} strain (17). Tol 5_{REK} carries mutations in the restriction-modification

system, making genetic manipulation easier, and also harbors a mutation in AtaA, resulting in

decreased cell adhesion and autoaggregation, thereby improving the accuracy of growth

evaluation through turbidity measurements. The growth abilities of Tol 5_{REK} and its $\Delta todC1$

and $\Delta fadL2$ mutant strains were compared in basal salt (BS) medium (11). When sodium

lactate was added as the sole carbon source to BS medium, all strains grew well. On the other

hand, when toluene was added as the sole carbon source, the Δ*todC1* mutant strain failed to

grow, while both Tol 5$_{REK}$ and the Δ*fadL2* mutant strain grew well (Fig. 3). These results

suggest that *todC1* is involved in toluene degradation, whereas *fadL2* is not essential.

The effect of mutation in *todC1* on the utilization of other carbon sources—benzene,

which is metabolized via the TOD pathway, and sodium benzoate, which is metabolized via

another pathway, the Ben or Xyl pathway (32, 33)—was evaluated by measuring cell growth.

In BS medium containing toluene or benzene as the sole carbon source, the Δ*todC1* mutant

strain showed no growth (Fig. 4). On the other hand, when sodium benzoate was added as the

sole carbon source, both Tol 5$_{REK}$ and the Δ*todC1* mutant strain grew well. These results

indicate that *todC1* plays a crucial role in the degradation of toluene and benzene but is not

involved in the utilization of benzoate and lactate. This also suggests that the TOD pathway is

the primary route for the degradation of both toluene and benzene in Tol 5.

Discussion

In this study, we discovered that Tol 5 possesses a gene cluster for toluene degradation highly

similar to the *tod* operon of *P. putida* (Fig. 1). Genomic analysis revealed that this gene island

encodes homologs of the TOD pathway enzymes (*todF, C1, C2, B, A, D, E, G, I, H*) and the

two-component regulatory system (*todS* and *todT*). Our experimental data showed that the

Δ*todC1* mutant strain lost the ability to grow on toluene and benzene (Fig. 3 and 4),

confirming that the TOD pathway is the primary route for Tol 5 to degrade those aromatic

hydrocarbons. This is the first report of a functional *Pseudomonas*-like TOD pathway

identified in *Acinetobacter*.

Bioinformatic analysis suggests that Tol 5 acquired its *tod* genes through horizontal

gene transfer. Transposon-like sequences identified at the boundary of the *tod* genes (Fig. 1A)

indicate transposition events facilitating their integration. Furthermore, BLAST searches show

that these *tod* genes are uncommon among *Acinetobacter* species, supporting the hypothesis

that the *tod* genes in Tol 5 were horizontally acquired. Indeed, horizontal gene transfer is

recognized as a key mechanism by which bacteria acquire gene clusters involved in the

degradation of organic pollutants (34). Horizontal transfer of such a large metabolic operon

could immediately confer new metabolic capabilities, explaining how Tol 5 came to degrade

toluene effectively.

In *P. putida*, *todX*, encoding an outer membrane protein thought to facilitate toluene

uptake, is co-transcribed as part of the *tod* operon (23). The Tol 5 genome encodes *fadL2*, a

putative outer membrane transporter, in place of *todX*. However, the sequence similarity of

FadL2 with TodX is relatively low (Table S3) and *fadL2* is transcribed independently of the

tod operon (Fig.2). In addition, unlike *todC1*, *fadL2* was not essential for toluene degradation

(Fig. 3 and 4). These findings suggest that Tol 5 may employ an alternative mechanism for

toluene uptake, possibly involving an unidentified transporter or passive diffusion across the

membrane.

The strong adhesion ability of Tol 5, due to AtaA, has been utilized for rapid

immobilization in bioreactors (14, 35). Now, with the identification of the *tod* pathway, Tol 5

could be a promising biocatalyst for transforming toluene into valuable compounds. Further

genetic modifications, such as disruption of *tod* genes, could enable the production of

industrially relevant intermediates (36). The combination of such genetic modifications and

cell immobilization techniques could lead to the development of efficient bioprocesses using

Tol 5 as an immobilized biocatalyst.

In conclusion, our findings provide a detailed genetic and functional characterization

of the *tod* operon in Tol 5, revealing its potential for industrial applications such as

bioremediation and the bioconversion of toluene into high-value chemicals.

Acknowledgements

The authors thank Michio Homma for helpful discussion. The authors also thank Eriko

Kawamoto for her technical assistance. This research was supported by the Graduate Program

of Transformative Chem-Bio Research at Nagoya University supported by MEXT (WISE

Program) to SI, the Japan Society for the Promotion of Science (JSPS) KAKENHI (Grant

Number JP24H00043) to KH and SY, and GteX Program Japan Grant number JPMJGX23B4

to KH.

References

1.   **Holmes, M. D. and Murray, B. P.:** Toluene Toxicity, StatPearls,
https://www.ncbi.nlm.nih.gov/books/NBK599523/ (2024).

2.   **Fedorov, K., Sun, X., and Boczkaj, G.:** Combination of hydrodynamic cavitation and
SR-AOPs for simultaneous degradation of BTEX in water, Chem Eng J, **417**, 128081
(2021).

3.   **El-Naas, M. H., Acio, J. A., and E., T. A. E.:** Aerobic biodegradation of BTEX:
Progresses and prospects, J Environ Chem Eng, **2**, 1104-1122 (2014).

4.   **Yang, C. P., Qian, H., Li, X., Cheng, Y., He, H. J., Zeng, G. M., and Xi, J. Y.:**
Simultaneous removal of multicomponent VOCs in biofilters, Trends Biotechnol, **36**,
673-685 (2018).

5.   **Jindrova, E., Chocová, M., Demnerová, K., and Brenner, V.:** Bacterial aerobic
degradation of benzene, toluene, ethylbenzene and xylene, Folia Microbiol, **47**, 83-93
(2002).

6.   **Parales, R. E., Parales, J. V., Pelletier, D. A., and Ditty, J. L.:** Diversity of microbial
toluene degradation pathways, Adv Appl Microbiol, **64**, 1-73 (2008).

7.   **Yeh, W. K., Gibson, D. T., and Liu, T. N.:** Toluene dioxygenase: A multicomponent
enzyme system, Biochem Bioph Res Co, **78**, 401-410 (1977).

8.   **Zylstra, G. J., Mccombie, W. R., Gibson, D. T., and Finette, B. A.:** Toluene
degradation by *Pseudomonas putida* F1: Genetic organization of the Tod operon, Appl
Environ Microb, **54**, 1498-1503 (1988).

9.   **Singh, P., Singh, V. K., Singh, R., Borthakur, A., Kumar, A., Tiwary, D., and**
Mishra, P. K.: Biological degradation of toluene by indigenous bacteria
Acinetobacter junii CH005 isolated from petroleum contaminated sites in India, Energ
Ecol Environ, **3**, 162-170 (2018).

10. **Arroyo-Caraballo, M. and Colón-Burgos, G.:** Kinetics of aerobic degradation of
toluene by a tropical isolated soil bacterium, pp. 247-257, in: Pepper D. W., Brebbia

C. A., and Zannetti P. (Ed.), Development and Application of Computer Techniques to Environmental Studies VII, WIT Press, Southampton (2000).

11. **Hori, K., Yamashita, S., Ishii, S., Kitagawa, M., Tanji, Y., and Unno, H.:** Isolation, characterization and application to off-gas treatment of toluene-degrading bacteria, J Chem Eng Jpn, **34**, 1120-1126 (2001).

12. **Ishii, S., Yoshimoto, S., and Hori, K.:** Single-cell adhesion force mapping of a highly sticky bacterium in liquid, J Colloid Interf Sci, **606**, 628-634 (2022).

13. **Ishikawa, M., Nakatani, H., and Hori, K.:** AtaA, a new member of the trimeric autotransporter adhesins from *Acinetobacter* sp. Tol 5 mediating high adhesiveness to various abiotic surfaces, PLoS One, **7**, e48830 (2012).

14. **Yoshimoto, S., Ohara, Y., Nakatani, H., and Hori, K.:** Reversible bacterial immobilization based on the salt-dependent adhesion of the bacterionanofiber protein AtaA, Microb Cell Fact, **16**, 123 (2017).

15. **Wang, Y., Wang, Z. P., Chen, Y., Hua, X. T., Yu, Y. S., and Ji, Q. J.:** A highly efficient CRISPR-Cas9-based genome engineering platform in *Acinetobacter baumannii* to understand the H_2O_2-sensing mechanism of OxyR, Cell Chem Biol, **26**, 1732-1742 (2019).

16. **Inoue, S., Yoshimoto, S., and Hori, K.:** A new target of multiple lysine methylation in bacteria, J Bacteriol, **207**, e00325-00324 (2024).

17. **Ishikawa, M. and Hori, K.:** The elimination of two restriction enzyme genes allows for electroporation-based transformation and CRISPR-Cas9-based base editing in the non-competent Gram-negative bacterium *Acinetobacter* sp. Tol 5, Appl Environ Microb, **90**, e0040024 (2024).

18. **Gabler, F., Nam, S. Z., Till, S., Mirdita, M., Steinegger, M., Soding, J., Lupas, A. N., and Alva, V.:** Protein sequence analysis using the MPI Bioinformatics Toolkit, Curr Protoc Bioinform, **72**, e108 (2020).

19. **Frickey, T. and Lupas, A.:** CLANS: a Java application for visualizing protein families based on pairwise similarity, Bioinformatics, **20**, 3702-3704 (2004).

20. **Ishikawa, M. and Hori, K.:** Complete genome sequence of the highly adhesive bacterium *Acinetobacter* sp. strain Tol 5, Microbiol Resour Ann, **10** (2021).

21. **Jiang, H. Y., Parales, R. E., Lynch, N. A., and Gibson, D. T.:** Site-directed mutagenesis of conserved amino acids in the alpha subunit of toluene dioxygenase: Potential mononuclear non-heme iron coordination sites, J Bacteriol, **178**, 3133-3139 (1996).

22. **Shingler, V.:** Integrated regulation in response to aromatic compounds: from signal sensing to attractive behaviour, Environ Microbiol, **5**, 1226-1241 (2003).

23. **Somboon, K., Doble, A., Bulmer, D., Baslé, A., Khalid, S., and van den Berg, B.:** Uptake of monoaromatic hydrocarbons during biodegradation by FadL channel-

mediated lateral diffusion, Nat Commun, **11**, 6331 (2020).

24. **Masai, E., Yamada, A., Healy, J. M., Hatta, T., Kimbara, K., Fukuda, M., and**
Yano, K.: Characterization of biphenyl catabolic genes of gram-positive
polychlorinated biphenyl degrader *Rhodococcus* sp. strain RHA1, Appl Environ
Microb, **61**, 2079-2085 (1995).

25. **Díaz, E., Ferrández, A., and García, J. L.:** Characterization of the *hca* cluster
encoding the dioxygenolytic pathway for initial catabolism of 3-phenylpropionic acid
in *Escherichia coli* K-12, J Bacteriol, **180**, 2915-2923 (1998).

26. **Chakraborty, J., Ghosal, D., Dutta, A., and Dutta, T. K.:** An insight into the origin
and functional evolution of bacterial aromatic ring-hydroxylating oxygenases, J
Biomol Struct Dyn, **30**, 419-436 (2012).

27. **Kahng, H. Y., Byrne, A. M., Olsen, R. H., and Kukor, J. J.:** Characterization and
role of *tbuX* in utilization of toluene by *Ralstonia pickettii* PKO1, J Bacteriol, **182**,
1232-1242 (2000).

28. **Agulló, L., Romero-Silva, M. J., Domenech, M., and Seeger, M.:** *p*-Cymene
promotes Its catabolism through the *p*-cymene and the *p*-cumate pathways, activates a
stress response and reduces the biofilm formation in *Burkholderia xenovorans* LB400,
Plos One, **12** (2017).

29. **Kasai, Y., Inoue, J., and Harayama, S.:** The TOL Plasmid pWW0 *xylN* gene product
from *Pseudomonas putida* Is involved in *m*-xylene uptake, J Bacteriol, **183**, 6662-
6666 (2001).

30. **Lacal, J., Guazzaroni, M. E., Busch, A., Krell, T., and Ramos, J. L.:** Hierarchical
binding of the TodT response regulator to its multiple recognition sites at the *tod*
pathway operon promoter, J Mol Biol, **376**, 325-337 (2008).

31. **Koh, S., Hwang, J., Guchhait, K., Lee, E. G., Kim, S. Y., Kim, S., Lee, S., Chung,**
J. M., Jung, H. S., Lee, S. J., and other 3 authors: Molecular Insights into Toluene
Sensing in the TodS/TodT Signal Transduction System, J Biol Chem, **291**, 8575-8590
(2016).

32. **Gibson, D. T. and Parales, R. E.:** Aromatic hydrocarbon dioxygenases in
environmental biotechnology, Curr Opin Biotech, **11**, 236-243 (2000).

33. **Marques, S. and Ramos, J. L.:** Transcriptional control of the *Pseudomonas putida*
TOL plasmid catabolic pathways, Mol Microbiol, **9**, 923-929 (1993).

34. **Springael, D. and Top, E. M.:** Horizontal gene transfer and microbial adaptation to
xenobiotics: new types of mobile genetic elements and lessons from ecological
studies, Trends Microbiol, **12**, 53-58 (2004).

35. **Hori, K., Ohara, Y., Ishikawa, M., and Nakatani, H.:** Effectiveness of direct
immobilization of bacterial cells onto material surfaces using the bacterionanofiber
protein AtaA, Appl Microbiol Biotechnol, **99**, 5025-5032 (2015).

36.   **Yamashita, S., Sameshima, Y., Konishi, M., Kato, J., Kishimoto, M., Honda, K.,**
365   **Omasa, T., and Ohtake, H.:** Integrated biooxidation and acid dehydration process for
366   monohydroxylation of aromatics, Process Biochem, **42**, 46-51 (2007).
367
368

Figure legends

Fig. 1. Sequence analysis of the *tod* operon. (A) Genetic map of the *tod* operon region of *P.*

putida (top) and *Acinetobacter* sp. Tol 5 (bottom). (B) Reactions catalyzed by the TOD

pathway for toluene degradation. (C,D) Clustering analysis of amino acid sequences similar to

those of TodC1 and FadL2. Large dots highlight TodC1 and FadL2 of Tol 5. (C) Each dot

represents amino acid sequences similar to that of Tol 5 TodC1. (D) Each dot represents

amino acid sequences similar to that of Tol 5 FadL2.

Fig. 2. Comparison of promoter regions of *tod* operons. The upstream sequence of *todX*

from *P. putida* DOT-T1E was compared with that of *todF* from Tol 5. The '+1' position

indicates the transcription initiation site. The integration host factor (IHF) binding site, two

pseudopalindromic sequences (box 1 and box 2), and a half-site (box 3), previously identified

in *P. putida*, are indicated. (B,C) RT-PCR analysis of total RNA isolated from Tol 5 cells.

Primer sets 1F/2R and 5F/6R and 9F/10R were used to examine the transcription of each

ORFs. Primer set 3F/4R was used to examine the cotranscription of *fadL2* and *todF*, whereas

primer set 7F and 8R was used to examine the cotranscription of *todF* and *todC1*.

Fig. 3. Assimilation test. Overnight culture of Tol 5$_{REK}$, $\Delta todC1$ mutant, and $\Delta fadL2$ mutant in BS medium with sodium lactate or toluene. As a control, Overnight culture of Tol 5$_{REK}$ without any carbon source (-) was also shown.

Fig. 4. Growth curves with different carbon sources. Tol 5_{REK} (open circle) and the $\Delta todC1$ mutant strain (closed circle) was inoculated to BS medium containing toluene, benzene, or sodium benzoate.