The plant-time-bender model: predicting yield through wheat's perception of time

Lukas Roth[1], Juan M. Herrera[2], Lilia Levy Häner[2], Didier Pellet[2], Dario Fossati[2], Mike Boss[1], Xiaoran Chen[3], Paraskevi Nousi[3], and Michele Volpi[3]

[1]ETH Zürich, Institute of Agricultural Sciences, Zürich, Switzerland
[2]Agroscope, Nyon, Switzerland
[3]ETH Zürich and EPFL, Swiss Data Science Center, Switzerland

May 1, 2025

Abstract

To address challenges in food security, a better understanding of crop performance under varying environmental conditions is required. Plant Time Bender (PTB) is a deep learning model that integrates high-throughput field phenotyping data with genomic and environmental information to predict wheat yield. PTB leverages image time series, genetic markers, and environmental covariates to learn genotype-specific responses to temperature and vapor pressure deficit. Compared to mere genomic prediction models, PTB demonstrates superior performance when predicting yield in unseen environments across 48 year-locations in Europe. The model captures non-linear growth responses varying with phenological stages and identifies distinct patterns associated with yield performance and stability. Specifically, varieties with higher yield stability exhibit reduced sensitivity to vapor pressure deficit around 1.5 kPa and distinctive temperature responses during emergence and senescence. PTB enables retrospective yield predictions across 20 years, providing a foundation for location-specific variety recommendations and targeted breeding strategies.

1 Introduction

Three major factors influence crop production systems, and thus impact our global food security: (1) the genetics (G) of plants, determined by the choice of variety a breeder selects and farmer grows, (2) the environment (E) that the growing plants experience, and (3) the management practices (M) applied by the farmer. In the breeding and variety testing context we address here, M is generally considered to be optimized beforehand and hence a constant, leaving us with G and E as model inputs to understand the dynamics of such systems. Considering potential interactions between these factors, a common starting point is to define a phenotype P to be the result of genotype effects G with complex environments E (Malosetti et al. 2016),

$$P = G \times E. \tag{1}$$

Describing G with molecular methods is nowadays feasible down to the whole-genome level (Walkowiak et al. 2020). Public weather station data to describe E are becoming increasingly available via governmental institutions that commit themselves to open data (e.g., Deutscher Wetterdienst (DWD) and

MeteoSwiss). In contrast, public variety testing data and other multi-environment trial (MET) data to describing Ps at a set of Es are yet rare.

However, while this lack of data is a restriction, it is not the main cause for the current limitations of predictive models. The key is that G and E are measured at massively different scales than P. While data sets comprising genetic marker data and time series of environmental covariates easily exceed 100k data points (Roth et al. 2024b), MET data barely reach the 10k level (Herrera et al. 2020). Taking the Swiss winter wheat variety testing as an example, the ten locations and 36 examined varieties per year would only add up to 3,600 data points if 10 years of data were used. The result is a typical 'large p, small n' problem, where one tries to model system dynamics with many variables (p) but few samples (n).

Field phenotyping technologies promise to address this problem by time-resolving P in P_t measurements that describe the phenotype P at time t (Roth et al. 2021). Hence, one can define P_t as time-integrated result of genotype effects G with complex and changing environments E_t,

$$P_t = \int_0^t G \times E_\tau \, d\tau \,. \tag{2}$$

Field phenotyping can deliver data sets with 150k data points and more (Roth et al. 2024b), which brings data set sizes of P_t to the level of G and E_t, allowing to find empirical solutions for Equation 2.

The question arises how to model the interactions between G and E. Historically, such $G \times E$ effects are considered to be the product ($\times$) of scalars describing Es (e.g., mean yield performance per E) and scalars describing Gs (e.g., factors of a Finlay-Wilkinson regression) (Fisher and Mackenzie 1923). Yet, it has been shown that genotype effects can be further detailed in small effects of deoxyribonucleic acid (DNA) polymorphism (Li et al. 2021) that can be integrated in whole-genome predictions approaches (Meuwissen, Hayes, and Goddard 2001). Similarly, environment effects can be further detailed using environmental covariates such as temperature and precipitation in whole-genome prediction approaches (Jarquín et al. 2014).

Small effects of G and E interactions are then additively combined to predict the performance of a specific genotype in a specific environment. Hence, a linear relationship between genetic and environment characteristics is implied. Yet, strong evidence point towards the fact that relations between environmental covariates and growth are far from being linear (Roth et al. 2024a). Additionally, these relationships change over time as crop phenology advances (Porter and Gawith 1999). Since crop growth and phenological development are primary drivers of overall crop performance, the assumption of constant linearity appears to be an oversimplification (Wang et al. 2017).

Consequently, it was suggested to combine whole-genome prediction approaches with crop growth models that model non-linear dependencies based on *a priori* knowledge (Diepenbrock et al. 2021). However, such approaches rely heavily on pre-calibrated model parameters (van Voorn et al. 2023). When crop models are used in a breeding context, their genotype-specific parameters are typically reduced to a handful, while the rest remain fixed (Chapman 2008). Moreover, the models may even require additional controlled experiments to further refine predictions to a specific set of environments (Millet et al. 2019). In this work, we hypothesize that the bias of linear models and the out-of-domain issues of pre-calibrated crop growth models can be avoided by directly learning crop growth models on time series of phenotyping data, that is on P_t data instead of P data.

Attempting to integrate P_t values in models raises the question of how to represent growth. While phenotyping data are usually multidimensional, e.g., imaging data or 3-D point clouds, common modelling approaches integrate growth as a scalar, e.g., a growth rate (Roth, Piepho, and Hund 2022; Roth et al. 2024a). Hence, a common step is to first apply feature extraction and dimensionality reduction techniques to multi-dimensional phenotyping data (Hund et al. 2019). Often, such approaches follow a domain-specific purpose, i.e., they try to approximate a physical trait such as for example number of plant organs, canopy structure, leaf area, and others. However, choosing which traits to track over time may create a strong bias towards identifying only well-established relationships.

To address these issues, we propose a novel end-to-end deep learning model called plant time bender (PTB) that is trained simultaneously on high-resolution red, green, and blue (RGB) images of hundreds

Figure 1: Plant time bender (PTB) model with P-state encoder, G×E-state encoder, and target trait decoder. Input to the P-state encoder are image time series, and to the G×E-state encoder environmental covariates (temperature, vapour pressure deficit (VPD)) and genetic marker data as single-nucleotide polymorphisms (SNPs).

of wheat genotypes grown in the field phenotyping facility 'FIP' of ETH Zurich, and on genetic marker data and environmental covariates (Figure 1). The model inherently learns (1) a representation of growth from images (Figure 1, Loss 1 and P-state), and (2) how to represent interactions between genetic marker and environmental covariates that explain this growth (Figure 1, Loss 2/3 and G×E-state).

The model is built around the philosophy of 'bending' time with environmental covariates in a way that linearizes genotype-specific growth (Roth et al. 2024a), and subsequently using this learned latent representation of growth to predict yield. Based on unseen data from 48 year-locations in France, Germany and Switzerland, we validate the model and test its ability to predict unseen environments, unseen genotypes, and unseen genotypes in unseen environments. Furthermore, we evaluate the model on 20 years of historic weather data and yield measurements on a country scale. Finally, we use the trained model to extract optimized phenotypes, so-called ideotypes, that promise higher and more stable wheat yields.

2 Results

2.1 Model performance

Common yield prediction models either utilize environmental covariates alongside genomic marker data that characterizes the genetic set-up of crop varieties (genomic prediction) or phenomic data that describes crop phenotypes (phenomic prediction). The proposed PTB model offers a hybrid approach that can be trained on both genomic and phenomic data, yet can generate predictions using genomic data alone. Hence, we trained the model on both phenotyping data (FIP 1.0 data set, MET data set) and genomic data (SNP data set) in combination with environmental data. The model was then tested on previously unseen MET data using only environmental and genomic data as inputs. This testing approach most closely resembles a genomic prediction framework and was therefore compared to a common genomic prediction approach.

Random regression models represent an established approach in genomic prediction that estimates genetic effects as continuous functions across environmental gradients (Jarquin et al. 2018). Such models can handle correlated environmental covariates through appropriate covariance structures (Tolhurst et al. 2022). We benchmarked our novel deep learning architecture against this conventional method.

The genomic prediction model performed well when tested on unseen genotypes, with a MSE of less than 1 t/ha (Figure 2C), corresponding to a MAPE of less than 10 % (Table 1). However, error sizes increased substantially when the test environment was unseen, with a MAPE of approximately 30 % for the GABI-MET data set and around 20 % for the FIP 1.0 and CH-MET data set. Correlations within environments for the MET data sets were weak to moderate (0.35–0.53), for the FIP 1.0 data set very weak to strong (0.15–0.69).

The suggested PTB model consistently outperformed the random regression baseline for the 48 year-locations of the MET data sets. Its performance was superior for both test scenarios, (1) unseen environments (test (E)) and (2) unseen genotypes in unseen environments (Figure 2C, test (G and E)). While the baseline struggled with the larger CH-MET dataset with 40 year-locations, exhibiting significantly higher mean squared errors (MSEs) compared to the smaller GABI-MET dataset with 8 year-locations, the PTB model effectively reduced MSEs for both data sets to similarly low levels.

The MSE can be influenced by different factors whose relative importance may vary among models. Gauch et al.'s MSE decomposition method provides a valuable framework for evaluating model performance by separating the mean squared error into three distinct components: squared bias (systematic error), non-unity slope (proportional deviation), and lack of correlation (random error) (Gauch, Hwang, and Fick 2003). Although the PTB model tended to increase the lack of correlation on test sets compared to the baseline, this drawback was outweighed by a reduced non-unity of slope and lower squared bias, leading to overall improved performance (Figure 2 and Extended Data Figure 5–7).

While the PTB model was superior for the MET data sets, it underperformed for the FIP 1.0 dataset. Higher bias and a slight increase in the lack of correlation contributed to an overall higher MSE than for the baseline across both test scenarios. Pearson's correlation coefficients and mean absolute percentage errors (MAPEs) confirmed the baseline's advantage, though the differences were relatively small (Table 1). For instance, in the model using the extended SNP marker set, correlation differences were less than 0.09. Notably, differences between the two SNP marker sets were negligible.

When evaluating within-environment correlations and MAPE for the MET data sets, the baseline demonstrated slightly better correlations than the PTB model. However, this gap narrowed when models were trained on combined FIP 1.0 and MET data sets. For the MAPE, the PTB model even surpassed the baseline when trained on combined data sets. When evaluation MAPE and correlations across environments, the PTB model was clearly superior to the baseline, provided again it was trained on combined data sets.

Figure 2: Training/test scenarios (A, B), test performance (C) for the three data sets field phenotyping platform 1.0 (FIP 1.0), GABI multi-environment trial (GABI-MET), and Swiss variety testing multi-environment trial (CH-MET), and 20-years predictions (D, E) of the proposed plant time bender (PTB) model. Test performance for the PTB model and baseline model (random regression) is reported as mean squared error (MSE) decomposed in its three model-prediction related components squared bias, non-unity of slope, and lack of correlation. 20-year predictions show average variety performance and standard deviation (sd) (D) and recommended varieties per location (E).

Table 1: Performance of suggested plant time bender (PTB) model and of baseline (random regression to environmental covariates) on independent test sets. Provided are Pearson's r (Cor) and mean absolute percentage error (MAPE) within environments (Within E) and across environments (Across E) for seen genotypes in unseen environments (E), unseen genotypes in unseen environments (G and E), and unseen genotypes in seen environments (G). Models were trained on either the field phenotyping platform 1.0 (FIP 1.0) data set train split, the full FIP 1.0 data set, or the FIP 1.0 data set plus one of the multi-environment trial (MET) data set, GABI multi-environment trial (GABI-MET) or Swiss variety testing multi-environment trial (CH-MET).

Context	Trained on	Model	Pearson's r (correlation)			MAPE		
FIP 1.0 test sets, GABI SNP set			(E)	(G and E)	(G)	(E)	(G and E)	(G)
Within E	FIP (train)	Baseline	**0.69**	**0.15**	**0.45**	**24.5**	**19.4**	**8.2**
		PTB	0.32	-0.08	0.35	31.5	24.8	9.4
Across E	FIP (train)	Baseline	-	-	**0.80**	-	-	-
		PTB	-	-	0.68	-	-	-
FIP 1.0 test sets, extended SNP set			(E)	(G and E)	(G)	(E)	(G and E)	(G)
Within E	FIP (train)	Baseline	**0.68**	**0.23**	**0.48**	**22.9**	**19.7**	**7.7**
		PTB	0.59	0.17	0.44	29.1	27.2	8.1
Across E	FIP (train)	Baseline	-	-	**0.80**	-	-	-
		PTB	-	-	0.71	-	-	-
GABI-MET test sets, extended SNP set			(E)	(G and E)	(G)	(E)	(G and E)	(G)
Within E	FIP (all)	Baseline	**0.35**	**0.37**	-	28.4	29.2	-
		PTB	0.15	0.01	-	**20.3**	**22.1**	-
	FIP & CH-MET	PTB	0.23	0.13	-	26.1	27.7	-
Within E (2010)	FIP & GABI-MET	Baseline	**0.53**	**0.53**	-	10.9	10.9	-
		PTB	0.41	0.25	-	**5.9**	**7.0**	-
Across E	FIP (all)	Baseline	**0.20**	**0.24**	-	28.4	29.2	-
		PTB	0.10	0.02	-	**20.3**	**22.1**	-
	FIP & CH-MET	PTB	0.08	-0.03	-	26.1	27.7	-
Across E (2010)	FIP & GABI-MET	Baseline	0.27	0.30	-	10.9	10.9	-
		PTB	**0.85**	**0.80**	-	**5.9**	**7.0**	-
CH-MET test sets, extended SNP set			(E / G and E)		(G)	(E / G and E)		(G)
Within E	FIP (all)	Baseline	**0.51**		-	**19.9**		-
		PTB	0.36		-	21.5		-
	FIP & GABI-MET	PTB	0.33		-	27.7		-
Within E (2013)	FIP & CH-MET	Baseline	**0.50**		-	43.7		-
		PTB	0.45		-	**11.2**		-
Across E	FIP (all)	Baseline	0.14		-	**20.5**		-
		PTB	**0.21**		-	23.8		-
	FIP & GABI-MET	PTB	0.09		-	27.8		-
Across E (2013)	FIP & CH-MET	Baseline	0.16		-	43.7		-
		PTB	**0.87**		-	**11.2**		-

2.2 20 years prediction and virtual variety testing recommendations

The architecture of the PTB model allowed to treat images as optional input. Hence, we could predict cultivar performance based on genetic marker data and historical weather data for a 20-year period across 10 locations in Switzerland. The PTB model demonstrated moderate alignment of average per-year yield predictions and measured yield reported by the Swiss interbranch organization Swiss granum (Figure 2D). However, certain years deviated from this trend: in 2016, the model predicted moderate yields despite exceptionally low observed values. The year 2016 was characterized by low-light conditions in Spring and an in general very wet season with low yields in all Europe (Le Gouis, Oury, and Charmet 2020), resulting in significant *Septoria tritici* pressure. Diseases are factors that the proposed PTB model can only incorporate indirectly via environmental covariates, which may lead to overestimations in situations with high disease pressure such as in 2016. In contrast, in 2011, 2014, and 2015, measured yields were notably higher than the model's moderate predictions, for which there is no obvious cause known to us.

PTB model predictions were not only year-specific, but also location and genotype specific. Hence, the model allowed to virtually test varieties at unseen locations and years using historical weather data. The result of such a 'virtual variety testing' closely matched the official list of recommended varieties in 2018 published by Agroscope (Courvoisier et al. 2017) across all locations (Figure 2E). However, a closer examination at the individual location level revealed notable shifts in genotype rankings. For instance, while the variety 'SIALA' (a sister-line of CH CLARO) ranked among the top three on both the official list and the PTB model's overall recommendation, its ranking varied significantly by location. For the location 'Delley', 'SIALA' was entirely outperformed by predictions for the variety 'ARNOLD', highlighting the potential for location-specific variety recommendations. However, the variety 'ARNOLD' also serves as an instructive example of how trial data may not always fully represent field-scale performance: Despite its initial promise on the recommended variety list in 2018, the variety was later removed due to its susceptibility to lodging under farming conditions.

2.3 P-state and G×E-state predictions

The PTB model extracts growth estimates from images (P-state) and from genotype-environment interactions based on temperature, VPD, and genetic marker data (G×E-state). Hence, the trained PTB can visualize those states and responses at inference, enabling an analysis and interpretation of learned genotype-characteristics.

The P-state encoder is based on images only and hence genotype-agnostic, allowing to examine the genotype-specificity of the extracted P-state with a heritability calculation. Heritability is a common measure in plant breeding to indicate the extent to which the variation in a phenotypic trait has a genetic origin, with values close to one indicating a high dominance of the phenotypic signal (Falconer and Mackay 1996). The P-states exhibited varying heritability over time, with peaks coinciding with phenological stages of emergence, tillering, heading, and senescence (Figure 3A). P-state values demonstrated wide ranges throughout the season. Similarly, G×E-states varied, revealing clear genotype-specific differences (Figure 3B).

The learned temperature responses indicated non-linear responses to temperature with low growth rates at temperatures around 0 °C, pronounced maximum growth rates at an optimum temperature around 20 °C, and, depending on the growth phase, decreasing growth responses with supraoptimal temperatures above 20 °C (Figure 3D). This response pattern changed throughout the season, with a tendency to higher optimum temperatures later in the season. In Winter (around 120 days after sowing (DAS)), an inversed response pattern emerged with higher responses at negative temperatures and lower responses at positive temperatures. Stress responses to VPD were characterised by an optimum VPD around 1.5 kPa and increased stress responses at lower and higher VPD values.

The fact that the GABI-MET data was unseen in training allowed to use this MET data to investigate genotype yield and genotype yield sensitivity using a Finlay-Wilkinson regression. Clustering of the learned genotype-specific responses to temperature and VPD resulted in groups that distinctly separated in this yield – yield sensitivity space (Figure 3C). Three clusters were visually sampled at the extremes

Figure 3: Predicted P-state and heritability for all varieties (A) and G×E-state of four selected varieties (B), and clustering of learned genotype responses to temperature and vapour pressure deficit (VPD) in relation to their yield and yield sensitivity (C) for three exemplary clusters (D).

of this space, one with high yield and low sensitivity, one with high yield and high sensitivity, and one with both low yield and low sensitivity. The clusters displayed distinguished response patterns to temperature and VPD (Figure 3D).

Genotypes in the high yield/low yield sensitivity group—the cluster that breeding is supposedly most interested in—were associated with reduced stress response to VPD around 1.5 kPa. Both genotype groups with high yield (insensitive and sensitive ones) were characterized by pronounced temperature responsiveness at emergence (DAS 21) and in Spring (DAS 183–222). Yet, during senescence, temperature responses of the high-yielding sensitive genotype cluster was more pronounces for temperatures below 20 °C and less pronounces above 20 °C than for the insensitive cluster.

3 Discussion

High-throughput field phenotyping platforms and multi-environment trials (METs) represent distinct data sources with complementary strengths: phenotyping platforms provide high temporal resolution (Kirchgessner et al. 2017), while METs offer high spatial resolution across geographical locations. METs can sample crucial combinations of climatic and soil conditions, but typically face a large p, small n problem. Field phenotyping platforms can deliver large data sets, but struggles with out-of-domain issues. Merging these two data sources has the potential to offset each other's limitations, thereby improving predictions for unseen environments.

Yet, integrating multi-modal data such as images, genetic marker data, and environmental covariates

is non-trivial. Our proposed PTB model achieves this integration by learning specific data encoders, with results demonstrating superior performance for unseen MET environments if compare to a state-of-the-art baseline.

Limited interpretability is one of the strongest limitations of most deep learning models. In contrast to this prevailing trend, the proposed PTB model allows visualization of growth responses related to temperature and VPD stress, hence providing insights in learned plant physiology mechanistics. The observed patterns aligned well with existing domain knowledge. Increasing stress due to VPD was identified starting from 2 kPa, and temperature responses showed seasonal variations in cardinal temperatures, which strongly agrees with previous research (Schoppach and Sadok 2013; Porter and Gawith 1999). The observed "inverted" temperature response in winter can be interpreted as vernalization requirements (Slafer 1996).

It was often assumed that evolutionary processes have limited variations in response patterns within species (Parent and Tardieu 2012). Yet, the PTB model identified clear genotype-specific deviations from the well-known general response patterns. Those patterns were not only related to individual genotypes, but also to clusters of high-yielding genotypes with low respectively high sensitivity to environment differences. For a given target population of environments (TPE) where one can determine yield sensitivity beforehand, the model allowed providing new insights into optimized ideotypes. For the investigated TPE in this work (Switzerland), a selection for more responsive temperature reactions at emergence and during the senescence phase may lead to higher yield. Yield stability might be enhanced by selecting for higher resistance to VPD around 1.5 kPa.

In addition to assisting in ideotype formulation, a deep learning model such as the PTB may also directly support selection decisions in breeding. The model's ability to genomically predict yield performance could enhance breeders' decisions when selecting corresponding genotypes in their breeding programs through a genomic selection approach.

However, for any application it is crucial to ensure that the TPE is well-represented in the training data. Training solely on the FIP 1.0 dataset and predicting "distant" environments in Germany and France showed significant bias and reduced agreement on genotype ranking. Adding MET data from an unrelated TPE did not improve results. This strongly indicates the difficulty of learning stress factors that are underrepresented in the training environment. We assume that combining multiple field phenotyping platform datasets that cover different stressors and growth conditions could substantially increase the power of such models. With the FIP 1.0 dataset, one such dataset is publicly available, and hopefully more will follow soon. Given recent developments in mobile and/or portable device phenotyping, breeders may have tools at hand to extend such data sets to better cover their TPE.

Variety testing data have previously demonstrated the potential for location-specific variety recommendations (Herrera et al. 2018). However, variation across years in such data is usually significant and the reason why variety testing typically spans at least three years in most countries. Our work demonstrates how this time span can be further extended to "virtually" test in arbitrary historical years via a learned model. This approach enables fine-grained genotype recommendations based on the frequency a variety results as a top-ranked genotype per year-location. Given the availability of genetic marker data, such a model could be incrementally extended each year, effectively leveraging the substantial work produced by variety testing.

Traditional variety testing involves experiments across numerous locations, yet recommendations are typically based on overall adjusted genotype mean values to ensure statistical robustness. The proposed PTB model offers promising potential to enhance these recommendations by making them location-specific, potentially increasing crop production sustainability through locally adapted agricultural practices. By integrating MET data with location-specific descriptive factors such as management practices and soil types, the model could effectively distinguish between relatively stable location effects and variable environmental factors that change year to year. This separation would likely improve prediction accuracy for location-specific recommendations, providing more tailored guidance to producers.

Looking beyond the present, the proposed PTB model has the potential to enable predictions related to climate change for future climate scenarios. We refrained from pursuing this in our work. One reason for this decision was that projected temperature trajectories for future climate scenarios may exceed the conditions in the training data, making predictions unreliable. Such out-of-domain effects

were observed in the results for German and French environments in general, and for specific years for the Swiss environment. The year 2016 for example with very low reported yield was characterized by low-light conditions in Spring and an in general very wet season in all Europe (Le Gouis, Oury, and Charmet 2020), conditions that may not be present in the FIP 1.0 data set used in training. Having field phenotyping data from environments that correspond to such condition could significantly help to address this limitation.

Another reason for not projecting results onto future climate scenarios was that the PTB model relies on hourly temperature and VPD values. Climate scenario projections on an hourly basis are still rare; for example, those from MeteoSwiss are currently available only on a daily basis (Fischer et al. 2022). However, this situation is expected to improve in the near-future.

In future works, predictions could be extended beyond yield to include quality traits such as grain protein content, Zeleny sedimentation values, and other relevant parameters. Such extensions would, of course, be dependent upon the availability of corresponding phenotypic data, a limitation of the data sets used in this work. Yet, the end-to-end deep learning approach exemplified by the PTB model could be readily adapted for multi-trait prediction with minimal additional complexity.

We conclude that the integration of high-throughput field phenotyping platform data with multi-environment trial data in deep learning models represents a significant advancement in crop performance prediction. Our PTB model successfully bridged these complementary data sources, enabling more accurate predictions across diverse environments, and revealing important genotype-specific responses to environmental stressors. The proposed approach offers valuable insights for developing optimized crop ideotypes and enhancing breeding decisions. While challenges remain in extrapolating to environments beyond the ones learned from the training data, the framework established here provides a foundation for future work, including potential applications in climate change adaptation strategies. As more comprehensive datasets become available and climate projections improve in resolution, the predictive power and applicability of such integrated models will continue to expand, ultimately contributing to more resilient and productive agricultural systems.

4 Online Methods

4.1 Data

Three different data sets were used to train and test the PTB model, (1) the FIP 1.0 data set (Roth et al. 2024b), (2) the GABI-MET data set (Gogna et al. 2022), and (3) the CH-MET data set (Herrera et al. 2020). Details of the three data sets are described in the following.

4.1.1 FIP 1.0 data set

The FIP 1.0 data set consisted of annotated image time series acquired with the Field Imaging Platform (FIP) at ETH Zurich in six years (Roth et al. 2024b). The open data set comprises approximately 4,000 image time series of wheat plots. Each image in the time series is capturing a canopy area of 1.0×1.5 m of a specific winter wheat genotype. Time series are annotated with the secondary/low-level traits canopy cover, plant height, wheat head counts, and senescence ratings, and with the target traits yield, protein content, heading date, and final height. Of the four low-level and four target traits, all low-level traits but only the target trait yield were used for the PTB model.

Also part of the FIP 1.0 data set are genetic marker data (90k SNP array) for the GABI-WHEAT panel based on the public data set published by Gogna et al. (2022). A separate set of additional marker data for genotypes that are not part of the GABI-WHEAT panel was generated using a 45k SNP array (non-public data). Hence, two sub-data sets were assembled: one consisting solely of GABI-WHEAT panel markers and another extended set that combined overlapping markers from both public and protected sources. Marker data filtering and imputation procedures are part of the data set preparation done by Roth et al. (2024b). The final GABI-WHEAT panel marker data set after filtering comprised 372 genotypes (GABI-WHEAT panel) and 18,846 markers. The extended data set after filtering consisted

Figure 4: Image input batch example, showing random cutout per image time series (rows), varying time series length (grey images), random masking (black images), and color jittering. Time series are cropped to length 20 to improve visual representation in this manuscript, original time series lengths are <=63.

of 844 genotypes (GABI-WHEAT panel, registered Swiss varieties, and F8 generation genotypes) and 11,943 markers.

Image time series in the FIP 1.0 data set are anntotated with environmental covariates. For the PTB model, air temperature and relative humidity were used, while vapour pressure deficit (VPD) was derived from temperature and relative humidity assuming an atmospheric pressure of 101 Pa.

4.1.2 GABI-MET data set

The GABI-MET data set consists of plot-based target traits and derived adjusted genotype values (best linear unbiased estimations (BLUEs)) for eight environments as part of a multi-environment trial (MET), and genetic marker data for the GABI-WHEAT panel (see previous section) (Gogna et al. 2022). BLUEs of MET data from the eight year-locations were used as first test set for unseen environments.

The MET data were enriched with hourly temperature and relative humidity data originating from the Climate Data Center (CDC) of the German Weather Service (for German environments), and from Meteo France (SYNOP, 3-hourly data only) (for French environments). The data set comprised 300 genotypes that were part of the FIP 1.0 data set and therefore seen in training, and 60 other genotypes that were unseen in training.

4.1.3 CH-MET data set

MET data from 40 year-locations (years 2010–2013, 10 locations) from the official Swiss variety testing (Agroscope, Switzerland) were used as second test set for unseen environments (Herrera et al. 2020). Experimental field designs followed a Latin square design with three replications and 36 genotypes per year-site (only 24 genotypes in 2010). Genotype overlaps between years were in minimum 15% (2010 to 2013) but otherwise between 25 and 50%. BLUEs were calculated based on plot-based values using ASReml-R (The VSNi Team 2023) with genotype and replicate as random factors and row and columns as auto-regression of order 1 (AR1)×AR1 residual model to account for gradients on fields. Hourly temperature and relative humidity originated from the MeteoSwiss data base interface Idaweb.

THE CH-MET data included 71 genotypes in total. For 35 genotypes, genetic marker data were available, 12 of these genotypes were also part of the FIP 1.0 data set and hence seen in training. The remaining 13 genotypes were unseen in training.

4.2 Model architecture

The PTB model consists of two encoder parts (P-state and G×E-state encoder) and one decoder part (target trait decoder) (Figure 1). The P-state encoder takes image time series as input and predicts time series of P-traits (canopy cover, plant height, wheat head counts, and senescence) and time series of P-states (latent representation of growth of the measured phenotype). The G×E-state encoder takes genetic marker data (SNPs) and time series of environmental covariates (temperature and VPD) as input and predicts time series of G×E-states (latent representation of growth of the modelled phenotype). The P-state from the P-state encoder corresponds to the G×E-state from the G×E-state encoder in regard to representing the same characteristic (growth) with different means (phenotyping data versus genomic and environmental data). The target trait decoder takes P-state/G×E-state time series and genetic marker data and predicts yield. The two encoders are trained in parallel to agree on the P-state/G×E-state, following a concept inspired by recent trends in self-supervised, non-contrastive learning from computer vision and large language model research (Tian, Chen, and Ganguli 2021).

The PTB model was implemented in pytorch (Paszke et al. 2017) and trained with pytorch-lightning (Falcon and team 2025). Common to all model parts are rectified linear unit (ReLU) activation functions (Glorot, Bordes, and Bengio 2011). Further details on specific model parts are described in the following.

4.2.1 P-state encoder

Feature embedding of images per point in time was achieved using a ResNet18 architecture, with weights pretrained on ImageNet (He et al. 2015), where the last classification layer was removed, yielding a 512-dimensional output vector. This representation served as an initial input for a transformer model.

The transformer model (Vaswani et al. 2023) consisted of 32 attention heads and 16 layers. Positional encoding was added to the input embeddings using sinusoidal functions as proposed in Vaswani et al. (2023). Subsequent to the transformer, four fully connected layers were applied with dimensions 512, 128, 6, and 5. A sigmoid activation function was used to normalize trait predictions (canopy cover, plant height, head counts, senescence, and P-state) into $[0, 1]$. While the four traits were trained in a supervised way using labels, the fifth output value, the P-state, was intended to represent a latent representation of growth. This trait was trained in a self-supervised manner by simultaneously predicting its correspondence, the G×E-state, with the G×E-state encoder based on environmental covariates and genetic marker data, as described in the following.

4.2.2 G×E-state encoder

Time series with hourly measurements of temperature and VPD and genetic marker data were input of this second encoder. As encoder for the genetic data based on SNPs, the DeepGS model architecture suggested by Ma et al. (2018) was used. Two separate DeepGS model parts were included in the encoder to estimate temperature growth response parameters and VPD stress response parameters. The DeepGS architecture followed the original publication, with an additional batch normalization layer before the first layer.

A first set of spline knot coordinates predicted by DeepGS was used to parametrize a 2D spline fitting growth response to temperature and days after sowing, i.e., changing responses to temperature with time. A second set of spline knot coordinates predicted by DeepGS was used to parametrize a 1D spline that fitted stress response to VPD, i.e., a constant penalization of growth with VPD.

The G×E-state GxE at point in time t was computed as a cumulated response to temperature and VPD,

$$GxE_t = \sum_{t_n=0}^{t} \left(\text{spl2D}(t_n, T_{t_n}) \times \text{spl1D}(V_{t_n}) \right), \tag{3}$$

where spl2D represents a 2D spline with 10×8 knots and spl1D a 1D spline with 8 knots, V denotes VPD, and T denotes temperature.

4.2.3 Target trait decoder

The estimated G×E-state values served as input for a transformer-based decoder trained to predict yield. Daily G×E-state values were retained up to 260 days after sowing as input tokens, with subsequent values discarded. An additional token was introduced for yield prediction. Positional encoding was applied to the sequence to annotate DAS.

A transformer model was employed with an embedding dimension of 128, 64 attention heads, and 16 layers. The output from the transformer was processed through three fully connected layers (dimensionality 128, 64, 64), yielding the final yield prediction based on G×E-state time series.

For the genetic main effect, one additional DeepGS models was included. The final yield prediction was obtained as the sum of the G×E effect and the genetic main effect.

4.3 Training procedure

The model was trained using a weighted combination of four loss terms (Figure 1), two for the supervised training part (L_1, L_4) and two for the self-supervised part (L_2, L_3),

$$L_{total} = L_1 + L_2 + L_3 + \lambda \cdot L_4 \,. \tag{4}$$

L_1 minimized the mean squared error (MSE) between predicted P-traits and labelled P-traits (canopy cover, plant height, wheat heads, and senescence) per image in image time series. L_2 minimized the MSE between predicted G×E-state values and predicted P-state values. L_3 maximized the range of the G×E-state per time series,

$$L_3 = (1 - \max(GxE))^2 + \min(GxE)^2 \,, \tag{5}$$

where GxE are G×E-state values in the range $[0, 1]$. This loss is required to prevent the model from minimizing L_2 by predicting all-zero for P-state and G×E-state, i.e., it enforces

$$\sum_t GxE_t > 0 \tag{6}$$

to prevent model collapse. L_4 minimized the MSE between predicted yield values and labelled yield values per image time series. A scaling factor of $\lambda = 10$ for L_4 was empirically determined to balance its contribution relative to the other loss components, ensuring that no single term would dominate the optimization process.

Training was conducted in two phases. In a first training phase for 16 epochs, the genetic main effect was set to zero, and the PTB trained using all available samples. In a second training phase for 28 epochs, genetic main effects were included while freezing the P-state prediction component. Only samples with available genetic marker data were used in this second phase.

If MET data were included in the training procedure (see next Section), an additional third training phase followed. In this phase, the additional MET data were used to fine-tune the model for another 125 (GABI-MET) respectively 250 (CH-MET) epochs together with the image data.

In contrast to the FIP 1.0 data, the MET data could not be augmented, as they do not contain images. Consequently, only two (for GABI-MET) respectively one (for CH-MET) augmentation per image data sample were used. This ensured a close-to 1:1 ratio between image and MET data.

4.3.1 Training variants

Models were trained on either the GABI-WHEAT SNP set (noted as context 'GABI') or the extended SNP set (noted as context 'Extended'). Three training and test scenarios were chosen. Models were either trained on (1) the pre-defined training/validation/test split of the FIP 1.0 data set (5 years, 1 location), (2) the full FIP 1.0 data set as training (6 years, 1 location), or (3) the full FIP 1.0 data set combined with either part of the GABI-MET data set or CH-MET data set (7 years, 4 locations, respectively 9 years, 11 locations) as training (Figure 2). If the full FIP 1.0 data set was used, tests were only performed on unseen MET data. If GABI-MET was additionally included in the training,

one year (2009) was included in the training while the other year (2010) served as test set. If CH-MET was additionally included, three years (2010–2012) were included in the training while the left-over year (2013) served as test set.

4.4 Data augmentation

Images captured with the field phenotyping platform (FIP) were subjected to a squared random cutout process, extracting $35\% \pm 10\%$ of the original image, resulting in a cropped area of approximately 1.3×0.875 m (Figure 4). The cutout position was randomly determined within the original image. Additionally, images were randomly rotated and resampled to 224×224 pixels. All random operations followed a uniform distribution. The cutout bounding box was kept constant within the image time series, resulting in image patches that showed approximately the same three plant rows throughout the season.

We also applied physics based color augmentation using Kornia's `RandomPlanckianJitter` that simulates illumination changes between scenes (probability of 0.8, in "blackbody" mode, selecting color temperatures from the range [4, 15]). Temporal masking of images within the time series was performed at a frequency of 0.1. Each sample underwent 12 augmentations, including the cutout and random color jittering.

4.5 Baseline

As baseline served a genomic prediction model based on a random regression to environmental covariates (Jarquín et al. 2014). The FIP 1.0 data set publications provides baselines for the proposed train/validation/test splits. For the additional test sets GABI-MET and CH-MET, identical models as in the FIP 1.0 paper were trained on the full FIP 1.0 data set and on combined FIP 1.0 and MET training sets.

Environmental covariates were the standardized precipitation and evapotranspiration index (SPEI) (Beguería et al. 2014), VPD, air temperature at 2.0 m above ground, and precipitation. For the SPEI and temperature, the mean, maximum and minimum value over the season were used, for VPD, the mean and maximum, and for precipitation, the sum of the values was used. Models were fitted with ASReml-R (The VSNi Team 2023).

4.6 Evaluation metrics

As test metric, mean squared error (MSE), Pearson's r (correlation) and mean absolute percentage error (MAPE) were used. MSE was decomposed in the three components squared bias, non-unity of slope, and lack of correlation as proposed by Gauch, Hwang, and Fick (2003). Pearson's r (correlation) and mean absolute percentage error (MAPE) were calculated per environment and then averaged ('Within E'), and across all environments simultaneously ('Across E').

P-state traits were predicted on a plot base, i.e., with replications of genotypes. This allowed to calculate heritability according to Oakey et al. (2006) with SpATS (Rodríguez-Álvarez et al. 2018).

4.7 Model analysis and interpretation

4.7.1 Ideotype clustering

To visualize optimized phenotypes in regard to yield (so-called ideotypes), the learned responses to temperature and VPD of the model trained on the full FIP 1.0 data set were clustered according to their shape. K-shape clustering (Paparrizos and Gravano 2016) with cluster sizes $k = [4, 5, 6, ..., 9, 10, 20, ..., 80]$ were evaluated regarding their Dunn index (Arbelaitz et al. 2013) with the R package dtwclust (Sarda-Espinosa 2023). Clusters built by the winning configuration ($k = 40$) were annotated with yield performance and yield sensitivity mean values evaluated on the GABI-MET data set with a Finlay-Wilkinson regression (Finlay and Wilkinson 1963) performed with the R package statgenGxE (van Rossum 2023). Three clusters that were at the extremes of the yield performance versus yield sensitivity space were

visually identified. The targeted ideotype, a cluster with high yield and low yield sensitivity, was contrasted two to other cluster, one with low yield and low yield sensitivity and one with high yield but high yield sensitivity.

4.7.2 Yield prediction and variety recommendations for 20 years

To demonstrate the potential of the learned PTB model to predict for unseen years, genotype performances were predicted for 20 years of environmental data (temperature and relative humidity) from MeteoSwiss. Predictions were performed for the 10 locations of the CH-MET variety testing data set for 14 varieties (quality class TOP and I) that were on the recommended variety list in 2018 (Courvoisier et al. 2017). Variety predictions were then ranked per year for number of times being the top 1 variety per location. Yield per location and variety were averaged to mean yield and variance of yield per year, and compared with published annual yield per area data by Swiss granum `https://www.swissgranum.ch/zahlen/inlandproduktion`. As Swiss granum data and variety testing data are characterized by the typical shift in absolute values between experimentally determined yield and yield measured on true agricultural fields, swiss granum data were aligned to variety testing data by correcting for a fixed offset (-0.63) and slope (2.26) determined in 2010–2014 where data of both types were available.

Data availability

Data from the field phenotyping platform 1.0 (FIP 1.0) platform have been released as the FIP1.0 dataset, `https://doi.org/10.1093/gigascience/giaf051`. The GABI-MET dataset is public, `https://doi.org/10.5447/ipk/2022/18`. The CH-MET dataset was at the time of writing subject to limitations that may have been resolved in the meantime. Please contact Agroscope (Juan Herrera) for further details.

Code availability

All code are available from the gitlab repository of ETH Zurich—Model code repository: `https://gitlab.ethz.ch/crop_phenotyping/PTB_model`; Data preparation code repository: `https://gitlab.ethz.ch/crop_phenotyping/PTB_annotate`. Model weights of trained models are provided with the code respository `https://gitlab.ethz.ch/crop_phenotyping/PTB_model` ('./scripts/weights'). Preprocessed input batches and all model weights are available upon request (> 1.5 TB).

References

Arbelaitz, Olatz, Ibai Gurrutxaga, Javier Muguerza, Jesús M. Pérez, and Iñigo Perona (2013). "An Extensive Comparative Study of Cluster Validity Indices". In: *Pattern Recognition* 46.1, pp. 243–256. DOI: `10.1016/j.patcog.2012.07.021`.

Beguería, Santiago, Sergio M. Vicente-Serrano, Fergus Reig, and Borja Latorre (2014). "Standardized Precipitation Evapotranspiration Index (SPEI) Revisited: Parameter Fitting, Evapotranspiration Models, Tools, Datasets and Drought Monitoring". In: *International Journal of Climatology* 34.10, pp. 3001–3023. DOI: `10.1002/joc.3887`.

Chapman, Scott C. (2008). "Use of Crop Models to Understand Genotype by Environment Interactions for Drought in Real-World and Simulated Plant Breeding Trials". In: *Euphytica* 161.1, pp. 195–208. DOI: `10.1007/s10681-007-9623-z`.

Courvoisier, Numa, Lilia Levy Häner, Mario Bertossa, Etienne Thévoz, Martin Anders, Peter Stoll, Thomas Weisflog, and Jacques Dugon (2017). *Liste Der Empfohlenen Getreidesorten Für Die Ernte 2018*.

Diepenbrock, Christine H, Tom Tang, Michael Jines, Frank Technow, Sara Lira, Dean Podlich, Mark Cooper, and Carlos Messina (2021). "Can We Harness Digital Technologies and Physiology to Hasten Genetic Gain in US Maize Breeding?" In: *Plant Physiology*, pp. 1–17. DOI: 10.1093/plphys/kiab527.

Falcon, William and The PyTorch Lightning team (2025). *PyTorch Lightning*. Zenodo. DOI: 10.5281/zenodo.15053754.

Falconer, Douglas S. and Trudy F. C. Mackay (1996). *Introduction to Quantitative Genetics*. Harlow, Essex: Longman Group Limited.

Finlay, K. W. and G. N. Wilkinson (1963). "The Analysis of Adaptation in a Plant-Breeding Programme". In: *Australian Journal of Agricultural Research* 14, pp. 742–754. DOI: 10.1071/AR9630742.

Fischer, A. M., K. M. Strassmann, M. Croci-Maspoli, A. M. Hama, R. Knutti, S. Kotlarski, C. Schär, C. Schnadt Poberaj, N. Ban, M. Bavay, U. Beyerle, D. N. Bresch, S. Brönnimann, P. Burlando, A. Casanueva, S. Fatichi, I. Feigenwinter, E. M. Fischer, M. Hirschi, M. A. Liniger, C. Marty, I. Medhaug, N. Peleg, M. Pickl, C. C. Raible, J. Rajczak, O. Rössler, S. C. Scherrer, C. Schwierz, S. I. Seneviratne, M. Skelton, S. L. Sørland, C. Spirig, F. Tschurr, J. Zeder, and E. M. Zubler (2022). "Climate Scenarios for Switzerland CH2018 – Approach and Implications". In: *Climate Services* 26, p. 100288. DOI: 10.1016/j.cliser.2022.100288.

Fisher, R. A. and W. A. Mackenzie (1923). "Studies in Crop Variation. II. The Manurial Response of Different Potato Varieties". In: *The Journal of Agricultural Science* 13.3, pp. 311–320. DOI: 10.1017/S0021859600003592.

Gauch, Hugh G., J. T. Gene Hwang, and Gary W. Fick (2003). "Model Evaluation by Comparison of Model-Based Predictions and Measured Values". In: *Agronomy Journal* 95.6, pp. 1442–1446. DOI: 10.2134/agronj2003.1442.

Glorot, Xavier, Antoine Bordes, and Yoshua Bengio (2011). "Deep Sparse Rectifier Neural Networks". In: *AISTATS*.

Gogna, Abhishek, Albert Wilhelm Schulthess Börgel, Marion Röder, Martin W Ganal, and Jochen Christoph Reif (2022). *The Genotypic Data of Elite European Cultivar Panel Comprising 358 Winter and 14 Summer Wheat Varieties Released from 1975 to 2007 at Different Marker Densities*. DOI: 10.5061/dryad.n02v6wwzc.

He, Kaiming, Xiangyu Zhang, Shaoqing Ren, and Jian Sun (2015). *Deep Residual Learning for Image Recognition*. DOI: 10.48550/arXiv.1512.03385. arXiv: 1512.03385 [cs].

Herrera, Juan M., Lilia Levy Häner, Annelie Holzkämper, and Didier Pellet (2018). "Evaluation of Ridge Regression for Country-Wide Prediction of Genotype-Specific Grain Yields of Wheat". In: *Agricultural and Forest Meteorology* 252. DOI: 10.1016/j.agrformet.2017.12.263.

Herrera, Juan M., Lilia Levy Häner, Fabio Mascher, Jürg Hiltbrunner, Dario Fossati, Cécile Brabant, Raphaël Charles, and Didier Pellet (2020). "Lessons From 20 Years of Studies of Wheat Genotypes in Multiple Environments and Under Contrasting Production Systems". In: *Frontiers in Plant Science* 10.1745. DOI: 10.3389/fpls.2019.01745.

Hund, Andreas, Lukas Kronenberg, Jonas Anderegg, Kang Yu, and Achim Walter (2019). "Non-Invasive Field Phenotyping of Cereal Development". In: *Advances in Breeding Techniques for Cereal Crops*. Ed. by Frank Ordon and Wolfgang Friedt. Cambridge, United Kingdom: Burleigh Dodds Science Publishing, Cambridge, UK, pp. 249–292. DOI: 10.19103/as.2019.0051.13.

Jarquin, Diego, Reka Howard, Alencar Xavier, and Sruti Das Choudhury (2018). "Increasing Predictive Ability by Modeling Interactions between Environments, Genotype and Canopy Coverage Image Data for Soybeans". In: *Agronomy* 8.4, p. 51. DOI: 10.3390/agronomy8040051.

Jarquín, Diego, José Crossa, Xavier Lacaze, Philippe Du Cheyron, Joëlle Daucourt, Josiane Lorgeou, François Piraux, Laurent Guerreiro, Paulino Pérez, Mario Calus, Juan Burgueño, and Gustavo de los Campos (2014). "A Reaction Norm Model for Genomic Selection Using High-Dimensional Genomic and Environmental Data". In: *Theoretical and Applied Genetics* 127.3, pp. 595–607. DOI: 10.1007/s00122-013-2243-1.

Kirchgessner, Norbert, Frank Liebisch, Kang Yu, Johannes Pfeifer, Michael Friedli, Andreas Hund, and Achim Walter (2017). "The ETH Field Phenotyping Platform FIP: A Cable-Suspended Multi-Sensor System". In: *Functional Plant Biology* 44, pp. 154–168. DOI: 10.1071/FP16165.

Le Gouis, Jacques, François-Xavier Oury, and Gilles Charmet (2020). "How Changes in Climate and Agricultural Practices Influenced Wheat Production in Western Europe". In: *Journal of Cereal Science* 93, p. 102960. DOI: 10.1016/j.jcs.2020.102960.

Li, Xianran, Tingting Guo, Jinyu Wang, Wubishet A. Bekele, Sivakumar Sukumaran, Adam E. Vanous, James P. McNellie, Laura E. Tibbs-Cortes, Marta S. Lopes, Kendall R. Lamkey, Mark E. Westgate, John K. McKay, Sotirios V. Archontoulis, Matthew P. Reynolds, Nicholas A. Tinker, Patrick S. Schnable, and Jianming Yu (2021). "An Integrated Framework Reinstating the Environmental Dimension for GWAS and Genomic Selection in Crops". In: *Molecular Plant* 14.6, pp. 874–887. DOI: 10.1016/j.molp.2021.03.010.

Ma, Wenlong, Zhixu Qiu, Jie Song, Jiajia Li, Qian Cheng, Jingjing Zhai, and Chuang Ma (2018). "A Deep Convolutional Neural Network Approach for Predicting Phenotypes from Genotypes". In: *Planta* 248.5, pp. 1307–1318. DOI: 10.1007/s00425-018-2976-9.

Malosetti, Marcos, Daniela Bustos-Korts, Martin P. Boer, and Fred A. van Eeuwijk (2016). "Predicting Responses in Multiple Environments: Issues in Relation to Genotype x Environment Interactions". In: *Crop Science* 56, pp. 2210–2222. DOI: 10.2135/cropsci2015.05.0311.

Meuwissen, T. H. E., B. J. Hayes, and M. E. Goddard (2001). "Prediction of Total Genetic Value Using Genome-Wide Dense Marker Maps". In: *Genetics* 157, pp. 1819–1829. DOI: 10.1534/genetics.116.189795.

Millet, Emilie J., Willem Kruijer, Aude Coupel-Ledru, Santiago Alvarez Prado, Llorenç Cabrera-Bosquet, Sébastien Lacube, Alain Charcosset, Claude Welcker, Fred van Eeuwijk, and François Tardieu (2019). "Genomic Prediction of Maize Yield across European Environmental Conditions". In: *Nature Genetics* 51.6, pp. 952–956. DOI: 10.1038/s41588-019-0414-y.

Oakey, Helena, Arunas Verbyla, Wayne Pitchford, Brian Cullis, and Haydn Kuchel (2006). "Joint Modeling of Additive and Non-Additive Genetic Line Effects in Single Field Trials". In: *Theoretical and Applied Genetics* 113, pp. 809–819. DOI: 10.1007/s00122-006-0333-z.

Paparrizos, John and Luis Gravano (2016). "K-Shape: Efficient and Accurate Clustering of Time Series". In: *SIGMOD Rec.* 45.1, pp. 69–76. DOI: 10.1145/2949741.2949758.

Parent, Boris and François Tardieu (2012). "Temperature Responses of Developmental Processes Have Not Been Affected by Breeding in Different Ecological Areas for 17 Crop Species". In: *New Phytologist* 194, pp. 760–774. DOI: 10.1111/j.1469-8137.2012.04086.x.

Paszke, Adam, Sam Gross, Soumith Chintala, Gregory Chanan, Edward Yang, Zachary DeVito, Zeming Lin, Alban Desmaison, Luca Antiga, and Adam Lerer (2017). "Automatic differentiation in PyTorch". In: *NIPS-W*.

Porter, John R. and Megan Gawith (1999). "Temperatures and the Growth and Development of Wheat a Review". In: *European Journal of Agronomy* 10, pp. 23–36. DOI: 10.1016/S1161-0301(98)00047-1.

Rodríguez-Álvarez, María Xosé, Martin P. Boer, Fred A. van Eeuwijk, and Paul H.C. Eilers (2018). "Correcting for Spatial Heterogeneity in Plant Breeding Experiments with P-splines". In: *Spatial Statistics* 23, pp. 52–71. DOI: 10.1016/j.spasta.2017.10.003.

Roth, Lukas, Martina Binder, Norbert Kirchgessner, Flavian Tschurr, Steven Yates, Andreas Hund, Lukas Kronenberg, and Achim Walter (2024a). "From Neglecting to Including Cultivar-Specific Per Se Temperature Responses: Extending the Concept of Thermal Time in Field Crops". In: *Plant Phenomics* 6, p. 0185. DOI: 10.34133/plantphenomics.0185.

Roth, Lukas, Mike Boss, Norbert Kirchgessner, Helge Aasen, Brenda Patricia Aguirre-Cuellar, Price Pius Atuah Akiina, Jonas Anderegg, Joaquin Gajardo Castillo, Xiaoran Chen, Simon Corrado, Krzysztof Cybulski, Beat Keller, Stefan Göbel Kortstee, Lukas Kronenberg, Frank Liebisch, Paraskevi Nousi, Corina Oppliger, Gregor Perich, Johannes Pfeifer, Kang Yu, Nicola Storni, Flavian Tschurr, Simon Treier, Michele Volpi, Hansueli Zellweger, Olivia Zumsteg, Andreas Hund, and Achim Walter (2024b). *The FIP 1.0 Data Set: Highly Resolved Annotated Image Time Series of 4,000 Wheat Plots Grown in Six Years.* DOI: 10.1101/2024.10.04.616624.

Roth, Lukas, Hans-Peter Piepho, and Andreas Hund (2022). "Phenomics Data Processing: Extracting Dose-Response Curve Parameters from High-Resolution Temperature Courses and Repeated Field-Based Wheat Height Measurements". In: *in silico Plants* 4.1. DOI: 10.1093/insilicoplants/diac007.

Roth, Lukas, María Xosé Rodríguez-Álvarez, Fred van Eeuwijk, Hans-Peter Piepho, and Andreas Hund (2021). "Phenomics Data Processing: A Plot-Level Model for Repeated Measurements to Extract the Timing of Key Stages and Quantities at Defined Time Points". In: *Field Crops Research* 274.108314. DOI: `10.1016/j.fcr.2021.108314`.

Sarda-Espinosa, Alexis (2023). *dtwclust: Time Series Clustering Along with Optimizations for the Dynamic Time Warping Distance*. R package version 5.5.12.

Schoppach, Rémy and Walid Sadok (2013). "Transpiration Sensitivities to Evaporative Demand and Leaf Areas Vary with Night and Day Warming Regimes among Wheat Genotypes". In: *Functional Plant Biology* 40.7, pp. 708–718. DOI: `10.1071/FP13028`.

Slafer, G. A. (1996). "Differences in Phasic Development Rate amongst Wheat Cultivars Independent of Responses to Photoperiod and Vernalization. A Viewpoint of the Intrinsic Earliness Hypothesis". In: *The Journal of Agricultural Science* 126.4, pp. 403–419. DOI: `10.1017/S0021859600075493`.

The VSNi Team (2023). *asreml: Fits Linear Mixed Models using REML*. R package version 4.2.0.332.

Tian, Yuandong, Xinlei Chen, and Surya Ganguli (2021). *Understanding Self-Supervised Learning Dynamics without Contrastive Pairs*. arXiv: `2102.06810 [cs]`.

Tolhurst, Daniel J., R. Chris Gaynor, Brian Gardunia, John M. Hickey, and Gregor Gorjanc (2022). "Genomic Selection Using Random Regressions on Known and Latent Environmental Covariates". In: *Theoretical and Applied Genetics* i. DOI: `10.1007/s00122-022-04186-w`.

van Rossum, Bart-Jan (2023). *statgenGxE: Genotype by Environment (GxE) Analysis*. R package version 1.0.6.

van Voorn, George A. K., Martin P. Boer, Sandra Huynh Truong, Nicholas A. Friedenberg, Shota Gugushvili, Ryan McCormick, Daniela Bustos Korts, Carlos D. Messina, and Fred A. van Eeuwijk (2023). "A Conceptual Framework for the Dynamic Modeling of Time-Resolved Phenotypes for Sets of Genotype-Environment-Management Combinations: A Model Library". In: *Frontiers in Plant Science* 14. DOI: `10.3389/fpls.2023.1172359`.

Vaswani, Ashish, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N. Gomez, Lukasz Kaiser, and Illia Polosukhin (2023). *Attention Is All You Need*. DOI: `10.48550/arXiv.1706.03762`. arXiv: `1706.03762 [cs]`.

Walkowiak, Sean, Liangliang Gao, Cecile Monat, Georg Haberer, Mulualem T. Kassa, Jemima Brinton, Ricardo H. Ramirez-Gonzalez, Markus C. Kolodziej, Emily Delorean, Dinushika Thambugala, Valentyna Klymiuk, Brook Byrns, Heidrun Gundlach, Venkat Bandi, Jorge Nunez Siri, Kirby Nilsen, Catharine Aquino, Axel Himmelbach, Dario Copetti, Tomohiro Ban, Luca Venturini, Michael Bevan, Bernardo Clavijo, Dal-Hoe Koo, Jennifer Ens, Krystalee Wiebe, Amidou N'Diaye, Allen K. Fritz, Carl Gutwin, Anne Fiebig, Christine Fosker, Bin Xiao Fu, Gonzalo Garcia Accinelli, Keith A. Gardner, Nick Fradgley, Juan Gutierrez-Gonzalez, Gwyneth Halstead-Nussloch, Masaomi Hatakeyama, Chu Shin Koh, Jasline Deek, Alejandro C. Costamagna, Pierre Fobert, Darren Heavens, Hiroyuki Kanamori, Kanako Kawaura, Fuminori Kobayashi, Ksenia Krasileva, Tony Kuo, Neil McKenzie, Kazuki Murata, Yusuke Nabeka, Timothy Paape, Sudharsan Padmarasu, Lawrence Percival-Alwyn, Sateesh Kagale, Uwe Scholz, Jun Sese, Philomin Juliana, Ravi Singh, Rie Shimizu-Inatsugi, David Swarbreck, James Cockram, Hikmet Budak, Toshiaki Tameshige, Tsuyoshi Tanaka, Hiroyuki Tsuji, Jonathan Wright, Jianzhong Wu, Burkhard Steuernagel, Ian Small, Sylvie Cloutier, Gabriel Keeble-Gagnère, Gary Muehlbauer, Josquin Tibbets, Shuhei Nasuda, Joanna Melonek, Pierre J. Hucl, Andrew G. Sharpe, Matthew Clark, Erik Legg, Arvind Bharti, Peter Langridge, Anthony Hall, Cristobal Uauy, Martin Mascher, Simon G. Krattinger, Hirokazu Handa, Kentaro K. Shimizu, Assaf Distelfeld, Ken Chalmers, Beat Keller, Klaus F. X. Mayer, Jesse Poland, Nils Stein, Curt A. McCartney, Manuel Spannagl, Thomas Wicker, and Curtis J. Pozniak (2020). "Multiple Wheat Genomes Reveal Global Variation in Modern Breeding". In: *Nature* 588.7837, pp. 277–283. DOI: `10.1038/s41586-020-2961-x`.

Wang, Enli, Pierre Martre, Zhigan Zhao, Frank Ewert, Andrea Maiorano, Reimund P. Rötter, Bruce A. Kimball, Michael J. Ottman, Gerard W. Wall, Jeffrey W. White, Matthew P. Reynolds, Phillip D. Alderman, Pramod K. Aggarwal, Jakarat Anothai, Bruno Basso, Christian Biernath, Davide Cammarano, Andrew J. Challinor, Giacomo De Sanctis, Jordi Doltra, Elias Fereres, Margarita Garcia-Vila, Sebastian Gayler, Gerrit Hoogenboom, Leslie A. Hunt, Roberto C. Izaurralde, Mohamed Jabloun, Curtis D. Jones, Kurt C. Kersebaum, Ann Kristin Koehler, Leilei Liu, Christoph Müller, Soora Naresh

Kumar, Claas Nendel, Garry O'Leary, Jørgen E. Olesen, Taru Palosuo, Eckart Priesack, Ehsan Eyshi Rezaei, Dominique Ripoche, Alex C. Ruane, Mikhail A. Semenov, Iurii Shcherbak, Claudio Stöckle, Pierre Stratonovitch, Thilo Streck, Iwan Supit, Fulu Tao, Peter Thorburn, Katharina Waha, Daniel Wallach, Zhimin Wang, Joost Wolf, Yan Zhu, and Senthold Asseng (2017). "The Uncertainty of Crop Yield Projections Is Reduced by Improved Temperature Response Functions". In: *Nature Plants* 3.17102. DOI: 10.1038/nplants.2017.102.

Acknowledgements

L.R. discloses support for the research of this work from Swiss Data Science Center [grant number PHENO-MINE C21-04]. The authors would like to thank the Crop Science Group of ETH Zurich, especially Achim Walter, Andreas Hund and Norbert Kirchgessner, for their massive support in carrying out the field trials and phenotyping experiments, and Delley Semences et Plants SA (DSP), especially Flavio Foiada, for their essential support in preparing seed batches for the Swiss varieties.

Authors contributions

L.R.: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data Curation, Visualization, Funding acquisition, Investigation, Writing – original draft. J.M.H.: Methodology, Validation, Resources, Data Curation, Writing - Review & Editing. L.L.H.: Resources, Data Curation, Writing - Review & Editing. D.P.: Project administration, Resources, Writing - Review & Editing. D.F.: Project administration, Resources, Writing - Review & Editing. M.B.: Methodology, Writing - Review & Editing X.C.: Methodology, Writing - Review & Editing P.N.: Methodology, Writing - Review & Editing M.V.: Methodology, Resources, Supervision, Project administration, Writing - Review & Editing

Competing interests

The authors declare no competing interests.

Extended data

Extended figures

The following figures visualize predictions versus observed values for the three data sets (1) field phenotyping platform 1.0 (FIP 1.0) (Figure 5), (2) GABI multi-environment trial (GABI-MET) (Figure 6) and (3) Swiss variety testing multi-environment trial (CH-MET) (Figure 7).

Figure 5: Predicted versus measured yield for the field phenotyping platform 1.0 (FIP 1.0) (Extended SNP marker) data set for the baseline (random regression, RR) and the plant time bender (PTB) model.

Figure 6: Predicted versus measured yield for the GABI multi-environment trial (GABI-MET) (2010) data set for the baseline (random regression, RR) and the plant time bender (PTB) model.

Figure 7: Predicted versus measured yield for the Swiss variety testing multi-environment trial (CH-MET) data (2013) set for the baseline (random regression, RR) and the plant time bender (PTB) model.

Acronyms

AR1 auto-regression of order 1. 8

ASReml-R statistical package that fits linear mixed models using Residual Maximum Like- lihood (REML). . 8, 12

BLUE best linear unbiased estimation. 8

CDC Climate Data Center. 8

CH-MET Swiss variety testing multi-environment trial. 2, 4–6, 8, 10, 11, 13, 15

DAS days after sowing. 5

DNA deoxyribonucleic acid. 1

DWD Deutscher Wetterdienst. 1

FIP field phenotyping platform. 8

FIP 1.0 field phenotyping platform 1.0. 2, 4–6, 8, 11, 13

GABI-MET GABI multi-environment trial. 2, 4–6, 8, 10, 11, 13, 14

GABI-WHEAT Genomanalyse im Biologischen System Pflanze - Weizen. 8, 11

MAPE mean absolute percentage error. 2, 4, 11

MET multi-environment trial. 1, 2, 4, 8, 10, 11

MeteoSwiss Federal Office of Meteorology and Climatology MeteoSwiss.. 1

MSE mean squared error. 2, 6, 10, 11

PTB plant time bender. 2–6, 9–11, 13–15

RGB red, green, and blue. 2

SNP single-nucleotide polymorphism. 2, 3, 8–11

SPEI standardized precipitation and evapotranspiration index. 11

SYNOP surface synoptic observation. 8

VPD vapour pressure deficit. 3, 5, 7–12