HIV-1 Amplifies IL-8 Response Of Human Stellate Cells To Gram-Positive Microbial

Products Via H4K5 Histone Acetylation

Lumin Zhang[1]*, Rabab O. Ali[1]*, Naichaya Chamroonkul[5], Parissa Tabrizian[2], Myron Schwartz[2],

Ganesh Gunasekaran[2], Thomas Schiano[3], Maria Isabel Fiel[4], Steven Ward[4], Meena B. Bansal[1]

[1]Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[2]Department of Surgery, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[3]Recanti-Miler Transplant Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[4]Department of Pathology, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[5]Gastroenterology and Hepatology Unit, Division of Internal Medicine, Faculty of Medicine, Prince of

Songkla University, Hatyai, Songkhla, Thailand

*The authors contribute equally to this work

Corresponding Author

Meena B. Bansal, MD, Icahn School of Medicine at Mount Sinai, 1425 Madison Avenue, Room

11-70; Box 1123, New York, NY 10029, E-mail: meena.bansal@mssm.edu, Fax: 212-849-2574,

Tel : 212-659-9510.

Financial support

This work was supported by research grant NIH R56DK92128 and R01DK108364 to MBB

Abstract

Background: Patients living with Human Immunodeficiency Virus (PLWH) develop accelerated liver fibrosis, but the exact mechanism remains unknown. Hepatic stellate cell (HSC) activation is the cornerstone for liver fibrosis and influenced by multiple factors, such as viral infection, hepatocellular injury, chronic immune activation, gut mucosal barrier dysfunction, and microbial translocation. Unlike gram-negative bacterial products, the impact of gram-positive microbial products in Human Immunodeficiency Virus 1 (HIV-1) associated liver inflammation and fibrosis remains poorly understood. In this study, we investigated the effect of lipoteichoic acid (LTA), a major gram-positive bacterial component, on HSCs in the context of HIV-1 infection. **Methods:** Human HSCs were isolated from the livers of both non-HIV and HIV-infected patients undergoing hepatic resection. The inflammatory responses of HSCs to LTA stimulation were measured by ELISA before or after HIV-1BaL ex vivo expision. Western blotting, ChIP-qPCR and RNA-seq were used to reveal the mechanisms contributing to the IL-8 response to LTA stimulation and HIV-1BaL exposure in HSCs. **Results:** While LTA stimulation modestly increased IL-8 production in HSCs, this effect was significantly amplified in HIV-1-infected HSCs. Elevated IL-8 levels were detected in HIV-1+ liver sections, and IL-8 treatment of cultured HSCs significantly increased α-SMA and COL1A1 expression, supporting IL-8 as a key inflammatory driver of liver fibrosis in HIV-1[+] individuals. Transcriptomic analyses implicated histone acetylation as a regulator of the enhanced IL-8 response of HSCs to LTA in HIV-1 infection. Consistent with this, treatment with Trichostatin A (TSA), a histone deacetylase (HDAC) inhibitor, further increased the IL-8 response in HIV-1-infected HSCs. ChIP-qPCR confirmed that histone H4K5 acetylation facilitated IL-8 promoter transactivation, rendering HIV-1-infected cells more responsive to LTA. **Conclusions:** HIV-1 infection sensitizes HSCs to LTA stimulation, leading to an amplified IL-8 response mediated through histone acetylation that may accelerate liver fibrosis progression in PLWH. As microbial translocation persists despite effective antiretroviral

48   therapy, these findings underscore the need for targeted strategies to mitigate liver fibrosis in

49   HIV-1 infected HIV-1[+] individuals.

51   **Abstract Word Count:** 318

Keywords

Human Immunodeficiency Virus, LTA, Inflammatory response, Fibrosis, Hepatic Stellate Cells, Microbial Translocation

Abbreviations

Antiretroviral therapy (ART), Bovine serum albumin (BSA), Chromatin immunoprecipitation with quantitative PCR (ChIP-qPCR), Cycle threshold (CT), C-C chemokine receptor type 5 (CCR5), C-X-C motif chemokine receptor 1 (CXCR), Differentially expressed genes (DEGs), Dulbecco's Modified Eagle Medium (DMEM), Enzyme-linked immunosorbent assay (ELISA), Hepatic stellate cells (HSC), Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Histone deacetylase (HDAC), Human Immunodeficiency Virus 1 (HIV-1), Immunohistochemistry (IHC), Interleukin-1β (IL-1β), Interleukin-8/CXCL8 (IL-8), Lipopolysaccharide (LPS), Lipoteichoic acid (LTA), Multiplicity of infection (MOI), Pathogen-associated molecular pattern (PAMP), Primary HSCs (pHSCs), Persons living with HIV (PLWH), RNA sequencing (RNA-seq), Sirtuin (SRT), Toll-Like Receptor (TLR), Trichostatin A (TSA), Transcription factor (TF), Tumor necrosis factor α (TNF-α), and Weighted Gene Co-Expression Network Analysis (WGCNA), Body Mass Index (BMI), Liver Stiffness Measurement (LSM), Platelet count (Plt), Aspartate Aminotransferase (AST), Alanine Aminotransferase (ALT), Alkaline Phosphatase (ALP), Albumin (Alb), Fasting Blood Sugar (FBS), Total cholesterol level (Chol), Triglycerides (TG), High-Density Lipoprotein (HDL), Low-Density Lipoprotein (LDL), Controlled Attenuation Parameter (CAP), Hepatic Steatosis Index (HIS), Fibrosis-4 Index (FIB-4).

Introduction

Human immunodeficiency virus 1 (HIV-1) infection is characterized by severe depletion of CD4+ T cells, leading to attenuated systemic immunity. Consequently, individuals living with HIV-1 experience increased susceptibility to opportunistic bacterial infections and community-acquired infections. For instance, the risk of *Streptococcus pneumoniae* infection is 100 times higher in HIV-positive individuals than in the general population[1]. Early during HIV infection, the depletion of gut associated CD4+ T cells disrupts the integrity of the intestinal barrier, facilitating microbial products translocation into the peripheral circulation[2]. Notably, microbial translocation remains elevated even in patients undergoing with antiretroviral therapy (ART), as ART does not restore gut permeability. Thus, microbial translocation has been identified as a major driver of sustained systemic immune activation in patients living with HIV (PLWH).

HIV-induced damage to the gastrointestinal mucosa leads to the translocation of both gram-positive and gram-negative microbial products from the gut lumen into the liver via the portal vein[3]. Although translocation of gram-negative bacterial products occurs more efficiently following mucosal injury[4], clinically relevant gram-positive bacteria can also frequently enter the peripheral circulation[5], either through shared routes or opportunistic infections. While extensive studies have addressed the impact of gram-negative microbial products, such as lipopolysaccharide (LPS), on chronic immune activation, liver injury and fibrogenesis in HIV infection[6,7], the role of gram-positive microbial products remains poor understudied. Lipoteichoic acid (LTA), a major component of the gram-positive bacterial cell walls, is recognized as pathogen-associated molecular pattern (PAMP) capable of significantly modulating the host immune immunity[8-10]. Functionally similar to LPS, LTA induces CD14-dependent nitric oxide release from vascular smooth muscle cells, macrophages, and mononuclear cells[11]. Recent studies have showed high serum LTA levels in HIV-positive individuals, associated with increased autoantibody secretion and enhanced immune responses[12]. Given the liver's unique

position as the first organ to interact with gut-derived microbial products via portal circulation,

elucidating the influence of LTA on liver cells , particularly hepatic stellate cells (HSCs), in HIV

associated chronic liver disease is critical but. remains inadequately explored.

End-stage liver disease contributes to represent a significant cause of mortality among HIV-

positive individuals despite effective ART, underscoring the persistent impact of HIV in liver

dysfunction and progressive fibrosis[13]. Several studies have established a positive correlation

between HIV RNA levels and advanced fibrosis in HIV mono-infected patient[14-16]. Furthermore,

HIV-1 co-infection with Hepatitis C Virus (HCV) accelerates liver fibrosis progression by

threefold compared to HCV mono-infection[17]. Although interactions between HSCs and liver

macrophages are crucial for HSC activation[18], HSCs remain central drivers of hepatic

fibrogenesis. Our previous studies[19,20] have shown that HIV-1 infection enhances the

inflammatory responses of liver macrophages to gram-negative microbial products, such as LPS,

resulting in elevated pro-inflammatory cytokine production including Interleukin-8/CXCL8 (IL-8).

However, the contributions of these cytokines to direct HSC activation require further

clarification.

In this study, we investigated the inflammatory responses of primary human HSCs isolated from

HIV-1 negative and HIV-1 positive individuals upon LTA stimulation. Our results demonstrated

that HIV-1 infection amplifies IL-8 responses to LTA stimulation in HSCs, and hepatic IL-8

directly exhibits pro-fibrogenic effects at the transcriptional level in HIV-1 positive liver tissues.

Additionally, we found histone acetylation, specifically H4K5 acetylation, mediates the

transactivation of IL-8 in HIV-1-infected HSCs, providing new mechanistic insights into HIV

related liver fibrosis.

Materials and Methods

Ethics Statements

De-identified liver resection specimens from HIV-1+ individuals (n=7) were provided by Drs.

Schwartz, Ganeskaran, and Fiel. In all cases, HIV was effectively suppressed with ART. Normal

liver tissue obtained from the Mount Sinai Biobank served as controls (HIV non-infected liver

tissues i.e., controls, n=10 with 9 from individuals undergoing resection for colon cancer

metastases, and 1 from a cholangiocarcinoma patient, n=10). Liver tissue samples were

obtained from patients undergoing clinical liver transplantation or tumor resection. Tissues were

formalin-fixed, frozen, or placed in Dulbecco's Modified Eagle Medium (DMEM). These studies

have been approved by Mount Sinai IRB (GCO #06-0523 and #10-1211). Tissues from non-

infected and HIV infected individuals were utilized for immunohistochemistry (IHC) staining of

IL-8. Blood samples were separately collected from HIV+ participants enrolled in a nonalcoholic

fatty liver disease (NAFLD) study. The characteristics of the study population in Table 2 are

analyzed by R package survey[21] and tableone[22].

Isolation of Human Stellate Cells from HIV Non-infected and HIV-1 Infected Liver tissues

Primary HSCs (pHSCs) were isolated from surgical resection specimens by performing

collagenase perfusion and mincing of the liver tissue. Cells were washed with PBS four times

and pelleted at 300 x G for 5 min. Cells were then re-suspended in DMEM followed by gradient

centrifugation using 25 and 50% Percoll (Pharmacia, Uppsala, Sweden), as described[23]. pHSCs

were enriched by adherence to tissue culture plates. After two weeks in culture, α-SMA

immunostaining was used to check the purity of pHSCs.

Tissue Immunohistochemistry Assay

The liver tissue sections were prepared in duplicate for each patient. After rehydrated and

treated for antigen retrieval, slides were incubated with anti-human IL-8 antibody (NOVUS, Cat

No NBP2-33819, 1:200 dilution) overnight at 4°C in a humidified chamber. The staining was

performed using ImmPress™ Duet Double staining polymer kit (Cat. MP-7724). Four random

visual areas were selected for each tissue section. The images were captured with Zen software

for LSM 880 laser air-scanning microscope (magnification: 100x). IL-8 signal intensity was

quantified by Fiji image J software (NIH, Bethesda, MD).

Flow Cytometry Analyses

CXCR1, CXCR2, and TLR2 expression in pHSCs was evaluated before and after gp120

stimulation (100ng/ml). Cells were re-suspended in PBS containing 0.5% bovine serum albumin

(BSA) at concentration 0.5- 1×10^6 cells/ml. incubated Fc-R blocking reagent, and stained with

anti-human CXCR1-APC (Biolegend, Cat No 320612), CXCR2-FITC (Biolegend, Cat No

320704), and TLR2-PE (Biolegend, Cat No 309708) for 15□min. After fixation and

permeabilization at 4^0C for 20 min, samples were analyzed on LSRII (BD Biosciences). Data

was analyzed using FCS express 6 software (De Novo software).

Histone Protein Extraction and Acetylation Protein ELISA

1×10^6 pHSCs cells and Lx2 were seeded in 10 cm dishes and infected with HIV-1$_{BaL}$ virus (NIH

AIDS Reagent program, Bethesda, MD) for 48 hrs. Total histone proteins were extracted by

EpiQuickTM total Histone extraction Kit (Cat. OP-0006) as per manufacturer instructions. 200

ng of histone protein was subjected to acetylation protein quantification using a histone

acetylation ELISA kit (Cat. P-4022-96) following manufacturer instructions.

RNA Extraction and RT-qPCR

Total RNA was extracted by using RNeasy spin columns (Qiagen, Cat No:74136). After

treatment with RNase-free DNase (Qiagen, Chatsworth, CA), RT was performed on RNA by

using Clontech RNA to cDNA EcoDry kit according to the manufacturer's protocol. RT-qPCR

was performed on Roche Lightcycle II 480. Reactions were carried out in 10 µL by using SYBR

Green PCR master mix according to the manufacturer's protocol (Bio-Rad, Cat No: 1708884).

The concentration of primer pairs was 300 nM. All primer sequences are shown in Table 1. All

reactions were performed in triplicate. To make comparisons between samples and controls, the

cycle threshold (CT), defined as the cycle number at which the fluorescence is above the fixed

threshold values were normalized to the CT of α-actin in each sample.

IL-8 ELISA Assay

Isolated HSCs were treated with different conditions for 24 hrs. IL-8 production in the

supernatant was collected at indicated time and measured by using human IL-8 ELISA Kit

(Affymetrix, Cat No: 88-8086-88) according to the manufacturer's protocol.

CHIP RT-PCR Assay

1x10^6 pHSCs cells were infected with HIV-1$_{BaL}$ virus (NIH AIDS Reagent program, Bethesda,

MD) for 48 hrs. Chromatin fragments were prepared with EZ-ZymeTM Chromatin Prep Kit (Cat.

17-375). Then CHIP assay was performed using Magna CHIPTM G chromatin

immunoprecipitation Kits (Cat. 17409). After DNA purification, IL-8 promoter activation was

examined by SYBR Green RT-PCR. CHIP-seq was analyzed using a pipeline of bowtie2[24],

macs callpeak[25], samtools[26], bedtools[27], and deeptools[28]. H4K5 peaks in CXCL8 were

visualized using IGV tools.

RNA-Seq Analysis

Total RNA was extracted from Lx2 cells treated with Medium, HIV-1$_{BaL}$ virus (MOI=0.1) and/or

LTA (1ug/ml) (n=3 for each). RNA quality was assessed by a Nano Drop 2000 Bioanalyzer, and

RNA-Seq was performed by Mount Sinai NGS core facility. RNA-Seq analysis was performed

using a pipeline of STAR[29], htseq[30], Deseq2[31] with batch adjustment, and hg19 annotation.

Pathway analysis of RNA-seq was conducted using Cytoscape with ClueGO 14 and R

packages: clusterProfiler[32], enrichplot[33], and Weighted Gene Co-Expression Network Analysis

(WGCNA)[34].

Western Blot

Tissue proteins from livers tissue samples from HIV non-infected controls and HIV-1 mono-

infected individuals were purified with T-PER Tissue protein Extraction Reagent (Thermo , Cat

No:78510). Equal amounts of protein were loaded in each lane for NuPAGE 4-12% Bis-tris gel

and then transferred to polyvinylidene difluoride membranes. The membranes were washed

with blotting buffer (1× PBS containing 0.1% Tween20) and then blocked for 60 min in blotting

buffer containing 10% low-fat powdered milk. Membranes were washed 3 times with blotting

buffer, incubated at 4^0 C overnight with primary antibody (1:1000) containing 5% lowfat

powdered milk, and incubated with secondary antibody (1:1000) at room temperature for 60 min.

Target proteins were developed with TMB 1-component membrane solution (SeraCare Life

Sciences Inc., Milford, MA). α-SMA (NOVUS, Cat No. NBP 1-45433 and NB100-56565),

Collagen I (NOVUS, Cat No: NBP 2-12446), MMP9 was purchased from Navius Bio. GAPDH

was purchased from Santa Cruz Biotechnology (Santa Cruz, Cat No. sc-66163). The blots were

detected with Amersham Imager 600 system. The relative expression of proteins was

normalized to beta-actin and analyzed using Image J.

Statistical Analysis

Data are presented as means ± standard error (SEM). Statistical analysis was evaluated by

GraphPad Prism 5.0 (La Jolla, CA). Comparisons between two groups were made using

parametric (Student's *t* test) or nonparametric (Wilcoxon signed ranks test) statistical tests, as

appropriate. One-way ANOVA with Kruskal-Wallis test was used to compare multiple

groups. For immunostaining, co-localization was analyzed by Fiji image J software (NIH,

Bethesda, MD) and presented using Pearson's R value. P values of <0.05 were considered

statistically significant and 0.05-0.09 as trending towards significance. Unless otherwise

specified, non-significant results indicate $p \geq 0.1$.

Results

HIV-1$_{BaL}$ exposition Increases IL-8 Response of pHSCs to LTA Stimulation

To evaluate the effect of lipoteichoic acid (LTA) stimulation on primary human hepatic stellate

cells (pHSCs), three pHSCs lines were isolated from human liver tissues without HIV-1 infection

(control pHSCs). Control pHSCs were stimulated with LTA, and the secretion of pro-

inflammatory cytokines IL-6, IL-8, IL-10, and TNF-α was measured both before and post HIV-

1$_{BaL}$ exposition for 48 hours. No significant induction of IL-6, TNF-α, or IL-10 was observed

following LTA treatment of control pHSCs, either with or without HIV-1$_{BaL}$ exposure

(Supplemental Figure 1). However, 48 hours post-HIV-1$_{BaL}$ exposure, LTA stimulation

significantly increased the release of IL-8 from control pHSCs (Figure 1A), compared to control

pHSCs stimulated with LTA alone.

To validate these findings, IL-8 secretion was measured following LTA stimulation in pHSCs

isolated from HIV-1-infected liver tissues (HIV$^+$ pHSCs). Two out of three HIV+ pHSCs exhibited

significantly increased IL-8 secretion upon LTA stimulation (Figure 1B). Consistent with these

findings, when Lx2 cells were co-stimulated with LTA and HIV-1$_{BaL}$, a synergistic increase in IL-8

secretion was observed (Figure 1C).

HIV-1$_{BaL}$ infection in Lx2 cells could be effectively suppressed by the C-C chemokine receptor

type 5 (CCR5) antagonist Maraviroc, leading to a subsequent reduction in p24 levels in cell

supernatant (Figure 1D). Also, maraviroc pre-treatment significantly reduced the enhanced IL-8

response to LTA stimulation in HIV-1$_{BaL}$ exposition Lx2 cells (Figure 1E).

To explore the mechanism underlying IL-8 secretion from HSCs stimulated by LTA, we

assessed cell surface receptor expression, including IL-8 receptors (C-X-C motif chemokine

receptors, CXCR1 and CXCR2) and the LTA ligand receptor (Toll-Like Receptor 2, TLR2) by

flow cytometry before and after control pHSCs stimulated with gp120, an HIV-1 envelope

protein. Only CXCR1 expression was upregulated following gp120 stimulation, whereas CXCR2

and TLR2 expression remained unchanged (Figure 2A). An IL-8 inhibitor failed to suppress IL-8

secretion in response to LTA stimulation (Figure 2B). Furthermore, treatment of Lx2 cells with

an IL-8 receptor inhibitor or a TLR2 inhibitor did not reverse the HIV-1$_{BaL}$-enhanced IL-8

secretion upon LTA stimulation (Figure 2C).

Collectively, these findings suggest that HIV-1 infection may sensitize the primary human HSCs

to LTA stimulation, promoting increased IL-8 secretion through a pathway independent of a IL8

receptor/TLR2 signaling.

IL-8 Expression is Elevated in HIV-1-Infected Human Liver Tissue

To further validate these findings, we analyzed the mRNA expression levels of pro-inflammatory

and fibrogenesis markers in liver tissues from PLWH. Liver tissues from PLWH exhibited

increased mRNA expression of IL-8, TGF-β and α-SMA compared to liver tissues from HIV-1

negative individuals (Figure 3A). Similarly, α-SMA protein expression was elevated in HIV-1-

infected liver tissues compared to controls (Figure 3B). Immunohistochemical (IHC) staining

analysis confirmed increased hepatic IL-8 expression in HIV-1 infected liver tissues compared to

HIV-1-negative liver tissues (Figure 3C). Moreover, ELISA data using serum from HIV[+]

individuals showed that individuals with HIV-1 viremia and liver fibrosis had significantly higher

IL-8 levels than those of aviremic individuals and viremic individuals without liver fibrosis.

Although there was no difference in serum LTA levels between individuals stratified by HIV-1

viremia and liver fibrosis, individuals with both HIV-1 viremia and liver fibrosis exhibited

markedly higher serum LTA concentrations (mean = 138.81 pg/ml), compared to those with

either liver fibrosis only (mean = 96.44 pg/ml) or those negative for both HIV-1 viremia and liver

fibrosis (mean = 19.28 pg/ml) (Figure 3D). These findings reinforce the role of IL-8 in liver

fibrosis associated with HIV-1 infection.

To investigate whether LTA and/or IL-8 have direct pro-fibrogenic effects, we treated Lx2 cells

with varying doses of LTA and IL-8. RT-PCR results revealed that LTA stimulation increased IL-

8 gene expression with a dose-dependent manner and without significantly affecting other pro-

fibrogenic genes in Lx2 cells (Figure 4A). Treatment with IL-8 repeatedly increased the protein

levels of pro-fibrogenic markers α-SMA and Collagen I in Lx2 cells (Figure 4B).

Collectively, these results underscore the potential pathological relevance of IL-8 response to

LTA in hepatic fibrosis in the context of HIV-1 infection.

RNA-seq Suggests Post-translational Regulation of Hepatic IL-8 Response to LTA in HIV-

1 Infection

To clarify how HIV-1 exposition primes HSCs for IL-8 response upon LTA exposition, we

conducted RNA sequencing (RNA-seq) analysis of Lx2 cells under three conditions: untreated,

exposed to HIV-1$_{BaL}$ alone, and co-exposed to HIV-1$_{BaL}$ plus LTA (Figure 5). Although statistical

significance was not reached in differential gene expression analysis (Figure 5A-B), IL-8 trended

higher in cells co-stimulated with HIV-1$_{BaL}$ and LTA compared to untreated Lx2 cells, or Lx2 cells

exposed with HIV-1$_{BaL}$ alone (p=0.059) (Figure 5C).

using WGCNA analysis, differentially expressed genes (DEGs) correlated with IL-8 (correlation

coefficient >0.2, p≤0.05) were analyzed by Cytoscape using ClueGO, revealing enrichment of

the tRNA aminoacylation (Figure 5D-E). Notably, Sirtuin 4 (SIRT4) was upregulated (Figure 5F),

whereas Sirtuin3 and the other aminoacylation related genes were relatively reduced. Given the

role of SIRT3 and SIRT4 in regulating protein acetylation and deacetylation under the conditions

of oxidative stress and inflammation[35], we further check other acetylation related genes. RNA-

seq analysis identified that genes associated with histone H4 acetylation and deacetylation—

such as HDAC2 and MORFL2[36,37]—were significantly regulated by the combined exposed to

HIV-1$_{BaL}$ and LTA (Figure 5G), indicating a histone acetylation potential involved in enhanced

expression of IL-8 in HIV-1 exposed Lx2 cells [35].

H4K5 Acetylation is Essential for IL-8 Transactivation in HIV-1 exposed HSCs

To investigate the role of histone acetylation in the HIV-1-mediated IL-8 response to LTA in

HSCs, we assessed IL-8 response in the presence of HDAC inhibitor, Trichostatin A (TSA). TSA

significantly increased IL-8 secretion in response to LTA stimulation in both normal HSCs

(Figure 6A) and HIV-1+ pHSCs (Figure 6B), specifically without affecting the production of IL-1β,

TNF-α, IL-6, or IL-10 (Supplemental Figure 1). Furthermore, HIV-1$_{BaL}$ infection increased total

acetylation levels in Lx2 cells (Supplemental Figure 2). Among the histone acetylation measured,

H4K5 acetylation levels were significantly elevated (Figure 6C), which was confirmed by

western blot results (Figure 6D). Additionally, no significant changes were observed in H3

acetylation (Supplemental Figure 3A-D).

CHIP-RT-PCR analysis demonstrated H4K5 acetylation at the IL-8 promoter in HIV-1 infected

Lx2 cells, with enrichment at NFκB, AP-1, and NF-IL16 binding sites (Figure 6E-F). In line with

this, CHIP-seq peaks in NF-kB and AP-1 binding motifs near IL-8(CXCL8) transcriptional start

sites confirmed these observations (Supplementary Figure 4A- B).

Collectively, these results show that H4K5 acetylation at the IL-8 promoter is crucial for IL-8

transactivation in response to LTA stimulation in HIV-1 infected hepatic stellate cells.

Discussion

End-stage liver disease is a leading cause of mortality in HIV-infected individuals, with several

studies confirming a positive correlation between HIV viremia and advanced fibrosis in HIV

mono-infection[13-16]. Chronic HIV infection can disrupt host immune homeostasis and accelerates

liver fibrosis, yet the mechanisms by which HIV-mediated inflammation contributes to hepatic

fibrogenesis remain unclear. Our *ex vivo* and *in vitro* analyses suggest that HIV-1-exposition of

hepatic stellate cells (HSCs) amplify their IL-8 response to a gram-positive microbial product

known as lipoteichoic acid (LTA). We identified histone acetylation-mediated transactivation of

IL-8 in HSCs as a key mechanism in this process.

Elevated serum IL-8 levels (≥46.5 pg/ml) are strongly associated with liver-related mortality and

liver transplantation[38,39]. While studies have shown increased IL-8 levels in HIV mono-infection[40],

the pathological role of IL-8 in HIV associated liver fibrosis remains unclear. Serological studies

reported no correlation between IL-8 levels and liver fibrosis in HIV/HCV co-infection, whereas

other studies suggest that elevated IL-8 can predict 3-year risk of fibrosis in HIV/HCV co-

infection[41,42]. In our cohort of HIV-1 positive individuals without HCV co-infection, serum IL-8

levels were significantly higher in viremic individuals with advanced fibrosis compared to non-

viremic individuals. Serum IL-8 levels differed by viremic status only among HIV-1+ individuals

with advanced fibrosis, suggesting a direct correlation between IL-8 expression and HIV-1

related pathogenesis during advanced fibrosis. Immunohistochemical analysis of HIV-1-infected

liver tissues confirmed increased hepatic IL-8 expression alongside enhanced pro-fibrogenic

gene expression (α-SMA and collagen I). Furthermore, *in vitro* analysis of stellate cell lines

demonstrated that IL-8 contributes directly to liver fibrosis by inducing pro-fibrogenic gene

expression in HSCs.

Similarly, studies on TNF-α pre-incubated HSCs suggest that IL-8 serves as a marker of acute

inflammatory response to liver injury by activated HSCs[43,44]. Moreover, when HSCs were primed

with TNF-alpha and IL-1, subsequent LTA stimulation significantly induced IL-8 production

through enhanced NF-kB activation [45]. In our study, without pre-treatment with TNF-α and IL-1β,

LTA exposure still led to increased IL-8 secretion from HSCs, an effect amplified in the

presence of HIV-1. Importantly, no induction of TNF-α and IL-1β was observed in HIV-1

exposed HSCs, underscoring the role of HIV in modulating HSCs' response to gram-positive

microbial products. Although the previous study have suggested that interactions among HIV-1,

immune cells/mediators, and HSCs via TLR pathway[46], our data demonstrate that exaggerated

IL-8 response to LTA in HIV-1 infected HSCs is independent of IL-8 receptors and TLR2

mediated pathways. Notably, RNA-seq results of Lx2 cells co-exposed to HIV-1 and LTA

revealed no activation of the canonical NF-κB signaling pathway in IL-8 induction

(Supplementary Figure 5), implying the involvement of an alternative transcriptional mechanism

contributing to IL-8 induction.

Our further transcriptomic analysis uncovered a correlation between the aminoacylation

pathway and IL8 transactivation. RNA-seq analysis also identified that genes related to H4

acetylation were significantly modulated upon co-exposition to HIV-1 and LTA. IL-8 promoter

methylation and acetylation has been implicated in the post-translational regulation of IL-8

activation[47]. Moreover, IL-8-driven chronic inflammation in various pathological conditions can

be reversed via acetylation inhibition by Histone deacetylases (HDACs)[48]. Our ex vivo

experiments using TSA to induce histone acetylation in HSCs specifically increased IL-8

response to LTA stimulation irrespective of HIV-1 exposition, while no effects on the induction of

IL-6, IL1beta and TNF production. Examinations by using ELISA, CHIP-seq and ChIP-qPCR,

we identified increased histone acetylation, specifically H4K5 acetylation, at the IL-8 promoter

during HIV-1 infection. This H4K5 facilitates chromatin remodeling and recruitment of the three

most critical binding sites in the IL-8 promoter i.e., transcription factors AP-1, NF-IL-6, and NF-

kB[49]. These findings align with observations in other diseases, such as cystic fibrosis where

histone hyperacetylation through use of histone deacetylase (HDAC) inhibitors promotes IL-8

transcription[50-53].Similarly, histone hyperacetylation promotes IL-8 transcription in hepatitis B

virus (HBV) infection, inducing an amplified IL-8 response to HBV replication in HepG2 cells[54].

The unique role of histone acetylation in modulating HSCs responses to microbial products in

HIV-associated liver disease has not been previously explored and warrants further

investigation. Our study is the first to suggest a direct role of IL-8 in HIV-1-associated HSCs

activation and liver fibrosis.

Conclusion

In conclusion, our *ex vivo* and *in vitro* analyses highlight the pathological significance of LTA in

hepatic inflammation and fibrogenesis during chronic HIV-1 infection. Elevated serum LTA

levels in HIV-1 individuals with fibrosis and viremia, coupled with an enhanced IL-8 response to

LTA in HSCs under HIV-1 exposure, underscore the critical role of gram-positive microbial

products in HIV associated chronic liver disease. Mechanistic insights from transcriptomic

analysis revealed that HIV-1 infection enhances histone H4K5 acetylation, amplifying the IL-8

response to LTA stimulation in HSCs. Analysis of liver tissues from HIV-1+ individuals illustrated

this exaggerated pro-inflammatory response of HSCs to LTA as direct pro-fibrogenic effects via

hepatic IL-8. Our findings advance understanding of liver fibrosis pathogenesis in HIV by linking

gut-derived microbial products, HSC activation, and IL-8 secretion via histone acetylation,

presenting important pathological and therapeutic implications for HIV-associated liver fibrosis.

Acknowledgement

We appreciate the reagents support from NIH AIDS Reagents Program.

Author's Contribution

Dr. Lumin Zhang and Dr. Meena B. Bansal conceptualized and designed the experiments. Dr.

Lumin Zhang performed the experiments and conducted bioinformatics analyses. Dr. Naichaya

Chamroonkul prepared serum samples. Dr. Lumin Zhang, Dr. Rabab O. Ali, and Dr. Meena B.

Bansal analyzed the data and drafted the manuscript. Dr. Parissa Tabrizian, Dr. Myron

Schwartz, Dr. Ganesh Gunasekaran, Dr. Thomas Schiano, and Dr. Maria Isabel Fiel provided

clinical samples and associated clinical data. All authors contributed to manuscript revisions and

approved the final version.

Funding

This work was supported by research grant NIH R56DK92128 and R01DK108364 to MBB

Conflict of Interest

The authors declare no conflicts of interest.

Data availability

Processed data files are available at github.com/lumin2003/HIV_LTA_files/

Declarations

The human study has been approved by Mount Sinai IRB (GCO #06-0523 and #10-1211).

Consent for publication

No individual patient information in this study.

Competing interests

All the authors have no competing interests in this study.

References

441   References

444   **Uncategorized References**

445   1   Zhang, L. *et al.* Humoral immune responses to Streptococcus pneumoniae in the setting

446   of HIV-1 infection. *Vaccine* **33**, 4430-4436 (2015).

447   https://doi.org/10.1016/j.vaccine.2015.06.077

448   2   Sandler, N. G. & Douek, D. C. Microbial translocation in HIV infection: causes,

449   consequences and treatment opportunities. *Nat Rev Microbiol* **10**, 655-666 (2012).

450   https://doi.org/10.1038/nrmicro2848

451   3   Ipp, H., Zemlin, A. E., Erasmus, R. T. & Glashoff, R. H. Role of inflammation in HIV-1

452   disease progression and prognosis. *Crit Rev Clin Lab Sci* **51**, 98-111 (2014).

453   https://doi.org/10.3109/10408363.2013.865702

454   4   Ljungdahl, M. *et al.* Bacterial translocation in experimental shock is dependent on the

455   strains in the intestinal flora. *Scand J Gastroenterol* **35**, 389-397 (2000).

456   https://doi.org/10.1080/003655200750023958

457   5   Dillon, S. M., Frank, D. N. & Wilson, C. C. The gut microbiome and HIV-1 pathogenesis:

458   a two-way street. *AIDS* **30**, 2737-2751 (2016).

459   https://doi.org/10.1097/QAD.0000000000001289

6 Kaspar, M. B. & Sterling, R. K. Mechanisms of liver disease in patients infected with HIV.

BMJ Open Gastroenterol **4**, e000166 (2017). https://doi.org/10.1136/bmjgast-2017-

000166

7 Debes, J. D., Bohjanen, P. R. & Boonstra, A. Mechanisms of Accelerated Liver Fibrosis

Progression during HIV Infection. *J Clin Transl Hepatol* **4**, 328-335 (2016).

https://doi.org/10.14218/JCTH.2016.00034

8 Ginsburg, I. Role of lipoteichoic acid in infection and inflammation. *Lancet Infect Dis* **2**,

171-179 (2002). https://doi.org/10.1016/s1473-3099(02)00226-8

9 Weidenmaier, C. & Peschel, A. Teichoic acids and related cell-wall glycopolymers in

Gram-positive physiology and host interactions. *Nat Rev Microbiol* **6**, 276-287 (2008).

https://doi.org/10.1038/nrmicro1861

10 Theilacker, C. *et al.* Protection against Staphylococcus aureus by antibody to the

polyglycerolphosphate backbone of heterologous lipoteichoic acid. *J Infect Dis* **205**,

1076-1085 (2012). https://doi.org/10.1093/infdis/jis022

11 Hattor, Y., Kasai, K., Akimoto, K. & Thiemermann, C. Induction of NO synthesis by

lipoteichoic acid from Staphylococcus aureus in J774 macrophages: involvement of a

CD14-dependent pathway. *Biochem Biophys Res Commun* **233**, 375-379 (1997).

https://doi.org/10.1006/bbrc.1997.6462

12 Siewe, B., Keshavarzian, A., French, A., Demarais, P. & Landay, A. A role for TLR

signaling during B cell activation in antiretroviral-treated HIV individuals. *AIDS Res Hum*

Retroviruses **29**, 1353-1360 (2013). https://doi.org/10.1089/AID.2013.0115

13 Sanmartin, R. *et al.* Progression of liver fibrosis in HIV/hepatitis C virus-coinfected

individuals on antiretroviral therapy with early stages of liver fibrosis at baseline. *HIV*

Med **15**, 203-212 (2014). https://doi.org/10.1111/hiv.12105

14   Blackard, J. T. *et al.* HIV mono-infection is associated with FIB-4 - A noninvasive index

of liver fibrosis - in women. *Clin Infect Dis* **52**, 674-680 (2011).

https://doi.org/10.1093/cid/ciq199

15   Stabinski, L. *et al.* High prevalence of liver fibrosis associated with HIV infection: a study

in rural Rakai, Uganda. *Antivir Ther* **16**, 405-411 (2011). https://doi.org/10.3851/IMP1783

16   Tahiri, M. *et al.* Risk factors for liver fibrosis among human immunodeficiency virus

monoinfected patients using the FIB4 index in Morocco. *World J Hepatol* **5**, 584-588

(2013). https://doi.org/10.4254/wjh.v5.i10.584

17   Mohsen, A. H. *et al.* Impact of human immunodeficiency virus (HIV) infection on the

progression of liver fibrosis in hepatitis C virus infected patients. *Gut* **52**, 1035-1040

(2003). https://doi.org/10.1136/gut.52.7.1035

18   Koyama, Y. & Brenner, D. A. Liver inflammation and fibrosis. *J Clin Invest* **127**, 55-64

(2017). https://doi.org/10.1172/JCI88881

19   Mosoian, A. *et al.* Frontline Science: HIV infection of Kupffer cells results in an amplified

proinflammatory response to LPS. *J Leukoc Biol* **101**, 1083-1090 (2017).

https://doi.org/10.1189/jlb.3HI0516-242R

20   Zhang, L. *et al.* HIV infection modulates IL-1beta response to LPS stimulation through a

TLR4-NLRP3 pathway in human liver macrophages. *J Leukoc Biol* **105**, 783-795 (2019).

https://doi.org/10.1002/JLB.4A1018-381R

21   T, L. Analysis of Complex Survey Samples. *journal of Statistical Software*, 1-19 (2004).

22   Panos, A. & Mavridis, D. TableOne: an online web application and R package for

summarising and visualising data. *Evid Based Ment Health* **23**, 127-130 (2020).

https://doi.org/10.1136/ebmental-2020-300162

23   Alabraba, E. B. *et al.* A new approach to isolation and culture of human Kupffer cells. *J*

Immunol Methods **326**, 139-144 (2007). https://doi.org/10.1016/j.jim.2007.06.014

24   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*

9, 357-359 (2012). https://doi.org/10.1038/nmeth.1923

25   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137

(2008). https://doi.org/10.1186/gb-2008-9-9-r137

26   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,

2078-2079 (2009). https://doi.org/10.1093/bioinformatics/btp352

27   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841-842 (2010).

https://doi.org/10.1093/bioinformatics/btq033

28   Ramirez, F., Dundar, F., Diehl, S., Gruning, B. A. & Manke, T. deepTools: a flexible

platform for exploring deep-sequencing data. *Nucleic Acids Res* **42**, W187-191 (2014).

https://doi.org/10.1093/nar/gku365

29   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21

(2013). https://doi.org/10.1093/bioinformatics/bts635

30   Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-

throughput sequencing data. *Bioinformatics* **31**, 166-169 (2015).

https://doi.org/10.1093/bioinformatics/btu638

31   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

https://doi.org/10.1186/s13059-014-0550-8

32   Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for comparing

biological themes among gene clusters. *OMICS* **16**, 284-287 (2012).

https://doi.org/10.1089/omi.2011.0118

33   Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.

Innovation (Camb) **2**, 100141 (2021). https://doi.org/10.1016/j.xinn.2021.100141

34   Gennady Korotkevich, V. S., Alexey Sergushichev. Fast gene set enrichment analysis.

bioRaxiv (2021).

35   Zou, Y. *et al.* The regulatory roles of aminoacyl-tRNA synthetase in cardiovascular

disease. *Mol Ther Nucleic Acids* **25**, 372-387 (2021).

https://doi.org/10.1016/j.omtn.2021.06.003

36   Kooblall, K. G. *et al.* miR-3156-5p is downregulated in serum of MEN1 patients and

regulates expression of MORF4L2. *Endocr Relat Cancer* **29**, 557-568 (2022).

https://doi.org/10.1530/ERC-22-0045

37   Becht, D. C. *et al.* MORF and MOZ acetyltransferases target unmethylated CpG islands

through the winged helix domain. *Nat Commun* **14**, 697 (2023).

https://doi.org/10.1038/s41467-023-36368-5

38   Zimmermann, H. W. *et al.* Interleukin-8 is activated in patients with chronic liver diseases

and associated with hepatic macrophage accumulation in human liver fibrosis. *PLoS*

One **6**, e21381 (2011). https://doi.org/10.1371/journal.pone.0021381

39   Umemura, T. *et al.* Association between serum soluble CD14 and IL-8 levels and clinical

outcome in primary biliary cholangitis. *Liver Int* **37**, 897-905 (2017).

https://doi.org/10.1111/liv.13316

40   Ellwanger, J. H. *et al.* Increased IL-8 levels in HIV-infected individuals who initiated ART

with CD4(+) T cell counts <350 cells/mm(3) - A potential hallmark of chronic

inflammation. *Microbes Infect* **22**, 474-480 (2020).

https://doi.org/10.1016/j.micinf.2020.05.019

41   Chen, X. *et al.* Plasma Inflammatory Biomarkers Associated with Advanced Liver

Fibrosis in HIV-HCV-Coinfected Individuals. *Int J Environ Res Public Health* **17** (2020).

https://doi.org/10.3390/ijerph17249474

42 Keating, S. M. *et al.* The effect of HIV infection and HCV viremia on inflammatory

mediators and hepatic injury-The Women's Interagency HIV Study. *PLoS One* **12**,

e0181004 (2017). https://doi.org/10.1371/journal.pone.0181004

43 Lang, A. *et al.* Somatostatin inhibits pro-inflammatory cytokine secretion from rat hepatic

stellate cells. *Liver Int* **25**, 808-816 (2005). https://doi.org/10.1111/j.1478-

3231.2005.01057.x

44 Schwabe, R. F., Schnabl, B., Kweon, Y. O. & Brenner, D. A. CD40 activates NF-kappa B

and c-Jun N-terminal kinase and enhances chemokine secretion on activated human

hepatic stellate cells. *J Immunol* **166**, 6812-6819 (2001).

https://doi.org/10.4049/jimmunol.166.11.6812

45 Paik, Y. H. *et al.* Hepatic stellate cells primed with cytokines upregulate inflammation in

response to peptidoglycan or lipoteichoic acid. *Lab Invest* **86**, 676-686 (2006).

https://doi.org/10.1038/labinvest.3700422

46 Del Corno, M. *et al.* HIV-1 gp120 signaling through TLR4 modulates innate immune

activation in human macrophages and the biology of hepatic stellate cells. *J Leukoc Biol*

100, 599-606 (2016). https://doi.org/10.1189/jlb.4A1215-534R

47 Takahashi, A., de Andres, M. C., Hashimoto, K., Itoi, E. & Oreffo, R. O. Epigenetic

regulation of interleukin-8, an inflammatory chemokine, in osteoarthritis. *Osteoarthritis*

Cartilage **23**, 1946-1954 (2015). https://doi.org/10.1016/j.joca.2015.02.168

48 Gatla, H. R. *et al.* Regulation of Chemokines and Cytokines by Histone Deacetylases

and an Update on Histone Decetylase Inhibitors in Human Diseases. *Int J Mol Sci* **20**

(2019). https://doi.org/10.3390/ijms20051110

49 Roebuck, K. A. Regulation of interleukin-8 gene expression. *J Interferon Cytokine Res*

19, 429-438 (1999). https://doi.org/10.1089/107999099313866

50 Ashburner, B. P., Westerheide, S. D. & Baldwin, A. S., Jr. The p65 (RelA) subunit of NF-

kappaB interacts with the histone deacetylase (HDAC) corepressors HDAC1 and

HDAC2 to negatively regulate gene expression. *Mol Cell Biol* **21**, 7065-7077 (2001).

https://doi.org/10.1128/MCB.21.20.7065-7077.2001

51   Vanden Berghe, W., De Bosscher, K., Boone, E., Plaisance, S. & Haegeman, G. The

nuclear factor-kappaB engages CBP/p300 and histone acetyltransferase activity for

transcriptional activation of the interleukin-6 gene promoter. *J Biol Chem* **274**, 32091-

32098 (1999). https://doi.org/10.1074/jbc.274.45.32091

52   Jundi, K. & Greene, C. M. Transcription of Interleukin-8: How Altered Regulation Can

Affect Cystic Fibrosis Lung Disease. *Biomolecules* **5**, 1386-1398 (2015).

https://doi.org/10.3390/biom5031386

53   Barnes, P. J., Adcock, I. M. & Ito, K. Histone acetylation and deacetylation: importance

in inflammatory lung diseases. *Eur Respir J* **25**, 552-563 (2005).

https://doi.org/10.1183/09031936.05.00117504

54   Pollicino, T. *et al.* Hepatitis B virus (HBV) induces the expression of interleukin-8 that in

turn reduces HBV sensitivity to interferon-alpha. *Virology* **444**, 317-328 (2013).

https://doi.org/10.1016/j.virol.2013.06.028

Figure Legends

Figure 1: HIV-1 exposition accentuates the IL-8 response to LTA stimulation in primary

isolated HSCs. A) Primary human hepatic stellate cells (pHSCs) were isolated from HIV-

uninfected liver tissues (n=3) and treated with LTA (1□µg/ml) either before or after exposition

with HIV-1BaL (MOI=0.1) for 48 hours. Supernatants were collected 24 hours after LTA

stimulation, and IL-8 levels were measured by ELISA. B) pHSCs from HIV-1-infected liver

tissues (n=6) were stimulated with LTA (1□µg/ml) for 24 hours, and IL-8 levels were quantified

by ELISA. C) Lx2 cells were treated with LTA alone or HIV-1 plus LTA. Supernatants were

collected and IL-8 was measured by ELISA (n=6). D) HIV-1 p24 antigen was quantified by

ELISA in supernatants from HIV-1BaL-exposed Lx2 cells (MOI=0.1) with or without Maraviroc

(1 mM) treatment (n=9). E) IL-8 levels were measured by ELISA in HIV-1BaL-exposed Lx2

cells treated with or without Maraviroc (1 mM) (n=6). HIV-1$_{BaL}$

Figure 2: IL-8 response of HSCs to LTA is independent of the IL-8 and TLR receptor

pathways. A) IL-8 receptor was analyzed by Flow cytometry before and after gp120 stimulation

(100ng/ml). B) Lx2 cells were stimulated with LTA(1 μg/ml) for 24 hours with or without an IL-8

receptor inhibitor. Supernatants were analyzed for IL-8 levels by ELISA (n=3). C) Lx2 cells were

exposed with HIV-1BaL for 48 hours and then treated with either an IL-8 receptor inhibitor or a

TLR2 inhibitor. Supernatants were analyzed for IL-8 levels by ELISA (n=3).

Figure 3: Hepatic IL-8 expression is elevated in HIV-1 infected liver tissues alongside

enhanced expression of pro-fibrogenic genes. A) Total RNA was extracted from HIV-

uninfected (n=7) and HIV-1-infected (n=7) liver tissues. mRNA expression of IL-8, α-SMA,

collagen I, and TGF-β was quantified by RT-PCR. B) Protein expression of α-SMA and collagen

I was examined by western blot; GAPDH served as a loading control. Band intensity was

quantified using ImageJ (n>3). C) IL-8 expression was evaluated by immunohistochemistry in

HIV-1 infected and HIV-1 non-infected tissues (n=32). The results were quantified by image J D)

Serum IL-8 and LTA concentration were measured in duplicate by ELISA and compared across

groups stratified by HIV-1 viremia and fibrosis status ($n^{HIV-1\ viremia-} = 9$, $n^{HIV-1\ viremia-\ Fib+} = 9$, n^{HIV-1}

$^{viremia+Fib+} = 10$).

Figure 4: IL-8 increases the expression of pro-fibrogenic genes in hepatic stellate cell

lines. A) Lx2 cells were treated with increasing concentrations of LTA for 24 hours. mRNA

expression of pro-fibrotic and inflammatory genes was quantified by SYBR Green RT-PCR.

B) Lx2 cells were repeatedly treated with IL-8 (200 pg/ml) or PBS for 3times. Protein

expression of α-SMA and collagen I was analyzed by western blot and normalized to GAPDH.

Band intensities were quantified using ImageJ (n=3).

Figure 5: RNA-seq analysis Suggests a link between histone acetylation and IL-8

Activation following HIV-1 and LTA Co-Stimulation in Lx2 Cells. A-B) Volcano plots

showing differentially expressed genes (DEGs) between control, HIV-1$_{BaL}$-exposed, and HIV-

1$_{BaL}$ plus LTA-treated Lx2 cells (p < 0.05, log2FC > 1.0). Blue and red dots represent up-

and down- regulated genes, respectively. C) Heatmap analysis of inflammatory genes across

treatment groups. D) ClueGO pathway enrichment of genes positively correlated with IL-8

expression (correlation coefficient > 0.2), highlighting the tRNA aminoacylation pathway. E)

Cytoscape-based network reconstruction of the tRNA aminoacylation pathway, showing CXCL8

(IL-8) enrichment. F) Heatmap of tRNA aminoacylation gene expression across the three

conditions. G) Heatmap analysis of Acetylation related genes across the three conditions.

Figure 6: H4K5 Acetylation is Essential for IL-8 Response to LTA stimulation in HIV-1

infected HSCs. A) Two pHSCs isolated from HIV-1 negative individuals were stimulated with

LTA with or without HDAC inhibitor (TSA) for 24hrs, IL-8 in supernatant was measured by

ELISA (n=6). B) Two pHSCs isolated from HIV-1 infected individuals were stimulated with LTA

with or without HDAC inhibitor (TSA) for 24hrs, IL-8 in supernatant was measured by ELISA

(n=6). C) H4K5 acetylation protein was measured by ELISA in both pHSCs (n=4) and Lx2 cell

lines (n=8) before and after HIV-1$_{BaL}$ exposition. D) H4K5 acetylation protein was examined by

western blotting and normalized to GAPDH. E) Binding motif analysis of IL-8 promoter. F)

Chromatin immunoprecipitation (ChIP) was performed using anti-H4K5 antibody. DNA was

purified and analyzed by SYBR RT-PCR for enrichment of NF-κB, AP-1, and NF-IL6 binding

motifs in the IL-8 promoter.

Tables

Table 1: Primers:

All primer sequences.

h-a-SMA R: 5'-AGGCACCCCTGAACCCCAA-3'
h-a-SMA R: 5'-CAGCACCGCCTGGATAGCC-3'
h-Collagen I F: 5'-GGCTTCCCTGGTCTTCCTGG-3'
h-Collagen I R: 5'-CCAGGGGGTCCAGCCAAT-3'
h-PDGF F: 5'-CCAGAAGCCATCAGCAGCAAG-3'
h-PDGF R: 5'-AGGCCCTGAGAGATCTGTGG-3'
h-MMP2 F: 5'- CCCCAAGCTCATCGCAGAT-3'
h-MMP2 R: 5'- GGTCCACGACGGCATCC-3'
h-TIMP1 F: 5'-TGACATCCGGTTCGTCTACA-3'
h-TIMP1 R: 5'-TGATGTGCAAGAGTCCATCC-3'
h-TIMP2 F: 5'- AAGCGGTCAGTGAGAAGGAA-3'
h-TIMP2 R: 5'-GGGGGCCGTGTAGATAAACT-3'
h-TGF-beta F: 5'-CAAGGGCTACCATGCCAACT-3'
h-TGF-beta R: 5'-AGGGCCAGGACCTTGCTG-3'
h-TGF-beta1 Receptor F: 5'-CGACTGCCAAATGAAGAGGACC-3'
h-TGF-beta1 Receptor R: 5'-AAACCTGAGCCAGAACCTGACG-3'
h-IL-8 F: 5'-TAGGACAAGAGCCAGGAAGA-3'
h-IL-8R: 5'-CTTCCTGATTTCTGCAGCTCT-3'
h-GAPDH F: 5'-CAATGACCCCTTCATTGACC-3'
h-GAPDH R: 5'- GACAAGCTTCCCGTTCTCAG -3'

667 **Table 2:** Characteristics of blood donors

	Overall	0	1	p test
n	28	11	17	
Sigfibrosis = 1 (%)	17.0 (60.7)	0.0 (0.0)	17.0 (100.0)	<0.001
gender = 2 (%)	5.0 (17.9)	2.0 (18.2)	3.0 (17.6)	0.972
Ethinic (%)				0.368
AA	7.0 (25.0)	2.0 (18.2)	5.0 (29.4)	
AS	1.0 (3.6)	0.0 (0.0)	1.0 (5.9)	
HS	13.0 (46.4)	4.0 (36.4)	9.0 (52.9)	
O	3.0 (10.7)	2.0 (18.2)	1.0 (5.9)	
W	4.0 (14.3)	3.0 (27.3)	1.0 (5.9)	
age (mean (SD))	52.89 (9.52)	53.55 (4.27)	52.47 (11.88)	0.732
bmi (mean (SD))	30.55 (6.62)	29.21 (6.42)	31.41 (6.79)	0.384

alcohol = 1 (%)	9.0 (32.1)	3.0 (27.3)	6.0 (35.3)	0.666
duradiag (mean (SD))	19.82 (8.69)	19.20 (8.94)	20.33 (8.85)	0.762
viremia = 1 (%)	19.0 (67.9)	11.0 (100.0)	8.0 (47.1)	0.005
cd4 (mean (SD))	649.18 (324.81)	627.00 (348.98)	663.53 (318.36)	0.777
fibroscan (mean (SD))	9.33 (5.21)	5.50 (1.23)	11.81 (5.31)	<0.001
plt (mean (SD))	197.71 (63.29)	200.73 (58.45)	195.76 (67.93)	0.835
alt (mean (SD))	30.46 (20.36)	23.00 (9.14)	35.29 (24.17)	0.065
ALTab = 1 (%)	10.0 (35.7)	2.0 (18.2)	8.0 (47.1)	0.138
alp (mean (SD))	91.18 (28.40)	82.55 (34.78)	96.76 (22.82)	0.23
alb (mean (SD))	3.86 (0.82)	4.01 (0.29)	3.77 (1.02)	0.366
fbs (mean (SD))	106.61 (37.03)	90.45 (22.32)	117.06 (41.31)	0.034
chol (mean (SD))	167.71 (41.52)	182.73 (48.59)	158.00 (34.33)	0.144
tg (mean (SD))	167.79 (133.44)	170.91 (122.65)	165.76 (143.64)	0.918
hdl (mean (SD))	41.78 (9.67)	42.20 (9.37)	41.53 (10.12)	0.86
lowHDL (mean (SD))	0.56 (0.51)	0.50 (0.53)	0.59 (0.51)	0.666
ldl (mean (SD))	94.18 (34.95)	110.04 (45.12)	83.92 (22.46)	0.079
CAP (mean (SD))	283.18 (61.62)	261.00 (66.65)	297.53 (55.45)	0.134
fattyliver = 1 (%)	23.0 (82.1)	8.0 (72.7)	15.0 (88.2)	0.313
AST (mean (SD))	28.38 (12.35)	25.27 (10.58)	30.67 (13.39)	0.253
HCV (mean (SD))	0.46 (0.51)	0.45 (0.52)	0.47 (0.51)	0.936
HBV (mean (SD))	0.04 (0.19)	0.00 (0.00)	0.06 (0.24)	0.321
HSI (mean (SD))	37.48 (13.26)	37.87 (7.23)	37.21 (16.43)	0.886
FIB4 (mean (SD))	1.67 (1.24)	1.59 (0.86)	1.74 (1.48)	0.738

669 Definition:

670 Active alcohol consumption = drinking more than 10 g of ethanol per day in females and 20 g

671 per day in males

672 Liver steatosis = evidence of CAP more than 248 dB/m

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A)

Lx2 cells

IL8 in supernatant (pg/ml)

p=0.008

Medium LTA TSA-LTA

pHSCs1

IL8 in supernatant (pg/ml)

p=0.0022

Medium LTA TSA-LTA

B)

HIV1⁺ pHSCs1

IL8 in supernatant (pg/ml)

p=0.004

Medium LTA TSA

HIV1⁺ pHSCs2

IL8 in supernatant (pg/ml)

p=0.0022

p=0.02

Medium LTA TSA

C)

p=0.031

Actylation of H4K5(ng/uM)

pHSCs   pHSC2+HIV1$_{BaL}$ Lx2   Lx2+HIV1$_{BaL}$

D)

H4K5

CTL BaL

GAPDH

E)

TGGAACCCAGATGCTCTTTCCACTAAGCATACAACTTTCCATTAGATAAACACCTCCCTCCCACCCCAACCA
AGCAGCTCCAGTGCACCACTTTCTGGAGCATAAACATACCTTAACTTTACAACTTGAGTGGCCTTGAATAC
TGTTCCTATCTGGAATGTGCTGTTCTCTTTCATCTTCCTCTATTGAAGCCCTCCTATTCCTCAATGCCTTG
CTCCAACTGCCTTTGGAAGATTCTGCTCTTATGCCTCCACTGGAATTAATGTCTTAGTACCACTTGTCTAT
TCTGCTATATAGTCAGTCCTACATTGCTTTCTTCTTCTGATAGACCAAACTCTTTAAGGACAAGTACCTA
GTCTTATCTATTTCTAGATCCCACATTACTTCAGAAAGTTACTCCATAAATGTTTGTGGAACTGATTTCT
ATGTGAAGCACATGTGCCCCTTCACTCTGTTAACATGCATTAGAAAACTAAATCTTTTGAAAAGTTGTAGT
ATGCCCCTAAGAGCAGTAACAGTTCCTAGAAACTCTCTAAAATGCTTAGAAAAAGATTTATTTTAAATTA
CCTCCCCAATAAAATGATTGGCTGGCTTATCTTCACCATCATGATAGCATCTGTAATTAACTGAAAAATAA
TAATTATGCCATTAAAAGAAAATCATCCATGATCTTGTTCTAACACCTGCCACTCTAGTACTATATCTGTC
ACATGGTACTATGATAAAGTTATCTAGAAATAAAAAAGCATACAATTGATAATTCACCAAATTGTGGAGCT
TCAGTATTTTAAATGTATATTAAAATTAAATTATTTTAAAGATCAAAGAAAACTTTCGTCATACTCCGTAT
TTGATAAGGAACAAATAGGAAGTGTGATGACTCAGGTTTGCCCTGAGGGGATGGGCCATCAGTTGCAAATC
GTGGAATTCCTCTGACATAATGAAAAGATGGGGTGCATAAGTTCTCTAGTAGGGTGATGATATAAAAAG
CACCGGAGCACTCCATAAGGCACAAACTTTCAGAGACAGCAGAGCACCAAGCTTCTAGGACAAGAGCCA
GGAAGAAACCACCGGAAGGAACCATCTCACTGTGTGTAAACAT

Red Bold :   NF-IL16:
Purple Bold : AP-1
Blue Bold :   NF-kB

F)

CHIP-RT-PCR

The relative level of H4K5 Acetylation

Input   NF-kB Sites   NF-IL16 Sites   AP-1 Sites