HIV-1 Amplifies IL-8 Response of Human Stellate Cells to Gram-Positive Microbial

Products Via H4K5 Histone Acetylation

Lumin Zhang[1]*, Rabab O. Ali[1]*, Naichaya Chamroonkul[5], Parissa Tabrizian[2], Myron Schwartz[2],

Ganesh Gunasekaran[2], Thomas Schiano[3], Maria Isabel Fiel[4], Steven C. Ward[4], Meena B.

Bansal[1]

[1]Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[2]Department of Surgery, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[3]Recanti-Miler Transplant Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[4]Department of Pathology, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[5]Gastroenterology and Hepatology Unit, Division of Internal Medicine, Faculty of Medicine, Prince of

Songkla University, Hatyai, Songkhla, Thailand

*The authors contribute equally to this work

Corresponding Author

Meena B. Bansal, MD, Icahn School of Medicine at Mount Sinai, 1425 Madison Avenue, Room

11-70; Box 1123, New York, NY 10029, E-mail: meena.bansal@mssm.edu, Fax: 212-849-2574,

Tel : 212-659-9510.

Financial support

This work was supported by research grant NIH R56DK92128 and R01DK108364 to MBB.

Abstract

Background: Patients living with human immunodeficiency virus 1 (PLWH) develop accelerated liver fibrosis, but the exact mechanism remains unknown. Activation of hepatic stellate cells (HSCs)—a cornerstone of fibrosis—is influenced by various factors, including viral infection, hepatocellular injury, chronic immune activation, gut barrier dysfunction, and microbial translocation. The role of gram-positive microbial products in human immunodeficiency virus 1 (HIV-1) infection–associated liver inflammation and fibrosis remains poorly understood. This study investigates the effect of lipoteichoic acid (LTA), a major gram-positive bacterial component, on HSCs in the context of HIV-1 infection. **Methods:** Human HSCs were isolated from liver tissues of both HIV-1–infected and uninfected individuals undergoing hepatic resection. Inflammatory responses of HSCs to LTA stimulation were measured ex vivo via ELISA before and after HIV-1$_{BaL}$ exposure. Western blotting, ChIP-qPCR and RNA-seq were used to reveal the mechanisms contributing to the IL-8 response to LTA stimulation and HIV-1$_{BaL}$ exposure in HSCs. **Results:** LTA modestly induced interleukin-8/CXCL8 (IL-8) production in HSCs, but this response was significantly heightened following HIV-1 exposure. Increased IL-8 levels were also observed in liver tissues from HIV-1–infected patients. In vitro, IL-8 treatment of HSCs elevated α-SMA and COL1A1 expression, implicating IL-8 in fibrosis progression in HIV-1 infection. Transcriptomic analysis pointed to histone acetylation as a key regulator of the IL-8 response of HSCs to LTA during HIV-1 infection. Supporting this, the histone deacetylase (HDAC) inhibitor Trichostatin A (TSA) further enhanced IL-8 expression in HIV-1–exposed HSCs. ChIP-qPCR confirmed that acetylation of histone H4K5 facilitated IL-8 promoter transactivation, sensitizing HSCs to LTA under HIV-1 influence. **Conclusions:** HIV-1 infection primes HSCs for an exaggerated response to LTA, driven by histone acetylation and resulting in elevated IL-8 production—potentially accelerating liver fibrosis in PLWH. Given the persistence of microbial translocation despite effective antiretroviral therapy, these findings highlight the need for targeted interventions to prevent or mitigate liver fibrosis in PLWH.

Abstract Word Count: 295

Keywords

Human Immunodeficiency Virus, LTA, Inflammatory response, Fibrosis, Hepatic Stellate Cells,

Microbial Translocation.

Abbreviations

Aspartate Aminotransferase (AST), Alanine Aminotransferase (ALT), Alkaline Phosphatase

(ALP), Albumin (Alb), Antiretroviral therapy (ART), Body Mass Index (BMI), Bovine serum

albumin (BSA), Controlled Attenuation Parameter (CAP), Chromatin immunoprecipitation with

quantitative PCR (ChIP-qPCR), Cycle threshold (CT), C-C chemokine receptor type 5 (CCR5),

C-X-C motif chemokine receptor 1 (CXCR), Differentially expressed genes (DEGs), Dulbecco's

Modified Eagle Medium (DMEM), Enzyme-linked immunosorbent assay (ELISA), Fasting Blood

Sugar (FBS), Fibrosis-4 Index (FIB-4), Hepatic stellate cells (HSC), Hepatitis B virus (HBV),

Hepatitis C virus (HCV), Hepatic Steatosis Index (HIS), High-Density Lipoprotein (HDL), Histone

deacetylase (HDAC), Human immunodeficiency virus 1 (HIV-1), Immunohistochemistry (IHC),

Interleukin-1β (IL-1β), Interleukin-8/CXCL8 (IL-8), Lipopolysaccharide (LPS), Lipoteichoic acid

(LTA), Liver Stiffness Measurement (LSM), Low-Density Lipoprotein (LDL), Metabolic

dysfunction-associated steatotic liver disease (MASLD), Multiplicity of infection (MOI),

Pathogen-associated molecular pattern (PAMP), Platelet count (Plt), Primary HSCs (pHSCs),

Patients living with human immunodeficiency virus 1 (PLWH), RNA sequencing (RNA-seq),

Sirtuin (SRT), Toll-Like Receptor (TLR), Total cholesterol level (Chol), Triglycerides (TG),

Trichostatin A (TSA), Transcription factor (TF), Tumor necrosis factor α (TNF-α), and Weighted

Gene Co-Expression Network Analysis (WGCNA).

Introduction

Human immunodeficiency virus 1 (HIV-1) infection is marked by profound depletion of CD4$^+$ T cells, leading to systemic immunosuppression. As a result, patients living with HIV-1 (PLWH) are more susceptible to opportunistic and community-acquired bacterial infections. For instance, the risk of *Streptococcus pneumoniae* infection is 100 times higher in HIV-1–infected individuals than in the general population[1]. Early during infection, the loss of gut-associated CD4$^+$ T cells compromises intestinal barrier integrity, facilitating the translocation of microbial products into systemic circulation[2]. Notably, this microbial translocation persists even in patients receiving antiretroviral therapy (ART), as ART fails to fully restore gut permeability—making microbial translocation a key driver of sustained immune activation in PLWH.

HIV-induced damage to the gastrointestinal mucosa leads to the translocation of both gram-positive and gram-negative microbial products from the gut lumen into the liver via the portal vein[3]. While gram-negative products typically translocate more efficiently following mucosal disruption[4], gram-positive bacteria also enter the circulation frequently[5], either through the same routes or secondary infections. Although extensive research has explored the effects of gram-negative components like lipopolysaccharide (LPS) on immune activation, liver injury, and fibrogenesis in HIV[6,7], the influence of gram-positive microbial products remains comparatively understudied. Lipoteichoic acid (LTA), a major component of the gram-positive bacterial cell walls, functions as a pathogen-associated molecular pattern (PAMP) capable of modulating host immunity[8-10]. Similar to LPS, LTA triggers CD14-dependent nitric oxide release in vascular smooth muscle cells, macrophages, and mononuclear cells[11]. Elevated serum LTA levels have been detected in PLWH, correlating with heightened autoantibody production and immune activation[12]. Given the liver's frontline exposure to gut-derived microbial products via the portal vein, the impact of LTA—particularly on hepatic stellate cells (HSCs)—in HIV-associated chronic liver disease remains insufficiently explored.

End-stage liver disease continues to be a major cause of mortality in HIV-positive patients

despite successful ART, highlighting ongoing hepatic dysfunction and progressive fibrosis[13].

Numerous studies have shown a positive correlation between circulating HIV RNA levels and

advanced fibrosis in HIV mono-infected patients[14-16]. Additionally, co-infection with hepatitis C

virus (HCV) accelerates fibrosis progression threefold compared to HCV mono-infection[17]. While

cross-talk between HSCs and liver macrophages is critical for HSC activation[18], HSCs are the

primary mediators of hepatic fibrogenesis. Our prior work[19,20] revealed that HIV-1 infection

enhances the sensitivity of liver macrophages to gram-negative microbial stimuli such as LPS,

promoting elevated production of pro-inflammatory cytokines including interleukin-8 (IL-

8/CXCL8). However, whether these cytokines directly stimulate HSC activation remains unclear.

In this study, we examined the inflammatory responses of primary human HSCs—isolated from

liver tissues of both HIV-1–infected and uninfected individuals —upon LTA stimulation. We

demonstrate that HIV-1 infection amplifies IL-8 production in response to LTA and that IL-8

exerts pro-fibrogenic effects in HIV-1–infected liver tissue. Furthermore, we identify histone

H4K5 acetylation as a mediator of IL-8 transactivation in HIV-1–exposed HSCs, offering

mechanistic insight into the epigenetic regulation of liver fibrosis in the context of HIV infection.

Materials and Methods

Ethics Statement

De-identified liver resection specimens from HIV-1–positive individuals (n = 7) with effectively

suppressed viral loads on ART were provided by Drs. Schwartz, Ganeskaran, and Fiel. Control

liver tissues (n = 10) were obtained from the Mount Sinai Biobank and included samples from

patients undergoing liver resection for colon cancer metastases (n = 9) or cholangiocarcinoma

(n = 1). All tissue samples were collected during liver transplantation or tumor resection

procedures and were processed as formalin-fixed, frozen, or fresh in Dulbecco's Modified Eagle

Medium (DMEM). The study was approved by the Mount Sinai Institutional Review Board (GCO

#06-0523 and #10-1211). Liver tissue samples from both HIV-1–infected and uninfected

individuals were used for IL-8 immunohistochemistry (IHC). In addition, blood samples were

collected from HIV-positive participants enrolled in a metabolic dysfunction-associated steatotic

liver disease (MASLD) study. Population characteristics (Table 2) were analyzed using the R

packages *survey*[21] and *tableone*[22].

Isolation of Primary Human Hepatic Stellate Cells

Primary HSCs (pHSCs) were isolated from surgical liver tissue specimens using collagenase

perfusion and tissue mincing. Cells were washed with PBS (4×) and centrifuged at 300 × g for

5 minutes. Following resuspension in DMEM, cells underwent density gradient centrifugation

with 25% and 50% Percoll (Pharmacia, Sweden) as described previously[23]. pHSCs were

enriched by adherence to tissue culture plates and cultured for two weeks. Purity was confirmed

via α-SMA immunostaining.

Tissue Immunohistochemistry

Liver tissue sections (prepared in duplicate) were rehydrated, processed for antigen retrieval,

and incubated overnight at 4°C with anti–human IL-8 antibody (NOVUS, Cat. NBP2-33819,

1:200 dilution) in a humidified chamber. IL-8 staining was detected using the ImmPress™ Duet

Double Staining Polymer Kit (Cat. MP-7724). Four random fields per section were imaged at

100× magnification using Zen software with an LSM 880 laser-scanning microscope. IL-8 signal

intensity was quantified using Fiji ImageJ (NIH, Bethesda, MD).

Flow Cytometry Analysis

Expression of CXCR1, CXCR2, and TLR2 was assessed in pHSCs before and after gp120

stimulation (100 ng/mL). Cells (0.5–1 × 10 /mL) were resuspended in PBS with 0.5% bovine

serum albumin (BSA), blocked with FcR reagent, and stained with anti-human CXCR1-APC

(Biolegend, Cat. 320612), CXCR2-FITC (Biolegend, Cat. 320704), and TLR2-PE (Biolegend,

Cat. 309708) for 15 minutes. After fixation and permeabilization (20 minutes at 4°C), samples

were analyzed on an LSRII flow cytometer (BD Biosciences). Data were processed using FCS

Express 6 (De Novo Software).

Histone Extraction and Acetylation ELISA

pHSCs and Lx2 cells (1 × 10) were seeded in 10 cm dishes and infected with HIV-1$_{BaL}$ virus

(NIH AIDS Reagent program, Bethesda, MD) for 48 hours. Total histones (200 ng) were

extracted using the EpiQuick™ Total Histone Extraction Kit (Cat. OP-0006) and quantified for

acetylation using a histone acetylation ELISA Kit (Cat. P-4022-96) per manufacturer protocols.

RNA Extraction and RT-qPCR

Total RNA was extracted using RNeasy spin columns (Qiagen, Cat. 74136) and treated with

RNase-free DNase (Qiagen, Chatsworth, CA). Reverse transcription was performed using the

Clontech RNA-to-cDNA EcoDry Kit. RT-qPCR was conducted on a Roche LightCycler 480 II

using SYBR Green PCR master mix (Bio-Rad, Cat. 1708884) with primers at 300 nM (primer

sequences in Table 1). Reactions were run in triplicate. Cycle thresholds (CTs) were normalized

to α-actin.

IL-8 ELISA

HSCs were treated under various conditions for 24 hours. Supernatants were collected at

designated time points, and IL-8 levels were measured using the Human IL-8 ELISA Kit

(Affymetrix, Cat. 88-8086-88).

CHIP RT-PCR Assay

pHSCs (1×10) were infected with HIV-1$_{BaL}$ virus (NIH AIDS Reagent program, Bethesda,

MD) for 48 hours. Chromatin was prepared using the EZ-Zyme™ Chromatin Prep Kit (Cat. 17-

375), and ChIP assays were performed with Magna ChIP™ G Kits (Cat. 17409). IL-8 promoter

activation was quantified using SYBR Green RT-PCR. CHIP-seq data was analyzed with

*Bowtie*2[24], *Macs callpeak*[25], *Samtools*[26], *Bedtools*[27], and *deepTools*[28]. H4K5 peaks were

visualized using IGV.

RNA-Seq Analysis

Total RNA from Lx2 cells treated with medium, HIV-1$_{BaL}$ (MOI $= 0.1$), and/or LTA (1 µg/mL)

was sequenced (n $= 3$ per group). RNA quality was assessed by a Nano Drop 2000

Bioanalyzer, and RNA-Seq was performed by Mount Sinai NGS Core. Reads were aligned with

STAR[29], quantified with HTSeq[30], and analyzed with DESeq2[31] using hg19 annotation. Pathway

analysis used Cytoscape with ClueGO 14 and R packages: *clusterProfiler*[32], *enrichplot*[33], and

Weighted Gene Co-Expression Network Analysis (WGCNA)[34].

Western Blot Analysis

Proteins from liver tissues were extracted with T-PER Reagent (Thermo, Cat. 78510). Equal

protein amounts were loaded onto NuPAGE 4–12% Bis-Tris gels, transferred to PVDF

membranes. Membranes were washed with blotting buffer (1× 0.1% Tween20 PBS) and

blocked for 60 minutes in blotting buffer containing 10% low-fat powdered milk. Membranes

were washed 3 times with blotting buffer. Blots were probed with primary antibodies (1:1000)

overnight at 4°C and secondary antibodies (1:1000) for 1 hour at room temperature. Target

protein signals were developed with TMB (SeraCare Life Sciences Inc., Milford, MA) and

visualized using the Amersham Imager 600. Antibodies included α-SMA (NOVUS, Cat. NBP 1-

45433 and NB100-56565), Collagen I (NOVUS, Cat. NBP 2-12446), MMP9 (NOVUS), and

GAPDH (Santa Cruz, Cat. sc-66163). Densitometry was performed using ImageJ and

normalized to β-actin.

Statistical Analysis

Data are reported as mean ± SEM. Statistical analyses were performed using GraphPad

Prism 5.0 (La Jolla, CA). Two-group comparisons used Student's t test or the Wilcoxon signed-

rank test, as appropriate. Multi-group comparisons used one-way ANOVA with Kruskal–Wallis

test. Co-localization in IHC was quantified using Fiji ImageJ and Pearson's r. $P < 0.05$ was

considered statistically significant; $P = 0.05–0.09$ was considered a trend. Non-significant

differences are noted as $P \geq 0.1$ unless stated otherwise.

Results

HIV-1$_{BaL}$ Exposure Amplifies IL-8 Response of pHSCs to LTA Stimulation

To examine how lipoteichoic acid (LTA) affects primary human hepatic stellate cells (pHSCs),

three independent control pHSC lines (from HIV-1 uninfected liver tissues) were treated with

LTA. Secretion of pro-inflammatory cytokines IL-6, IL-8, IL-10, and TNF-α was assessed before

and after 48-hour exposure to HIV-1$_{BaL}$. LTA treatment alone did not significantly induce IL-6,

TNF-α, or IL-10 secretion in either condition (Supplemental Fig. 1). However, IL-8 secretion was

significantly elevated in HIV-1$_{BaL}$–exposed pHSCs following LTA stimulation compared to LTA

alone (Fig. 1A).

To validate these findings, pHSCs isolated from HIV-1–infected livers were stimulated with LTA.

Two out of three HIV+ pHSC lines showed significantly increased IL-8 secretion with LTA (Fig.

1B). Similarly, Lx2 cells co-stimulated with LTA and HIV-1$_{BaL}$ displayed synergistic IL-8 induction

(Fig. 1C). Lastly, the C-C chemokine receptor type 5 (CCR5) antagonist Maraviroc successfully

suppressed HIV-1$_{BaL}$ infection in Lx2 cells, as indicated by reduced p24 levels (Fig. 1D), and

significantly attenuated IL-8 production in response to LTA (Fig. 1E).

Flow cytometry analysis of control pHSCs revealed that among IL-8/C-X-C motif chemokine

receptors, CXCR1 expression increased upon gp120 stimulation— an HIV-1 envelope protein—

while CXCR2 and LTA ligand receptor (TLR2) remained unchanged (Fig. 2A). Notably, neither

IL-8 receptor nor TLR2 inhibition altered the heightened IL-8 response to LTA in HIV-1$_{BaL}$–

exposed cells (Figs. 2B–2C). Collectively, these findings suggest that HIV-1 sensitizes pHSCs

to LTA promoting IL-8 secretion via a pathway independent of IL-8 receptor or TLR2 signaling.

Elevated IL-8 Expression in HIV-1–Infected Liver Tissue

To confirm the in vitro observations, liver tissues from PLWH were analyzed. IL-8, TGF-β, and

α-SMA mRNA levels were significantly higher in HIV-1–infected tissues compared to uninfected

controls (Fig. 3A). α-SMA protein levels were also elevated in HIV-1–infected tissues (Fig. 3B).

Immunohistochemistry confirmed higher hepatic IL-8 expression in HIV-1–infected tissues (Fig.

3C).

Serum IL-8 levels were elevated in individuals with HIV viremia and liver fibrosis compared to

those with fibrosis alone. Although mean serum LTA levels showed no groupwise significance,

individuals with both HIV viremia and fibrosis exhibited the highest LTA concentrations (mean:

138.81 pg/mL), compared to those with fibrosis only (96.44 pg/mL) or no fibrosis/viremia (19.28

pg/mL) (Fig. 3D).

To test whether LTA or IL-8 exert direct fibrogenic effects, Lx2 cells were treated with increasing

doses of LTA and IL-8. IL-8 mRNA expression increased dose-dependently following LTA

exposure in Lx2 cells, with no change in other pro-fibrogenic genes (Fig. 4A). IL-8 treatment

elevated α-SMA and Collagen I protein levels in Lx2 cells (Fig. 4B), supporting its fibrogenic

potential in HIV-associated liver disease.

RNA-Seq Reveals Epigenetic Regulation of IL-8 in HIV-1 and LTA Co-Stimulation

To investigate how HIV-1 exposure sensitizes HSCs to LTA–induced IL-8 expression, RNA-seq

was conducted on Lx2 cells under three experimental conditions: untreated, treated with HIV-

1_{BaL} alone, or co-treated with HIV-1_{BaL} and LTA (Fig. 5). Although global differential gene

expression did not reach statistical significance (Figs. 5A–5B), IL-8 expression trended higher in

cells co-stimulated with HIV-1_{BaL} and LTA (p = 0.059) (Fig. 5C).

WGCNA followed by ClueGO pathway analysis of differentially expressed genes (DEGs)

correlated with IL-8 (correlation coefficient r>0.2, p≤0.05) linked IL-8 to genes involved in tRNA

aminoacylation (Figs. 5D–5E), including an upregulation of *SIRT4* and downregulation of *SIRT3*

and related genes (Fig. 5F). Given the role of SIRT3 and SIRT4 in acetylation regulation during

oxidative stress and inflammation[35], we further explored these findings. This revealed altered

expression of genes associated with histone H4 acetylation and deacetylation—such as HDAC2

and MORFL2[36,37]—in response to combined HIV-1_{BaL} and LTA exposure (Fig. 5G), suggesting

an epigenetic mechanism in IL-8 amplification in LTA and HIV-1 co-stimulated Lx2 cells[35].

H4K5 Acetylation is Required for IL-8 Transactivation in HIV-1–Exposed HSCs

To test the role of histone acetylation, the histone deacetylase (HDAC) inhibitor Trichostatin A

(TSA) was applied during LTA stimulation. TSA significantly increased IL-8 secretion in both

control and HIV+ pHSCs (Figs. 6A–6B) without affecting IL-1β, IL-6, IL-10, or TNF-α

(Supplemental Fig. 1). HIV-1_{BaL} infection also elevated total acetylation in Lx2 cells

(Supplemental Fig. 2), with H4K5 acetylation specifically increased (Fig. 6C), confirmed by

Western blot (Fig. 6D). H3 acetylation remained unchanged (Supplemental Fig. 3A–D).

ChIP-RT-PCR identified H4K5 acetylation enrichment at IL-8 promoter regions in HIV-1–

infected Lx2 cells, particularly at NFκB, AP-1, and NF-IL6 binding sites (Figs. 6E–6F). ChIP-seq

further confirmed strong peaks in NF-kB and AP-1 binding motifs near the IL-8 transcriptional

start sites (Supplemental Fig. 4A–B). Together, these results demonstrate that H4K5 acetylation

is a critical mediator of IL-8 transcriptional activation in response to LTA in HIV-1–infected

hepatic stellate cells.

Discussion

End-stage liver disease remains a major cause of mortality among PLWH, with multiple studies

showing a positive correlation between HIV viremia and advanced fibrosis in HIV mono-

infection[13-16]. Although chronic HIV infection disrupts immune homeostasis and accelerates liver

fibrosis, the specific mechanisms linking HIV-mediated inflammation to hepatic fibrogenesis

have not been fully elucidated. Our ex vivo and in vitro findings demonstrate that HIV-1

exposure amplifies the IL-8 response of HSCs to LTA—the gram-positive microbial

component—with histone acetylation–mediated IL-8 transactivation identified as a key

underlying mechanism.

Elevated serum IL-8 levels (≥46.5 pg/mL) have been independently associated with liver-related

mortality and the need for liver transplantation[38,39]. Although IL-8 is elevated in mono-infection[40],

its pathological role in HIV-associated liver fibrosis has remained uncertain. Some studies have

found no correlation between IL-8 levels and fibrosis in HIV/HCV co-infection, while others

suggest elevated IL-8 predicts a greater risk of fibrosis progression[41,42]. In our HIV-1 mono-

infected cohort, IL-8 levels were significantly higher in viremic individuals with advanced fibrosis

compared to non-viremic counterparts. Interestingly, this distinction was present only in

individuals with advanced fibrosis, suggesting a specific link between IL-8 expression and HIV-

mediated fibrogenesis. Immunohistochemistry of HIV-1–infected liver tissues confirmed

increased hepatic IL-8 expression, which was accompanied by upregulated pro-fibrogenic

markers α-SMA and collagen I. In vitro, stellate cell lines responded to IL-8 stimulation with

increased expression of these same fibrosis-related genes, supporting a direct fibrogenic role

for IL-8 in HSCs.

Previous studies have identified IL-8 as a marker of inflammatory response to liver injury in

HSCs pretreated with TNF-α[43,44]. Furthermore, LTA has been shown to promote IL-8 production

through NF-κB activation in HSCs pre-primed with TNF-α and IL-1β[45]. Notably, in our study, LTA

alone was sufficient to stimulate IL-8 secretion in the absence of such priming, and this effect

was significantly enhanced by HIV-1 exposure. Importantly, HIV-1–exposed HSCs did not

exhibit increased TNF-α or IL-1β production, reinforcing the possibility that HIV infection

independently alters HSC sensitivity to microbial products. Although prior work implicated TLR-

mediated interactions between HIV-1, immune cells, and HSCs[46], our data show that the

enhanced IL-8 response is independent of IL-8 receptors or TLR2 signaling. Consistently, RNA-

seq analyses revealed no activation of the canonical NF-κB pathway following HIV-1 and LTA

co-exposure (Supplementary Fig. 5), suggesting that IL-8 induction involves an alternative

regulatory mechanism.

Transcriptomic analysis linked IL-8 transactivation to the aminoacylation pathway and identified

differential expression of genes related to histone H4 acetylation. Previous research has shown

that IL-8 promoter acetylation contributes to its post-translational regulation[47]and that HDAC

inhibitors can suppress IL-8–driven inflammation in various disease contexts[48]. In our ex vivo

experiments, treatment with the HDAC inhibitor TSA selectively increased IL-8 responses to

LTA, regardless of HIV-1 exposure, without affecting other cytokines such as IL-1β, IL-6, or

TNF-α. Through ELISA, ChIP-seq, and ChIP-qPCR, we identified specific enrichment of H4K5

acetylation at the IL-8 promoter in HIV-1–infected cells. This modification promotes chromatin

remodeling and recruitment of critical binding sites in the IL-8 promoter i.e., transcription factors

AP-1, NF-IL6, and NF-κB[49].

These findings align with prior studies in cystic fibrosis[50-53]and hepatitis B virus infection[54],

where histone hyperacetylation was shown to drive IL-8 expression in epithelial or

hepatocellular cells. Our work highlights the unique role of histone acetylation in modulating

HSC responses to microbial stimuli in HIV-associated liver disease. This discovery opens up

new avenues for exploring the epigenetic regulation of fibrosis in HIV and underscores the

potential therapeutic relevance of targeting gram-positive microbial products or histone

acetylation pathways.

Conclusion

Our ex vivo and in vitro findings underscore the pathological contribution of LTA to hepatic

inflammation and fibrogenesis during chronic HIV-1 infection. Elevated serum LTA levels in HIV-

1–infected individuals with both viremia and fibrosis, alongside amplified IL-8 secretion from

HSCs in response to LTA under HIV exposure, point to gram-positive microbial translocation as

a critical driver of disease progression. Transcriptomic analysis revealed that HIV-1 enhances

histone H4K5 acetylation, thereby increasing IL-8 promoter activity in response to LTA

stimulation, and contributing to sustained fibrogenic signaling in HSCs. These findings link gut-

derived microbial products, IL-8–driven inflammation, and epigenetic regulation of HSC

activation, providing important mechanistic insight and potential therapeutic targets for

managing HIV-associated liver fibrosis.

Acknowledgement

We appreciate the reagents support from NIH AIDS Reagents Program.

Authors' Contribution

Drs. Zhang and Bansal conceptualized and designed the experiments. Dr. Zhang performed the

experiments and conducted bioinformatics analyses. Drs. Zhang, Ali, and Bansal analyzed the

data and drafted the manuscript. Dr. Chamroonkul prepared serum samples. Drs. Tabrizian,

Schwartz, Gunasekaran, Schiano, and Fiel provided clinical samples and associated clinical

data. All authors contributed to the manuscript revisions and approved the final version.

Funding

This work was supported by research grant NIH R56DK92128 and R01DK108364 to MBB.

Conflict of Interest

The authors declare no conflicts of interest.

Data Availability

Processed data files are available at github.com/lumin2003/HIV_LTA_files/.

Declarations

The human study has been approved by Mount Sinai IRB (GCO #06-0523 and #10-1211).

Consent for Publication

There is no individual patient information in this study.

Competing Interests

None of the authors have competing interests in this study.

References

1   Zhang, L. *et al.* Humoral immune responses to Streptococcus pneumoniae in the setting

of HIV-1 infection. *Vaccine* **33**, 4430-4436 (2015). https://doi.org/10.1016/j.vaccine.2015.06.077

2   Sandler, N. G. & Douek, D. C. Microbial translocation in HIV infection: causes,

consequences and treatment opportunities. *Nat Rev Microbiol* **10**, 655-666 (2012).

https://doi.org/10.1038/nrmicro2848

3   Ipp, H., Zemlin, A. E., Erasmus, R. T. & Glashoff, R. H. Role of inflammation in HIV-1

disease progression and prognosis. *Crit Rev Clin Lab Sci* **51**, 98-111 (2014).

https://doi.org/10.3109/10408363.2013.865702

4   Ljungdahl, M. *et al.* Bacterial translocation in experimental shock is dependent on the

strains in the intestinal flora. *Scand J Gastroenterol* **35**, 389-397 (2000).

https://doi.org/10.1080/003655200750023958

5   Dillon, S. M., Frank, D. N. & Wilson, C. C. The gut microbiome and HIV-1 pathogenesis:

a two-way street. *AIDS* **30**, 2737-2751 (2016).

https://doi.org/10.1097/QAD.0000000000001289

6   Kaspar, M. B. & Sterling, R. K. Mechanisms of liver disease in patients infected with HIV.

BMJ Open Gastroenterol **4**, e000166 (2017). https://doi.org/10.1136/bmjgast-2017-

000166

7   Debes, J. D., Bohjanen, P. R. & Boonstra, A. Mechanisms of Accelerated Liver Fibrosis

Progression during HIV Infection. *J Clin Transl Hepatol* **4**, 328-335 (2016).

https://doi.org/10.14218/JCTH.2016.00034

8   Ginsburg, I. Role of lipoteichoic acid in infection and inflammation. *Lancet Infect Dis* **2**,

171-179 (2002). https://doi.org/10.1016/s1473-3099(02)00226-8

9   Weidenmaier, C. & Peschel, A. Teichoic acids and related cell-wall glycopolymers in

Gram-positive physiology and host interactions. *Nat Rev Microbiol* **6**, 276-287 (2008).

https://doi.org/10.1038/nrmicro1861

10   Theilacker, C. *et al.* Protection against Staphylococcus aureus by antibody to the

polyglycerolphosphate backbone of heterologous lipoteichoic acid. *J Infect Dis* **205**,

1076-1085 (2012). https://doi.org/10.1093/infdis/jis022

11   Hattor, Y., Kasai, K., Akimoto, K. & Thiemermann, C. Induction of NO synthesis by

lipoteichoic acid from Staphylococcus aureus in J774 macrophages: involvement of a

CD14-dependent pathway. *Biochem Biophys Res Commun* **233**, 375-379 (1997).

https://doi.org/10.1006/bbrc.1997.6462

12   Siewe, B., Keshavarzian, A., French, A., Demarais, P. & Landay, A. A role for TLR

signaling during B cell activation in antiretroviral-treated HIV individuals. *AIDS Res Hum*

Retroviruses **29**, 1353-1360 (2013). https://doi.org/10.1089/AID.2013.0115

13   Sanmartin, R. *et al.* Progression of liver fibrosis in HIV/hepatitis C virus-coinfected

individuals on antiretroviral therapy with early stages of liver fibrosis at baseline. *HIV*

Med **15**, 203-212 (2014). https://doi.org/10.1111/hiv.12105

14   Blackard, J. T. *et al.* HIV mono-infection is associated with FIB-4 - A noninvasive index

of liver fibrosis - in women. *Clin Infect Dis* **52**, 674-680 (2011).

https://doi.org/10.1093/cid/ciq199

15   Stabinski, L. *et al.* High prevalence of liver fibrosis associated with HIV infection: a study

in rural Rakai, Uganda. *Antivir Ther* **16**, 405-411 (2011). https://doi.org/10.3851/IMP1783

16   Tahiri, M. *et al.* Risk factors for liver fibrosis among human immunodeficiency virus

monoinfected patients using the FIB4 index in Morocco. *World J Hepatol* **5**, 584-588

(2013). https://doi.org/10.4254/wjh.v5.i10.584

17   Mohsen, A. H. *et al.* Impact of human immunodeficiency virus (HIV) infection on the

progression of liver fibrosis in hepatitis C virus infected patients. *Gut* **52**, 1035-1040

(2003). https://doi.org/10.1136/gut.52.7.1035

18   Koyama, Y. & Brenner, D. A. Liver inflammation and fibrosis. *J Clin Invest* **127**, 55-64

(2017). https://doi.org/10.1172/JCI88881

19   Mosoian, A. *et al.* Frontline Science: HIV infection of Kupffer cells results in an amplified

proinflammatory response to LPS. *J Leukoc Biol* **101**, 1083-1090 (2017).

https://doi.org/10.1189/jlb.3HI0516-242R

20   Zhang, L. *et al.* HIV infection modulates IL-1beta response to LPS stimulation through a

TLR4-NLRP3 pathway in human liver macrophages. *J Leukoc Biol* **105**, 783-795 (2019).

https://doi.org/10.1002/JLB.4A1018-381R

21   T, L. Analysis of Complex Survey Samples. *journal of Statistical Software*, 1-19 (2004).

22   Panos, A. & Mavridis, D. TableOne: an online web application and R package for

summarising and visualising data. *Evid Based Ment Health* **23**, 127-130 (2020).

https://doi.org/10.1136/ebmental-2020-300162

23   Alabraba, E. B. *et al.* A new approach to isolation and culture of human Kupffer cells. *J*

Immunol Methods **326**, 139-144 (2007). https://doi.org/10.1016/j.jim.2007.06.014

24   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*

9, 357-359 (2012). https://doi.org/10.1038/nmeth.1923

25   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137

(2008). https://doi.org/10.1186/gb-2008-9-9-r137

26   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,

2078-2079 (2009). https://doi.org/10.1093/bioinformatics/btp352

27   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841-842 (2010).

https://doi.org/10.1093/bioinformatics/btq033

28   Ramirez, F., Dundar, F., Diehl, S., Gruning, B. A. & Manke, T. deepTools: a flexible

platform for exploring deep-sequencing data. *Nucleic Acids Res* **42**, W187-191 (2014).

https://doi.org/10.1093/nar/gku365

29   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21

(2013). https://doi.org/10.1093/bioinformatics/bts635

30   Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-

throughput sequencing data. *Bioinformatics* **31**, 166-169 (2015).

https://doi.org/10.1093/bioinformatics/btu638

31   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

https://doi.org/10.1186/s13059-014-0550-8

32   Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for comparing

biological themes among gene clusters. *OMICS* **16**, 284-287 (2012).

https://doi.org/10.1089/omi.2011.0118

33   Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.

Innovation (Camb) **2**, 100141 (2021). https://doi.org/10.1016/j.xinn.2021.100141

34   Gennady Korotkevich, V. S., Alexey Sergushichev. Fast gene set enrichment analysis.

bioRaxiv (2021).

35   Zou, Y. *et al.* The regulatory roles of aminoacyl-tRNA synthetase in cardiovascular

disease. *Mol Ther Nucleic Acids* **25**, 372-387 (2021).

https://doi.org/10.1016/j.omtn.2021.06.003

36   Kooblall, K. G. *et al.* miR-3156-5p is downregulated in serum of MEN1 patients and

regulates expression of MORF4L2. *Endocr Relat Cancer* **29**, 557-568 (2022).

https://doi.org/10.1530/ERC-22-0045

37   Becht, D. C. *et al.* MORF and MOZ acetyltransferases target unmethylated CpG islands

through the winged helix domain. *Nat Commun* **14**, 697 (2023).

https://doi.org/10.1038/s41467-023-36368-5

38   Zimmermann, H. W. *et al.* Interleukin-8 is activated in patients with chronic liver diseases

and associated with hepatic macrophage accumulation in human liver fibrosis. *PLoS*

One **6**, e21381 (2011). https://doi.org/10.1371/journal.pone.0021381

39   Umemura, T. *et al.* Association between serum soluble CD14 and IL-8 levels and clinical

outcome in primary biliary cholangitis. *Liver Int* **37**, 897-905 (2017).

https://doi.org/10.1111/liv.13316

40   Ellwanger, J. H. *et al.* Increased IL-8 levels in HIV-infected individuals who initiated ART

with CD4(+) T cell counts <350 cells/mm(3) - A potential hallmark of chronic

inflammation. *Microbes Infect* **22**, 474-480 (2020).

https://doi.org/10.1016/j.micinf.2020.05.019

41   Chen, X. *et al.* Plasma Inflammatory Biomarkers Associated with Advanced Liver

Fibrosis in HIV-HCV-Coinfected Individuals. *Int J Environ Res Public Health* **17** (2020).

https://doi.org/10.3390/ijerph17249474

42   Keating, S. M. *et al.* The effect of HIV infection and HCV viremia on inflammatory

mediators and hepatic injury-The Women's Interagency HIV Study. *PLoS One* **12**,

e0181004 (2017). https://doi.org/10.1371/journal.pone.0181004

43   Lang, A. *et al.* Somatostatin inhibits pro-inflammatory cytokine secretion from rat hepatic

stellate cells. *Liver Int* **25**, 808-816 (2005). https://doi.org/10.1111/j.1478-

3231.2005.01057.x

44   Schwabe, R. F., Schnabl, B., Kweon, Y. O. & Brenner, D. A. CD40 activates NF-kappa B

and c-Jun N-terminal kinase and enhances chemokine secretion on activated human

hepatic stellate cells. *J Immunol* **166**, 6812-6819 (2001).

https://doi.org/10.4049/jimmunol.166.11.6812

45   Paik, Y. H. *et al.* Hepatic stellate cells primed with cytokines upregulate inflammation in

response to peptidoglycan or lipoteichoic acid. *Lab Invest* **86**, 676-686 (2006).

https://doi.org/10.1038/labinvest.3700422

46   Del Corno, M. *et al.* HIV-1 gp120 signaling through TLR4 modulates innate immune

activation in human macrophages and the biology of hepatic stellate cells. *J Leukoc Biol*

100, 599-606 (2016). https://doi.org/10.1189/jlb.4A1215-534R

47   Takahashi, A., de Andres, M. C., Hashimoto, K., Itoi, E. & Oreffo, R. O. Epigenetic

regulation of interleukin-8, an inflammatory chemokine, in osteoarthritis. *Osteoarthritis*

Cartilage **23**, 1946-1954 (2015). https://doi.org/10.1016/j.joca.2015.02.168

48   Gatla, H. R. *et al.* Regulation of Chemokines and Cytokines by Histone Deacetylases

and an Update on Histone Decetylase Inhibitors in Human Diseases. *Int J Mol Sci* **20**

(2019). https://doi.org/10.3390/ijms20051110

49   Roebuck, K. A. Regulation of interleukin-8 gene expression. *J Interferon Cytokine Res*

19, 429-438 (1999). https://doi.org/10.1089/107999099313866

50   Ashburner, B. P., Westerheide, S. D. & Baldwin, A. S., Jr. The p65 (RelA) subunit of NF-

kappaB interacts with the histone deacetylase (HDAC) corepressors HDAC1 and

HDAC2 to negatively regulate gene expression. *Mol Cell Biol* **21**, 7065-7077 (2001).

https://doi.org/10.1128/MCB.21.20.7065-7077.2001

51   Vanden Berghe, W., De Bosscher, K., Boone, E., Plaisance, S. & Haegeman, G. The

nuclear factor-kappaB engages CBP/p300 and histone acetyltransferase activity for

transcriptional activation of the interleukin-6 gene promoter. *J Biol Chem* **274**, 32091-

32098 (1999). https://doi.org/10.1074/jbc.274.45.32091

52   Jundi, K. & Greene, C. M. Transcription of Interleukin-8: How Altered Regulation Can

Affect Cystic Fibrosis Lung Disease. *Biomolecules* **5**, 1386-1398 (2015).

https://doi.org/10.3390/biom5031386

53 Barnes, P. J., Adcock, I. M. & Ito, K. Histone acetylation and deacetylation: importance

in inflammatory lung diseases. *Eur Respir J* **25**, 552-563 (2005).

https://doi.org/10.1183/09031936.05.00117504

54 Pollicino, T. *et al.* Hepatitis B virus (HBV) induces the expression of interleukin-8 that in

turn reduces HBV sensitivity to interferon-alpha. *Virology* **444**, 317-328 (2013).

https://doi.org/10.1016/j.virol.2013.06.028

Figure Legends

Figure 1: HIV-1 exposure accentuates IL-8 response to LTA stimulation in primary

isolated HSCs. A) Primary human hepatic stellate cells (pHSCs) were isolated from HIV-1–

uninfected liver tissues (n=3) and treated with LTA (1□µg/ml) either before or after exposure

with HIV-1$_{BaL}$ (MOI=0.1) for 48 hours. Supernatants were collected 24 hours after LTA

stimulation, and IL-8 levels were measured by ELISA. B) pHSCs from HIV-1–infected liver

tissues (n=6) were stimulated with LTA (1□µg/ml) for 24 hours, and IL-8 levels were quantified

by ELISA. C) Lx2 cells were treated with LTA alone or HIV-1 plus LTA. Supernatants were

collected and IL-8 was measured by ELISA (n=6). D) HIV-1 p24 antigen was quantified by

ELISA in supernatants from HIV-1$_{BaL}$-exposed Lx2 cells (MOI=0.1) with or without Maraviroc

(1□mM) treatment (n=9). E) IL-8 levels were measured by ELISA in HIV-1$_{BaL}$-exposed Lx2 cells

treated with or without Maraviroc (1□mM) (n=6). HIV-1$_{BaL}$

Figure 2: IL-8 response of HSCs to LTA is independent of IL-8 and TLR receptor

pathways. A) IL-8 receptor was analyzed by Flow cytometry before and after gp120 stimulation

(100ng/ml). B) Lx2 cells were stimulated with LTA (1□µg/ml) for 24 hours with or without IL-8

receptor inhibitor. Supernatants were analyzed for IL-8 levels by ELISA (n=3). C) Lx2 cells were

exposed to HIV-1$_{BaL}$ for 48 hours and then treated with either an IL-8 receptor inhibitor or a

TLR2 inhibitor. Supernatants were analyzed for IL-8 levels by ELISA (n=3).

Figure 3: Hepatic IL-8 and pro-fibrogenic gene expression is elevated in HIV-1–infected

liver tissues. A) Total RNA was extracted from HIV-1–uninfected (n=7) and HIV-1–infected

(n=7) liver tissues. mRNA expression of IL-8, α-SMA, collagen I, and TGF-β was quantified by

RT-PCR. B) Protein expression of α-SMA and collagen I was examined by western blot;

GAPDH served as a loading control. Band intensity was quantified using ImageJ (n>3). C) IL-8

expression was evaluated by immunohistochemistry in HIV-1–uninfected and HIV-1–infected

liver tissues (n=32). The results were quantified by image J. D) Serum IL-8 and LTA

concentration were measured in duplicate by ELISA and compared across groups stratified by

HIV-1 viremia and fibrosis ($n^{HIV\text{-}1\ viremia\text{-}} = 9$, $n^{HIV\text{-}1\ viremia\text{-}\ Fib+} = 9$, $n^{HIV\text{-}1\ viremia+Fib+} = 10$).

Figure 4: IL-8 increases pro-fibrogenic gene expression in hepatic stellate cell lines. A)

Lx2 cells were treated with increasing concentrations of LTA for 24 hours. mRNA expression of

pro-fibrotic and inflammatory genes was quantified by SYBR Green RT-PCR. B) Lx2 cells were

treated with IL-8 (200 pg/ml) or PBS x 3. Protein expression of α-SMA and collagen I was

analyzed by western blot and normalized to GAPDH. Band intensities were quantified using

ImageJ (n=3).

Figure 5: RNA-seq suggests a link between histone acetylation and IL-8 activation

following HIV-1 and LTA co-stimulation in Lx2 Cells. A-B) Volcano plots showing

differentially expressed genes (DEGs) between control, HIV-1$_{BaL}$-exposed, and HIV-1$_{BaL}$ plus

LTA-treated Lx2 cells (p < 0.05, log2FC > 1.0). Blue and red dots represent up- and down-

regulated genes, respectively. C) Heatmap analysis of pro-inflammatory genes across treatment

groups. D) ClueGO pathway enrichment of genes positively correlated with IL-8 expression

(correlation coefficient *r* > 0.2), highlighting the tRNA aminoacylation pathway. E) Cytoscape-

based network reconstruction of the tRNA aminoacylation pathway, showing IL-8/CXCL8

enrichment. F) Heatmap of tRNA aminoacylation gene expression across three experimental

conditions. G) Heatmap analysis of acetylation related genes across three experimental

conditions.

Figure 6: H4K5 acetylation is essential for IL-8 response to LTA stimulation in HIV-1

infected HSCs. A) Two pHSCs isolated from HIV-1–uninfected individuals were stimulated with

LTA with or without HDAC inhibitor (TSA) for 24hrs, IL-8 in supernatant was measured by

ELISA (n=6). B) Two pHSCs isolated from HIV-1–infected individuals were stimulated with LTA

with or without HDAC inhibitor (TSA) for 24hrs, IL-8 in supernatant was measured by ELISA

(n=6). C) H4K5 acetylation protein was measured by ELISA in both pHSCs (n=4) and Lx2 cell

lines (n=8) before and after HIV-1$_{BaL}$ exposure. D) H4K5 acetylation protein was examined by

western blot and normalized to GAPDH. E) Binding motif analysis of IL-8 promoter. F)

Chromatin immunoprecipitation (ChIP) was performed using anti-H4K5 antibody. DNA was

purified and analyzed by SYBR RT-PCR for enrichment of NF-κB, AP-1, and NF-IL6 binding

motifs in the IL-8 promoter.

Tables

Table 1: Primers:

All primer sequences.

h-a-SMA R: 5'-AGGCACCCCTGAACCCCAA-3'
h-a-SMA R: 5'-CAGCACCGCCTGGATAGCC-3'
h-Collagen I F: 5'-GGCTTCCCTGGTCTTCCTGG-3'
h-Collagen I R: 5'-CCAGGGGGTCCAGCCAAT-3'
h-PDGF F: 5'-CCAGAAGCCATCAGCAGCAAG-3'
h-PDGF R: 5'-AGGCCCTGAGAGATCTGTGG-3'

h-MMP2 F: 5'- CCCCAAGCTCATCGCAGAT-3'
h-MMP2 R: 5'- GGTCCACGACGGCATCC-3'
h-TIMP1 F: 5'-TGACATCCGGTTCGTCTACA-3'
h-TIMP1 R: 5'-TGATGTGCAAGAGTCCATCC-3'
h-TIMP2 F: 5'- AAGCGGTCAGTGAGAAGGAA-3'
h-TIMP2 R: 5'-GGGGGCCGTGTAGATAAACT-3'
h-TGF-beta F: 5'-CAAGGGCTACCATGCCAACT-3'
h-TGF-beta R: 5'-AGGGCCAGGACCTTGCTG-3'
h-TGF-beta1 Receptor F: 5'-CGACTGCCAAATGAAGAGGACC-3'
h-TGF-beta1 Receptor R: 5'-AAACCTGAGCCAGAACCTGACG-3'
h-IL-8 F: 5'-TAGGACAAGAGCCAGGAAGA-3'
h-IL-8R: 5'-CTTCCTGATTTCTGCAGCTCT-3'
h-GAPDH F: 5'-CAATGACCCCTTCATTGACC-3'
h-GAPDH R: 5'- GACAAGCTTCCCGTTCTCAG -3'

609 **Table 2:** Characteristics of blood donors.

	Overall	0	1	p test
n	28	11	17	
Sigfibrosis = 1 (%)	17.0 (60.7)	0.0 (0.0)	17.0 (100.0)	<0.001
gender = 2 (%)	5.0 (17.9)	2.0 (18.2)	3.0 (17.6)	0.972
Ethinic (%)				0.368
AA	7.0 (25.0)	2.0 (18.2)	5.0 (29.4)	
AS	1.0 (3.6)	0.0 (0.0)	1.0 (5.9)	
HS	13.0 (46.4)	4.0 (36.4)	9.0 (52.9)	
O	3.0 (10.7)	2.0 (18.2)	1.0 (5.9)	
W	4.0 (14.3)	3.0 (27.3)	1.0 (5.9)	
age (mean (SD))	52.89 (9.52)	53.55 (4.27)	52.47 (11.88)	0.732
bmi (mean (SD))	30.55 (6.62)	29.21 (6.42)	31.41 (6.79)	0.384
alcohol = 1 (%)	9.0 (32.1)	3.0 (27.3)	6.0 (35.3)	0.666
duradiag (mean (SD))	19.82 (8.69)	19.20 (8.94)	20.33 (8.85)	0.762
viremia = 1 (%)	19.0 (67.9)	11.0 (100.0)	8.0 (47.1)	0.005
cd4 (mean (SD))	649.18 (324.81)	627.00 (348.98)	663.53 (318.36)	0.777
fibroscan (mean (SD))	9.33 (5.21)	5.50 (1.23)	11.81 (5.31)	<0.001
plt (mean (SD))	197.71 (63.29)	200.73 (58.45)	195.76 (67.93)	0.835
alt (mean (SD))	30.46 (20.36)	23.00 (9.14)	35.29 (24.17)	0.065
ALTab = 1 (%)	10.0 (35.7)	2.0 (18.2)	8.0 (47.1)	0.138
alp (mean (SD))	91.18	82.55	96.76	0.23

	(28.40)	(34.78)	(22.82)	
alb (mean (SD))	3.86 (0.82)	4.01 (0.29)	3.77 (1.02)	0.366
fbs (mean (SD))	106.61 (37.03)	90.45 (22.32)	117.06 (41.31)	0.034
chol (mean (SD))	167.71 (41.52)	182.73 (48.59)	158.00 (34.33)	0.144
tg (mean (SD))	167.79 (133.44)	170.91 (122.65)	165.76 (143.64)	0.918
hdl (mean (SD))	41.78 (9.67)	42.20 (9.37)	41.53 (10.12)	0.86
lowHDL (mean (SD))	0.56 (0.51)	0.50 (0.53)	0.59 (0.51)	0.666
ldl (mean (SD))	94.18 (34.95)	110.04 (45.12)	83.92 (22.46)	0.079
CAP (mean (SD))	283.18 (61.62)	261.00 (66.65)	297.53 (55.45)	0.134
fattyliver = 1 (%)	23.0 (82.1)	8.0 (72.7)	15.0 (88.2)	0.313
AST (mean (SD))	28.38 (12.35)	25.27 (10.58)	30.67 (13.39)	0.253
HCV (mean (SD))	0.46 (0.51)	0.45 (0.52)	0.47 (0.51)	0.936
HBV (mean (SD))	0.04 (0.19)	0.00 (0.00)	0.06 (0.24)	0.321
HSI (mean (SD))	37.48 (13.26)	37.87 (7.23)	37.21 (16.43)	0.886
FIB4 (mean (SD))	1.67 (1.24)	1.59 (0.86)	1.74 (1.48)	0.738

611 **Definition:**

612 Active alcohol consumption = drinking more than 10 g of ethanol per day in females and 20 g

613 per day in males; Liver steatosis = evidence of CAP more than 248 dB/m.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A)

Lx2 cells

p=0.008

IL8 in supernatant (pg/ml)

800 — 600 — 400 — 200 — 0

Medium   LTA   TSA-LTA

pHSCs1

p=0.0022

IL8 in supernatant (pg/ml)

400 — 300 — 200 — 100 — 0

Medium   LTA   TSA-LTA

B)

HIV1⁺ pHSCs1

p=0.004

IL8 in supernatant (pg/ml)

10000 — 8000 — 6000 — 4000 — 2000 — 0

Medium   LTA   TSA

HIV1⁺ pHSCs2

p=0.0022

p=0.02

IL8 in supernatant (pg/ml)

1000 — 800 — 600 — 400 — 200 — 0

Medium   LTA   TSA

C)

p=0.031

Actylation of H4K5(ng/uM)

500 — 400 — 300 — 200 — 100 — 0

pHSCs   pHSC2+HIV1$_{BaL}$ Lx2   Lx2-HIV1$_{BaL}$

D)

H4K5

CTL   BaL

GAPDH

E)

TGGAACCCAGATGCTCTTTCCACTAAGCATACAACTTTCCATTAGATAAACACCTCCCTCCCACCCCAACCA
AGCAGCTCCAGTGCACCACTTTCTGGAGCATAAACATACCTTAACTTTACAACTTGAGTGGCCTTGAATAC
TGTTCCTATCTGGAATGTGCTGTTCTCTTT**CATCTTCCT**CTATTGAAGCCCTCCTATTCCTCAATGCCTTG
CTCCAACTGCCTTTGGAAGATTCTGCTCTTATGCCTCCACTGGAATTAATGTCTTAGTACCACTTGTCTAT
TCTGCTATATAGTCAGTCCT**ACATTGCTTTCTTCT**TCTGATAGACCAAACTCTTTAAGGACAAGTACCTA
GTCTTATCTATTTCTAGATCCCCACATTACTTCAGAAAGTTACTCCATAAATGTTTGTGGAACTGATTTCT
ATGTGAAGCACATGTGCCCCTTCACTCTGTTAACATGCATTAGAAAACTAAATCTTTTGAAAAGTTGTAGT
ATGCCCCTAAGAGCAGTAACAGTTCCTAGAAACTCTCTAAAATGCTTAGAAAAAGATTTATTTTAAATTA
CCTCCCCAATAAAATGATTGGCTGGCTTATCTTCACCATCATGATAGCATCTGTAATTAACTGAAAATAAA
TAATTATGCCATTAAAAGAAAATCATCCATGATCTTGTTCTAACACCTGCCACTCTAGTACTATATCTGTC
ACATGGTACTATGATAAAGTTATCTAGAAATAAAAAGCATACAATTGATAATTCACCAAATTGTGGAGCT
TCAGTATTTTAAATGTATATTAAAATTAAATTATTTTAAAGATCAAAGAAAACTTTCGTCATACTCCGTAT
TTGATAAGGAACAAATAGGAAGTGTG**ATGACTCAG**GTTTGCCCTGAGGGG**ATGGGCCAT**CAGTTGCAAATC
GT**GGAATTTCCT**CTGACATAATGAAAAGAT[G]GGGGTGCATAAGTTCTCTAGTAGGGTGATGATATAAAAAG
CACCGGAGCACTCCATAAGGCACAAACTTTCAGAGACAGCAGAGCCACCAAGCTTCTAGGACAAGAGCCA
GGAAGAAACCACCGGAAGGAACCATCTCACTGTGTGTAAACAT

Red Bold :   NF-IL16:
Purple Bold :   AP-1
Blue Bold :   NF-kB

F)

CHIP-RT-PCR

The relative level of H4K5 Acetylation

200 — 150 — 100 — 50 — 30 — 20 — 10 — 0

Input   NF-kB Sites   NF-IL16 Sites   AP-1 Sites