Phytoplankton and giant virus dynamics during different monsoon seasons, a fish kill, and a toxic bloom in a eutrophic mariculture area

Andrian P. Gajigan[1,2*], Gianina Cassandra May B. Apego[3], Eunice Lois D. Gianan[3], Rachel B. Francisco[3], Jun Jet Tai[4], Garry A. Benico[5], Charina Lyn A. Repollo[3], Cesar L. Villanoy[3], Aletta T. Yñiguez[3], Cecilia Conaco[3], Grieg F. Steward[1,2*]

[1]Department of Oceanography, School of Ocean and Earth Science and Technology, University of Hawaiʻi at Mānoa, USA
[2]Center for Microbial Oceanography Research and Education, University of Hawaiʻi at Mānoa, USA
[3]Marine Science Institute, University of the Philippines Diliman, Philippines
[4]Institute for Future Transport and Cities, Coventry University, UK
[5]Department of Biological Sciences, Central Luzon State University, Philippines
*Corresponding authors: agajigan@hawaii.edu, grieg@hawaii.edu

Abstract

Harmful algal blooms (HABs) pose a significant public health concern and can cause severe economic losses. In Bolinao, Philippines, intensive mariculture has led to diatom- or dinoflagellate-dominated HABs since the 2000s, raising risks of paralytic shellfish poisonings and fish kills. In the context of ongoing HAB monitoring and mitigation efforts, we investigated the phytoplankton and associated giant viruses during two seasons (Nov-Dec 2021 and Apr-May 2022). Our sampling encompassed a saxitoxin episode lasting 2.5 months (April-June 2022), and a fish kill (May 15-16, 2022). We used a low-cost flow-through camera (PlanktoScope) and 18S rRNA gene amplicon sequencing to identify the dominant phytoplankton. We detected the presence of *Alexandrium* sp., a known saxitoxin-producing alga, coincident with a toxin alert that resulted in a shellfish ban, even though it was not the dominant taxon. Using electron microscopy, we observed diverse morphologies of virus-like particles (VLPs), including "giant" icosahedral VLPs (~200 nm capsids). We assembled giant virus genomes from metagenomic data collected at different bloom phases. Phylogenetic and functional analyses suggest that the recovered genomes are primarily from viruses in the orders *Imitervirales* and *Algavirales*, encoding diverse auxiliary metabolic genes involved in nutrient transport and carbohydrate metabolism. The diversity and spatiotemporal dynamics of plankton and their viruses described in this study contributed to our understanding of the broader microbial context of HABs.

Importance

HABs are widespread phenomena that can be detrimental to livelihoods and health in coastal communities. Understanding what drives their formation, maintenance, and crash could aid in predicting, and potentially alleviating these events. Here, we report on the diversity and temporal dynamics of phytoplankton and giant viruses during a HAB event. Our findings revealed a diverse array of giant viruses belonging to thirteen families in a eutrophic coastal environment. We found evidence of associations between several phytoplankton-giant virus pairs and found a temporal/seasonal influence on phytoplankton-giant virus community structure. While future research is needed to establish definitive links between the giant viruses and HABs species succession, this study lays the groundwork for future mechanistic studies.

Keywords: harmful algal bloom, PlanktoScope, 18S sequencing, metagenomics, NCLDV

Abstract (219 words), Importance (114 words), Main Text (4,994 words), Figures (6)

Introduction

Harmful algal blooms (HABs), exacerbated by intensive fish farming, are a recurring problem in the coastal waters of Bolinao, NW Philippines. The Bolinao mariculture of milkfish (*Chanos chanos*) started in 1970 and intensified in 1995 [1]. The proliferation of fish pens restricts circulation which, together with nutrient and organic loading from the use of excessive fish feed, contributes to eutrophication that fuels HABs since 2000s [2–5]. Microbial respiration of organic matter from excess fish feed and crashing phytoplankton blooms contributes to severe oxygen depletion. Stratification resulting from freshwater runoff and surface warming can aggravate the problem by restricting the mixing of oxygen into waters below the thin surface layer, resulting in hypoxia and fish kills. Fish kills in Bolinao have been frequent, with the most notable events occurring in 2002 and 2010 [6–8]. The 2002 *Prorocentrum minimum* bloom lasted five days, resulting $9 million loss due to fish kill [6]. Dissolved oxygen (DO) was < 2mg/L, far from the standard ~5-6 mg/L [3]. In response, the local government enforced restrictions on the number of fish structures; however, nutrient concentrations remained high [9]. In 2010, blooms of *Alexandrium* sp. and *Skeletonema costatum* lasted two months reducing DO to < 0.5 mg L^{-1}. This cycle of eutrophication, HABs, hypoxia, and fish kills continues to be a problem, and, in 2018, Bolinao suffered a $2 million loss from another fish kill [10]. The recurrent HABs provides an opportunity to examine phytoplankton and viral dynamics during these devastating bloom events.

Viruses, which infect diverse phytoplankton[11], grazers [12], and bacteria [13] can influence HAB dynamics both directly and indirectly. Viruses that infect HAB-forming species may act directly to suppress initiation or cause termination of a HAB. Viruses are major contributors to phytoplankton mortality [14], often causing loss rates similar in magnitude to zooplankton grazing[15,16]. In some cases, viruses may be the primary cause of abrupt bloom termination [17]. Lysis of a dominant phytoplankton population drives algal species successions [18,19] and the concomitant release of dissolved organic material alters the composition and activity of the heterotrophic microbial food web [20,21]. Viral infections could also indirectly favor HAB blooms by lysing a dominant non-HAB phytoplankton and reducing resource competition. For HAB species adept at using organic nutrients, blooms might be promoted by organic nutrients released by viral lysis [22]. Thus, viruses may stimulate bloom formation by top-down (control on predators) or bottom-up (increasing nutrient supply) mechanisms. Despite the potentially significant and multifaceted contributions of viruses to bloom ecology, no studies have examined the viral communities associated with Bolinao HABs.

This study focuses on large dsDNA viruses (Phylum *Nucleocytoviricota*), a group of viruses also referred to as Nucleocytoplasmic Large DNA Viruses (NCLDV). There are many smaller RNA- or DNA-containing viruses that could also have a significant influence on HAB dynamics, but we found the NCLDVs, and members of the *Imitervirales* in particular, to be of interest because they are abundant, exceptionally diverse [23], and have large genomes with extensive metabolic genes[24,25]. Using environmental sequencing, plankton imaging via low-cost device (PlanktoScope [26]) and conventional microscopy, we documented phytoplankton and associated giant viruses dynamics across seasons and a saxitoxin HAB episode. This research provides context for future studies and contributes to ongoing efforts to create an ecological model and a HAB early-warning system for the area.

Materials and methods

Study sites and sampling

The study was conducted in Guiguiwanen Channel, Bolinao, with samples collected at six stations (**Figure 1**). Two stations were selected to leverage historical and on-going efforts: Station 1_CCMS (16.3868°N 119.9252°E) is the site of a floating platform (Continuous and Comprehensive Monitoring System) established in 2011 within an area densely packed with fish pens [9], and Station 2_BML (16.3815°N 119.9125°E) is the Bolinao Marine Laboratory monitoring station established in 1995.

Additional stations include Station 3_CH_MOUTH (16.39538°N 119.9052°E) at the northern mouth of
the channel and Station 4_CH_OUT (16.4038°N 119.9022°E) even further north. These latter stations
correspond, respectively, to the stations Guig2 and Guig1in a previous study [27]. Hydrographic data were
routinely collected at these four stations, but two additional stations further south, within less dense fish
pens (**Figure S1)** were also sampled on occasion: Station 5_Panaen and Station 6_Siapar. A Hanna
multiparameter probe (HI98194) was deployed at 1m depth to measure temperature, salinity, DO, total
dissolved solids (TDS), and pH. Sampling occurred in November-December 2021 and April-May 2022,
when HABs are most likely to occur [28]. The first sampling was done during the Northeast (NE) monsoon
(typically October to March), while the second sampling corresponds to the transition period (April) and
the onset of the Southwest (SW) monsoon (typically starts mid-May) [29,30]. A Gratuitous Permit (#0228-
22) and Export Clearance (#2022-54 and 55) were secured from the Philippines' Bureau of Fisheries and
Aquatic Resources (BFAR).
116
Phytoplankton and virus collection
Phytoplankton samples were collected using a 30-cm wide, 10-μm mesh plankton net, vertically towed
from 5 meters to the surface. Discrete seawater samples (~1 m depth) for sequencing and chlorophyll-a
analyses were obtained using a 5 L Niskin bottle. Biomass for 18S rRNA gene sequencing was collected
by filtration of ~800 mL of seawater onto 0.2 μm polycarbonate filters (Cytiva). Material for
metagenomic analysis of the total community, including viruses, was collected by filtration of ~370 mL
of seawater on 0.02 μm aluminum oxide filters (Anotop, Cytiva). Pre-filtration through 200 μm nylon
mesh was performed for all samples before PlanktoScope imaging, microscopy and biomass collection.
Phytoplankton were counted within 2 hours after collection using the PlanktoScope (live imaging) and
microscopy (preserved in Lugol's iodine).
127
128 *Chlorophyll-a measurement*
129 Seawater from each site was transported in the dark back to the lab (< 2 h) then 100 mL was filtered
130 through a 25 mm glass fiber filter (GF/F; 0.7 um) and stored in the dark at -20°C until processing.
131 Chlorophyll-a was extracted with 90% acetone and quantified by fluorometry [31].
132
PlanktoScope imaging and analysis
We built and replicated a PlanktoScope v2.1 as previously described [26]. The 2021 fieldwork served as a
trial deployment since the standardized protocol was not yet available. In 2022, imaging was done
following the standardized protocol v1[32] wherein the pump volume and image size were calibrated.
Hence, only the 2022 counts are reported, but both datasets provided insights into morphological features
that can discriminate plankton groups. On average, we collected 1000 images per sample, totaling
~42,900 images covering 23 sampling trips. The machine records the total imaged volume used for cell
count estimates. We labeled 1250 representative images as the training set. Plankton identification was
based on comparing cell gross morphology to images in existing identification guides [33,34] with an
emphasis on discriminating total diatom vs. total dinoflagellates which were the most distinctive groups
and the groups responsible for HABs in this region. We implement image contrast normalization to
ensure backgrounds with consistent color. We trained a Residual U-Net [35] to perform semantic
segmentation on the labeled data using pixel-level BCELoss, followed by noise filtering, contour-finding
algorithm [36] and entity counting. We performed the counting using the trained model seven times to get
diatom and dinoflagellate aggregate estimates.
148
Phytoplankton microscopy counting
Microscopy counting was performed in parallel with PlanktoScope counting on eight select samples (four
from station 1, four from station 4) during the April-May 2022 season. Additional microscopy counts
were conducted on a different set of samples collected biweekly at Station 1_CCMS (Feb-Apr 2022; Jan-
Oct 2023). For microscopy, a 1-mL aliquot of sample was deposited onto a Sedgewick Rafter slide and

cell counts were performed thrice per sample bottle and then averaged. Identification was made down to
the genus level based on existing guides [33,34].
156
Electron microscopy of Bolinao VLPs
Viruses in 2 mL of the 200 μm pre-filtered seawater were concentrated using Amicon Ultra-0.5 30kDa
(12/04/2021) and 16 mL in Amicon Ultra-4 100kDa (06/06/2022) centrifugal ultrafilters (Millipore
Sigma). The concentrated samples were preserved in 2% glutaraldehyde at -20^{o}C until processing. Grids
(200 mesh copper with carbon-stabilized formvar) were rendered hydrophilic by glow discharge. A 4 μL
sample was applied to the grid and viruses allowed to adsorb for 45 seconds. Sample was then wicked
away with filter paper followed by washing with distilled water and negative staining with 2% uranyl
acetate. The grid was viewed on a Hitachi HT7700 TEM at 100 kV and photographed with an AMT XR-
41B 2k x 2k CCD camera.
166
CCMS-AWQMS data collection
The Automated Water Quality Monitoring Station (AWQMS) at Station 1 is equipped with sensors for
conductivity, temperature, and depth (CTD), and DO. In situ measurements were recorded hourly.
Temperature, salinity, and chlorophyll were measured by sensors that sampled seawater through an
inflow tubing, while the DO sensor was directly submerged. These measurements complemented the data
collected using the handheld multiparameter probe noted above.
173
18S rRNA gene amplicon analysis
DNA was extracted from the 0.2 μm polycarbonate filters using a MoBio PowerSoil DNA Isolation kit.
The V9 region of the 18S rRNA gene was amplified with the Earth Microbiome Project primers (1391f-
GTACACACCGCCCGTC and EukBr-TGATCCTTCTGCAGGTTCACCTAC) then sequenced on the
Illumina MiSeq platform to generate 300 bp paired-end (PE) reads [37,38]. Sequencing was done at the
University of Hawaii Advanced Studies in Genomics, Proteomics and Bioinformatics Sequencing
Facility. Using the software package Mothur v1.48 [39], PE reads were assembled, quality filtered, and
aligned to the SILVA database v132 [40]. Sequences were checked for chimeras and clustered into
operational taxonomic units (OTUs) at a 97% similarity. Sequences were taxonomically classified using
SILVA v132. Community comparisons and statistical analysis were implemented in Mothur using
subsampled libraries to keep the number of sequences the same across samples. Additional analyses at the
level of amplicon sequence variants (ASVs) or phylotype (genus) were done. Non-metric
multidimensional scaling (NMDS) plots illustrating the relationships among samples were made using the
vegan package in R.
188
Metagenome preparation
DNA was extracted from 0.02 μm Anotops using a Zymo DNA Microprep kit, but modifying the protocol
to include proteinase K and a backflushing of the lysis buffer through the filter in the housing[41]. Lysis
buffer recovered from the filter housing with a syringe was deposited into a microtube and the remainder
of the purification steps followed the manufacturer's protocol. The resulting DNA samples were sent to
Azenta, NJ, USA, for 150 bp PE Illumina HiSeq metagenomic sequencing. Raw reads were subjected to
adapter trimming and quality filtering using Trimmomatic [42]. Reads were normalized using bbnorm
(minimum 5x and average depth of 50x) (https://sourceforge.net/p/bbmap). A mixed co-assembly
approach was implemented using MEGAHIT [43]. Reads in the same station and month were combined for
co-assembly (**Table S1 & S2**). All the resulting assemblies were combined in one file, and contigs less
than 1000 bp were removed using SeqKit [44]. Contigs were clustered at 95% nucleotide similarity using
CD-HIT [45] to remove the redundancy, resulting in 2.28 million contigs. Before the downstream analysis,
the assembly file was re-formatted to simplify names using Anvio [46]. 28,462 bins/MAGs were produced
through Bowtie2 mapping [47] and MetaBAT2 binning [48]. DAS tool was used to identify the contigs-to-bin
memberships [49] and SeqKit was used to determine the length statistics [44].
204

NCLDV identification, characterization, and co-occurrence network
Viral contigs were identified using geNomad [50] producing 1,739 candidate NCLDV bins. Only those bins
with sizes between 100kb - 3Mbp were retained, corresponding to the known smallest (103 kbp [51]) and
largest (2.77 Mbp [52,53]) NCLDVs. We retained only those bins with < 40 contigs, similar to a previous
study [54]. We also included only those bins with the longest contig >10,000 bp. These length cutoffs
resulted in 190 bins with a lower degree of fragmentation. ViralRecall [55] was used to identify the five
highly conserved NCLDV marker genes, including MCP (major capsid protein), polB (DNA polymerase
B), A32 (A32-like packaging ATPase), VLTF3 (virus-like late transcription factor), and SFII
(superfamily II helicase). Only the bins with 4 out of 5 marker genes were retained, resulting in 90 bins.
Additional NCLDV marker genes were identified, including RNAPL, RPANS (RNA polymerase large
and small subunits, mRNAc (mRNA capping enzyme), RNR (ribonucleotide reductase) and D5 (D5
primase/helicase). Only those contigs with positive ViralRecall scores were retained in each bin. In
addition, we removed bins with greater than 7 MCP copies to lessen the chance of analyzing chimeric
bins. It is reported that NCLDV can contain multiple MCP copies, for instance, BpV1 has 5 (NCBI:
NC_014765.1), PBCV-1 has 7 (NC_000852.5), OtV5 has 8 (NC_010191.2). These criteria led to a final
set of 82 bins. Genome completeness is estimated using CheckV [56] and rRNA genes were identified using
barrnap (https://github.com/tseemann/barrnap). No rRNA genes were found in the final set of 82 GV-
MAGs. Genome annotation was done using NuMP (https://github.com/BenMinch/NuMP; accessed Jan
2024) and ViralRecall [55]. KEGG metabolic module completeness was calculated using MicrobeAnnotator
via DIAMOND blastp [57]. MCP and PolB phylogenetic trees were constructed using NuPhylo
(https://github.com/ BenMinch/NuPhylo; accessed Jan 2024) and were visualized in iTOL [58]. Taxonomic
identification was done using the phylogenetic trees and confirmed using TIGTOG [59]. Anvio was used to
calculate the mean coverage[46]. The normalized relative abundance of bins is calculated by dividing the
number of recruited reads by the total number of reads. The number of recruited reads is calculated by
multiplying Anvio's mean coverage by bin length divided by read length. Bakta was used for genome
visualization [60]. Finally, after removing diatom and zooplankton 18S OTUs, a co-occurrence network
between remaining phytoplankton and GV MAGs was made using SpiecEasi in R [61]. The resulting
network was visualized using igraph and ggplot.

Results

Environmental condition, saxitoxin episode, and fish kill
Intensive mariculture activities within the Guiguiwanen Channel in Bolinao **(Figures 1 and S1)** creates a
DO and pH gradient over a short distance from the fish cages to the channel mouth **(Figures 2 and S2)**.
Our sampling covers different seasons and in 2021, the Philippine weather agency reported the onset of
the NE monsoon on October 25 [62] with a surge reported on December 2, leading to a decrease in surface
seawater temperature **(Figure 2)** [63]. Otherwise, surface temperature, salinity, and TDS were similar across
sites and seasons **(Figures 2 and S2)**, except at the start of the saxitoxin (STX) episode (April 13, 2021),
when a drop in salinity to ~26 PSU was observed. Chlorophyll-a, a rough proxy for phytoplankton
biomass, varied from 1–20 ug L^{-1} but was relatively higher during NE monsoon **(Figures 2 and S2)**.
During the second sampling, there was a 2.5-month period of paralytic shellfish poisoning (PSP) alert
triggered by elevated STX levels in shellfish tissues. Persistent detection of STX in the shellfish started
on April 7, 2022. The toxin level was reported at 249 µg STX per 100 g shellfish at the start of the ban,
beyond the 60 µg STX per 100 g shellfish limit, as reported by the local monitoring agency (BFAR) [64].
The shellfish bulletin (herein referred to as the BFAR toxin alert) was only cleared on June 17, 2022. A
fish kill event was recorded on May 15, 2022, with images from May 16 showing extensive fish mortality
near Station 1 **(Figure S3)**. Surface DO at Station 1 remained consistently low for months, sometimes
reaching hypoxia (<2 mg/L) **(Figures 2 and S4)**.

Phytoplankton monitoring and composition
Phytoplankton monitoring was conducted using three methods: [1] 18S rRNA amplicon sequencing, [2] a
PlanktoScope **(Figure S5)**, and [3] conventional microscopy identification and counting. The 18S rRNA
sequencing provided relative abundances of dominant phytoplankton with high taxonomic resolution,
while PlanktoScope provided cell count estimates with relatively low taxonomic resolution (**Figure S6**).
Our analysis focused on Stations 1 and 4 because of logistical constraints. The PlanktoScope captured
images of dominant plankton and our taxa of interest—dinoflagellates and diatoms.

Plankton diversity and succession revealed by 18S rRNA gene sequencing
Diatoms frequently dominated the amplicon pool, comprising 5–97% (66% on average) of 18S rRNA
gene sequences over the sampling period. However, on April 29 and May 5, 2022, dinoflagellate
amplicons (31% and 32% of the sequences) were more abundant than those of diatoms at Station 4
(**Figure 2 and S7**). Phytoplankton community composition significantly varied by season and, to some
degree, station **(Figure 3A)**, with consequent influences on giant virus composition **(Figure 3B)** as
discussed in the next section. Species richness and diversity increased from the NE monsoon to SW
monsoon **(Figure 3C)**. Specifically, we observed a conspicuous diatom succession from *Skeletonema*
(early November), *Chaetoceros* (late November), unclassified Mediophyceaea (end of November),
Thalassiosira (December) back to *Skeletonema* (April), then unclassified Diatomea (early May), and
mostly *Chaetoceros* (end of May) **(Figure S8)**. Succession in the dinoflagellate community was difficult
to resolve as most sequences belonged to unclassified Dinophyceae and Gymnodiniphycidae **(Figure S9)**.
Still, several dinoflagellate genera were evident, such as *Gyrodinium*, *Karlodinium*, *Scrippsiella*,
Heterocapsa, *Protoperidinium*, and *Alexandrium* **(Figure S9)**. Other notable plankton included
Chlorophyta (0.29-34%) represented by *Ostreococcus*, *Micromonas*, and *Tetraselmis* **(Figure S7)**. Non-
phytoplankton groups that were captured by 18S rRNA gene sequencing include copepods (0-35%),
tunicates (0-2.8%), and ciliates (0.14-11.6%) **(Figure S7)**.

Alexandrium sequences constituted a small fraction of the plankton community (at most 0.1% of 18S
rRNA gene sequences). Of the three STX-producing genera, *Alexandrium, Pyrodinium*, and
Gymnodinium [65], only *Alexandrium* coincides well with the BFAR STX alert **(Figure S10)**. *Pyrodinium*
was absent throughout the time series. Further analysis at the OTU and ASV levels suggested the
presence of multiple *Alexandrium* represented by two OTUs (Otu00725 & Otu01700) and seven
Alexandrium ASVs (two are singletons not included in the figure) **(Figure S11)**. Microscopy data
supported these findings, showing that *Alexandrium* sp. was present at 83–111 cells/L during the start of
the shellfish ban but represented only 0.15–0.38% of the total phytoplankton community **(Figure S12)**.

PlanktoScope performance
The correspondence of PlanktoScope and microscopy counts is better for diatoms (linear regression, $r^2 =$
0.69, n=8) than the small dinoflagellates (<40 µm) **(Figure 2)**. This is likely due to the limits of
PlanktoScope optical resolution which is best suited for plankton in the 40–200 µm range [32].
Nevertheless, we can still detect the essential features of dinoflagellates, such as girdles and horns, but not
the thecal plates **(Figure S6)**. Microscopy identified 33 plankton taxa during the 2022 sampling **(Table
S3)**. Station 1 had seven dinoflagellate genera, eleven diatom genera, and four zooplankton taxa, with
diatoms—especially *Pseudo-nitzschia, Skeletonema, and Thalassiosira*—being the most abundant.
Station 4 had ten dinoflagellate genera, twenty-five diatom genera, one cyanobacterium, and six
zooplankton taxa, with *Pseudo-nitzschia* also dominating, followed by *Chaetoceros* and *Skeletonema*.
Other notable forms in both stations were the dinoflagellate *Peridinium* sp. and zooplankton forms,
bivalve larvae, and tintinnids.

Diversity and dynamics of Bolinao giant viruses
Giant viruses (GVs) were investigated using metagenomics to examine their diversity and potential
interactions with phytoplankton**.** Based on our filtering criteria, which includes genome size and presence

of NCLDV markers, we recovered 82 taxonomically diverse GV MAGs spanning five orders:
Algavirales, *Imitervirales*, *Pimascovirales*, *Pandoravirales*, and the recently suggested new class/order
Proculvirales/-*viricetes* (PC_01) **(Figures 4A, S13, S14; Table S4)**. The GV MAGs were further
classified into 13 families: AG_01 (Prasinoviruses), AG_04, IM_01 (*Mesomimiviridae*), IM_09
(*Schizomimiviridae*), IM_12 (*Allomimiviridae*), IM_13, IM_16 (*Mimiviridae*), PM_01, PM_06, PM_07
(*Pithoviridae*), PV_03 and PV_05 (*Coccolithoviridae*) **(Table S4)**. One GV MAG (GV78) was putatively
classified within newly identified taxa, PC_01, thought to be exclusive in the Southern and the Arctic
Oceans [66]. These GV MAGs have varying CheckV quality and completeness from high quality (n=18)
and medium quality (n=41) to low quality (n=24) **(Table S4)**, with the largest (GV18) being 960 kb and
smallest (GV82) with a 101 kb genome. Electron microscopy confirmed the presence of virus-like
particles (VLPs), with diverse morphologies including ovoid, filamentous, isometric, spindled, and tailed
VLPs **(Figure S15)** and large icosahedral giant virus-like particles (~170-200 nm) **(Figure 4B).**
319
These GV-MAGs exhibit extensive metabolic capacities, with genes involved in glycolysis,
gluconeogenesis, tricarboxylic acid (TCA) cycle, light harvesting/photosynthesis, cytoskeleton,
transporter, and nutrient metabolism **(Figure 5)**. One GV-MAG code for an ammonium transporter
(GV17) and six coded for phosphate transporters (GV11, GV19, GV21, GV28, GV57, and GV71).
(Figure 6A). The relative abundance of GV-MAGs varied seasonally, with three main clusters,
representing roughly the months of sampling and monsoon season—November (Cluster 1; NE monsoon),
April (Cluster 2; Transition period), and May (Cluster 3; onset of SW monsoon) **(Figure 6A)**. We further
examined the functional capacity of our GV-MAGs by estimating the presence and completeness of
KEGG metabolic modules **(Figure S16)**. Present in the transition period and SW monsoon clusters are
the beta-lactam biosynthesis pathway, cysteine and methionine metabolism and drug resistance pathways
(Figure S16). Among all the GV-MAGs, the largest (GV18) had the most extensive metabolic capacity
and the most copies of glycolysis and pentose phosphate pathway genes **(Figure S17)**. GV01 and GV02
were notable for containing several copies of 'chitin response' and lipid metabolism genes. GV01
contains a rhodopsin and a chlorophyll AB binding protein, while GV02 contains carbonate anhydrase.
GV02 also encodes for a sugar transporter (MtN3_slv) and apoptosis inhibition-associated gene (Bax1-I).
335
We analyzed the co-occurrence of phytoplankton and giant viruses to gain insight into putative host-virus
associations linking 495 phytoplankton OTUs with 82 giant viruses **(Figure 6B and S18; Table S5).** We
focused on phytoplankton hosts known to be infected by giant viruses; thus, we removed diatom and
zooplankton OTUs. Known virus-host pairs are captured in the network. The high frequency of green
algae and dinoflagellate co-occurring with *Algavirales* and *Imitervirales* (known viral orders infecting
algae) is of interest. For instance, *Ostreococcus* (Otu00039) co-occurs with GV53, family AG_01, the
known Prasinovirus family. Another example is Prymnesiophyceae (*Chrysochromulina*; Otu00104 &
Otu01410), which co-occurs with IM_01 (in which Chrysochromulina parva virus belongs). One
Tetraselmis (Otu000169) co-occurs with three Algavirales GV15, GV54, and GV51. The relative
abundance of *Alexandrium* was too low to be captured by the network, but several dinoflagellate OTUs,
such as *Gyrodinium* (Otu00034), *Karlodinium* (Otu00209), *Pelagodinium* (Otu00906), and *Heterocapsa*
(Otu00213), were found to be associated with GVs belonging to *Imitervirales* families.
348
349 **Discussion**
350
The Bolinao mariculture area has experienced repeated eutrophication-induced HABs for years [3,9] as a
result of the high concentrations of fish structures and nutrient input from excess fish feed. Consequently,
DO and pH vary over a short distance from mariculture-intensive area to the channel mouth likely due to
the decomposition of organic matter that consumes oxygen and produces carbon dioxide that lowers pH
[67]. During this study, the causative toxic HAB species was likely *Alexandrium* sp., as supported by our
observations (~0.1% 18S sequences, ~0.35% phytoplankton counts, around 83–111 cells L^{-1}) and
consistent with historical accounts [5,7,68]. This is in contrast to many other areas of the Philippines (e.g.

Sorsogon Bay, and Eastern Visayas embayments, and Dumanquillas Bay to name a few) where
Pyrodinium is the primary STX-producing species [69–72]. Even low concentrations of toxin-producing cells
can result in high risks of PSP. In a 2014 PSP episode involving *Alexandrium tamiyavanichii,* cells
reached only 840 cells L^{-1} and resulted in 3,500 μg STX per 100 g shellfish in Kuantan port, Malaysia [73].
Likewise, in Bolinao, *Alexandrium minutum* concentrations of only 5–7 cells L^{-1} were associated with
accumulation of 930–2,894 μg STX per 100 g shellfish meat [74]. This is consistent with our observation
that, despite a PSP alert, *Alexandrium* was a minor component of the phytoplankton community at the
time. This highlights the need for rapid, sensitive, and specific methods for detecting toxin-producing
species.
367
Typically, a drop in salinity and DO is observed during HABs and fish kill in Bolinao. The 2010
Alexandrium toxin bloom was associated with low salinity (26–27) and low DO (<2.8 mg L^{-1}, as low as
0.3 mg L^{-1} at the bottom) [8], consistent with the current study. *Alexandrium* species are known to survive
and produce more toxins at low salinity [75]. The observed DO decline during this study was also associated
with a fish kill. These DO estimates are surface measurements; thus, we expect an even lower DO at the
deeper portion of the water column, which could be lethal for fish, especially demersal species. DO
declined several times in a 2-year period, but not every drop in DO leads to a fish kill (**Figure S4**); thus,
the May 15 fish kill event may have other, or at least additional, causes such as toxins, ammonia
poisoning, or gill obstruction.
377
The overall phytoplankton community observed in this study is typical of eutrophic coastal areas
dominated by diatoms and dinoflagellates [76,77] along with other common marine taxa such as chlorophytes
and haptophytes [78]. A 2018 Bolinao survey reported similar composition, with dinoflagellates (~35–65%)
and diatoms (~30–60%) as the dominant taxa in the inner Guiguiwanen channel [27]. Monsoon seasons
were the major driver of community changes consistent with previous work [79,80]. The relative uniformity
across stations within a season may reflect the close spacing between the sampling sites (a few km).
384
Aside from the 18S amplicon approach, we tested the utility of PlanktoScope, a community-driven, open-
access, low-cost camera [26]. We illustrated the applicability of the machine for estimating diatoms but had
less success counting and identifying dinoflagellates. Increasing the optical resolution, and perhaps
refining the detection algorithm, could improve this situation. A newer, more stable version of the
PlanktoScope machine (v2.6) became available after completion of this study (we used v2.1). Adoption of
this newer model might also improve count accuracy. The PlanktoScope, being the most economical and
accessible machine (compared to FlowCam and Imaging FlowCytobot), might be a means to improve
monitoring coverage, especially in resource-poor settings. The one described in this study represents our
first attempt at utilizing PlanktoScope in monitoring HABs and has shown promise.
394
In addition to phytoplankton, we examined GV diversity and its spatiotemporal variation. As expected,
the Bolinao GV MAGs encoded extensive suites of metabolic genes, including nutrient transport that
could transiently alter the nitrogen and phosphorus acquisition capabilities of the infected host cell (the
virocell). Such a virus-mediated alteration of host phenotype was observed during infections of the green
alga *Ostreococcus tauri* with the virus OtV6. In that study, the virus-encoded ammonium transporter
expressed during infection appears to have a higher affinity, but lower V_{max}, for methyl-ammonium than
the host-encoded transporter, which is presumed to be advantageous in nitrogen-limited waters [81]. The
Bolinao GV MAGs encoding for ammonium (n=1) and phosphate transporters (n=6), may similarly
reflect adaptations to nitrogen or phosphate limitation in Bolinao waters [3,9]. The higher frequency of
phosphate transporters may be a reflection of the high nucleic acid content of virions, which increases
their relative P requirement compared to that of the cells in which they replicate[82]. The Bolinao GV
MAGs also encode a whole suite of energy-generating metabolic enzymes related to photosynthesis,
glycolysis, pentose phosphate pathway, and TCA cycle, which presumably either augment or replace host
functions during energy-demanding processes of viral replication. One GV MAG encodes for carbonic

anhydrase that can influence the carbon concentrating mechanism and perhaps aid photosynthesis in the
phytoplankton virocell [83,84]. In the clusters representing the transition period and SW monsoon samples,
KEGG pathways related to antioxidants are enriched, which may reflect selection for viruses that have
capacity to counter oxidative stress during infection [85,86]. In addition, KEGG pathways related to
multidrug resistance efflux pumps are also enriched during these months, which might be related to
removal of antiviral toxins [87].
415
The giant viruses described here were likely not involved in direct bloom termination, as the
phytoplankton community was dominated by diatoms which seem to be infected only by small viruses [88].
However, the putative phytoplankton-GV associations described here, and their changes over time,
suggest that the GV are involved in community dynamics during Bolinao HABs, similar to what is
observed in HABs elsewhere [89]. GV could be involved in the demise and succession of HABs at other
times, especially for blooms dominated by dinoflagellates for which some giant viruses are known. In
such cases, viruses might be more efficient in killing HAB species than zooplankton which are usually
deterred by toxins [90,91]. Several single species blooms have occurred in Bolinao. For instance, in 2002, a
mono-species bloom of *P. minimum* (99%) lasted several days, succeeded by diatoms (87%) after 4 days.
In 2010, an *Alexandrium* sp. bloom (~86%) succeeded by a *S. costatum* bloom (99%) was also recorded.
In 2018, a mono-species *Takayama* sp. bloom was associated with a fish kill event [27]. Aside from the
direct impact on the bloom, viral infection will have pronounced consequences on carbon export and
microbial succession. For instance, it was shown that viral infection causes a 2-4-fold increase in carbon
release and a rise of not only bacteria but also eukaryotic heterotrophic organic matter recyclers [92].
430
By focusing only on large dsDNA-containing viruses, we did not account for other viruses that may have
influenced bloom dynamics, especially those infecting diatoms. Most diatom viruses are small and
contain ssRNA or ssDNA [88,93,94]. Nonetheless, this study expands our knowledge on giant virus diversity
and spatiotemporal dynamics in the context of an STX episode and across a eutrophic coastal
environment and monsoonal seasons. Some research groups are exploring viruses as HAB mitigation.
One study has shown a significant reduction in the density of the harmful algae, *H. circularisquama*, with
the application of either virus alone or virus-sediment mix in a mesocosm experiment [95]. This underscores
the urgency and importance of understanding phytoplankton-virus interaction and dynamics in various
contexts to fully comprehend the impact of such mitigation efforts.
440
441 **Conclusion**
442
This study examines phytoplankton diversity and identifies associated giant viruses during different
monsoon seasons, a toxic bloom, and a fish kill, contributing to the growing literature on HAB dynamics.
We used a complementary method for phytoplankton/HABs tracking that provided data on *Alexandrium*
sp. appearance coinciding with STX accumulation in shellfish. Further fine-tuning of PlanktoScope to
better detect and count dinoflagellates in addition to diatoms will increase the efficiency of HAB
monitoring. We documented the diversity of giant viruses, but further work is needed to link specific
viruses and their hosts and determine the influence of viral lysis on the bloom dynamics. Bolinao HABs
remains a recurring problem and detailed investigations into the various sources of algal mortality may
shed light on the variables influencing bloom severity and duration.

References

1. Verceles, L. F., Talaue-McManus, L. & Aliño, P. M. Participatory Monitoring and Feedback System: An Important Entry Towards Sustainable Aquaculture in Bolinao, Northern Philippines. *Science Diliman* **12**, 78–87 (2000).

2. David, L. T. *et al.* The saga of community learning: Mariculture and the Bolinao experience. *Aquatic Ecosystem Health and Management* **17**, 196–204 (2014).

3. San Diego-McGlone, M. L., Azanza, R. V., Villanoy, C. L. & Jacinto, G. S. Eutrophic waters, algal bloom and fish kill in fish farming areas in Bolinao, Pangasinan, Philippines. *Marine pollution bulletin* **57**, 295–301 (2008).

4. Obliosca, J. M., Jacinto, G. S. & San Diego-McGlone, M. L. Preliminary results on the development of marine environmental quality criteria for mariculture areas. *Philippine Science Letters* **40**, 73–87 (2003).

5. Yñiguez, A. T. *et al.* Over 30 years of HABs in the Philippines and Malaysia: What have we learned? *Harmful Algae* (2020) doi:10.1016/j.hal.2020.101776.

6. Azanza, R. V., Fukuyo, Y., Yap, L. G. & Takayama, H. Prorocentrum minimum bloom and its possible link to a massive fish kill in Bolinao, Pangasinan, Northern Philippines. *Harmful Algae* **4**, 519–524 (2005).

7. Azanza, R. V. & Benico, G. A. Toxic Alexandrium blooms in Fish Farming Sites in Bolinao, Pangasinan. *Journal of Environmental Science and Management* 44–49 (2013).

8. Escobar, Ma. T. L. *et al.* Eutrophic conditions during the 2010 fish kill in Bolinao and Anda, Pangasinan, Philippines. *Journal of Environmental Science and Management* 29–35 (2013).

9. Ferrera, C. M. *et al.* Phosphorus as a driver of nitrogen limitation and sustained eutrophic conditions in Bolinao and Anda, Philippines, a mariculture-impacted tropical coastal area. *Marine Pollution Bulletin* **105**, 237–248 (2016).

10. Austria, H. & Pasion, A. Anda, Bolinao suffer P107-M fish kill losses. *Philippine News Agency* https://www.pna.gov.ph/articles/1037212 (2018).

11. Short, S. M., Staniewski, M. A., Chaban, Y. V., Long, A. M. & Wang, D. Diversity of viruses infecting eukaryotic algae. *Current Issues in Molecular Biology* **39**, 29–62 (2020).

12. Fromm, A. *et al.* Single-cell RNA-seq of the rare virosphere reveals the native hosts of giant viruses in the marine environment. *Nature Microbiology* **9**, 1619–1629 (2024).

13. Riemann, L. & Middelboe, M. Viral lysis of marine bacterioplankton: Implications for organic matter cycling and bacterial clonal composition. *Ophelia* **56**, 57–68 (2002).

14. Brussaard, C. P. D. & Martínez, J. Algal Bloom Viruses. *Plant viruses* **2**, 1–10 (2008).

15. Baudoux, A. C., Noordeloos, A. A. M., Veldhuis, M. J. W. & Brussaard, C. P. D. Virally induced mortality of *Phaeocystis globosa* during two spring blooms in temperate coastal waters. *Aquatic Microbial Ecology* **44**, 207–217 (2006).

16. Brussaard, C. P. D., Kuipers, B. & Veldhuis, M. J. W. A mesocosm study of *Phaeocystis globosa* population dynamics: I. Regulatory role of viruses in bloom control. *Harmful Algae* **4**, 859–874 (2005).

17. Nagasaki, K., Ando, M., Itakura, S., Imai, I. & Ishida, Y. Viral mortality in the final stage of *Heterosigma akashiwo* (Raphidophyceae) red tide. *Journal of Plankton Research* **16**, 1595–1599 (1994).

18. Bratbak, G., Egge, J. K. & Heldal, M. Viral mortality of the marine alga *Emiliania huxleyi* (Haptophyceae) and termination of algal blooms. *Marine Ecology Progress Series* **93**, 39–48 (1993).

19. Pound, H. L. *et al.* The "Neglected Viruses" of Taihu: Abundant Transcripts for Viruses Infecting Eukaryotes and Their Potential Role in Phytoplankton Succession. *Frontiers in Microbiology* **11**, 1–11 (2020).

20. Brussaard, C. P. D., Mari, X., Bleijswijk, J. D. L. V. & Veldhuis, M. J. W. A mesocosm study of *Phaeocystis globosa* (Prymnesiophyceae) population dynamics: II. Significance for the microbial community. *Harmful Algae* **4**, 875–893 (2005).

21. Sheik, A. R. *et al.* Responses of the coastal bacterial community to viral infection of the algae Phaeocystis globosa. *ISME Journal* **8**, 212–225 (2014).

22. Gobler, C. J. *et al.* Ecology of phytoplankton communities dominated by *Aureococcus anophagefferens*: the role of viruses, nutrients, and microzooplankton grazing. *Harmful Algae* **3**, 471–483 (2004).

23. Mihara, T. *et al.* Taxon Richness of "Megaviridae" Exceeds those of Bacteria and Archaea in the Ocean. *Microbes and environments* **33**, 162–171 (2018).

24. Brahim Belhaouari, D. *et al.* Metabolic arsenal of giant viruses: Host hijack or self-use? *eLife* **11**, e78674
(2022).
25. Moniruzzaman, M. *et al.* Virologs, viral mimicry, and virocell metabolism: the expanding scale of cellular
functions encoded in the complex genomes of giant viruses. *FEMS Microbiology Reviews* **47**, fuad053
(2023).
26. Pollina, T. *et al.* PlanktoScope: Affordable Modular Quantitative Imaging Platform for Citizen
Oceanography. *Front. Mar. Sci.* **9**, 949428 (2022).
27. Albelda, R. L. *et al.* Summer phytoplankton community structure and distribution in a mariculture-
affected coastal environment. *Philippine Science Letters* **12**, 157–166 (2019).
28. Yñiguez, A. T. *et al.* Technical Terminal Report: Detection and Mitigation Technology and Early
Warning System for Philippines Harmful Algal Blooms (HABTech). 71 (2015).
29. Matsumoto, J., Olaguera, L. M. P., Nguyen-Le, D., Kubota, H. & Villafuerte, M. Q. Climatological
seasonal changes of wind and rainfall in the Philippines. *Intl Journal of Climatology* **40**, 4843–4857
(2020).
30. Wang, B. & LinHo. Rainy Season of the Asian–Pacific Summer Monsoon*. *J. Climate* **15**, 386–398
(2002).
31. Parsons, T. R., Maita, Y., Lalli, C. M. & LALLI, C. M. Determination of Chlorophylls and Total
Carotenoids: Spectrophotometric Method. in *A Manual of Chemical & Biological Methods for Seawater*
Analysis (eds. PARSONS, T. R., MAITA, Y. & LALLI, C. M. B. T.) 101–104 (Pergamon, Amsterdam,
1984). doi:https://doi.org/10.1016/B978-0-08-030287-4.50032-3.
32. Lombard, F. Planktoscope protocol for plankton imaging V.1.
dx.doi.org/10.17504/protocols.io.bp2l6bq3zgqe/v1 (2022).
33. *Marine Phytoplankton of the Western Pacific*. (Tokyo : Kouseisha Kouseikaku Co., Ltd., 2012).
34. *Identifying Marine Phytoplankton*. (Academic Press, San Diego, 1997).
35. Ronneberger, O., Fischer, P. & Brox, T. U-Net: Convolutional Networks for Biomedical Image
Segmentation. in *Medical Image Computing and Computer-Assisted Intervention – MICCAI 2015* (eds.
Navab, N., Hornegger, J., Wells, W. M. & Frangi, A. F.) 234–241 (Springer International Publishing,
Cham, 2015).
36. Suzuki, S. & Abe, K. Topological structural analysis of digitized binary images by border following.
Computer Vision, Graphics, and Image Processing **30**, 32–46 (1985).
37. Amaral-Zettler, L. A., McCliment, E. A., Ducklow, H. W. & Huse, S. M. A method for studying protistan
diversity using massively parallel sequencing of V9 hypervariable regions of small-subunit ribosomal
RNA Genes. *PLoS ONE* **4**, 1–9 (2009).
38. Stoeck, T. *et al.* Multiple marker parallel tag environmental DNA sequencing reveals a highly complex
eukaryotic community in marine anoxic water. *Molecular Ecology* **19**, 21–31 (2010).
39. Schloss, P. D. *et al.* Introducing mothur: open-source, platform-independent, community-supported
software for describing and comparing microbial communities. *Appl Environ Microbiol* **75**, (2009).
40. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing and web-
based tools. *Nucleic Acids Res* **41**, (2013).
41. Mueller, J. A., Culley, A. I. & Steward, G. F. Variables Influencing Extraction of Nucleic Acids from
Microbial Plankton (Viruses, Bacteria, and Protists) Collected on Nanoporous Aluminum Oxide Filters.
Appl Environ Microbiol **80**, 3930–3942 (2014).
42. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina Sequence Data.
Bioinformatics (2014) doi:10.1093/bioinformatics/btu170.
43. Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-node solution for
large and complex metagenomics assembly via succinct *de Bruijn* graph. *Bioinformatics* **31**, 1674–1676
(2015).
44. Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File
Manipulation. *PLoS ONE* **11**, e0163962 (2016).
45. Huang, Y., Niu, B., Gao, Y., Fu, L. & Li, W. CD-HIT Suite: a web server for clustering and comparing
biological sequences. *Bioinformatics* **26**, 680–682 (2010).
46. Eren, A. M. *et al.* Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ* **3**, e1319
(2015).

47. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nature Methods* **9**, 357–359
(2012).
48. Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient genome
reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).
49. Sieber, C. M. K. *et al.* Recovery of genomes from metagenomes via a dereplication, aggregation and
scoring strategy. *Nat Microbiol* **3**, 836–843 (2018).
50. Camargo, A. P. *et al.* Identification of mobile genetic elements with geNomad. *Nat Biotechnol* (2023)
doi:10.1038/s41587-023-01953-y.
51. Tidona, C. A. & Darai, G. The Complete DNA Sequence of Lymphocystis Disease Virus. *Virology* **230**,
207–216 (1997).
52. Philippe, N. *et al.* Pandoraviruses: Amoeba viruses with genomes up to 2.5 Mb reaching that of parasitic
eukaryotes. *Science* **341**, 281–286 (2013).
53. Queiroz, V. F. *et al.* Amoebae: Hiding in Plain Sight: Unappreciated Hosts for the Very Large Viruses.
Annu. Rev. Virol. **9**, 79–98 (2022).
54. Moniruzzaman, M., Martinez-Gutierrez, C. A., Weinheimer, A. R. & Aylward, F. O. Dynamic genome
evolution and complex virocell metabolism of globally-distributed giant viruses. *Nature Communications*
11, 1–11 (2020).
55. Aylward, F. O. & Moniruzzaman, M. ViralRecall—A Flexible Command-Line Tool for the Detection of
Giant Virus Signatures in 'Omic Data. *Viruses* **13**, 150 (2021).
56. Nayfach, S. *et al.* CheckV assesses the quality and completeness of metagenome-assembled viral
genomes. *Nat Biotechnol* **39**, 578–585 (2021).
57. Ruiz-Perez, C. A., Conrad, R. E. & Konstantinidis, K. T. MicrobeAnnotator: a user-friendly,
comprehensive functional annotation pipeline for microbial genomes. *BMC Bioinformatics* **22**, 11 (2021).
58. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and
annotation. *Nucleic Acids Research* **49**, W293–W296 (2021).
59. Ha, A. D. & Aylward, F. O. Automated classification of giant virus genomes using a random forest model
built on trademark protein families. *npj Viruses* **2**, 9 (2024).
60. Schwengers, O. *et al.* Bakta: rapid and standardized annotation of bacterial genomes via alignment-free
sequence identification: Find out more about Bakta, the motivation, challenges and applications, here.
Microbial Genomics **7**, (2021).
61. Kurtz, Z. D. *et al.* Sparse and Compositionally Robust Inference of Microbial Ecological Networks. *PLoS*
Comput Biol **11**, e1004226 (2015).
62. DOST-PAGASA. Onset of the Northeast Monsoon. (2021).
63. DOST-PAGASA. Special Weather Outlook for the Bonifacio Day (November 30) and National COVID-
19 Vaccination Dats (November 29-December 01). (2021).
64. DA BFAR. Local Shellfish Advisory No.1: Coastal Waters of Bolinao, Pangasinan Positive of Paralytic
Shellfish Poison. (2022).
65. Beasley, V. R. Harmful Algal Blooms (Phycotoxins). in *Reference Module in Earth Systems and*
Environmental Sciences B9780124095489112758 (Elsevier, 2020). doi:10.1016/B978-0-12-409548-
9.11275-8.
66. Gaïa, M. *et al.* Mirusviruses link herpesviruses to giant viruses. *Nature* **616**, 783–789 (2023).
67. Isah, R. R., Enochs, I. C. & San Diego-McGlone, M. L. Sea surface carbonate dynamics at reefs of
Bolinao, Philippines: Seasonal variation and fish mariculture-induced forcing. *Front. Mar. Sci.* **9**, 858853
(2022).
68. Baula, I. U., Azanza, R. V., Fukuyo, Y. & Siringan, F. P. Dinoflagellate cyst composition, abundance and
horizontal distribution in Bolinao, Pangasinan, Northern Philippines. *Harmful Algae* **11**, 33–44 (2011).
69. Azanza, R. V. *et al.* Expansion of Toxic Algal Blooms in Coastal and Marine Areas in the Philippines and
Malaysia: Is It Climate Change Related? *Sustainability* **16**, 3304 (2024).
70. Ravelo, S. F. *et al.* A Snapshot on the Distribution of Coastal Phytoplankton Communities in Five HAB-
Affected Bays in Eastern Visayas, Philippines. *Front. Mar. Sci.* **9**, 730518 (2022).
71. Usup, G. & Azanza, R. V. Physiology and bloom dynamics of the tropical dinoflagellate *Pyrodinium*
bahamense. in *In Physiological Ecology of Harmful Algal Bloom* (1998).

72. Yñiguez, A. T. *et al.* Investigating the roles of intrinsic and extrinsic factors in the blooms of *Pyrodinium*
bahamense var. compressum using an individual-based model. *Procedia Environmental Sciences* **13**,
1462–1476 (2012).
73. Mohammad-Noor, N. *et al.* First report of paralytic shellfish poisoning (PSP) caused by *Alexandrium*
tamiyavanichii in Kuantan Port, Pahang, East Coast of Malaysia. *Phycological Research* **66**, 37–44
(2018).
74. Bajarias, F. *et al.* Paralytic shellfish poisoning due to *Alexandrium minitum* Halim in Nortwestern
Philippines. T. in *The First Joint Seminar in Coastal Oceanography* (Thailand, 2003).
75. Lim, P.-T. & Ogata, T. Salinity effect on growth and toxin production of four tropical Alexandrium
species (Dinophyceae). *Toxicon* **45**, 699–710 (2005).
76. Annabi-Trabelsi, N. *et al.* Effects of Eutrophication on Plankton Abundance and Composition in the Gulf
of Gabès (Mediterranean Sea, Tunisia). *Water* **14**, 2230 (2022).
77. Bužančić, M., Ninčević Gladan, Ž., Marasović, I., Kušpilić, G. & Grbec, B. Eutrophication influence on
phytoplankton community composition in three bays on the eastern Adriatic coast. *Oceanologia* **58**, 302–
316 (2016).
78. Pierella Karlusich, J. J., Ibarbalz, F. M. & Bowler, C. Phytoplankton in the *Tara* Ocean. *Annu. Rev. Mar.*
Sci. **12**, 233–265 (2020).
79. De La Cruz, M. A. M., Hingpit, B. W., Guillou, L. & Onda, D. F. L. Effects of monsoons and storms on
the structuring and diversity of picoeukaryotic microbial communities in a tropical coastal environment.
Deep Sea Research Part II: Topical Studies in Oceanography **209**, 105294 (2023).
80. Yap, L. G., Azanza, R. V. & Talaue-McManus, L. The community composition and production of
phytoplankton in fish pens of Cape Bolinao, Pangasinan: A field study. *Marine Pollution Bulletin* **49**,
819–832 (2004).
81. Monier, A. *et al.* Host-derived viral transporter protein for nitrogen uptake in infected marine
phytoplankton. *Proc. Natl. Acad. Sci. U.S.A.* **114**, (2017).
82. Jover, L. F., Effler, T. C., Buchan, A., Wilhelm, S. W. & Weitz, J. S. The elemental composition of virus
particles: implications for marine biogeochemical cycles. *Nat Rev Microbiol* **12**, 519–528 (2014).
83. Badger, M. R. The Role of Carbonic Anhydrase in Photosynthesis. *Limnol. Oceanogr.* (1994).
84. Tortell, P. D. Evolutionary and ecological perspectives on carbon acquisition in phytoplankton. *Limnology*
& Oceanography **45**, 744–750 (2000).
85. Kang, M. *et al.* Chlorovirus PBCV-1 Encodes an Active Copper-Zinc Superoxide Dismutase. *J Virol* **88**,
12541–12550 (2014).
86. Lartigue, A. *et al.* The Megavirus Chilensis Cu,Zn-Superoxide Dismutase: the First Viral Structure of a
Typical Cellular Copper Chaperone-Independent Hyperstable Dimeric Enzyme. *J Virol* **89**, 824–832
(2015).
87. Nishino, K., Yamasaki, S., Nakashima, R., Zwama, M. & Hayashi-Nishino, M. Function and Inhibitory
Mechanisms of Multidrug Efflux Pumps. *Front. Microbiol.* **12**, 737288 (2021).
88. Arsenieff, L., Kimura, K., Kranzler, C. F., Baudoux, A.-C. & Thamatrakoln, K. Diatom Viruses. in *The*
Molecular Life of Diatoms (eds. Falciatore, A. & Mock, T.) 713–740 (Springer International Publishing,
Cham, 2022). doi:10.1007/978-3-030-92499-7_24.
89. Moniruzzaman, M. *et al.* Diversity and dynamics of algal Megaviridae members during a harmful brown
tide caused by the pelagophyte, Aureococcus anophagefferens. *FEMS Microbiology Ecology* **92**, 1–10
(2016).
90. Lürling, M. Grazing resistance in phytoplankton. *Hydrobiologia* **848**, 237–249 (2021).
91. Zheng, Y., Gong, X. & Gao, H. Selective grazing of zooplankton on phytoplankton defines rapid algal
succession and blooms in oceans. *Ecological Modelling* **468**, 109947 (2022).
92. Vincent, F. *et al.* Viral infection switches the balance between bacterial and eukaryotic recyclers of
organic matter during coccolithophore blooms. *Nat Commun* **14**, 510 (2023).
93. Tomaru, Y. *et al.* Isolation and characterization of a single-stranded RNA virus that infects the marine
planktonic diatom Chaetoceros sp. (SS08-C03). *Phycological Research* **61**, 27–36 (2013).
94. Tomaru, Y. *et al.* New single-stranded DNA virus with a unique genomic structure that infects marine
diatom *Chaetoceros setoensis*. *Scientific Reports* **3**, (2013).

95. Nakayama, N., Hamaguchi, M., Yamaguchi, H., Masuda, K. & Fujiwara, M. Evaluation of a virus-based
control method to protect cultured oysters from the harmful dinoflagellate *Heterocapsa circularisquama*.
Aquaculture **529**, (2020).
665

Acknowledgments

This project was funded by the National Geographic Early Career Grant (EC-KOR-66418R-21), East-West Center Field Research Grant, Experiment's Ocean Solutions Grant, and Uehiro Fellowship to APG. We are indebted to Tina Carvalho at the Microscopy Facility at the PBRC for all the electron microscopy training and usage. We thank the Rappé lab (UHM) and Dr. Charissa Ferrera (UPD) for providing some materials and lending equipment. We are thankful for the Microbial Oceanography Lab, led by Dr. Deo Onda (UPD) through the UPGRADE CIA project funded by the National Security Council for providing supplies and a DNA extraction kit. The PlanktoScope was made with the help of Jie Zhou (Fabrication Lab, UHM College of Engineering), Ian Quiño Fernandez, Eric Kolb, Dr. Manu Prakash, Dr. Adam Larson, Thibaut Pollina, and the rest of the PlanktoScope community. Niño Posadas, Jake Baquiran, Dr. Malou San Diego-McGlone, Miraflor Sanchez, Dr. Josefa Pante, Dr. Van Rodriguez, and Abby Melendres were pivotal in completing the required paperwork, research agreements, and permits. Erika Gernato and Raffi Isah assisted in obtaining materials, reagents, and instruments. We are also grateful to Cha Caalim, Rubie Esmolo, Angel Doctor, and the rest of the Bolinao Marine Lab research assistants and boatman for the marine lab access, sampling, and conducting community outreach in Bolinao.

Data Accessibility

The data used in this paper have been deposited in the NCBI Short Read Archive as BioProject PRJNA1237401 for 18S sequences and BioProject PRJNA1238504 for metagenomic sequences. The GV-MAGs assembly and annotation are available at FigShare https://figshare.com/s/79c4bebb28187961279b. The codes for PlanktoScope counting are available at https://github.com/jjshoots/PlanktonSegmentationTrainer.

Author Contributions

APG, CC, ATY, and GFS conceived of the study. APG and GCMBA performed the sampling and experiments. APG, GCMBA, JJT, and GAB performed the analysis. ELDG, RBF, CLAR, and CLV contributed data. APG and GFS wrote the paper. All authors edited and approved the manuscript.

Conflict of Interest

The authors declare no conflict of interest.

Figures and Legends

Figure 1. Bolinao map and sampling sites. Bolinao is situated in NW Philippines. We focused our sampling on the Guiguiwanen channel, covering mariculture sites (Stations 1 and 2) and relatively offshore sites (Stations 3 and 4). We sampled inward the channel (5 and 6) whenever logistics and time permits.

724
725

726
727
728
Figure 2. Time series of physical, chemical, and phytoplankton data. [From top to bottom] Plots showing similar
surface temperature and salinity ranges. A decrease in salinity was also observed at the onset of saxitoxin alert. In
contrast, a gradient of dissolved oxygen was observed. Chlorophyll-a fluctuates, reaching up to 20ug/L, and is
relatively higher in Nov-Dec 2021. Diatoms are the most abundant phytoplankton; however, in some cases,
dinoflagellates reach relatively high abundance (Station 4; April 29 & May 18). *Alexandrium* appearance coincides
with BFAR saxitoxin alert. No *Pyrodinium* sequences were detected, while *Gymnodinium* was present and in low
18S quantity. Finally, PlanktoScope counts agree with microscopy counting for diatoms (r^2=0.69) but had difficulty
counting dinoflagellates accurately because of their relatively small size. Moreover, a fish kill event happened on
May 15-16, 2022.
738
739
740

749
750
Figure 3. Community structure and diversity of phytoplankton and giant viruses. Non-metric multidimensional
scaling plots illustrate seasonal and spatial variations in (A) phytoplankton communities based on 18S rRNA gene
amplicons and (B) giant viruses based on metagenome-assembled genomes (GV-MAGs). Symbols are colored
according to season and symbol shapes reflect the stations from which samples were collected. (C) The richness and
diversity indices of phytoplankton derived from 18 S rRNA gene amplicon data show an increasing trend from NE
monsoon, though the transitional period, to the SW monsoon. P-value based on ANOVA test.

Figure 4. Phylogeny and morphology of Bolinao giant viruses. (A) Phylogeny of GV MAGs based on PolB sequences. Highlighted in black are the Bolinao GV MAGs. Written in dark blue are representative sequences from different NCDLV families, as provided by the NuPhylo package. For complete taxonomy, see supplementary materials. **(B)** A survey of viral-like particles (VLPs) in Bolinao waters shows VLPs with sizes ~170-200 nm.

Figure 5. Auxiliary metabolic genes and virologs of Bolinao GV MAGs. Curated auxiliary metabolic genes or virologs (colored based on pathways). The bubble size is based on the number of copies in the genome. The gene set is based on Ha et al. 2021 and implemented in the NuMP package.

791 **Figure 6. Temporal GV dynamics and phytoplankton-GV associations.** (A) Heatmap showing the relative distribution of GV MAGs and (B) Frequency of
792 plankton-GV associations based on their corresponding taxa.