Title: Non-redundant cardiolipin synthases shape membrane composition and support stress resilience in *Bacteroides fragilis*

Authors: Matthew K. Schnizlein[1], BongJin Hong[1,2], Jennifer N.T. Nguyen[3], Katarina Jones[4,5], Alyssa I. Rodriguez[3], Aretha Fiebig[1,6], Shawn R. Campagna[4,5], Marcy J. Balunas[3,7], Thomas V. O'Halloran[1,2,8], Sean Crosson[1,6,*]

*corresponding author: crosson4@msu.edu

Affiliations

[1]Department of Microbiology, Genetics & Immunology, Michigan State University, East Lansing, MI, USA

[2]Elemental Health Institute, Michigan State University, East Lansing, MI, USA

[3]Department of Medicinal Chemistry, University of Michigan, Ann Arbor, MI, USA

[4]Biological and Small Molecule Mass Spectrometry Core, University of Tennessee-Knoxville, TN, USA

[5]Department of Chemistry, University of Tennessee-Knoxville, TN, USA

[6]BioMolecular Science Gateway, Michigan State University, East Lansing, MI, USA

[7]Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI, USA

[8]Department of Chemistry, Michigan State University, East Lansing, MI, USA

Abstract: *Bacteroides fragilis* is an anaerobic resident of the human gut known to tolerate the toxic effects of host-produced and microbially-modified bile acids. Two conserved genes, *clsA* and *clsB*, encode putative cardiolipin synthases that have been linked to bile acid tolerance, but their physiological roles remain undefined. Phylogenetic analysis indicates that *Bacteroides* spp. ClsA

and ClsB diverge from the well-characterized cardiolipin synthases of Gammaproteobacteria and

Firmicutes. Here, we show that these enzymes have distinct cardiolipin synthase activities and

make non-redundant contributions to *B. fragilis* fitness under gut-relevant stress conditions,

including osmotic stress, disruption of membrane potential, and exposure to the bile acid

deoxycholate. Although deoxycholate treatment perturbed K^+/Na^+ homeostasis in *B. fragilis*,

deletion of *clsA* or *clsB* did not significantly alter intracellular ion levels, suggesting that cardiolipin

loss does not substantially impact ion balance under standard cultivation conditions. High-

resolution lipidomic analyses showed that cardiolipin comprises less than 1% of *B. fragilis*

membranes and that ClsA and ClsB produce distinct cardiolipin products with unique acyl chain

lengths and levels of unsaturation. Deletion of either *cls* gene led to Cls-specific remodeling of *B.*

fragilis envelope lipid content, which was also associated with shifts in non-lipid metabolites

indicative of stress-induced metabolic changes. These results define distinct roles for ClsA and

ClsB in shaping *B. fragilis* membrane composition, metabolism, and stress resilience, and

highlight cardiolipin as a key determinant of fitness under bile acid stress.

Importance: Inflammatory bowel diseases are a growing global health concern, motivating efforts

to understand how inflammation-associated microbes, such as *B. fragilis*, adapt to and persist in

stress conditions typical of the intestinal environment. Our targeted genetic analyses reveal that

the cardiolipin synthases ClsA and ClsB make non-redundant contributions to *B. fragilis* fitness

under bile acid exposure and other gut-relevant stress conditions. Complementary omics-based

approaches show that these enzymes produce chemically distinct cardiolipin species and that

their loss leads to specific remodeling of membrane lipid composition and cellular metabolite

profiles. Together, our results indicate that cardiolipin synthases contribute to membrane and

metabolic adaptation in *B. fragilis*, supporting stress resilience in the presence of bile acids and

other gut-relevant stressors.

INTRODUCTION

To successfully occupy gut niches and maintain homeostasis, bacteria must preserve membrane integrity to resist phage attack, host immune pressures, and membrane-disrupting molecules such as bile acids (1-7). The functional properties of membranes are shaped by the diverse lipids classes of which they are made, each characterized by an array of structural species (8-10). For example, bacteria can alter their membrane fluidity and/or permeability by modulating the length of phospholipid acyl tails in response to extracellular conditions (8, 11). Bacteria also adjust the abundance of different phospholipid types or modify lipid headgroups through reactions such as aminoacylation (12-16).

In addition to protecting against external threats, membranes also help support cellular metabolism by maintaining a potential energy source in the form of ion gradients, such as hydrogen (protons, H^+), sodium (Na^+), and potassium (K^+) (17). One of the best-known functions of proton gradients is to power rotary ATPase complexes (i.e., F_oF_1) associated with aerobic and anaerobic respiration (18-22). Prokaryotes as well as eukaryotes maintain Na^+ and K^+ at intracellular concentrations that are lower and higher, respectively, relative to their environment, which drives the transport of metals or other molecules across the membrane (23, 24).

Cardiolipins, which make up 1–5% of the bacterial membrane, play an integral role in maintaining ion gradients (25-30). This class of membrane lipid comprises two covalently bonded phospholipids, such as phosphatidylglycerol (PG) or in some cases phosphatidylethanolamine (PE) (31). While cardiolipins increase proton permeability across model liposome membranes (28, 32), they also contribute to membrane stabilization by enhancing hydrogen bonding, reducing water permeability and sustaining membrane potential during stress-induced membrane depolarization (32, 33). In addition, several protein complexes require cardiolipin for proper function, including some two-component systems, protein translocation machinery, and respiratory complexes (34-39).

Escherichia coli has three well-studied genes responsible for cardiolipin biosynthesis: *clsA*, *clsB* and *clsC* (31, 40). Each gene makes distinct contributions to *E. coli* physiology depending on growth phase, with *clsA* being the primary *cls* used during logarithmic (log)-phase growth and all three used during stationary phase (26, 31). While ClsA and ClsB act on PG as the sole substrate, ClsC synthesizes cardiolipin from PE as well as PG (31, 41-43). The Cls enzymes encoded by these genes are thought to synthesize three structural families of cardiolipin, which have four (cardiolipin), three (monolysocardiolipin) and two acyl tails (dilysocardiolipin), respectively (for chemical structures, see Fig. 6A) (44). However, mechanistic insight into bacterial cardiolipin biosynthesis and function largely comes from studies in *E. coli* and other Enterobacteriaceae. Expanding this research across diverse bacterial lineages is essential for uncovering the broader physiological roles of this important lipid class.

The gut microbe *Bacteroides fragilis* is a gram-negative facultative anaerobe that must maintain membrane integrity for gut fitness (45-47). While *B. fragilis* strains, particularly those that carry the fragilysin toxin, are linked to colorectal cancer and other diseases (48), it constitutes about 1–5% of the microbiota in otherwise healthy humans (49, 50). We recently identified two cardiolipin synthases as critical for the survival of the non-toxigenic *B. fragilis* strain P207 in the presence of physiologically relevant levels of deoxycholate, a secondary bile acid (51). *B. fragilis* P207 was isolated from the ileoanal pouch of a patient with ulcerative colitis (UC) and is representative of *B. fragilis* populations that can dominate prior to the onset of pouchitis (52), a condition characterized by UC-like inflammation of the ileoanal pouch following colectomy (53). In this study, we build on our previous findings by investigating the roles of the two *B. fragilis* cardiolipin synthases across a broader range of membrane stress conditions. Since cardiolipin synthases are largely uncharacterized in the Bacteroidota, our work provides foundational knowledge about this important class of enzymes in a dominant group of gut microbes (54). Using a combination of approaches, we demonstrate that the two *cls* genes in *B. fragilis* P207, *clsA* and *clsB*, are evolutionarily and functionally distinct from *cls* homologs in *E. coli* and other well-studied

bacterial phyla. These genes encode non-redundant enzymes that contribute structurally unique

cardiolipin species to the membrane, conferring distinct physiological advantages under stress.

We further show that *clsA* and *clsB* differ in their expression profiles and that mutants lacking

each gene produce distinct cardiolipin species as well as distinct metabolomic profiles, supporting

complementary roles in maintaining membrane homeostasis and adapting to diverse

environmental conditions.

RESULTS

B. fragilis *P207 ClsA and ClsB are cardiolipin synthases*

The two *B. fragilis* P207 genes, ptos_000612 (*clsA*) and ptos_003252 (*clsB*), encode

proteins with predicted cardiolipin synthase domains (TIGR04265, PFAM13091, cd09112),

though *clsA* and *clsB* have low nucleotide (52.4%) and amino acid sequence (29.6%) identity to

each other (Fig. 1A). The low primary structure identity (≤30%) of ClsA and ClsB to *E. coli* Cls

proteins limited our ability to predict their specific enzymatic activities based on sequence alone

(Fig. 1A). Furthermore, *E. coli* ClsC did not align well with ClsA or ClsB from either *B. fragilis* or

E. coli (Fig. 1A, B). We therefore excluded ClsC from further comparisons.

ClsA and ClsB from *B. fragilis* and *E. coli* exhibit high amino acid conservation in two

regions around their predicted active sites but share limited identity across the remainder of the

protein (Fig. 1B, C). The Cls proteins from both species harbor two phospholipase-D active site

motifs H-X-K-X(4)-D (26, 42), indicative of phospholipase-D activity (Fig. 1D). Indeed, deleting

either *clsA* (Δ*clsA*) or *clsB* (Δ*clsB*) in *B. fragilis* resulted in a partial loss of cellular cardiolipin

accumulation relative to the wild-type (WT) strain; deleting both genes (Δ*clsA* Δ*clsB*) completely

abolished cardiolipin accumulation (Fig. 1E). We conclude that *clsA* and *clsB* in *B. fragilis* P207

encode *bona fide* cardiolipin synthases. A more comprehensive analysis of how deletion of *clsA*

and *clsB* affects the membrane lipid composition of the *B. fragilis* envelope is presented in

sections below.

The cardiolipin synthases of Bacteroides *sp. occupy a distinct phylogenetic cluster among*

dominant gut bacterial phyla

To evaluate the relatedness between *Bacteroides* Cls proteins and those encoded by other

gut microbes, we performed pairwise alignments of Cls sequences from dominant bacterial phyla

present in the human gut. These sequences clustered according to their evolutionary heritage,

except for the Bacillota, which had two clusters (Fig. 1F). Based on this result, we conclude that

each cardiolipin synthase originated from a proto-lineage, rather than having been horizontally

transferred.

We next evaluated Cls protein sequences within the *Bacteroides* genus. We aligned 301

Bacteroides protein sequences containing the TIGR04265 domain (i.e. cardiolipin synthases).

Most genomes encoded two cardiolipin synthases that partitioned into distinct clusters (Fig. S1).

Notably, the *B. thetaiotaomicron* genome encodes a third Cls that is distinct from the other two

Bacteroides Cls clusters (Fig S1) indicating an evolutionarily distant introduction. We designate

cluster 1 homologs as ClsA based on our observation that both *B. fragilis* and *E. coli* ClsA proteins

are predicted to have two N-terminal transmembrane helices, suggesting a similar mode of post-

translational processing (55, 56). We named the cluster 2 homologues ClsB, as they represent

proteins from the second and only other annotated *cls* gene in most *Bacteroides*. We left the third

cls gene, which we identified only in *B. thetaiotaomicron*, unnamed and did not further

characterize it.

Within the *Bacteroides* genus, the chromosomal gene neighborhoods around *clsA* (Fig.

S2A) and *clsB* (Fig. S2B) are highly conserved; this conservation is lost outside the *Bacteroides*.

Thus, the genomic context of *clsA* and *clsB* provides some indication of the identity of each *cls*

gene, specifically within the *Bacteroides* lineage. This evidence also supports the hypothesis that

taxa inherited each *cls* gene in two distinct regions of the chromosome. Considering this

phylogeny, our analysis of the *B. fragilis cls* genes presented below offer insight into the function

of the two broadly distributed cardiolipin synthases across the genus *Bacteroides*.

clsA *and* clsB *deletion mutants have growth defects under membrane stress*

To explore the functions of the two cardiolipin synthases in *B. fragilis* P207, we generated

mutant strains harboring in-frame deletions of *clsA* (Δ*clsA*) and *clsB* (Δ*clsB*), and a strain lacking

both *cls* genes (Δ*clsA* Δ*clsB*). We also generated corresponding genetic complementation strains

bearing *clsA* or *clsB* integrated at a neutral heterologous site in the chromosome. We assessed

the growth characteristics of these *B. fragilis* strains under stress conditions that perturb various

aspects of membrane biology (Fig. 2). The treatments in this study included: 1) the detergents

deoxycholate and sodium dodecyl sulfate (SDS); 2) low pH; 3) the salts NaCl and KCl; and 4) the

ionophores monensin A (for Na^+) and nigericin (for K^+ and H^+) to disrupt transmembrane ion

gradients. Treatment concentrations were chosen based on their effect on WT strain growth.

In the absence of treatment, all strains grew equally well (Figs. 2A, S3A, B). Treatment

with the detergents deoxycholate (at 0.01% [w/v]; Fig. 2A, S3) or SDS (at 0.002% [w/v]) (Fig. 2B)

limited the growth of the *cls* mutant strains to a greater extent than WT. Deoxycholate affected

maximum OD_{600} in the Δ*clsA* Δ*clsB* att::*clsA* strain, and both max OD_{600} and growth rate the Δ*clsA*

Δ*clsB* strain. SDS treatment mainly affected lag time, decreasing the max OD_{600} values in the first

20 h of growth. Doubling time defects appeared as little or no growth, rather than slower growth

(Fig. 2B). Under both detergent conditions, the deletion of *clsB* had a larger effect on fitness than

deletion of *clsA* (Figs. 2A, B, S3) and deletion of both genes (Δ*clsA* Δ*clsB*) had an additive effect.

300 mM K^+ or Na^+ (Fig. 2C, D) also lowered the growth rate and maximum OD_{600} of the mutant

strains relative to WT (Fig. S3). Both Δ*clsA* and Δ*clsB* strains exhibited similar phenotypes, with

the Δ*clsA* Δ*clsB* strain showing more pronounced effects. Lowering the medium pH from 6.9 to

pH 5.5 (i.e., higher [H^+]) had a detrimental effect on all *B. fragilis* strains, with no clear *cls-*

dependent growth defects (Fig. 2E).

The ionophores nigericin and monensin A dissipate ion gradients by mediating the

exchange of ions across the inner membrane (57, 58). Specifically, nigericin is a potassium-

selective ionophore and a H^+-K^+ antiporter, erasing both gradients by acidifying the cytoplasm and

moving K^+ out of the cell (57). Monensin A is a sodium-selective ionophore with a 16-fold

preference for Na^+ over K^+, but can also affect H^+ gradients (57-59). Under either nigericin or

monensin A treatment, *clsB* deletion decreased strain fitness relative to the WT (Fig. 2F, G).

However, strains harboring *clsB* but lacking *clsA* (Δ*clsA* and Δ*clsA* Δ*clsB* att::*clsB*) exhibited a

higher maximum OD_{600} relative to the WT (Fig. S3).

We conclude that deleting *clsB* results in greater fitness defects than loss of *clsA* under

membrane perturbations. Furthermore, a functional ClsA limits growth under nigericin and

monensin A treatment, as we observed increased culture density in the Δ*clsA* strains. These

growth phenotypes provide evidence that ClsA and ClsB have distinct roles in *B. fragilis*

membrane physiology.

Loss of cardiolipin changes cell shape

Given that in many bacteria, cardiolipin influences cell division, morphology, and

membrane fluidity (60-67), we asked whether loss of Cls function in *B. fragilis* would affect cell

shape. Deletion of either *cls* gene did not markedly affect overall rod-like cell shape or cellular

length distribution (Fig. S4A, B). However, cells lacking *clsA* (i.e., Δ*clsA* and Δ*clsA* Δ*clsB* strains;

Figs. 3A, B and S4C) were narrower. While cells lacking *clsB* were not significantly narrower than

WT, this cell population did have a broader cell width distribution compared to WT, suggesting a

role for ClsB in cell size regulation (F-test, WT to Δ*clsB* p = 0.019, WT to Δ*clsA* Δ*clsB* att:*clsA* p <

0.001, WT variances compared to Δ*clsA* and Δ*clsA* Δ*clsB* were not significantly different; Figs. 3A

and S4C). Both phenotypes could be genetically complemented. These results demonstrate that

ClsA and ClsB influence cell width in *B. fragilis* (Fig. 3A, B).

B. fragilis *cardiolipin synthase genes have distinct expression profiles*

Considering the unique phenotypes of the Δ*clsA* and Δ*clsB* mutant strains, we

hypothesized that these genes have distinct expression patterns. To test this idea, we quantified

clsA and *clsB* transcripts levels across growth phases and stress conditions. *clsB* transcript levels

were four-fold higher during logarithmic (log) phase growth than in stationary phase, while *clsA*

expression was two-fold higher in stationary phase (Fig. 4A). During logarithmic growth, deletion

of either *cls* gene did not affect the transcript levels of the other (Fig. 4B). However, in stationary

phase, the deletion of *clsA* led to an 8-fold increase in *clsB* transcripts. No change in *clsA*

transcripts were observed upon *clsB* deletion (Fig. 4B). We conclude that the two cardiolipin

synthase genes of *B. fragilis* have distinct, growth-phase dependent expression profiles and that

clsB transcript upregulation in the Δ*clsA* strain may compensate for loss of ClsA activity.

To test if membrane stress influences the expression of the *cls* genes, we subjected cells

to acute stress for 20 minutes and harvested RNA. While SDS treatment did not affect *cls*

transcript levels, we observed a two-fold increase in *clsA* expression and a small decrease in *clsB*

transcripts after 0.01% deoxycholate and 300 mM K$^+$ treatments (Fig. 4C) (51). Acidic pH did not

alter *cls* transcript levels, but basic pH did decrease transcript levels of both genes (Fig. 4C).

Finally, we detected differences under ionophore treatment, although the biological significance

of these smaller changes is unclear (Fig. 4C).

Loss of cardiolipin synthase genes does not have a major impact on homeostasis of major

cellular elements

As cardiolipins play an important role in maintaining membrane barrier integrity (25-27,

68, 69), we used inductively coupled plasma mass spectrometry (ICP-MS) to quantitatively define

amounts of certain elements in *B. fragilis* cells and measure the effect of ClsA and/or ClsB or

stress treatment on those elements. While cell washing is routinely used to prepare samples for

metal analysis, it can disrupt intracellular levels of elements (Fig. S5A). Therefore, we used a

wash-free ICP-MS assay to minimize perturbation during sample preparation, adapted here for

bacterial cultures (70). Using this approach, we found that *B. fragilis* P207 cells grown in

supplemented brain heart infusion (BHIS) medium contained 550 mM phosphorus (P), 430 mM

potassium (K), 98 mM sulfur and 63 mM sodium (Na) (Fig. S5C, medium elemental concentrations

shown in S5B).

To compare strains, we normalized intracellular element levels to phosphorus content, an

accepted practice in this type of analysis (71). This revealed no significant differences across the

ΔclsA, ΔclsB, or double-deletion strains in the relative levels of elements detected (Fig. 5A, B,

and S5C). Notably, given the lack of strain-dependent defects in CFU viability, we observed shifts

in relative phosphorus content per CFU across *cls* mutants that was consistent with differences

in the OD_{600}/CFU ratios (Fig. S5E-G). Furthermore, sulfur levels scaled proportionally with

phosphorus across samples (Fig. S5D and S5H). While P and S abundance can vary across

physiological states of the cell, they are tightly regulated (72, 73). Together, the elemental

composition and OD_{600} data suggest that *cls* deletion can alter the ability of *cls* deletion strains to

consistently form colonies.

Like most organisms, *B. fragilis* maintains a higher concentration of intracellular K^+ (+400

ΔmM) and lower concentration of intracellular Na^+ (-67 ΔmM) than the surrounding environment

(Fig. 5C, D). Exposure to deoxycholate, a microbially-modified bile acid, led to decreased

intracellular K^+ and increased Na^+ contents, providing evidence that this treatment indeed disrupts

transmembrane ion gradients in *B. fragilis* (Figs. 5C, D & S5I-J). The observed increase in

intracellular P per CFU (Fig. S5K) suggests that some of these changes could result from the

presence of non-viable cells (Fig. S5L), as these cells would retain the S and P mass associated

with large molecules like proteins and nucleotides, while smaller, more permeable elements would

be lost across damaged membranes (74, 75). Treatment with the sodium selective ionophore,

monensin A, did not affect K^+ levels but caused a 32 mM increase in intracellular Na^+ (Fig. 5C,

D). By contrast, treatment with the potassium selective ionophore, nigericin, caused no detectable

effect on either K^+ or Na^+ content, suggesting that nigericin may require more than 20 min to act

on *B. fragilis*, or that the cells quickly adapt to the imposed ionic disturbance (Fig. 5C, D).

Membrane lipid composition of WT* B. fragilis *P207

Through tandem mass spectrometric-based lipidomics (MS/MS), we characterized lipid

species present in *B. fragilis* P207 membranes using both negative and positive electrospray

ionization modes. To get an overall picture of WT cell membranes, we calculated the relative peak

area abundance for lipids detected in negative ionization mode. This analysis revealed that WT

cell membranes were predominantly composed of phospholipids (47% of total peak areas) and

ceramides (42%). The remaining abundance consisted of saccharolipids, diacylglycerols and *N*-

acyl lipids (Figs. 6A & S6A). The major phospholipids were species of PE (37% of total peak

areas), PS (7.2%) and PG (1.5%) (Fig. 6A, B, Table S3). Cardiolipin species accounted for

approximately 0.33% of membrane lipids detected (Fig. 6B, C). Dominant ceramides were

ceramide phosphoethanolamine (31.7% of total peak area) and ceramide beta-hydroxy fatty acid-

dihydrosphingosine (8.6%). These results are consistent with previous studies on *Bacteroides*

membrane lipid contents (76-78).

Since cardiolipin synthases use phospholipids as substrates, we investigated the types of

PA, PE, PS and PG species. For PA, PS and PE, the dominant lipid types were fully saturated

(30:0), comprising 85.8%, 60.7% and 57.1% of their respective classes (Fig. S6B-C). The most

prevalent PG species was 32:0 (54.5%, Fig. S6B). *B. fragilis* membranes also contained

lysophospholipids at lower levels than their two-acyl tailed versions (Fig. 6A), the most abundant

species of which were fully saturated at C_{15} chains for lysophosphatidylethanolamine (LPE;

76.9%) and lysophosphatidylserine (LPS; 54.9%), and unsaturated at 18:3 for lysophosphatidic

acid (LPA; 70.5%) (Fig. S6D). Ether-linked lipids and cardiolipin lipid pools showed much greater

diversity in their constituent acyl chains (Fig. S6E–J).

clsA *and* clsB *mutants have distinct cardiolipins profiles*

We compared the membrane lipid profiles of the *B. fragilis* P207 WT strain and the derived

Δ*clsA*, Δ*clsB*, and Δ*clsA* Δ*clsB* mutant strains (Fig. 1E). While the overall abundance of the

dominant lipid classes did not vary significantly among the WT and *cls* mutant strains (Fig. 6B),

abundances of cardiolipin and monolysocardiolipin were lower in the *cls* deletion mutants (CL and

MLCL; Fig. 6C). Total cardiolipin levels were lower in the Δ*clsA*, Δ*clsB*, and Δ*clsA* Δ*clsB* strains

compared to WT, with the double knockout strain having no measurable cardiolipin (Fig. 6C). We

detect lower MLCL levels only in strains lacking *clsB*, indicating that ClsB is primarily responsible

for their biosynthesis (Fig. 6C). Notably, the deletion of both *clsA* and *clsB* led to no change in the

detected dilysocardiolipin (DLCL), indicating that in contrast to *E. coli* where Cls proteins

synthesize these lipids, a different enzymatic pathway is responsible for biosynthesis of DLCLs in

B. fragilis (Figs. 6C and S6K) (31, 41).

Principal component analysis (PCA) of membrane lipid peak areas revealed that each

mutant strain clustered separately, indicating that ClsA and ClsB independently shape membrane

lipid composition, while the double knockout reflected the combined loss of both functions (Fig.

6D). K-means clustering of lipid changes in *cls* deletion mutants relative to WT identified four

distinct lipid subpopulations (Fig. 6E). Clusters 1 and 2 were composed almost entirely of

cardiolipin, monolysocardiolipin and hexosylceramides (i.e., cluster 1 = AHexCer, cluster 2 =

AHexCer, Hex2Cer, HexCer_HDS), and were absent in the Δ*clsA* Δ*clsB* double deletion strain.

On average, the peak area of lipids in cluster 3 was significantly lower in strains lacking ClsA,

while cluster 4 lipids were significantly higher in strains lacking ClsB (Fig. 7A). Lipids from cluster

1 were largely missing is strains without *clsB*, and lipids from cluster 2 were largely missing in

strains without *clsA*. We therefore refer to these as *clsB*- and *clsA*-associated lipid clusters,

respectively. From these results, we conclude that ClsA and ClsB catalyze the production of

distinct lipid species in *B. fragilis*, consistent with the unique phenotypes observed in their

respective deletion mutants (Fig. 2).

ClsA and ClsB differentially influence cardiolipin saturation and membrane composition

To better understand the roles of ClsA and ClsB in cardiolipin biosynthesis, we analyzed

the acyl chain length and degree of unsaturation in CL and MLCL species from lipid clusters 1

and 2. The cardiolipins associated with ClsA and with ClsB exhibited a different distribution of

chain lengths, including a subset of longer species that were only detected when ClsB was

present (Figs. 7B and S7A). The major differences between ClsA- and ClsB-associated lipids

were in acyl chain unsaturation: ClsA-associated cardiolipins were enriched in species with 3 and

11 double bonds, whereas ClsB-associated cardiolipins were enriched with 1, 2, and 14 double

bonds (Fig. 7C & S7B). These unique lipid profiles provide evidence that ClsA and ClsB have

distinct substrate preferences.

To determine whether *cls* deletions alter the membrane lipidome globally, we performed

PCA on the lipid profiles of each sample excluding cardiolipins. This analysis showed that each

strain still clustered separately with an additive phenotype, suggesting broad ranging, *cls*-specific

changes (Fig. S7C). Upon loss of cardiolipin biosynthesis in *B. fragilis*, we observe altered

membrane lipid composition, including shortening the acyl chain length of

phosphatidylethanolamine (PE), phosphatidylglycerol (PG) and phosphatidylserine (PS) species

(Figs. 7D and S7D). This shift was coupled with increases in lipid acyl chains with a high number

of total unsaturations (Figs. 7E and S7E). The combined effect of shorter acyl chains and

increased unsaturation is predicted to alter membrane fluidity and permeability, potentially

compensating for cardiolipin loss. Additional remodeling occurred in non-phospholipid lipid

classes, with lipid families such as fatty acids (FAHFA) and hexosylceramides (HexCer-HDS)

more affected by *clsA* deletion, while *N*-acyl lipids (NAOrn, NAE, NAGlySer), phosphoceramides

(CerP), other hexosylceramides (AHexCer, Hex3Cer), and ether-linked glycolipids (EtherMGDG)

were more affected by *clsB* deletion (Fig. 7F–H & S7F-G). These patterns were confirmed by

elastic net regression, which distinguished the double mutant from WT based reductions in Cluster

1 (CL 75:3, AHexCer [O-22:6]51:9:2O) and Cluster 3 (PE O-38:0:2O) lipid species, as well as

increases in two Cluster 4 lipids (NAGlySer 29:0:1O, PC O-24:4:1O). Together, these results

provide evidence that ClsA and ClsB have distinct biosynthetic activities that contribute to

cardiolipin diversity, and that the loss of either enzyme perturbs the membrane lipidome in unique

ways. These broader shifts in membrane composition upon *cls* gene deletion likely underlie the

distinct phenotypes observed in each mutant strain.

ClsA and ClsB loss lead to distinct changes in non-lipid metabolites

Since we observed distinct shifts in non-cardiolipin lipids, we hypothesized that these

shifts might extend to other *B. fragilis* metabolites under non-stressed conditions. Using LC-

tandem mass spectrometry (LC-MS/MS), we found that each strain had unique metabolite

profiles, with Δ*clsB* being the most distinct (Figs. 8A-C). Using publicly available databases, 56%

of all features were annotated, with 83% of those matches from small molecule databases.

Approximately 27% of the metabolomic features were not shared between all the strains, with

~14% of the features only depleted in Δ*clsA*, Δ*clsB*, or Δ*clsA* Δ*clsB* (Fig. 8B). Notably, we observed

that the effect of *clsA* and *clsB* deletion is not additive, with *clsB* deletion showing the most

profound alterations on the metabolomic profiles by hierarchical clustering (Fig. 8C).

To determine metabolites that are associated with *cls* deletions, we performed pathway

enrichment analysis comparing feature abundances between the WT and double knockout strain

(Tables S6, S7). As with the lipidomics analysis above, we identified fatty amides as the most

differentially enriched pathway, followed by the fatty esters and sphingolipids (Fig. S8). Several

non-lipid classes were also identified, including alkaloids, amino acids, amino sugars, benzoic

acid esters, and pyrimidines as the most differentially enriched pathways with high enrichment

factors and significant p-values (Fig. 8D). Analysis of amino acids and related compounds

identified lipid-conjugated homoserine lactones (HSLs) and fatty amides to be substantially

changed in the mutant strains (Figs 8E–F). The lipid-conjugated HSLs, which are small signaling

molecules crucial in quorum sensing (79), were mainly associated with *clsB* deletion. In these

mutants, levels of long-chain HSLs (e.g., *N*-hexadecanoyl HSL) were reduced, while shorter lipid

conjugates (e.g. *N*-tetradecanoyl- and *N*-pentadecanoyl HSL) were enriched. Changes in fatty

amides were associated with both *clsA* and *clsB* deletions, with many metabolites showing

opposing trends between mutant strains. Levels of several *N*-acyl amides were affected by both

clsA and *clsB* deletion, including *N*-hexadecyl-L-hydroxyproline, *N*-(3-hydroxy-14-

methyldecanoyl) glycine, *N*-palmitoyl threonine, and *N*-stearoyl tryptophan. Some *N*-acyl amides

showed gene specific changes such as *N*-(3-hydroxy-1-oxodecanoyl) glycine, which was primarily

affected by *clsA* deletion, and *N*-palmitoyl GABA, which was impacted by *clsB* deletion (Fig. 8F).

N-acyl amides may serve as additional carbon sources and/or signaling molecules between

bacteria or to the host, such as through G-protein coupled receptors (GPCRs) (80, 81). Molecular

network analysis identified *N*-acyl amides in larger clusters of chemically similar but unannotated

metabolites, suggesting that *cls* modifications may lead to larger global metabolomic shifts (Fig.

S9A-B). These findings demonstrate that cardiolipin synthases play essential non-redundant

roles in maintaining metabolic homeostasis in *B. fragilis* P207.

DISCUSSION

In this study, we demonstrate that *B. fragilis* P207 has two functionally non-redundant

cardiolipin synthases, which we named ClsA and ClsB. Deleting either gene leads to distinct

defects in *B. fragilis* fitness under stress, with the loss of *clsB* generally being more detrimental

than *clsA* (Fig. 9). We note that *clsA* deletion results in elevated *clsB* expression, which may

partially compensate for and minimize the impact of its loss. We present evidence that ClsA and

ClsB synthesize cardiolipin species that differ in acyl chain length and unsaturation levels,

suggesting that each enzyme has unique substrate specificity and contributes distinct lipids to the

membrane. Furthermore, loss of either enzyme has knock-off effects on diverse cellular metabolic

processes.

This study is significant for several reasons. First, we offer an approach that integrates

metallomics-, lipidomics-, metabolomics-, and classical genetics-based techniques to investigate

cardiolipin synthase function. We employ tandem mass spectrometry to assess how *cls* deletion

affects the lipid composition of the *B. fragilis* envelope. To our knowledge, previous studies have

not leveraged high-resolution mass detection to differentiate the specific lipid species produced

by individual Cls proteins (26, 31, 41, 42, 82). While some have used MS/MS to characterize

membranes of diverse taxa, including *Bacteroides*, these approaches have not been applied to

strains with deletion of individual *cls* genes, limiting the identification of products associated with

each enzyme (76, 77, 83-86). By combining genetic and analytical approaches, we distinguish

the unique contributions of each Cls to the cardiolipin pool and intracellular ion gradient

homeostasis. Second, we provide functional characterization of *cls* genes in the *Bacteroides*

taxonomic group, particularly genes from a novel clinical isolate from the IBD gut. As important

members of the human gut ecosystem, *Bacteroides* species play key roles in microbiota

metabolism and host immune health (46). Understanding how these microbes regulate membrane

composition and homeostasis is essential for understanding their fitness in the context of the gut.

Future research should continue to explore cardiolipin biosynthesis in these organisms,

particularly in *B. thetaiotaomicron*, which possesses an additional *cls* not found in most other

Bacteroides species.

Due to the high diversity of bacterial *cls* genes, we struggled with the best approach to

name the *Bacteroides* group *cls* homologs (34, 82, 87, 88). While new letter codes (e.g. *clsX* or

clsN) might help distinguish *cls* gene diversity across bacteria, this naming scheme might have

become onerous upon characterization of other homologs. We therefore decided on the naming

conventions used for *E. coli cls* and other taxa, acknowledging that *cls* homologs across different

bacterial phyla may have distinct functional roles (89, 90).

The *B. fragilis cls* genes have some similarity to *E. coli clsA* and *clsB* as the genes from

both taxa have strong alignment with cardiolipin synthase domain models. However, at the amino

acid sequence level, no clear phylogenetic relationship exists between these homologs. In each

taxon, the ClsA homolog is predicted to have two N-terminal transmembrane helices. While these

domains in ClsA are thought to be cleaved post-translationally in *E. coli*, it is likely that this does

not occur in *Bacillus subtilis* (26, 90). Therefore, the functional consequences of these putative

transmembrane domains in *B. fragilis* remain unknown, particularly their role in the cellular

localization of these enzymes (90, 91). In addition, across these two taxa, the *cls* expression

profiles are distinct. *E. coli* relies almost exclusively on *clsA* during log phase, and expression of

clsB and *clsC* is largely limited to the stationary phase (26, 31, 41, 42). Under our growth

conditions, *B. fragilis* expresses both *cls* genes during log phase, with *clsA* expression further

enhanced during stationary phase. To determine if the two *B. fragilis* Cls have overlapping

enzymatic activities with the Cls of these other taxa, comparative biochemical analysis of the lipid

products produced by the Cls homologs of each species is needed.

Maintaining ion gradients across membranes is essential for bacterial survival, as these

gradients provide a reservoir of potential energy. The loss of cardiolipin biosynthesis under non-

stressed conditions largely does not affect ion homeostasis, as seen in our *B. fragilis* metallomic

studies. Bacteria generally tightly regulate intracellular K^+ content due to its physiological

importance, with some adjusting it by several fold depending on extracellular conditions (24, 92).

For example, *B. subtilis* and *E. coli* both increase intracellular K^+ levels in response to elevated

osmolarity (93, 94). However, *S. aureus* does not modulate K^+ content in this manner (95). While

one might expect ion gradient perturbations given previous work showing improved membrane

stability with the presence of cardiolipin (88, 96, 97), the resilience of intracellular K^+ and Na^+

levels in the absence of cardiolipin suggests that otherwise unperturbed cells maintain membrane

integrity. Even if loss of cardiolipin changed the inherent membrane permeability of either K^+ or

Na^+, our data indicate that *B. fragilis* leverages membrane remodeling and several compensatory

ion shuttling systems to achieve and maintain intracellular homeostasis (Fig. 5C-D) (24, 92, 98-

103).

While *B. fragilis clsA* is not essential and may even diminish fitness during selective ion

permeabilization, both *cls* genes are beneficial when the bacterium faces broader membrane

stress conditions. Although unlikely, the absence of ClsA-associated cardiolipins in the Δ*clsA*

strain may provide stress resistance by impairing ionophore translocation across the membrane

(59, 104, 105). Our favored hypothesis is that Δ*clsA*-dependent membrane remodeling enhances

resilience to ion gradient loss, possibly through altered membrane protein or lipid function. Beyond

general effects like membrane fluidity, cardiolipins can influence bacterial physiology through

multiple pathways. For example, some proteins that influence cell shape and cell division, such

as the actin homolog MreB, rely on cardiolipin for proper folding at the membrane (106, 107).

Interactions of cardiolipin with membrane-bound or membrane-associated cell division/shape

proteins may also underlie our observation that the loss of cardiolipin biosynthesis results in *B.*

fragilis cell shape defects (60-66). The cls-specific cell size effects may be shaped by

asymmetrical distribution of cardiolipin species across inner and outer leaflets of the inner

membrane as well as inner leaflet of the outer membrane (16, 108-110).

The catalytic products of *B. fragilis* ClsA and ClsB, as inferred from our lipidomics data,

have distinct acyl chain lengths and unsaturation levels. In response to environmental changes,

bacteria can adjust these lipid properties, which are key factors influencing membrane fluidity and

permeability (111). For example, under conditions that increase membrane rigidity, such as lower

temperature or high pressure, microbes increase the abundance of highly unsaturated fatty acids

to restore membrane fluidity (112, 113). Exposure to membrane destabilizing agents, such as

alcohols, can result in decreased acyl chain unsaturation (114). Exposure to phenols, which can

also disrupt membranes, cause some anaerobes to increase their membrane fluidity by

incorporating fatty acids with trans-double bonds (115). Although *B. fragilis* apparently lacks

enzymes required for this particular mechanism of introducing double bonds, phylogenetically

diverse bacteria have other regulatory mechanisms for the introduction of double bonds into acyl

chains already present in the membrane (116, 117). Having two unique pools of cardiolipins with

corresponding membrane lipids that respond to cardiolipin fluctuations, such as hexosylceramide

for ClsA and *N*-acyl glycylserine for ClsB, would provide bacteria like *B. fragilis* with greater

adaptability in the face of diverse selective pressures in the gut.

The distinct metabolomic alterations observed in *B. fragilis* ClsA and ClsB mutants

underscore the non-redundant and critical roles of cardiolipin synthases in bacterial metabolic

homeostasis. Our results demonstrate that loss of either ClsA or ClsB leads to unique changes in

the abundance of key lipid and non-lipid metabolites. These changes likely reflect the broader

disruptions in membrane composition and cellular signaling, as well as shifts in central metabolic

pathways. Importantly, the non-additive nature of the double mutant phenotype suggests that ClsA

and ClsB may influence partially overlapping yet distinct biosynthetic networks. The pronounced

depletion and enrichment of specific metabolites, such as lipid-conjugated HSLs and *N*-acyl

amides, further indicates that cardiolipin synthase activity influences both membrane structure

and intercellular communication. Given the essential role of membrane integrity and metabolic

adaptability in bacterial survival within the gut environment, these findings suggest that ClsA and

ClsB are pivotal for the ecological fitness of *B. fragilis*.

In summary, *B. fragilis* relies on cardiolipin synthases to maintain barrier integrity against

diverse stresses that it encounters in the gut, such as bile acids and osmotic variation. However,

much remains to be discovered regarding *B. fragilis* Cls-mediated physiology, including protein

domain function, cellular localization, and the mechanism by which each enzyme produces

distinct cardiolipins. Furthermore, the overlapping, yet possibly distinct, functional roles that this

diverse enzyme class plays across the prokaryotic tree of life is ripe with potential. The data we

present here serve as a foundation for a deeper understanding of *B. fragilis* physiology specifically

as well as that of the Bacteroidota generally, potentially leading to the engineering of probiotic gut

species with improved gut fitness and therapeutic outcomes.

METHODS (see supplemental methods for additional detail)

Bacterial Growth and Strains

Strain generation

E. coli strains were grown aerobically at 37˚ C in Miller LB broth. As appropriate, E. coli

cultures were supplemented with filter-sterilized 300 µM diaminopimelic acid (Sigma-Aldrich) to

grow auxotrophic strains and/or 100 µg/ml carbenicillin to select for plasmid containing cells. All

B. fragilis strains were grown in supplemented brain heart infusion (BHIS) medium. As

appropriate, cultures were supplemented with 5 µg/mL erythromycin for vector selection, 100

ng/mL anhydrotetracycline (aTC) for inducible ssbfe1 counter-selection (118) and 50 ng/mL aTC

to induce expression from aTC-inducible complementation vectors (118). *B. fragilis* strains were

manipulated aerobically and grown at 37˚C in an anaerobic chamber (Coy Laboratory Products)

filled with 5% CO_2, 2.5% hydrogen and 92.5% nitrogen.

Mutant strains were generated by allelic exchange using modified vectors described

previously (118). For conjugation into *B. fragilis* P207, plasmids were transformed into the donor

E. coli strain WM3064. Allele exchange was conducted as previously described (118). The Δ*clsA*

mutant (lacking ptos_000612) was complemented by integration of an aTC-inducible *clsA*

construct. The Δ*clsB* mutant (lacking ptos_003215) was complemented by integration of *clsB*

under the control of a constitutive *Bacteroides* promoter (P*$_{rpoD}$*, bt_1311⁄BT_RS06635).

Complementation constructs were integrated into the chromosome at the *att* site using an intN1

integrase (119). Primers, vectors, and strains are listed in Tables S1.

Cls conservation analysis

Alignment. The *B. fragilis* P207 ClsA and ClsB gene and protein sequences, as well as *E.*

coli K-12 MG1655 (NC_000913) ClsA, ClsB, and ClsC were aligned using default settings of the

Geneious alignment algorithm (Geneious Prime 2024) (51, 52).

DeepTMHMM. The ClsA and ClsB amino acid sequences from *B. fragilis* P207 and *E. coli*

K-12 MG1655 were uploaded to DeepTMHMM (v. 1.0.42, https://dtu.biolib.com/DeepTMHMM) to

predict transmembrane domains (120).

Inter-Phylum Tree. Representative taxa were selected from the top five phyla in the human

gut (metadata listed in Table S2) (121, 122). Annotated *cls* genes were obtained from genomes

on NCBI and protein aligned in Geneious. One genome per species was used, aside from the two

B. fragilis strains. The phylogenetic tree was reconstructed using neighbor-end joining via

Geneious and then visualized using R.

Intra-Bacteroides *Tree.* TIGR04265 domains in *B. fragilis* ClsA and ClsB were identified

using the conserved domain database of NCBI (123). *Bacteroides* protein sequences containing

this domain were obtained by querying the Interpro database. Alignment, tree reconstruction, and

tree visualization were performed as with the inter-phylum tree, except that *E. coli* K-12 MG1655

ClsA was specified as an outgroup.

Genomic Neighborhoods. WebFlags2 was used to query genomic neighborhoods around

B. fragilis P207 *clsA* and *clsB* genes (124).

Growth curve assays

Stationary phase *B. fragilis* strain cultures were back-diluted to an optical density at 600

nm (OD_{600}) of 0.05 and grown to OD_{600} 0.3 in the presence of 50 ng/mL aTC inducer. Each strain

was then diluted again to 0.025–0.05 for growth curves in a 96-well plate format. OD_{600} was

measured with a Tecan Infinite M Nano plate reader (Tecan, Männedorf, Switzerland) for 24 h.

Three biological replicates, each in technical triplicate, were performed for each growth condition.

To expose cells to stress conditions, BHIS medium was prepared as above but modified.

For pH assays, cells were pelleted and then resuspended in aTC-containing BHIS medium

adjusted to the appropriate pH. For deoxycholate and SDS assays, 0.01% (w/v) deoxycholate

(Sigma) or 0.002% (w/v) SDS (Lab Scientific) was added to cultures from filter-sterilized 1% (w/v)

stocks dissolved in UltraPure water. For Na^+ and K^+ assays, NaCl (Fisher Scientific) and KCl

(Fisher Scientific) were dissolved to a $4\times$ concentration of each cation in BHIS medium and then

filter sterilized before being added to cells to yield the final concentration of cells and cation (i.e.,

300 mM). To determine how much additional NaCl or KCl to add, the average Na^+ and K^+ content

of our BHIS medium was determined using ICP-MS (see below for methods), yielding 130 mM

for Na^+ and 33 mM for K^+ (Table S4). For ionophore assays, a concentrated ionophore (monensin

A or nigericin; Sigma-Aldrich) was added to cultures to yield final concentrations of 0.8 µg/mL

monensin A or 0.1 µg/mL nigericin, with equivalent volumes of ethanol added to non-treated

conditions as a vehicle control. We calculated maximum OD600 and doubling time using R

(ipolygrowth; see supplemental methods) (125).

Microscopy

B. fragilis P207 cell cultures were grown overnight under antibiotic selection and then

back-diluted into medium containing inducer (aTC) without antibiotic selection to reach log-phase

growth. All cultures were removed from the 37ºC incubator but were maintained under anaerobic

conditions, with only two or three cultures removed for microscopy at a time. Cells were visualized

by phase-contrast microscopy on a Leica DMI6000 B inverted microscope with a Hamamatsu

ORCA-R2 10600 camera with a $63\times$ Plan Apo objective (126). Each strain was visualized on four

different days to generate four biological replicates. The two images per replicate from each

strain/day were chosen for analysis. Chosen images were processed using the MicrobeJ plugin

(v. 5.13I) cell size analysis software in Fiji (v. 1.54k) (127). Called cells were manually curated and

resulting cell lengths and widths were exported and analyzed in R to calculate cell volume and

Epanechnikov kernel densities.

Gene expression assays

Stress condition exposure: For comparison of *cls* gene expression at log and stationary

phases, *B. fragilis* WT, $\Delta clsA$ and $\Delta clsB$ strain log phase cultures were back diluted to an OD_{600}

of 0.05 without inducer. Two and 24 h later, cell samples were pelleted and resuspended in 1 mL

of Trizol (Invitrogen). For acute toxicity assays, back-diluted WT *B. fragilis* cultures that reached

OD_{600} ~0.3–0.5 were exposed to a specified condition (i.e., pH, SDS, deoxycholate, K^+, monensin

A, or nigericin, as described for the growth assays) and incubated for 20 min before cell sample

collection.

RNA extraction: RNA was extracted as described previously (51). RNeasy Extraction Kit

(Qiagen) was used to treat samples with Turbo DNase (Invitrogen) using the manufacturers

protocol.

RT-qPCR: Samples were run in technical triplicate. *Ct* values were determined using a

QuantStudio5 (Applied Biosystems) instrument and a Luna One-Step RT-qPCR kit (NEB) for

target genes in each sample (51). Using *B. fragilis dnaN* (ptos_002600, encoding DNA

polymerase subunit beta) as a normalization gene.

Inductively coupled plasma mass spectrometry

Log phase cells were removed from the anaerobic chamber for sample collection. If

exposed to a stress, cells were incubated for 20 min aerobically at room temperature with either

0.01% (w/v) deoxycholate, 0.8 µg/mL monensin A or 0.1 µg/mL nigericin. Cultures were spiked

with gadolinium-DOTA (Gd-DOTA, Macrocyclics) to a final concentration of 40 µM. Washing cells,

even with the BHIS medium cells were grown in, perturbs intracellular concentrations of small

alkali metals like Na^+ and K^+ making quantification difficult (Fig. S5A). Using a Gd-DOTA spike

enabled us to remove washing steps in the ICP-MS sample preparation process (128, 129).

For the wash experiment, wash condition cells were washed twice with either 250 mM

sucrose or BHIS media. Then, unwashed cells (i.e., no previous centrifugation) and resuspended

washed cells were transferred to a metal-free tube, given a Gd-DOTA spike to 40 µM, and then

pelleted. For *cls* and treatment comparison experiments, cells were unwashed.

Weighed cell pellets and supernatant were dried overnight, then digested using 70% HNO_2

acid, before diluting to 3% nitric acid. All standards, blanks, and ICP-MS samples were prepared

using ultra-trace metal grade nitric acid (70%, Fisher), ultrapure water (Millipore), metal free

polypropylene conical tubes (15 and 50 mL, Labcon), and trace metal grade pipette tips (Labcon).

All solutions were weighed for accurate elemental determination using the XSR205 DU semi-

analytical balance (Mettler Toledo).

Samples were analyzed using an Agilent 8900 Triple Quadrupole ICP-MS (Agilent

Technologies). To prepare ICP-MS standards, the multi element standard IV-65024 (Inorganic

Ventures) was diluted with 3% (v/v) nitric acid in ultrapure water. Internal standardization was

accomplished inline using a 200 ppb internal standard solution in 3% (v/v) nitric acid in ultrapure

water (IV-ICPMS-71D, Inorganic Ventures). The isotopes selected for analysis were ^{23}Na, ^{31}P, ^{32}S,

^{39}K and ^{157}Gd with ^{45}Sc, ^{89}Y, ^{115}In, and ^{159}Tb used for internal standardization. For accurate

quantification of ^{31}P and ^{32}S, these isotopes were measured in the oxygen mode. To quantify

intracellular elemental concentration, the average strain-specific cell volume was determined and

then the molar concentration calculated for each element from the number of atoms per cell. The

ΔmM concentrations were calculated by subtracting the concentration of an element in the

medium from that in the intracellular space.

Lipidomics assays

WT, $\Delta clsA$, $\Delta clsB$, and $\Delta clsA$ $\Delta clsB$ strains were backdiluted, grown to OD_{600} 0.7, and then

submitted on dry ice to the University of Tennessee-Knoxville Biological and Small Molecule Mass

Spectrometry Core (BSMMSC; RRID: SCR_021368) for targeted lipidomic analysis of membrane

lipids. Lipids were extracted as previously described (see supplemental methods for details) (130,

131). The lipidomic analyses were performed using a previously validated method (131) on an

ultra high performance liquid chromatography system coupled to a high resolution mass

spectrometer (UHPLC-HRMS). The chromatographic separations were carried out using

Vanquish Horizon LC system (Thermo Scientific) and reversed phase separations as described

previously (131). The eluent was introduced to an Exploris 120 mass spectrometer (Thermo

Scientific) via electrospray ionization for high resolution analysis. Following the analysis, the lipids

were annotated and the peak area integrated using MS-DIAL (132-135). Peak area for a

metabolite was first normalized to the optical density of a sample, then relative lipid abundance

was determined by normalizing to the total peak area of all lipids detected in each

UHPLC−MS/HRMS mode (i.e., positive versus negative). Elastic net regression performed using

the glmnet R package (136, 137).

Metabolomic assays

WT, $\Delta clsA$, $\Delta clsB$, and $\Delta clsA$ $\Delta clsB$ strains were grown in 50mL BHIS medium and

metabolites were extracted with ethyl acetate (see supplemental methods for details). Untargeted

metabolomics data was acquired using a Bruker timsTOF Pro2 (Bruker-Daltonics, Billerica, MA,

USA) coupled to an Agilent 1290 Infinity II Bio UHPLC (Agilent, Santa Clara, CA, USA). The

chromatographic separations were carried out using an Acquity UPLC HSST3 column (2.1 x

150mm, 1.8 μM) (see supplemental methods for details). Following data acquisition, mass

spectrometry (MS) data were pre-processed using Bruker MetaboScape® version 9.0.1 (Bruker-

Daltonics, Billerica, MA, USA) and filtered using mpactR (see supplemental methods for details)

(138). Peak intensity for each metabolite was normalized to the total peak intensity of all features

detected in the sample. Molecular networking was generated as previously described (139) using

GNPS2: Global Natural Products Social Molecular Networking 2 (http://gnps2.org/). A minimum

cosine similarity score of 0.7 with at least six matching peaks, parent mass tolerance of 2 Da, and

fragment ion tolerance 0.5 Da were selected to generate consensus spectra. Files were imported

into CytoScape 3.10.3 and nodes were arranged with yFiles organic layout plugin (140). Pathway

analysis was performed with MetaboAnalyst 6.0 functional analysis [LC-MS] module (Tables S6,

S7) (141). A formatted peak intensity table was uploaded to the web server

(https://metaboanalyst.ca), then normalized by sum and log transformed. Default setting of 5.0

ppm mass tolerance, mummichog algorithm v2.0 with p-value cutoff of 0.0005, against the lipid

and non-lipid pathway libraries with at least 3 entries in each pathway. Metabolites from enriched

pathways were plotted using R (v4.5.0).

Data analysis and reproducibility

All data analyses were performed using R (v4.3.3 unless otherwise specified) and several

packages (125, 136, 137, 142-164). Code used and original datasets for all data analysis and

figure generation, except for the WebFlags neighborhoods, is published on GitHub

(https://github.com/mschnizlein/bfrag_cardiolipin). Lipidomics data alongside appropriate

metadata were deposited on MetaboLights (MTBLS11891

www.ebi.ac.uk/metabolights/MTBLS11891) (165). Untargeted metabolomics were deposited to

MassIVE (https://massive.ucsd.edu) (deposit in progress at time of submission).

Acknowledgements

We would like to thank the Biological and Small Molecule Mass Spectrometry Core (BSMMSC)

at the University of Tennessee-Knoxville and the Q-Beam Center at Michigan State University for

their work processing and analyzing samples for lipid and elemental analyses, respectively. We

thank Drs. Kiwon Ok, Keith MacRenaris and Aaron Sue from the O'Halloran Lab for their advice

on using the ICP-MS an analyzing output. We also would like to thank the entire Crosson Lab for

their advice as the project progressed, particularly Dr. Melene Alakavuklar for her feedback on

this manuscript and figure presentation, respectively. We used Biorender to generate the model

presented in Figure 9. We thank two undergrads who piloted several experiments: Damone

Charleston and Timothy Baldwin. We thank Kathryn McBride for strain culturing in the Balunas

Lab. *E. coli* strain WM3064 was a gift from William Metcalf at UIUC. Work on this project in the

S.C. lab was supported in part by NIDDK (RC2DK122394) and NIGMS (R35GM131762), in the

T.V.O lab by NIGMS (P41GM135018), and in the M.J.B. lab by NCI (U01CA264071).

REFERENCES

1. Schnizlein MK, Young VB. 2022. Capturing the environment of the *Clostridioides difficile* infection cycle. Nature Reviews Gastroenterology & Hepatology 19:508-520.
2. Shortt C, Hasselwander O, Meynier A, Nauta A, Fernández EN, Putz P, Rowland I, Swann J, Türk J, Vermeiren J, Antoine J-M. 2018. Systematic review of the effects of the intestinal microbiota on selected nutrients and non-nutrients. European Journal of Nutrition 57:25-49.
3. Kern L, Abdeen SK, Kolodziejczyk AA, Elinav E. 2021. Commensal inter-bacterial interactions shaping the microbiota. Current Opinion in Microbiology 63:158-171.
4. Runge S, Rosshart SP. 2021. The mammalian metaorganism: A holistic view on how microbes of all kingdoms and niches shape local and systemic immunity. Frontiers in Immunology 12.
5. He X, McLean JS, Guo L, Lux R, Shi W. 2014. The social structure of microbial community involved in colonization resistance. ISME J 8:564-574.
6. Wennerström H, Oliveberg M. 2022. On the osmotic pressure of cells. QRB Discovery 3:e12.
7. Denyer SP, Maillard JY. 2002. Cellular impermeability and uptake of biocides and antibiotics in Gram-negative bacteria. Journal of Applied Microbiology 92:35S-45S.
8. Zhang Y-M, Rock CO. 2008. Membrane lipid homeostasis in bacteria. Nature Reviews Microbiology 6:222-233.
9. Strahl H, Errington J. 2017. Bacterial membranes: Structure, domains, and function. Annual Review of Microbiology 71:519-538.
10. Sohlenkamp C, Geiger O. 2015. Bacterial membrane lipids: Diversity in structures and pathways. FEMS Microbiology Reviews 40:133-159.
11. Blevins MS, James VK, Herrera CM, Purcell AB, Trent MS, Brodbelt JS. 2020. Unsaturation elements and other modifications of phospholipids in bacteria: New insight from ultraviolet photodissociation mass spectrometry. Analytical Chemistry 92:9146-9155.
12. Roy H, Dare K, Ibba M. 2009. Adaptation of the bacterial membrane to changing environments using aminoacylated phospholipids. Molecular Microbiology 71:547-550.
13. Cox E, Michalak A, Pagentine S, Seaton P, Pokorny A. 2014. Lysylated phospholipids stabilize models of bacterial lipid bilayers and protect against antimicrobial peptides. Biochimica et Biophysica Acta (BBA) - Biomembranes 1838:2198-2204.
14. Donohue TJ, Cain BD, Kaplan S. 1982. Alterations in the phospholipid composition of *Rhodopseudomonas sphaeroides* and other bacteria induced by Tris. Journal of Bacteriology 152:595-606.
15. Lin T-Y, Weibel DB. 2016. Organization and function of anionic phospholipids in bacteria. Applied Microbiology and Biotechnology 100:4255-4267.
16. Bogdanov M, Pyrshev K, Yesylevskyy S, Ryabichko S, Boiko V, Ivanchenko P, Kiyamova R, Guan Z, Ramseyer C, Dowhan W. 2020. Phospholipid distribution in the cytoplasmic membrane of Gram-negative bacteria is highly asymmetric, dynamic, and cell shape-dependent. Science Advances 6:eaaz6333.
17. Poolman B. 2023. Physicochemical homeostasis in bacteria. FEMS Microbiology Reviews 47.
18. Zharova TV, Grivennikova VG, Borisov VB. 2023. $F_1 \cdot F_o$ ATP synthase/ATPase: Contemporary view on unidirectional catalysis. International Journal of Molecular Sciences 24:5417.
19. Haddock BA, Jones CW. 1977. Bacterial respiration. Bacteriol Rev 41:47-99.
20. Lu Z, Imlay JA. 2021. When anaerobes encounter oxygen: Mechanisms of oxygen toxicity, tolerance and defence. Nature Reviews Microbiology 19:774-785.

21. Ito T, Gallegos R, Matano LM, Butler NL, Hantman N, Kaili M, Coyne MJ, Comstock LE,
Malamy MH, Barquera B. 2020. Genetic and biochemical analysis of anaerobic
respiration in *Bacteroides fragilis* and its importance *in vivo*. mBio 11:e03238-19.
22. Butler NL, Ito T, Foreman S, Morgan JE, Zagorevsky D, Malamy MH, Comstock LE,
Barquera B. 2023. *Bacteroides fragilis* maintains concurrent capability for anaerobic and
nanaerobic respiration. Journal of Bacteriology 205:e00389-22.
23. Beagle SD, Lockless SW. 2021. Unappreciated roles for K$^+$ channels in bacterial
physiology. Trends in Microbiology 29:942-950.
24. Stautz J, Hellmich Y, Fuss MF, Silberberg JM, Devlin JR, Stockbridge RB, Hänelt I. 2021.
Molecular mechanisms for bacterial potassium homeostasis. Journal of Molecular
Biology 433:166968.
25. Schlame M. 2008. *Thematic Review Series: Glycerolipids.* Cardiolipin synthesis for the
assembly of bacterial and mitochondrial membranes. Journal of Lipid Research 49:1607-
1620.
26. Tropp BE. 1997. Cardiolipin synthase from *Escherichia coli*. Biochimica et Biophysica
Acta (BBA) - Lipids and Lipid Metabolism 1348:192-200.
27. Pinkas D, Fišer R, Kozlík P, Dolejšová T, Hryzáková K, Konopásek I, Mikušová G. 2020.
Bacillus subtilis cardiolipin protects its own membrane against surfactin-induced
permeabilization. Biochimica et Biophysica Acta (BBA) - Biomembranes 1862:183405.
28. Chen Q-P, Li Q-T. 2001. Effect of cardiolipin on proton permeability of phospholipid
liposomes: The role of hydration at the lipid–water interface. Archives of Biochemistry
and Biophysics 389:201-206.
29. Domitin S, Puff N, Pilot-Storck F, Tiret L, Joubert F. 2025. Role of cardiolipin in proton
transmembrane flux and localization. Biophysical Journal 124:408-416.
30. Shibata A, Ikawa K, Shimooka T, Terada H. 1994. Significant stabilization of the
phosphatidylcholine bilayer structure by incorporation of small amounts of cardiolipin.
Biochimica et Biophysica Acta (BBA) - Biomembranes 1192:71-78.
31. Jeucken A, Helms JB, Brouwers JF. 2018. Cardiolipin synthases of *Escherichia coli* have
phospholipid class specific phospholipase D activity dependent on endogenous and
foreign phospholipids. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology
of Lipids 1863:1345-1353.
32. Domitin S, Puff N, Pilot-Storck F, Tiret L, Joubert F. 2025. Role of cardiolipin in proton
transmembrane flux and localization. Biophys J 124:408-416.
33. Shibata A, Ikawa K, Shimooka T, Terada H. 1994. Significant stabilization of the
phosphatidylcholine bilayer structure by incorporation of small amounts of cardiolipin.
Biochim Biophys Acta 1192:71-8.
34. Yeo W-S, Dyzenhaus S, Torres VJ, Brinsmade SR, Bae T. 2023. Regulation of bacterial
two-component systems by cardiolipin. Infection and Immunity 91:e00046-23.
35. Ryabichko S, Ferreira VdM, Vitrac H, Kiyamova R, Dowhan W, Bogdanov M. 2020.
Cardiolipin is required *in vivo* for the stability of bacterial translocon and optimal
membrane protein translocation and insertion. Scientific Reports 10:6296.
36. Musatov A, Sedlák E. 2017. Role of cardiolipin in stability of integral membrane proteins.
Biochimie 142:102-111.
37. Hong Y, Muenzner J, Grimm SK, Pletneva EV. 2012. Origin of the conformational
heterogeneity of cardiolipin-bound cytochrome c. Journal of the American Chemical
Society 134:18713-18723.
38. Arias-Cartin R, Grimaldi S, Arnoux P, Guigliarelli B, Magalon A. 2012. Cardiolipin binding
in bacterial respiratory complexes: Structural and functional implications. Biochimica et
Biophysica Acta (BBA) - Bioenergetics 1817:1937-1949.

39. Planas-Iglesias J, Dwarakanath H, Mohammadyani D, Yanamala N, Kagan Valerian E,
776   Klein-Seetharaman J. 2015. Cardiolipin interactions with proteins. Biophysical Journal
777   109:1282-1294.
778 40. Hirschberg CB, Kennedy EP. 1972. Mechanism of the enzymatic synthesis of cardiolipin
779   in *Escherichia coli*. Proceedings of the National Academy of Sciences 69:648-651.
780 41. Tan BK, Bogdanov M, Zhao J, Dowhan W, Raetz CR, Guan Z. 2012. Discovery of a
781   cardiolipin synthase utilizing phosphatidylethanolamine and phosphatidylglycerol as
782   substrates. Proceedings of the National Academy of Sciences 109:16504-16509.
783 42. Guo D, Tropp BE. 2000. A second *Escherichia coli* protein with CL synthase activity.
784   Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids 1483:263-
785   274.
786 43. Li C, Tan BK, Zhao J, Guan Z. 2016. *In Vivo* and *in Vitro* synthesis of
787   phosphatidylglycerol by an *Escherichia coli* cardiolipin synthase. Journal of Biological
788   Chemistry 291:25144-25153.
789 44. Homma H, Nojima S. 1982. Synthesis of various phospholipids from 2-acyl
790   lysophospholipids by *Escherichia coli* extract. The Journal of Biochemistry 91:1103-1110.
791 45. Patrick S. 2022. A tale of two habitats: *Bacteroides fragilis*, a lethal pathogen and
792   resident in the human gastrointestinal microbiome. Microbiology 168.
793 46. Wexler HM. 2007. *Bacteroides*: The good, the bad, and the nitty-gritty. Clin Microbiol Rev
794   20:593-621.
795 47. Tally FP, Stewart PR, Sutter VL, Rosenblatt JE. 1975. Oxygen tolerance of fresh clinical
796   anaerobic bacteria. J Clin Microbiol 1:161-4.
797 48. Valguarnera E, Wardenburg JB. 2020. Good Gone Bad: One Toxin Away From Disease
798   for Bacteroides fragilis. J Mol Biol 432:765-785.
799 49. Scott N, Whittle E, Jeraldo P, Chia N. 2022. A systemic review of the role of enterotoxic
800   *Bacteroides fragilis* in colorectal cancer. Neoplasia 29:100797.
801 50. Sun F, Zhang Q, Zhao J, Zhang H, Zhai Q, Chen W. 2019. A potential species of next-
802   generation probiotics? The dark and light sides of *Bacteroides fragilis* in health. Food
803   Res Int 126:108590.
804 51. Fiebig A, Schnizlein MK, Pena-Rivera S, Trigodet F, Dubey AA, Hennessy MK, Basu A,
805   Pott S, Dalal S, Rubin D, Sogin ML, Eren AM, Chang EB, Crosson S. 2024. Bile acid
806   fitness determinants of a *Bacteroides fragilis* isolate from a human pouchitis patient.
807   mBio 15:e02830-23.
808 52. Vineis JH, Ringus DL, Morrison HG, Delmont TO, Dalal S, Raffals LH, Antonopoulos DA,
809   Rubin DT, Eren AM, Chang EB, Sogin ML. 2016. Patient-specific *Bacteroides* genome
810   variants in pouchitis. mBio 7:e01713-16.
811 53. Akiyama S, Rai V, Rubin DT. 2021. Pouchitis in inflammatory bowel disease: a review of
812   diagnosis, prognosis, and treatment. Intest Res 19:1-11.
813 54. Sitaraman R. 2013. Phospholipid catabolism by gut microbiota and the risk of
814   cardiovascular disease. J Med Microbiol 62:948-950.
815 55. Romantsov T, Gonzalez K, Sahtout N, Culham DE, Coumoundouros C, Garner J, Kerr
816   CH, Chang L, Turner RJ, Wood JM. 2018. Cardiolipin synthase A colocalizes with
817   cardiolipin and osmosensing transporter ProP at the poles of *Escherichia coli* cells.
818   Molecular Microbiology 107:623-638.
819 56. Quigley BR, Tropp BE. 2009. *E. coli* cardiolipin synthase: Function of N-terminal
820   conserved residues. Biochimica et Biophysica Acta (BBA) - Biomembranes 1788:2107-
821   2113.
822 57. Antonenko YN, Yaguzhinsky LS. 1988. The ion selectivity of nonelectrogenic ionophores
823   measured on a bilayer lipid membrane: Nigericin, monensin, A23187 and lasalocid A.
824   Biochim Biophys Acta 938:125-30.

58. Li J, Ding L, Wang E, Dong S. 1996. The ion selectivity of monensin incorporated
phospholipid/alkanethiol bilayers. Journal of Electroanalytical Chemistry 414:17-21.

59. Łowicki D, Huczyński A. 2013. Structure and antimicrobial properties of monensin A and
its derivatives: Summary of the achievements. BioMed Research International
2013:742149.

60. Lin T-Y, Gross William S, Auer George K, Weibel Douglas B. 2019. Cardiolipin alters
Rhodobacter sphaeroides cell shape by affecting peptidoglycan precursor biosynthesis.
mBio 10.

61. Elías-Wolff F, Lindén M, Lyubartsev AP, Brandt EG. 2019. Curvature sensing by
cardiolipin in simulated buckled membranes. Soft Matter 15:792-802.

62. Luchini A, Cavasso D, Radulescu A, D'Errico G, Paduano L, Vitiello G. 2021. Structural
organization of cardiolipin-containing vesicles as models of the bacterial cytoplasmic
membrane. Langmuir 37:8508-8516.

63. Labajová N, Baranova N, Jurásek M, Vácha R, Loose M, Barák I. 2021. Cardiolipin-
containing lipid membranes attract the bacterial cell division protein DivIVA. International
Journal of Molecular Sciences 22:8350.

64. Renner LD, Weibel DB. 2011. Cardiolipin microdomains localize to negatively curved
regions of *Escherichia coli* membranes. Proceedings of the National Academy of
Sciences 108:6264-6269.

65. Huang KC, Mukhopadhyay R, Wingreen NS. 2006. A curvature-mediated mechanism for
localization of lipids to bacterial poles. PLOS Computational Biology 2:e151.

66. Bailey BWA, Stansfeld PJ. 2024. The bacterial cytoskeleton alters the cell membrane in
molecular dynamics simulations. Biophysical Journal 123:374a.

67. Rowlett VW, Mallampalli VKPS, Karlstaedt A, Dowhan W, Taegtmeyer H, Margolin W,
Vitrac H. 2017. Impact of membrane phospholipid alterations in *Escherichia coli* on
cellular function and bacterial stress adaptation. Journal of Bacteriology
199:10.1128/jb.00849-16.

68. Duncan AL, Robinson AJ, Walker JE. 2016. Cardiolipin binds selectively but transiently
to conserved lysine residues in the rotor of metazoan ATP synthases. Proc Natl Acad Sci
U S A 113:8687-92.

69. Mehdipour AR, Hummer G. 2016. Cardiolipin puts the seal on ATP synthase. Proc Natl
Acad Sci U S A 113:8568-70.

70. Shapiro JS, Chang H-C, Tatekoshi Y, Zhao Z, Waxali ZS, Hong BJ, Chen H, Geier JA,
Bartom ET, De Jesus A, Nejad FK, Mahmoodzadeh A, Sato T, Ramos-Alonso L, Romero
AM, Martinez-Pastor MT, Jiang S-C, Sah-Teli SK, Li L, Bentrem D, Lopaschuk G, Ben-
Sahra I, O'Halloran TV, Shilatifard A, Puig S, Bergelson J, Koivunen P, Ardehali H. 2023.
Iron drives anabolic metabolism through active histone demethylation and mTORC1.
Nature Cell Biology 25:1478-1494.

71. Lane A, Gokhale A, Werner E, Roberts A, Freeman A, Roberts B, Faundez V. 2022.
Sulfur- and phosphorus-standardized metal quantification of biological specimens using
inductively coupled plasma mass spectrometry. STAR Protoc 3:101334.

72. Schaechter M, MaalØe O, Kjeldgaard NO. 1958. Dependency on medium and
temperature of cell size and chemical composition during balanced growth of *Salmonella
typhimurium*. Microbiology 19:592-606.

73. Kjeldgaard NO, MaalØe O, Schaechter M. 1958. The transition between different
physiological states during balanced growth of *Salmonella typhimurium*. Microbiology
19:607-616.

74. Mulyukin AL, Sorokin VV, Loiko NG, Suzina NE, Duda VI, Vorob'eva EA, El'-Registan GI.
2002. Comparative study of the elemental composition of vegetative and resting
microbial cells. Microbiology 71:31-40.

75. Sivakumar P, Fernández-Bravo A, Taleh L, Biddle JF, Melikechi N. 2015. Detection and classification of live and dead *Escherichia coli* by laser-induced breakdown spectroscopy. Astrobiology 15:144-153.

76. Sartorio Mariana G, Valguarnera E, Hsu F-F, Feldman Mario F. 2022. Lipidomics analysis of outer membrane vesicles and elucidation of the inositol phosphoceramide biosynthetic pathway in *Bacteroides thetaiotaomicron*. Microbiology Spectrum 10:e00634-21.

77. Ryan E, Gonzalez Pastor B, Gethings LA, Clarke DJ, Joyce SA. 2023. Lipidomic analysis reveals differences in *Bacteroides* species driven largely by plasmalogens, glycerophosphoinositols and certain sphingolipids. Metabolites 13:360.

78. Rizza V, Tucker Anne N, White David C. 1970. Lipids of *Bacteroides melaninogenicus*. Journal of Bacteriology 101:84-91.

79. Withers H, Swift S, Williams P. 2001. Quorum sensing as an integral component of gene regulatory networks in Gram-negative bacteria. Current Opinion in Microbiology 4:186-193.

80. Otaru N, Ye K, Mujezinovic D, Berchtold L, Constancias F, Cornejo FA, Krzystek A, de Wouters T, Braegger C, Lacroix C, Pugin B. 2021. GABA production by human intestinal Bacteroides spp.: Prevalence, regulation, and role in acid stress tolerance. Frontiers in Microbiology 12.

81. Cohen LJ, Kang HS, Chu J, Huang YH, Gordon EA, Reddy BV, Ternei MA, Craig JW, Brady SF. 2015. Functional metagenomic discovery of bacterial effectors in the human microbiome and isolation of commendamide, a GPCR G2A/132 agonist. Proc Natl Acad Sci U S A 112:E4825-34.

82. Koprivnjak T, Zhang D, Ernst CM, Peschel A, Nauseef WM, Weiss JP. 2011. Characterization of *Staphylococcus aureus* cardiolipin synthases 1 and 2 and their contribution to accumulation of cardiolipin in stationary phase and within phagocytes. Journal of Bacteriology 193:4134-4142.

83. Mirretta Barone C, Heaver SL, Gruber L, Zundel F, Vu DL, Ley RE. 2024. Spatially resolved lipidomics shows conditional transfer of lipids produced by Bacteroides thetaiotaomicron into the mouse gut. Cell Host & Microbe 32:1025-1036.e5.

84. An D, Na C, Bielawski J, Hannun YA, Kasper DL. 2011. Membrane sphingolipids as essential molecular signals for Bacteroides survival in the intestine. Proceedings of the National Academy of Sciences 108:4666-4671.

85. Frankfater CF, Sartorio MG, Valguarnera E, Feldman MF, Hsu F-F. 2023. Lipidome of the Bacteroides Genus Containing New Peptidolipid and Sphingolipid Families Revealed by Multiple-Stage Mass Spectrometry. Biochemistry 62:1160-1180.

86. Hines KM, Xu L. 2019. Lipidomic consequences of phospholipid synthesis defects in *Escherichia coli* revealed by HILIC-ion mobility-mass spectrometry. Chemistry and Physics of Lipids 219:15-22.

87. Czolkoss S, Fritz C, Hölzl G, Aktas M. 2016. Two distinct cardiolipin synthases operate in *Agrobacterium tumefaciens*. PLOS ONE 11:e0160373.

88. Tsai M, Ohniwa RL, Kato Y, Takeshita SL, Ohta T, Saito S, Hayashi H, Morikawa K. 2011. *Staphylococcus aureus* requires cardiolipin for survival under conditions of high salinity. BMC Microbiology 11:13.

89. Liu J, Ryabichko S, Bogdanov M, Fackelmayer OJ, Dowhan W, Krulwich TA. 2014. Cardiolipin is dispensable for oxidative phosphorylation and non-fermentative growth of alkaliphilic Bacillus pseudofirmus OF4. J Biol Chem 289:2960-71.

90. Kusaka J, Shuto S, Imai Y, Ishikawa K, Saito T, Natori K, Matsuoka S, Hara H, Matsumoto K. 2016. Septal localization by membrane targeting sequences and a conserved sequence essential for activity at the COOH-terminus of *Bacillus subtilis* cardiolipin synthase. Research in Microbiology 167:202-214.

91. Shibuya I, Yamagoe S, Miyazaki C, Matsuzaki H, Ohta A. 1985. Biosynthesis of novel
acidic phospholipid analogs in *Escherichia coli*. Journal of Bacteriology 161:473-477.
92. Walderhaug MO, Dosch DC, Epstein W. 1987. Potassium Transport in Bacteria, p 85-
130. *In* Rosen BP, Silver S (ed), Ion Transport in Prokaryotes doi:10.1016/B978-0-12-
596935-2.50005-0. Academic Press.
93. Whatmore AM, Chudek JA, Reed RH. 1990. The effects of osmotic upshock on the
intracellular solute pools of *Bacillus subtilis*. Microbiology 136:2527-2535.
94. Epstein W. 1986. Osmoregulation by potassium transport in *Escherichia coli*. FEMS
Microbiology Reviews 2:73-78.
95. Christian JHB, Waltho JA. 1964. The composition of *Staphylococcus aureus* in relation
to the water activity of the growth medium. Microbiology 35:205-213.
96. Jagannathan S. 1994. Characterization of proton and water permeability across model
membrane vesicles and proteoliposomes. 9600672. Clemson University, South Carolina,
USA.
97. Kato S, Tobe H, Matsubara H, Sawada M, Sasaki Y, Fukiya S, Morita N, Yokota A. 2019.
The membrane phospholipid cardiolipin plays a pivotal role in bile acid adaptation by
Lactobacillus gasseri JCM1131T. Biochimica et Biophysica Acta (BBA) - Molecular and
Cell Biology of Lipids 1864:403-412.
98. Dimroth P. 1994. Bacterial sodium ion-coupled energetics. Antonie Van Leeuwenhoek
65:381-95.
99. Mulkidjanian AY, Dibrov P, Galperin MY. 2008. The past and present of sodium
energetics: may the sodium-motive force be with you. Biochim Biophys Acta 1777:985-
92.
100. Ren Q, Paulsen IT. 2009. Transport, Solute, p 529-544. *In* Schaechter M (ed),
Encyclopedia of Microbiology (Third Edition) doi:10.1016/B978-012373944-5.00107-3.
Academic Press, Oxford.
101. van de Vossenberg JL, Ubbink-Kok T, Elferink MG, Driessen AJ, Konings WN. 1995. Ion
permeability of the cytoplasmic membrane limits the maximum growth temperature of
bacteria and archaea. Mol Microbiol 18:925-32.
102. Deamer DW, Bramhall J. 1986. Permeability of lipid bilayers to water and ionic solutes.
Chemistry and Physics of Lipids 40:167-188.
103. Hopfer U, Lehninger AL, Lennarz WJ. 1970. The effect of the polar moiety of lipids on
the ion permeability of bilayer membranes. The Journal of Membrane Biology 2:41-58.
104. Russell JB. 1987. A proposed mechanism of monensin action in inhibiting ruminant
bacterial growth: Effects on ion flux and proton motive force. Journal of Animal Science
64:1519-1525.
105. Guffanti AA, Davidson LF, Mann TM, Krulwich TA. 1979. Nigericin-induced death of an
acidophilic bacterium. Microbiology 114:201-206.
106. Ouellette SP, Fisher-Marvin LA, Harpring M, Lee J, Rucks EA, Cox JV. 2022. Localized
cardiolipin synthesis is required for the assembly of MreB during the polarized cell
division of *Chlamydia trachomatis*. PLOS Pathogens 18:e1010836.
107. Kurita K, Kato F, Shiomi D. 2020. Alteration of membrane fluidity or phospholipid
composition perturbs rotation of MreB complexes in *Escherichia coli*. Frontiers in
Molecular Biosciences 7.
108. Dong H, Zhang Z, Tang X, Huang S, Li H, Peng B, Dong C. 2016. Structural insights into
cardiolipin transfer from the Inner membrane to the outer membrane by PbgA in Gram-
negative bacteria. Scientific Reports 6:30815-30815.
109. Bogdanov M. 2023. The power and challenge of lipid (a)symmetry across the membrane
and cell. Emerging Topics in Life Sciences 7:1-6.

110. Bogdanov M. 2023. Renovating a double fence with or without notifying the next door
and across the street neighbors: why the biogenic cytoplasmic membrane of Gram-
negative bacteria display asymmetry? . Emerging Topics in Life Sciences 7:137-150.
111. Lande MB, Donovan JM, Zeidel ML. 1995. The relationship between membrane fluidity
and permeabilities to water, solutes, ammonia, and protons. Journal of General
Physiology 106:67-84.
112. Neidleman SL. 1987. Effects of temperature on lipid unsaturation. Biotechnology and
Genetic Engineering Reviews 5:245-268.
113. Yano Y, Nakayama A, Ishihara K, Saito H. 1998. Adaptive changes in membrane lipids of
barophilic bacteria in response to changes in growth pressure. Applied and
Environmental Microbiology 64:479-485.
114. Huffer S, Clark ME, Ning JC, Blanch HW, Clark DS. 2011. Role of alcohols in growth,
lipid composition, and membrane fluidity of yeasts, bacteria, and archaea. Applied and
Environmental Microbiology 77:6400-6408.
115. Keweloh H, Heipieper HJ. 1996. Trans unsaturated fatty acids in bacteria. Lipids 31:129-
137.
116. Altabe SG, Aguilar P, Caballero GM, de Mendoza D. 2003. The *Bacillus subtilis* acyl lipid
desaturase is a delta5 desaturase. J Bacteriol 185:3228-31.
117. Schweizer HP, Choi K-H. 2011. *Pseudomonas aeruginosa* aerobic fatty acid desaturase
DesB is important for virulence factor production. Archives of Microbiology 193:227-234.
118. García-Bayona L, Comstock LE. 2019. Streamlined genetic manipulation of diverse
Bacteroides and *Parabacteroides* isolates from the human gut microbiota. mBio 10.
119. Rajeev L, Segall A, Gardner J. 2007. The *Bacteroides* NBU1 integrase performs a
homology-independent strand exchange to form a holliday junction intermediate. J Biol
Chem 282:31228-37.
120. Hallgren J, Tsirigos KD, Pedersen MD, Almagro Armenteros JJ, Marcatili P, Nielsen H,
Krogh A, Winther O. 2022. DeepTMHMM predicts alpha and beta transmembrane
proteins using deep neural networks. bioRxiv
doi:10.1101/2022.04.08.487609:2022.04.08.487609.
121. Bäckhed F, Ley RE, Sonnenburg JL, Peterson DA, Gordon JI. 2005. Host-bacterial
mutualism in the human intestine. Science 307:1915-1920.
122. Adak A, Khan MR. 2019. An insight into gut microbiota and its functionalities. Cellular
and Molecular Life Sciences 76:473-493.
123. Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, Gwadz M, Hurwitz DI,
Marchler GH, Song JS, Thanki N, Yamashita RA, Yang M, Zhang D, Zheng C, Lanczycki
CJ, Marchler-Bauer A. 2020. CDD/SPARCLE: The conserved domain database in 2020.
Nucleic Acids Res 48:D265-d268.
124. Saha CK, Sanches Pires R, Brolin H, Delannoy M, Atkinson GC. 2020. FlaGs and
webFlaGs: Discovering novel biology through the analysis of gene neighbourhood
conservation. Bioinformatics 37:1312-1314.
125. Wang J, Barger K. 2024. ipolygrowth: An R package to calculate individual growth curve
parameters from bacterial time series data. Microbiology Resource Announcements
13:e00662-24.
126. McLaughlin M, Fiebig A, Crosson S. 2023. XRE transcription factors conserved in
Caulobacter and φCbK modulate adhesin development and phage production. PLOS
Genetics 19:e1011048.
127. Ducret A, Quardokus EM, Brun YV. 2016. MicrobeJ, a tool for high throughput bacterial
cell detection and quantitative analysis. Nature Microbiology 1:16077.
128. Sakol N, Egawa A, Fujiwara T. 2020. Gadolinium complexes as contrast agent for
cellular NMR spectroscopy. Int J Mol Sci 21:4042.

129. Aime S, Cabella C, Colombatto S, Geninatti Crich S, Gianolio E, Maggioni F. 2002.
Insights into the use of paramagnetic Gd(III) complexes in MR-molecular imaging
investigations. Journal of Magnetic Resonance Imaging 16:394-406.
130. Basso JTR, Jones KA, Jacobs KR, Christopher CJ, Fielland HB, Campagna SR, Buchan
A. 2022. Growth Substrate and Prophage Induction Collectively Influence Metabolite and
Lipid Profiles in a Marine Bacterium. mSystems 7:e0058522.
131. Woodall B, Fozo EM, Campagna SR. 2023. Dual stable isotopes enhance lipidomic
studies in bacterial model organism Enterococcus faecalis. Anal Bioanal Chem
415:3593-3605.
132. Tsugawa H, Ikeda K, Takahashi M, Satoh A, Mori Y, Uchino H, Okahashi N, Yamada Y,
Tada I, Bonini P, Higashi Y, Okazaki Y, Zhou Z, Zhu Z-J, Koelmel J, Cajka T, Fiehn O,
Saito K, Arita M, Arita M. 2020. A lipidome atlas in MS-DIAL 4. Nature Biotechnology
38:1159-1163.
133. Tsugawa H, Nakabayashi R, Mori T, Yamada Y, Takahashi M, Rai A, Sugiyama R,
Yamamoto H, Nakaya T, Yamazaki M, Kooke R, Bac-Molenaar JA, Oztolan-Erol N,
Keurentjes JJB, Arita M, Saito K. 2019. A cheminformatics approach to characterize
metabolomes in stable-isotope-labeled organisms. Nature Methods 16:295-298.
134. Lai Z, Tsugawa H, Wohlgemuth G, Mehta S, Mueller M, Zheng Y, Ogiwara A, Meissen J,
Showalter M, Takeuchi K, Kind T, Beal P, Arita M, Fiehn O. 2018. Identifying metabolites
by integrating metabolome databases with mass spectrometry cheminformatics. Nature
Methods 15:53-56.
135. Tsugawa H, Cajka T, Kind T, Ma Y, Higgins B, Ikeda K, Kanazawa M, VanderGheynst J,
Fiehn O, Arita M. 2015. MS-DIAL: data-independent MS/MS deconvolution for
comprehensive metabolome analysis. Nat Methods 12:523-6.
136. Friedman JH, Hastie T, Tibshirani R. 2010. Regularization paths for generalized linear
models via coordinate descent. Journal of Statistical Software 33:1 - 22.
137. Tay JK, Narasimhan B, Hastie T. 2023. Elastic net regularization paths for all generalized
linear models. Journal of Statistical Software 106:1 - 31.
138. Mason Allison R, Johnson JG, Krampen J, Nguyen Jennifer NT, Balunas Marcy J,
Schloss Patrick D. 2025. mpactR: An R adaptation of the metabolomics peak analysis
computational tool (MPACT) for use in reproducible data analysis pipelines. Microbiology
Resource Announcements 14:e00997-24.
139. Wang M, Carver JJ, Phelan VV, Sanchez LM, Garg N, Peng Y, Nguyen DD, Watrous J,
Kapono CA, Luzzatto-Knaan T, Porto C, Bouslimani A, Melnik AV, Meehan MJ, Liu W-T,
Crüsemann M, Boudreau PD, Esquenazi E, Sandoval-Calderón M, Kersten RD, Pace
LA, Quinn RA, Duncan KR, Hsu C-C, Floros DJ, Gavilan RG, Kleigrewe K, Northen T,
Dutton RJ, Parrot D, Carlson EE, Aigle B, Michelsen CF, Jelsbak L, Sohlenkamp C,
Pevzner P, Edlund A, McLean J, Piel J, Murphy BT, Gerwick L, Liaw C-C, Yang Y-L,
Humpf H-U, Maansson M, Keyzers RA, Sims AC, Johnson AR, Sidebottom AM, Sedio
BE, et al. 2016. Sharing and community curation of mass spectrometry data with Global
Natural Products Social Molecular Networking. Nature Biotechnology 34:828-837.
140. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski
B, Ideker T. 2003. Cytoscape: A software environment for integrated models of
biomolecular interaction networks. Genome Res 13:2498-504.
141. Pang Z, Lu Y, Zhou G, Hui F, Xu L, Viau C, Spigelman Aliya F, MacDonald Patrick E,
Wishart David S, Li S, Xia J. 2024. MetaboAnalyst 6.0: Towards a unified platform for
metabolomics data processing, analysis and interpretation. Nucleic Acids Research
52:W398-W406.
142. Team RDC. 2013. R: A language and environment for statistical computing, R
Foundation for Statistical Computing, http://www.R-project.org/.

143. Wickham H. 2016. ggplot2: Elegant graphics for data analysis, Springer-Verlag, New
York. https://ggplot2.tidyverse.org.
144. Wong B. 2011. Points of view: Color blindness. Nature Methods 8:441-441.
145. Ou J. 2021. colorBlindness: Safe color set for color blindness, https://CRAN.R-
project.org/package=colorBlindnes.
146. Neuwirth E. 2022. RColorBrewer: ColorBrewer palettes, https://CRAN.R-
project.org/package=RColorBrewer.
147. Vu VQ, Friendly M. 2024. ggbiplot: A grammar of graphics implementation of biplots,
https://CRAN.R-project.org/package=ggbiplot.
148. Pedersen TL. 2024. ggforce: Accelerating 'ggplot2', https://CRAN.R-
project.org/package=ggforce.
149. Gu Z, Gu L, Eils R, Schlesner M, Brors B. 2014. circlize implements and enhances
circular visualization in R. Bioinformatics 30:2811-2.
150. Gu Z, Eils R, Schlesner M. 2016. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. Bioinformatics 32:2847-9.
151. Neitmann T. 2020. mdthemes: Markdown themes for 'ggplot2', https://CRAN.R-
project.org/package=mdthemes.
152. C W, B W. 2022. ggtext: Improved text rendering support for 'ggplot2', https://CRAN.R-
project.org/package=ggtext.
153. Wickham H, Bryan J. 2023. readxl: Read excel files, https://CRAN.R-
project.org/package=readxl.
154. Wickham H, Vaughan D, Girlich M. 2024. tidyr: Tidy messy data, https://CRAN.R-
project.org/package=tidyr.
155. Bengtsson H. 2024. matrixStats: Functions that apply to rows and columns of matrices
(and to

vectors), https://CRAN.R-project.org/package=matrixStats.
156. Barrett T, Dowle M, Srinivasan A, Gorecki J, Chirico M, Hocking T. 2024. data.table:
Extension of `data.frame`, https://CRAN.R-project.org/package=data.table.
157. Bache SM, Wickham H. 2022. magrittr: A forward-pipe operator for R, https://cran.r-
project.org/web/packages/magrittr/vignettes/magrittr.html.
158. Wickham H, Pedersen TL, Seidel D. 2023. scales: Scale functions for visualization,
https://CRAN.R-project.org/package=scales.
159. Wang L-G, Lam TT-Y, Xu S, Dai Z, Zhou L, Feng T, Guo P, Dunn CW, Jones BR, Bradley
T, Zhu H, Guan Y, Jiang Y, Yu G. 2019. Treeio: An R package for phylogenetic tree input
and output with richly annotated and associated data. Molecular Biology and Evolution
37:599-603.
160. Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2017. ggtree: An R package for visualization
and annotation of phylogenetic trees with their covariates and other associated data.
Methods in Ecology and Evolution 8:28-36.
161. Xu S, Li L, Luo X, Chen M, Tang W, Zhan L, Dai Z, Lam TT, Guan Y, Yu G. 2022. Ggtree:
A serialized data object for visualization of a phylogenetic tree and annotation data.
iMeta 1:e56.
162. Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using
lme4. J of Statistical Software 67:1-48.
163. Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. lmerTest package: Tests in linear
mixed effects models. J of Statistical Software 82:1-26.
164. A K. 2023. ggpubr: 'ggplot2' based publication ready plots, https://CRAN.R-
project.org/package=ggpubr.

165. Yurekten O, Payne T, Tejera N, Amaladoss FX, Martin C, Williams M, O'Donovan C.
2023. MetaboLights: open data repository for metabolomics. Nucleic Acids Research
52:D640-D646.
166. Vingadassalom D, Kolb A, Mayer C, Rybkine T, Collatz E, Podglajen I. 2005. An unusual
primary sigma factor in the Bacteroidetes phylum. Mol Microbiol 56:888-902.
167. Loferer-Krössbacher M, Klima J, Psenner R. 1998. Determination of bacterial cell dry
mass by transmission electron microscopy and densitometric image analysis. Appl
Environ Microbiol 64:688-94.
168. Epanechnikov VA. 1969. Non-parametric estimation of a multivariate probability density.
Theory of Probability & Its Applications 14:153-158.
169. Poynton EF, van Santen JA, Pin M, Contreras MM, McMann E, Parra J, Showalter B,
Zaroubi L, Duncan Katherine R, Linington RG. 2024. The Natural Products Atlas 3.0:
Extending the database of microbially derived natural products. Nucleic Acids Research
53:D691-D699.
170. Wang F, Allen D, Tian S, Oler E, Gautam V, Greiner R, Metz Thomas O, Wishart
David S. 2022. CFM-ID 4.0 – a web server for accurate MS-based metabolite
identification. Nucleic Acids Research 50:W165-W174.
171. Conroy MJ, Andrews RM, Andrews S, Cockayne L, Dennis Edward A, Fahy E, Gaud C,
Griffiths William J, Jukes G, Kolchin M, Mendivelso K, Lopez-Clavijo Andrea F, Ready C,
Subramaniam S, O'Donnell Valerie B. 2023. LIPID MAPS: update to databases and
tools for the lipidomics community. Nucleic Acids Research 52:D1677-D1682.
1144

1145

FIGURES AND TABLES

Figure 1. Cardiolipin synthases diverge across the dominant gut bacterial phyla. A. Percentage

identity from pairwise amino acid alignment of cardiolipin synthases in *B. fragilis* P207 (*Bf*) and *E. coli* K12

MG1655 (*Ec*). **B.** Alignment of ClsA and ClsB from *B. fragilis* P207 and *E. coli* MG1655, with the two HKD

domains marked in blue. **C.** Percentage identity from amino acid alignment in panel B colored based on the

thresholds <30% (pink), <100% (orange) and 100% (blue). **D.** Consensus and individual sequences of the

regions flanking the two HKD active sites of *B. fragilis* and *E. coli* Cls proteins. HKD residues are colored

in blue; the N- and C-terminal positions are shown. **E.** Relative abundance of cardiolipin in *B. fragilis* P207

wild-type (WT), Δ*clsA*, Δ*clsB* and Δ*clsA* Δ*clsB* strains measured by mass spectrometry. The horizontal line

indicates the median, and the whiskers 1.5× the interquartile range outside the 25th and 75th percentiles of

n = 3 samples. (linear regression; unadjusted p-values are shown; *** indicates *p* < 0.001). **F.** Neighbor-

joining tree of a Cls alignment from representative taxa of major bacterial phyla in the gut, using only one

genome per species. For each sequence in the tree, taxa names are followed by genome IDs at the tip of

each branch; genome accession numbers and other metadata are provided in Table S2. Bootstrap values

are shown only if >80%. Bacterial taxa are colored according to the phylum to which they belong (red,

Pseudomonadota – 3 classes with 1, 1, and 3 families; brown, Actinomycetota – 1 class; blue, Bacillota –

3 classes; violet, Fusobacteriota – 1 class; green, Bacteroidota – 2 classes, respectively). Key indicates the

rate of amino acid substitutions per site.

1165

1166

Figure 2. Loss of cardiolipin synthase activity results in growth defects under stress. Optical density

(OD_{600}) measurements of *B. fragilis* P207 wild-type (WT) and *cls* single or double mutant strains exposed

to the detergents (**A.**) 0.01% deoxycholate or (**B.**) 0.002% SDS; the salts (**C.**) 300 mM K^+ or (**D.**) 300 mM

Na^+; (**E.**) medium adjusted to pH 5.5; and the ionophores (**F.**) 0.1 µg/mL nigericin or (**G.**) 0.8 µg/mL

monensin A. Untreated (solid lines) and treated cultures (dashed lines) are plotted by strain, with blue,

green and orange/purple indicating *clsA*-, *clsB*- and *clsA clsB*-related strains, respectively. Black indicates

WT. The shaded areas around each line indicate $2\times$ the standard error of the mean of n = 3 biological

replicates.

1176

1177

Figure 3. Loss of cardiolipin synthase activity is associated with defects in cell width. A. Summary

of the cell width distributions. Horizontal line lengths indicate the 25% and 75% quartiles of the distribution;

circles indicate the median; triangles indicates the mode (i.e., the peak of the distribution). **B.** Cell width

data for all cells shown in A. plotted by strain (*n* = 323 cells/strain). The gray or black bars indicate the

mean. The table at right shows the results of a mixed-effect linear model comparing strain width to the

widths of either WT cells or Δ*clsA* Δ*clsB* double mutant (ΔΔ) cells, as indicated at the top of the column,

with a random intercept of biological replicate (i.e., cell width ~ strain + 1|biological replicate). Unadjusted

p-values are shown; *, *p* < 0.05, *** *p* < 0.001.

1186

1187

Figure 4. Cardiolipin synthase genes have distinct expression patterns. A. RT-qPCR analysis of

relative levels of *clsA* and *clsB* transcripts, shown as the $\log_2$ ratios between log- and stationary-phase

growth. **B.** Expression levels of *clsA* or *clsB* when the other gene is knocked out in log- or stationary-phase

growth. **C.** *clsA* and *clsB* expression levels in WT *B. fragilis* P207 after 20 min exposure to of 0.01%

deoxycholate, 0.002% SDS, 300 mM K, pH 5.5, pH 9.0, 0.8 μg/mL monensin A, or 0.1 μg/mL nigericin. For

all panels, values are means ± standard deviation from three biological replicates, each biological replicate

value being the mean of three technical replicates. The vertical brackets for statistical analysis assess the

significance of the difference between conditions in A and C, between mutant and WT in B. The horizontal

brackets for statistical analysis assess the significance of the difference between ΔΔCt values for *clsA* and

clsB (Students *t*-test). Genes amplified are colored in blue (*clsA*) or green (*clsB*). Error bars indicate

standard deviation. Unadjusted p-values are shown; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

1199

1200

1201

1202

Figure 5. Stress but not loss of Cls function disrupts membrane ion gradients. A., B. Abundance of
(**A.**) potassium (K) and (**B.**) sodium (Na) normalized to intracellular phosphorus (P) in *B. fragilis* P207 WT,
Δ*clsA*, Δ*clsB* and complemented strains as well as the Δ*clsA* Δ*clsB* strain, as detected by inductively
coupled plasma mass spectrometry (ICP-MS). **C., D.** Differential concentrations of (**C.**) K$^+$ and (**D.**) Na$^+$ in
WT strain cells after a 20 min exposure to 0.01% deoxycholate, 0.8 µg/mL monensin A or 0.1 µg/mL nigericin
(linear regression). The intracellular concentrations (i.e., normalized to cell volume and number of colony-
forming units) with the extracellular concentrations subtracted are shown. For all boxplots, gray or black
lines indicate the median; the top and bottom edges of the box indicate the 25th and 75th percentiles of the
data; the error bars indicate standard deviation (10 biological replicates for untreated conditions and three
for the treatment conditions). Unadjusted p-values are shown; *, p < 0.05; **, p < 0.01; ***, p < 0.001.

1213

1214

Figure 6. Paralogous cardiolipin synthases produce unique lipid products. A. Structures of lipid

families detected by UHPLC-MS/HRMS. For fatty acids, the R-group is H; for saccharolipids, a sugar (i.e.,

lipid A); and for acyl carnitines, a carnitine. The R-group is H for phosphatidic acid and ethanolamine for

phosphatidylethanolamine. Phospholipids are shown as members of three categories: ester-linked (i.e.,

classical phospholipids), ether-linked and lysophospholipids. Diacylglycerols and monogalactosyl /

monoglucosyl-diacyglycerols have H or a hexose ring as an R-group, respectively. Ceramides, which can

be conjugated to a hexose (i.e., hexosylceramide), are synthesized by *N*-acylating a sphingosine. As

phosphorylated ceramides, sphingolipids have similar R-groups as phospholipids. Length and unsaturation

of the acyl tails are not shown. **B.** Relative abundance of lipid species across WT, Δ*clsA,* Δ*clsB and* Δ*clsA*

Δ*clsB* strains (three biological replicates). Lipid families are abbreviated as follows: PC indicates

phosphatidylcholine; PA, phosphatidic acid; PE, phosphatidylethanolamine; PG, phosphatidylglycerol; PS,

phosphatidylserine; CL, cardiolipin. L-, ML- and DL- indicate lyso-, monolyso- and dilyso-, respectively. **C.**

Relative abundance of cardiolipin, monolyso-cardiolipin and dilysocardiolipin (linear regression).

Unadjusted p-values are shown; ***, $p < 0.001$. **D.** Principal component analysis plot of lipid species in each

strain, with the minimum volume–enclosing ellipses estimated by the Khachiyan algorithm. **E.** Heatmap

representation of the $\log_2$(fold change) of lipid species in each knockout strain compared to WT. Columns

are clustered by similarity and rows by K-means. The left side of the heatmap shows lipid families and K-

means clusters.

1233

1234

Figure 7. Cardiolipin synthases shape the length and unsaturation level of membrane lipid acyl chains. A. log$_2$(fold change) of the total peak areas for all lipids in k-means clusters compared to the mean of WT peak areas of that particular cluster (linear regression of peak areas compared to WT). **B., C.** Relative

distribution of sum of acyl chain (**B.**) carbons and (**C.**) unsaturations in cardiolipin and monolysocardiolipin

species of Km clusters 1 (ClsB-associated) and 3 (ClsA-associated) in WT cells. **D., E.** $\log_2$(fold change) of

total two acyl-tailed phospholipids in $\Delta clsA$ $\Delta clsB$ strain cells compared to WT shown by (**D.**) acyl chain

length and (**E.**) unsaturations. Student's *t*-test comparing values at designated x-axis values); nd, not

detected (i.e., 0) in the $\Delta clsA$ $\Delta clsB$ strain. **F., G., H.** $\log_2$(fold change) of lipids detected in negative (**F.**) and

positive (**G., H.**) ionization mode UHPLC-MS/HRMS comparing each strain to the mean WT peak area.

AHexCer was detected in only 2 and 1 replicates of $\Delta clsB$ and $\Delta clsA$ $\Delta clsB$, respectively. Lipid families are

abbreviated as follows: DG, diacylglycerol; MG, monogalactosyl/monoglucosyl conjugation; NA, *N*-acyl,

with substituent conjugations indicated for glycyl- (Gly), glycylseryl- (GlySer, ornithine- (Orn), and

ethanolamine- (E); CER, ceramide, with A indicating acyl-, P indicating phospho- and Hex, Hex2 and Hex3

indicating hexosyl-, dihexosyl-, and trihexosyl-; HDS, hydroxy fatty acid-dihydrosphingosine; SM,

sphingomyelin; Ether-, ether-linked (linear regression compared to WT). Across all panels, error bars

indicate standard deviation. Unadjusted p-values are shown; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

1251

Figure 8. Cardiolipin synthases disrupt metabolic homeostasis. A. NMDS of metabolomics data

colored by strain. **B.** Upset plot with the main bar plot color coded for the metabolomic features annotation

sources (dark green = lipid databases, purple = small molecule databases, grey = not annotated within

5ppm error) and the lower section color coded for metabolomic features only present or absent in each

strain (grey = WT, blue = Δ*clsA,* green = Δ*clsB*, orange = Δ*clsA* Δ*clsB*). Inset is a rescaled axis of

metabolites missing in at least one strain. **C.** Heatmap of row normalized abundance of each metabolite in

each strain, clustering by metabolomic features on the vertical axis and samples on the horizontal axis

using Euclidean distance and Ward clustering (3 technical replicates shown for each strain). **D.** Metabolic

pathway analysis with nonlipid sub-categories of *clsAclsB* knockout compared to wildtype (≥3

features/pathway). The x-axis represents the enrichment factor computed from pathway topological

analysis on non-lipid species, and the y-axis is the log of p-value obtained from pathway enrichment

analysis. The pathways that were most significantly changed are characterized by both a high-log(p) value

and high enrichment factor (top right region). **E., F.** Abundance of compounds in enriched pathways for (E)

lipid-conjugated homoserine lactones and (F) fatty amides (top enriched pathways) (mean and standard

deviation of 3 technical replicates shown; linear regression compared to WT). Across all panels, error bars

indicate standard deviation. Unadjusted p-values are shown; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

1269

1270

Figure 9. Cardiolipins mitigate the fitness cost of membrane stress in *B. fragilis*. In unperturbed cells, ion gradients are maintained across the inner membrane, with higher extracellular concentrations of H$^+$ and Na$^+$ and elevated intracellular K$^+$ (top left). The outer membrane is generally permeable to these ions due to the presence of porins. In the absence of cardiolipin, *B. fragilis* remodels its membrane lipid composition, which presumably helps to maintain ion gradients (bottom left). Chemical membrane (or membrane ion gradient) perturbation alters *cls* gene expression and ion homeostasis (top right). In the absence of cardiolipins, cells are more susceptible to chemical membrane perturbations (bottom right). The larger blue star indicates the greater fitness cost associated with membrane perturbation in cardiolipin-deficient cells, particularly *clsB* mutants. In contrast, the smaller red star reflects reduced stress susceptibility in cells that retain cardiolipin synthesis.

1281

Figure S1. Neighbor-joining tree of cardiolipin synthases from *Bacteroides* species. Clusters 1

(ClsA), 2 (ClsB) and 3 (not named) of *Bacteroides* Cls proteins are indicated. *E. coli* ClsA was designated

as an outgroup. ClsA and ClsB of *B. fragilis* P207 are colored in blue and green, respectively. Other strains

1285   of *B. fragilis, B. thetaiotaomicron*, and *E. coli* are colored in gold, purple and light blue, respectively. Taxa

1286   labels with similar tree positions are combined for clarity at times (i.e., *B. xylanisolvens/caecimuris*). Key

1287   indicates the rate of amino acid substitutions per site. Bootstrap values are generally shown only if ≥90.

1288

1290   **Figure S2. Genomic neighborhood synteny indicates the ancestral introduction of *clsA* and *clsB***

1291   **into the *Bacteroides.*** Curated WebFlags2 output of (**A.**) *clsA* and (**B.**) *clsB* genomic neighborhoods. *B.*

1292   *fragilis* is colored blue. Genera of interest with similar levels of gene conservation are indicated at right.

1293

1294

1295

Figure S3. Loss of cardiolipin synthase activity results in growth defects under stress. A., B. $\log_2$(fold

change) of (**A.**) doubling time and (**B.**) maximum OD_{600} comparing *B. fragilis* wild-type (WT) and *cls* mutants

to the average of WT in that condition. Metrics were calculated from 24-h growth curves in BHIS medium

alone or containing 0.01% deoxycholate (Dc), 0.002% sodium dodecyl sulfate (SDS), 300 mM sodium

(Na^+), 300 mM potassium (K^+), and 0.01 µg/mL nigericin and 0.8 µg/mL monensin A. Any doubling time

≥100 h was considered as no growth. The dashed horizontal line indicates no change from the average of

WT. Strain are colored with blue (*clsA-related*), green (*clsB*-related) and orange/purple (*clsA clsB*-related).

Black indicates WT. The vertical lines indicate mean; the ends of the whiskers ± standard deviation (linear

regression compared to WT). Three biological replicates for each condition were run, each run in technical

triplicate. nf, no accurate fit of growth data by the polynomial was found. Unadjusted p-values are shown;

*, p < 0.05; **, p < 0.01; ***, p < 0.001.

1307

1308

1309 **Figure S4. Loss of cardiolipin synthase activity is associated with defects in cell width. A.** Phase-

1310 contrast microscopy images of *B. fragilis* P207 WT, Δ*clsA*, Δ*clsB* and Δ*clsA* Δ*clsB* strains. Scale bars, 5 µm.

1311 **B., C.** Distributions estimated by kernel density of (**B.**) cell length and (**C.**) cell width. Results are plotted by

1312 strain with blue, green and orange/purple indicating *clsA*-, *clsB*- and *clsA clsB*-related strains, respectively.

1313 Black indicates WT.

1314

Figure S5. Washing but not loss of *cls* genes disrupts intracellular elemental concentrations. Inductively coupled plasma mass spectrometry (ICP-MS) analysis using wash-free sample preparation to quantify intracellular elemental concentrations. **A.** $\log_2$(fold change) of elements comparing unwashed cells to cell pellets washed with either BHIS growth medium or an isotonic sucrose solution. Error bars indicate standard deviation. **B.** Concentrations of detected elements in BHIS growth medium. **C.** Intracellular concentrations of detected elements in WT *B. fragilis* P207. **D.** Intracellular sulfur in *B. fragilis* P207 strains normalized to P. **E.** Intracellular phosphorus (P) per 10^6 colony-forming units (CFU) in *B. fragilis* P207 wild-type (WT), $\Delta clsA$, $\Delta clsB$ and $\Delta clsA$ $\Delta clsB$ strains. **F.** OD_{600} normalized to CFU per mL of strain cultures used in ICP-MS input. **G.** Raw CFU/mL of each strain shown in panels F and G. **H.** Intracellular P content compared to sulfur (S) content across *cls* strains and treatment conditions. **I., J.** Intracellular concentrations of (**I.**) K^+ and (**J.**) Na^+ after a 20 min exposure to 0.01% deoxycholate, 0.8 µg/mL monensin A and 0.1 µg/mL nigericin. Each treatment had the same culture input. **K.** Intracellular P concentrations of *B. fragilis* WT cells

after 20 min of exposure to the indicated stress conditions. **L.** CFU/mL of strain cultures after exposure to

the treatments in **I–L**. For all boxplots, the gray or black middle lines indicate the median; the top and

bottom edges of the box indicate the 25th and 75th percentiles of n = 3 replicates for wash and treatment

experiments and n = 10 replicates for rest. Whiskers indicate 1.5× the interquartile range (IQR). For the

linear regression, the shade ribbon indicates standard error of the mean. All statistical comparisons were

made using linear regression, with unadjusted p-values shown; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

1334

1335

1336

Figure S6. Cardiolipin synthases shape the membrane lipid population. A-J. Relative abundance of

lipid species in membranes across *B. fragilis* WT, Δ*clsA*, Δ*clsB*, and Δ*clsA* Δ*clsB* strains in positive (**A.**) or

negative (**B-J.**) electrospray ionization modes for UHPLC–MS/HRMS. For **A,** colors designate lipid family.

Positive-mode lipid families are abbreviated as follows: BMP, bismonoacylglycerophosphate; SL,

saccharolipids; DG, diacylglycerol; EtherMG, ether-linked with monogalactosyl/monoglucosyl conjugation;

NA, *N*-acyl, with substituent conjugations indicated for glycyl- (Gly), glycylserine- (GlySer), and

ethanolamine- (E); CAR, acylcarnitine; CER, ceramide, AP indicating alpha-hydroxy fatty acid-

phytospingosine and Hex indicating hexosyl-; SM, sphingomyelin; DHSph, sphinganine (please see the

legend for panel E for other abbreviations). For **B-J**, species are colored based on the total number of

carbons in their acyl chains. Lipids at <1% abundance are colored gray, with the number of lipid species in

this category indicated in the figure. Relative abundance plots are grouped as (**B.**) two or (**C.**) one acyl-

tailed phospholipid species, with oxidized and ether-linked lipids shown in **E.-G.**, or (**H.**) dilysocardiolipin,

(**I.**) monolysocardiolipin or (**J.**) cardiolipin species. Negative-mode lipid families are abbreviated as follows:

PA, phosphatidic acid; PE, phosphatidylethanolamine; PS, phosphatidylserine; PG, phosphatidylglycerol;

PC, phosphatidylcholine; CL, cardiolipin. L-, ML- and DL- indicate lyso-, monolyso- and dilyso-, respectively.

Ox and Ether indicate oxidized and ether-linked, respectively. **K.** $\log_2$(fold change) peak area of

dilysocardiolipin species comparing $\Delta clsA$ $\Delta clsB$ to WT (linear regression, $p < 0.00029$; $\log_{10}$[peak area] ~

strain + DLCL species). Error bars indicate standard deviation.

1355

Figure S7. Cardiolipin synthases produce unique lipid products and their loss remodels overall

membrane lipid composition. A., B. Relative abundance of lipids from K-means clusters 1 and 2 across

B. fragilis WT, Δ*clsA*, Δ*clsB* and Δ*clsA* Δ*clsB* strains plotted by (**A.**) acyl chain length or (**B.**) acyl chain

unsaturations. **C.** Principal component analysis plot of lipids in the *B. fragilis* membrane excluding

cardiolipin and monolysocardiolipin species. Minimum volume-enclosing ellipses were estimated using

the Khachiyan algorithm. **D., E.** $\log_2$(fold change) of Pa, PE, PG and PS lipids in Δ*clsA* Δ*clsB* cells

compared to WT plotted by the sum of acyl chain (**D.**) carbons or (**E.**) unsaturations. **F., G.** $\log_2$(fold

change) of lipids detected in (**F.**) negative or (**G.**) positive ionization modes with values for each strain

compared to the mean abundance in WT cells. See figures 7 and S6 for lipid family abbreviations. Error

bars indicate standard deviation. Unadjusted p-values are shown.

1368

1369

1370 **Figure S8. Pathway enrichment analysis for main lipid categories.** Pathway analysis is performed

1371 with MetaboAnalyst 6.0, with default mummichog parameters with p-value cutoff of 0.0005 and lipid –

1372 main chemical class as the selected metabolite sets containing at least 3 entries.

1373

1374

1375

Figure S9. Molecular networking shows chemically related metabolites in *B. fragilis* strains. A.

Feature-based molecular networking of all features (excluding singletons) from WT, Δ*clsA*, Δ*clsB*, Δ*clsA*

Δ*clsB*, and extracted BHIS media. Each feature is a node, connected by an edge calculated from cosine

similarity score. **B.** Subset of features connected by an edge to fatty amides highlighted in Fig. 8 (red

border). Each pie chart in the node is colored by relative abundance of the feature in each strain.

Table S1. Primers, vectors and strains used in this study.

Table S2. Metadata for genomics analyses.

Table S3. Raw data for the lipidomics analyses.

Table S4. Raw data for the ICP-MS analyses.

Table S5. Raw data for untargeted metabolomics analyses.

Table S6. Pathway enrichment analysis for main lipid categories.

Table S7. Pathway enrichment analysis for sub non-lipid categories.

1388