Hematopoietic Single Cell Atlas Reveals a Diverse Repertoire of Lymphoid Cells in Larval Zebrafish

Anastasia Nizhnik[1,2], Bianca A. Ulloa[1,2], Kevyn Jackson[5], Deyou Zheng[5,6,7], Teresa V. Bowman[#1,2,3,4]

1. Albert Einstein College of Medicine, Department of Developmental and Molecular Biology Department, Bronx, NY, USA
2. Albert Einstein College of Medicine, Gottesman Institute of Stem Cell Biology and Regenerative Medicine, Bronx, NY, USA
3. Albert Einstein College of Medicine and Montefiore Medical Center, Department of Oncology, Bronx, NY, USA
4. Albert Einstein College of Medicine and Montefiore Medical Center, Department of Medicine, Bronx, NY, USA
5. Albert Einstein College of Medicine, Department of Genetics, Bronx, NY, USA
6. Albert Einstein College of Medicine and Montefiore Medical Center, The Saul R. Korey Department of Neurology, Bronx, NY, USA
7. Albert Einstein College of Medicine, Department of Neuroscience, Bronx, NY, USA

[#] To whom correspondence should be addressed.

Address: 1300 Morris Park Avenue, Bronx, NY 10461
Email: teresa.bowman@einsteinmed.edu
Phone: (718) 430-4001
Fax: (718) 430-8567

Word count: 5,997
Abstract/first paragraph word count: 149
Number of figures: 7 main, 7 supplemental figures, 6 supplemental tables
Number of references: 76

Keywords:
Hematopoietic stem cells, HSCs, HSPCs, zebrafish, differentiation, lineage tracing, single cell RNA sequencing, Innate Lymphoid Cells, ILCs, hematopoietic ontogeny

SUMMARY

During embryogenesis, hematopoietic stem cells (HSC) and HSC-independent progenitor cells form concurrently and produce overlapping and distinct immune cell types. The full repertoire of immune cells generated during development and their origin across vertebrates is incompletely elucidated. Here, we performed temporal lineage tracing of the emerging hematopoietic system during zebrafish larval development followed by single cell RNA sequencing and identified a greater diversity of lymphoid cells than previously recognized. The cells are comprised of T-lymphocytes and Innate Lymphoid-like Cells (ILCs) that are Runx1-dependent. Both T-lymphocytes and ILC-like cells depended on Il2rγ, while only T-lymphocytes depended on Rag1. The larval ILC-like cells were detected in lymphoid and mucosal organs and were responsive to viral mimicry-induced stimulation, indicating their functionality in early vertebrate life. The work provides new fundamental knowledge on the early establishment of immune hierarchies and opens the zebrafish model to broader exploration of lymphoid immunity origination and function.

INTRODUCTION

The hematopoietic system is comprised of an array of blood cells carrying out a variety of functions ranging from oxygenation to pathogen defense. Embryonic development is a critical stage for the establishment of hematopoietic and immune cell types, where multiple sequential and overlapping 'waves' of hematopoiesis produce progenitors and mature cell types[1-3]. The variety of cells generated at each wave serve a particular purpose for the proper establishment and function of lifelong hematopoietic and immune systems. Currently the field refers to two main waves of hematopoiesis: primitive and definitive[3]. The 'primitive wave' gives rise largely to macrophages and, erythrocytes[4], and neutrophils[5]. The 'definitive wave' is subdivided into an earlier pro-definitive phase and a definitive phase[6]. The pro-definitive phase progenitors generate myeloid lineage cells, megakaryocytes, NK cells[7], and B-cells. The definitive phase produces Hematopoietic Stem Cells (HSCs) that can sustain lifelong hematopoiesis.

During the definitive wave, hemogenic endothelial cells give rise to Hematopoietic Stem and Progenitor Cells (HSPCs) which then migrate to secondary sites of hematopoiesis like the fetal liver and thymus[8-10]. The emerging HSPCs are not uniform, but rather heterogeneous[11-13]. Within the pool of HSPCs, there are embryonic HSC-independent progenitors that emerge simultaneously with the definitive HSCs. Lineage tracing studies in mice and zebrafish are establishing the connections between emerging HSPCs and the mature cell types that derive from them during development[14-16]. However, HSCs and HSC-independent progenitors' contribution to immune and hematopoietic systems remain poorly resolved due to lack of specific genetic markers. The establishment and layering of the hematopoietic and immune systems during development remains to be fully elucidated.

Embryonic HSCs and HSC-independent progenitors differentiate to produce a diverse repertoire of adaptive and innate immune cells, encompassing various cells of the myeloid and lymphoid lineages[2,17]. While the HSC-independent origins of tissue-resident macrophages and B-1a lymphocytes are studied, origins of many other immune cell types such as the Innate Lymphoid Cells (ILCs) are poorly resolved. ILCs belong to the lymphoid lineage and resemble T-helper lymphocytes, however ILCs lack the adaptive antigen-specific receptors characteristic of T lymphocytes[18]. ILCs are grouped into three categories: ILC1, ILC2, and ILC3, paralleling the T-helper groups Th1, Th2, and Th17[19]. Generally, ILC1s express *Eomes* and *T-bet,* ILC2s express *Il13, Il4* and *Gata3*, while ILC3s express *Il17, Il22* and *Rorc*[20]. In response to infections, ILCs can secrete cytokines and recruit other immune cells[21]. The various groups of ILCs were identified in humans, mice, and zebrafish models, however their embryonic origins remain unknown in zebrafish and incompletely characterized in other model systems.

We sought to resolve the cell types generated by HSCs and HSC-independent progenitors in zebrafish using a temporally inducible fluorescent reporter-based lineage tracing system followed by single cell RNA sequencing (scRNA-seq). We find that zebrafish HSPCs can be temporally resolved with discrete functions, whereby early HSPCs produce a burst of lymphoid lineage differentiation, and late emerging HSPCs generate transcriptionally different macrophages and neutrophils. Furthermore, we

uncovered that HSPCs make a diverse set of previously undescribed immune cells in

zebrafish, including various groups of ILCs and several other novel immune cells. The

ILC-like cells in zebrafish are present as early as 3 days-post-fertilization (dpf) and require

the function of the transcription factor Runx1. Moreover, we demonstrated that viral

mimicry stimulated the larval immune system with strong responses from NK/ILC1-like

cells, macrophages and il2rb+T cells, which expand in numbers when treated with the

TLR7/8 receptor agonist R848. To aid in future inquiries of developmental hematopoiesis

and immunology, we share all scRNA-seq datasets as a resource for the research

communities.

RESULTS

Lineage tracing and scRNA-seq were employed for resolving early, mid and late

HSPCs in zebrafish.

During embryonic and larval development, long-lived HSCs and HSC-independent

progenitors display distinct lymphoid-myeloid differentiation kinetics[14-16]. HSC-

independent progenitors are the first to differentiate into mature immune cells, while HSC-

derived progeny appear later[14]. While prior studies established this new paradigm, they

were limited in the cell types they could examine due to technical limitations. The

hematopoietic system is diverse, with numerous immune cell types that serve an array of

functions and are found localized in many tissues. In a previous zebrafish lineage tracing

study[14], neutrophils and thymic cells were examined, but it is unclear if the differentiation

repertoire and kinetics that distinguish HSC and HSC-independent progenitors in larvae

is unique to these cell types or more generalizable throughout the hematopoietic system.

To address this question, we used an inducible Cre-Lox-based genetic lineage tracing

approach that fluorescently labels HSPCs and their progeny, coupled with scRNA-seq to

delineate the identities of labeled cells (**Figure 1A**). The Cre-Lox system is comprised of

two components: *Tg(draculin:creER^{T2})*, which allows hematopoietic-selective expression

of a tamoxifen-inducible Cre recombinase, and the *Tg(ubi:loxP-GFP-loxP-stop-mCherry),*

which allows for ubiquitous expression of fluorescent proteins[22,23]. Upon addition of

tamoxifen, Cre recombinase excises the loxP-GFP-loxP cassette, allowing for the

expression of mCherry. The result is permanent mCherry labeling of *draculin+* cells,

including HSPCs and their differentiated progeny. Previously, we determined that varying

the window of tamoxifen exposure during embryonic development helped to resolve

HSCs from the HSC-independent progenitors[14]. Here, tamoxifen-mediated Cre induction

of mCherry labelling is begun at either 1 dpf, 2 dpf, or 3 dpf, with hydroxytamoxifen added

for 20 hours (**Figure 1A**). We refer to the labeling at 1 dpf as "early-HSPCs", at 2 dpf as

"mid-HSPCs" and at 3 dpf as "late-HSPCs". These time points were chosen as they

represent early, middle, and late time points during HSPC specification and expansion

during zebrafish development[1]. Based on our prior study, we hypothesized that early-

HSPCs would be enriched for HSC-independent progenitors and mid/late-HSPCs would

be enriched for long-lived HSCs[14]. To identify what cells are labeled at early, mid, and

late induction time points, we performed scRNA-seq on mCherry-expressing cells isolated

from larvae 1-day-post labelling, so at 2, 3, and 4 dpf, respectively. To assess the

subsequent differentiated hematopoietic progeny, mCherry-expressing cells were

sequenced from 6 and 10 dpf larvae (**Figure 1A**). After filtering out low quality cells based

on standard quality control parameters in the field[24], we clustered cells in Seurat[25]. In

total, about 75% of mCherry-expressing cells were identified as endothelial or

hematopoietic, while the remaining 25% were non-hematopoietic based on top marker

gene expression and thus were filtered out (**Supplemental Figure 1**). As a result, 25

hematopoietic and immune clusters were detected across all timepoints and traces,

including HSCs, progenitors, and other cells of the major mature hematopoietic lineages,

including lymphoid, myeloid, and erythroid cells (**Figure 1B-C**, **Supplemental Figure 2,**

Tables S1). To facilitate ease of access for the community, this full dataset and all others

described below are available to search via a shinyapp webtool made with ShinyCell[26] at

https://sites.google.com/view/tvbowmanlab/developmental-hematopoiesis-atlas.

The HSPC clusters were identified based on the expression of *myb*, which is a

marker gene commonly used to assess HSPCs by *in situ* hybridization[10,27]. Expression of

myb across all time points was localized to the four stem and progenitor clusters identified

as Myeloid Progenitors (MyPs), Erythroid Progenitors (EryPs), Lymphoid Progenitors

(LyPs), and HSCs (**Figure 2A-B**). Differential expression analysis comparing HSCs to

EryPs, MyPs and LyPs showed distinct transcriptomes across the four clusters (**Figure 2C, Table S2**). Consistent with the lineage-enriched expression, EryPs express higher levels of erythroid genes such as *klf17, alas2*, and *hemgn*, the MyP cluster express higher levels of genes critical for the myeloid lineage such as *csf2rb* and *mmp9*, and the LyP cluster expresses higher levels of genes associated with lymphopoiesis such as *runx3, tcf7,* and *ccr9b* (**Table S2**). HSCs were enriched for genes encoding stem cell transcription factors like *runx1*, *drl*, *tal1,* and *blf*. The abundance of each HSPC cluster increased over developmental time, consistent with reported expansion of *myb*-expressing HSPCs[10] (**Figure 2D**). To identify transcriptional changes occurring in HSPCs as they mature during development, we did gene enrichment analysis comparing each cluster over developmental time (**Figure 2E, Table S2**). We observed that the HSC transcriptome was enriched for genes involved in RNA splicing and metabolism at 1-day-post labeling and 10 dpf, with elevated expression of genes involved with hypoxia at 6 dpf. In contrast, LyPs and MyPs showed more similar changes with elevated expression of genes involved with protein translation at 1-day-post labeling and protein folding at 10 dpf. These observations are in line with the rapid growth and protein production typically associated with hematopoietic progenitors[28].

To visualize the early-, mid-, and late-HSPCs *in vivo*, we did confocal imaging of the larval zebrafish HSPC niche termed the Caudal Hematopoietic Tissue (CHT) at 1-day-post labelling (**Figure 2F**). The CHT contained the highest number of mCherry-labeled cells in the early-HSPCs trace and the lowest number in the late-HSPCs trace (**Figure 2F**). To identify if there were differences in the composition of HSPCs labeled by each trace, we analyzed the HSPC subsets at 2, 3, and 4 dpf in the scRNA-seq dataset (**Figure 2G**). Although all four populations, MyPs, LyPs, EryPs and HSCs, were detected in the early-, mid-, and late-traces, the frequency of the populations was distinct. LyPs were most abundant in the early-HSPC trace, HSCs were enriched in the mid-HSPC trace, while EryPs were most frequent in the late-HSPC trace. These data show different distributions of HSPCs across the early-, mid-, and late-HSPC pools during zebrafish development.

HSPCs differentiate into a large repertoire of immune and hematopoietic cell types during development.

By 6 and 10 dpf, HSPCs differentiate into various immune and hematopoietic lineages. Consistent with our prior study[14], there is a substantial contribution of the early-HSPCs to the CHT and thymus at 6 dpf, with minimal contributions from mid- and late-HSPC traces, as observed by confocal imaging (**Figure 3A,A'**). The precise identity of the hematopoietic cells arising in the hematopoietic tissues was unclear, so we leveraged the scRNA-seq data to identify the immune cells derived from early-, mid-, and late-HSPC lineage traces. Lymphoid and myeloid lineage clusters were subset and analyzed by developmental timepoint (**Figure 3B**). When examined as all lymphoid clusters in aggregate, a burst of differentiation from the early-HSPCs, and a concomitant decrease in the proportion of LyPs, potentially due to ongoing differentiation, was observed (**Figure 3B-C**). Compared to mid- and late-HSPCs, the early-HSPCs showed a stronger propensity for differentiation into lymphoid cluster 3 identified as ILC3/il2rb+Tcells. This observation parallels confocal imaging, where early-HSPCs contributed significantly more to the thymus than mid- or late-HSPCs (**Figure 3A**). Not all lymphoid clusters behaved the same. For example, cluster 13 rag+ T cells arose at similar frequency from early- and mid-HSPCs, while late-HSPCs did not generate rag+ T cells by 10 dpf (**Figure 3B-C**).

In contrast to the "burst" of lymphoid production, analysis of all myeloid lineage clusters in aggregate showed a steady level of differentiation from all three early-, mid-, and late-HSPCs traces (**Figure 3B-C'**). However, the production of myeloid lineage cells by the three HSPC pools was not entirely homogenous. Late-HSPCs showed a bias toward making more MyPs at 6 and 10 dpf compared to early- or mid-HSPCs, suggesting that hematopoiesis maybe skewed toward myeloid lineage differentiation during later larval development. Furthermore, macrophages and Antigen Presenting Cells (APCs) showed different enrichments among the three traces (**Figure 3C**).

To determine if the immune cells derived from early-, mid- and late-HSPCs are different, we did differential gene expression and functional enrichment analyses. The 10 dpf macrophage and neutrophil clusters were compared and showed significant differences across traces (**Figure 3D,D', Table S3**). Macrophages derived from early-

and mid-HSPCs expressed genes involved in differentiation and NFkB-signaling, while
those from late-HSPCs showed an upregulation of genes in chemotaxis GO terms
(**Figure 3D'**). Similarly, neutrophils derived from early- and mid-HSPCs upregulated
oxygen transport and NFkB-signaling, while those from late-HSPCs upregulated actin
polymerization. To further identify genes that may be driving differentiation of
macrophages and neutrophils from the early-, mid- and late-HSCPs, we did pseudotime
analysis using psupertime, a method that takes into account the real biological timepoints
in pseudotime inference (**Supplemental Figure 3**)[29]. The analysis found that the cytokine
il1b appeared to be a driver of both macrophage and neutrophils differentiation from early-
HSPCs (**Supplemental Figure 3A,D**). The finding further indicates that macrophages
and neutrophils arising from early-HSPCs are different than those arising from late-
HSPCs.

In summary, our data uncovers how early-, mid- and late-HSPCs differentiate to
the lymphoid and myeloid lineages during larval development. These data suggest early-
HSPCs, which are more enriched for LyP compared to MyP, show higher propensity
towards the lymphoid lineage. This finding reveals layered ontogeny of lymphoid lineage
cells in zebrafish that is in line with other lineage tracing studies[15,16]. While all three HSPC
pools differentiate into macrophages and neutrophils, the gene expression analyses
suggest there are differences in function depending on ontogeny. The findings challenge
current models that infer myelopoiesis dominates earlier HSPC differentiation with
lymphopoietic differentiation potential arising later during developmental hematopoiesis.
203

HSPCs differentiate to ILC-like and Natural Killer-like cells.
In developing zebrafish, immature rag+ T-cells are the best-defined
lymphocytes[30,31]. However, mice and humans are known to have a larger repertoire of
lymphoid cells emerging during embryogenesis and fetal life. For instance, the
developmental formation and maturation of NK cells, ILC2s and ILC3s, as well as B1 and
B2 B-cell progenitors were studied in mouse models[32-34]. Non-T lymphocytes like NK cells
and ILCs were detected in juvenile and adult zebrafish but are not characterized in larval
developmental stages[24,35]. Our lineage tracing and scRNA-seq approaches allowed for a

more agnostic assessment of hematopoietic cells found throughout the entire larvae and

permitted enrichment and detection of a wider diversity of lymphoid cells than currently

described in larval zebrafish publications. Consistent with previous zebrafish work[31], we

detected rag+ T lymphocytes expressing *lck, cd4-1, cd8b, tcra,* and *tcrb* in cluster 9 of the

lymphoid lineage subset of cells in the scRNA-seq data (**Figure 4A-B, Table S4**). Other

less defined populations in larval zebrafish expressed top marker genes associated with

mammalian ILCs and NK cells. Cluster 0 expresses a mix of ILC3-like markers such as

il7r, il23r, il17ra1a and *tnfa*, as well as T-cell marker genes like *tcra* and *lck*. Cluster 2

expresses *eomesa*, *nkl.1, tbx21 (t-bet)*, and *gzm3.3* reminiscent of NK/ILC1-like cells, and

given the low abundance of mature markers of NK-cells like lysins and perforin, the cluster

may represent an immature state of NK/ILC1-like cells. Cluster 3 expresses *il13, il4,*

gata3, mafa (MAF) consistent with an ILC2-like identity. Differential expression analysis

between clusters 0, 2, and 3, representative of the three groups of ILCs, further highlights

their unique transcriptional identities (**Figure 4C, Table S4**).

Aside from ILC-like identity, the lymphoid subset contains a variety of T-cells,

progenitors, and other previously uncharacterized immune cells in larval zebrafish

(**Figure 4A-B**). Cluster 4 appears to be immature T cells, expressing *tcf7, mxd, cd28,* and

zap70. Cluster 8 expresses *tcrg, trdc,* indicating a possible presence of gamma-delta T-

cells. Cluster 7 highly expresses *gata3,* suggesting these could be ILC progenitors. The

putative identity of other clusters is less clear. For example, cluster 6 expresses *blnk*,

ccl36.1, and *il12rb2* possibly indicating presence of a blnk+ B-cell progenitor although

classic markers such as *pax5* are not detected. Cluster 5 expresses genes like *il12bb,*

tox, cd59, and *rorb*, that are expressed in many immune cell types making their identity

less clear thus we refer to them as cd59+ Immune Cells.

The various ILC types have prototypic tissue locations. For example, ILC2s can be

found residing in mucosal tissues like lungs and intestine and in the skin[36]. To determine

if larval zebrafish ILC-like cells are in tissues that align with their hypothesized identity,

we did whole mount *in situ* hybridization for top marker genes of ILC-like cells in 6 and 8

dpf larvae. We selected *eomesa* for the NK/ILC1-like cluster and *il7r* for the

ILC3/il2rb+Tcells. To detect ILC2-like cells, we selected the gene *si:dkey21e2.16*, which

is a predicted <u>s</u>erine-<u>t</u>ype endopeptidase, and named it *swift* (**Figure 4D**). The expression

of *eomesa* puncta is detectable around the thymic area, consistent with thymic NK cell

localization[24] (**Figure 4E**). The expression of *swift* is detectable at 6 & 8 dpf along the

pharyngeal arches and intestines, which is consistent with mucosal localization of

mammalian ILC2s (**Figure 4E**). Some *swift* expressing cells were detectable in the CHT

(**Supplemental Figure 3G**). Lastly, *il7r* expression was localized to the thymus at 6 & 8

dpf, but some puncta were present outside the thymus, along the pharyngeal arches. The

two different locations are consistent with *il7r* marking multiple pots of cells, possibly

ILC3s and il2rb+Tcells. These data illustrate a previously unappreciated variety of

lymphoid lineage immune cells that are detectable in hematopoietic and non-

hematopoietic zebrafish tissues similar with their mammalian counterparts.

Zebrafish ILC-like cells require the function of Runx1.

Given the limitations of identifying novel cells by scRNA-seq alone, we functionally

tested if larval ILC-like cells behave consistently with their predicted identity. To determine

if ILC-like cells in zebrafish parallel the development of mammalian ILCs, we assessed

them in several mutant contexts.

Mammalian ILCs can emerge from 'definitive wave' embryonic progenitors that

pass thorough the fetal liver and genes like *Il7*, *Gata3*, *Tbx21* and *Eomes* were shown to

control ILC embryonic development[37]. However, it's unclear if the genetic program

regulating HSPC emergence is also needed for subsequent ILC development. We sought

to determine if Runx1 is required for ILC-like cells during embryogenesis. Runx1 is a

transcription factor essential for the functionality of some cells of the primitive wave and

for the generation of the entire definitive wave including both pro-definitive and definitive

phases[38-40]. Animals lacking Runx1 have a paucity of definitive HSPCs, erythrocytes,

macrophages, neutrophils, and T lymphocytes in both mice and zebrafish[41-43]. To

determine if ILC-like cells require the function of Runx1, we performed hematopoietic

lineage tracing in the *runx1^{W84X}* zebrafish mutants[41]. Heterozygous *runx1^{W84X}* fish

containing either *Tg(drl:creER^{T2})* or *Tg(ubi:loxP-GFP-loxP-stop-mCherry)* were crossed

to obtain *runx1* wild-type, heterozygous, and homozygous mutant embryos containing

both the tamoxifen-inducible Cre enzyme and the switch cassette
Tg(drl:creER^(T2),ubi:Switch) (**Figure 5A**). Given that the largest proportion of lymphoid
lineage cells were detected in the early-HSPC cohort, we did confocal imaging and
scRNA-seq using the early-HSPCs labeling time point with final analysis at 6 dpf.

By confocal imaging, early-HSPC-derived mCherry-expressing cells were
significantly reduced in hematopoietic sites like the thymus and caudal hematopoietic
tissue, as well as along the intestine in the *runx1^(W84X)* homozygous mutants (**Figure 5B**).
Loss of thymic signal suggested that lymphoid lineage cells were depleted in mutants,
and loss of signal in the caudal hematopoietic tissue suggested a depletion of other
lineages like erythroid, myeloid, and HSPCs.

To determine which specific hematopoietic and immune cells are dependent on
Runx1, we performed scRNA-seq on mCherry+ cells isolated from *runx1^(W84X)*
homozygous mutants and their siblings at 6 dpf (**Figure 5C, Supplemental Figure 4A-**
E, Table S5). To identify the homozygous mutants for sequencing, larvae were scored
for a loss of thymic mCherry signal at 6 dpf by imaging. We selected *runx1^(W84X)*
homozygous mutants and unaffected siblings [mix of wildtype (+/+) and heterozygous (+/-
)] based on this metric and confirmed the genotype of single cell suspensions using a
SNP-based genotyping strategy. The scRNA-seq data revealed that *runx1^(W84X)*
homozygous mutants lack the following lineages: HSPCs, neutrophils and macrophages,
and all the lymphoid lineage cells including rag+ T cells, and ILC-like clusters (**Figure 5C-**
D, Supplemental Figure 4E). As anticipated, the clusters identified in a wild-type control
dataset from prior experiment were present in the sibling controls (**Figure 5C**). Despite
the depletion of most hematopoietic cells in *runx1* homozygous mutants, some
erythrocytes and thrombocytes were still detectable in 6 dpf mutants. Erythrocytes are
longer lived cells that can arise from both primitive and definitive hematopoietic waves,
and prior work in murine models also suggest that megakaryopoiesis can occur from
primitive wave hematopoietic progenitors[44,45]. It is likely that the primitive erythrocytes and
thrombocytes are directly labeled in the early-HSPC trace due to the expression of
Tg(drl:creER^(T2)), and appear Runx1-independent[14]. Whole mount *in situ* RNA
hybridization for *eomesa, swift,* and *il7r* in *runx1^(W84X)* mutants corroborated the significant

reduction of NK/ILC1-like, ILC2-like, and ILC3/il2rb+Tcells in the homozygous *runx1^W84X*

mutants at 6 dpf (**Figure 5E, Supplemental Figure 4F**). These findings show the Runx1

dependency of the newly identified lymphocytes including ILC-like cells.

Zebrafish ILC-like cells show similar dependencies on Il2rγ and Rag1 as
mammalian ILCs.

ILCs and adaptive T-lymphocytes, specifically *cd4+* T-helper cells, express similar

genes[18]. Therefore, identification of ILCs solely based on gene expression is inadequate

to distinguish ILCs from T-helper cells. ILCs and T-lymphocytes show some common and

unique genetic dependencies. In humans and in mice, both ILCs and T-lymphocytes

depend on the function of interleukin 2 common gamma chain (IL2Rγ) because the

immune cells express various interleukins that require IL2RG as a co-receptor[46]. Unlike

T-lymphocytes, ILCs lack antigen-specific receptors, and therefore do not require the

function of Rag1 and Rag2 recombinase enzymes for their development[19]. To better

parse out ILCs from T-cells in larval zebrafish, we assessed the cells in *il2rga^Y91fs*

homozygous mutants[47] and *rag1* mutagenized zebrafish. We anticipated that zebrafish

ILC-like cells would be diminished in an *il2rga* mutant and unaffected by mutagenesis of

rag1.

In zebrafish, Il2rγ is encoded by the duplicated genes *il2rga* and *il2rgb*, however

only *il2rga* functions during developmental lymphopoiesis[48]. We used the *il2rga^Y91fs*

zebrafish mutants to determine if larval ILCs require the function of Il2rγ. Heterozygous

il2rga^Y91fs fish containing either *Tg(drl:creER^T2)* or *Tg(ubi:loxP-GFP-loxP-stop-mCherry)*

were crossed to obtain *il2rga* wild-type, heterozygous, and homozygous embryos with

both the tamoxifen-inducible Cre enzyme and the switch cassette (**Figure 6A**). Early-

HSPCs were labeled at 1 dpf and the mCherry expressing progeny were analyzed at 8

dpf using confocal imaging. In homozygous *il2rga^Y91fs* mutants, mCherry-expressing cells

were significantly reduced in the thymus and intestine compared with the wild-type and

heterozygous siblings (**Figure 6B**). However, mCherry-expressing cells in the CHT were

unaffected by the loss of *il2rga* consistent with undetectable expression of ILC1 and ILC3

markers (data not shown) and minimal ILC2 marker expression in this hematopoietic

tissue (**Supplemental Figure 4F**). To further parse out if ILC-like cells are dependent on *il2rga*, we did *in situ* hybridization for *eomesa, swift,* and *il7r* at 6 dpf and showed reduced expression of all three markers in *il2rga^Y91fs^* homozygous mutants compared to wild-type and heterozygous siblings (**Figure 6C, Supplemental Figure 5A**).

To distinguish ILCs from T-lymphocytes, we mutagenized the *rag1* locus using CRISPR/Cas9 in *Tg(drl:creER^T2^,ubi:Switch)* zebrafish larvae and then assessed T-lymphocytes and ILCs **(Supplemental Figure 5B-D)***.* To control for off-target effects from mutagenesis, we compared *rag1* crispants to larvae mutagenized at the *slc24a5 golden* locus, a non-essential gene which is required for melanin pigmentation in zebrafish[49]. In *rag1* crispants, T-cells were decreased when assessed by imaging of the T-cell fluorescent reporter *Tg(rag2:mCherry)* and by *in situ* hybridization for *rag1* and *tcra*, functionally confirming robust *rag1* mutagenesis (**Supplemental Figure 5C-D**).

To assess all lymphopoiesis in *rag1* and control *slc24a5* crispants, we performed scRNA-seq on the early-HSPC trace-derived mCherry-expressing cells isolated from 8 dpf larvae (**Figure 6A**). We hypothesized that Rag1-independent cells would show no change in abundance in the *rag1* crispants, while Rag1-dependent cells like T-lymphocytes would be diminished. As anticipated, cluster 8 identified as *rag*-expressing T-lymphocytes were diminished in *rag1* crispants (**Figure 6D,E, Table S6**). The T-cell identity of the newly discovered il2rb+ T cells was confirmed by their decreased abundance in *rag1* crispants. Based on gene expression, clusters 1 and 7 appear to be the ILC3s and T-cells observed as cluster 0 in the original HSPC trace dataset (**Figure 4A**), but possibly due to the *rag1* genetic perturbation, the two cell types separated in this dataset (**Figure 6D**). Clusters 2, 6, 7, 9, and 11 were identified as ILC-like based on top marker gene expression, and were still present in high abundance in *rag1* crispants in accordance with ILCs not requiring Rag enzymes for their development (**Figure 6D,E, Supplemental Figure 5E,F**). ILC2-like cells are represented by two clusters in this dataset, clusters 6 and 9. Concordant with a recent mammalian study[50], we noted an increase in Cluster 9, which has more *il13*-expressing ILC2s than in the control. Clusters 3 and 4 identified as cd59+ and blnk+ immune cells, respectively, are also Rag1-independent, indicating that they are likely not T-lymphocytes but rather another type of

specialized immune cells. *In situ* hybridization for markers of ILC1/NK-like and ILC2-like

cells, *eomesa*, and *swift* respectively, confirmed Rag1 independence (**Figure 6F,**

Supplemental Figure 5G). However, expression of *il7r* by *in situ* decreased in *rag1*

crispants, indicating that the cells marked by *il7r* include T-lymphocytes consistent with

the expression of *il7r* in both il2rb+ Tcell cluster 1 and ILC3-like cluster 7 (**Table S6**). In

summary, transcriptionally detected ILC-like cells in zebrafish show genetic

dependencies consistent with their proposed identity, and in agreement with mammalian

ILCs.

Viral mimicry stimulates ILC-like expansion in zebrafish larvae.

Innate lymphocytes participate in the immune system response to pathogenic

insults by viruses, bacteria, and helminths[21]. We sought to test the functionality of larval

ILC-like cells as well as other immune cells by simulating a viral infection in zebrafish

larvae. In the scRNA-seq datasets, the ILC1/NK-like cluster expresses TLR receptor 8

(*tlr8b*) which is a receptor known to selectively sense single-stranded RNA (ssRNA)

viruses like SARS-CoV2 and influenza[51] (**Supplemental Figure 6E**). To mimic a ssRNA

viral infection, we chose the TLR7/8 agonist R848 (Resiquimod) because studies showed

adult mammalian NK-cells respond to this stimulant[52]. Furthermore, R848 can induce a

strong immune response by prompting activation of macrophages[53,54], however the effect

of R848 agonist on the larval lymphoid lineage in zebrafish is unknown.

First, we sought to test the effect of R848 on larval immune cells broadly using

imaging. Early-HSPCs were labeled with mCherry and then larvae were treated with 28.5

μM R848 at 6 dpf and subjected to confocal imaging at 8 dpf. (**Figure 7A**). The treatment

window of 6 – 8 dpf was selected to capture the potentially more mature ILCs as they had

higher abundance in the scRNA-seq at the older timepoints. At 8 dpf, the R848-treated

larvae had a significant increase in mCherry signal in the thymus and CHT as well as an

increased number of puncta in the intestine as compared to untreated controls (**Figure**

7B).

Next, to determine which immune cells and ILC types respond to the viral mimic,

we did scRNA-seq on early-HSPC-derived mCherry-expressing cells from 8 dpf *slc24a5*

crispant controls and *rag1* crispants treated with R848 (**Supplemental Figure 6A-D**). As

anticipated, there is a significant expansion of macrophage clusters 5, 9, and 18 in R848-

treated larvae at 8 dpf (**Figure 7C-D, Supplemental Figure 7B**). Gene enrichment

analysis revealed R848 induced transcriptional changes in these clusters that are

concordant with metabolic changes and defense response activation (**Supplemental**

Figure 7A, Table S6). Based on expression of *tlr8b*, we expected that R848 would elicit

a response from the ILC1/NK-like cluster (**Supplemental Figure 6E,F**). Indeed, there is

an expansion of cluster 12 identified as ILC1/NK-like (**Figure 7C-D, Supplemental**

Figure 7B) in both *slc24a5* controls and in *rag1* crispants treated with R848. Gene

enrichment analysis of differentially expressed genes in ILC1/NK-like cells between

control and R848 treated larvae showed an upregulation of actin polymerization and

motility GO terms, consistent with a study showing increased migration of NK cells in

responses to R848[55](**Figure 7E, Table S6**). We also observed an expansion of cluster 6

il2rb+ T cells, in all R848-treated samples. Lastly, cluster 21 identified as ILC2-like was

diminished in R848-treated larvae, with about a 3-fold decrease in the number of cells

between controls (153 cells) and R848 treated (45 cells). Our observation is in agreement

with a study showing suppression of ILC2s in response to R848[56](**Figure 7C,D**).

To confirm the cellular effects of R848 as observed by scRNA-seq, we did *in situ*

hybridization in whole larvae at 8 dpf for ILC-like marker genes, *eomesa*, *swift*, and *il7r*.

(**Figure 7F**). Expression of *eomesa* was significantly increased by R848 treatment,

including along the pharyngeal arches, suggesting that ILC1/NK-like cells are responsive

during larval development. The expression of ILC2 marker gene *swift* was unchanged

between controls and R848 treated, counter to what was observed in scRNA-seq (**Figure**

7F, Supplemental Figure 7C). Lastly, the expression of *il7r* in the thymus was

significantly increased in response to R848 agonist (**Figure 7F**). The *in situ* hybridization

result supports the observed expansion of cluster 6 il2rb+Tcells in the sequencing data.

Together, our findings indicate that larval ILC-like cells in zebrafish have the same

dependencies as established in murine and human studies. Larval zebrafish ILCs require

Il2rγ but not Rag1 and are responsive to a viral-like immune challenge. Identification of

the various immune system populations equivalent to mammalian systems shows that

larval zebrafish represent a valid and tractable model for addressing broader questions

about immunity and disease.

424
425 **DISCUSSION**
426

In this paper, we show that HSPCs arising at different developmental windows

have distinctive kinetics of differentiation to the lymphoid and myeloid lineages. Using

scRNA-seq, we uncovered embryonic progenitors that have myeloid, lymphoid, or

erythroid biases. The broad characterization of cells made by HSPCs during development

revealed the existence of a wider variety of immune cells in larvae. We also observed

numerous previously unknown lymphoid populations including ILCs. Zebrafish ILCs have

similar genetic dependencies as mammalian counterparts, and larval ILCs can respond

to a viral mimic by at least 6 dpf. The scRNA-seq datasets are available via an online

search tool for the zebrafish, hematology, and immunology communities to facilitate

further study of the various immune cells identified.

The layered nature of developmental hematopoiesis makes it challenging to

examine the distinct qualities of each wave of stem and progenitor cells as well as their

progeny. The early, mid and late HSPC lineage tracing results help illuminate these

features and uncovered distinct differentiation dynamics. Early HSPCs showed a more

rapid differentiation into lymphoid cells, especially il2rb+ T cells and ILC3s. Meanwhile

late-HSPCs differentiate at a slower pace and with a more limited collection of lymphoid

cells compared to early and mid HSPCs. Murine studies are in line with our finding, where

an early wave of ILC3s arising from the hemogenic endothelium and seed the nascent

thymus during embryogenesis[57,58]. From the myelopoietic perspective, all three HSPC

traces appeared to produce myeloid cells at a similar steady rate. It could be interpreted

that the early and late wave myeloid cells are derived from primitive and definitive waves,

respectively. However, early HSPC-derived myeloid cells were depleted in *runx1[W84X]*

homozygous mutants, which do not display decreases in primitive myeloid cells[41]. This

finding indicates the myeloid cells detected in the early HSPC trace are likely of definitive

origin implying the difference between early and late HSPC derived cells is something

new. The increase in MyPs in the late-HSPCs trace suggests distinctive definitive myeloid

differentiation across zebrafish larval development. Further supporting this model is the

observation that the myeloid cells derived from early *versus* late HSPCs have distinct

gene expression profiles. The late-HSPC-derived macrophages and neutrophils showed

different gene enrichment from those derived from early and mid HSPCs in concordance

with other work in the field indicating origin informs functional properties[59]. Future

assessments of early, mid, and late HSPC-derived myeloid progeny across diverse

hematopoietic and immune organs in adulthood could help elucidate how different

embryonic waves shape the adult immune and hematopoietic systems.

Prior to this study, the array of cells routinely examined in 1-5 dpf larvae using

fluorescent reporter lines and *in situ* RNA hybridization were HSPCs, neutrophils,

macrophages, erythrocytes, thrombocytes, rag+ T cells, and most recently eosinophils[60].

Others have detected gene expression in larvae characteristic of NK and ILC cells via

bulk methods like RT-PCR[61], but advancement in sequencing, combined with the

unbiased lineage tracing approach we took, allowed for detection of numerous new cell

types with improved precision and development of new RNA probes for cell type detection

in situ.

Two other scRNA-seq studies identified ILCs in the intestines and thymus of adult

zebrafish[24,35,62]. The top marker genes of intestinal ILC3s align with those detected at

larval stages, including *nitr* genes, *tnfa*, and *il17a/f3*[35,62]. Thymic ILC2s detected in

juveniles also express similar marker genes to those we observed in larvae including *il13*,

il4, *gata3*, and *si:dkey-21e2.10*[24]. The marker gene *eomesa* of the ILC1/NK-like

population was also detected in adult thymus NK-like cells using scRNA-seq[24]. There

remains some ambiguity on the absolute identity of some of cell types we identified as

the cells within some clusters have mixed identities. For example, in our assessment of

wildtype larvae, ILC3-like cells are found intermixed with il2rb+Tcells, but upon loss of

rag1, the populations become distinct. Similarly, ILC2-like cells are intermingled in a

single cluster in unperturbed conditions but become separated into two populations upon

loss of *rag1* and R848 treatment. Long-term lineage tracing studies are needed to clarify

how the newly identified larval populations separate and distribute among adult organ

system.

Much remains unknown about the regulation and function of the larval ILCs. Many

of the top marker genes in ILC populations remain uncharacterized. For example, the

ILC2 gene *swift* (aka *si:dkey-21e2.16*) is unannotated but has homology with granzyme

and chymase, which are more characteristic of cytotoxic immune cells like mast cells. A

recent study in humans showed that ILC2s express granzyme[63], indicating that ILC2s

may have more cytotoxic functions than previously appreciated. This finding hints at how

the unknown genes identified in the new immune effector cell clusters could reveal novel

biology of these lymphocytes. Moreover, the highly expressed and cell-type-restricted

marker genes for ILCs can be exploited to generate new transgenic lines to visualize how

larval and adult ILC populations shape organogenesis, immune responses, and diseases.

We also illustrated that larval ILCs have genetic dependencies consistent with their

innate lymphoid identity. We showed that ILCs expressing *eomesa, swift,* and *il7r* depend

on the functions of Il2rga and are independent of Rag1, consistent with murine studies[46].

Beyond ILCs, we uncovered novel Rag1-independent cells characterized by *blnk* and

cd59 expression although their mammalian counterparts are unclear. To our knowledge,

this study is the first demonstration of the requirement for Runx1 for ILC development. As

we performed whole body loss-of-function studies, it is unclear if Runx1 is required

specifically within the stem and progenitor cells giving rise to the newly described

lymphoid populations or intrinsically within the mature cells. Factors like Runx1 show dual

requirements during the genesis of the definitive hematopoietic wave and within

differentiated immune effectors such as neutrophils and in megakaryocytes[38,64,65]. More

work is needed to identify if ILCs in zebrafish also require factors like *Eomes*, *Id2, Tcf7,*

Gata3, Tox and *Zbtb16* shown to be essential for development in other models[20]. Further

manipulation of the genes within each ILC cluster can uncover mechanisms controlling

the genesis of ILC lineages as well as the keys to their tissue distribution and functionality.

In general, ILCs participate in the immune responses to various pathogens like

bacteria, viruses, fungi, and helminths. Concordant with prior work, we showed that

ssRNA viral mimicry with the TLR7/8 agonist R848 expanded NK/ILC1-like cells[52]. Our

sequencing results showed a decrease in ILC2-like cells with R848 treatment, however

in situ hybridization did not recapitulate the result. We speculate that a portion of ILC2-

like cells may be dying in response to R848 and therefore are removed during cell sorting

prior to sequencing, while *in situ* hybridization may be capturing ILC2-like cells during the

process of dying. Lastly, R848 agonist also caused an expansion of T lymphocytes

although they appear to lack expression of the targeted Tlr7/8 receptors in our datasets.

It is possible that like in other model systems, interferon secreted by NK-cells recruits the

adaptive T lymphocytes via antigen presenting cells[66]. Further studies are needed to

determine if the mechanistic interplay between the immune cells is conserved in

zebrafish.

In summary, we identified how HSPCs differentiate to myeloid and lymphoid

lineages, including the previously unidentified larval ILCs in zebrafish. Studies on

mammalian adult ILCs are often complicated by the presence of T-helper cells which

closely resemble ILCs, so experiments are often done in a *Rag1/2* mutant context where

T lymphocytes are depleted. The zebrafish larval stage may be a unique time window for

studying ILCs in a non-perturbed, non-mutant context, pending further confirmation that

T-helper cell diversification to Th1, Th2, Th17 occurs later in zebrafish life. The discovery

of a wider variety of immune cells in larvae will permit the use of zebrafish for studying a

broader range of questions in the fields of immunology and disease.

Limitations of the study

We chose to use the *(drl:creER^{T2})* transgene based lineage tracing assay, because it is

established that it labels all cells within the adult kidney marrow[67]. Not all definitive

immune effectors derive from marrow hematopoiesis. Tissue-resident macrophages and

ILCs can seed peripheral organs independent of marrow hematopoiesis. Although we

detected a wide variety of immune effectors including those in mucosal tissues, it is

possible that the *(drl:creER^{T2})* system does not capture all HSPC subsets generating all

hematopoietic cells so we could be under sampling the immune landscape in larval

zebrafish by this labeling method.

Acknowledgements

T.V.B. was supported by grants from the National Institutes of Health (NIH) R01DK121738, R01DK131445, R01DK141169, and the Edward P. Evans Foundation. AN was supported by NIH F31HL167600, NYSTEM award C34874GG (PI Frenette), and a Liang Zhu Memorial Fellowship. BAU was supported by NIH F31HL152562 and T32GM007288 (PI Akabas), KJ was supported by T32GM145438 (PI Query) and R01HL153920 (PI Zheng). Instrumental feedback to the development of this project was provided by Dr. Kira Gritsman, Dr. Ulrich Steidl, and Dr. Meelad Dawlady. We thank the Langenau lab for sharing *il2rga^{Y91fs}* mutant lines generated as part of R24OD016761. We thank the core facilities instrumental for this work that are supported by the Einstein National Cancer Institute's grant number P30CA013330, including: Flow Cytometry Core Facility at Einstein and core members Yu (Joey) Zhang and Ming Liu for assistance with FACS; the Einstein Genomics Core and core member David Reynolds for scRNA library preparation; the Analytical Imaging Facility. We thank Dr. Roshan Sharma for meaningful feedback on scRNA-seq data and Dr. Eliseo Castillo and Dr. Keir Balla for fruitful discussions on innate lymphoid cells. Thank you to Dr. Noura Ghazale for meaningful discussions and technical assistance, and Dr. Kathryn Potts for meaningful review and discussion of the manuscript. Graphics were created in BioRender with copyright https://BioRender.com/rsdc34d and https://BioRender.com/riwx66c

Author Contributions

Conceptualization, TVB, AN, BAU; Methodology, TVB, AN, BAU, KJ, DZ; Investigation, TVB, AN, BAU; Writing—Original Draft, TVB, AN; Writing—Review & Editing, TVB, AN, BAU, KJ, DZ; Funding Acquisition, TVB, AN; Supervision, DZ, TVB.

Declaration of Interests

The authors declare no competing interests.

MATERIALS AND METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER

Antibodies		
Anti-Digoxigenin-AP (Anti-DIG)	Sigma	11093274910
Chemicals, peptides, and recombinant proteins		
(Z)-4-Hydroxytamoxifen	Sigma	H7904
4'6-Diamidino-2-phenlyindole (DAPI)	Sigma	D8417
R848 (Resiquimod)	InvivoGen	tlrl-r848-1
Liberase TM Research Grade	Sigma	5401127001
NBT/BCIP Color Development Substrate	Fisher	PR-S3771
Proteinase K, recombinant, PCR Grade	Sigma	3115887001
T7 Endonuclease I	NEB	M0302L
Alt-R™ S.p. Cas9 Nuclease V3	IDT	1081058
T7 RNA Polymerase	Sigma	10881775001
T3 RNA Polymerase	Sigma	11031163001
DIG RNA Labeling Mix	Sigma	11277073910
Deposited data		
Raw and processed scRNA-seq data	This paper	NCBI Accession: GSE292726
Experimental models: Organisms/strains		
Zebrafish *Tg(drl:CreER^T2^)*	Mosimann et al., 2015[23]	ZDB-TGCONSTRCT-160129-2
Zebrafish *Tg(ubi:loxP-GFP-loxP-stop-mCherry)*	Mosimann et al., 2011[22]	ZDB-ALT-110124-1
Zebrafish *Tg(rag2:mCherry)*	Harrold et al., 2016[68]	ZDB-TGCONSTRCT-160329-2
Zebrafish *runx1^W84X^*	Jin et al., 2009[41]	ZDB-ALT-091019-1
Zebrafish *il2rga^Y91fs^*	Tang et al., 2017[47]	ZDB-ALT-170710-3
Zebrafish AB (wild type)	ZIRC	ZL1
Oligonucleotides		
Alt-R® CRISPR-Cas9 tracrRNA	IDT	1072532
rag1 sgRNA #1: GGACTGCTGCCATAAGGGGA	ZFIN	ZDB-CRISPR-201209-4
rag1 sgRNA #2: AATGATGCCCACATCCCAGG	This paper	NA
rag1 sgRNA #3: TGACCGGCAAGTAACCATCA	This paper	NA
slc24a5 sgRNA: GGACAGACGTGTTTCTCCAG	Avagyan et al., 2021[69]	NA
Primer1 for *il7r* RNA probe: TAATACGACTCACTATAGGGGGCCTTCCTTTCACTTTCAG	This paper	NA
Primer2 for *il7r* RNA probe: CATTAACCCTCACTAAAGGGAACATGGATTAATGTCTATAGCAGAGGA	This paper	NA
SNP probe for *runx1^W84X^* target sequence: GGTTCTGGTACCTTGAAGGCGATGGGCAGGGTCTTGTTGCAGCG[C/T]CAGTGTGTCGGCAGGACGGAGCAGAGGAAGTTCGGGCTGTCGG	Life Technologies	TaqMan® SNP Genotyping Assays
Recombinant DNA		

TCRalpha-L1 plasmid	Danilova et al., 2004[70]	Addgene Plasmid #11262
pTWIST-Kan-HighCopy-v2-eomesa plasmid	This paper	Addgene Plasmid #236391
pTWIST-Kan-HighCopy-v2-swift plasmid	This paper	Addgene Plasmid #236392
pENTR™5'-il7r plasmid	This paper	Addgene Plasmid #236393
Software and algorithms		
Cell Ranger v8.0.1 and v7.0.1	10XGenomics	https://support.10xgenomics.com/
Seurat 5.2	Hao et al., 2024[25]	https://satijalab.org/seurat/
clusterProfiler: enrichGO	Yu et al., 2012[71]	DOI: 10.18129/B9.bioc.clusterProfiler
Image Analysis: Fiji	Schindelin et al., 2012[72]	doi:10.1038/nmeth.2019
Photoshop CC 2022	Adobe	https://www.adobe.com/
Adobe Illustrator CC 2022	Adobe	https://www.adobe.com/
GraphPad Prism version 10.2.3	GraphPad	https://www.graphpad.com/
Other		
Resource website for scRNA-seq atlas	This paper	https://sites.google.com/view/tvbowmanlab/developmental-hematopoiesis-atlas

RESOURSE AVAILABILITY

Lead Contact

Further information and requests for resources, reagents, and other supporting data should be made to the Lead Contact, Teresa Bowman at teresa.bowman@einsteinmed.edu

Materials Availability

Reagents used in this study will be made available upon request. Plasmids for making RNA probes are deposited at Addgene.

Data and Code Availability

The sequencing data is available via the GEO accession number: GSE292726

The processed data can be viewed at:

https://sites.google.com/view/tvbowmanlab/developmental-hematopoiesis-atlas

Codes can be made available upon request.

METHOD DETAILS

Zebrafish Husbandry

Zebrafish were bred and maintained as described previously[73]. All fish were maintained

according to Institutional Animal Care and Use Committee (IACUC)-approved protocols

in accordance with the Albert Einstein College of Medicine research guidelines.

Transgenic and Mutant Lines

For lineage tracing, we used *Tg(drl:creER^{T2})* and the *Tg(ubi:loxP-GFP-loxP-stop-*

mCherry) also known as *Tg(ubi:Switch)*[22,23]. For all experiments, we crossed fish with two

copies of *Tg(drl:creER^{T2},ubi:Siwtch)* transgenes and crossed to WT AB fish so all

experimental animals were hemizygous. We used the published *runx1^{W84X}* mutants and

il2rga^{Y91fs} mutants[41,47]. Genotyping of *runx1^{W84X}* mutants was done using a SNP probe

and qPCR, and *il2rga^{Y91fs}* genotyping was done as described in Tang et.al[47]. To validate

the functional consequences of *rag1* crispants on T-cells, we used the

Tg(rag2:mCherry)[68].

Lineage Tracing Assay

To induce fluorescent-based lineage tracing of HSPCs, embryos containing

Tg(drl:creER^{T2},ubi:Siwtch) were incubated in a 28°C incubator with 12 μM 4-

hydroxytamoxifen for 20 hours, starting at 30 hpf for early HSPCs, at 54 hpf for mid

HSPCs, and at 86 hpf for late HSPCs. Hydroxytamoxifen was washed out three times

with embryo water then larvae were kept in clean dishes until endpoint.

Single Cell RNA Sequencing

Cell isolation and Library Preparation

For the early-HSPCs and mid-HSPCs cohorts, 200-250 larvae were used at one-day-post

labelling, 6 dpf, and 10 dpf. To obtain enough fluorescent cells from the late-HSPCs

derived cohort at the three timepoints 4, 6, and 10 dpf, we processed 1,100, 200, and 170

zebrafish larvae, respectively. For the *runx1^{W84X}* mutant experiment, 14 larvae were

processed for each condition. Homozygous *runx1* mutants were identified prior to

sequencing by scoring for absence of thymic mCherry signal and morphologically

observable thymic involution. Genotype was confirmed post-processing using left over

cells (data not shown). For the *rag1* crispants/R848 experiment, 35 larvae were

processed per condition. For processing of all samples, whole larvae were minced with a

razor blade and digested with 5 mg/mL liberase for 15 minutes at 37°C, followed by FBS

inactivation as previously described[14]. The mCherry-expressing cells were sorted out

using the BD FACSAria II machine into buffer made of 500 μL IMDM + 10% FBS at room

temperature machine setting. The cells were immediately processed for cell capture by

the Einstein genomics facility, using the "Chromium Next GEM Single Cell 3′ Kit v3.1" for

samples in Figures 1-5 and the "Chromium Next GEM Single Cell 3′ Kit v4" kit for *rag1*

crispant/R848 samples in Figures 6 & 7. Cell numbers for each experiment are listed in

Tables S1-7.

Quality Control and Analysis

Fastq files for *rag1* crispant/R848 experiment were processed in Cell Ranger 8.0.1 using

the count function and all other experiments were processed in Cell Ranger 7.0.1. To

generate the counts matrix in Cell Ranger, we made a custom reference genome based

on *Danio rerio* GRCz11 with the addition of mCherry, and T cell specific genes: TCRaC,

TCRbC1 and TCRbC2. All subsequent analysis was done in Seurat version 5.2.0[25]. No

batch effect was apparent between samples, so all analysis was done without batch

corrections **(Figure S1D, S2E).** For all experiments, cells with less than 200 genes and

more than 10% mitochondrial DNA were filtered out **(Figure S1, S2, S4, S5)**. For the

lineage tracing dataset in Figure 1, we used the 4,000 top variable features for initial

dimensional reduction, with 30 principal components (PC) and at cluster resolution of 1.

After non-hematopoietic cells were removed, the remaining hematopoietic cells were re-

analyzed with top 2,000 variable features, 30 PCs and at cluster resolution of 1 **(Figure**

S1A,F). Two more lower quality clusters were identified and removed to obtain the final

clustering using 2,000 variable features, 20 PCs, and cluster resolution of 1. For the

lymphoid only subset analysis in Figure 4A, we used 2,000 variable genes, 25 PCs, and

0.3 cluster resolution. For the *runx1* mutant dataset in Figure 5, we included a wild-type

control dataset at 6 dpf derived from the early-HSPCs trace to check for common

identification of clusters between experiments. For initial clustering we used 4,000

variable genes, 30 PCs, and cluster resolution of 1. We removed non-hematopoietic

clusters based on top marker genes, and the final dataset was generated using 2,000

variable genes, with 20 PCs, and at a cluster resolution of 0.7. For *rag1* crispant/R848

dataset in Figures 6-7, initial clustering was done using 4,000 variable genes, 30 PCs,

and cluster resolution of 1. After removing non-hematopoietic cells, final clustering was

done using 2,000 variable genes, 20 PCs, and a cluster resolution of 1. For analysis of

the lymphoid only subset, we used 1,000 variable genes, 15 PCs, and 0.2 cluster

resolution. Cluster identities were assigned based on literature and top marker genes

obtained using the FindAllMarkers function in Seurat which compares the target cluster

to other clusters using the Wilcoxon Rank Sum test.

Differential Expression and GO Enrichment Analysis

For differential gene expression analysis, two clusters of interest were compared using

the FindMarkers function in Seurat using standard settings for Wilcoxon Rank Sum test.

For gene enrichment analysis, the list of genes obtained from differential expression

analysis was filtered to include only those with adjusted p-value of less that 0.05, and an

average fold-change greater than 1.5. Using the enrichGO function available in

ClusterProfiler package, we identified overrepresented GO terms that belong to the

Biological Processes category. Displayed in the Figures 2E, 3C', and S6D are selected

GO categories from the output list.

Pseudotime Analysis

The software Psupertime[29] was used to temporally order each cluster of cells in the

hematopoietic atlas dataset from the different lineage traces. We include in the

supplement results from psupertime on two clusters of cells identified as macrophages

and neutrophils (Figure S3). Psupertime was run with default settings with the exception

of "min_expression" set to 0.00 instead of 0.01 to include genes that were expressed in

any cell rather than at least 1% to ensure that rare lineage genes were not missed due to

low abundance of some cell types in some traces.

Confocal Microscopy and Analysis

The CSU-W1 Nikon spinning disc confocal was used for all high-resolution imaging of

lineage tracing in wild-type and mutant conditions and in drug-treated larvae. For live

imaging, larvae were anesthetized with 0.01% tricaine and mounted in 4%

methylcellulose on glass bottom dished. Images were acquired using 10x lens with the

following parameters: 600 millisecond exposure, and 561λ laser at 65% power and the z-

stack was set to 6 μm size steps. Images were analyzed in Fiji (ImageJ) version 2.1.0. To

calculate mCherry signal in the thymus and CHT, we used Corrected Total Cell

Fluorescence formular: (Integrated density of region of interest) - (Area of region of

interest x Mean fluorescence of 6 different background regions). All quantification of live

imaging was done blindly to genotype and/or condition, with the exception of images in

Figure 2F and Figure 3A.

***In situ* Hybridization**

The *tcra* RNA probe was made from Addgene plasmid #11262 gifted by Lisa Steiner[70].

The *il7r* RNA probe was made off a PCR template that was amplified from cDNA collected

from 10 dpf whole larvae. Transcription of the anti-sense *il7r* RNA off the PCR template

was done using T3 RNA polymerase. To make *eomesa* and *swift* probes, the DNA

sequences were synthesized and put into pTWIST-kan high copy vector by TWIST

Bioscience. For *eomesa* probe, a 971 bp fragment containing exon 7 - 3'UTR region and

flanked by T7 and T3 primers was synthesized. For *swift* probe, a 922 bp fragment was

made containing the entire coding region of the gene. Gene-specific primers that

contained T3 RNA polymerase sequence (forward) and T7 RNA polymerase sequence
(reverse) at the 5'end were then used to generate a linear PCR-amplified template for *in*
vitro transcription using T7 RNA polymerase. Whole mount *in situ* RNA hybridization was
done based on established zebrafish protocols with modifications for older larvae[74]. For
6 & 8 dpf stages, pigment was removed by bleaching with a solution of 0.8%KOH, 0.9%
H_2O_2, and 0.1% Tween20 for 35 & 45 min, respectively. Proteinase K working
concentration was increased to 15 mg/mL, and digestion time was extended to 35 & 40
min for 6 & 8 dpf, respectively. After digestion, samples were fixed with 0.25%
glutaraldehyde for 25 and 30 minutes for 6 & 8 dpf, respectively. Incubation in RNA probes
was done for two nights at 70°C to maximize detection. To visualize the RNA probe signal,
we used anti-DIG followed by colorimetric detection using NBT/BCIP. Samples were
incubated in NBT/BCIP for 1-3 days for *eomesa* or *il7r*, and 8 hours for *swift,* all at room
temperature. For experiments with *runx1^{W84X}* and *il2rga^{Y91fs}* mutants, each embryo was
imaged, quantified, and then genotyped. All quantification was done blindly to genotype
and condition, except for experiments in Figure 6C and 7E for *eomesa* and *il7r* probes.
Full images of *swift* expression are shown in the supplemental figures.
718

CRISPR/Cas9 Mutagenesis
The crispant approach was used to mutagenize the *rag1* locus, where a single guide RNA
and Cas9 protein were assembled into a complex and injected into one-cell stage
embryos[75]. The single guide RNA was assembled by annealing a tracr RNA with the locus
specific crRNA for 5 min at 95°C, and cooled at room temperature for 10 min. Three
guides were used to target the *rag1* locus, two of them were designed in CHOPCHOP,
and one sequence that is deposited in ZFIN **(Figure S5)**[76]. As a control for mutagenesis,
the *slc26a4* (golden) guide RNA was used[69]. Each single guide RNA was assembled into
a complex with Cas9 separately. For the control, *slc24a5* targeting complex alone was
injected, and for *rag1* crispants, the four guide complexes (3 *rag1* guides + 1 *slc24a5*
guide) were mixed in equal volumes. A T7 endonuclease assay was used to confirm
mutagenesis, where a cutting occurs only in the presence of heteroduplex DNA indicating
mutagenesis **(Figure S5)**.

Resiquimod R848 Drug Treatment

The 500 μg of R848 powder was obtained from InvivoGen and reconstituted in 500 μL of water for a final concentration of 1 μg/μL or 2.85 mM. To treat larvae, the stock was diluted in embryo water to a working concentration of 28.5 μM. Larvae were incubated with R848 starting at 6 dpf and analyzed at 8 dpf.

Statistical Analysis

Analysis was done in GraphPad Prism 10.2.3. For all experiments with 3 groups, a one-way ANOVA with Tukey multiple testing corrections was used. For 2 groups, an unpaired t-test was used as indicated in the figure legends. All error bars show standard deviation from the mean. A Chi-square test was used to compare number of cells per cluster in the stacked bar plots. In all plots, one point is a single zebrafish larva. For single-cell RNA sequencing, one replicate was done per condition. Confocal imaging experiments were done with at least 3 independent experiments, and in situ experiments were done at least 2 independent times. In all figures: ns = not significant, * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, **** = $p < 0.0001$.

REFERENCES

1   Gore, A. V., Pillay, L. M., Venero Galanternik, M. & Weinstein, B. M. The zebrafish: A fintastic model for hematopoietic development and disease. *Wiley Interdiscip Rev Dev Biol* **7**, e312, doi:10.1002/wdev.312 (2018).

2   Kobayashi, M. & Yoshimoto, M. Multiple waves of fetal-derived immune cells constitute adult immune system. *Immunol Rev* **315**, 11-30, doi:10.1111/imr.13192 (2023).

3   Jagannathan-Bogdan, M. & Zon, L. I. Hematopoiesis. *Development* **140**, 2463-2467, doi:10.1242/dev.083147 (2013).

4   McGrath, K. E., Frame, J. M. & Palis, J. Early hematopoiesis and macrophage development. *Semin Immunol* **27**, 379-387, doi:10.1016/j.smim.2016.03.013 (2015).

5   Le Guyader, D. *et al.* Origins and unconventional behavior of neutrophils in
developing zebrafish. *Blood* **111**, 132-141, doi:10.1182/blood-2007-06-095398
(2008).

6   Canu, G. & Ruhrberg, C. First blood: the endothelial origins of hematopoietic
progenitors. *Angiogenesis* **24**, 199-211, doi:10.1007/s10456-021-09783-9 (2021).

7   Dege, C. *et al.* Potently Cytotoxic Natural Killer Cells Initially Emerge from
Erythro-Myeloid Progenitors during Mammalian Development. *Dev Cell* **53**, 229-
239 e227, doi:10.1016/j.devcel.2020.02.016 (2020).

8   Kissa, K. & Herbomel, P. Blood stem cells emerge from aortic endothelium by a
novel type of cell transition. *Nature* **464**, 112-115, doi:10.1038/nature08761
(2010).

9   Bertrand, J. Y. *et al.* Haematopoietic stem cells derive directly from aortic
endothelium during development. *Nature* **464**, 108-111, doi:10.1038/nature08738
(2010).

10   Murayama, E. *et al.* Tracing hematopoietic precursor migration to successive
hematopoietic organs during zebrafish development. *Immunity* **25**, 963-975,
doi:10.1016/j.immuni.2006.10.015 (2006).

11   Chen, M. J. *et al.* Erythroid/myeloid progenitors and hematopoietic stem cells
originate from distinct populations of endothelial cells. *Cell Stem Cell* **9**, 541-552,
doi:10.1016/j.stem.2011.10.003 (2011).

12   Dignum, T. *et al.* Multipotent progenitors and hematopoietic stem cells arise
independently from hemogenic endothelium in the mouse embryo. *Cell Rep* **36**,
109675, doi:10.1016/j.celrep.2021.109675 (2021).

13   Ghersi, J. J. *et al.* Haematopoietic stem and progenitor cell heterogeneity is
inherited from the embryonic endothelium. *Nat Cell Biol* **25**, 1135-1145,
doi:10.1038/s41556-023-01187-9 (2023).

14   Ulloa, B. A. *et al.* Definitive hematopoietic stem cells minimally contribute to
embryonic hematopoiesis. *Cell Rep* **36**, 109703,
doi:10.1016/j.celrep.2021.109703 (2021).

15   Tian, Y. *et al.* The first wave of T lymphopoiesis in zebrafish arises from aorta
endothelium independent of hematopoietic stem cells. *J Exp Med* **214**, 3347-
3360, doi:10.1084/jem.20170488 (2017).

16   He, S. *et al.* In vivo single-cell lineage tracing in zebrafish using high-resolution
796   infrared laser-mediated gene induction microscopy. *Elife* **9**,
797   doi:10.7554/eLife.52024 (2020).

798   17   Neo, W. H., Lie, A. L. M., Fadlullah, M. Z. H. & Lacaud, G. Contributions of
799   Embryonic HSC-Independent Hematopoiesis to Organogenesis and the Adult
800   Hematopoietic System. *Front Cell Dev Biol* **9**, 631699,
801   doi:10.3389/fcell.2021.631699 (2021).

802   18   Cherrier, D. E., Serafini, N. & Di Santo, J. P. Innate Lymphoid Cell Development:
803   A T Cell Perspective. *Immunity* **48**, 1091-1103,
804   doi:10.1016/j.immuni.2018.05.010 (2018).

805   19   Spits, H. *et al.* Innate lymphoid cells--a proposal for uniform nomenclature. *Nat*
806   *Rev Immunol* **13**, 145-149, doi:10.1038/nri3365 (2013).

807   20   Klose, C. S. N. & Artis, D. Innate lymphoid cells control signaling circuits to
808   regulate tissue-specific immunity. *Cell Res* **30**, 475-491, doi:10.1038/s41422-
809   020-0323-8 (2020).

810   21   Yuan, T. *et al.* Innate lymphoid cells and infectious diseases. *Innate Immun* **30**,
811   120-135, doi:10.1177/17534259241287311 (2024).

812   22   Mosimann, C. *et al.* Ubiquitous transgene expression and Cre-based
813   recombination driven by the ubiquitin promoter in zebrafish. *Development* **138**,
814   169-177, doi:10.1242/dev.059345 (2011).

815   23   Mosimann, C. *et al.* Chamber identity programs drive early functional partitioning
816   of the heart. *Nat Commun* **6**, 8146, doi:10.1038/ncomms9146 (2015).

817   24   Rubin, S. A. *et al.* Single-cell analyses reveal early thymic progenitors and pre-B
818   cells in zebrafish. *J Exp Med* **219**, doi:10.1084/jem.20220038 (2022).

819   25   Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-
820   cell analysis. *Nat Biotechnol* **42**, 293-304, doi:10.1038/s41587-023-01767-y
821   (2024).

822   26   Ouyang, J. F., Kamaraj, U. S., Cao, E. Y. & Rackham, O. J. L. ShinyCell: simple
823   and sharable visualization of single-cell gene expression data. *Bioinformatics* **37**,
824   3374-3376, doi:10.1093/bioinformatics/btab209 (2021).

27 Soza-Ried, C., Hess, I., Netuschil, N., Schorpp, M. & Boehm, T. Essential role of
c-myb in definitive hematopoiesis is evolutionarily conserved. *Proc Natl Acad Sci*
U S A **107**, 17304-17308, doi:10.1073/pnas.1004640107 (2010).

28 Gao, X., Xu, C., Asada, N. & Frenette, P. S. The hematopoietic stem cell niche:
from embryo to adult. *Development* **145**, doi:10.1242/dev.139691 (2018).

29 Macnair, W., Gupta, R. & Claassen, M. psupertime: supervised pseudotime
analysis for time-series single-cell RNA-seq data. *Bioinformatics* **38**, i290-i298,
doi:10.1093/bioinformatics/btac227 (2022).

30 Langenau, D. M. *et al.* In vivo tracking of T cell development, ablation, and
engraftment in transgenic zebrafish. *Proc Natl Acad Sci U S A* **101**, 7369-7374,
doi:10.1073/pnas.0402248101 (2004).

31 Langenau, D. M. & Zon, L. I. The zebrafish: a new model of T-cell and thymic
development. *Nat Rev Immunol* **5**, 307-317, doi:10.1038/nri1590 (2005).

32 Dorshkind, K. & Crooks, G. Layered immune system development in mice and
humans. *Immunol Rev* **315**, 5-10, doi:10.1111/imr.13198 (2023).

33 Tang, Y. *et al.* Emergence of NK-cell progenitors and functionally competent NK-
cell lineage subsets in the early mouse embryo. *Blood* **120**, 63-75,
doi:10.1182/blood-2011-02-337980 (2012).

34 Schneider, C. *et al.* Tissue-Resident Group 2 Innate Lymphoid Cells Differentiate
by Layered Ontogeny and In Situ Perinatal Priming. *Immunity* **50**, 1425-1438
e1425, doi:10.1016/j.immuni.2019.04.019 (2019).

35 Hernandez, P. P. *et al.* Single-cell transcriptional analysis reveals ILC-like cells in
zebrafish. *Sci Immunol* **3**, doi:10.1126/sciimmunol.aau5265 (2018).

36 Naito, M. & Kumanogoh, A. Group 2 innate lymphoid cells and their surrounding
environment. *Inflamm Regen* **43**, 21, doi:10.1186/s41232-023-00272-8 (2023).

37 Hernandez-Torres, D. C. & Stehle, C. Embryonic ILC-poiesis across tissues.
Front Immunol **13**, 1040624, doi:10.3389/fimmu.2022.1040624 (2022).

38 Yzaguirre, A. D., de Bruijn, M. F. & Speck, N. A. The Role of Runx1 in Embryonic
Blood Cell Formation. *Adv Exp Med Biol* **962**, 47-64, doi:10.1007/978-981-10-
3233-2_4 (2017).

39   Kalev-Zylinska, M. L. *et al.* Runx1 is required for zebrafish blood and vessel
development and expression of a human RUNX1-CBF2T1 transgene advances a
model for studies of leukemogenesis. *Development* **129**, 2015-2030,
doi:10.1242/dev.129.8.2015 (2002).

40   North, T. E. *et al.* Runx1 expression marks long-term repopulating hematopoietic
stem cells in the midgestation mouse embryo. *Immunity* **16**, 661-672,
doi:10.1016/s1074-7613(02)00296-0 (2002).

41   Jin, H. *et al.* Definitive hematopoietic stem/progenitor cells manifest distinct
differentiation output in the zebrafish VDA and PBI. *Development* **136**, 647-654,
doi:10.1242/dev.029637 (2009).

42   Sasaki, K. *et al.* Absence of fetal liver hematopoiesis in mice deficient in
transcriptional coactivator core binding factor beta. *Proc Natl Acad Sci U S A* **93**,
12359-12363, doi:10.1073/pnas.93.22.12359 (1996).

43   Wang, Q. *et al.* Disruption of the Cbfa2 gene causes necrosis and hemorrhaging
in the central nervous system and blocks definitive hematopoiesis. *Proc Natl
Acad Sci U S A* **93**, 3444-3449, doi:10.1073/pnas.93.8.3444 (1996).

44   Tober, J. *et al.* The megakaryocyte lineage originates from hemangioblast
precursors and is an integral component both of primitive and of definitive
hematopoiesis. *Blood* **109**, 1433-1441, doi:10.1182/blood-2006-06-031898
(2007).

45   Potts, K. S. *et al.* A lineage of diploid platelet-forming cells precedes polyploid
megakaryocyte formation in the mouse embryo. *Blood* **124**, 2725-2729,
doi:10.1182/blood-2014-02-559468 (2014).

46   Colonna, M. Innate Lymphoid Cells: Diversity, Plasticity, and Unique Functions in
Immunity. *Immunity* **48**, 1104-1117, doi:10.1016/j.immuni.2018.05.013 (2018).

47   Tang, Q. *et al.* Dissecting hematopoietic and renal cell heterogeneity in adult
zebrafish at single-cell resolution using RNA sequencing. *J Exp Med* **214**, 2875-
2887, doi:10.1084/jem.20170976 (2017).

48   Sertori, R. *et al.* Conserved IL-2Rgammac Signaling Mediates Lymphopoiesis in
Zebrafish. *J Immunol* **196**, 135-143, doi:10.4049/jimmunol.1403060 (2016).

49   Lamason, R. L. *et al.* SLC24A5, a putative cation exchanger, affects
pigmentation in zebrafish and humans. *Science* **310**, 1782-1786,
doi:10.1126/science.1116238 (2005).

50   Ver Heul, A. M. *et al.* RAG suppresses group 2 innate lymphoid cells. *bioRxiv*,
doi:10.1101/2024.04.23.590767 (2024).

51   Veneziani, I., Alicata, C., Moretta, L. & Maggi, E. Human toll-like receptor 8
(TLR8) in NK cells: Implication for cancer immunotherapy. *Immunol Lett* **261**, 13-
16, doi:10.1016/j.imlet.2023.07.003 (2023).

52   Veneziani, I. *et al.* Toll-like receptor 8 agonists improve NK-cell function primarily
targeting CD56(bright)CD16(-) subset. *J Immunother Cancer* **10**, doi:10.1136/jitc-
2021-003385 (2022).

53   Anfray, C. *et al.* Intratumoral combination therapy with poly(I:C) and resiquimod
synergistically triggers tumor-associated macrophages for effective systemic
antitumoral immunity. *J Immunother Cancer* **9**, doi:10.1136/jitc-2021-002408
(2021).

54   Lopez-Cuevas, P. *et al.* Reprogramming macrophages with R848-loaded artificial
protocells to modulate skin and skeletal wound healing. *J Cell Sci* **137**,
doi:10.1242/jcs.262202 (2024).

55   Zhou, J., Nouri-Shirazi, M., Tang, H., Shen, Y. & Zeng, M. R848/TLR7-Mediated
Stronger CD8+ T Immunity Is Dependent on DC-NK Cell Interactions. *Int Arch
Allergy Immunol* **183**, 860-875, doi:10.1159/000522364 (2022).

56   Okuzumi, S. *et al.* TLR7 Agonist Suppresses Group 2 Innate Lymphoid Cell-
mediated Inflammation via IL-27-Producing Interstitial Macrophages. *Am J Respir
Cell Mol Biol* **65**, 309-318, doi:10.1165/rcmb.2021-0042OC (2021).

57   Elsaid, R. *et al.* A wave of bipotent T/ILC-restricted progenitors shapes the
embryonic thymus microenvironment in a time-dependent manner. *Blood* **137**,
1024-1036, doi:10.1182/blood.2020006779 (2021).

58   Simic, M. *et al.* Distinct Waves from the Hemogenic Endothelium Give Rise to
Layered Lymphoid Tissue Inducer Cell Ontogeny. *Cell Rep* **32**, 108004,
doi:10.1016/j.celrep.2020.108004 (2020).

59   Garcia-Lopez, J. P. *et al.* Ontogenetically distinct neutrophils differ in function
and transcriptional profile in zebrafish. *Nat Commun* **14**, 4942,
doi:10.1038/s41467-023-40662-7 (2023).

60   Li, G. *et al.* Cebp1 and Cebpbeta transcriptional axis controls eosinophilopoiesis
in zebrafish. *Nat Commun* **15**, 811, doi:10.1038/s41467-024-45029-0 (2024).

61   Pereiro, P. *et al.* Zebrafish Nk-lysins: First insights about their cellular and
functional diversification. *Dev Comp Immunol* **51**, 148-159,
doi:10.1016/j.dci.2015.03.009 (2015).

62   Jones, L. O. *et al.* Single-cell resolution of the adult zebrafish intestine under
conventional conditions and in response to an acute Vibrio cholerae infection.
Cell Rep **42**, 113407, doi:10.1016/j.celrep.2023.113407 (2023).

63   Li, Z. *et al.* Therapeutic application of human type 2 innate lymphoid cells via
induction of granzyme B-mediated tumor cell death. *Cell* **187**, 624-641 e623,
doi:10.1016/j.cell.2023.12.015 (2024).

64   Huang, Z. *et al.* Runx1 regulates zebrafish neutrophil maturation via synergistic
interaction with c-Myb. *J Biol Chem* **296**, 100272, doi:10.1016/j.jbc.2021.100272
(2021).

65   Zezulin, A. U. *et al.* RUNX1 is required in granulocyte-monocyte progenitors to
attenuate inflammatory cytokine production by neutrophils. *Genes Dev* **37**, 605-
620, doi:10.1101/gad.350418.123 (2023).

66   Iwasaki, A. & Medzhitov, R. Toll-like receptor control of the adaptive immune
responses. *Nat Immunol* **5**, 987-995, doi:10.1038/ni1112 (2004).

67   Henninger, J. *et al.* Clonal fate mapping quantifies the number of haematopoietic
stem cells that arise during development. *Nat Cell Biol* **19**, 17-27,
doi:10.1038/ncb3444 (2017).

68   Harrold, I. *et al.* Efficient transgenesis mediated by pigmentation rescue in
zebrafish. *Biotechniques* **60**, 13-20, doi:10.2144/000114368 (2016).

69   Avagyan, S. *et al.* Resistance to inflammation underlies enhanced fitness in
clonal hematopoiesis. *Science* **374**, 768-772, doi:10.1126/science.aba9304
(2021).

70   Danilova, N. *et al.* T cells and the thymus in developing zebrafish. *Dev Comp*
Immunol **28**, 755-767, doi:10.1016/j.dci.2003.12.003 (2004).

71   Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for
comparing biological themes among gene clusters. *OMICS* **16**, 284-287,
doi:10.1089/omi.2011.0118 (2012).

72   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis.
Nat Methods **9**, 676-682, doi:10.1038/nmeth.2019 (2012).

73   Lawrence, C. in *Methods in Cell Biology* Vol. 104 (eds H. William Detrich, Monte
Westerfield, & Leonard I. Zon) 429-451 (Academic Press, 2011).

74   Thisse, C. & Thisse, B. High-resolution in situ hybridization to whole-mount
zebrafish embryos. *Nat Protoc* **3**, 59-69, doi:10.1038/nprot.2007.514 (2008).

75   Burger, A. *et al.* Maximizing mutagenesis with solubilized CRISPR-Cas9
ribonucleoprotein complexes. *Development* **143**, 2025-2037,
doi:10.1242/dev.134809 (2016).

76   Labun, K. *et al.* CHOPCHOP v3: expanding the CRISPR web toolbox beyond
genome editing. *Nucleic Acids Res* **47**, W171-W174, doi:10.1093/nar/gkz365
(2019).
962

Figure 1. **Single cell atlas of hematopoietic and immune cells detected during zebrafish development.** (**A**) Hematopoietic-selective and tamoxifen-inducible Cre-Lox labelling system was activated by addition of 12 μM hydroxy-tamoxifen at either 1, 2, or 3 days-post-fertilization (dpf) which we termed the "early-HSPCs", "mid-HSPCs" and "late-HSPCs" respectively (2). The mCherry-expressing progeny of each trace was captured at three developmental timepoints using single cell RNA sequencing (scRNA-seq), at one-day-post labelling (1 dpl), 6 dpf, and at 10 dpf (3). (**B**) UMAP

showing clusters detected at 1 dpl, 6 dpf, and 10 dpf derived from any of the HSPC tracing paradigms. The dotted line highlights all HSPC populations including LyPs, MyPs, EryPs and HSCs, while solid lines highlight mature cells by lineage. (**C**) Dot plot displaying top marker genes of erythrocytes, thrombocytes, myelocytes, lymphocytes, stem & progenitor cells, endothelium, and proliferating cells that were obtained using the FindAllMarkers function in Seurat with Wilcoxon Rank Sum testing.

Figure 2. Comparison of contemporaneous hematopoietic progenitors and HSCs during zebrafish development. (**A**) UMAP showing expression of the gene *myb* across all cells, with enrichment in HSPCs clusters (dotted line). The *myb* expressing population contains proliferating cells representing a mix of identities. (**B**) UMAP of the HSPC subset circled in (A). (**C**) Heatmap of differentially expressed genes of HSCs compared to EryPs (left), MyPs (middle), and LyPs (right). (**D**) UMAP showing the HSPC populations at 1 dpl, 6 dpf, and 10 dpf. (**E**) Gene enrichment analysis of HSCs, LyPs, and MyPs at the developmental timepoints indicated using EnrichGO for Biological Processes. Number in parentheses indicate the number of genes differentially expressed. Comparison was done between two timepoints, and top DEGs with greater than 1.5-fold effect were used for the enrichment analysis. (**F**) Confocal imaging of mCherry signal in the CHT at one-day-post labeling for early-, mid-, and late-HSPC showing highest signal from the early labeling (2 dpf) compared to later. Quantification of the number of mCherry+ cells in the CHT shown on right. (**G**) UMAP showing HSPCs at one-day post labeling (1 dpl, left) and their relative percentages in the bar plot (right).

Figure 3. Lympho-myelo lineage differentiation kinetics vary across HSPC development. (**A**) Confocal imaging of mCherry signal in the CHT (left) and Thymus (right) derived from early, mid, and late HSPCs at 6 dpf and are circled with dotted white lines. (A') Quantification shows highest signal in thymus and CHT derived from early-HSPCs. (**B-B''**) UMAP showing the lymphoid-myeloid subset at 1 dpl, 6 dpf, and 10 dpf from the early (top), mid (middle), or late (bottom) HSPC trace. Dotted lines circle the myeloid lineage cells or lymphoid lineage cells. Solid lines circle the clusters of interest analyzed in B'-B'. In B' (lymphoid) and B'' (myeloid), plots show changes in percentages of cells per cluster in early- (gray), mid- (light green), and late-HSPC (dark green) traces. (**C**) To determine the differences between myeloid progeny derived from early, mid, and late HSPCs, we did differential gene expression followed by GO analyses of cells from 10 dpf larvae. (C') EnrichGO for Biological Processes enrichment analysis of upregulated genes in macrophages and neutrophils from the comparisons shown in C.

Figure 4. HSPCs generate a variety of lymphoid lineage cells during development. (A) UMAP showing lymphoid subset of data from Figure 1B. Circled populations highlight clusters containing the three types of ILC-like cells. **(B)** Dot plot shows selected top marker genes of each cluster generated

using the FindAllMarkers function in Seurat with Wilcoxon Rank Sum testing. (**C**) Heatmap of genes differentially expressed comparing the three ILC clusters to each other. (**D**) UMAP showing expression of *eomesa, si:dkey21e2.16, il7r*, that were selected for *in situ* hybridization testing for the three ILC-like populations. (**E**) Images of *in situ* RNA hybridization for selected marker genes of ILC-like cells at 6 and 8 dpf. Triangles point to puncta and dotted line circles the thymus. Inset shows enlarged image around the thymus.

Figure 5. Zebrafish ILCs depend on the function of Runx1. (**A**) Outline of experiment, where early-HSPC lineage tracing was done in *runx1^W84X^* homozygous mutants and in sibling controls. Early-HSPCs were labeled at 1 dpf by addition of tamoxifen. At 6 dpf, larvae were either imaged or processed for scRNA-seq. (**B**) Confocal images showing mCherry signal derived from early-HSPCs in CHT (left), thymus (middle), and the intestines (right). Signal from early HSPC progeny is reduced in *runx1^W84X^* homozygous mutants in all three locations. Quantification displayed below, where each dot represents the signal for an individual larva. (**C**) UMAP showing hematopoietic and endothelial cells from 6 dpf wild-type controls (early HSPC data from Figure 1), *runx1^W84X^* siblings and homozygous mutants. Lymphoid lineage cells including ILCs are circled. (**D**) Bar plot shows percentage of cells when grouped by major hematopoietic categories. Erythroid includes clusters (7)Thrombocytes and (0,1,2,18)Erythroid. Myeloid category includes (8)Macrophages, (17)APCs, and (11)Neutrophils. Lymphoid category includes all the circled cells on the UMAP, clusters 3,10,12,15,16. HSPCs category includes clusters 4, 5, and 6. The "Other" category includes (13,14)Endothelial and (9)Proliferating cells. (**E**) *In situ* RNA hybridization of *eomesa*, *swift*, and *il7r* in *runx1^W84X^* homozygous mutants. Images (left) are representative of the mean signal. Triangles point to quantified puncta, while dotted line circles the thymus. Quantification (right) shows significant reduction of all three ILC genes. Statistics: in (B) average ± std and p-values are calculated with an unpaired t-test.

Figure 6. Zebrafish ILCs are Il2rga-dependent and Rag1-independent. (**A**) Outline of experiment of early-HSPC lineage tracing in *il2rga^Y91fs* mutants and in *rag1* crispants. The *il2rga^Y91fs* mutants were analyzed by confocal imaging and crispants were subject to scRNA-seq at 8 dpf. Both groups were

analyzed by *in situ* hybridization. As a control for CRISPR/Cas9 mutagenesis, we targeted the *slc24a5 golden* locus for the crispants experiment. (**B**) Confocal imaging of mCherry signal derived from early-HSPCs in siblings and *il2rga*^{Y91fs} homozygous mutants at 8 dpf in the thymus (left), intestine (middle), and CHT (right). Quantification shows significant decreases in all three locations in the *il2rga* mutants. (**C**) *In situ* RNA hybridization for ILC marker genes in 8 dpf larvae of all three *il2rga*^{Y91fs} genotypes. Triangles point to quantified puncta, while dotted line circles the thymus. Images (left) are representative of the mean signal. Quantification (right) shows all three ILC genes were decreased. (**D,D'**) UMAP showing the lymphoid subset of clusters detected in controls and *rag1* crispants in D. (D') The *rag1*-independent clusters are circled. Quantification of the percentage of cells in each cluster within the lymphoid subset shown in (D). See supplemental for absolute numbers. (**E**) *In situ* RNA hybridization for ILC marker genes in 8 dpf in controls and *rag1* crispants. Images (left) are representative of the mean signal. Triangles point to quantified puncta, while dotted line circles the thymus. Quantification (right) shows *eomesa* was unchanged, while *swift* was increased and *il7r* was decreased in the *rag1* crispants. Statistics: in (B,E) average ± std and p-values are calculated with an unpaired t-test.

A Cre-Lox system

drl | creER^T2

Tg(drl:creER^T2, ubi:loxP-GFP-loxP-stop-mCherry) in **slc24a5 crispant controls** or **rag1 crispants**

Induce labeling of early-HSPCs

early ↓ 4-OHT ...... + 28.5 µM R848
1   2   3   4   5   6   7   8
day-post-fertilization(dpf)

Treatment with Resiquimod (R848)

Confocal imaging or scRNA-seq of **mCherry+** cells at 8 dpf:
slc24a5 crispants — Control
slc24a5 crispants | rag1 crispants — +R848

B *Tg(draculin:creER^T2, ubi:loxP-GFP-loxP-stop-mCherry)* in **Control or R848 treated**

Thymus — Control 8 dpf / R848 8 dpf — mCherry CTCF in Thymus — ****
H2O control | R848 treated

Intestine — Control 8 dpf / R848 8 dpf — # of puncta along intestine — ****
H2O control | R848 treated

CHT — Control 8 dpf / R848 8 dpf — mCherry CTCF in CHT — ****
H2O control | R848 treated

C slc24a5 crispants control — 8 dpf / slc24a5 crispants control +R848 — 8 dpf / rag1 crispants +R848 — 8 dpf

ILC1/NK-like, ILC2-like, il2rb+T-cells, Macrophages

0:Erythroid — 5:Macrophages — 10:Erythroid Progenitors — 15:Endothelial — 20:ILC3-like — 25:Erythroid
1:Erythroid — 6:il2rb+Tcells — 11:Proliferating — 16:blnk+Immune Cells — 21:ILC2-like — 26:Neutrophils
2:Erythroid Progenitors — 7:MyPs — 12:ILC1/NK-like — 17:Immature T-cells — 22:rag+T-cells — 27:APCs
3:Neutrophils — 8:Thrombocytes — 13:Neutrophils — 18:Macrophages — 23:LyPs — 28:il22+Immune Cells
4:myb+HSCs — 9:Macrophages — 14:cd59+Immune Cells — 19:Proliferating — 24:ILC2-like

D Macrophages
5:Macs. / 9:Macs. / 18:Macs.
slc24a5 crisp.ctr. / slc24a5 crisp.ctr. +R848 / rag1 crispant +R848

ILCs and T cells
6:il2rb+Tcells / 12:ILC1/NK-like / 21:ILC2-like / 24:ILC2-like / 20:ILC3-like / 22:rag+T-cells
**** / **** / ****
slc24a5 crisp.ctr. / slc24a5 crisp.ctr. +R848 / rag1 crispant +R848

E 12:ILC1/NK-like
ribosome biogenesis
rRNA processing
mRNA modification
ragulation of macroautophagy
tRNA processing
regulation of actin filament polymerization
actin filament polymerization
Arp2/3 complex-mediated actin nucleation
humoral immune response mediated by circulating immunoglobulin
complement activation
Ctrl. (42) / R848 (10)
p.adjust / GeneRatio

F *eomesa* RNA at 8 dpf
Control 8 dpf n=19 / R848 8 dpf n=17 — # of eomesa+ puncta — ****
H2O Control | R848 treated

swift RNA at 8 dpf
Control 8 dpf n=59 / R848 8 dpf n=57 — # of swift+ puncta — ns
H2O Control | R848 treated

il7r RNA at 8 dpf
Control 8 dpf n=20 / R848 8 dpf n=21 — Signal inThymus — ***
H2O Control | R848 treated

Figure 7. Viral mimicry stimulates ILC-like cell expansion in zebrafish larvae. (A) Outline of experiment exploring the impact of viral mimicry on the larval immune cell landscape. Crispants for *slc24a5* (control) and *rag1* were treated with either vehicle control or 28.5μM R848 from 6-8 dpf and mCherry+ cells from each condition were sequenced at 8 dpf. (**B**) Confocal imaging of early-HSPCs derived mCherry+ cells in the thymus (left), intestine (middle), and CHT (right) at 8 dpf in controls and R848-treated larvae. Images (left) are representative of the mean signal. Quantification (right) shows increased mCherry+ signal in all three locations. (**C**) UMAP showing all hematopoietic clusters detected at 8 dpf per condition. Clusters showing substantial expansion upon R848 treatment are circled. Clusters 11 and 19 are not circled as they contains proliferating cells representing a mix of identities. (**D**) Bar plots show the percentages of cells in macrophage or the ILC/T-cell clusters. Percentages are relative to all the cells shown in UMAP in (C), see supplemental for absolute numbers. (**E**) Gene enrichment analysis using EnrichGO for Biological Processes on genes upregulated in ILC1/NK-like cluster in response to R848 treatment showing enrichment in cytokine and chemokine activity. (**F**) *In situ* RNA hybridization for ILC genes in control and R848 treated 8 dpf larvae. Triangles point to quantified puncta, while dotted line circles the thymus. Images (left) are representative of the mean signal. Quantification (right) shows no change in *swift* levels and significant increases in *eomesa* and *il7r*. Statistics: in (B,F) average ± std and p-values calculated with an unpaired t-test.