Honeybees collect pollen from the buzz-pollinated flowers of invasive

Solanum elaeagnifolium in Northern Greece

Kantsa, A.[1]; Nakas, G.[2]; Kariyat, R.R.[1,3]; Lambert, H.[1]; Petanidou, T.[2]; De Moraes, C.M.[1]; Mescher, M.C.[1]*

Corresponding author

[1] Department of Environmental Systems Science, ETH Zurich, Switzerland.

[2] Department of Geography, University of the Aegean, Greece.

[3] Department of Entomology & Plant Pathology, University of Arkansas, Fayetteville AR, USA.

ORCID IDs and e-mails:

Kantsa, A.: a.kantsa@gmail.com; 0000-0002-4586-1109

Nakas, G.: nakas.g@geo.aegean.gr; 0000-0003-3023-583

Kariyat, R.R.: rkariyat@uark.edu; 0000-0002-6565-6276

Lambert, H.: harriet.lambert@wur.nl 0000-0001-7427-597X

Petanidou, T.: t.petanidou@aegean.gr; 0000-0003-1883-0945

De Moraes, C.M.: consuelo.demoraes@usys.ethz.ch; 0000-0001-6737-9842

Mescher, M.C.: mescher@usys.ethz.ch; 0000-0002-7908-3309

Abstract

Around 6% of flowering plants, including most solanaceous species, have flowers

specialized for buzz-pollination (sonication). The poricidal anthers of these flowers release

pollen in response to frequency-specific vibrations, which buzz-pollinating bees, including

bumble bees and many solitary bees, generate via rapid oscillation of flight muscles.

Honeybees are incapable of sonication and typically avoid visiting buzz-pollinated plants,

although they have previously been observed to scavenge residual pollen from petals of buzz-

pollinated flowers in natural settings. Here, we report a novel interaction in which honeybees

consistently visit and extract pollen from flowers of *Solanum elaeagnifolium* (silverleaf

nightshade) within a large invasive plant population in Northern Greece. We postulate that

honeybee foraging effort on *S. elaeagnifolium* reflects the shortage of alternative floral

resources at the peak of the Mediterranean summer, consistent with observations for pollen-

starved bees in captivity. This newly established behavioral shift of honeybees may convey

important implications for plant invasions, particularly under climate change, and requires

further investigation for untangling its ultimate drivers.

Keywords

Agroecology; *Apis mellifera*; Buzz-pollination; Plant invasions; Pollen resources; Solanaceae;

Weed ecology

Introduction

"...the polylectic honeybee does not seem able to learn the process."

Linsley (1978)

Buzz-pollinated flowers occur in around 20,000 species across various angiosperm

clades (~6% of species), including key crops such as tomato, eggplant, blueberry, cranberry,

and kiwi (Buchmann 1983, De Luca and Vallejo-Marín 2013, Pritchard and Vallejo-Marín 2020,

Cooley and Vallejo-Marín 2021). Owing to their unique floral morphology, the poricidal anthers

of these flowers require vibration at specific frequencies (also known as sonication) to release

pollen. Buzz pollination is hypothesized to serve as an adaptation for preventing pollen access

by relatively inefficient, non-specialist pollinators (Pritchard and Vallejo-Marín 2020), although

the morphology of buzz-pollinated flowers may also deter herbivory by making pollen less

accessible (Hargreaves et al. 2009). A capacity for sonication via rapid oscillation of the thoracic

indirect flight muscles has evolved independently ca. 45 times within bees (Cardinal et al.

2018), enabling approximately 58% of bee species (at least 74 genera) to efficiently extract

pollen from buzz-pollinated flowers (Buchmann 1983, De Luca and Vallejo-Marín 2013,

Pritchard and Vallejo-Marín 2020, Jankauski et al. 2022). However, honeybees, the most

frequent floral visitors in both natural and agricultural systems worldwide (Hung et al. 2018), are

incapable of sonication, and their limited interaction with buzz-pollinated plant species has

been widely discussed by pollination biologists (Linsley 1978, Buchmann 1983, De Luca and

Vallejo-Marín 2013, Vallejo-Marin et al. 2024).

The proximate mechanisms preventing honeybees from effectively collecting pollen

from buzz-pollinated flowers appear to be based in physiological constraints—specifically

relating to the maximum sustained frequency of thoracic vibrations (King and Buchmann

2003)—rather than a lack of behavioral flexibility or learning capacity, as had previously been

suggested (Linsley 1978). In any case, honeybees typically avoid buzz-pollinated flowers in

natural settings, and particularly those of species that do not offer nectar, as in the genus

Solanum (Michener 1962, Linsley 1978, Buchmann 1983, Buchmann and Cane 1989, Tscheulin

et al. 2009, Tscheulin and Petanidou 2013, Petanidou et al. 2018, Tayal and Kariyat 2021). On

rare occasions in which honeybees have been observed to visit buzz-pollinated flowers in the

wild, they have primarily been reported to glean leftover pollen from petals following visits by

buzz-pollinators (Wille 1963, Buchmann 1983, Buchmann and Cane 1989, Gross and Mackay

1998, Solís-Montero et al. 2015, Mesquita-Neto et al. 2018, Petanidou et al. 2018, Rocha et al.

2023), and sometimes to steal already deposited pollen from the stigma, presumably reducing

plant fitness (e.g., Gross and Mackay 1998).

Despite their limited ability to utilize buzz-pollinated flowers, honeybees can be induced

to visit such flowers under conditions of pollen deprivation and are sometimes utilized for

pollination of buzz-pollinated agricultural crops. In the case of buzz-pollinated flowers that do

not offer nectar (e.g., tomato, eggplant, potato, tamarillo), managed honeybees will visit only

when confined into greenhouses or flight cages with no access to other pollen sources

(Neiswander 1956, Cane et al. 1993, Cane and Schiffhauer 2003, Miyamoto et al. 2006).

However, honeybees will spontaneously visit only the nectar-offering buzz-pollinated crops

(e.g., blueberry, cranberry), in which case they can be used in open environments (Cane et al.

1993, Dogterom and Winston 1999). The reported pollinating efficiency of honeybees on buzz-

pollinated crops ranges from zero (avoided the flowers, although pollen-deprived; e.g, Sanford

and Hanneman 1981, Banda and Paxton 1990, Amala and Shivalingaswamy 2017, Zhang et al.

2020, Rijo et al. 2022), to inconsistent (Higo et al. 2004, Dos Santos et al. 2009, Courcelles et al.

2013, Cooley and Vallejo-Marín 2021), or sometimes satisfactory provided that hives are used in

high densities and honeybee workers do not steal nectar (Neiswander 1956, Cribb et al. 1993,

MacKenzie 1994, Delaplane and Mayer 2000, Cane and Schiffhauer 2003, Sabara and Winston

2003, Miyamoto et al. 2006, Courcelles et al. 2013, Cooley and Vallejo-Marín 2021).

The observation that managed honeybees will visit and attempt to collect pollen from

buzz-pollinated flowers under pollen deprivation, raises the possibility that similar behaviors

may occur in nature, particularly in environments where honeybees have limited access to other

pollen sources or, conversely, where other potential pollinators are absent or rare. Both

conditions might arise in the context of plant invasion (Albrecht et al. 2016, Davis et al. 2018).

Yet, as noted above, reports of honeybee visitation of buzz-pollinated flowers in nature are rare

and generally limited to observations of bees collecting residual pollen from petals, following

visitation by other, buzz-pollinating species. In the current study, we report the large-scale

extraction and utilization by honeybees of pollen from a buzz-pollinated plant (*Solanum*

elaeagnifolium), which forms large monocultures in its invaded (Mediterranean) range and

produces flowers in a seasonal period when other flowers are scarce. We also provide evidence

that this honeybee behavior has become widespread within the last decade and discuss

potential ecological and evolutionary implications within the context of plant invasion.

Materials and Methods

Study species

Solanum elaeagnifolium (silverleaf nightshade) is currently among the top-noxious

invasive plants worldwide (Uludag et al. 2016, Roberts and Florentine 2022). A perennial

herbaceous species originating in N Mexico and SW USA, *S. elaeagnifolium* has been rapidly

expanding to almost all continents over the past century, posing serious threats to native

biodiversity and causing significant losses in agricultural and livestock production in many

countries (Uludag et al. 2016, Roberts and Florentine 2022). Its uncontrollable expansion has

severely affected the Mediterranean Basin, making it one of the most impacted regions (Uludag

et al. 2016, Krigas et al. 2021). In croplands and urban areas of N Greece, this species forms

numerous extensive monospecific stands (of >10,000 individuals) (Petanidou et al. 2018) (Figure

1).

Bee observations

Bee observations on *S. elaeagnifolium* were performed in July-August 2015, 2016, 2017,

and 2023 in the Metropolitan Area of Thessaloniki, Greece (Table 1; Figure 1), at sites with either

urban or agricultural land use, which had been monitored in the context of previous research of

this species in the area in 2006-2007 (Petanidou et al. 2012, Petanidou et al. 2018).

Observations were made during 30-min random walks across each study site in the morning

hours (09:00-12:00). Honeybees that visited the flowers and tried to extract pollen using any of

the behaviors described above were photographed and video-recorded, and samples were

brought to the lab in order to identify the pollen they carried.

Scanning electron microscopy (SEM)

Honeybees foraging on *S. elaeagnifolium* were captured and brought to the lab to

confirm the presence of pollen from this plant in their corbiculae and on their bodies. For this,

we compared SEM images of the pollen of this species and of the pollen found on the honeybee

specimens [see also Du et al. (2018)]. SEM was performed at the Scientific Center for Optical

and Electron Microscopy of ETH Zurich (ScopeM), using a multi-purpose analytical high

resolution field emission gun scanning electron microscope (HR-FEG-SEM Quanta200F).

Images were acquired at a 10 kV accelerating voltage under low vacuum conditions (100 Pa),

with the secondary electron (SE) signal recorded using the LFD detector. The field of view (FOV)

was 149 µm or 298 µm, depending on the imaging setup. Since charging was suppressed by

using the low vacuum mode, no metal coating was applied on the bees before introducing them

to the SEM vacuum chamber (Symondson and Williams 1997).

Figure 1. Invasive population stands in a cultivated field (up) and in an urban residential area (down)

in the Metropolitan area of Thessaloniki, N Greece (photos: R.R. Kariyat, G. Nakas).

Table 1. Study sites, surveys, and honeybee observations on the flowers of *Solanum elaeagnifolium*
in Thessaloniki. The years of the first and the last surveys performed at each site, as well as the year
in which the interaction between the plant and honeybees was first observed are shown.

Site ID	Site name	Coordinates	Land use	First survey	Last survey	First observation
S01	Sindos	40° 41' 6.0" N, 22° 43' 40.8" E	Agricultural	2007	2016	-
S02	Technological Institute	40° 40' 12.0" N, 22° 47' 20.4" E	Agricultural	2007	2023	-
S03	Anchialos	40° 43' 32.2" N, 22° 41' 39.1" E	Waste land	2007	2016	-
S04	Redaistos	40° 30' 58.88" N, 23° 4' 21.37" E	Agricultural	2007	2017	2015
S05	Panorama	40° 35' 26.72" N, 23° 2' 38.96" E	Semi-natural	2007	2015	-
S06	Asvestochori	40° 40' 15.6" N, 23° 3' 0.0" E	Semi-natural	2007	2015	-
S07	Thermi	40° 32' 6.29" N, 22° 59' 45.12" E	Agricultural	2016	2023	2017
S08	Airport	40° 31' 10.9" N, 22° 58' 15.2" E	Semi-natural	2023	2023	2023
S09	City center	40° 37' 20.22" N, 22° 57' 17.89" E	Urban parks	2023	2023	2023
S10	Stavroupoli	40° 39' 32.8" N, 22° 55' 8.8" E	Urban parks	2023	2023	-

Results and Discussion

Since 2015, beginning in the E–SE part of the Thessaloniki metropolitan area and
gradually expanding towards the city center (Figure 2), we have frequently observed honeybees
visiting *S. elaeagnifolium* flowers and actively extracting pollen from the poricidal anthers using
three distinct behaviors: (a) shaking/drumming the anthers using their forelegs; (b) biting off
pieces of anthers with their mandibles; (c) holding the anthers with their forelegs and inserting
their proboscis through the pores of the anthers (Videos S1-S3). As a result, pollen of silverleaf

nightshade adheres on the bodies of honeybees and is used to fill up the corbiculae (Videos S1-

S3) (Figure 3). These modes of behavior have previously been reported for honeybees trying to

exploit buzz-pollinated flowers in cultivation, and Cane et al. (1993) hypothesized that the

foreleg drumming behavior is a permutation of pollen-grooming movements that bees perform

to transfer pollen grains from their upper body parts to the corbiculae.

Figure 2. Study sites of *Solanum elaegnifolium* in the Metropolitan Area of Thessaloniki. In the sites

where the honeybees have been observed to interact with the flowers, the year of the first

observation is shown. The city center is marked with dark grey. See Table 1 for details on the sites.

As documented in agricultural settings, honeybees spend more time on visitation of

buzz-pollinated flowers than they usually do on other types of flowers, presumably due to the

low efficiency of pollen extraction (Kendall et al. 2022). However, honeybee interactions with *S.*

elaeagnifolium entail a high level of settings, physical effort and an exceptionally long-duration

of floral visits. Indeed, in Thessaloniki, we recorded visits on single flowers lasting as long as 224

s (Video S1), whereas known maximum visitation times of honeybees on buzz-pollinated

species are 100 s, observed on blueberry flowers (Kendall et al. 2022). Instead, for non-buzz-

pollinated species, handling times are considerably shorter, typically ranging from 2.6 s per

flower (oregano) and 4 s (lavender) (Hennessy et al. 2021) to 40 s (melon) (Ribeiro et al. 2017).

Figure 3. (a-b) Honeybees foraging for pollen on *Solanum elaeagnifolium*; their corbiculae (pollen

baskets) are replete with the pollen of the plant (photos: R.R. Kariyat). (c-d) SEM images showing the

pollen grains, and how these accumulate among the bristles on the body of the bee.

Furthermore, especially during the latter part of the flowering season (August),

honeybees in Thessaloniki regularly visited *S. elaeagnifolium* flowers that were in poor

condition, (e.g., heavily damaged by florivores, wilting, or deformed), which otherwise is known

to reduce attractiveness to pollinators (e.g., Sõber et al. 2010, Vega-Polanco et al. 2020). Finally,

it should be noted that honeybees are now by far the most frequent visitors of these flowers,

outnumbering buzz-pollinating bees in the area (i.e., *Bombus, Pseudapis, Xylocopa,* and

Amegilla) (RRK and GN personal observations). To our knowledge, this is the first report of

honeybees consistently extracting pollen from a buzz-pollinated plant in a natural community.

The striking similarity of the behavior reported here with that previously reported for

pollen-starved honeybees in confined agricultural settings (see above), suggests that it might be

linked to severe limitation of other pollen sources. This is highly plausible given the magnitude

of the invasion of silverleaf nightshade in the study area (Petanidou et al. 2018). It has been

estimated that *S. elaeagnifolium* reached current population sizes in and around Thessaloniki

city within 50 years after its first unintentional introduction (most probably from Texas) in the

1930s, forming one of the most heavily invaded areas in Europe (Krigas and Kokkini 2004,

Uludag et al. 2016, Roberts and Florentine 2022). While we cannot be certain about when the

widespread utilization of *S. elaeagnifolium* flowers by honeybees in Thessaloniki began, we can

be confident that it started later than 2007, based on previous extensive surveys of pollinator

activity in this area (Petanidou et al. 2012, Petanidou et al. 2018). While these surveys reported

that other wild bees were scarce (Petanidou et al. 2018), honeybee interactions with *S.*

elaeagnifolium were also rare, and restricted to gleaning leftover pollen from petals, presumably

following visits by other pollinators (Petanidou et al. 2012, Petanidou et al. 2018). Similar

behavior has been repeatedly reported for silverleaf nightshade both in Greece (Tscheulin et al.

2009, Tscheulin and Petanidou 2013) and in its native range, i.e., Texas and Arizona (Buchmann

and Cane 1989, Tayal and Kariyat 2021). Besides, this fits the typical interaction of honeybees

with *Solanum*-type flowers in the wild, where they have been reported to either avoid (*S.*

wedlandii) (Michener 1962), or glean pollen leftovers (*S. rostratum*) (Solís-Montero et al. 2015)

from flowers of *Solanum* species.

Since the successful establishment of silverleaf nightshade in Thessaloniki, local

honeybee populations have presumably been exposed to a novel "operative environment"

(Spomer 1973, Agosta and Klemens 2008), including a super-abundant pollen source, which,

despite being challenging to exploit, is available during the most adverse time of year. Given the

relative scarcity of other insect-pollinated species in flower in the lowlands of the

Mediterranean at the peak of the dry summer (Kantsa et al. 2023b), silverleaf nightshade is by

far the dominant pollen source in the invaded habitats. Therefore, we assume that the observed

utilization of *S. elaeagnifolium* is a consequence of plant invasion in an area with limited

alternative pollen sources, although it remains unclear why the behavior appears to have

become widespread only in recent years.

The well-known aversion of honeybees towards buzz-pollinated flowers suggests that

the utilization of nectarless, buzz-pollinated flowers is not adaptive in typical environments,

likely because the ratio of pollen rewards obtained to effort expended is unattractive, relative to

that provided by other flower types (Buchmann and Cane 1989). If utilization of *S.*

elaeagnifolium flowers has been adaptive under the novel conditions in Thessaloniki, a question

is whether this arises due to pre-existing behavioral flexibility or entails genetic adaptation. The

apparently similar exploitation of buzz-pollinated flowers by pollen-starved honeybees in

agricultural settings suggests that the former is plausible; the employment of different

techniques by the honeybees (viz., drumming, "milking", or biting; Videos S1-S3) is consistent

with ecological fitting (Janzen 1985, Agosta and Klemens 2008), suggesting that the eventual

mode of action depends on individual decision-making ability, agility, or both. These behaviors

may thus represent a notable example of behavioral plasticity of the honeybees, facilitating

exploration of novel areas of fitness space created by the massive invasive populations of *S.*

elaeagnifolium in Northern Greece.

Regarding why the reported behavior was not observed prior to 2015, it may be useful to

explore the potential preference of relevant colony-level heritable traits (e.g., pollen hoarding) in

the local beekeeping market. For instance, Cane and Schiffhauer (2001, 2003) suggested that

when honeybee queens are genetically predisposed for pollen hoarding (Rueppell 2014) pollen-

foraging trips on buzz-pollinating flowers are significantly more frequent. Furthermore, the

previously mentioned hypothesis by Cane et al. (1993) that the movement of foreleg drumming

in honeybees is a permutation of pollen-grooming movements that bees perform to transfer

pollen grains from their upper body parts to the corbiculae, suggests the possibility of an

existing pre-adaptation. This could indicate a case of exaptation, where a trait originally evolved

for one function (pollen grooming) has been co-opted for another (buzz-pollination). A similar

example is the woodpecker finches (*Camarhynchus pallidus*, Thraupidae) at the Galápagos

Islands, in which beak morphology originally suited for cracking seeds or eating insects has

been co-opted for tool utilization, a behavior which varies according to habitat type (i.e.,

operative environment), is not acquired with social learning (but possibly with trial-and-error

learning at a young age), and is believed to be based on genetic predisposition (Tebbich et al.

2001). In any case, whether honeybees have established a novel interaction circumventing

adaptive processes (through phenotypic plasticity), or we are witnessing the real-time

establishment of an exapted behavior, possibly conveying a more subtle and complex genetic

background (e.g., pleiotropic traits), this phenomenon requires in-depth investigation.

Honeybees are globally abundant, frequently utilized in agriculture, and highly

competitive with wild bees, often with adverse effects on native bee faunae (Goulson 2003,

Lázaro et al. 2021). Given that they are generally among the first pollinators to interact with

newly introduced plants in natural communities (Kantsa et al. 2023a), these behavioral shifts

can be directly linked with invasion status and dynamics, with *S. elaeagnifolium* providing a

characteristic example. In general, the rapid geographical expansion of this species is thought

to be promoted largely by its ability to propagate vegetatively with root fragments (Roberts and

Florentine 2022). The invasive populations of silverleaf nightshade are known to be self-

incompatible, and (at least until 2007) pollen-limited (Petanidou et al. 2012, Petanidou et al.

2018). Based on these observations, it had been suggested that sexual reproduction was less

important for the expansion of this plant in the invaded area. Yet, if this condition is overturned,

and honeybee visitation can now convey the advantages of outcrossing, the genetic robustness

of this vast invasive population is expected to be even further enhanced, with potentially even

more disastrous effects on the native wildlife and landscape. For example, in the genus

Solanum, outcrossing maintains high chemodiversity of volatile emissions, and it relates to

better performance in chemically repelling herbivores and in recruiting predators (Kariyat et al.

2012a, Kariyat et al. 2012b, Kariyat et al. 2013).

Outlook

Here we report a new foraging behavior in honeybees, which has emerged during the

past decade, probably owing to the substantial expansion of *S. elaeagnifolium*, resulting in an

overabundance of pollen during the most adverse period of the year in the Mediterranean in

terms of floral resources. We have argued that our observation, likely involving a complex

interplay of honeybee biology and floral ecology, has important implications both for the

invasion status of silverleaf nightshade and for honeybee biology.

The enormous invasive populations of silverleaf nightshade in Northern Greece (and

probably elsewhere) in fact represent large-scale natural experiments, which can reveal

fascinating aspects of the evolution of plant–pollinator interactions. For instance, we know that

buzz-pollination is not a dead-end, and novel floral forms (that do not even require bees) can

evolve from bee-buzz-pollinated flowers (Dellinger et al. 2019). In the system examined here, we

see the archetypical buzz-pollinated flower form (the *Solanum*-type) potentially experiencing a

pollinator shift with apparently no phenotypic transition. Further investigation is needed to

determine if this indeed represents a functional transition, the extent floral phenotype has been

modified, and the drivers (e.g., floral abundance, abiotic conditions, honeybee plasticity, etc.) of

these transitions.

Beyond basic research questions, it should also be kept in mind that silverleaf

nightshade is a top-noxious weed that is adversely affecting natural, agricultural-, and urban

ecosystems across the world. Moreover, its invasive potential remains high, especially in the

context of global change: spatial models simulating climatic suitability for this plant indicate

numerous currently invadable regions in all continents, with poleward expansions predicted in a

warmer atmosphere (Kriticos et al. 2010). Given that honeybees occur with generally high

population densities in almost all continents, their impact on the realized fitness of invasive

populations of *S. elaeagnifolium*, and the broader implications for natural communities should

be explored.

Acknowledgements

GN and TP would like to thank Chrysoula Tananaki and Georgios Goras for support during fieldwork.

Conflict of interest statement

The Authors declare no conflict of interest.

Author contributions

MCM, CMDM, and RRK designed the study. RRK, HL, TP, and GN performed field observations. RRK performed scan microscopy. AK processed the observations, curated media, and wrote the first draft. All co-authors revised the manuscript.

References

Agosta, S. J., and J. A. Klemens. 2008. Ecological fitting by phenotypically flexible genotypes: implications for species associations, community assembly and evolution. Ecology Letters **11**:1123-1134.

Albrecht, M., M. R. Ramis, and A. Traveset. 2016. Pollinator-mediated impacts of alien invasive plants on the pollination of native plants: the role of spatial scale and distinct behaviour among pollinator guilds. Biological Invasions **18**:1801-1812.

Amala, U., and T. Shivalingaswamy. 2017. Role of native buzz pollinator bees in enhancing fruit and seed set in tomatoes under open field conditions. Journal of Entomology and Zoology Studies **5**:1742-1744.

Banda, H., and R. Paxton. 1990. Pollination of greenhouse tomatoes by bees. Pages 194-198 *in* VI International Symposium on Pollination 288.

Buchmann, S. L. 1983. Buzz pollination in angiosperms. Pages 73-113. Van Nostrand Reinhold Company, New York, NY.

Buchmann, S. L., and J. H. Cane. 1989. Bees assess pollen returns while sonicating *Solanum* flowers. Oecologia **81**:289-294.

Cane, J. H., K. Mackenzie, and D. Schiffhauer. 1993. Honey bees harvest pollen from the porose
anthers of cranberries (*Vaccinium macrocarpon*) (Ericaceae). American Bee Journal
133:293-295.

Cane, J. H., and D. Schiffhauer. 2001. Pollinator genetics and pollination: do honey bee colonies
selected for pollen-hoarding field better pollinators of cranberry *Vaccinium macrocarpon*?
Ecological Entomology **26**:117-123.

Cane, J. H., and D. Schiffhauer. 2003. Dose-response relationships between pollination and fruiting
refine pollinator comparisons for cranberry (*Vaccinium macrocarpon* [Ericaceae]). American
Journal of Botany **90**:1425-1432.

Cardinal, S., S. L. Buchmann, and A. L. Russell. 2018. The evolution of floral sonication, a pollen
foraging behavior used by bees (Anthophila). Evolution **72**:590-600.

Cooley, H., and M. Vallejo-Marín. 2021. Buzz-Pollinated Crops: A Global Review and Meta-analysis of
the Effects of Supplemental Bee Pollination in Tomato. Journal of Economic Entomology
114:505-519.

Courcelles, D. M. M., L. Button, and E. Elle. 2013. Bee visit rates vary with floral morphology among
highbush blueberry cultivars (*Vaccinium corymbosum* L.). Journal of Applied Entomology
137:693-701.

Cribb, D. M., D. W. Hand, and R. N. Edmondson. 1993. A comparative study of the effects of using the
honeybee as a pollinating agent of glasshouse tomato. Journal of Horticultural Science
68:79-88.

Davis, E. S., R. Kelly, C. A. Maggs, and J. C. Stout. 2018. Contrasting impacts of highly invasive plant
species on flower-visiting insect communities. Biodiversity and Conservation **27**:2069-2085.

De Luca, P. A., and M. Vallejo-Marín. 2013. What's the 'buzz' about? The ecology and evolutionary
significance of buzz-pollination. Current Opinion in Plant Biology **16**:429-435.

Delaplane, K. S., and D. F. Mayer. 2000. Crop pollination by bees. CABI publishing.

Dellinger, A. S., M. Chartier, D. Fernández-Fernández, D. S. Penneys, M. Alvear, F. Almeda, F. A.
Michelangeli, Y. Staedler, W. S. Armbruster, and J. Schönenberger. 2019. Beyond buzz-
pollination – departures from an adaptive plateau lead to new pollination syndromes. New
Phytologist **221**:1136-1149.

Dogterom, M. H., and M. L. Winston. 1999. Pollen storage and foraging by honey bees (Hymenoptera:
Apidae) in highbush blueberries (Ericaceae), cultivar Bluecrop. The Canadian Entomologist
131:757-768.

Dos Santos, S., A. Roselino, M. Hrncir, and L. Bego. 2009. Pollination of tomatoes by the stingless bee
Melipona quadrifasciata and the honey bee *Apis mellifera* (Hymenoptera, Apidae). Genet.
Mol. Res **8**:751-757.

Du, T., C. Zhao, and J. Liu. 2018. The pollen of *Solanum* L. and its systematic significance. Palynology
42:291-310.

Goulson, D. 2003. Effects of Introduced Bees on Native Ecosystems. Annual Review of Ecology,
Evolution, and Systematics **34**:1-26.

Gross, C. L., and D. Mackay. 1998. Honeybees reduce fitness in the pioneer shrub *Melastoma affine*
(Melastomataceae). Biological Conservation **86**:169-178.

Hargreaves, A. L., L. D. Harder, and S. D. Johnson. 2009. Consumptive emasculation: the ecological
and evolutionary consequences of pollen theft. Biological Reviews **84**:259-276.

Hennessy, G., C. Harris, L. Pirot, A. Lefter, D. Goulson, and F. L. W. Ratnieks. 2021. Wind slows play:
increasing wind speed reduces flower visiting rate in honey bees. Animal Behaviour **178**:87-
93.

Higo, H. A., N. D. Rice, M. L. Winston, and B. Lewis. 2004. Honey bee (Hymenoptera: Apidae)
distribution and potential for supplementary pollination in commercial tomato greenhouses
during winter. Journal of Economic Entomology **97**:163-170.

Hung, K.-L. J., J. M. Kingston, M. Albrecht, D. A. Holway, and J. R. Kohn. 2018. The worldwide
importance of honey bees as pollinators in natural habitats. Proceedings of the Royal
Society B: Biological Sciences **285**:20172140.

Jankauski, M., R. Ferguson, A. Russell, and S. Buchmann. 2022. Structural dynamics of real and
modelled *Solanum* stamens: implications for pollen ejection by buzzing bees. Journal of the
Royal Society Interface **19**:20220040.

Janzen, D. H. 1985. On Ecological Fitting. Oikos **45**:308-310.

Kantsa, A., C. M. De Moraes, and M. C. Mescher. 2023a. Global change and plant–pollinator
communities in Mediterranean biomes. Global Ecology and Biogeography **32**:1893-1913.

Kantsa, A., J. E. Garcia, R. A. Raguso, A. G. Dyer, R. Steen, T. Tscheulin, and T. Petanidou. 2023b.
Intrafloral patterns of color and scent in *Capparis spinosa* L. and the ghosts of its selection
past. American Journal of Botany **110**:e16098.

Kariyat, R. R., C. M. De Moraes, A. G. Stephenson, and M. C. Mescher. 2012a. Inbreeding increases
susceptibility to powdery mildew (*Oidium neolycopersici*) infestation in horsenettle
(*Solanum carolinense* L.). Plant Signaling & Behavior **7**:803-806.

Kariyat, R. R., K. E. Mauck, C. M. Balogh, A. G. Stephenson, M. C. Mescher, and C. M. De Moraes.
2013. Inbreeding in horsenettle (*Solanum carolinense*) alters night-time volatile emissions
that guide oviposition by *Manduca sexta* moths. Proc Biol Sci **280**:20130020.
Kariyat, R. R., K. E. Mauck, C. M. De Moraes, A. G. Stephenson, and M. C. Mescher. 2012b. Inbreeding
alters volatile signalling phenotypes and influences tri-trophic interactions in horsenettle
(*Solanum carolinense* L.). Ecology Letters **15**:301-309.
Kendall, L. K., J. R. Stavert, V. Gagic, M. Hall, and R. Rader. 2022. Initial floral visitor identity and
foraging time strongly influence blueberry reproductive success. Basic and Applied Ecology
60:114-122.
King, M. J., and S. L. Buchmann. 2003. Floral Sonication by Bees: Mesosomal Vibration by *Bombus*
and *Xylocopa*, but Not *Apis* (Hymenoptera: Apidae), Ejects Pollen from Poricidal Anthers.
Journal of the Kansas Entomological Society **76**:295-305.
Krigas, N., and S. Kokkini. 2004. A Survey of the Alien Vascular Flora of the Urban and Suburban Area
of Thessaloniki, N Greece. Willdenowia **34**:81-99.
Krigas, N., M. A. Tsiafouli, G. Katsoulis, N.-E. Votsi, and M. van Kleunen. 2021. Investigating the
Invasion Pattern of the Alien Plant *Solanum elaeagnifolium* Cav. (Silverleaf Nightshade):
Environmental and Human-Induced Drivers. Plants **805**.
Kriticos, D. J., N. D. Crossman, N. Ota, and J. K. Scott. 2010. Climate change and invasive plants in
South Australia. Report for the South Australian Department of Water, Land and Biodiversity
Conservation. CSIRO Climate Adaptation Flagship, Canberra.
Lázaro, A., A. Müller, A. W. Ebmer, H. H. Dathe, E. Scheuchl, M. Schwarz, S. Risch, A. Pauly, J.
Devalez, T. Tscheulin, C. Gómez-Martínez, E. Papas, J. Pickering, N. M. Waser, and T.
Petanidou. 2021. Impacts of beekeeping on wild bee diversity and pollination networks in
the Aegean Archipelago. Ecography **44**:1353-1365.
Linsley, E. G. 1978. Temporal Patterns of Flower Visitation by Solitary Bees, with Particular Reference
to the Southwestern United States. Journal of the Kansas Entomological Society **51**:531-546.
MacKenzie, K., E. 1994. The foraging behaviour of honey bees (*Apis mellifera* L) and bumble bees
(*Bombus* spp) on cranberry (*Vaccinium macrocarpon* Ait). Apidologie **25**:375-383.
Mesquita-Neto, J. N., N. Blüthgen, and C. Schlindwein. 2018. Flowers with poricidal anthers and their
complex interaction networks—Disentangling legitimate pollinators and illegitimate visitors.
Functional Ecology **32**:2321-2332.
Michener, C. D. 1962. An Interesting Method of Pollen Collecting by Bees from Flowers with Tubular
Anthers. Revista De Biologia Tropical **10**:167–175.

Miyamoto, M., M. Ono, M. Sasaki, H. Abe, and I. Kenmochi. 2006. Development of Fruiting Promotion

System Using Honeybees in Semi-forced Eggplant Culture I. Effectiveness of Pollination by

the European Honeybee. Japanese Journal of Applied Entomology and Zoology **50**:297-304.

Neiswander, R. 1956. Pollination of greenhouse tomatoes by honey bees. Journal of Economic

Entomology **49**:436-437.

Petanidou, T., R. C. Godfree, D. S. Song, A. Kantsa, Y. L. Dupont, and N. M. Waser. 2012. Self-

compatibility and plant invasiveness: Comparing species in native and invasive ranges.

Perspectives in Plant Ecology Evolution and Systematics **14**:3-12.

Petanidou, T., M. V. Price, J. L. Bronstein, A. Kantsa, T. Tscheulin, R. Kariyat, N. Krigas, M. C. Mescher,

C. M. De Moraes, and N. M. Waser. 2018. Pollination and reproduction of an invasive plant

inside and outside its ancestral range. Acta Oecologica **89**:11-20.

Pritchard, D. J., and M. Vallejo-Marín. 2020. Buzz pollination. Curr Biol **30**:R858-R860.

Ribeiro, M. F., E. M. S. Silva, L. H. P. Kiill, K. M. M. Siqueira, M. P. Silva, and M. S. Coelho. 2017.

Foraging of Honeybees (*Apis mellifera*) on Flowers of Yellow Melon (*Cucumis melo*): Duration

of Visits. Journal of Agricultural Science **9**.

Rijo, G., D. Alameda, and A. Barro. 2022. Tomato (*Solanum lycopersicum*) specialized pollination is

isolated from neighboring plants and pollinators. Journal of Pollination Ecology **31**:29-38.

Roberts, J., and S. Florentine. 2022. Biology, distribution and management of the globally invasive

weed *Solanum elaeagnifolium* Cav (silverleaf nightshade): A global review of current and

future management challenges. Weed Research **62**:393-403.

Rocha, F. H., D. N. Peraza, S. Medina, and J. J. G. Quezada-Euán. 2023. Pollination service provided

by honey bees to buzz-pollinated crops in the Neotropics. PLoS ONE **18**:e0280875.

Rueppell, O. 2014. The architecture of the pollen hoarding syndrome in honey bees: implications for

understanding social evolution, behavioral syndromes, and selective breeding. Apidologie

45:364-374.

Sabara, H. A., and M. L. Winston. 2003. Managing honey bees (Hymenoptera: Apidae) for greenhouse

tomato pollination. Journal of Economic Entomology **96**:547-554.

Sanford, J. C., and R. E. Hanneman. 1981. The use of bees for the purpose of inter-mating in potato.

American Potato Journal **58**:481-485.

Sõber, V., M. Moora, and T. Teder. 2010. Florivores decrease pollinator visitation in a self-

incompatible plant. Basic and Applied Ecology **11**:669-675.

Solís-Montero, L., C. H. Vergara, and M. Vallejo-Marín. 2015. High incidence of pollen theft in natural

populations of a buzz-pollinated plant. Arthropod-Plant Interactions **9**:599-611.

Spomer, G. G. 1973. The Concepts of 'Interaction' and 'Operational Environment' in Environmental
Analyses. Ecology **54**:200-204.

Symondson, W., and I. Williams. 1997. Low-vacuum electron microscopy of carabid
chemoreceptors: a new tool for the identification of live and valuable museum specimens.
Entomologia Experimentalis Et Applicata **85**:75-82.

Tayal, M., and R. Kariyat. 2021. Examining the Role of Buzzing Time and Acoustics on Pollen
Extraction of *Solanum elaeagnifolium*. Plants **10**:2592.

Tebbich, S., M. Taborsky, B. Fessl, and D. Blomqvist. 2001. Do woodpecker finches acquire tool-use
by social learning? Proceedings of the Royal Society of London. Series B: Biological
Sciences **268**:2189-2193.

Tscheulin, T., and T. Petanidou. 2013. The presence of the invasive plant *Solanum elaeagnifolium*
deters honeybees and increases pollen limitation in the native co-flowering species
Glaucium flavum. Biological Invasions **15**:385-393.

Tscheulin, T., T. Petanidou, S. G. Potts, and J. Settele. 2009. The impact of *Solanum elaeagnifolium*,
an invasive plant in the Mediterranean, on the flower visitation and seed set of the native co-
flowering species *Glaucium flavum*. Plant Ecology **205**:77-85.

Uludag, A., G. Gbehounou, J. Kashefi, M. Bouhache, M.-C. Bon, C. Bell, and A. L. Lagopodi. 2016.
Review of the current situation for *Solanum elaeagnifolium* in the Mediterranean Basin.
EPPO Bulletin **46**:139-147.

Vallejo-Marin, M., D. L. Field, J. Fornoni, D. Montesinos, C. A. Dominguez, I. Hernandez, G. C. Vallejo,
C. Woodrow, R. Ayala Barajas, and N. Jafferis. 2024. Biomechanical properties of non-flight
vibrations produced by bees. Journal of Experimental Biology **227**.

Vega-Polanco, M., L. A. Rodríguez-Islas, R. Y. Escalona-Domenech, L. Cruz-López, J. C. Rojas, and L.
Solís-Montero. 2020. Does florivory affect the attraction of floral visitors to buzz-pollinated
Solanum rostratum? Arthropod-Plant Interactions **14**:41-56.

Wille, A. 1963. Behavioral adaptations of bees for pollen collecting from *Cassia* flowers. Revista De
Biologia Tropical **11**:205–210.

Zhang, H., S. Shan, S. Gu, X. Huang, Z. Li, A. Khashaveh, and Y. Zhang. 2020. Prior Experience with
Food Reward Influences the Behavioral Responses of the Honeybee *Apis mellifera* and the
Bumblebee *Bombus lantschouensis* to Tomato Floral Scent. Insects **11**.

Electronic Supplementary Material

Video S1. A honeybee worker attempting to extract pollen from the flowers of silverleaf

nightshade by biting the anthers. Videographer: Georgios Nakas.

Video S2. A honeybee worker attempting to extract pollen from the flowers of silverleaf

nightshade by holding the anthers with their forelegs and inserting their proboscis through the

pores of the anthers ("milking" behavior). Videographer: Georgios Nakas.

Video S3. A honeybee worker attempting to extract pollen from the flowers of silverleaf

nightshade by shaking the anthers. Videographer: Georgios Nakas.