An ATM-PPM1D Circuit Controls the Processing and Restart of DNA Replication Forks

Yiting Cao[1#], Yingzheng Wang[1#], Jumana Badar[1#], Khoula Jaber[1], Vitor Marcel Faça[2], Tyler C.

Huang[1], Eliz Çelik[1], and Marcus B. Smolka[1*]

[1] Department of Molecular Biology and Genetics, Weill Institute for Cell and Molecular Biology,

Cornell University, Ithaca, NY, 14850, USA

[2] Department of Biochemistry and Immunology and Cell-Based Therapy Center, Ribeirao Preto

Medical School, University of Sao Paulo, Ribeirao Preto, SP, 14049-900, Brazil

# These authors contributed equally to this work.

*To whom correspondence should be addressed: Marcus B. Smolka, e-mail:

mbs266@cornell.edu

ABSTRACT

In response to DNA replication stress, DNA damage signaling kinases inhibit origin firing and

promote the remodeling and stabilization of replication forks, leading to a systemic reduction in

DNA synthesis, which collectively protects genomic integrity. Little is understood about the

regulatory mechanisms of replication stress recovery, including the mechanisms involved in the

restart of stalled replication forks. Here, we identify the oncogenic phosphatase PPM1D/WIP1

as a regulator of replication fork restart. During recovery from replication stress, PPM1D

prevents excessive MRE11- and DNA2-dependent nucleolytic degradation of stalled forks. Loss

of PPM1D function leads to defects in RAD51 recruitment to chromatin and impairs RAD51-

dependent fork restart. Phosphoproteomic analysis reveals that PPM1D regulates a network of

ATM substrates, several of which are phosphorylated at an $S/T-Q-(E/D)^n$ motif. Strikingly,

inhibition of ATM suppresses the deleterious consequences of impaired PPM1D function at

replication forks, enabling timely fork restart. The dominant effect of ATM hyper-signaling in

suppressing fork restart occurs, in part, through the excessive engagement of 53BP1 and

consequent RAD51 antagonization. These findings uncover a new mode of ATM signaling

responding to fork stalling and highlight the need for PPM1D to restrain ATM signaling and

facilitate proper fork restart.

INTRODUCTION

DNA replication occurs within a highly complex and dynamic genomic landscape. The progression of DNA replication forks is often slowed or impeded by various obstacles, such as DNA lesions or adducts, secondary DNA structures, transcriptional intermediates, compacted chromatin, and the shortage of nucleotides and other essential replication factors (*1*, *2*). When replication forks stall, they become vulnerable structures prone to degradation by nucleases, which can lead to the extensive single-stranded DNA (ssDNA) accumulation and/or the formation of double-stranded breaks (DSBs) (*3*). The ability of cells to maintain the integrity of stalled replication forks largely relies on the ATR kinase, a member of the phosphatidylinositol 3-kinase–related kinase (PIKK) family, which senses the exposure of ssDNA at replication forks (*4*). Upon fork stalling, ATR and its downstream effector CHK1 suppress late origin firing to prevent the exhaustion of key replication factors, including nucleotides, replication protein A (RPA), and the RFC/PCNA complex (*5*, *6*). In parallel, ATR signaling restrains the activity of nucleases, such as MRE11, EXO1, and MUS81, to limit excessive fork processing (*7–9*). Cells also rely on structural remodeling of replication forks to avoid nucleolytic degradation (*10*). For instance, a replication fork regresses into a four-way DNA junction, known as fork reversal, allowing nascent DNA strands to be stabilized by protective factors such as RAD51, BRCA1, and BRCA2 (*3*, *11*). This reversal process is catalyzed by DNA translocases, including SMARCAL1, HLTF, and ZRANB3 (*12*), and is tightly regulated by the DNA damage response kinases ATR and DNA-PKcs (*13*, *14*).

While fork reversal and protection are important for stabilizing replication forks, the restart of stalled replication forks is essential for the timely completion of DNA replication and the preservation of genomic integrity (*15*). Stalled forks that undergo regression can be restarted by branch migration, via the RECQ1 helicase, which is considered the primary choice for fork

restart given the minimal chance for error-prone DNA processing (*16*). Alternatively, reversed

forks can be restarted through regulated processing of the regressed ends by helicases and

nucleases, such as WRN and DNA2, enabling RAD51-mediated homologous recombination

(HR) transactions for restoring a functional replication fork (*17*). In addition, forks that collapse

may undergo HR-mediated repair transactions to restore a functional fork, albeit with an

increased risk of initiating mutagenic processes such as break-induced replication (BIR) (*18*).

Engagement of NHEJ-mediated repair into fork-associated DSBs also has the potential of

generating deleterious outcomes, such as chromosomal translocations, especially when single-

ended DSBs are generated (*19, 20*). Moreover, processes that prevent the stalling of a

replication fork, such as PrimPol-mediated repriming or translesion synthesis, can mitigate the

potential deleterious outcomes associated with stalled or collapsed forks, although they also

introduce mutations and other genomic instabilities (*21*). Overall, the ability of cells to restart

replication forks in a regulated and error-free manner is crucial to protect genome integrity upon

replication stress. Since replication fork restart mechanisms need to counteract the mechanisms

that stabilize stalled forks, including fork reversal and inhibition of nucleolytic processing, cells

must tightly control the balance of fork-stabilizing and fork-restarting mechanisms during

replication stress recovery.

While ATR is the central kinase in the response to stalled forks, the other PIKKs, ATM and

DNA-PKcs, were also reported to engage in the response to various forms of replication stress.

For example, in response to replication stress induced by the B-family DNA polymerase inhibitor

aphidicolin, ATM was shown to phosphorylate several proteins, including H2AX, Claspin,

BRCA1, MCM3, MCM6 and RIF1 (*22*), in a manner dependent on RAD18, WRNIP1, and

ATMIN (*23*). Consistent with its major role in the double-strand break (DSB) response, ATM is

robustly activated by agents that cause replication fork-associated DSBs, such as the

topoisomerase inhibitors camptothecin and etoposide (*24, 25*). Differences in the cellular

response to stalled versus broken replication forks have been characterized at the proteomic

level, which revealed a role for ATM in promoting HR-mediated repair specifically at broken

forks by impairing recruitment of the anti-resection BRCA1-A complex (*25*). DNA-PKcs, which

also has canonical roles in the DSB response, was shown to be recruited to both ongoing and

stalled replication forks, where it mediates drug-induced replication fork reversal and promotes

fork stability through mechanisms that remain unknown (*14, 26–28*).

Since PIKK signaling halts core cellular processes such as DNA replication and cell cycle

progression (*29, 30*), mechanisms for the deactivation of PIKK signaling must be in place to

enable proper recovery from replication stress. Multiple phosphatases have been implicated in

antagonizing PIKK signaling, including PP1, PP2A, PPM1G and PPM1D (*31–33*). The

phosphatase PPM1D (protein phosphatase Mn^{2+}/Mg^{2+}-dependent 1D) has been shown to play

prominent roles in dephosphorylating and antagonizing ATM, p53 and CHK2 (*34–37*). PPM1D is

overexpressed in multiple tumor types, including breast, ovarian, and glioblastoma tumors (*38*),

which can occur through gene amplification or truncating mutations that remove a degron at its

C-terminal domain (*39, 40*). Elevated PPM1D protein levels counteract DNA damage signaling

and are proposed to play a major role in reducing p53-mediated apoptosis (*39, 41, 42*). Not

surprisingly, PPM1D overexpression has been implicated in chemotherapy resistance (*39, 43*).

Here we find that PPM1D/WIP1 is required for the restart of stalled replication forks. Upon

recovery from replication stress, PPM1D prevents the excessive nucleolytic degradation of

stalled forks. Loss of PPM1D function leads to defects in RAD51 recruitment to chromatin and

impairs RAD51-dependent fork restart. Phosphoproteomic analysis reveals that PPM1D

regulates a network of proteins involved in ATM signaling, several of which are targeted by ATM

at an S/T-Q-(E/D)^n motif. Strikingly, inhibition of ATM suppresses the deleterious consequences

of impaired PPM1D function at replication forks, enabling timely fork restart. The dominant effect

of ATM hyper-signaling in suppressing fork restart occurs through the excessive engagement of

53BP1 and consequent RAD51 antagonization. These findings uncover a new mode of ATM

signaling responding to fork stalling and highlight the need for PPM1D to limit ATM signaling

and enable proper fork restart.

RESULTS

PPM1D is required for proper restart of stalled replication forks

In search for phosphatases required for proper fork restart, we focused on the oncogenic

phosphatase PPM1D (PP2Cδ/WIP1), since it was previously shown to antagonize PIKK

signaling (*34*, *36*, *44–47*). PPM1D is a monomeric phosphatase for which a potent and selective

inhibitor is available (GSK2830371; herein referred to as PPM1Di) (*48*), which allowed us to

assess the importance of PPM1D during recovery from replication stress. U2OS cells were

treated with hydroxyurea (HU) for 30 minutes, a condition previously reported to promote

replication fork stalling without causing fork collapse (*49*). Cells were then washed and released

in the presence or absence of PPM1Di, and DNA synthesis was assessed by monitoring the

incorporation of the nucleotide analog EdU (Fig. 1a). As shown in Figs. 1a-b, PPM1Di reduced

the ability of cells to incorporate EdU, suggesting a defect in replication fork restart and/or origin

firing. A similar result was observed in PPM1D-knockout U2OS cells, supporting that the

observed effects were not due to off-target inhibitor activity (Figs. 1c and S1a-b). DNA combing

assay in U2OS cells revealed that the reduction in EdU incorporation caused by PPM1Di was

associated with an increase in stalled forks and a reduction in fork restart (Figs. 1d and S1c).

Consistent results were observed using DNA fiber assays in both U2OS and HCT116 cells (Fig.

S1f). Additionally, both knockdown and knockout of PPM1D significantly increased the

frequency of stalled forks. Notably, in PPM1D knockout cells, siPPM1D neither increases fork

stalling nor decreases fork restart, supporting the specificity of the siPPM1D-mediated depletion

(Figs. 1e and S1d). PPM1Di did not reduce origin firing (Fig. S1f), consistent with a defect in

fork restart being the major cause of the observed reduction in DNA synthesis during recovery

from replication stress. This notion is also supported by the additive effect of PPM1Di in

combination with the CDC7 inhibitor, an inhibitor of origin firing, in reducing DNA synthesis (Fig.

S1g). Moreover, the overall abundance of PPM1D is increased in S-phase cells (Figs. 1f-g),

consistent with previously published results (*50*), and supporting a potential role for PPM1D in

DNA replication. Taken together, these results reveal a role for PPM1D in the control of

replication fork restart.

PPM1D prevents the nucleolytic processing of stalled replication forks during replication

stress recovery

To gain insights into how loss of PPM1D function impairs replication forks during replication

stress recovery, we monitored RPA32 foci as a proxy for ssDNA accumulation. U2OS cells were

treated with HU for 30 minutes and released in the presence or absence of PPM1D function. As

shown in Figs. 2a-d, both PPM1Di and PPM1D knockout caused an increase in RPA32 foci in

PCNA-positive cells. We also monitored the accumulation of ssDNA at nascent DNA strands

during release from HU arrest using BrdU labeling followed by non-denaturing

immunofluorescence and observed a significant increase in nascent ssDNA exposure in cells

treated with PPM1Di (Figs. 2e and S2a-b). We interpret the increased BrdU signal on nascent

DNA strands as likely originating from ssDNA at regressed replication forks. Consistent with this

notion, cells lacking SMARCAL1, a translocase involved in fork reversal, did not exhibit

significantly increased levels of RPA32 foci or stalled forks upon PPM1Di treatment during HU

release (Figs. 2f-g and S2c). Moreover, inhibition of the nucleases MRE11 or DNA2 suppressed

RPA32 foci accumulation (Figs. 2h-i), supporting the model that, upon PPM1Di treatment during

HU release, the action of these nucleases generates excessive ssDNA at stalled replication

forks. Notably, DNA2 seemed to play a more prominent role, with DNA2i completely rescuing

the effect of PPM1Di, consistent with previous reports showing that this is the key nuclease

involved in the hyper-processing of reversed replication forks (*17*). Overall, these results point to

a role for PPM1D in preventing the aberrant nucleolytic processing of stalled replication forks

during recovery from replication stress.

PPM1D antagonizes ATM signaling during recovery from fork stalling

To delineate the contribution of PPM1D to signaling downregulation during recovery from

replication stress, we compared the phosphoproteomes of mock-treated versus PPM1Di-treated

cells during release from HU arrest. The generated dataset was crossed with data from a

separate analysis comparing the phosphoproteomes of cells arrested by HU with the

phosphoproteome of cells that were HU-arrested and then released, yielding a dataset of

10,833 overlapping phosphosites identified in both phosphoproteomic experiments (Fig. 3a,

Table S1). PPM1Di caused extensive changes in signaling during replication stress recovery,

resulting in at least a two-fold increase in the abundance of about 18% of the phosphorylation

events found to be downregulated during release from HU arrest (Figs. 3a-b). While this finding

defines PPM1D as a prominent phosphatase acting during replication stress recovery, it also

implies that additional phosphatases must contribute to the dephosphorylation of other signaling

events downregulated during replication stress recovery. Notably, out of the phosphorylation

events upregulated by at least 2-fold upon treatment with PPM1Di, 22.5% contained a

phosphorylation event at the SQ/TQ consensus motif (Fig. 3c), suggesting that PPM1D displays

some level of selectivity for antagonizing PIKK signaling. Consistent with previous reports, we

found that PPM1D displays a preference for targeting phosphorylation at an $S/T-Q-(E/D)^n$ motif

(Fig. 3d) (*47, 51*). Notably, the phosphorylation events at $S/T-Q-(E/D)^n$ motifs most highly

regulated by PPM1D included several known substrates of the ATM kinase, such as RAP80,

NUMA1, SMC1A, and 53BP1 (*52–55*), suggesting that PPM1D plays a prominent role in

antagonizing ATM signaling. We confirmed this prediction by monitoring the phosphorylation of

KAP1 at serine 824, a well-established ATM substrate within the $S/T-Q-(E/D)^n$ motif (*56, 57*). As

shown in Figs. 3e-g, while PPM1D inhibition during release from HU treatment caused a drastic

increase in KAP1 phosphorylation, it led to a more subtle increase in CHK1 phosphorylation,

consistent with the model that PPM1D plays a more prominent role in antagonizing ATM

signaling compared to ATR signaling. Notably, we did not observe changes in the level of ATM

autophosphorylation (Fig. 3h), suggesting that PPM1D acts on ATM downstream substrates as

opposed to acting on ATM itself. Overall, our results establish PPM1D as a key phosphatase

antagonizing ATM signaling during replication stress recovery.

Unrestrained ATM signaling causes aberrant fork processing and fork restart defects

upon loss of PPM1D function

Since inhibition of PPM1D resulted in the increased phosphorylation of multiple ATM substrates

during replication stress recovery, we next tested whether inhibition of ATM could suppress the

defects we observed with fork restart and nucleolytic processing. PPM1Di alone did not cause a

strong reduction in EdU incorporation when compared to its effect during replication stress

release (Fig. S3a). Strikingly, inhibition of ATM during release from HU treatment fully rescued

the increased accumulation of RPA32 foci (Figs. 4a-b) and restored EdU incorporation in

PPM1Di-treated cells (Figs. 4c-d and S3a). ATMi also rescued the accumulation of stalled forks

and fork degradation caused by PPM1Di, as monitored by DNA fiber assays (Figs. 4e-f and

S3b-c). Furthermore, inhibition of ATM, but not ATR, was able to rescue the proliferation defect

caused by PPM1Di (Fig. 4g), reinforcing the hypothesis that PPM1D plays a key role in

selectively antagonizing ATM-dependent signaling. Taken together, these results support a

model in which PPM1D dephosphorylates ATM substrates during recovery from replication

stress, thereby preventing their toxic accumulation and the consequent hyperactivation of

nucleases whose action results in ssDNA accumulation and defects in fork restart.

PPM1D controls RAD51 chromatin recruitment and RAD51-mediated fork restart

To better characterize the effects of unrestrained ATM signaling during replication stress

recovery, we monitored RAD51, a recombinase protein with separable roles in protecting

reversed forks from degradation and mediating fork restart via homologous recombination (58).

Immunofluorescence-based analysis revealed that the number of RAD51 foci was significantly

reduced in cells treated with PPM1Di during recovery from replication stress (Figs. 5a-b). This

finding is consistent with a recent report showing that PPM1D inhibition suppresses RAD51 foci

formation in DIPG cells, as well as in U2OS cells following exposure to ionizing radiation (*59,*

60). In addition, chromatin fractionation also revealed a reduction of RAD51 in the chromatin-

bound fraction upon PPM1D inhibition during recovery from replication stress (Fig. 5c). We

hypothesized that PPM1D promotes RAD51-dependent fork restart during replication stress

recovery. To test this, we performed the DNA fiber assay to monitor persistent fork stalling in

cells treated with the RAD51 inhibitor B02, in the presence or absence of PPM1Di (Figs. 5d and

S4). Consistent with previous reports highlighting the importance of RAD51 for fork restart (*49*),

RAD51 inhibition alone increased the incidence of stalled forks. Cells co-treated with RAD51i

and PPM1Di during recovery from HU arrest did not display increased incidence of stalled forks

compared to cells treated with RAD51i alone. These findings suggest that PPM1D and RAD51

function through the same pathway to facilitate fork restart. The degree of fork stalling in

RAD51i-treated cells was higher than in PPM1Di-treated cells, congruent with a model whereby

PPM1D functions as a regulator of RAD51 roles in ensuring proper fork restart. Such regulation

is likely to involve PPM1D's role in antagonizing ATM signaling as inhibition of ATM prevents

reduction of RAD51 foci in PPM1D-KO cells (Fig. 5e). Additionally, we observed that PPM1D

inhibition caused a rapid increase in 53BP1 foci during the recovery from replication stress, and

that the increase was dependent on ATM (Fig. 5f). 53BP1 is a bona-fide ATM target (*61, 62*)

and one of the top PPM1D-regulated targets identified in our phosphoproteomic screen (Fig.

3d). Recent studies have shown that 53BP1 counteracts RAD51 loading at replication forks

independently of its role in antagonizing resection (*63*). Therefore, we reasoned that the

hyperphosphorylation of 53BP1 upon PPM1D inhibition causes its accumulation at stalled

replication forks, which in turn counteracts RAD51 engagement, thereby impairing proper fork

restart. Consistent with this model, PPM1D inhibition did not cause a decrease in RAD51

engagement in 53BP1-KO cells (Fig. 5g).

Increased phosphorylation status of ATM targets upon loss of PPM1D is not correlated

with an increase in detectable DSBs

ATM is canonically activated by DSBs (*64*). However, neutral comet assay did not detect

PPM1Di-dependent increase in DSBs within 30 minutes following recovery from replication

stress, a time point in which PPM1D inhibition resulted in a significant increase in the ATM-

dependent phosphorylation of KAP1 at serine 824 (Figs. 6a-c). These findings suggest that

ATM signaling and the observed increases in the phosphorylation status of ATM substrates are

independent of DSB formation during early replication stress recovery. Our finding that the

autophosphorylation of ATM is not increased during release from replication stress is also

consistent with this notion (Fig. 3h). To provide further evidence that PPM1D inhibition is not

causing strong induction of DSBs early (~30 minutes) during recovery from replication stress,

we engineered a ProKAS biosensor (*65*) containing two sensors that differentiate ATM signaling

regulated by PPM1D (ATM-PPM1D sensor) from ATM signaling that is not controlled by PPM1D

(ATM sensor) (Figs. 6d-e). As shown in Figs. 6f and S5, CPT robustly increased the

phosphorylation of both sensors, consistent with the canonical role of ATM in responding to

DSBs. The ATM sensor did not report increased ATM activity during recovery from replication

stress, in the absence or presence of PPM1D inhibitor, further supporting the absence of DSB-

induced ATM signaling under these conditions. Notably, the ATM-PPM1D sensor reported a

drastic increase in ATM signaling upon PPM1D inhibition during recovery from replication

stress. These findings with the biosensor are congruent with the model that PPM1D inhibition is

not causing DSBs, and suggest that in the context of replication fork stalling and restart, ATM

may be activated by alternative DNA structures, such as the nascent DNA ends of reversed

forks.

DISCUSSION

The ability of cells to restart stalled replication forks is crucial for the faithful completion of DNA replication under conditions of replication stress. Despite its importance, the regulatory mechanisms that govern fork restart remain incompletely understood. Here, we identify the phosphatase PPM1D as a regulator of replication fork recovery. Our findings reveal that attenuation of ATM-mediated signaling is a central function of PPM1D during replication stress resolution. In our working model (Fig. 6g), ATM is activated at stalled forks and phosphorylates substrates such as 53BP1, which in turn interferes with RAD51 loading. Persistent ATM activity antagonizes HR-mediated fork restart and also leads to fork degradation due to reduced RAD51-mediated fork protection. PPM1D counteracts the block in RAD51 loading by dephosphorylating ATM targets. Thus, the ATM–PPM1D axis acts as a dynamic rheostat that modulates ATM signaling at stalled forks to coordinate the balance between fork protection and fork restart.

A key unresolved question is how PPM1D is recruited to stalled forks. It is possible that the build-up of ATM-dependent phosphorylation events triggers PPM1D recruitment, establishing a feedback mechanism to restrain excessive ATM signaling at forks. Future studies will be needed to define the mechanisms driving PPM1D localization and whether recruitment is influenced by specific post-translational modifications, replisome-associated factors, or fork-associated DNA structures. Moreover, our results suggest that PPM1D primarily acts on ATM substrates rather than directly on ATM itself or ATR substrates. This is consistent with PPM1D's preference for dephosphorylating S/T-Q motifs followed by acidic residues [S/T-Q-(D/E)n motifs], a motif also preferred by ATM (66).

While ATM canonically responds to DSBs, our work suggests that ATM can also respond to

replication stress independently of DSB formation. Under conditions of PPM1D inhibition, we

observe elevated ATM signaling without detectable DSBs by neutral comet assay. Additionally,

ProKAS biosensor analyses show that the branch of ATM signaling regulated by PPM1D is

selectively induced during replication stress recovery, while other branches remain unchanged.

Although the precise DNA structures activating ATM at stalled forks remain unclear, we favor

the model that ATM senses unprocessed DNA ends at reversed forks. Nevertheless, given

previous reports of ATM activation in response to altered chromatin states or oxidative stress

(*67*, *68*), we cannot exclude the possibility that chromatin remodeling or oxidative stress during

replication stress recovery also contributes to ATM activation.

Another outstanding question is whether ATM signaling at stalled forks plays beneficial roles in

maintaining genome stability or supporting replication.One possibility is that early ATM-

mediated 53BP1 recruitment to stalled forks, and the consequent suppression of RAD51

loading, allows nucleases such as MRE11 and DNA2 to perform limited resection during

recovery from replication stress. This limited ssDNA production is needed for subsequent HR-

mediated fork restart.Therefore, PPM1D is needed later in the recovery process to attenuate

ATM signaling, reduce 53BP1 engagement and increase RAD51 loading, facilitating the

transition from fork processing to HR-mediated fork restart. Importantly, the ability of 53BP1 to

antagonize RAD51 loading, a role for 53BP1 also recently demonstrated by others (*63*), may

prevent inappropriate or premature recombination events that could cause chromosomal

rearrangements. Thus, by orchestrating these dynamic regulatory steps, the ATM–PPM1D axis

could ensure a timely, error-free, and balanced coordination of fork protection, processing, and

HR-mediated restart. Future investigations will be required to define the spatio-temporal

regulation of PPM1D and ATM action at replication forks.

Alternatively, ATM activation at stalled forks may largely reflect an unintended response to

replication intermediates resembling DSBs, such as the DNA end of a regressed fork arm. In

this scenario, unchecked ATM signaling may represent a maladaptive response, with PPM1D

acting as a safeguard to dampen inappropriate ATM signaling and enable proper fork restart

and recovery from replication stress. In the future, it will be important to better characterize the

impact of PPM1D and ATM inhibition on the accumulation of replication stress-induced genomic

instabilities, such as chromosomal aberrations. Detection of HR-driven chromosomal

rearrangements upon ATM inhibition would point to an important role for ATM signaling in

promoting HR quality control during replication stress responses. Moreover, delayed RAD51

loading and impaired HR-mediated fork restart in the absence of PPM1D may lead to increased

utilization of error-prone pathways, such as non-homologous end joining (NHEJ) or

microhomology-mediated end joining (MMEJ), resulting in elevated frequencies of chromosomal

translocations and radial chromosomes, which can be revealed by metaphase spread analysis.

These insights have important implications for cancer biology. PPM1D is frequently

overexpressed or mutated in cancers, often leading to elevated phosphatase activity. Such

dysregulation may enhance cancer cell survival under endogenous or therapy-induced

replication stress by facilitating fork restart and recovery (*59*). Pharmacological targeting of

PPM1D could synergize with therapies that impair fork remodeling or HR, offering potential

strategies to selectively eliminate tumor cells with high levels of replication stress. Going

forward, it will be important to assess the generality of PPM1D's role in fork restart across

different cell types and biological contexts. Of note, the cell lines used in this study, such as

U2OS and HCT116, carry hyper-stabilized forms of PPM1D. It is possible that the dependency

on PPM1D varies with the level of ATM signaling, with cells exhibiting higher ATM activity

relying more heavily on PPM1D for efficient fork restart, thus exerting selective pressure for

mutations that stabilize or increase PPM1D expression.

In conclusion, this study defines a function for the ATM–PPM1D signaling axis in regulating

replication fork restart. By limiting ATM activity at stalled forks, PPM1D enables RAD51

recruitment and HR-mediated fork restart and recovery from replication stress. These findings

underscore the importance of dynamic phosphorylation control at replication forks and establish

PPM1D as a central player in replication stress response pathways with broad relevance to

cancer biology and therapeutic intervention.

METHODS

Cell lines and cell culture

All cells were maintained at 37 °C with 5% CO_2 in a humidified incubator and cultured in

Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% bovine calf serum

(BCS), 1% non-essential amino acids, 100 U/mL penicillin, and 100 µg/mL streptomycin. U2OS

and HCT116 cells were obtained from the American Type Culture Collection (ATCC). Parental

U2OS and U2OS-PPM1D-KO cells were kindly provided by Macurek lab. Parental U2OS and

U2OS-SMARCAL1-KO cells were kindly provided by Vindigni lab. Parental U2OS and U2OS-

53BP1-KO cells were kindly provided by Durocher lab. All cell lines were regularly tested for

mycoplasma contamination using the Universal Mycoplasma Detection Kit (ATCC).

Drugs and reagents

Hydroxyurea (HU, Sigma-Aldrich, #H8627), PPM1D inhibitor (GSK2830371, Sigma-Aldrich,

#SML1048), CDC7 inhibitor (XL413, Selleck Chemicals, #S7547), DNA2 inhibitor (C5, MCE,

#HY-128729), MRE11 inhibitor (Mirin, Selleck Chemicals, #S8096), ATM inhibitor (AZD0156,

MedChemExpress, #HY-100016), ATR inhibitor (AZD6738, MedChemExpress, #HY-19323),

and RAD51 inhibitor (B02, Selleck Chemicals, #S8434) were used as indicated in the

experiments. Sodium L-ascorbate (Sigma-Aldrich, #A7631) and azide-fluor 488 (Sigma-Aldrich,

#760765) were used for the EdU incorporation assay. Dharmacon siGENOME non-targeting

control siRNA (siCTL, Horizon Discovery, #D-001210-02-05) and siGENOME human PPM1D

(siPPM1D, Horizon Discovery, #D-004554-01-0002) were used for siRNA transfection. 5-

Bromo-2'-deoxyuridine (BrdU, Sigma-Aldrich, #10280879001), 5-Iodo-2'-deoxyuridine (IdU,

Sigma-Aldrich, #I7125), and 5-Chloro-2'-deoxyuridine (CldU, Sigma-Aldrich, #C6891) were used

for DNA fiber and combing assays. 2-morpholinoethanesulfonic acid (MES, GoldBio,

#M095250), N-lauroylsarcosine sodium salt (Sarkoysl, Sigma-Aldrich, #L5125), and ß-agarase

(NEB #M0392L) were used for DNA combing assay. Propidium Iodide (PI, Thermo Fisher

Scientific, #P3566) and antifade mounting media with DAPI (Vectashield, #H-1200-10) were

used for flow cytometry and imaging experiments as specified.

Antibodies

Primary antibodies include: β-actin (Proteintech, #66009-1-Ig), β-tubulin (Proteintech, #66240-1-

Ig), Histone H4 (Bethyl Laboratories, #A300-647A), KAP1-pS824 (Bethyl Laboratories, #304-

146A), Gamma-H2AX (Bethyl Laboratories, #A300-081A), RPA32 (Bethyl Laboratories, #A300-

244A), PCNA (Santa Cruz Biotechnology, #sc-56), CHK1-pS345 (133D3, Cell Signaling

Technology, #2348), PPM1D/WIP1 (Santa Cruz Biotechnology, #sc-376257), ATM-pS1981

(Cell Signaling Technology, #4526), 53BP1 (Cell Signaling Technology #4937), RAD51

(Millipore Sigma, #PC130), Rat anti-BrdU (BU1/75 (ICR1) (Abcam, #6326), Mouse anti-BrdU

(B44) (BD Biosciences, #347580). Secondary antibodies include: HRP-conjugated goat anti-

mouse IgG (Invitrogen #31430), HRP-conjugated donkey anti-rabbit IgG (Cytiva, #NA934V),

Alex Fluor 594 goat anti-mouse IgG (H+L) (Invitrogen, #A11005), Alexa Fluor 555 F(ab')2-Goat

anti-rabbit IgG (H+L) (Invitrogen, #A21430), Alexa Fluor 488 goat anti-rabbit IgG (H+L)

(Invitrogen, #A11008), Alexa Fluor 488 goat anti-mouse IgG (H+L) (Invitrogen, #A11001).

Immunofluorescence

For immunofluorescence experiments, cells (25,000 per well) were seeded onto coverslips in

12-well plates. The following day, cells were treated as indicated. Cells were then washed twice

with 1 mL ice-cold 1x PBS. Cells were pre-extracted with ice-cold buffer (25 mM HEPES, pH

7.5, 50 mM NaCl, 1 mM EDTA, 3 mM $MgCl_2$, 300 mM sucrose, 0.5% Triton X-100) for 5

minutes, followed by a PBS wash. Fixation was performed with 4% paraformaldehyde in PBS

for 10 minutes at room temperature (RT), followed by three 5-minute washes in PBS. Cells were

permeabilized with 0.25% Triton X-100 in PBS for 5 minutes and washed three times in PBS.

Cells were blocked with 5% BSA in 1x PBS for 1 hour at RT. Coverslips were then incubated

with primary antibodies (1:300 dilution in 5% BSA) for 1 hour at RT or overnight at 4°C. After

three washes with 1x PBS, cells were incubated with fluorophore-conjugated secondary

antibodies (1:500 dilution in 5% BSA) for 1 hour at RT in the dark. Following three final PBS

washes, coverslips were mounted with DAPI-containing antifade mounting medium, sealed with

nail polish, and stored at 4°C. Images were acquired using a Leica DMi8 inverted fluorescence

microscope and analyzed using Fiji/ImageJ. Statistical analyses were performed using

GraphPad Prism v9.

EdU incorporation assay

For the EdU incorporation experiment, U2OS cells (25,000 cells per well) were seeded onto

coverslips in 12-well plates. The following day, cells were treated as indicated and then

processed until permeabilization as immunofluorescence assay. Then, cells were washed by

3% BSA in 1x PBS twice at room temperature (RT). Click reactions were performed with pre-

mixed reagents (10 mM Na-L-Ascorbate; 0.1 mM Alexa Fluor Azide; 5 mM $CuSO_4$) for 30

minutes at RT. Cells were washed by 3% BSA in 1x PBS twice at RT. Cells were blocked and

incubated with primary and secondary antibodies as indicated in the immunofluorescence

assay. Following three final 1x PBS washes, coverslips were mounted with a DAPI-containing

antifade mounting medium, sealed with nail polish, and stored at 4°C. Images were acquired

using a Leica DMi8 inverted fluorescence microscope and analyzed using Fiji/ImageJ. Statistical

analyses were performed using GraphPad Prism v9.

DNA fiber assay

For DNA fiber experiments to monitor replication fork restart, cells were first labeled with 25 µM

CldU for the indicated time, then treated with 4 mM HU for 30 minutes to induce fork stalling.

After two washes with prewarmed media, cells were incubated in fresh warm media containing

250 μM IdU and 10 μM PPM1D inhibitor, with or without additional inhibitors, for the indicated

time. For the fork protection assay, cells were sequentially labeled with 25 μM CldU for 30

minutes and 250 μM IdU for 30 minutes, followed by treatment with 2 mM HU and/or 10 μM

PPM1Di or 100 nM ATMi for 90 minutes. Following treatment, cells were harvested by

trypsinization, washed, and resuspended in 1x PBS. Approximately 1,000 cells were pipetted

onto clean glass microscope slides and lysed with DNA fiber lysis buffer (200 mM Tris-HCl at

pH 7.4, 50 mM EDTA, 0.5% SDS) for 2 minutes. Slides were immediately tilted at a 15º angle to

stretch DNA fibers, air-dried for 5 minutes, and fixed in a 3:1 methanol/acetic acid solution for 10

minutes at RT. DNA fibers were denatured in 2.5 M HCl for 1 hour, neutralized with 0.4 M Tris-

HCl (pH 7.4) for 5 minutes, and washed three times in 1xPBS for 5 minutes each. Slides were

blocked for 1 hour in PBS containing 10% goat serum and 10% BSA. Slides were then

incubated with mouse anti-BrdU (BD Biosciences, #347580, 1:50) and rat anti-BrdU (Abcam,

#ab6326, 1:200) antibodies in PBS with 3% BSA for 1 hour at 37 °C in a humidified chamber.

After three 5-min PBS washes, slides were incubated with Alexa Fluor 488-conjugated anti-

mouse and Alexa Fluor 594-conjugated anti-rat secondary antibodies (1:500) for 1 hour at RT.

Slides were further washed three times in PBS, mounted with Fluoromount-G (Thermo Fisher

Scientific), and sealed with nail polish. Images were acquired using a Leica DMi8 inverted

fluorescence microscope and analyzed with Fiji/ImageJ software. For fork restart analysis,

ongoing forks, bi-directional forks, and terminating forks are classified as restarted forks. The

percentages of stalled forks, restarted forks, and newly fired forks were calculated accordingly.

Representative images of the different fork types are shown in Fig. S1e. Statistical analyses

were performed using GraphPad Prism v9.

Western blotting

Cells were lysed in ice-cold modified RIPA buffer (50 mM Tris-HCl, pH 7.5; 150 mM NaCl; 1%

Tergitol; 0.25% sodium deoxycholate; 5 mM EDTA) supplemented with complete EDTA-free

protease inhibitor cocktail (Roche), 1 mM PMSF, and 5 mM NaF. Lysis was performed for 30

minutes at 4 ºC and sonicated. Protein concentration was determined using the Bradford assay

(Bio-Rad), and 10-15 µg of total protein was resolved by SDS-PAGE. Proteins were transferred

to polyvinylidene difluoride (PVDF) membranes and blocked with 5% non-fat milk in TBST (Tris-

buffered saline with 0.05% Tween-20) for 1 hour at RT. Membranes were incubated with

primary antibodies overnight at 4 ºC. The following day, membranes were washed three times

with TBST and incubated with HRP-conjugated secondary antibodies for 1 hour at RT. After

three additional washes with TBST, membranes were developed using Clarity Western ECL

substrate (Bio-Rad, #1705060). Chemiluminescent signals were acquired using a ChemiDoc

Imaging System (Bio-Rad) and analyzed with ImageLab software (Bio-Rad).

Chromatin fractionation

Cells were harvested by trypsinization, washed with PBS, and resuspended in Solution A (10

mM HEPES, pH 7.9; 10 mM KCl; 1.5 mM $MgCl_2$; 0.34 M sucrose; 10% glycerol; 0.1% Triton X-

100; 1 mM DTT; 10 mM NaF; 1x protease inhibitor cocktail (ThermoFisher, #A32965); 1x

PMSF). Cells were incubated on ice for 8 min, then centrifuged at 1,300 x g for 5 minutes at 4

ºC. The supernatant was harvested as the cytosolic fraction. The pellet was washed once with

solution A, then resuspended in solution B (3 mM EDTA; 0.2 mM EGTA; 1 mM DTT; 1x

protease inhibitor cocktail) and incubated on ice for 30 minutes. After centrifugation at 1,700 x g

for 5 minutes at 4 ºC, the supernatant was collected as the nucleoplasmic fraction. The

remaining pellet was washed once with Solution B and resuspended in the 2x sample buffer for

analysis as chromatin fraction. Next, chromatin-associated proteins were analyzed by Western

blotting.

Clonogenic assay

U2OS cells (800 cells per well) were seeded in 6-well plates and treated with the indicated

drugs for 10 days. At the end of the experiment period, cells were washed once with 1x PBS,

fixed in 3.7% paraformaldehyde for 10 minutes at RT, and stained with 0.01% crystal violet in 1x

PBS solution. Plates were air-dried and scanned, and colony areas were quantified using

ImageJ software.

Neutral comet assay

Cells were treated as indicated and harvested by trypsinization, followed by a wash with ice-cold

1x PBS. Cells were resuspended at a concentration of $1x10^5$/mL and mixed with molten

CometAssay® LMAgarose (R&D Systems, #4250-050-02) at a 1:10 (v/v) ratio. A 50 µL aliquot

of the mixture was immediately applied onto CometSlide® (R&D Systems, #4250-050-03) and

allowed to solidify at 4 ºC for 10 min. Slides were then immersed in lysis solution (R&D

Systems, #4250-050-01) for 1 hour at 4 ºC. Following lysis, electrophoresis was carried out

under neutral conditions at 1V/cm for 30 minutes at 4 ºC in the dark. Slides were then incubated

in DNA precipitation solution (1M ammonium acetate in 82% ethanol) for 30 minutes at RT,

followed by a 30-minute incubation in 70% ethanol. Slides were air-dried at 37 °C and stained

with SYBR™ Gold (Thermo Fisher Scientific, #S11494). Images were captured at 20x

magnification using a Leica DMi8 inverted fluorescence microscope. DNA damage was

assessed by measuring the comet tail moment using OpenComet software (v1.3.1).

Sample preparation for phosphoproteomic analysis

HCT116 cells were cultured for at least five passages in either "light" or "heavy" SILAC DMEM

medium (Thermo Fisher Scientific, #88364) supplemented with 10% dialyzed FBS (Sigma-

Aldrich, #F0392), 100 U/mL penicillin, and 100 µg/mL streptomycin (Gibco, #15140-122).

Labeling conditions were inverted in biological replicates to filter out phosphopeptides

influenced by isotopic bias. After experimental treatments, cells were washed with ice-cold

1xPBS and harvested using a cell scraper. Cell pellets were collected by centrifugation at 1,000

x g for 5 minutes at 4 °C, washed once in PBS, and stored at -80 °C overnight. Frozen pellets

were thawed and resuspended in a hypotonic buffer (20 mM Tris-HCl, pH 7.4; 10 mM NaCl; 3

mM MgCl$_2$; 0.5 mM DTT) and incubated on ice for 20 min. Cells were centrifuged at 4,500 x g

for 4 minutes at 4 °C and the supernatant was discarded. Pellets were resuspended in lysis

buffer (50 mM Tris-HCl, pH 8.0; 0.2% Tergitol; 150 mM NaCl; 5 mM EDTA) supplemented with

protease inhibitors (ThermoFisher, #A32965), 1 mM PMSF, and 1x PhosSTOP phosphatase

inhibitor cocktail (Roche). Lysis proceeded on ice for 30 min, followed by sonication using a

Branson 450 digital sonifier with a 102-C probe (15% amplitude, 3s on/ 2s off pulses) for 50

seconds. Lysates were clarified by centrifugation at 45,000 x g for 30 minutes at 4°C.

Protein concentrations were determined using the Bradford assay (BioRad). Equal amounts (2

mg each) of light- and heavy-labeled lysates were combined. Samples were denatured and

reduced with 1% SDS and 5 mM DTT at 65 °C for 10 minutes. Cysteines were alkylated with 25

mM iodoacetamide for 15 min in the dark**.** Proteins were precipitated by adding 3 volumes of

cold precipitation solution (50% acetone, 49.9% ethanol, 0.1% acetic acid) and incubated on ice

for at least 20 minutes. Pellets were recovered by centrifugation at 4,500 x g for 5 minutes at

4°C, washed once with 8 M urea solution and deionized water, and centrifuged at 13,000 x g for

1 minute. Residual water was carefully removed without disturbing the pellet. Pellets were

resuspended in 400 µL of 8 M urea in 100 mM Tris-HCl (pH 8.0), then diluted to 2 M urea by

adding 1,200 µL of Tris/NaCl solution (50 mM Tris-HCl, pH 8.0; 150 mM NaCl). Proteins were

digested overnight at 37 °C with 4 µg of TPCK-treated trypsin. Peptides were acidified to a final

concentration 0.2% trifluoroacetic acid (TFA) and formic acid (FA), and desalted using a 200 mg

Sep-Pak C18 cartridge (Waters). Briefly, the C18 column was pre-conditioned with buffer B

(80% acetonitrile, 0.1% TFA) followed by buffer A (0.1% TFA). Samples were loaded, washed

with buffer C (0.1% acetic acid), and eluted with buffer D (80% acetonitrile, 0.1% acetic acid),

then dried in a SpeedVac.

Phosphopeptides were enriched using the High-Select Fe-NTA phosphopeptide enrichment kit

(Thermo Fisher Scientific, #A32992) following the manufacturer's protocol. Eluted

phosphopeptides were immediately dried and resuspended in 0.1% TFA, followed by a clean-up

using a small-scale C18 spin column (Thermo Fisher Scientific, #89879) packed with C18 slurry.

Columns were conditioned with buffer B and A, samples loaded, washed with 3% acetonitrile in

buffer A, and eluted with buffer D. Eluted peptides were dried in a SpeedVac.

Dried phosphopeptides were resuspended in 15 µL LC-grade H_2O, 10 µL 10% FA, and 60 µL

HPLC-grade acetonitrile. A total of 80 µL was injected into an UltiMate 3000 system and

fractionated by HILIC (hydrophilic interaction liquid chromatography) using a TSKgel Amide-80

column (2 mm × 150 mm, 5 µm; Tosoh Bioscience). The following gradient buffers were used:

buffer A (90% acetonitrile), buffer B (75%acetonitrile and 0.005% TFA), and buffer C (0.025%

TFA). The gradient program was as follows: 100% buffer A (0 min); 94% buffer B/6% buffer C (3

min); 65.6% buffer B/34.4% buffer C (30 minutes); and 5% buffer B/95% buffer C (32 - 40

minutes). Flow rate: 150 µL/min. Fractions were collected every 30 seconds from minute 5 to

minute 24. All fractions were dried, resuspended in 5 µL of 0.1P (1 pmole/µL angiotensin in

0.1% TFA), and subjected to LC-MS/MS analysis.

ProKAS biosensor sample preparation

Stable U2OS cell lines expressing the ProKAS construct were generated using lentiviral

transduction, followed by selection with 1 µg/mL puromycin. Following treatment, cells cultured

in 6-well tissue culture plates were lysed using a modified radioimmunoprecipitation assay

buffer (mRIPA) buffer supplemented with 0.2% SDS (SDS lysis buffer). After aspirating the

culture media, 175 µL of SDS lysis buffer was added to each well. Plates were placed on a hot

plate set to 200°C for 10 seconds and immediately removed and agitated to promote the lysis.

This heating and agitation step was repeated 2-3 times to ensure complete lysis. Following

lysis, 525 µL of mRIPA buffer was added to each well to dilute the final SDS concentration to

below 0.1%. Lysates were collected into pre-labeled tubes and sonicated using a Branson 1/8"

probe tip sonifier (5 seconds at 12% amplitude) to shear genomic DNA and reduce viscosity.

Samples were then centrifuged at 16,000 x g for 3 minutes at 4°C to pellet debris. For affinity

purification, 4 µL of ALFA magnetic resin (NanoTag, #N1516) was used per lysate. Resin was

equilibrated by resuspending in 1 mL mRIPA buffer, and then distributed evenly to LoBind

Eppendorf tubes placed on magnetic racks. Afterwards, the buffer was aspirated, and the

lysates were transferred to the tubes containing the immobilized resin. The samples were

incubated for 30 minutes at 4°C on a nutating mixer. After incubation, beads were immobilized

on magnetic racks and supernatants were aspirated. Beads were washed twice with 500 µL

mRIPA buffer and once with 1 mL sterile deionized water, with each wash involving nutation for

1 minute at RT. The beads were then immobilized and 80 µL of acidic elution buffer (0.1 M

Glycine-HCl, 150 mM NaCl, pH 2.2) was added to each tube. The tubes were agitated for 2

minutes at RT to aid in elution of the bait protein. Eluted material was collected into new LoBind

Eppendorf tubes. For digestion, each elution was adjusted by adding 20 µL of 1M Tris (pH 8.0),

50 µL of 8 M Urea, and 50 µL of 150 mM NaCl-50 mM Tris (pH 8.0) containing 100 ng Trypsin

Gold (Promega, #V5280). Samples were incubated at 37 °C on a nutating mixer for 6-18 hours

to allow tryptic digestion of immunoprecipitated proteins. Peptide samples were acidified by

adding 10 µL of 10% Trifluoroacetic acid (TFA) and 10 µL of 10% Formic Acid. Acidified digests

were desalted using 20 mg of C18 resin (Sep-Pak cartridge; Waters, WAT054945) packed into

Pierce Micro Spin Columns (ThermoFisher Scientific, #89879). The resin was conditioned with

200 µL of 80% acetonitrile (ACN) containing 0.1% TFA, followed by 200 µL 0.1% TFA. Acidified

samples were then loaded onto the resin, washed with 350 µL of 0.1% acetic acid, and peptides

were eluted into LoBind Eppendorf tubes using 100 µL of 80% ACN containing 0.1% acetic

acid. Eluted peptides were then dried in a vacuum concentrator and immediately resuspended

in 10 µL of 250 mM HEPES (pH 8.0) for tandem mass tag (TMT) labeling. A 50 µg aliquot of

TMTsixplex reagent (ThermoFisher Scientific, #90061) was used to label up to 6 different

sample conditions. For each TMT channel, 2 µL of ACN was added to the reagent, followed by

the immediate addition of 8 µL of resuspended peptide. Labeling reactions were incubated for

60 minutes at RT and quenched by adding 2 µL of 5% hydroxylamine. Subsequently, 25 µL of

0.1% TFA was added to each channel. Samples from all channels were then pooled, and an

additional 30 µL of 10% TFA was added to the combined sample. The pooled TMT-labeled

peptides were desalted again using C18 cleanup as described above. The dried samples were

resuspended in LC-grade water prior to MS injection.

DNA combing assay

For the DNA combing assay, 200,000 cells were seeded and cultured for 24 hours before

labeling. Cells were first labeled with 25 µM CldU for 20 minutes, followed by treatment with 4

mM HU for 30 minutes. Cells were washed twice with pre-warmed 1x PBS and incubated in pre-

warmed complete medium containing 250 µM IdU, with or without 10 µM PPM1Di for 40

minutes. Cells were harvested and resuspended in 1.2 % low-melting-temperature (LMT)

agarose (Sigma-Aldrich, #A4018) pre-warmed to 55 ºC and allowed to solidify at 4 ºC for 1 hour.

Cell-containing agarose was incubated in ESP solution (200 µL 0.5 M EDTA, pH 8.0; 25 µL 10%

Sarkosyl; 50 µL Proteinase K) at 50 ºC for 16 hours. The ESP solution was then discarded, and

cells were washed six times with a modified TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0).

Agarose plugs were melted in 1 mL MES buffer (1.22 g MES dissolved in 12.5 mL ddH$_2$O, pH

5.5) at 68 ºC for 20 minutes and further incubated at 42 ºC for 20 minutes. Subsequently, 1.0 µL

ß-agarase was added to each sample and digestion was performed at 42 ºC for 12 hours. DNA

solutions were transferred into disposable reservoirs (Genomic Vision) and loaded onto the

molecular combing system (Genomic Vision). Functionalized microarray coverslips (PolyAn,

#10490804) were used for DNA stretching. After combing, coverslips were baked at 65ºC for 2

hours and fixed in methanol for 10 minutes at RT. DNA fibers were denatured, blocked,

incubated with primary and secondary antibodies as described for the DNA fiber assay. Images

were acquired and analyzed as DNA fiber assay. Representative images of different combing

conditions are shown in Fig. 1d.

siRNA transfection

For siRNA transfection performed in EdU incorporation assay, 10,000 cells per condition were

seeded on 6cm plates and incubated at 37 °C until 60%-70% confluent. Then, solution A (3 µL

of siRNA (20 pmols) in 250 µL Opti-MEM) and solution B (9 µL of Lipofectamine in 250 µL Opti-

MEM) were incubated for 5 minutes at room temperature before mixing. After incubating for 20

minutes, the transfection mixture was added drop-wise on cells. Transfection reagents were

removed after 6 hours and cells were cultured with complete medium for at least 48 hours,

before being treated as indicated in each experiment.

MS data acquisition

Phosphoproteomics

Phosphopeptide fractions were resuspended in 0.1% trifluoroacetic acid and analyzed by liquid

chromatography-tandem mass spectrometry (LC-MS/MS) using an UltiMate 3000 HPLC system

coupled to a Q-Exactive HF mass spectrometer (Thermo Fisher Scientific). Peptides were

separated on a 35 cm in-house packed C18 column (inner diameter of 100 µM) using reversed-

phase chromatography. The Q-Exactive HF was operated in data-dependent acquisition (DDA)

mode. Full MS scans were acquired in the Orbitrap over a mass-to-charge range of 380-1800

m/z at a resolution of 60,000. The 20 most abundant precursor ions with charge states of +2,

+3, and +4 were selected for MS/MS fragmentation using higher-energy collisional dissociation

(HCD) with a normalized collision energy (NCE) of 28. Fragment ion spectra were acquired in

the Orbitrap at a resolution of 15,000 with an AGC target of 1×10^5 and a maximum injection time

of 100 ms. Dynamic exclusion was enabled with a 30-second exclusion window to prevent

repeated sequencing of previously fragmented ions.

ProKAS Sensor

To check the TMT labeling efficiency and retention time of labeled peptides, data-dependent

analyses were performed on the Q Exactive HF mass spectrometer using a 70-minute method

containing a 60,000 scanning resolution, AGC target of 3×10^6, maximum inject time of 30 ms,

and scan range between 380 and 1800 m/z. The MS2 parameters included a 15,000 resolution,

AGC target of 1×10^5, maximum inject time of 100 ms, isolation window of 2.0 m/z and

normalized collision energy (NCE) of 36. Once the mass of target peptides and their

phosphorylated forms were identified, parallel reaction monitoring was performed utilizing a 70-

minute method with 15,000 resolution, AGC of 2×10^5, an inject time of 120 ms, and an isolation

window of 0.4 m/z. Labeling efficiency was determined to be between 85-90%, and parallel

reaction monitoring libraries were determined based on Comet searches to design an inclusion

list with peptide masses.

Phosphoproteomic data analysis

Raw LC-MS/MS data were analyzed using the Comet search engine. The following search

parameters were applied: precursor mass tolerance of ±10 ppm; semi-tryptic digestion; variable

modifications of +8.0142 Da (Lysine, heavy SILAC), +10.00827 Da (Arginine, heavy SILAC),

+79.966331 Da (phosphorylation of Ser/Thr/Tyr), and +57.021465 (carbamidomethylation of

cysteine). Searches were performed against the Uniprot human proteome database (74,304

entries). Peptide identification was refined using PeptideProphet, with the following parameters:

minimum probability of 0.05, minimum peptide length of 7 residues, accurate mass binning,

charge states restriction to +2, +3, and +4, and phospho-site scoring enabled ("use Phospho-

information"). Peptide quantification was performed using Xpress, with a mass tolerance of

0.005 Da, requiring a minimum of one point for quantitation and one isotopic peak per peptide.

Phosphosite localization confidence was determined using PTMProphet. In-house R-scripts

were used to further process the data, including: grouping adjacent phosphorylatable residues

into phosphosite clusters when exact localization could not be determined; calculating the

average and median Xpress ratio for each peptide; computing statistical significance using the

Mann-Whitney U test.

ProKAS Data Analysis with Skyline

To construct the spectral libraries utilized for ProKAS parallel reaction monitoring, raw MS/MS

data generated from the DDA runs were searched on the Comet search engine to obtain

peptide masses. The search was conducted using the following parameters: semi-tryptic

peptide identification, mass tolerance within 15 ppm, variable modifications for STY

phosphorylation (79.966331 Da). The search was performed against the Uniprot human

proteome database in addition to ProKAS peptide sequences, as well as the TMT condition.

PTMprophet was utilized to determine phosphorylation site localization, search results were

imported into Skyline, containing masses of the unphosphorylated and phosphorylated forms of

the target ProKAS peptides. Once a library was generated, PRM data was imported into Skyline

for analysis by conversion to mzXML format using msConvert. Transitions were verified

manually before generating a transitions report from Skyline containing the Protein, Peptide

Modified Sequence, Replicate, Precursor Mz, Precursor Charge, Product Mz, Fragment Ion,

Retention Time, and Area. Data in these reports were processed for each TMT fragment ion in

each replicate by normalization of phosphorylated peptide ratios to their unphosphorylated

counterparts. Then, the normalized ratios in each condition were compared against the

normalized ratios in the untreated condition to calculate changes in phosphorylation levels for

698 each peptide. Each condition was run in three replicates and error bars in associated figures

699 reflect the standard deviation of the replicates in each condition.

701 **Statistical analysis**

702 Statistical analyses were performed using GraphPad Prism. When comparing three or more

703 groups, statistical significance was assessed using one-way ANOVA. For comparisons between

704 two groups, a two-tailed unpaired Student's *t*-test was used. Exact *p*-values, statistical tests, and

705 replicate numbers are detailed in the corresponding figure legends.

707 **Data Availability**

708 Mass spectrometry data generated for this study are available through PRIDE, wherein

709 phosphoproteomics data can be found under PXD identifier PXD045462

710 (https://www.ebi.ac.uk/pride/)(*69*).

ACKNOWLEDGEMENTS

We thank Libor Macurek for providing U2OS WT and PPM1D-KO cells, Daniel Durocher for providing U2OS WT and 53BP1-KO cells, and Alessandro Vindigni for providing U2OS WT and SMARCAL1-KO cells. We acknowledge the Weill Institute for providing access to the ChemiDoc Imaging system used in this study. We thank Catalina Pereira and Munisha Mujingjiang for providing supplies and technical support to perform DNA combing experiments. We thank Beatriz Almeida for technical assistance and all members of the Smolka lab for helpful discussions and technical support. M.B.S. was supported by NIH grant R35-GM563209. Y.W. was supported by a postdoctoral fellowship from the Fleming Research Foundation.

AUTHOR CONTRIBUTIONS

JB and MBS led the conceptualization of the main study, defining the problem statement and research objectives. YC, YW, and MBS contributed to the conceptualization and refinement of the final study design. JB performed and analyzed immunofluorescence, western blot, chromatin fractionation, comet assays, clonogenic assays, and DNA fiber assays for the initial study. YC performed and analyzed the final experiments for all immunofluorescence, EdU incorporation, western blot, siRNA transfection, and DNA combing assays. YW conducted and analyzed the final experiments for BrdU incorporation, neutral comet assays, and DNA fiber assays. YW and TCH jointly performed and analyzed the clonogenic survival assays. JB and VMF carried out and analyzed the phosphoproteomic experiments. KJ performed and analyzed the ProKAS experiments. EÇ contributed to DNA fiber data analysis for the pilot study. JB, YC, and YW prepared the data visualizations. MBS and JB wrote the initial version of the manuscript, with YC and YW contributing to the final version. YC, YW, KJ, and MBS reviewed and edited the manuscript. MBS secured funding and supervised the project. All authors contributed to and approved the final version of the manuscript.

DECLARATION OF INTERESTS

738   The authors declare no conflicts of interest.

REFERENCES

1. M. K. Zeman, K. A. Cimprich, Causes and consequences of replication stress. *Nat. Cell Biol.* **16**, 2–9 (2014).

2. S. Saxena, L. Zou, Hallmarks of DNA replication stress. *Mol. Cell* **82**, 2298–2314 (2022).

3. P. Pasero, A. Vindigni, Nucleases acting at stalled forks: How to reboot the replication program with a few shortcuts. *Annu. Rev. Genet.* **51**, 477–499 (2017).

4. E. A. Nam, D. Cortez, ATR signalling: more than meeting at the fork. *Biochem. J* **436**, 527–536 (2011).

5. L. I. Toledo, M. Altmeyer, M.-B. Rask, C. Lukas, D. H. Larsen, L. K. Povlsen, S. Bekker-Jensen, N. Mailand, J. Bartek, J. Lukas, ATR prohibits replication catastrophe by preventing global exhaustion of RPA. *Cell* **155**, 1088–1103 (2013).

6. A. P. Bertolin, B. Canal, M. Yekezare, J. Zeng, R. Instrell, M. Howell, J. F. X. Diffley, The mechanism of checkpoint-dependent DNA replication fork stabilization in human cells, *bioRxiv* (2024). https://doi.org/10.1101/2024.11.01.621514.

7. W. Leung, A. Simoneau, S. Saxena, J. Jackson, P. S. Patel, M. Limbu, A. Vindigni, L. Zou, ATR protects ongoing and newly assembled DNA replication forks through distinct mechanisms. *Cell Rep.* **42**, 112792 (2023).

8. M. El-Shemerly, D. Hess, A. K. Pyakurel, S. Moselhy, S. Ferrari, ATR-dependent pathways control hEXO1 stability in response to stalled forks. *Nucleic Acids Res.* **36**, 511–519 (2008).

9. S. Li, Z. Lavagnino, D. Lemacon, L. Kong, A. Ustione, X. Ng, Y. Zhang, Y. Wang, B. Zheng, H. Piwnica-Worms, A. Vindigni, D. W. Piston, Z. You, Ca2+-Stimulated AMPK-Dependent Phosphorylation of Exo1 Protects Stressed Replication Forks from Aberrant Resection. *Mol. Cell* **74**, 1123–1137.e6 (2019).

10. M. Berti, D. Cortez, M. Lopes, The plasticity of DNA replication forks in response to clinically relevant genotoxic stress. *Nat. Rev. Mol. Cell Biol.* **21**, 633–651 (2020).

11. A. Quinet, D. Lemaçon, A. Vindigni, Replication Fork Reversal: Players and Guardians. *Mol. Cell* **68**, 830–833 (2017).

12. M. B. Adolph, D. Cortez, Mechanisms and regulation of replication fork reversal. *DNA Repair* **141**, 103731 (2024).

13. K. Mutreja, J. Krietsch, J. Hess, S. Ursich, M. Berti, F. K. Roessler, R. Zellweger, M. Patra, G. Gasser, M. Lopes, ATR-Mediated Global Fork Slowing and Reversal Assist Fork Traverse and Prevent Chromosomal Breakage at DNA Interstrand Cross-Links. *Cell Rep.* **24**, 2629–2642.e5 (2018).

14. D. Dibitetto, S. Marshall, A. Sanchi, M. Liptay, J. Badar, M. Lopes, S. Rottenberg, M. B. Smolka, DNA-PKcs promotes fork reversal and chemoresistance. *Mol. Cell* **82**, 3932–3942.e6 (2022).

15. E. C. Minca, D. Kowalski, Replication fork stalling by bulky DNA damage: localization at
active origins and checkpoint modulation. *Nucleic Acids Res.* **39**, 2610–2623 (2011).

16. D. V. Bugreev, R. M. Brosh Jr, A. V. Mazin, RECQ1 possesses DNA branch migration
activity. *J. Biol. Chem.* **283**, 20231–20242 (2008).

17. S. Thangavel, M. Berti, M. Levikova, C. Pinto, S. Gomathinayagam, M. Vujanovic, R.
Zellweger, H. Moore, E. H. Lee, E. A. Hendrickson, P. Cejka, S. Stewart, M. Lopes, A.
Vindigni, DNA2 drives processing and restart of reversed replication forks in human cells. *J.*
Cell Biol. **208**, 545–562 (2015).

18. L. Costantino, S. K. Sotiriou, J. K. Rantala, S. Magin, E. Mladenov, T. Helleday, J. E.
Haber, G. Iliakis, O. P. Kallioniemi, T. D. Halazonetis, Break-induced replication repair of
damaged forks induces genomic duplications in human cells. *Science* **343**, 88–91 (2014).

19. P. J. Hastings, J. R. Lupski, S. M. Rosenberg, G. Ira, Mechanisms of change in gene copy
number. *Nat. Rev. Genet.* **10**, 551–564 (2009).

20. M. Ensminger, M. Löbrich, One end to rule them all: Non-homologous end-joining and
homologous recombination at DNA double-strand breaks. *Br. J. Radiol.* **93**, 20191054
(2020).

21. Z. Gyüre, Á. Póti, E. Németh, B. Szikriszt, R. Lózsa, M. Krawczyk, A. L. Richardson, D.
Szüts, Spontaneous mutagenesis in human cells is controlled by REV1-Polymerase ζ and
PRIMPOL. *Cell Rep.* **42**, 112887 (2023).

22. A. Mazouzi, A. Stukalov, A. C. Müller, D. Chen, M. Wiedner, J. Prochazkova, S.-C. Chiang,
M. Schuster, F. P. Breitwieser, A. Pichlmair, S. F. El-Khamisy, C. Bock, R. Kralovics, J.
Colinge, K. L. Bennett, J. I. Loizou, A Comprehensive Analysis of the Dynamic Response to
Aphidicolin-Mediated Replication Stress Uncovers Targets for ATM and ATMIN. *Cell Rep.*
15, 893–908 (2016).

23. N. Kanu, T. Zhang, R. A. Burrell, A. Chakraborty, J. Cronshaw, C. DaCosta, E. Grönroos,
H. N. Pemberton, E. Anderton, L. Gonzalez, S. Sabbioneda, H. D. Ulrich, C. Swanton, A.
Behrens, RAD18, WRNIP1 and ATMIN promote ATM signalling in response to replication
stress. *Oncogene* **35**, 4009–4019 (2016).

24. S. A. Wagner, H. Oehler, A. Voigt, D. Dalic, A. Freiwald, H. Serve, P. Beli, ATR inhibition
rewires cellular signaling networks induced by replication stress. *Proteomics* **16**, 402–416
(2016).

25. K. Nakamura, G. Kustatscher, C. Alabert, M. Hödl, I. Forne, M. Völker-Albert, S. Satpathy,
T. E. Beyer, N. Mailand, C. Choudhary, A. Imhof, J. Rappsilber, A. Groth, Proteome
dynamics at broken replication forks reveal a distinct ATM-directed repair response
suppressing DNA double-strand break ubiquitination. *Mol. Cell* **81**, 1084–1099.e6 (2021).

26. C. Ribeyre, R. Zellweger, M. Chauvin, N. Bec, C. Larroque, M. Lopes, A. Constantinou,
Nascent DNA Proteomics Reveals a Chromatin Remodeler Required for Topoisomerase I
Loading at Replication Forks. *Cell Rep.* **15**, 300–309 (2016).

27. A. K. Ashley, M. Shrivastav, J. Nie, C. Amerin, K. Troksa, J. G. Glanzer, S. Liu, S. O.
Opiyo, D. D. Dimitrova, P. Le, B. Sishc, S. M. Bailey, G. G. Oakley, J. A. Nickoloff, DNA-PK

phosphorylation of RPA32 Ser4/Ser8 regulates replication stress checkpoint activation, fork restart, homologous recombination and mitotic catastrophe. *DNA Repair (Amst)* **21**, 131–139 (2014).

28. S. Ying, Z. Chen, A. L. Medhurst, J. A. Neal, Z. Bao, O. Mortusewicz, J. McGouran, X. Song, H. Shen, F. C. Hamdy, B. M. Kessler, K. Meek, T. Helleday, DNA-PKcs and PARP1 Bind to Unresected Stalled DNA Replication Forks Where They Recruit XRCC1 to Mediate Repair. *Cancer Res* **76**, 1078–1088 (2016).

29. M. C. Lanz, D. Dibitetto, M. B. Smolka, DNA damage kinase signaling: checkpoint and repair at 30 years. *EMBO J.* **38**, e101801 (2019).

30. D. P. Waterman, J. E. Haber, M. B. Smolka, Checkpoint responses to DNA double-strand breaks. *Annu. Rev. Biochem.* **89**, 103–133 (2020).

31. D.-H. Lee, D. Chowdhury, What goes on must come off: phosphatases gate-crash the DNA damage response. *Trends Biochem. Sci.* **36**, 569–577 (2011).

32. M. Shimada, M. Nakanishi, Response to DNA damage: why do we need to focus on protein phosphatases? *Front. Oncol.* **3**, 8 (2013).

33. A. K. Freeman, A. N. Monteiro, Phosphatases in the cellular response to DNA damage. *Cell Commun. Signal.* **8**, 27 (2010).

34. X. Lu, B. Nannenga, L. A. Donehower, PPM1D dephosphorylates Chk1 and p53 and abrogates cell cycle checkpoints. *Genes Dev.* **19**, 1162–1174 (2005).

35. X. Lu, T.-A. Nguyen, X. Zhang, L. A. Donehower, The Wip1 phosphatase and Mdm2: cracking the "Wip" on p53 stability. *Cell Cycle* **7**, 164–168 (2008).

36. S. Shreeram, O. N. Demidov, W. K. Hee, H. Yamaguchi, N. Onishi, C. Kek, O. N. Timofeev, C. Dudgeon, A. J. Fornace, C. W. Anderson, Y. Minami, E. Appella, D. V. Bulavin, Wip1 phosphatase modulates ATM-dependent signaling pathways. *Mol. Cell* **23**, 757–764 (2006).

37. L. Macůrek, A. Lindqvist, O. Voets, J. Kool, H. R. Vos, R. H. Medema, Wip1 phosphatase is associated with chromatin and dephosphorylates gammaH2AX to promote checkpoint inhibition. *Oncogene* **29**, 2281–2291 (2010).

38. X. Lu, T.-A. Nguyen, S.-H. Moon, Y. Darlington, M. Sommer, L. A. Donehower, The type 2C phosphatase Wip1: an oncogenic regulator of tumor suppressor and DNA damage response pathways. *Cancer Metastasis Rev.* **27**, 123–135 (2008).

39. J. D. Kahn, P. G. Miller, A. J. Silver, R. S. Sellar, S. Bhatt, C. Gibson, M. McConkey, D. Adams, B. Mar, P. Mertins, S. Fereshetian, K. Krug, H. Zhu, A. Letai, S. A. Carr, J. Doench, S. Jaiswal, B. L. Ebert, PPM1D-truncating mutations confer resistance to chemotherapy and sensitivity to PPM1D inhibition in hematopoietic cells. *Blood* **132**, 1095–1105 (2018).

40. P. Kleiblova, I. A. Shaltiel, J. Benada, J. Ševčík, S. Pecháčková, P. Pohlreich, E. E. Voest, P. Dundr, J. Bartek, Z. Kleibl, R. H. Medema, L. Macurek, Gain-of-function mutations of PPM1D/Wip1 impair the p53-dependent G1 checkpoint. *J. Cell Biol.* **201**, 511–521 (2013).

41. M. Stoyanov, A. S. Martinikova, K. Matejkova, K. Horackova, P. Zemankova, K. Burdova, Z. Zemanova, P. Kleiblova, Z. Kleibl, L. Macurek, PPM1D activity promotes cellular

transformation by preventing senescence and cell death. *Oncogene* **43**, 3081–3093 (2024).

42. M. Burocziova, P. Danek, A. Oravetzova, Z. Chalupova, M. Alberich-Jorda, L. Macurek,
Ppm1d truncating mutations promote the development of genotoxic stress-induced AML.
Leukemia **37**, 2209–2220 (2023).

43. J. I. Hsu, T. Dayaram, A. Tovy, E. De Braekeleer, M. Jeong, F. Wang, J. Zhang, T. P.
Heffernan, S. Gera, J. J. Kovacs, J. R. Marszalek, C. Bristow, Y. Yan, G. Garcia-Manero,
H. Kantarjian, G. Vassiliou, P. A. Futreal, L. A. Donehower, K. Takahashi, M. A. Goodell,
PPM1D Mutations Drive Clonal Hematopoiesis in Response to Cytotoxic Chemotherapy.
Cell Stem Cell **23**, 700–713.e6 (2018).

44. X. Lu, T.-A. Nguyen, L. A. Donehower, Reversal of the ATM/ATR-mediated DNA damage
response by the oncogenic phosphatase PPM1D. *Cell Cycle* **4**, 1060–1064 (2005).

45. H. Cha, J. M. Lowe, H. Li, J.-S. Lee, G. I. Belova, D. V. Bulavin, A. J. Fornace Jr, Wip1
directly dephosphorylates gamma-H2AX and attenuates the DNA damage response.
Cancer Res. **70**, 4112–4122 (2010).

46. H. Yamaguchi, S. R. Durell, D. K. Chatterjee, C. W. Anderson, E. Appella, The Wip1
phosphatase PPM1D dephosphorylates SQ/TQ motifs in checkpoint substrates
phosphorylated by PI3K-like kinases. *Biochemistry* **46**, 12594–12603 (2007).

47. J. F. Gräf, I. Mikicic, X. Ping, C. Scalera, K. Mayr, L. S. Stelzl, P. Beli, S. A. Wagner,
Substrate spectrum of PPM1D in the cellular response to DNA double-strand breaks.
iScience **25**, 104892 (2022).

48. A. G. Gilmartin, T. H. Faitg, M. Richter, A. Groy, M. A. Seefeld, M. G. Darcy, X. Peng, K.
Federowicz, J. Yang, S.-Y. Zhang, E. Minthorn, J.-P. Jaworski, M. Schaber, S. Martens, D.
E. McNulty, R. H. Sinnamon, H. Zhang, R. B. Kirkpatrick, N. Nevins, G. Cui, B. Pietrak, E.
Diaz, A. Jones, M. Brandt, B. Schwartz, D. A. Heerding, R. Kumar, Allosteric Wip1
phosphatase inhibition through flap-subdomain interaction. *Nat. Chem. Biol.* **10**, 181–187
(2014).

49. E. Petermann, M. L. Orta, N. Issaeva, N. Schultz, T. Helleday, Hydroxyurea-stalled
replication forks become progressively inactivated and require two different RAD51-
mediated pathways for restart and repair. *Mol. Cell* **37**, 492–502 (2010).

50. L. Macurek, J. Benada, E. Müllers, V. A. Halim, K. Krejčíková, K. Burdová, S. Pecháčková,
Z. Hodný, A. Lindqvist, R. H. Medema, J. Bartek, Downregulation of Wip1 phosphatase
modulates the cellular threshold of DNA damage signaling in mitosis. *Cell Cycle* **12**, 251–
262 (2013).

51. P. Khadka, Z. J. Reitman, S. Lu, G. Buchan, G. Gionet, F. Dubois, D. M. Carvalho, J. Shih,
S. Zhang, N. F. Greenwald, T. Zack, O. Shapira, K. Pelton, R. Hartley, H. Bear, Y. Georgis,
S. Jarmale, R. Melanson, K. Bonanno, K. Schoolcraft, P. G. Miller, A. L. Condurat, E. M.
Gonzalez, K. Qian, E. Morin, J. Langhnoja, L. E. Lupien, V. Rendo, J. Digiacomo, D. Wang,
K. Zhou, R. Kumbhani, M. E. Guerra Garcia, C. E. Sinai, S. Becker, R. Schneider, J.
Vogelzang, K. Krug, A. Goodale, T. Abid, Z. Kalani, F. Piccioni, R. Beroukhim, N. S.
Persky, D. E. Root, A. M. Carcaboso, B. L. Ebert, C. Fuller, O. Babur, M. W. Kieran, C.
Jones, H. Keshishian, K. L. Ligon, S. A. Carr, T. N. Phoenix, P. Bandopadhayay, PPM1D
mutations are oncogenic drivers of de novo diffuse midline glioma formation. *Nat. Commun.*

13, 604 (2022).

52. J. Yan, X.-P. Yang, Y.-S. Kim, A. M. Jetten, RAP80 responds to DNA damage induced by
both ionizing radiation and UV irradiation and is phosphorylated at Ser 205. *Cancer Res* **68**,
4269–4276 (2008).

53. B. Sobhian, G. Shao, D. R. Lilli, A. C. Culhane, L. A. Moreau, B. Xia, D. M. Livingston, R. A.
Greenberg, RAP80 targets BRCA1 to specific ubiquitin structures at DNA damage sites.
Science **316**, 1198–1202 (2007).

54. S.-T. Kim, B. Xu, M. B. Kastan, Involvement of the cohesin protein, Smc1, in Atm-
dependent and independent responses to DNA damage. *Genes Dev.* **16**, 560–570 (2002).

55. N. Salvador Moreno, J. Liu, K. M. Haas, L. L. Parker, C. Chakraborty, S. J. Kron, K.
Hodges, L. D. Miller, C. Langefeld, P. J. Robinson, S. A. Lelièvre, P.-A. Vidi, The nuclear
structural protein NuMA is a negative regulator of 53BP1 in DNA double-strand break
repair. *Nucleic Acids Res.* **47**, 2703–2715 (2019).

56. Y. Ziv, D. Bielopolski, Y. Galanty, C. Lukas, Y. Taya, D. C. Schultz, J. Lukas, S. Bekker-
Jensen, J. Bartek, Y. Shiloh, Chromatin relaxation in response to DNA double-strand
breaks is modulated by a novel ATM- and KAP-1 dependent pathway. *Nat. Cell Biol.* **8**,
870–876 (2006).

57. H. Jaiswal, J. Benada, E. Müllers, K. Akopyan, K. Burdova, T. Koolmeister, T. Helleday, R.
H. Medema, L. Macurek, A. Lindqvist, ATM/Wip1 activities at chromatin control Plk1 re-
activation to determine G2 checkpoint duration. *EMBO J.* **36**, 2161–2176 (2017).

58. A. M. Kolinjivadi, V. Sannino, A. de Antoni, H. Técher, G. Baldi, V. Costanzo, Moonlighting
at replication forks - a new life for homologous recombination proteins BRCA1, BRCA2 and
RAD51. *FEBS Lett.* **591**, 1083–1100 (2017).

59. Z. Wang, C. Xu, B. H. Diplas, C. J. Moure, C.-P. J. Chen, L. H. Chen, C. Du, H. Zhu, P. K.
Greer, L. Zhang, Y. He, M. S. Waitkus, H. Yan, Targeting mutant PPM1D sensitizes diffuse
intrinsic pontine glioma cells to the PARP inhibitor olaparib. *Mol. Cancer Res.* **18**, 968–980
(2020).

60. K. Burdova, R. Storchova, M. Palek, L. Macurek, WIP1 promotes homologous
recombination and modulates sensitivity to PARP inhibitors. *Cells* **8**, 1258 (2019).

61. I. Rappold, K. Iwabuchi, T. Date, J. Chen, Tumor suppressor p53 binding protein 1 (53BP1)
is involved in DNA damage-signaling pathways. *J. Cell Biol.* **153**, 613–620 (2001).

62. L. Feng, N. Li, Y. Li, J. Wang, M. Gao, W. Wang, J. Chen, Cell cycle-dependent inhibition of
53BP1 signaling by BRCA1. *Cell Discov.* **1**, 15019 (2015).

63. R. Pavani, V. Tripathi, K. B. Vrtis, D. Zong, R. Chari, E. Callen, A. V. Pankajam, G. Zhen,
G. Matos-Rodrigues, J. Yang, S. Wu, G. Reginato, W. Wu, P. Cejka, J. C. Walter, A.
Nussenzweig, Structure and repair of replication-coupled DNA breaks. *Science* **385**,
eado3867 (2024).

64. A. N. Blackford, S. P. Jackson, ATM, ATR, and DNA-PK: The Trinity at the heart of the
DNA damage response. *Mol. Cell* **66**, 801–817 (2017).

65.   W. J. Comstock, M. V. Navarro, D. V. Maybee, Y. Rho, M. Wagner, Y. Wang, M. B.
Smolka, Proteomic sensors for quantitative, multiplexed and spatial monitoring of kinase
signaling, *bioRxivorg* (2024). https://doi.org/10.1101/2024.12.16.628391.

66.   J. L. Johnson, T. M. Yaron, E. M. Huntsman, A. Kerelsky, J. Song, A. Regev, T.-Y. Lin, K.
Liberatore, D. M. Cizin, B. M. Cohen, N. Vasan, Y. Ma, K. Krismer, J. T. Robles, B. van de
Kooij, A. E. van Vlimmeren, N. Andrée-Busch, N. F. Käufer, M. V. Dorovkov, A. G.
Ryazanov, Y. Takagi, E. R. Kastenhuber, M. D. Goncalves, B. D. Hopkins, O. Elemento, D.
J. Taatjes, A. Maucuer, A. Yamashita, A. Degterev, M. Uduman, J. Lu, S. D. Landry, B.
Zhang, I. Cossentino, R. Linding, J. Blenis, P. V. Hornbeck, B. E. Turk, M. B. Yaffe, L. C.
Cantley, An atlas of substrate specificities for the human serine/threonine kinome. *Nature*
613, 759–766 (2023).

67.   C. J. Bakkenist, M. B. Kastan, DNA damage activates ATM through intermolecular
autophosphorylation and dimer dissociation. *Nature* **421**, 499–506 (2003).

68.   Z. Guo, S. Kozlov, M. F. Lavin, M. D. Person, T. T. Paull, ATM activation by oxidative
stress. *Science* **330**, 517–521 (2010).

69.   Y. Perez-Riverol, J. Bai, C. Bandla, D. García-Seisdedos, S. Hewapathirana, S.
Kamatchinathan, D. J. Kundu, A. Prakash, A. Frericks-Zipper, M. Eisenacher, M. Walzer, S.
Wang, A. Brazma, J. A. Vizcaíno, The PRIDE database resources in 2022: a hub for mass
spectrometry-based proteomics evidences. *Nucleic Acids Res* **50**, D543–D552 (2022).

955
956

 FIGURE LEGENDS

Figure 1. PPM1D regulates replication fork restart.

(a) EdU incorporation analysis in U2OS cells. Top: experimental scheme illustrating treatment

with the PPM1D inhibitor (PPM1Di; GSK2830371; 10 µM). Bottom: dot plot showing EdU

intensity in PCNA-positive cells pooled from three independent biological replicates (at least 104

cells per condition per replicate). The mean EdU intensity from each replicate is indicated by

black dots. *p*-values were determined by one-way ANOVA with Tukey's multiple comparisons

test. Error bars represent standard deviation. (b) Representative immunofluorescence images

from the EdU incorporation assay shown in (a). Scale bar, 10 µm. (c) EdU incorporation

analysis in U2OS wild-type (WT) and PPM1D-knockout (KO) cells. Top: experimental scheme.

Bottom: dot plot showing EdU intensity in PCNA-positive cells pooled from three independent

biological replicates (at least 83 cells per condition per replicate). The mean EdU intensity from

each replicate is indicated by black dots. *p*-values were determined by one-way ANOVA with

Tukey's multiple comparisons test. Error bars represent standard deviation. (d) DNA combing

assay in U2OS cells. Top: experimental scheme. Bottom left: Representative images of DNA

combing assay. Bottom right: bar graph showing the percentages of stalled replication forks in

cells treated with or without PPM1Di (10 µM). A minimum of 2,720 fibers were quantified per

condition from three independent biological replicates. *p*-values were calculated using Student's

t-test. Error bars represent standard deviation. (e) DNA combing assay in U2OS cells

transfected with control and PPM1D siRNA (siCTL and si*PPM1D*). Top: experimental scheme.

Bottom: bar graph showing the percentages of stalled replication forks in U2OS wild-type (WT)

and PPM1D-knockout (KO) cells. A minimum of 1,644 fibers were quantified per condition from

two independent biological replicates. *p*-values were calculated using one-way ANOVA with

Tukey's multiple comparisons test. Error bars represent standard deviation. (f) Representative

immunofluorescence images showing co-staining of PPM1D (red) and EdU (green) detected via anti-PPM1D antibody and click chemistry, respectively. Scale bar, 10 µm. (g) Dot plot showing the PPM1D intensity in EdU-negative and EdU-positive U2OS cells. At least 328 cells were analyzed per condition. *p*-value was calculated using Student's t-test. Error bars represent standard deviation.

Figure 2. PPM1D counteracts nucleolytic processing of stalled replication forks.

(a) RPA32 foci analysis in U2OS cells. Top: experimental scheme illustrating treatment with the PPM1D inhibitor (PPM1Di, 10 µM). Bottom: dot plot showing the number of RPA32 foci in PCNA-positive cells pooled from three independent biological replicates (at least 101 cells per condition per replicate). The mean number of RPA32 foci from each replicate is indicated by black dots. *p*-value was calculated using Student's t-test. Error bars represent standard deviation. (b) Representative immunofluorescence images corresponding to (a). Scale bar, 10 µm. (c) RPA32 foci analysis in U2OS wild-type (WT) and PPM1D-knockout (KO) cells. Top: experimental scheme. Bottom: dot plot shows the number of RPA32 foci in PCNA-positive cells pooled from three independent biological replicates (at least 101 cells per condition per replicate). The mean number of RPA32 foci from each replicate is indicated by black dots. *p*-value was calculated using Student's t-test. (d) Representative immunofluorescence images corresponding to (c). Scale bar, 10 µm. (e) Analysis of nascent single-stranded DNA (ssDNA) in U2OS cells. Top: experimental scheme showing BrdU immunostaining under non-denaturing conditions following the indicated treatment. Bottom: dot plot shows the number of BrdU foci in PCNA-positive cells from one representative experiment. Two biological replicates were performed; results from the second biological replicate are shown in Fig. S2b. *p*-values were calculated using one-way ANOVA. (f) RPA32 foci analysis in U2OS WT and SMARCAL1-KO cells. Top: experimental scheme illustrating treatment with the PPM1Di (10 µM). Bottom: dot plot shows the number of RPA32 foci in PCNA-positive cells pooled from three independent

biological replicates (at least 73 cells per condition per replicate). The mean number of RPA32

foci from each replicate is indicated by black dots. *p*-values were calculated using one-way

ANOVA with Tukey's multiple comparisons test. (g) DNA fiber assay in U2OS WT and

SMARCAL1-KO cells. Top: experimental scheme. Bottom: bar graph shows the percentage of

stalled replication forks in the presence or absence of PPM1Di (10 μM). At least 942 fibers per

condition were quantified across three independent biological replicates. *p*-values were

calculated using one-way ANOVA with Tukey's multiple comparisons test. Error bars represent

standard deviation. (h-i) RPA32 foci analysis in U2OS WT and PPM1D-KO cells treated with or

without MRE11 inhibitor (Mirin; 10 μM) (h) or DNA2 inhibitor (DNA2i; 20 μM) (i), as shown in the

experimental schemes. Dot plots show the number of RPA32 foci in PCNA-positive cells pooled

from three independent biological replicates (at least 97 (h) or 99 (i) cells per condition per

replicate). The mean number of RPA32 foci from each replicate is indicated by black dots. *p*-

values were calculated using one-way ANOVA with Tukey's multiple comparisons test. Error

bars represent standard deviation.

1022

Figure 3. PPM1D antagonizes ATM signaling during replication stress recovery.

(a) Scatter plot showing all phosphosites downregulated during replication stress recovery.

Phosphosites from the HU Release/HU Arrest comparison are categorized along the *y*-axis as

either PPM1D-independent (left and center) or PPM1D-dependent (right). (b) Pie chart

summarizing all phosphosites downregulated during replication stress recovery, illustrating the

proportions of PPM1D-independent and PPM1D-dependent sites. (c) Pie chart highlighting the

subset of S/T-Q phosphosites within the PPM1D-independent and PPM1D-dependent groups

shown in (b). (d) Bar graph demonstrating the enrichment of the consensus phospho-motifs

(S/T-Q or S/T-Q-(D/E)$^{2+}$) in cells treated with or without PPM1D inhibitor (PPM1Di). (e)

Immunoblot analysis of U2OS cells treated with hydroxyurea (HU; 4 mM), PPM1Di (10 μM),

and/or ATMi (100 nM) for the indicated times. (f-g) Quantification of relative protein levels of

pKAP1 (f) and pCHK1 (g) based on immunoblot analysis in (e). Bar graphs show normalized

protein expression levels relative to β-actin across three independent biological replicates. The

normalized protein expression level from each replicate is indicated by black dots. *p*-values

were calculated using one-way ANOVA with Tukey's multiple comparisons test. Error bars

represent standard deviation. (h) Phospho-ATM (pATM, S1981) foci analysis in U2OS cells

treated with or without PPM1Di. Dot plot shows the number of pATM foci in PCNA-positive cells

pooled from three independent biological replicates (at least 91 cells per condition per replicate).

The mean number of pATM foci from each replicate is indicated by black dots. *p*-values were

calculated using one-way ANOVA with Tukey's multiple comparisons test. Error bars represent

standard deviation.

1044

Figure 4. ATM inhibition rescues defects in replication fork processing and restart

caused by loss of PPM1D function.

(a) RPA32 foci analysis in U2OS cells. Top: experimental scheme showing treatment with

PPM1Di and ATM inhibitor (ATMi; AZD0156; 100 nM). Bottom: dot plot shows the number of

RPA32 foci in PCNA-positive cells pooled from three independent biological replicates (at least

93 cells per condition per replicate). The mean number of RPA32 foci from each replicate is

indicated by black dots. *p*-values were calculated using one-way ANOVA with Tukey's multiple

comparisons test. Error bars represent standard deviation. (b) Representative

immunofluorescence images corresponding to the RPA32 foci analysis shown in (a). Scale bar,

10 μm. (c) EdU incorporation analysis in U2OS cells. Top: experimental scheme showing

treatment with PPM1Di and ATMi. Bottom: dot plot shows the EdU intensity in PCNA-positive

cells pooled from three independent biological replicates (at least 95 cells per condition per

replicate). The mean EdU intensity from each replicate is indicated by black dots. *p*-values were

determined by one-way ANOVA with Tukey's multiple comparisons test. Error bars represent

standard deviation. (d) Representative immunofluorescence images from the EdU incorporation

assay shown in (c). Scale bar, 10 µm. (e) DNA fiber assay in U2OS cells. Top: experimental

scheme showing treatment with PPM1Di and ATMi, as indicated. Bottom: bar graph shows the

percentage of stalled replication forks under each treatment condition. A minimum of 2,929

fibers per condition were analyzed from three independent biological replicates. *p*-values were

calculated using one-way ANOVA. Error bars represent standard deviation. (f) DNA protection

assay in U2OS cells. Top: experimental scheme. Bottom: dot plot showing the ratio of IdU/CldU

tract lengths as an indicator of replication fork degradation in one representative experiment. At

least 150 fibers were analyzed per condition. Results from the second biological replicate are

shown in Fig. S3c. *p*-values were calculated using one-way ANOVA. (g) Clonogenic survival

assay of U2OS cells co-treated with ATMi, ATRi, and PPM1Di. Colony area was normalized to

the respective untreated PPM1Di control group for each treatment condition. Data represent

three independent biological replicates. *p*-values were assessed using Student's t-test.

1072

Figure 5. PPM1D regulates RAD51-mediated replication fork restart.

(a) RAD51 foci analysis in U2OS cells. Top: experimental scheme. Bottom: dot plot shows the

number of RAD51 foci in PCNA-positive cells pooled from three independent biological

replicates (at least 96 cells per condition per replicate). The mean number of RAD51 foci from

each replicate is indicated by black dots. *p*-value was calculated using Student's t-test. Error

bars represent standard deviation. (b) Representative immunofluorescence images

corresponding to (a). Scale bar, 10 µm. (c) Immunoblot analysis of PPM1D, RAD51, and histone

H4 in chromatin fractions from U2OS cells treated with or without PPM1Di. Ponceau S staining

was used as a loading and total protein normalization control. (d) DNA fiber assay in U2OS

cells. Top: experimental scheme showing treatment with PPM1Di and RAD51 inhibitor (RAD51i;

B02; 20 µM). Bottom: bar graph shows the percentage of stalled replication forks under each

condition. A minimum of 2,959 fibers per condition were quantified from three independent

biological replicates. *p*-values were calculated using one-way ANOVA. Error bars represent

standard deviation. (e) RAD51 foci analysis in U2OS wild-type (WT) and PPM1D-knockout (KO)

cells. Top: experimental scheme. Bottom: dot plot shows the number of RAD51 foci in PCNA-

positive cells pooled from three independent biological replicates (at least 102 cells per

condition per replicate). The mean number of RAD51 foci from each replicate is indicated by

black dots. *p*-values were calculated using one-way ANOVA with Tukey's multiple comparisons

test. Error bars represent standard deviation. (f) 53BP1 foci analysis in U2OS cells. Top:

experimental scheme. Bottom: dot plot shows the number of 53BP1 foci in PCNA-positive cells

pooled from three independent biological replicates (at least 96 cells per condition per replicate).

The mean number of 53BP1 foci from each replicate is indicated by black dots. *p*-values were

calculated using one-way ANOVA with Tukey's multiple comparisons test. Error bars represent

standard deviation. (g) RAD51 foci analysis in U2OS WT and 53BP1-knockout (KO) cells. Top:

experimental scheme. Bottom: dot plot shows the number of RAD51 foci in PCNA-positive cells

pooled from three independent biological replicates (at least 88 cells per condition per replicate).

The mean number of RAD51 foci from each replicate is indicated by black dots. *p*-values were

calculated using one-way ANOVA with Tukey's multiple comparisons test. Error bars represent

standard deviation.

1102

Figure 6. Increased ATM signaling during fork recovery in cells with impaired PPM1D

function is not correlated with an increase in detectable DSB.

(a) Neutral comet assay in U2OS cells. Top: experimental scheme. Bottom: dot plot shows the

tail moment for each condition. At least 102 cells were quantified per condition. Treatment with 1

µM camptothecin (CPT) for 30 minutes was used as a positive control. *p*-values were calculated

using one-way ANOVA. (b) Immunoblot analysis of phosphorylated KAP1 (pKAP1) in U2OS

cells treated with HU (4 mM) and/or PPM1Di (10 µM) for the indicated durations. The bar graph

at the bottom shows the normalized pKAP1 expression levels relative to β-tubulin from three

independent biological replicates. The normalized pKAP1 expression level from each replicate

is indicated by black dots. *p*-values were calculated using one-way ANOVA with Tukey's

multiple comparisons test. Error bars represent standard deviation. (c) pKAP1 (S824) foci

analysis in U2OS cells. Top: experimental scheme. Bottom: dot plot shows the number of

pKAP1 foci in PCNA-positive cells pooled from three independent biological replicates (at least

125 cells per condition per replicate). The mean number of pKAP1 foci from each replicate is

indicated by black dots. *p*-values were calculated using one-way ANOVA with Tukey's multiple

comparisons test. Error bars represent standard deviation. (d) ProKAS (Proteomic Kinase

Activity Sensor) design. Peptide sequences used in the ATM sensor (PPM1D-untargeted) and

ATM-PPM1D sensor (PPM1D targeted) are shown. (e) Experimental scheme using ProKAS for

detecting ATM and PPM1D activities. (f) Quantification of ATM and PPM1D activity using

ProKAS sensors during replication stress recovery with or without PPM1Di. Bar graphs

represent the normalized ATM activity (left) and PPM1D acitivity (right) relative to untreated

condition from three independent biological replicates. *p*-values were calculated using one-way

ANOVA with Tukey's multiple comparisons test. Error bars represent standard deviation. (g)

Proposed model illustrating how PPM1D promotes replication fork restart and prevents ssDNA

accumulation by counteracting ATM signaling at stalled replication forks.

 SUPPLEMENTAL FIGURE LEGENDS

 Figure S1. PPM1D modulates DNA replication.

 (a) Representative immunofluorescence images from the EdU incorporation assay shown in Fig.

 1c. Scale bar, 10 µm. (b) EdU incorporation analysis in U2OS wild-type (WT) and PPM1D-

 knockout (KO) cells transfected with control or PPM1D siRNA (siCTL and si*PPM1D*). Top:

 experimental scheme. Bottom: dot plot showing EdU intensity in PCNA-positive cells from one

 representative experiment. A minimum of 400 cells were quantified per condition from two

 independent biological replicates. *p*-values were determined by one-way ANOVA with Tukey's

 multiple comparisons test. Error bars represent standard deviation. (c) DNA combing assay in

 U2OS cells. Top: experimental scheme. Bottom: bar graph showing the percentages of

 restarted replication forks in cells treated with or without PPM1Di (10 µM). A minimum of 2,720

 fibers were quantified per condition from three independent biological replicates. *p*-values were

 calculated using Student's t-test. Error bars represent standard deviation. (d) DNA combing

 assay in U2OS wild-type (WT) and PPM1D-knockout (KO) cells transfected with control or

 PPM1D siRNA (siCTL and si*PPM1D*). Top: experimental scheme. Bottom: bar graph showing

 the percentages of restarted replication forks. A minimum of 1,644 fibers were quantified per

 condition from two independent biological replicates. *p*-values were calculated using one-way

 ANOVA with Tukey's multiple comparisons test. Error bars represent standard deviation. (e)

 Representative images of different fork types in the DNA fiber assay. (f) DNA fiber assay in

 U2OS and HCT116 cells. Top: experimental scheme. Bottom: bar graphs show the percentages

 of restarted, stalled and newly fired replication forks in cells treated with or without PPM1Di (10

 µM). A minimum of 471 fibers (U2OS) and 641 fibers (HCT116) were quantified per condition

 from three independent biological replicates. *p*-values were calculated using Student's t-test.

 Error bars represent standard deviation. (g) EdU incorporation analysis in U2OS cells. Top:

 experimental scheme illustrating treatment with the PPM1D inhibitor (10 µM) and/or CDC7

1154   inhibitor (CDC7i; XL413; 20 µM). Bottom: dot plot showing EdU intensity in PCNA-positive cells

1155   pooled from three independent biological replicates (at least 73 cells per condition per replicate).

1156   The mean EdU intensity from each replicate is indicated by black dots. *p*-values were

1157   determined by one-way ANOVA with Tukey's multiple comparisons test. Error bars represent

1158   standard deviation.

1159

Figure S2. PPM1D regulates RPA32 foci accumulation and replication-associated ssDNA

formation.

(a) Representative immunofluorescence images corresponding to Fig. 2e. (b) Analysis of

nascent single-stranded DNA (ssDNA) in U2OS cells using BrdU immunostaining under non-

denaturing conditions, as shown in the treatment scheme. Dot plot shows the number of BrdU

foci in PCNA-positive cells from a second biological replicate. At least 113 PCNA-positive cells

were quantified per condition. *p*-values were calculated using one-way ANOVA. (c) DNA fiber

assay in U2OS WT and SMARCAL1-KO cells. Bar graphs showing the percentage of restarted

(left) or newly fired (right) replication forks in the presence or absence of PPM1D inhibitor

(PPM1Di). At least 942 fibers per condition were quantified per condition across three

independent biological replicates. *p*-values were calculated using one-way ANOVA with Tukey's

multiple comparisons test. Error bars represent standard deviation. A schematic of the fiber

assay condition is shown at the top of the panel.

1173

1174
1175   **Figure S3. ATM inhibition rescues replication fork restart.**
1176
(a) EdU incorporation assay in U2OS cells. Top: experimental schemes showing treatment with

and without HU (4 mM), PPM1Di (Low dose: 3 µM, High dose: 10 µM) and ATMi (100 nM).

Bottom: dot plot shows the EdU intensity in PCNA-positive cells from one representative

experiment (at least 196 cells per condition). *p*-values were determined by one-way ANOVA

with Tukey's multiple comparisons test. Error bars represent standard deviation. (b) DNA fiber

assay in U2OS cells treated with or without PPM1Di (10 μM) and ATMi (100 nM). Bar graphs

showing the percentage of restarted (left) or newly fired (right) replication forks under each

treatment condition across three independent biological replicates. *p*-values were calculated

using one-way ANOVA with Tukey's multiple comparisons test. Error bars represent standard

deviation. A schematic of the fiber assay condition is shown at the top of the panel. (c) DNA

protection assay in U2OS cells treated with or without PPM1Di (10 μM) and ATMi (100 nM), as

indicated. Dot plot showing the ratio of IdU/CldU tract lengths as indicator of replication fork

degradation from a second biological replicate. At least 144 fibers were analyzed per condition.

p-values were calculated using one-way ANOVA.

1191
1192   **Figure S4. PPM1D regulates RAD51-mediated replication fork restart.**
1193
DNA fiber assay in U2OS cells treated with or without PPM1Di (10 μM) and RAD51 inhibitor

(RAD51i; B02; 20 μM). Bar graphs showing the percentage of newly fired (left) or restarted

(right) replication forks under each treatment condition across three independent biological

replicates. *p*-values were calculated using one-way ANOVA with Tukey's multiple comparisons

test. Error bars represent standard deviation. A schematic of the fiber assay condition is shown

at the top of the panel.

1200
1201   **Figure S5. PPM1D inhibition induces phosphorylation of ProKAS sensors in an ATM-**
1202   **dependent manner.**
1203
1204   Quantification of ATM and PPM1D activity using ProKAS sensors during replication stress

1205   recovery in the presence of PPM1Di (10 μM), with or without ATMi (100 nM). Bar graphs

1206   represent the normalized ATM activity (left) and PPM1D acitivity (right) relative to untreated

1207   condition from three independent biological replicates. *p*-values were calculated using Student's

1208   t-test. Error bars represent standard deviation.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1

Figure S2

a

b

c

Figure S3

a

b

c

Figure S4

Figure S5