Tenotomized Achilles tendons of MRL/MpJ mice show scar-mediated healing, but have superior
recovery of biomechanical properties

Ningfeng Tang[1], Kimberly Wilson[1], Mehari M. Weldemariam[2], Hongying Tian[1], Christoph D.
Hart[3], Prishasai Ramnath[3], Divya S. Sivalingam[3], Jacob Glazier[3], Jie Jiang[1], Natalie Leong[1], Tao
Lowe[3,4], Maureen A. Kane[2], Masahiro Iwamoto[1], Catherine K. Kuo[1,3*], Motomi Enomoto-
Iwamoto[1*]

[1]Department of Orthopaedics, University of Maryland School of Medicine, Baltimore, Maryland,
USA

[2]Department of Pharmaceutical Sciences, University of Maryland School of Pharmacy,
Baltimore, Maryland, USA

[3]Fischell Department of Bioengineering, A. James Clark School of Engineering, University of
Maryland, College Park, Maryland, USA

[4]Department of Oral and Maxillofacial Surgery, School of Dentistry, University of Maryland,
Baltimore, Maryland, USA

15
16 *Corresponding Authors:
17
Contact corresponding author
Motomi Enomoto-Iwamoto
Department of Orthopaedics
University of Maryland School of Medicine
670 W Baltimore St, HSFIII 7171
Baltimore, MD 21201, USA
Motomi.iwamoto@som.umaryland.edu
25
Catherine K. Kuo
Fischell Department of Bioengineering
A. James Clark School of Engineering
University of Maryland
4108 A. James Clark Hall, 8278 Paint Branch Dr
College Park, MD 20742, USA
ckk@umd.edu
33

34 Key words: tendon, injury, C57BL/6J, MRL/MpJ, proteomics

Abstract

Tendon healing in Murphy Ross Large (MRL/MpJ) mice was examined using an Achilles tendon tenotomy model, a clinically relevant but more severe model compared to a patella focal injury model in which tendons in MRL/MpJ mice showed regenerative healing. After complete transection, the Achilles tendons of MRL/MpJ mice healed via scar formation regardless of sex, similar to those of C57BL/6J mice. Tensile testing found that MRL/MpJ male exhibited higher Achilles tendon stiffness and Young's modulus than C57BL/6J males after 4 weeks of healing. Proteomics analysis revealed similar alterations to signaling pathways in injured tendons of MRL/MpJ and C57BL/6J mice during the remodeling phase of healing, 4 weeks after injury. Among the altered pathways, actin cytoskeleton and integrin signaling pathways were two of the top pathways that were more prominently activated in C57BL/6J males than in MRL/MpJ males. These findings indicate that tendons of MRL/MpJ mice do not regenerate when severely injured, but still have better recovery of mechanical properties. Control of actin cytoskeleton and integrin signaling pathways may be important in improving the mechanical properties of injured tendons.

1. Introduction

Tendons have limited repair capacity. Injured tendons do not usually regain their original levels of biological, structural, and biomechanical characteristics, and instead can undergo chronic degeneration [1-5]. In contrast to embryonic and neonatal tendons [6, 7], adult tendons heal via scar formation characterized by high cell density, disorganized cell arrangement, thin and randomly aligned collagen fibers, high vascularity, and mucoid matrix [5, 8]. Scar formation inhibits recovery of biomechanical properties. Failure of full functional recovery may result in chronic tendon disorders and re-rupture, leading to poor quality of life and work disability [5].

Murphy Roths Large (MRL/MpJ) mice have strong repair capacity in musculoskeletal tissues

including auricular cartilage, articular cartilage, muscle and bone [9]. MRL/MpJ mice also

exhibit regenerative healing in the focal injury of the patellar tendon in adults, whereas C57BL/6

mice show scar-mediated healing after the same injury [10, 11]. Determining the molecular

mechanism that supports the reparative ability of the MRL/MpJ is crucial to develop new

modalities to promote effective tendon healing. In this study, we investigated whether tendons in

MRL/MpJ mice undergo scarless healing even in an Achilles tendon transection model, which is

more severe and clinically relevant compared to a focal injury model, and used proteomics

analyses to further examine which signaling pathways may be involved in improved tendon

repair by MRL/MpJ mice.

2. Materials and Methods

2.1. Mouse experiments

All animal studies were conducted with approval from the Institutional Animal Care and Use

Committee at the University of Maryland, Baltimore (0120007 and 1122006). Animals were

housed in a University Laboratory Animal Resources supervised animal facility with a 12-hour

light/dark cycle and controlled temperature of 22 ± 1ºC and humidity of $55 \pm 5\%$. Animals were

provided hygienic animal bedding of wood shavings and cotton pads. The health status of each

animal was monitored throughout the experiments by investigators, animal veterinary

technicians, and veterinarians, as per institutional guidelines. The mice were free of all viral,

bacterial, and parasitic pathogens during the experimental schedule.

A complete transverse transection (tenotomy), without attempt at repair, was made at the

midpoint of the left Achilles tendon in 12-week-old female and male C57BL/6J (#000646, The

Jackson Laboratory, Bar Harbor, ME) and MRL/MpJ (#000486, The Jackson Laboratory) mice

under anesthesia. The analgesic carprofen, 4.5 mg/Kg was administrated pre- and post-surgery.

The mice receiving the tenotomy surgery were euthanized 4 or 10-weeks post-surgery. The right

and left hind limbs were collected for histological analysis and mechanical tests. The mid-

substance of Achilles tendons were snap frozen in liquid nitrogen and stored at -80°C until use.

The criteria for animal exclusion during the experiment, determined prior to the experiment,

were to exclude mice that received analgesics and/or antibiotics and/or were euthanized due to

infection or distress. No mice were excluded in this study. No exclusion criteria were set for data

points during the analysis. No data points were excluded. MRL/MpJ and C57BL/6j mice were

maintained in the same location and compared, but randomization was not used and confounders

were not controlled. A total of 102 mice were used: 3 female C57BL/6J, 3 male C57BL/6J, 3

female MRL/MpJ, and 3 male MRL/MpJ mice for histological examination at 4 weeks post-

surgery; 8 female C57BL/6J, 5 female MRL/MpJ, 8 male C57BL/6J, and 7 female MRL/MpJ

mice for mechanical testing at 4 weeks post-surgery; 5 C57BL/6J and 5 MRL/MpJ male mice for

μCT analysis at 10 weeks post-surgery; 18 male C57BL/6J mice (making 5 replicates of

uninjured C57BL/6J tendon group), 10 male C57BL/6J mice (making 5 male replicates of

injured C57BL/6J tendon group), 10 male MRL/MpJ mice (making 4 replicates of uninjured

MRL/MpJ tendon group), and 8 male MRL/MpJ mice (making 4 replicates of injured MRL/MpJ

tendon group) for proteomics at 4-weeks post-surgery; and 3 C57BL/6J and 3 MRL/MpJ male

mice for immunoblot at 4 weeks post-surgery. Healing ability comparisons between MRL/MpJ

and C57BL/6J mice were based on mechanical properties (primary outcome measure) and

histology (secondary outcome measure).

2.2. Histology

The hind limbs were fixed with 4% paraformaldehyde for 24 hours at 4°C and decalcified with

10% ethylenediaminetetraacetic acid at pH 7.2. Serial longitudinal sections (5 μm) of the injured

or uninjured Achilles tendons were made. Sections were stained with hematoxylin-eosin and

picrosirius red. Hematoxylin-eosin staining images were captured using light microscopy (BZX-

700, Keyence, Tokyo, Japan). Picrosirius red staining images were captured using brightfield

microscopy and polarized light microscopy (20X objective, ZEISS Axio Scan, Germany) [7, 12-

14].

2.3. Micro-computed tomography (microCT) analysis

Samples were scanned using a Skyscan 1172 microCT scanner (Bruker, Billerica, MA, USA)

while immersed in phosphate buffered saline without Ca^{2+} or Mg^{2+} (PBS). Specifically, images

were acquired using previously reported conditions [15]: 6.5 µm/pixel; 0.2 mm Al filter using a

55 kV X-ray source; resolution at 2 K with step rotation of 0.51; and two average frames using

360° rotation. Image reconstruction was performed using NRecon software (Bruker). We

manually identified regions of heterotopic ossification, for which volume analyses were

performed.

2.4. Mass spectrometry-based proteomics

We collected tendon samples and stored them at -80 ºC until assay. Lysis buffer containing 5%

SDS (sodium dodecyl sulfate) and 50 mM triethylammonium bicarbonate (1 M, pH 8.0) was

added to tendons and the samples were then homogenized using bead beating with a Precellys

CK14 lysing kit (Bertin Corp., Rockville, MD). Next, protein extraction and digestion were

performed as described [15] using S-trap micro columns (ProtiFi, NY). Digested peptides were

quantified using a BCA assay kit (Thermo Fisher Scientific, 23275) to determine peptide

concentration. LC-MS/MS-based proteomic analysis was conducted on a nanoACQUITY Ultra-

Performance Liquid Chromatography system (Waters Corporation, Milford, MA USA) coupled

to an Orbitrap Fusion Lumos Tribrid mass spectrometer (Thermo Scientific, San Jose, CA USA).

Peptide separation was effected on a nanoACQUITY Ultra-Performance Liquid Chromatography

(UPLC) analytical column (BEH130 C18, 1.7 µm, 75 µm x 200 mm; Waters Corporation,

Milford, MA, USA) using a 185-min linear gradient with 3-40% acetonitrile and 0.1% formic

acid. Mass spectrometry conditions were as follows: Full MS scan resolution of 240,000,

precursor ions fragmentation by high-energy collisional dissociation of 35%, and a maximum

cycle time of 3 seconds. The resulting mass spectra were processed using Thermo Proteome

Discoverer ((PD, version 3.0.0.757), Thermo Fisher Scientific) and searched against a UniProt

mouse reference proteome (release 2024.03, 17180 entries) using Sequest HT algorithm. Search

parameters include carbamidomethylation of cysteines (+57.021 Da) as a static modification,

methionine oxidation (+15.995 Da) as a dynamic modification, precursor mass tolerance of 20

ppm, fragment mass tolerance of 0.5 Da, and trypsin as a digestion enzyme. Tryptic missed

cleavages were restricted to a maximum of two, with peptide lengths set between 6 and 144

residues. For protein quantification, the Minora feature detector, integrated in the PD, was used

as described previously [15]. To ensure high data quality, proteins were further filtered to a 1%

false discovery rate (FDR) threshold, calculated with the Percolator algorithm. Next, protein

abundance values exported from PD were post-processed using Perseus software (version

1.6.14.0). Proteins with missing values were excluded to improve data quality. Differentially

expressed proteins (DEPs) were identified using a two-tailed Student's t-test (adjusted p-value <

0.05). Ingenuity Pathway Analysis (IPA) software (Qiagen, Germantown, MD) was used to

identify dysregulated pathways and biological processes. The data have been deposited to the

ProteomeXchange Consortium via the PRIDE [16] partener repository with the dataset identifier

PXD057712.

2.5. Immunoblot analysis

Frozen tendon fragments were placed on a prechilled carbon steel block and crushed into powder

by a steel hammer, lysed in RIPA buffer, and centrifuged at 15,000 rpm for 10 min. Supernatants

(12 µg protein/lane) were subjected to SDS-PAGE (4-12%, NuPage Bis-Tris, Thermo Fisher

Scientific) followed by immunoblotting with antibodies against alpha-Actinin (11313-2-AP,

Proteintech Group, Inc, Rosment, IL) and talin-1 (14168-1-AP, Proteintech Group, Inc). The

antibodies bound to the membrane were visualized using the VECTASTATIN Elite Universal

PLUS kit, peroxidase (Vector Laboratories), and SuperSignal West Dura (Thermo Fisher

Scientific).

2.6. Mechanical Testing

Achilles tendons with calcanei were fine dissected to ensure removal of all musculature and

surrounding soft tissue. Samples were hydrated in PBS immediately after dissection and

throughout the course of mechanical testing. Prior to mechanical testing, the dissected tendons

were scanned by microCT and the tendon cross-sectional area was calculated. Mechanical

testing was performed, as previously described [17], on a DMA850 dynamic mechanical

analyzer (TA Instruments, New Castle, DE) with TRIOS Software. Tendons were gripped

between the calcaneus and myotendinous junction and underwent a uniaxial tensile testing

protocol consisting of a preload to 0.02 N, hold at 5% strain for 5 min with 1 min recovery, and

then ramp to failure test at 0.1% strain per second.

2.7. Statistical analysis

Power analysis (G*Power v.3.1, Heinrich Heine Universität Düsseldorf, Germany) performed

using experimental data, with power of 0.80 and significance level α=0.05, determined that a

minimum sample size of N=5 biological replicates (different mice) per experimental group was

needed. Mechanical testing and heterotopic ossification results were analyzed using Prism 10

(10.2.2, Graphpad, San Diego, CA) by Mann Whitney U test (one tailed) and two-way ANOVA

with uncorrected Fisher's LSD, respectively, to identify differences between groups. The

threshold for significance for all tests was set as $p<0.05$. The data met the assumptions of the

statistical approach.

3. Results

3.1. Both C57BL/6J and MRL/MpJ mouse Achilles tendons presented with scar formation after

tenotomy.

C57BL/6J and MRL/MpJ mice received unilateral tenotomy of the Achilles tendon. Four weeks

after surgery, injured Achilles tendons from both male and female mice presented as scar tissue

rich in fibroblasts and blood vessels, whereas uninjured Achilles tendons from age-, sex-, and

strain-matched mice appeared dense with extracellular matrix and aligned tenocytes. (Figs. 1A-

1H). The picrosirius red-stained sections showed disorganized immature collagen fibers in

injured tendons with brightfield (Figs. 1I, 1K, 1Q, 1S) and polarized (Figs. 1M, 1O, IU, 1W)

imaging, compared to aligned collagen fibers in uninjured tendons with brightfield (Figs. 1J, 1L,

1R, 1T) and polarized (Figs. 1N, 1P, 1V, 1X) imaging. There were no apparent differences in

histology between C57BL/6j and MRL/MpJ injured tendons.

3.2. MRL/MpJ male Achilles tendons showed better recovery of mechanical properties compared

to C57BL/6J male Achilles tendons.

Cross-sectional area measurements were similar between C57BL/6J and MRL/MpJ mouse

injured tendons regardless of sex (Fig. 2A). However, stiffness and Young's modulus were

significantly greater for MRL/MpJ male tendons than for C57BL/6J male tendons (Figs 2B and

2C, Male). Stiffness and Young's modulus of MRL/MpJ female tendons trended higher than for

C57L/6J female tendons, but were not significantly different (Figs 2B and 2C, Female).

3.3. MRL/MpJ male Achilles tendons possessed lower volume of heterotopic ossification

compared to C57BL/6J male Achilles tendons.

The mouse Achilles tendon tenotomy model has been studied as a heterotopic ossification model

based on the established phenomenon of chondroid degeneration followed by ectopic bone

formation in injured Achilles tendons by later timepoints of healing [18]. Injury-induced

heterotopic ossification was analyzed by micro CT scan ten weeks after surgery. Both C57BL/6J

and MRL/MpJ injured Achilles tendons induced heterotopic ossification (Fig. 3A). The total

volume of heterotopic ossified regions was smaller in MRL/MpJ male tendons than that in

C57BL/6J male tendons (Fig. 3B).

3.4. Activation of actin cytoskeleton and integrin signaling pathways was less pronounced in

MRL/MpJ male Achilles injured tendons.

Proteomic analysis of the samples obtained from C57BL/6J and MRL/MpJ male injured and

uninjured Achilles tendons revealed that 537 proteins and 78 proteins were up-regulated in

C57BL/6J injured tendons and MRL/MpJ injured tendons, respectively, whereas 594 proteins

and 54 proteins were down-regulated in C57BL/6J injured tendons and MRL/MpJ injured

tendons, respectively. Comparison between C57BL/6J and MRL/MpJ male injured tendons

showed that 221 proteins were up-regulated and 44 proteins were down-regulated in C57BL/6J

injured tendons compared to MRL/MpJ injured tendons. Based on IPA analysis, C57BL/6J and

MRL/MpJ injured tendons both activated the same top 5 pathways, including actin cytoskeleton

signaling and integrin signaling pathways relative to respective uninjured groups (Fig. 4A).

Interestingly, actin cytoskeleton signaling and integrin signaling pathways were also listed

amongst the top 5 pathways that were activated in C57BL/6J injured tendons relative to

MRL/MpJ injured tendons (Fig. 4B). This was also true in C57BL/6J uninjured tendons relative

to MRL/MpJ uninjured tendons (Fig. 4B). Volcano plots showed that, relative to uninjured

tendons, alpha-actinin and talin 1 protein levels were significantly higher in both C57BL/6J and

MRL/MpJ injured tendons (Fig. 4C). Alpha-actinin and talin 1 protein levels were also

significantly lower in MRL/MpJ injured tendons than in C57BL/6J injured tendons (Fig. 4C).

Immunoblot analysis corroborated proteomics data, showing that injured tendons (Injury)

contained higher levels of a-actinin and talin 1 than uninjured tendons (Normal) of both

C57BL/6J and MRL/MpJ mice. Talin 1 antibodies recognized intact and cleaved products as

previously reported [19]. The amount of talin1 tended to be higher in C57BL/6J injured tendons

than MRL/MpJ injured tendons (Fig. 4D).

4. Discussion

Our results indicate that tenotomized Achilles tendons in MRL/MpJ mice do not heal completely,

but have improved recovery of mechanical properties compared to those of C57BL/6J mice.

Further, this improved recovery may be associated with lesser activation of actin cytoskeleton

and integrin signaling pathways. These surprising and impactful findings demonstrate MRL/MpJ

mice possess limited capacity to regenerate severely injured tendons, but provide valuable data to

support study of the regulation of cytoskeleton and integrin signaling to promote tendon repair.

Superior healing by the MRL/MpJ mouse has been previously reported for patellar tendon

injuries [20]. After a 0.75 to 1 mm full-thickness defect in the patellar tendon, MRL/MpJ mice

exhibit full recovery of mechanical properties [10] and superior collagen fiber realignment [20],

compared to C57BL/6J mice. The patella tendons in the MRL/MpJ mice have also demonstrated

improved recovery of mechanical strength after full-length and full-thickness injuries [11],

although pathology of their healing process has not been reported. Kallenbach et al. [21] have

demonstrated that MRL tendons have reduced fibrotic adhesion in a flexor tendon injury model

with partial laceration. In the Achilles tendon tenotomy model that we used here, C57BL/6j mice

exhibit scar-mediated healing [18]. Neo-tissue forms in the gap of the transected tendon in the

first week after surgery, and the newly formed tissue gradually remodels over the next several

weeks, with a decrease in cell density and increase in longitudinal alignment of cells [18].

However, the remodeled tissue still contains poorly aligned, thin collagen fibers [8] and

undergoes chondroid degeneration and progressive heterotopic ossification 8 to 10 weeks after

injury [18, 22]. Here, we found that after complete transection, MRL/MpJ injured Achilles

tendons demonstrated similar pathology as observed in C57BL/6J mice, but the ectopic ossified

mass was smaller in MRL/MpJ mice than in C57BL/6J mice. These results show that healing of

the Achilles tendons after complete transection is scar-mediated, even in MRL/MpJ mice. A

complete transection of the Achilles tendon is more severe than the focal injury model in the

patellar tendon that previously show regenerative healing [10, 20, 23], suggesting that the

regenerative capacity of tendons in the MRL/MpJ mouse may be limited depending on the type

and severity of injury as well as the anatomical location of the tendon. Failure by the MRL/MpJ

mouse to regenerate after injury has been also demonstrated in other severely injured tissues,

including the heart after ischemia-reperfusion injury [9] and the skin after excisional injury [24].

Although injured Achilles tendons in the MRL/MpJ mice showed fibrotic healing, their

mechanical properties were superior compared to the injured Achilles tendons of the C57BL/6J

male mice. To examine potential underlying mechanisms, we performed proteomics analysis of

injured tendons during the remodeling phase, 4 weeks after injury, because previous studies

investigated the differences in the cells, matrix proteins, and signaling molecules between

MRL/MpJ and C57BL/6J mouse patellar tendons during the inflammation and proliferation

phases of early healing [10, 11, 20, 21, 23]. Here, the altered signaling pathways in injured

tendons during the remodeling phase were similar between MRL/MpJ and C57BL/6J mice,

suggesting that the remodeling molecular mechanism of injured tendons is similar in these

strains. Intriguingly, we found that the MRL/MpJ male tendons showed less activation of the

actin cytoskeleton and integrin signaling pathways than the C57BL/6J male tendons, results

which were validated by expression of alpha-actinin and talin-1, both of which are important

components of actin cytoskeleton and integrin signaling pathways [25, 26].

These pathways govern cell adhesion, migration, proliferation, and matrix synthesis and

organization [27]. Upregulation of these pathways in response to injury is expected and is likely

purposeful, especially during early healing phases after injury because inflammation and

angiogenesis occur during the inflammatory phase, and reparative cells such as tendon

progenitors migrate into and proliferate in the injured site during the proliferation phase [28]. In

regenerative healing, it would be expected that progenitor cells or other cells that are recruited

and/or proliferate at the injury site differentiate into tenocytes and then decline in number,

followed by the deposition of well aligned, relatively thick collagen fibers. In contrast, in scar-

mediated healing, the cells at the injury site persist without differentiating into tenocytes, and a

fibrous scar forms with randomly arranged, relatively thin collagen fibers. Although this scar-

mediated healing process has not been fully elucidated, myofibroblasts are key cells that

synthesize the fibrous tissue [29]. Myofibroblasts appear in wounded tissues during an early

phase and support and restore the integrity of damaged or defective tissues by depositing

extracellular matrix proteins [30]. Prolonged presence and activation of myofibroblasts result in

inhibition of regeneration of damaged tissue and stimulation of fibrosis [30]. Our findings

suggest that actin cytoskeletal signaling and integrin signaling are involved in regulation of

myofibroblast function in fibrous tissue formation because actin cytoskeletal proteins and

integrin adhesion complexes are essential molecules in regulation of migration, differentiation,

matrix synthesis, and organization of myofibroblasts [31]. Higher activity of these signaling

pathways may be caused by or be associated with higher activity of fibrous tissue formation,

leading to greater inhibition of structural recovery of collagen fibers and functional recovery of

mechanical properties in C57BL/6J mice.

Future studies should investigate the differences in the tendon healing processes between

MRL/MpJ and C57BL/6J mice over a longer period in this tenotomy injury model. Identification

of specific molecules that regulate the increased activity of the actin cytoskeleton and integrin

signaling pathways and validation of their role in injured C57BL/6J tendon healing is also

needed. Despite the unexpected discovery of limited MRL/MpJ tendon healing capacity

observed in this study, these data suggest that control of actin cytoskeleton and integrin signaling

pathways may be important in improving the mechanical properties of injured tendons.

Contributions: CKK and MEI designed the experiments. NT, KW, MW, HT, CDH, PR, and MEI

conducted experiments; NT, DSS, JJ, NL, MAK, CKK, and MEI evaluated the results; and CKK

and MEI prepared the manuscript. All authors have read and approved the final submitted

manuscript.

Funding: This work is supported by NIH / National Institute of Arthritis and Musculoskeletal and

Skin Diseases Grant/Award Numbers: R01AR070099 (MEI), R21AR080996 (MEI),

R01AR072886 (CKK), and R21AR077756 (CKK), University of Maryland Baltimore, Institute

for Clinical &Translational Research Pilot grant at #457 (MEI), and the United States (U.S.)

Department of Veterans Affairs Biomedical Laboratory R&D (BLRD) Service/ Career

Development Award Number IK2 BX004879 (NL).

Acknowledgments: We thank D. Muthusamy and E. Mirdamadi for assistance with the

DMA850.

References

[1] P. Sharma, N. Maffulli, Tendon injury and tendinopathy: healing and repair, J Bone Joint Surg

Am. 87(1)(2005) 187-202. 87/1/187 [pii]

10.2106/JBJS.D.01850

[2] J.D. Rees, A.M. Wilson, R.L. Wolman, Current concepts in the management of tendon

disorders, Rheumatology (Oxford). 45(5)(2006) 508-21. kel046 [pii]

10.1093/rheumatology/kel046

[3] H. Aslan, N. Kimelman-Bleich, G. Pelled, D. Gazit, Molecular targets for tendon

neoformation, J Clin Invest. 118(2)(2008) 439-44. 10.1172/JCI33944

[4] G. Nourissat, F. Berenbaum, D. Duprez, Tendon injury: from biology to tendon repair, Nat

Rev Rheumatol. 11(4)(2015) 223-33. 10.1038/nrrheum.2015.26

[5] L.M. Galatz, L. Gerstenfeld, E. Heber-Katz, S.A. Rodeo, Tendon regeneration and scar

formation: The concept of scarless healing, J Orthop Res. 33(6)(2015) 823-31.

10.1002/jor.22853

[6] K. Howell, C. Chien, R. Bell, D. Laudier, S.F. Tufa, D.R. Keene, et al., Novel Model of

Tendon Regeneration Reveals Distinct Cell Mechanisms Underlying Regenerative and Fibrotic

Tendon Healing, Sci Rep. 72017) 45238. 10.1038/srep45238

[7] P.K. Nguyen, C. Hart, K. Hall, I. Holt, C.K. Kuo, Establishing in vivo and ex vivo chick

embryo models to investigate fetal tendon healing, Sci Rep. 13(1)(2023) 9600. 10.1038/s41598-

023-35408-w

[8] S. Izumi, T. Oichi, S.S. Shetye, K. Zhang, K. Wilson, M. Iwamoto, et al., Inhibition of

glucose use improves structural recovery of injured Achilles tendon in mice, J Orthop Res.

40(6)(2022) 1409-19. 10.1002/jor.25176

[9] M.F. Rai, L.J. Sandell, Regeneration of articular cartilage in healer and non-healer mice,

Matrix Biol. 392014) 50-5. 10.1016/j.matbio.2014.08.011

[10] N.S. George, R. Bell, J.J. Paredes, P.J. Taub, N. Andarawis-Puri, Superior mechanical

recovery in male and female MRL/MpJ tendons is associated with a unique genetic profile, J

Orthop Res. 39(6)(2020) 1344-54. 10.1002/jor.24705

[11] A.L. Lalley, N.A. Dyment, N. Kazemi, K. Kenter, C. Gooch, D.W. Rowe, et al., Improved

biomechanical and biological outcomes in the MRL/MpJ murine strain following a full-length

patellar tendon injury, J Orthop Res. 33(11)(2015) 1693-703. 10.1002/jor.22928

[12] J.E. Marturano, J.D. Arena, Z.A. Schiller, I. Georgakoudi, C.K. Kuo, Characterization of

mechanical and biochemical properties of developing embryonic tendon, Proc Natl Acad Sci U S

A. 110(16)(2013) 6370-5. 10.1073/pnas.1300135110

[13] P.K. Nguyen, A. Jana, C. Huang, A. Grafton, I. Holt, M. Giacomelli, et al., Tendon

mechanical properties are enhanced via recombinant lysyl oxidase treatment, Front Bioeng

Biotechnol. 102022) 945639. 10.3389/fbioe.2022.945639

[14] S.H. Korntner, A. Jana, E. Kinnard, E. Leo, T. Beane, X. Li, et al., Craniofacial tendon

development-Characterization of extracellular matrix morphology and spatiotemporal protein

distribution, Front Cell Dev Biol. 102022) 944126. 10.3389/fcell.2022.944126

[15] M.M. Weldemariam, P.R. Sudhir, J. Woo, Q. Zhang, Effects of multiple stressors on

pancreatic human islets proteome reveal new insights into the pathways involved, Proteomics.

23(19)(2023) e2300022. 10.1002/pmic.202300022

[16] Y. Perez-Riverol, J. Bai, C. Bandla, D. Garcia-Seisdedos, S. Hewapathirana, S.

Kamatchinathan, et al., The PRIDE database resources in 2022: a hub for mass spectrometry-

based proteomics evidences, Nucleic Acids Res. 50(D1)(2022) D543-D52.

10.1093/nar/gkab1038

[17] I. Kurtaliaj, M. Golman, A.C. Abraham, S. Thomopoulos, Biomechanical Testing of Murine

Tendons, J Vis Exp. (152)(2019)   10.3791/60280

[18] S. Asai, S. Otsuru, M.E. Candela, L. Cantley, K. Uchibe, T.J. Hofmann, et al., Tendon

progenitor cells in injured tendons have strong chondrogenic potential: the CD105-negative

subpopulation induces chondrogenic degeneration, Stem Cells. 32(12)(2014) 3266-77.

10.1002/stem.1847

[19] N. Bate, A.R. Gingras, A. Bachir, R. Horwitz, F. Ye, B. Patel, et al., Talin contains a C-

terminal calpain2 cleavage site important in focal adhesion dynamics, PLoS One. 7(4)(2012)

e34461. 10.1371/journal.pone.0034461

[20] J. Paredes, D.A. Shiovitz, N. Andarawis-Puri, Uncorrelated healing response of tendon and

ear injuries in MRL highlight a role for the local tendon environment in driving scarless healing,

Connect Tissue Res. 59(5)(2018) 472-82. 10.1080/03008207.2018.1485665

[21] J.G. Kallenbach, M.A.T. Freeberg, D. Abplanalp, R.G. Alenchery, R.E. Ajalik, S. Muscat, et

al., Altered TGFB1 regulated pathways promote accelerated tendon healing in the superhealer

MRL/MpJ mouse, Sci Rep. 12(1)(2022) 3026. 10.1038/s41598-022-07124-4

[22] K. Zhang, S. Asai, M.W. Hast, M. Liu, Y. Usami, M. Iwamoto, et al., Tendon mineralization

is progressive and associated with deterioration of tendon biomechanical properties, and requires

BMP-Smad signaling in the mouse Achilles tendon injury model, Matrix Biol. 52-542016) 315-

24. 10.1016/j.matbio.2016.01.015

[23] J. Paredes, A. Pekmezian, N. Andarawis-Puri, MRL/MpJ tendon matrix-derived therapeutic

promotes improved healing outcomes in scar-mediated canonical tendon healing, J Orthop Res.

39(7)(2020) 1548-60. 10.1002/jor.24754

[24] A.S. Colwell, T.M. Krummel, W. Kong, M.T. Longaker, H.P. Lorenz, Skin wounds in the

MRL/MPJ mouse heal with scar, Wound Repair Regen. 14(1)(2006) 81-90. 10.1111/j.1524-

475X.2005.00092.x

[25] M. Nakamura, J. Hui, S.M. Parkhurst, Bending actin filaments: twists of fate, Fac Rev.

122023) 7. 10.12703/r/12-7

[26] Z. Baster, L. Russell, Z. Rajfur, A Review of Talin- and Integrin-Dependent Molecular

Mechanisms in Cancer Invasion and Metastasis, Int J Mol Sci. 26(5)(2025) 1798.

10.3390/ijms26051798

[27] R.O. Hynes, Integrins: bidirectional, allosteric signaling machines, Cell. 110(6)(2002) 673-

87. S0092867402009716 [pii]

[28] S. Thomopoulos, W.C. Parks, D.B. Rifkin, K.A. Derwin, Mechanisms of tendon injury and

repair, J Orthop Res. 33(6)(2015) 832-9. 10.1002/jor.22806

[29] J.E. Ackerman, S.N. Muscat, E. Adjei-Sowah, A. Korcari, A.E.C. Nichols, M.R. Buckley, et

al., Identification of Periostin as a critical niche for myofibroblast dynamics and fibrosis during

tendon healing, Matrix Biol. 1252024) 59-72. 10.1016/j.matbio.2023.12.004

[30] F.S. Younesi, B. Hinz, The Myofibroblast Fate of Therapeutic Mesenchymal Stromal Cells:

Regeneration, Repair, or Despair?, Int J Mol Sci. 25(16)(2024)   10.3390/ijms25168712

[31] F.S. Younesi, B. Hinz, The Myofibroblast Fate of Therapeutic Mesenchymal Stromal Cells:

Regeneration, Repair, or Despair?, Int J Mol Sci. 25(16)(2024) 8712.   10.3390/ijms25168712

Figure legend

Figure 1. Comparison of histology of injured tendons between C57BL/6J and MRL/MpJ mice.

A complete transverse incision was made at the midpoint of the left Achilles tendon in 12-week-

old female and male C57BL/6J and MRL/MpJ mice (n=3 per sex and strain). Longitudinal

sections of the injured and contralateral uninjured Achilles tendons four weeks after surgery were

made and stained with hematoxylin-eosin (HE) (A-H) and Picrosirius red (PSR) (I-L and Q-T,

bright field; M-P and U-X, polarized image). Bar represents 800 μm for HE staining and 1,500

μm for Picrosirius staining.

Figure 2. Comparison of biomechanical properties of injured tendons between C57BL/6J and

MRL/MpJ mice. A complete transverse transection was made at the midpoint of the left Achilles

tendon in 12-week-old female and male C57BL/6J (B6, blue; n=8 per sex) and MRL/MpJ (MRL,

red; n=5 for female and n=7 for male) mice. Four weeks after surgery, injured (left) and

contralateral uninjured (right) limbs were harvested. Dissected tendons were mechanically

assessed using tensile testing. (A) Cross-sectional area (CSA), (B) Stiffness, (C)Young's

modulus. Data points represent individual biological replicates, with averages and standard

deviations indicated. CSA averages and standard deviations were 0.601+/-0.0987 for C57BL/6J

female, 0.616+/-0.111 for C57BL/6J male, 0.571+/-0.0687 for MRL/MpJ female and 0.719+/-

0.128 mm^2 for MRL/MpJ male. Stiffness averages and standard deviations were 1.90+/-0.586

for C57BL/6J female, 2.93 +/- 1.02 for C57BL/6J male, 2.86 +/- 1.22 for MRL/MpJ female and

4.74 +/- 1.57 N/mm for MRL/MpJ male. Young's modulus averages and standard deviations

were 34.9+/-11.6 for C57BL/6J female, 48.8+/-13.3 for C57BL/6J male, 47.7+/-6.47 for

MRL/MpJ female and 72.4+/-23.1 MPa for MRL/MpJ male.

Figure 3. Comparison of heterotopic ossification in injured tendons between C57BL/6J and

MRL/MpJ mice. A complete transverse transection was made at the midpoint of the left Achilles

tendon in 12-week-old male C57BL/6J (blue) and MRL/MpJ (red) mice (n=5 per strain). Ten

weeks after surgery, injured (left) limbs were harvested. (A) Heterotopic ossification was imaged

by microCT. Bar represents 1 mm. (B) Total volume of ossified regions within the injured tendon

was analyzed. Data points represent individual biological replicates, with averages and standard

deviations indicated. Averages and standard deviations were 1.65+/-1.219 for C57BL/6j,

0.588+/-0.348 mm^3 for MRL/MpJ mice.

Figure 4. Proteomics analysis of injured tendons of C57BL/6J and MRL/MpJ mice. A complete

transverse transection was made at the midpoint of the left Achilles tendon in 12-week-old male

C57BL/6J (blue) and MRL/MpJ (red) mice. Four weeks after surgery, injured (left) and

contralateral uninjured (right) tendons were harvested, and proteomics analysis was performed.

The top 5 most altered signaling pathways were identified based on IPA analysis of differentially

expressed proteins between injured and uninjured tendons of C57BL/6J and MRL/MpJ mice (A),

and between C57BL/6J and MRL/MpJ tendons (B). (C) Volcano plot of proteins that differ

between injured and uninjured tendons in C57BL/6J (left) and MRL/MpJ (middle), and between

injured tendons of C57BL/6J and MRL/MpJ mice (right). (D) Immunoblots of α-actinin and

talin 1 in injured and uninjured tendons of C57BL/6J (C57) and MRL/MpJ (MRL) mice. Talin 1

antibodies recognized intact and cleaved products (239 and 190 kD products, respectively). The

same amount of protein was loaded for each group. Bars indicate the positions of molecular size

markers for α-actinin and talin 1 blots, and arrowheads indicate α-actinin and talin 1,

respectively.

Figure 1

Figure 2

Figure 3

A. Injured tendons to uninjured tendons

Ingenuity Canonical Pathways	-log(p-value)	z-score
C57BL/6j		
EIF2 Signaling	2.73E+01	3.9
Remodeling of Epithelial Adherens Junctions	2.14E+01	2.828
Actin Cytoskeleton Signaling	1.94E+01	2.744
Regulation of eIF4 and p70S6K Signaling	1.79E+01	0.707
Integrin Signaling	1.60E+01	3.053
MRL/MpJ		
EIF2 Signaling	2.62E+01	3.9
Remodeling of Epithelial Adherens Junctions	1.95E+01	2.828
Regulation of eIF4 and p70S6K Signaling	1.72E+01	1.414
Actin Cytoskeleton Signaling	1.68E+01	2.263
Integrin Signaling	1.62E+01	3.286

B. C57BL/6j to MRL/MpJ

Ingenuity Canonical Pathways	-log(p-value)	z-score
Injured tendon		
Actin Cytoskeleton Signaling	1.36E+01	2.294
Integrin Signaling	1.32E+01	2.985
Signaling by Rho Family GTPases	1.10E+01	2.828
ILK Signaling	1.09E+01	1.147
RHOGDI Signaling	1.02E+01	-1.387
Uninjured tendons		
Actin Cytoskeleton Signaling	8.04E+00	2.673
Regulation of eIF4 and p70S6K Signaling	7.92E+00	0.816
Epithelial Adherens Junction Signaling	7.80E+00	1.387
Remodeling of Epithelial Adherens Junctions	7.36E+00	NaN
Integrin Signaling	7.15E+00	3.207

C. Volcano plot

D. Immunoblot