Achilles tendons of MRL/MpJ mice show scar-mediated healing after tenotomy

Ningfeng Tang[1], Divya S. Sivalingam[2], Kimberly Wilson[1], Mehari M. Weldemariam[3], Hongying Tian[1], Christoph D. Hart[2], Prishasai Ramnath[2], Jacob Glazier[2], Jie Jiang[1], Natalie Leong[1], Tao Lowe[2,4], Maureen A. Kane[3], Masahiro Iwamoto[1], Catherine K. Kuo[1,2*], Motomi Enomoto-Iwamoto[1*]

[1]Department of Orthopaedics, University of Maryland School of Medicine, Baltimore, Maryland, USA

[2]Fischell Department of Bioengineering, A. James Clark School of Engineering, University of Maryland, College Park, Maryland, USA

[3]Department of Pharmaceutical Sciences, University of Maryland School of Pharmacy, Baltimore, Maryland, USA

[4]Department of Oral and Maxillofacial Surgery, School of Dentistry, University of Maryland, Baltimore, Maryland, USA

*Corresponding Authors:

Contact corresponding author
Catherine K. Kuo
Fischell Department of Bioengineering
A. James Clark School of Engineering
University of Maryland
4108 A. James Clark Hall, 8278 Paint Branch Dr
College Park, MD 20742, USA
ckk@umd.edu

Motomi Enomoto-Iwamoto
Department of Orthopaedics
University of Maryland School of Medicine
670 W Baltimore St, HSFIII 7171
Baltimore, MD 21201, USA
Motomi.iwamoto@som.umaryland.edu

Key words: tendon, injury, C57BL/6J, MRL/MpJ, proteomics

Abstract

Tendon healing in Murphy Ross Large (MRL/MpJ) mice was examined using an Achilles tendon

tenotomy model, a full transection model, compared to a patella focal injury model in which

tendons in MRL/MpJ mice showed regenerative healing. After complete transection, the Achilles

tendons of MRL/MpJ mice healed via scar formation, regardless of sex, as did C57BL/6J mice.

Tensile testing found that mechanical properties of injured tendons of both MRL/MpJ female and

male mice exhibited comparable levels to those of C57BL/6J female and male mice, respectively

after 4 weeks of healing, which is during the remodeling phase. After 10 weeks of healing,

injured tendons of MRL/MpJ mice possessed smaller heterotopic ossification volumes than those

of C57BL/6J mice. Proteomics analysis revealed similar alterations to signaling pathways in

injured tendons of MRL/MpJ and C57BL/6J male mice after 4 weeks of healing. However,

among the altered pathways, actin cytoskeleton and integrin signaling pathways were two of the

top pathways that were more prominently activated in C57BL/6J males than in MRL/MpJ males.

These findings indicate that MRL/MpJ mice possess limited capacity to regenerate injured

tendons after complete rupture, and that tenotomized Achilles tendons of MRL/MpJ male mice

have lesser induction of heterotopic ossification and lower activation of the signaling pathways

also induced with injury in C57BL/6J male mice.

1. Introduction

Tendons have limited repair capacity. Injured tendons do not usually regain their original levels

of biological, structural, and mechanical characteristics, and instead can undergo chronic

degeneration [1-5]. In contrast to embryonic and neonatal tendons [6,7], adult tendons heal via scar

formation characterized by high cell density, disorganized cell arrangement, thin and randomly

aligned collagen fibers, high vascularity, and mucoid matrix [5,8]. Scar formation inhibits recovery

of mechanical properties. Failure of full functional recovery may result in chronic tendon

disorders and re-rupture, leading to poor quality of life and work disability [5]. Murphy Roths

Large (MRL/MpJ) mice have strong repair capacity in musculoskeletal tissues including articular

cartilage, muscle, and bone [9]. MRL/MpJ mice also exhibit more regenerative healing of focal

injuries of the patellar tendon in adults, based on better recovery of mechanical properties and

improved extracellular matrix and cell alignment when compared to healing of the same injury in

C57BL/6 mice [10,11]. A flexor tendon injury model involving partial laceration has also shown that

MRL/MpJ tendons heal with reduced fibrotic adhesion [12].

Based on these compelling prior works, we were interested in investigating whether tendons in

MRL/MpJ mice undergo regenerative healing even in an Achilles tendon complete transection

model, which is different from the focal injury models previously used to study tendon repair in

MRL/MpJ mice [10-12]. Determining the molecular mechanism that supports the reparative ability

of the MRL/MpJ is crucial to develop new modalities to promote effective tendon healing, and

thus we also used proteomics analyses to examine which signaling pathways may be involved in

improved tendon repair by MRL/MpJ mice.

2. Materials and Methods

2.1. Mouse experiments

All animal studies were conducted with approval from the Institutional Animal Care and Use

Committee at the University of Maryland, Baltimore (0120007 and 1122006). Animals were

housed in a University Laboratory Animal Resources supervised animal facility with a 12-hour

light/dark cycle and controlled temperature of $22 \pm 1^\circ$C and humidity of $55 \pm 5\%$. Animals were

provided hygienic animal bedding of wood shavings and cotton pads. The health status of each

animal was monitored throughout the experiments by investigators, animal veterinary

technicians, and veterinarians, as per institutional guidelines. The mice were free of all viral,

bacterial, and parasitic pathogens during the experimental schedule.

A complete transverse transection (tenotomy), without attempt at repair, was made at the

midpoint of the left Achilles tendon in 12-week-old female and male C57BL/6J (#000646, The

Jackson Laboratory, Bar Harbor, ME) and MRL/MpJ (#000486, The Jackson Laboratory) mice

under anesthesia. The analgesic carprofen, 4.5 mg/Kg was administrated pre- and post-surgery.

The mice receiving the tenotomy surgery were euthanized 4 or 10-weeks post-surgery, which

was during the early and late remodeling phase, respectively. The right and left hind limbs were

collected for histological analysis and mechanical tests. The mid-substance of Achilles tendons

was snap frozen in liquid nitrogen and stored at -80°C until use. The criteria for animal

exclusion during the experiment, determined prior to the experiment, were to exclude mice that

received analgesics and/or antibiotics and/or were euthanized due to infection or distress. No

mice were excluded in this study. No exclusion criteria were set for data points during the

analysis. No data points were excluded. MRL/MpJ and C57BL/6J mice were maintained in the

same location and compared, but randomization was not used and confounders were not

controlled. A total of 109 mice were used: 3 female C57BL/6J, 3 male C57BL/6J, 3 female

MRL/MpJ, and 3 male MRL/MpJ mice for histological examination at 4 weeks post- surgery; 8

female C57BL/6J, 5 female MRL/MpJ, 8 male C57BL/6J, and 7 female MRL/MpJ mice for

mechanical testing at 4 weeks post-surgery; 5 male C57BL/6J, 5 male MRL/MpJ, 4 female

C57BL/6J, and 3 female MRL/MpJ mice for μCT analysis at 10 weeks post-surgery; 18

uninjured male C57BL/6J mice (for 5 unique biological replicates), 10 injured male C57BL/6J

mice (for 5 unique biological replicates), 10 uninjured male MRL/MpJ mice (for 4 unique

biological replicates), and 8 injured male MRL/MpJ mice (for 4 unique biological replicates) for

proteomics at 4-weeks post-surgery; and 3 C57BL/6J and 3 MRL/MpJ male mice for

immunoblotting at 4 weeks post-surgery. Healing ability comparisons between MRL/MpJ and

C57BL/6J mice were based on mechanical properties (primary outcome measure) and histology

(secondary outcome measure).

2.2. Histology

The hind limbs were fixed with 4% paraformaldehyde for 24 hours at 4°C and decalcified with

10% ethylenediaminetetraacetic acid at pH 7.2. Serial sagittal sections (5 μm) of the injured or

contralateral uninjured Achilles tendons were made. Sections were stained with hematoxylin-

eosin and picrosirius red. Hematoxylin-eosin staining images were captured using light

microscopy (BZX-700, Keyence, Tokyo, Japan). Picrosirius red staining images were captured

using polarized light microscopy (20X objective, ZEISS Axio Scan, Germany) [7,13-15].

2.3. Micro-computed tomography (microCT) analysis

Samples were scanned using a Skyscan 1172 microCT scanner (Bruker, Billerica, MA, USA)

while immersed in phosphate buffered saline without Ca^{2+} or Mg^{2+} (PBS). Specifically, images

were acquired using previously reported conditions [16]: 6.5 μm/pixel; 0.2 mm Al filter using a 55

kV X-ray source; resolution at 2 K with step rotation of 0.51; and two average frames using 360°

rotation. Image reconstruction was performed using NRecon software (Bruker). We manually

identified regions of heterotopic ossification by setting threshold between 80 and 255, for which

volume analyses were performed using CTAn software (Bruker). Rendering images were

obtained using CTVOX (Bruker).

2.4. Mass spectrometry-based proteomics

We collected tendon samples and stored them at -80 ºC until assay. Lysis buffer containing 5%

SDS (sodium dodecyl sulfate) and 50 mM triethylammonium bicarbonate (1 M, pH 8.0) was

added to tendons and the samples were then homogenized using bead beating with a Precellys

CK14 lysing kit (Bertin Corp., Rockville, MD). Next, protein extraction and digestion were

performed as described [17] using S-trap micro columns (ProtiFi, NY). Digested peptides were

quantified using a BCA assay kit (Thermo Fisher Scientific, 23275) to determine peptide

concentration. LC-MS/MS-based proteomic analysis was conducted on a nanoACQUITY Ultra-

Performance Liquid Chromatography system (Waters Corporation, Milford, MA USA) coupled

to an Orbitrap Fusion Lumos Tribrid mass spectrometer (Thermo Scientific, San Jose, CA USA).

Peptide separation was effected on a nanoACQUITY Ultra-Performance Liquid Chromatography

(UPLC) analytical column (BEH130 C18, 1.7 μm, 75 μm x 200 mm; Waters Corporation,

Milford, MA, USA) using a 185-min linear gradient with 3-40% acetonitrile and 0.1% formic

acid. Mass spectrometry conditions were as follows: Full MS scan resolution of 240,000,

precursor ions fragmentation by high-energy collisional dissociation of 35%, and a maximum

cycle time of 3 seconds. The resulting mass spectra were processed using Thermo Proteome

Discoverer ((PD, version 3.0.0.757), Thermo Fisher Scientific) and searched against a UniProt

mouse reference proteome (release 2024.03, 17180 entries) using Sequest HT algorithm. Search

parameters include carbamidomethylation of cysteines (+57.021 Da) as a static modification,

methionine oxidation (+15.995 Da) as a dynamic modification, precursor mass tolerance of 20

ppm, fragment mass tolerance of 0.5 Da, and trypsin as a digestion enzyme. Tryptic missed

cleavages were restricted to a maximum of two, with peptide lengths set between 6 and 144

residues. For protein quantification, the Minora feature detector, integrated in the PD, was used

as described previously [17]. To ensure high data quality, proteins were further filtered to a 1%

false discovery rate (FDR) threshold, calculated with the Percolator algorithm. Next, protein

abundance values exported from PD were post-processed using Perseus software (version

1.6.14.0). Proteins with missing values were excluded to improve data quality. Differentially

expressed proteins (DEPs) were identified using a two-tailed Student's t-test (adjusted p-value <

0.05). Ingenuity Pathway Analysis (IPA) software (Qiagen, Germantown, MD) was used to

identify dysregulated pathways and biological processes. The data were deposited in the

ProteomeXchange via the PRIDE database, with the project accession number PXD057712.

2.6. Mechanical Testing

Achilles tendons with calcanei were fine dissected to ensure removal of all musculature and

surrounding soft tissue. Injured and contralateral uninjured tendons were hydrated in PBS

immediately after dissection and throughout the course of mechanical testing. Prior to

mechanical testing, the dissected tendons were suspended in a tube pinching the calcaneus with

the lid and scanned using a Skyscan 1172 microCT scanner (Camera pixel size, 9.00; Rotation

range, 360 deg; Image rotation, 3 deg; scanning time, about 9 mins). PBS was placed in the

bottom of the tube to prevent the tendon from drying. The volume and length of the tendon were

measured using CTAn software. The cross-sectional area (CSA) was then calculated as the ratio

of volume to length. Mechanical testing was performed, as previously described [18], on a

DMA850 dynamic mechanical analyzer (TA Instruments, New Castle, DE) with TRIOS

Software. Tendons were gripped between the calcaneus and myotendinous junction and

underwent a uniaxial tensile testing protocol consisting of a preload to 0.02 N, and then ramp to

failure test at 0.1% strain per second. Mechanical (ultimate force, stiffness) and material (elastic

modulus, ultimate stress, ultimate strain) properties were determined, as previously described [19].

2.7. Statistical analysis

Power analysis (G*Power v.3.1, Heinrich Heine Universität Düsseldorf, Germany) performed

using experimental data, with power of 0.80 and significance level α=0.05, determined that a

minimum sample size of N=5 biological replicates (different mice) per experimental group was

needed. Data was assumed to be drawn from a normally distributed population. Normality was

analyzed by the Shapiro-Wilk test. Non-parametric tests were selected when the assumption of

normality could not be met. Mechanical testing and heterotopic ossification results were

analyzed using Prism 10 (10.6.0, Graphpad, San Diego, CA). Material and mechanical properties

were compared between sexes, mouse strains, and uninjured and injured tendons using a one-

way ANOVA with Tukey's post-hoc. Heterotopic ossification results were analyzed by Mann-

Whitney or Kologorov-Smirnov test to identify differences between groups. The threshold for

significance for all tests was set as $p < 0.05$. The data met the assumptions of the statistical

approach.

3. Results

3.1. Both C57BL/6J and MRL/MpJ mouse Achilles tendons presented with scar formation after

tenotomy.

C57BL/6J and MRL/MpJ mice received unilateral tenotomy of the Achilles tendon. To

investigate an early remodeling phase, we examined the recovery of extracellular matrix

organization at 4 weeks after surgery. Sagittal sections of injured Achilles tendons from both

female and male mice were acquired. Figure 1 shows sagittal sections of polarized images

stained with picrosirius red from the distal calcaneal insertion (left, single arrow) to the proximal

myotendinous junction (right, double arrow). Injured C57BL/6J tendons showed disorganized

immature collagen fibers (Fig. 1A and 1E), compared to aligned collagen fibers in uninjured

tendons of age- and sex-matched mice (Fig. 1B and 1F). Injured Achilles tendons of both female

and male MRL/MpJ mice also showed disorganized immature collagen fibers with polarized

imaging (Fig. 1C and 1G), in contrast to uninjured tendons (Fig. 1D and 1H). The hematoxylin

and eosin-stained sections of injured tendons presented as scar tissue rich in fibroblasts and

blood vessels (Fig. 2A-2H), whereas uninjured Achilles tendons from age-, sex-, and strain-

matched mice appeared dense with extracellular matrix and aligned tenocytes. (Fig. 2I-2L).

3.2. Recovery of material and mechanical properties of injured MRL/MpJ Achilles tendons were

comparable to that of C57BL/6J Achilles tendons.

Injured Achilles tendons of female and male MRL/MpJ mice possessed lower stiffness, elastic

modulus, ultimate load, and ultimate stress, but greater ultimate strain compared to their

corresponding uninjured MRL/MpJ Achilles tendons (Fig. 3, S1, and S2). Similarly, injured

female and male C57BL/6J tendon stiffness, elastic modulus, ultimate load, and ultimate stress

were lower, whereas ultimate strain was higher, than their corresponding uninjured Achilles

tendons (Fig. 3 and S2). In addition, the injury-induced changes in stiffness, elastic modulus,

ultimate load, ultimate stress, and ultimate strain were similar between MRL/MpJ and C57BL/6J

tendons (Fig. 3 and S2), suggesting that recovery of material and mechanical properties of

injured MRL/MpJ tendons is comparable to that of injured C57BL/6J tendons. Further, sex-

specific differences in material and mechanical properties in either injured or uninjured tendons

of either strain were not found (Fig. 3 and S2).

3.3. MRL/MpJ Achilles tendons possessed lower volume of heterotopic ossification compared to

C57BL/6J Achilles tendons.

The mouse Achilles tendon tenotomy model has been studied as a heterotopic ossification model

based on the established phenomenon of chondroid degeneration followed by ectopic bone

formation in injured Achilles tendons by later timepoints of healing [20]. Injury-induced

heterotopic ossification was analyzed by microCT scan ten weeks after surgery, the time when

heterotopic ossification is evident in this model [20]. Injury induced heterotopic ossification in

both C57BL/6J and MRL/MpJ Achilles tendons (Fig. 4A). The total volume of heterotopic

ossified regions was smaller in MRL/MpJ male tendons than that in C57BL/6J male tendons

(Fig. 4B, left). Combining male and female results also revealed significantly smaller volumes of

heterotopic ossification in MRL/MpJ mice (Fig. 4B, right).

3.4. Activation of actin cytoskeleton and integrin signaling pathways was less pronounced in

MRL/MpJ male Achilles injured tendons.

We performed proteomic analysis of the samples obtained from C57BL/6J and MRL/MpJ male

injured and uninjured Achilles tendons. Differentially expressed proteins (DEPs) were identified

using a two-tailed Student's t-test with a criterion of adjusted p-value < 0.05 (Fig. 5A). We found

that 538 proteins and 595 proteins were up-regulated in C57BL/6J injured tendons and

MRL/MpJ injured tendons, respectively, compared with the mouse strain-matched uninjured

tendons, whereas 79 proteins and 54 proteins were down-regulated in C57BL/6J injured tendons

and MRL/MpJ injured tendons, respectively. Comparison between C57BL/6J and MRL/MpJ

male injured tendons showed that 335 proteins were up-regulated and 70 proteins were down-

regulated in C57BL/6J injured tendons compared to MRL/MpJ injured tendons. In spite of

differences in DEPs between C57BL/6J and MRL/MpJ samples, based on IPA, C57BL/6J and

MRL/MpJ injured tendons revealed similar alterations of the canonical pathways relative to their

corresponding uninjured tendons (Supplementary Table 1). When the top 100 pathways were

compared between C57BL/6J and MRL/MpJ groups, the top 50 pathways overlapped

completely, and exhibited differences only in the rank order (common pathways in

Supplementary Table 1, highlighted). Both C57BL/6J and MRL/MpJ activated the same top 5

pathways with positive z-scores, including actin cytoskeleton signaling and integrin signaling

pathways relative to respective uninjured groups (Fig. 5B). Interestingly, actin cytoskeleton

signaling and integrin signaling pathways were also listed amongst the top 5 pathways in

C57BL/6J injured tendons relative to MRL/MpJ injured tendons (Fig. 5C, z-score 2.294 and

2.985, respectively). This was also observed in C57BL/6J uninjured tendons relative to

MRL/MpJ uninjured tendons (Fig. 5C).

4. Discussion

Our results indicate that tenotomized Achilles tendons of MRL/MpJ mice heal similarly as those

of C57BL/6J mice, but with reduced heterotopic ossification than in C57BL/6J tendons by late

stages of healing. Further, the injured tendons in MRL/MpJ male mice had lesser activation of

actin cytoskeleton and integrin signaling pathways. These findings demonstrate MRL/MpJ mice

possess limited capacity to regenerate injured tendons after complete rupture, but with reduced

heterotopic ossification and activation of some molecular signaling pathways during tendon

healing, such as cytoskeleton and integrin signaling, compared to C57BL/6J mice.

Superior healing by the MRL/MpJ mouse has been previously reported for patellar tendon

injuries [21]. After a 0.75 to 1 mm full-thickness focal defect in the patellar tendon, MRL/MpJ

mice exhibit greater recovery of mechanical properties [10] and superior extracellular matrix

realignment [21], compared to C57BL/6J mice. MRL/MpJ mice have also demonstrated improved

recovery of mechanical strength after full-length and full-thickness central patellar tendon

injuries [11], although pathology of their healing process has not been reported. In the Achilles

tendon tenotomy model that we used here, C57BL/6J mice exhibit scar-mediated healing [20].

Neo-tissue forms in the gap of the transected tendon in the first week after surgery, and the newly

formed tissue gradually remodels over the next several weeks, with a decrease in cell density and

increase in sagittal alignment of cells [20]. However, the remodeled tissue still contains poorly

aligned, thin collagen fibers [8] and undergoes chondroid degeneration and progressive heterotopic

ossification 8 to 10 weeks after injury [20,22]. Here, we found that after complete transection,

MRL/MpJ injured Achilles tendons demonstrated fibrotic healing as observed in C57BL/6J mice,

but the ectopic ossified mass was smaller in MRL/MpJ mice than in C57BL/6J mice.

Furthermore, by 4 weeks of healing, there was no difference in the extent of recovery of

mechanical properties between MRL/MpJ mice and C57Bl/6J mice regardless of sex. It appears

that completely transected Achilles tendons in MRL/MpJ mice undergo scar-mediated healing,

and do not have superior healing ability compared to C57BL/6J mice. However, additional

investigations are needed, including quantitative analysis of histology, as well as mechanical

testing and histological analysis at later stages.

In this study, we used a tendon injury model of complete transection of the Achilles tendon

whereas previous studies have used a focal injury model of the patellar tendon that has shown

regenerative healing [10,21,23]. Considering the many differences between the tenotomized Achilles

tendon injury model versus the patella tendon focal injury model, such as tendon type, injury

size, and instability of injury site by motion, it is not possible to make generalized conclusions

about the healing ability of MRL/MpJ tendons. In addition, the Achilles tendon injury model has

limitations, including the development of ectopic bone formation, and the lack of maintaining the

tendon ends in apposition following transection and during healing. Nevertheless, our results

suggest that the regenerative capacity of tendons in the MRL/MpJ mouse may be limited

depending on the type of injury as well as the anatomical location of the tendon. It has been

suggested by others that MRL/MpJ regenerative healing extent depends on injury type or

severity [11]. Failure by the MRL/MpJ mouse to fully regenerate after injury has been

demonstrated in other injured tissues, including the heart after ischemia-reperfusion injury [9,24],

the skin after excisional injury [25], and articular cartilage after full-thickness injury [26]. Here, we

did not observe sex-specific differences in elastic modulus, ultimate stress, and ultimate strain of

injured Achilles tendons 4 weeks after injury. This is consistent with previous findings that

MRL/MpJ patellar tendon focal defects do not exhibit sex-specific differences in mechanical or

material properties [27]. However, male MRL/MpJ mice exhibit superior healing of articular

cartilage full-thickness injuries compared to female MRL/MpJ mice [26]. Interestingly, the

heterotopic calcified volume in the injured Achilles tendons trended larger in males than in

females, regardless of strain: $0.838 +/- 0.600$ mm^3 versus $1.648 +/- 0.122$ mm^3 for C57BL/6J

female versus male mice, respectively; and $0.209 +/- 0.189$ mm^3 versus $0.588 +/- 0.348$ mm^3 for

MRL/MpJ female versus male mice, respectively. In injured mouse Achilles tendons, heterotopic

bone formation occurs via endochondral ossification at late healing stage [20]. Taken together, male

MRL/MpJ mice may have a stronger ability to generate cartilage tissue than female mice,

resulting in the induction of larger volumes of cartilage-mediated ectopic bone in healing

tendons.

To examine the molecular mechanisms involved in tendon healing, we performed proteomics

analysis of injured tendons during the remodeling phase, 4 weeks after injury, because previous

studies investigated the differences in the cells, matrix proteins, and signaling molecules between

MRL/MpJ and C57BL/6J mouse patellar tendons during the inflammation and proliferation

phases of early healing [10-12,21,23]. Here, the altered signaling pathways in injured tendons during

the remodeling phase were similar between MRL/MpJ and C57BL/6J mice, suggesting that in

both strains, similar signaling pathways are altered upon injury, and similar molecular

mechanisms are involved in scar formation and matrix remodeling. Intriguingly, we found that

the MRL/MpJ male tendons showed less activation of the actin cytoskeleton and integrin

signaling pathways than the C57BL/6J male tendons. These pathways govern cell adhesion,

migration, proliferation, and matrix synthesis and organization [28]. Upregulation of these

pathways in response to injury is expected and is likely purposeful, especially during early

healing phases after injury because inflammation and angiogenesis occur during the

inflammatory phase, and reparative cells such as tendon progenitors and myofibroblasts migrate

into and proliferate in the injured site during the proliferation phase [29]. Actin cytoskeletal

proteins and integrin adhesion complexes are essential molecules in regulation of migration,

differentiation, matrix synthesis, and organization of myofibroblasts [30]. It is of note that early

upregulation of integrins and cellular adhesion genes have also been implicated in superior

MRL/MpJ patellar tendon focal defect healing [27]. Higher expression of *Itgb1* during the first 7

days post-injury was postulated to lead to improved cell and matrix alignment in MRL/MpJ

healing patellar tendons, compared to C57BL/6J tendons. In contrast, during the remodeling

phase, higher activity of these signaling pathways may be caused by or be associated with higher

activity of fibrous tissue formation, leading to greater inhibition of structural recovery of

collagen fibers and functional recovery of mechanical properties in C57BL/6J mice.

Future studies should investigate the differences in the tendon healing processes between

MRL/MpJ and C57BL/6J mice over a longer period in this tenotomy injury model. In particular,

material property characterizations – including elastic modulus, ultimate strength, and collagen

crosslink density – should be determined after additional longer healing durations to more fully

evaluate long-term recovery of function and restoration of tissue quality. Quantitative

assessments of histological characterizations, including fibrillar collagen density and alignment,

cell and nuclear morphology, and integrin binding are needed to better inform the extent of

scarless healing in MRL/MpJ mice. Identification of specific molecules that regulate the

increased activity of the actin cytoskeleton and integrin signaling pathways and validation of

their role in injured C57BL/6J tendon healing is also needed. Despite the unexpected discovery

of limited MRL/MpJ tendon healing capacity observed in this study, these data suggest that

lower activity of the actin cytoskeleton and integrin signaling pathways may be associated with a

lower degree of heterotopic ossification. Complementing our current data with future proteomic

characterizations of healing MRL/MpJ versus C57BL/6J female Achilles tendons, along with

validation of specific protein targets, will provide more comprehensive insight into the sex-

specific MRL/MpJ regenerative healing capacity for tendon injuries.

Contributions: CKK and MEI designed the experiments. NT, KW, MW, HT, CDH, PR, and MEI

conducted experiments; NT, DSS, JJ, NL, MAK, CKK, and MEI evaluated the results; and CKK

and MEI prepared the manuscript. All authors have read and approved the final submitted

manuscript.

Funding: This work is supported by NIH / National Institute of Arthritis and Musculoskeletal and

Skin Diseases Grant/Award Numbers: R01AR070099 (MEI), R21AR080996 (MEI),

R01AR072886 (CKK), and R21AR077756 (CKK), University of Maryland Baltimore, Institute

for Clinical &Translational Research Pilot grant at #457 (MEI).

Acknowledgments: We thank D. Muthusamy and E. Mirdamadi for assistance with the

DMA850.

References

[1]   Sharma P, Maffulli N. Tendon injury and tendinopathy: healing and repair. J Bone Joint

Surg Am. Jan 2005;87(1):187-202. doi:87/1/187 [pii]

10.2106/JBJS.D.01850

[2]   Rees JD, Wilson AM, Wolman RL. Current concepts in the management of tendon

disorders. Rheumatology (Oxford). May 2006;45(5):508-21. doi:kel046 [pii]

10.1093/rheumatology/kel046

[3]   Aslan H, Kimelman-Bleich N, Pelled G, Gazit D. Molecular targets for tendon

neoformation. J Clin Invest. Feb 2008;118(2):439-44. doi:10.1172/JCI33944

[4]   Nourissat G, Berenbaum F, Duprez D. Tendon injury: from biology to tendon repair. Nat

Rev Rheumatol. Apr 2015;11(4):223-33. doi:10.1038/nrrheum.2015.26

[5]   Galatz LM, Gerstenfeld L, Heber-Katz E, Rodeo SA. Tendon regeneration and scar

formation: The concept of scarless healing. J Orthop Res. Jun 2015;33(6):823-31.

doi:10.1002/jor.22853

[6]   Howell K, Chien C, Bell R, et al. Novel Model of Tendon Regeneration Reveals Distinct

Cell Mechanisms Underlying Regenerative and Fibrotic Tendon Healing. Sci Rep. Mar 23

2017;7:45238. doi:10.1038/srep45238

[7]   Nguyen PK, Hart C, Hall K, Holt I, Kuo CK. Establishing in vivo and ex vivo chick

embryo models to investigate fetal tendon healing. Sci Rep. Jun 13 2023;13(1):9600.

doi:10.1038/s41598-023-35408-w

[8]   Izumi S, Oichi T, Shetye SS, et al. Inhibition of glucose use improves structural recovery

of injured Achilles tendon in mice. J Orthop Res. Jun 2022;40(6):1409-1419.

doi:10.1002/jor.25176

[9]   Rai MF, Sandell LJ. Regeneration of articular cartilage in healer and non-healer mice.

Matrix Biol. Oct 2014;39:50-5. doi:10.1016/j.matbio.2014.08.011

[10] George NS, Bell R, Paredes JJ, Taub PJ, Andarawis-Puri N. Superior mechanical

recovery in male and female MRL/MpJ tendons is associated with a unique genetic profile. J

Orthop Res. Apr 30 2020;39(6):1344-1354. doi:10.1002/jor.24705

[11] Lalley AL, Dyment NA, Kazemi N, et al. Improved biomechanical and biological

outcomes in the MRL/MpJ murine strain following a full-length patellar tendon injury. J Orthop

Res. Nov 2015;33(11):1693-703. doi:10.1002/jor.22928

[12] Kallenbach JG, Freeberg MAT, Abplanalp D, et al. Altered TGFB1 regulated pathways

promote accelerated tendon healing in the superhealer MRL/MpJ mouse. Sci Rep. Feb 22

2022;12(1):3026. doi:10.1038/s41598-022-07124-4

[13] Marturano JE, Arena JD, Schiller ZA, Georgakoudi I, Kuo CK. Characterization of

mechanical and biochemical properties of developing embryonic tendon. Proc Natl Acad Sci U S

A. Apr 16 2013;110(16):6370-5. doi:10.1073/pnas.1300135110

[14] Nguyen PK, Jana A, Huang C, et al. Tendon mechanical properties are enhanced via

recombinant lysyl oxidase treatment. Front Bioeng Biotechnol. 2022;10:945639.

doi:10.3389/fbioe.2022.945639

[15] Korntner SH, Jana A, Kinnard E, et al. Craniofacial tendon development-Characterization

of extracellular matrix morphology and spatiotemporal protein distribution. Front Cell Dev Biol.

2022;10:944126. doi:10.3389/fcell.2022.944126

[16] Garcia SA, Wilson K, Tang N, et al. Analysis of the Actions of RARγ Agonists on

Growing Osteochondromas in a Mouse Model. Int J Mol Sci. Jul 11 2024;25(14).

doi:10.3390/ijms25147610

[17] Weldemariam MM, Sudhir PR, Woo J, Zhang Q. Effects of multiple stressors on

pancreatic human islets proteome reveal new insights into the pathways involved. Proteomics.

Oct 2023;23(19):e2300022. doi:10.1002/pmic.202300022

[18] Kurtaliaj I, Golman M, Abraham AC, Thomopoulos S. Biomechanical Testing of Murine

Tendons. J Vis Exp. Oct 15 2019;(152)doi:10.3791/60280

[19] Navarro J, Korcari A, Nguyen P, et al. Method development and characterization of chick

embryo tendon mechanical properties. J Biomech. Mar 2022;133:110970.

doi:10.1016/j.jbiomech.2022.110970

[20] Asai S, Otsuru S, Candela ME, et al. Tendon progenitor cells in injured tendons have

strong chondrogenic potential: the CD105-negative subpopulation induces chondrogenic

degeneration. Stem Cells. Dec 2014;32(12):3266-77. doi:10.1002/stem.1847

[21] Paredes J, Shiovitz DA, Andarawis-Puri N. Uncorrelated healing response of tendon and

ear injuries in MRL highlight a role for the local tendon environment in driving scarless healing.

Connect Tissue Res. Sep 2018;59(5):472-482. doi:10.1080/03008207.2018.1485665

[22]   Zhang K, Asai S, Hast MW, et al. Tendon mineralization is progressive and associated

with deterioration of tendon biomechanical properties, and requires BMP-Smad signaling in the

mouse Achilles tendon injury model. Matrix Biol. May-Jul 2016;52-54:315-24.

doi:10.1016/j.matbio.2016.01.015

[23]   Paredes J, Pekmezian A, Andarawis-Puri N. MRL/MpJ tendon matrix-derived therapeutic

promotes improved healing outcomes in scar-mediated canonical tendon healing. J Orthop Res.

May 22 2020;39(7):1548-1560. doi:10.1002/jor.24754

[24]   Abdullah I, Lepore JJ, Epstein JA, Parmacek MS, Gruber PJ. MRL mice fail to heal the

heart in response to ischemia-reperfusion injury. Wound Repair Regen. 2005;13(2):205-8.

doi:10.1111/j.1067-1927.2005.130212.x

[25]   Colwell AS, Krummel TM, Kong W, Longaker MT, Lorenz HP. Skin wounds in the

MRL/MPJ mouse heal with scar. Wound Repair Regen. Jan-Feb 2006;14(1):81-90.

doi:10.1111/j.1524-475X.2005.00092.x

[26]   Fitzgerald J, Rich C, Burkhardt D, Allen J, Herzka AS, Little CB. Evidence for articular

cartilage regeneration in MRL/MpJ mice. Osteoarthritis Cartilage. Nov 2008;16(11):1319-26.

doi:10.1016/j.joca.2008.03.014

[27]   George NS, Bell R, Paredes JJ, Taub PJ, Andarawis-Puri N. Superior mechanical

recovery in male and female MRL/MpJ tendons is associated with a unique genetic profile. J

Orthop Res. Jun 2021;39(6):1344-1354. doi:10.1002/jor.24705

[28]   Hynes RO. Integrins: bidirectional, allosteric signaling machines. Cell. Sep 20

2002;110(6):673-87. doi:S0092867402009716 [pii]

[29]   Thomopoulos S, Parks WC, Rifkin DB, Derwin KA. Mechanisms of tendon injury and

repair. J Orthop Res. Jun 2015;33(6):832-9. doi:10.1002/jor.22806

[30]   Younesi FS, Hinz B. The Myofibroblast Fate of Therapeutic Mesenchymal Stromal Cells:

Regeneration, Repair, or Despair? Int J Mol Sci. Aug 9 2024;25(16):8712.

doi:10.3390/ijms25168712

<u>Figure Legends</u>

Fig. 1. Comparison of picrosirius red (PSR)-stained sections of injured tendons between

C57BL/6J and MRL/MpJ mice. A complete transverse transection was made at the midpoint of

the left Achilles tendon in 12-week-old female and male C57BL/6J (A, B, E, and F) and

MRL/MpJ (C, D, G, and H) mice (n=3 per sex and strain). Sagittal sections of the injured (A, C,

E, and G) and contralateral uninjured (B, D, F, and H) Achilles tendons four weeks after surgery

were stained with PSR. Bar represents 1.5 mm. Arrow and double arrow show distal calcaneal

insertion and proximal myotendinous junction of the Achilles tendon, respectively.

Fig. 2. Comparison of hematoxylin-eosin-stained sections of injured tendons between C57BL/6J

and MRL/MpJ mice. A complete transverse transection was made at the midpoint of the left

Achilles tendon in 12-week-old female and male C57BL/6J (A-D, I and J) and MRL/MpJ (E-H,

K and L) mice (n=3 per sex and strain). Sagittal sections of the injured (A-H) and contralateral

uninjured (I-L) Achilles tendons four weeks after surgery were stained with hematoxylin-eosin

(HE).

Fig. 3. Comparison of biomechanical properties of injured tendons between C57BL/6J and

MRL/MpJ mice. A complete transverse transection was made at the midpoint of the left Achilles

tendon in 12-week-old female (circles; F) and male (squares; M) C57BL/6J (blue; n=7 for female

and n=8 for male) and MRL/MpJ (red; n=5 for female and n=7 for male) mice. Four weeks after

surgery, injured (Inj; left limb) and contralateral uninjured (Uninj; right limb) limbs were

harvested. Dissected tendons were mechanically assessed using tensile testing. (A) Cross-

sectional area (CSA), (B) stiffness, (C) ultimate load, (D) elastic modulus, (E) ultimate stress,

and (F) ultimate strain were compared between uninjured and injured tendons of C57BL/6J

female (blue circles), MRL/MpJ female (red circles), C57BL/6J male (blue squares), and

MRL/MpJ male (red squares). Data points represent individual biological replicates. For (A-F),

p-values of statistically significant comparisons between uninjured and injured of the same sex

and strain, between sexes of the same injury status and strain, and between strains of the same

injury status and sex are shown. Averages and standard deviations of absolute values are

summarized in Fig. S2 of Supplementary Material. (G-I) Injured values were normalized to

average uninjured values of the respective group and (G) elastic modulus, (H) ultimate stress,

and (I) ultimate strain were compared between female and male C57BL/6J and MRL/MpJ mice.

Data points represent individual biological replicates, with averages and standard deviations

indicated. Normalized elastic modulus averages and standard deviations were 0.123 +/- 0.065 for

C57BL/6J female, 0.160 +/- 0.090 for MRL/MpJ female, 0.145 +/- 0.079 for C57BL/6J male,

and 0.238 +/- 0.159 for MRL/MpJ male. Normalized ultimate stress averages and standard

deviations were 0.189 +/- 0.056 for C57BL/6J female, 0.310 +/- 0.168 for MRL/MpJ female,

0.250 +/- 0.083 for C57BL/6J male, and 0.349 +/- 0.149 for MRL/MpJ male. Normalized

ultimate strain averages and standard deviations were 1.722 +/- 0.334 for C57BL/6J female,

2.177 +/- 0.466 for MRL/MpJ female, 1.826 +/- 0.431 for C57BL/6J male, and 1.838 +/- 0.455

for MRL/MpJ male. No statistically significant differences were found between normalized

mechanical properties between strains of the same sex or sexes of the same strain.

Fig. 4. Comparison of heterotopic ossification in injured tendons between C57BL/6J and

MRL/MpJ mice. A complete transverse transection was made at the midpoint of the left Achilles

tendon in 12-week-old male (square) and female (circle) C57BL/6J (blue) and MRL/MpJ (red)

mice (n=5 per strain). Ten weeks after surgery, injured (left) limbs were harvested. (A)

Heterotopic ossification was imaged by microCT. Bar represents 1 mm. (B) Total volume of

ossified regions within the injured tendon was analyzed. Data points represent individual

biological replicates, with averages and standard deviations indicated. Averages and standard

deviations were 1.65+/-1.219 for C57BL/6J male, 0.588+/-0.348 for MRL/MpJ male, 1.24+/-

0.573 for C57BL/6J male and female and 0.400+/-0.268 mm^3 MRL/MpJ male and female mice.

Fig. 5. Proteomics analysis of injured tendons of C57BL/6J and MRL/MpJ mice. A complete

transverse transection was made at the midpoint of the left Achilles tendon in 12-week-old male

C57BL/6J (blue) and MRL/MpJ (red) mice. Four weeks after surgery, injured (left) and

contralateral uninjured (right) tendons were harvested, and proteomics analysis was performed.

(A) The top 50 upregulated and downregulated proteins in injured tendons samples in both

C57BL/6J and MRL/MpJ male mice were well-clustered based on t-test comparisons. (B, C).

The top 5 most altered signaling pathways were identified based on IPA analysis of differentially

expressed proteins between injured and uninjured tendons of C57BL/6J and MRL/MpJ mice (B),

and between C57BL/6J and MRL/MpJ tendons (C). Positive and negative z-scores represent

upregulation and downregulation of the corresponding pathway, respectively.

Supplemental Figure Captions

Fig. S1. Representative force-displacement curves for uninjured and injured tendons of both

strains and sexes.

Fig. S2. Averages and standard deviations of absolute values for cross-sectional area (CSA),

stiffness, ultimate load, elastic modulus, ultimate stress, and ultimate strain of uninjured and

injured tendons of C57BL/6J female, MRL/MpJ female, C57BL/6J male, and MRL/MpJ male

mice.

517
518

Figure 1

Figure 2

Injured

C57BL/6J	A · 2 mm	B · 2 mm
	C · 500 µm	D · 500 µm
MRL/MpJ	E · 2 mm	F · 2 mm
	G · 500 µm	H · 500 µm

Uninjured

C57BL/6J	I · 500 µm	J · 500 µm
MRL/MpJ	K · 500 µm	L · 500 µm

Female   Male

Figure 4

A

B

Figure 5

A   T-test comparisons

C57BL/6J

MRL/MpJ

B   Injured tendons to uninjured tendons

C57BL/6J

C   C57BL/6J to MRL/MpJ

Injured tendon

MRL/MpJ

Supplementary Material (Fig. S1 and S2)

Achilles tendons of MRL/MpJ mice show scar-mediated healing after tenotomy

Ningfeng Tang[1], Divya S. Sivalingam[2], Kimberly Wilson[1], Mehari M. Weldemariam[3], Hongying Tian[1], Christoph D. Hart[2], Prishasai Ramnath[2], Jacob Glazier[2], Jie Jiang[1], Natalie Leong[1], Tao Lowe[2,4], Maureen A. Kane[3], Masahiro Iwamoto[1], Catherine K. Kuo[1,2]*, Motomi Enomoto-Iwamoto[1]*

*Corresponding Authors

Fig. S1. Representative force-displacement curves for uninjured and injured tendons of both strains and sexes.

Mean ± standard deviation of mechanical properties								
	Female				Male			
	C57BL/6J		MRL/MpJ		C57BL/6J		MRL/MpJ	
	Uninjured	Injured	Uninjured	Injured	Uninjured	Injured	Uninjured	Injured
CSA (mm²)	0.17 ± 0.01	0.58 ± 0.13	0.27 ± 0.04	0.57 ± 0.07	0.20 ± 0.03	0.62 ± 0.11	0.29 ± 0.05	0.72 ± 0.13
Stiffness (N/mm)	5.32 ± 0.83	2.33 ± 0.85	10.17 ± 2.62	3.04 ± 1.74	6.72 ± 2.96	2.92 ± 1.03	9.57 ± 2.67	5.16 ± 2.96
Ultimate load (N)	6.23 ± 1.28	4.14 ± 1.08	7.90 ± 0.83	5.19 ± 2.67	7.89 ± 2.28	6.15 ± 1.94	8.82 ± 1.57	7.90 ± 4.12
Elastic Modulus (MPa)	169.32 ± 50.35	20.78 ± 10.96	194.92 ± 16.68	31.17 ± 17.50	197.88 ± 84.48	28.79 ± 15.59	176.84 ± 48.43	42.08 ± 28.16
Ultimate Stress (MPa)	37.07 ± 7.91	7.01 ± 2.08	30.21 ± 4.33	9.36 ± 5.08	40.89 ± 15.68	10.21 ± 3.38	31.67 ± 7.16	11.04 ± 4.73
Ultimate Strain (mm/mm)	0.37 ± 0.04	0.64 ± 0.12	0.24 ± 0.02	3.04 ± 0.11	0.32 ± 0.09	0.59 ± 0.14	0.29 ± 0.06	0.53 ± 0.13

Fig. S2. Averages and standard deviations of absolute values for cross-sectional area (CSA), stiffness, ultimate load, elastic modulus, ultimate stress, and ultimate strain of uninjured and injured tendons of C57BL/6J female, MRL/MpJ female, C57BL/6J male, and MRL/MpJ male mice.