Intensity Modulation of Trichromatic Split Fluorescent Proteins for Live Cell Mapping

Mamoru Ishii[1,2*], Tomoaki Kinjo[3], Yohei Kondo[4,5], Kenta Terai[6], Kazuhiro Aoki[1], Brian Kuhlman[3,7], Michiyuki Matsuda[1,8,9*]

[1]Graduate School of Biostudies, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan.

[2]Institute of Industrial Science, The University of Tokyo, Tokyo, Japan

[3]Department of Biochemistry and Biophysics, University of North Carolina School of Medicine, Chapel Hill, NC, USA

[4]Center for One Medicine Innovative Translational Research (COMIT), Nagoya University, Nagoya, Aichi, Japan

[5]Graduate School of Medicine, Nagoya University, Tsurumai-cho, Nagoya, Aichi 466-8550, Japan

[6]Graduate School of Medicine, Tokushima University, Shinkura-cho, Tokushima 770-8501, Japan

[7]Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[8]Affiliate Graduate School, Graduate School of Medicine, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan.

[9]Integrated Graduate School of Medicine, Engineering, and Agricultural Sciences, University of Yamanashi, Chuo-shi, Yamanashi 409-3898 Japan.

*Corresponding authors. Center for Living Systems Information Science, Graduate School of Biostudies, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan

Lead correspondence: matsuda.michiyuki.2c@kyoto-u.ac.jp (M.M.), mishii@iis.u-tokyo.ac.jp (M.I.)

Running title

Multiplexed cell labeling with split FPs

Keywords

Split fluorescent proteins, Multiplexed spectral labeling, Cell labeling, Fluorescence microscopy, Protein tagging

Abstract

Multiplexed live cell-labeling technologies are essential tools for investigating complex biological systems, yet existing approaches face limitations in achieving reliable discrimination of multiple cell populations. Here, we present color and intensity modulation with split fluorescent proteins for cell labeling (Caterpie), a live cell labeling system that enables high-fidelity identification of multiple cell populations through rational design of split fluorescent proteins. Drawing inspiration from human trichromatic vision, we engineered optimized split fluorescent protein variants—split CFP2, split mNG3A, and split sfCherry3C—and developed a comprehensive library of the 11th β-strand (FP_{11}) tags comprising variable numbers of tandem repeats. Structure-guided protein engineering enhanced the performance of split mNG3A and split sfCherry3C through targeted modifications of their C-terminal sequences and binding interfaces, respectively. By systematically varying the composition and arrangement of FP_{11} repeats, we selected 20 distinct FP_{11} tags that enable robust cell population discrimination with 97% accuracy using conventional fluorescence microscopy. In these FP_{11} tag-expressing cells, we expressed multiple EGF ligands and their receptors and cultured them in bulk to examine the effects of EGF signaling on chemotaxis within the cell population. The results showed that chemotaxis changes according to the intensity of EGFR signaling. The Caterpie system holds significant potential for applications requiring precise identification and tracking of multiple cell populations in complex biological systems.

Introduction

Multiplexed spectral labeling techniques represent essential tools across diverse biological disciplines, including neuroscience, developmental biology, and immunology. These techniques enable critical applications such as discrimination of adjacent cells, specific cell populations labeling, and lineage tracing. The Brainbow system exemplifies this approach, utilizing multiple gene cassettes designed to stochastically express one of three fluorescent proteins (FPs), thereby generating distinct spectral signatures (*1*). While this system has proven particularly valuable for morphological analyses including connectomics, it presents limitations for specific cell labeling and lineage tracing due to potential variations in the relative FP expression levels across cellular generations. The development of Bitbow addressed certain limitations of earlier systems by enabling the generation of up to 31 (2^5-1) distinct color variations through random toggling of five FP genes within a single gene cassette (*2*). However, this approach faces challenges due to biased gene selection partners, resulting in fewer achievable color discrimination than theoretically predicted. These limitations underscore the advantages of rationally designed color palettes. However, conventional approaches to expanding color diversity through serial concatenation of multiple FPs face practical constraints due to increasing gene sizes. Split fluorescent proteins emerge as an elegant solution to these technical challenges.

Split fluorescent proteins (split FPs) are generated through strategic dissection of β-barrel fluorescent proteins (FPs) into two components: the 11th β-strand (FP_{11}) and a complementary segment comprising the first ten β-strands (FP_{1-10}) (*3*). While neither fragment exhibits fluorescence independently, their co-expression facilitates spontaneous reassembly, enabling chromophore maturation and subsequent fluorescence emission. This complementation system has been successfully employed for protein labeling applications, wherein target proteins are tagged with FP_{11} in cells expressing FP_{1-10} (*4, 5*). A particularly advantageous feature of this system emerges under conditions of abundant FP_{1-10} expression: enhanced fluorescence intensity can be achieved by increasing the number of FP_{11} tags conjugated to the protein of interest, while maintaining relatively modest gene size requirements (*6–9*).

Drawing inspiration from human color perception, which discriminates diverse spectral signatures through the integration of three primary color intensities: blue, green, and red, we developed a novel multiplexed cellular labeling approach. This system exploits engineered arrays of FP_{11} fragments derived from cyan, green, and red split fluorescent proteins, each designed to complement with their cognate FP_{1-10} fragments. This method, named color and intensity modulation with split fluorescent proteins for cell labeling (Caterpie), achieves robust discrimination of 20 distinct cell populations with 97% accuracy using conventional epi-fluorescence microscopy.

Results

Split fluorescent proteins for the trichromatic color palette

We aimed to establish a trichromatic color palette through the systematic engineering of tandem FP_{11} repeats derived from cyan, green, and red split fluorescent proteins, which undergo fluorescence complementation when co-expressed with their cognate FP_{1-10} fragments (Fig. 1A). For this, we sought to select split fluorescent proteins (split FPs) that achieve brightness comparable to their full-length FP ($FP_{Full-length}$) and exhibit fluorescence signal amplification through FP_{11} repeats. A critical consideration in designing this system is the requirement for sequence divergence among the trichromatic split FPs to prevent cross-complementation between fragments from different fluorescent protein pairs (*10*). To address this constraint, we evaluated split FPs derived from distinct evolutionary lineages: sfGFP, mNG2, sfCherry2, and mRuby4 (Fig. 1B). To quantitatively assess the relative brightness of $FP_{Full-length}$, $FP_{11}(x1)$, and $FP_{11}(x4)$ of reported split FPs, we developed a dual-reporter system. We engineered expression constructs encoding Histone H2B-tagged $FP_{Full-length}$, $FP_{11}(x1)$, or $FP_{11}(x4)$ fused to EBFP2-nls (nuclear localization signal) through a self-cleaving P2A site, enabling normalization of expression levels through nuclear EBFP2 fluorescence. These constructs were co-transfected into HeLa cells with a secondary plasmid encoding FP_{1-10} linked to iRFP670 via an internal ribosome entry site (IRES) at a DNA ratio of 1:4 (Fig. 1C).

After 48 hours, cells were observed under a confocal microscope, and split FP fluorescence intensities were quantified in iRFP670 positive cells (Fig. 1D). For each fluorescent protein variant, relative fluorescence intensity values were normalized to the full-length FP. In most cases, (x1) shows decreased fluorescence brightness compared to full-length, but sfGFP, mNG3A, and mNG3K were almost equivalent fluorescence brightness. Comparative analysis of (x1) versus (x4) constructs revealed that all sfGFP-derived split FPs exhibited more than two-fold enhancement in median brightness, with CFP2 demonstrating the most substantial increase of more than three-fold. Consistent with previous findings (*11*), split mNG3A demonstrated superior amplification from (x1) to (x4) compared to split mNG3K. Split sfCherry3C$_{11}$(x1) exhibited diminished fluorescence intensity relative to sfCherry3C$_{Full-length}$, attributable to reduced association efficiency, corroborating earlier observations (*12*). The truncated mRuby4$_{11}$ showed markedly decreased fluorescence intensity, highlighting the critical role of its C-terminal unstructured polypeptide chain (*10*). These preliminary findings indicated that while the combination of CFP2, mNG3A, and sfCherry3C provides a promising foundation for the trichromatic system, although further engineering optimization of mNG3A and sfCherry3C is necessary to achieve optimal performance across all spectral channels.

Fluorescence intensity-based cell classification using tandem CFP2 β-strand repeats

Before the engineering of mNG3A and sfCherry3C, we established a platform for cell identification through fluorescence intensity modulation using CFP2$_{11}$ repeats. To this end, we

constructed plasmids encoding repeated sequences of CFP2$_{11}$ (Fig. 2A). Initially, we

engineered a plasmid encoding a Histone H2B-CFP2$_{11}$(x1) fusion construct, in which the

CFP2$_{11}$(x1) sequence was flanked by a BglII restriction site at its 5' terminus and BamHI and

NotI restriction sites at its 3' terminus. Through sequential ligation of BglII/NotI-digested

fragments (insert) with BamHI/NotI-digested fragments (vector), we successfully amplified

CFP2$_{11}$ to create (x2), (x4), and (x8) variants. Based on previous research (6), we

incorporated GGSGG linker sequences between CFP2$_{11}$ fragments. At the BglII-BamHI

junction, the six nucleotides encode glycine and serine residues, thereby serving as an integral

part of the linker sequence. The CFP2$_{11}$ repeats [(x1), (x2), (x4), or (x8)] fused with Histone

H2B were introduced into HeLa cells stably expressing CFP2$_{1-10}$ (HeLa$^{CFP2\ 1-10}$) (Fig. 2B). As

a control, we expressed Histone H2B-tagged CFP2$_{Full-length}$ in the same HeLa$^{CFP2\ 1-10}$ cell line.

When normalized to nuclear mCherry fluorescence, the CFP2 fluorescence intensity increased

across consecutive constructs with a factor of approximately 1.7-fold, which was slightly

lower than the theoretically expected 2-fold increment (Fig. 2C).

To identify cells expressing different numbers of CFP2$_{11}$ repeats, we implemented a

Gaussian mixture model (GMM) for cell classification (Fig. 2D). The fluorescence intensities

of mCherry and CFP2 were transformed into polar coordinates, where the Angle$_{mCherry\ vs\ CFP2}$

(ranging from 0° to 90°) represented the CFP2/mCherry intensity ratio. The angular data were

divided into training and test datasets. We trained a GMM on the training dataset to establish

classification parameters, determining the decision boundary of the Angle$_{mCherry\ vs\ CFP2}$. The

model performance was validated using the test dataset, and the predicted repeat numbers

were compared with the actual repeat numbers. The model achieved an overall accuracy of

96% (Fig. 2E). Among all constructs, CFP2$_{11}$(x8) exhibited the lowest accuracy (93.9%), with

6.1% of (x8)-expressing cells being misclassified as (x4).

To validate the system's discriminative capacity in a heterogeneous context, we analyzed

mixed populations of cells expressing four different CFP2$_{11}$ repeat variants using fluorescence

microscopy and generated CFP2/mCherry ratio images (Fig. 2F). The trained GMM

demonstrated robust performance in classifying individual cells into (x1), (x2), (x4), or (x8)

populations with 96% accuracy, which was validated through visual inspection (Fig. 2G).

Furthermore, the GMM provided membership probabilities for each cell, enabling

quantitative assessment of classification reliability (Fig. 2H). These results established that

four-level intensity modulation of a single fluorescent protein is sufficient for reliable

population discrimination.

Computational design of split sfCherry3C for enhanced β-barrel stability and split-strand complementation

To optimize split sfCherry3C toward fluorescence intensities close to those of the full-

length protein, we modelled the sfCherry3C$_{1-10}$/sfCherry3C$_{11}$ complex with ColabFold (13)

(based on AlphaFold-Multimer) (14), and used the structure model for subsequent design with

Rosetta modeling suite (*15*). Because both global β-barrel stability and β_{1-10}/β_{11} interface

affinity are crucial for fluorescence of split FPs, we pursued two design strategies in parallel

(*16*). First, to enhance global β-barrel stability, we employed a Rosetta script adapted from the

previously described "Point-Mutation" (PM) workflow (*16*) and performed *in-silico* site-

saturation mutagenesis (SSM) across all 224 residues, ranking each variant by the change in

calculated energy (ΔE) calculated from the Rosetta *total_score* term (fig. S1). Next, to refine

the β_{1-10}/β_{11} interface, we repurposed the PM script for interface design; with the revised

script, we performed SSM at every β_{11} position and calculated the binding energy of each

single mutant from *dG_separated* score obtained with InterfaceAnalyzer protocol (Fig. 3A).

Each single-mutant model was subsequently processed by the "Mutation Cluster (MC)" (*16*)

workflow modified for interface design, in which FastDesign protocol with InterfaceDesign

relax script stochastically samples sequence and rotamer space at residues within 5–7 Å of the

"seed mutation" through Monte-Carlo design/minimization cycles, yielding multi-residue

mutants that are re-evaluated with InterfaceAnalyzer (Fig. 3B). After filtering by the scores

such as *total_score*, *dG_separated*, and mutation number, we conducted visual inspection to

retain the variants that reinforce interface contacts while excluding ones that disrupted the

chromophore-forming triad M67–Y68–G69 or introduced additional aromatic residues into

β_{11} which could promote aggregation of multimeric β_{11} repeat. Following these criteria, we

selected twenty-two candidate designs for subsequent in-cell validation.

Upon stable co-expression of β_{1-10} and β_{11} in HeLa cells, three variants (M1, M2, M3)

outperformed the parental split sfCherry3C in fluorescent intensity (Fig. 3C, D). We then

combined the mutations from the three top performing variants to generate an optimized

variant, named split sfCherry3Csp, which achieved a 2.5-fold enhancement in fluorescent

intensity compared to the parental split sfCherry3C. The M1 and M3 mutations locate at the

β_{1-10}/β_{11} interface, where the models suggest the improved association of β_{1-10} and β_{11} through

the formation of reinforced side-chain contacts. The M2 mutation lies on the opposite face of

the interface, implying an effect on β-barrel stability (Fig. 3E). The fluorescence spectra of

split sfCherry3Csp were indistinguishable to those of the parental split sfCherry3C, indicating

that the incorporated mutations enhanced the β-barrel stability and β_{1-10}/β_{11} association

without affecting the chromophore maturation (Fig. 3F).

197 Enhancement of complementation efficiency through structure-guided engineering of

198 split mNG3A

As shown in Fig. 1D, split mNG3A demonstrated only modest signal amplification, with a

1.3-fold increase in fluorescence intensity from (x1) to (x4) variants. To gain insight into the

structural factors underlying this limitation, we modelled the octameric $mNG3A_{11}$ peptide

using ColabFold (*13*) (based on AlphaFold2) (*17*). The predicted structures suggested that

$mNG3A_{11}$ repeat adopts a loosely helical conformation, with hydrophobic residues (F5, W8,

F12, M15, M16) converging to form a continuous hydrophobic interaction, despite only

moderate model confidence. These features implied that multimeric mNG3A$_{11}$ may self-

aggregate via its hydrophobic surfaces (Fig. 4A) (*17*). Additionally, inspection of the parental

mNG crystal structure (PDB: 5LTP) and the structure model of split mNG3A suggested that

the C-terminal residues D14–M15–M16 contribute minimally to the β-barrel fold (Fig. 4B).

We therefore reasoned that removing these C-terminal residues could mitigate the aggregation

propensity without perturbing β-barrel and chromophore maturation. Guided by these

insights, we implemented three C-terminal modification approaches: (1) deletion of the C-

terminal D14-M15-M16, (2) addition of one or two aspartic acid residue to preserve charge

that could electrostatically repel neighboring fragments, and (3) substitution of F12 with

tyrosine to introduce polarity (Fig. 4C). To evaluate these modifications, we expressed 8

repeats of the mNG3A$_{11}$ fragments fused to Histone H2B in HeLa cells expressing mNG3A$_{1-}$

$_{10}$ (Fig. 4D). Nuclear mNG3A fluorescence was quantified through microscopy, with

expression levels normalized to nuclear mTagBFP2 fluorescence intensity (Fig. 4E). Among

the modifications tested, we chose the mNG3Asp$_{11}$ variant for further analysis because of its

highest fluorescence intensity, which was 9.1-fold brighter than the parental mNG3A$_{11}$. The

fluorescence spectra obtained using the mNG3Asp$_{11}$ fragment were identical to those

observed with the mNG3A$_{11}$ fragment (Fig. 4F).

Performance evaluation of split mNG3Asp and split sfCherry3Csp was conducted in

parallel with split CFP2, following our previously established experimental framework (Fig.

4G). Quantitative analysis revealed that mNG3Asp$_{11}$(x4) achieved a 2.3-fold enhancement in

fluorescence intensity compared to its monomeric counterpart. The engineered sfCherry3Csp

demonstrated robust performance, with its monomeric variant exhibiting 80% of the

fluorescence intensity of the full-length protein, while the tetrameric construct showed a 3-

fold signal amplification relative to the monomer. These significant improvements in signal

amplification efficiency indicate that both engineered variants, split mNG3Asp and split

sfCherry3Csp, now achieve performance metrics comparable to split CFP2, providing the

basis for trichromatic imaging.

Optimization of the fusion partner for the FP$_{11}$ tag

Prior to establishing the FP$_{11}$ tag library, we aimed to reduce construct size by identifying

alternatives to the relatively large Histone H2B fusion tag. We evaluated multiple fusion

protein configurations, comparing Histone H2B (126 amino acids) with two smaller

alternatives: GB1 (56 amino acids) (*18*) and ΔSUMOstar (74 amino acids), a truncated

version of SUMOstar (*19*), as well as constructs lacking fusion tags (fig. S2A). ΔSUMOstar

was engineered by removing the unstructured polypeptide regions from both termini of

SUMOstar; the resulting sequence is detailed in Figure S2B. In agreement with previous

studies (*20*), constructs lacking fusion proteins demonstrated markedly reduced fluorescence

intensity, an effect that persisted even in the octameric (x8) variant (fig. S2C). Comparative

analysis of fusion partners revealed that ΔSUMOstar-tagged constructs achieved substantially

higher fluorescence intensity than GB1-tagged variants in octameric configurations,

demonstrating the importance of fusion partner selection for optimal signal amplification. The

ΔSUMOstar-tagged constructs exhibited consistent signal enhancement, with fluorescence

intensity increasing 1.6-fold for each doubling of $CFP2_{11}$ repeats, ultimately enabling 93%

discrimination accuracy between variants (fig. S2, D to F).

To optimize the modular arrangement of the three FP_{11} variants ($CFP2_{11}$, $mNG3Asp_{11}$,

and $sfCherry3Csp_{11}$), we conducted a systematic analysis of possible configurations. In initial

experiments, we generated ΔSUMOstar fusion constructs containing six total repeats, wherein

each FP_{11} fragment was represented in duplicate (fig. S3A). The complete set of six possible

permutations was generated and stably expressed in HeLa cells constitutively expressing all

three FP_{1-10} fragments. Quantitative analysis of relative fluorescence intensities was

performed using EBFP2 fluorescence for normalization of expression levels (fig. S3B).

Among all configurations evaluated, the $CFP2_{11}(x2)$-$sfCherry3Csp_{11}(x2)$-$mNG3Asp_{11}(x2)$

arrangement demonstrated marginally superior fluorescence intensity across all spectral

channels, leading to its selection for subsequent studies.

Selection of 20 optimal FP_{11} tags

Based on these preliminary findings, we established a comprehensive library of fusion

constructs combining ΔSUMOstar with $CFP2_{11}$, $sfCherry3Csp_{11}$, and $mNG3Asp_{11}$ fragments

in this order (Fig. 5A). Each fragment was represented in varying copy numbers [(x0), (x1),

(x2), (x4), or (x8)] for each. In the first screening, we limited the FP_{11} tag combinations to 96

variants, with the total number of repeats not exceeding (x12). The integrity of all FP_{11} repeat

sequences in the 96 plasmid constructs was verified by Sanger sequencing, thereby

establishing a comprehensive FP_{11} tag library.

To generate a stable cellular platform for evaluating the FP_{11} tag library, we engineered an

expression vector encoding all three FP_{1-10} fragments (CFP_{1-10}, $sfCherry3Csp_{1-10}$, and

$mNG3A_{1-10}$) separated by self-cleaving P2A peptides, followed by an IRES and iRFP670

(Fig. 5B). Through fluorescence-activated cell sorting (FACS) based on iRFP670 expression

levels, we established a HeLa cell clone ($HeLa^{FP1-10x3\_iRFP670}$) expressing high levels of three

FP_{1-10} fragments.

The complete library of 96 FP_{11} tags was stably expressed in $HeLa^{FP1-10x3\_iRFP670}$ cells,

with resultant populations analyzed by confocal microscopy. To quantitatively assess the

fluorescence distribution patterns, we implemented spherical coordinate transformation to

calculate two angular parameters: $Angle_{CvsG}$ and $Angle_{RvsCG}$, each ranging from 0° to 90° (Fig.

5C, left). To evaluate the discriminative power of these angular parameters, we performed

pairwise population analysis using selected cell populations from the library. Cell

classification was achieved using an unsupervised two-component Gaussian mixture model

without prior training data (Fig. 5C, right). Classification accuracy was determined by

comparing model-predicted clusters against known population identities. Superior

classification accuracy correlated with greater separation of populations in the Angle_{CvsG} –

Angle_{RvsCG} parameter space. This analytical framework was systematically applied to all

possible pairwise combinations within the 96 FP_{11}-tagged cell library (Fig. 5D).

To determine the number of distinguishable cell populations, we implemented an iterative

sampling approach due to the vast number of possible patterns (approximately 2×10^{20}) when

selecting 20 FP_{11}-tagged cells from a pool of 96 (Fig. 5E). Random subsets of predetermined

sizes were drawn from the 96 FP_{11}-tagged cell populations, with pairwise classification

accuracies extracted from the comprehensive analysis presented in Fig. 5D. For each FP_{11}-

tagged subset, we identified the minimum classification accuracy, reflecting the subset's

ability to distinguish the most similar cell population pairs. This sampling process was

repeated 10^7 times, and the maximum achievable minimum accuracy was plotted as a function

of subpopulation size (Fig. 5F). The analysis revealed that a subset of 20 cell populations

maintained robust discrimination with a minimum pairwise classification accuracy of 90%.

However, expanding to 30 populations resulted in a substantial decrease in minimum

accuracy to 60%. Based on these quantitative insights, we selected an initial panel of 30

candidate FP_{11} tags (Fig. 5A, 1st) for subsequent refinement to establish the final optimized set

of 20 tags (Fig. 5A 2nd) that would ensure maximal discriminative power.

We conducted comprehensive evaluation of the 30 candidate populations using both

confocal microscopy and flow cytometry, with three independent experimental replicates (fig.

S4). The relationship between population subset size and maximum achievable minimum

pairwise classification accuracy was reassessed using these orthogonal detection methods

(Fig. 5G). While fluorescence microscopy demonstrated marginally superior classification

accuracy, flow cytometry maintained robust discrimination capabilities, achieving >85%

accuracy across a set of top 20 FP_{11} tags. The final optimized set of 20 FP_{11} tags and their

comprehensive pairwise discrimination analysis is presented in Figure S5. Here, to optimize

the fluorescence signal intensity of single-color reference populations, we refined the copy

number composition (Fig. 5A Final): $CFP2_{11}$(x8) and $mNG3Asp_{11}$(x8) were implemented at

maximum repeat numbers, while $sfCherry3Csp_{11}$ was limited to (x4) to circumvent

aggregation phenomena observed at higher copy numbers (fig. S6).

High-fidelity discrimination of twenty cell populations using optimized FP_{1-10} expression

platform

The FP_{1-10} expression system was also refined by replacing the iRFP670 reporter with a

puromycin resistance cassette. Through stringent puromycin selection, we established a HeLa

clone cell ($HeLa^{FP1-10x3}$) expressing elevated levels of $CFP2_{1-10}$, $mNG3Asp_{1-10}$, and

$sfCherry3Csp_{1-10}$. The optimized set of 20 FP_{11} tags was stably expressed in $HeLa^{FP1-10x3}$ cells,

with individual cell populations analyzed by confocal microscopy. Fluorescence was

visualized with distinct spectral channels: CFP2 (blue), mNG3Asp (green), and sfCherry3Csp

(red). Merged images revealed unique signatures based on both color composition and

intensity distribution (Fig. 6A).

Cell segmentation was performed on maximum intensity projections of composite

fluorescence signals using Cellpose (*21*). Mean fluorescence intensities were quantified

across all three channels for each segmented cell. Following the methodology established in

Figure 5C, we implemented spherical coordinate transformation to calculate $Angle_{CvsG}$ and

$Angle_{RvsCG}$, generating a two-dimensional angular representation of relative fluorescence

intensities (Fig. 6B, left). These parameters were used to train a Gaussian mixture model

(GMM), which established both label assignment distributions and label membership

probability distributions in $Angle_{CvsG}$ and $Angle_{RvsCG}$ (Fig. 6B, center and right). Validation of

the trained GMM on independent test data demonstrated exceptional classification

performance, achieving 97% accuracy in discriminating all 20 cell populations (Fig. 6C).

Using unsupervised k-means clustering on the 20 components achieved 96% accuracy in

distinguishing between the 20 cell populations, even without requiring pre-training data for

individual fluorescence profiles (fig. S7).

To evaluate system performance under practical conditions, we used pooled samples

containing all 20 cell populations and acquired tiled images (Fig. 6D). Application of the

trained GMM enabled robust classification of individual cells, with 78% of cells being

classified with more than 95% probability (Fig. 6E).

Interaction dynamics in multi-population cell cultures of MDCK cells expressing EGF-

family ligands and their receptors

Finally, as an application of this system—named color and intensity modulation with split

fluorescent proteins for cell labeling (Caterpie)—we studied the contribution of heterotypic

cell interactions mediated by EGF-family ligands and their cognate receptors. For this, we

used Madin-Darby Canine Kidney (MDCK) cells, which are widely used to study collective

cell migration and cell competition regulated by the EGF signaling pathway. MDCK cells

were labeled with the aforementioned FP_{1-10} and 20 FP_{11} tags with 96% accuracy (fig. S8A,

B). Mixed cells were classified by applying a trained Gaussian mixture model (fig. S8C), 75%

of cells being classified with more than 95% probability. The cell densities of twenty different

labeled cell types at 72 hours post-labeling ranged within a 1.4-fold difference between the

minimum and maximum values (fig. S9).

Among the four EGF-family ligands expressed in MDCK cells, we focused on HBEGF

and EREG. HBEGF, a high-affinity ligand, binds to heparan sulfate proteoglycans to provide

strong signals restricted to short distances, promoting localized cell migration. In contrast,

EREG, a low-affinity ligand, diffuses quickly and remotely, propagating signals

approximately four times faster than high-affinity ligands and more efficiently to distant cells

(*22*). Both HBEGF and EREG bind to EGFR (ErbB1) and ErbB4. Using lentiviral

transduction, we introduced EREG, HBEGF, EGFR (ErbB1), or ErbB4 in fluorescently

labeled MDCK cells (Fig. 7A). To investigate how EGF ligands and receptors with different

affinities interact between heterotypic cells and generate spontaneous patterns, these four cell

types, along with wild-type cells, were co-cultured for 8 days under 3% low-serum

conditions. The resulting cell population was analyzed by fluorescence microscopy to identify

each cell type (Fig. 7B).

Before analyzing the positional relationships between heterotypic cells, we first examined

the distribution patterns of homotypic cells, as cell types with limited motility and dispersal

capability affect intercellular interactions with heterotypic cells. To examine homotypic cell

clustering, each cell type is individually displayed (Fig. 7C). Dispersion of each cell type was

analyzed by the Clark-Evans index; values less than 1 indicate cellular clustering. We did not

find significant homotypic cell clustering in any cell types (Fig. 7D). Then, we investigated

the frequency of adjacent heterotypic cells. When examining receptor-expressing cells

adjacent to EREG ligand or HBEGF-expressing cells (Fig. 7E), we found EGFR cells

displayed significantly higher adjacency to EREG cells than wild-type cells. Conversely,

when examining ligand-expressing cells adjacent to EGFR receptor or ErbB4-expressing

cells, HBEGF cells exhibited higher adjacency to EGFR cells and to a lesser extent to ErbB4

in comparison to wild-type cells. These results suggest that EREG and EGF function as

chemoattractants, with EREG weakly attracting distant cells, while HBEGF strongly attracts

nearby cells.

In conclusion, the Caterpie method provides a versatile platform for analyzing cell

populations that may exhibit homotypic and/or heterotypic clustering.

Discussion

We have developed color and intensity modulation with split fluorescent proteins for cell labeling (Caterpie), a novel cell identification platform that achieves robust identification of 20 distinct cell populations with 96-97% accuracy. This system leverages fluorescence intensity modulation through engineered arrays of split fluorescent protein (split FP) fragments. The technological foundation of Caterpie comprises three complementary split FPs: split CFP2 and two newly engineered variants - split mNG3Asp and split sfCherry3Csp. To implement this system, we established a comprehensive toolkit consisting of 20 distinct FP_{11} tags, each constructed by fusing truncated SUMOstar protein to the 11th β-strand of the three split fluorescent proteins in varying tandem repeat configurations. These engineered FP_{11} tags demonstrate highly specific labeling of FP_{1-10}-expressing cells, enabling population discrimination with 96-97% accuracy through Gaussian mixture model (GMM) analysis of the resulting multidimensional fluorescence signatures.

Current multicolor fluorescent labeling approaches can be broadly categorized into two distinct methodological frameworks. The first relies on stochastic recombination events mediated by site-specific recombinase systems such as Cre/loxP and Flp/FRT (*1, 2, 23–27*). The second approach exploits the inherent randomness of transfection efficiencies or genomic integration sites (*28–32*). While both established approaches depend on stochastic processes to generate fluorescent protein expression patterns, Caterpie represents a paradigm shift through its implementation of rationally designed color palettes. This deterministic approach enables precise targeting of specific cell populations and lineages. The system holds particular promise for applications requiring high-fidelity identification and longitudinal tracking of distinct cell types within complex biological contexts. However, the full realization of Caterpie's potential necessitates further optimization of delivery methodologies. Specifically, development of refined strategies is required to achieve selective incorporation of distinct FP_{11} tag variants into target cell populations while maintaining variant segregation and preventing cross-contamination.

The theoretical prediction of an 8-fold signal enhancement for $FP_{11}(x8)$ constructs relative to their monomeric counterparts was not fully realized in experimental measurements. Our studies with split CFP2 demonstrated approximately 5-fold enhancement in median fluorescence intensity (Fig. 2C). Further investigations revealed that the choice of fusion protein significantly impacts the signal amplification efficiency of octameric constructs (fig. S2C). In research using sfGFP, when x3, x4, or x7 $sfGFP_{11}$ was fused to β-tubulin or Lamin A/C, length and brightness correlated well, nearly achieving theoretical limits (*4, 6*). However, when Teneurin-m was fused to x7 $sfGFP_{11}$, only a 4-fold increase in brightness was achieved (*9*). With the goal of reducing gene size, which is an advantage of our approach, we changed from the initially used Histone H2B to ΔSUMOstar and achieved almost the same amplification rate as Histone H2B, x1.6 by each repetition (fig. S2). This collective evidence

suggests that achieving linear brightness amplification through multimerization of the 11th β-
strand requires careful optimization of the fusion partner.

The application of computational protein structure prediction proved instrumental in
enhancing split fluorescent protein performance (Fig. 3). This approach represents a
significant departure from conventional optimization strategies, which typically rely on
random mutagenesis (*33*). Simultaneous introduction of mutations in both β_{1-10} and β_{11}
regions that enhance their mutual interactions proved challenging through random
mutagenesis, because the statistical probability of obtaining cooperative mutations is
exceedingly low. Our successful implementation of structure prediction-guided optimization
suggests a broader applicability of this methodology across other split fluorescent protein
systems.

The discriminative capacity of the Caterpie system holds significant potential for
expansion through two primary avenues: the development of novel split fluorescent protein
variants and the integration of subcellular targeting sequences. Furthermore, compatibility
with the Landing Pad system (*34*), which enables precise single-copy genomic integration,
presents opportunities for multiplexed cellular labeling. These technological developments
collectively position Caterpie as a versatile platform for investigating complex biological
processes requiring simultaneous tracking of multiple distinct cell populations.

Materials and Methods

Plasmids

The following cDNAs were synthesized with optimized codons by GeneArt (Thermo Fisher Scientific, Waltham, MA): sfGFP$_{1-10}$ (*3*), CFP2$_{1-10}$ (*35*), mNeonGreen3A$_{1-10}$ (*11*), sfCherry3Csp$_{1-10}$, mRuby4$_{1-10}$ (*10*), GB1 (*18*), ΔSUMOstar (*19*). Additional split fluorescent protein variants were generated through site-directed mutagenesis of existing templates: Cerulean$_{1-10}$ (*10*) and YFP1$_{1-10}$ (*35*) from sfGFP$_{1-10}$, mNeonGreen3K$_{1-10}$ (*11*) from mNeonGreen3A$_{1-10}$, and sfCherry3C$_{1-10}$ (*12*) from sfCherry2$_{1-10}$ (Addgene: #82603). The Histone H2B coding sequence was subcloned from Addgene (plasmid # 11680 for Histone H2B; Cambridge, MA).

Vector Construction for Stable Cell Line Generation

A base Tol2 transposon vector (pT2A-IRES-iRFP670) was constructed by subcloning iRFP670 cDNA (*36*) with an internal ribosome entry site (IRES) (*37*) into the pT2A vector (*38*). CFP2$_{1-10}$ cDNA was subsequently PCR-amplified and inserted into this base vector using In-Fusion assembly (Clontech, Mountain View, CA) to generate pT2A_CFP2$_{1-10}$-IRES-iRFP670. Two distinct vector backbones, pT2A_IRES-iRFP670 and pT2ADW (containing IRES-puro cassette) (*39*), were used for multicistronic construct assembly. The following elements were PCR-amplified and assembled into both vectors using In-Fusion: sfCherry3Csp$_{1-10}$, mNeonGreen3Asp$_{1-10}$ with self-cleaving P2A peptide (*40*), and CFP2$_{1-10}$ with P2A peptide. Stable cell lines were generated through co-transfection of the constructed pT2A vectors with pCS-TP (obtained from Kawakami et al., 2004 (*38*)).

To generate pPBpuro_EF1a constructs containing CFP2$_{11}$(x1), CFP2$_{11}$(x2), CFP2$_{11}$(x4), CFP2$_{11}$(x8) -mCherry-NLS, cDNAs encoding Histone H2B, self-cleaving peptide P2A sequences, mCherry, and the nuclear localization signal (NLS) of the SV40 large T antigen (PKKKRKV) (*41*) were PCR-amplified and assembled using In-Fusion into pPBpuro_EF1a vectors (a kind gift from K. Yusa), yielding pPBpuro_EF1a_Histone H2B-P2A-mCherry-NLS. Either synthesized CFP2$_{11}$ or CFP2$_{Full-length}$ cDNA (created by PCR amplification of CFP2$_{1-10}$ cDNA and CFP2$_{11}$) was then assembled into pPBpuro_EF1a_Histone H2B-P2A-mCherry-NLS, resulting in pPBpuro_EF1a_CFP2$_{Full-length}$/CFP2$_{11}$(x1)-mCherry-NLS. Restriction enzyme sites (BglII before FP$_{11}$, BamHI and NotI after FP$_{11}$) were introduced to facilitate the construction of tandem repeats. To generate CFP2$_{11}$(x2), the CFP2$_{11}$(x1) insert and vector were digested with BglII/NotI and BamHI/NotI, respectively, and then ligated. Subsequently, CFP2$_{11}$(x4) and CFP2$_{11}$(x8) were generated using the same approach. By substituting the CFP2$_{11}$ insert with sfCherry3Csp$_{11}$ or mNeonGreen3Asp$_{11}$ inserts, multiple tandem repeats of these fragments were constructed using the same strategy. Various modifications were made to customize the constructs for specific experimental requirements: the puromycin-resistance gene (puro) was replaced with the blasticidin S-resistance gene (bsr); Histone H2B was substituted with GB1, ΔSUMOstar; mCherry was replaced with

EBFP2 or mTagBFP2; and the NLS was substituted with the nuclear export signal (NES) of

the HIV-1 rev protein (LPPLERLTLD) (*42*).

Mutant variants of sfCherry3C$_{1-10}$ were generated using overlap extension PCR and

assembled into pPB-based vectors (*43*) containing IRES-bsr (blasticidin S-resistance gene)

using In-Fusion. Additional mutant variants of mNeonGreen3A$_{11}$ and sfCherry3C$_{11}$ were

synthesized and inserted into pPB-based vectors (*43*) containing IRES-puro (puromycin-

resistance gene) using Ligation High Ver. 2 (TOYOBO). Stable cell lines were established

through co-transfection of these constructs with pCMV-mPBase (obtained from the Wellcome

Trust Sanger Institute).

Cell culture and establishment of cell lines

HeLa cells and Lenti-X 293T cells were obtained from the Human Science Research

Resources Bank and Clontech, respectively. Both cell lines were maintained in DMEM (Wako

Pure Chemical Industries) supplemented with 10% fetal bovine serum (Sigma-Aldrich) and

1% penicillin/streptomycin (Nacalai Tesque). MDCK (ECACC 84121903) cells were

purchased from the European Collection of Authenticated Cell Cultures (ECACC) through the

RIKEN BioResource Center (no. RCB0995) and maintained in DMEM (Wako Pure Chemical

Industries) supplemented with 10% fetal bovine serum (Sigma-Aldrich), 1%

penicillin/streptomycin (Nacalai Tesque).

For transposon-mediated gene transfer, pT2A_CFP2$_{1-10}$-IRES-iRFP670, or pT2A_FP$_{1-10}$x3-IRES-iRFP670 was cotransfected with pCS-TP into HeLa cells by using 293fectin

(Thermo Fisher Scientific, Waltham, MA). These obtained HeLa cells were sorted using an

FACS Aria IIu cell sorter (Becton Dickinson, Franklin Lakes, NJ) based on iRFP670

fluorescence to achieve a high expression level of the CFP2$_{1-10}$ or FP$_{1-10}$x3. Single-cell

cloning of these sorted populations to yield HeLa$^{CFP2\ 1-10\_iRFP670}$ cells and HeLa$^{FP1-10x3\_iRFP670}$

cells.

pT2A_FP$_{1-10}$x3-IRES-iRFP670 was cotransfected with pCS-TP using either 293fectin

(Thermo Fisher Scientific, Waltham, MA) for HeLa cells or electroporation with an Amaxa

nucleofector (Lonza, Basel, Switzerland) for MDCK cells. The transfected cells were selected

with 5 μg mL^{-1} puromycin (no. P-8833; Sigma-Aldrich), followed by single-cell cloning to

yield HeLa$^{FP1-10x3}$ cells or MDCK$^{FP1-10x3}$ cell lines.

Fluorescence imaging

For evaluation of split fluorescent proteins, HeLa cells in 24-well plates were co-transfected

with 100 ng of FP$_{11}$ plasmid [FP$_{Full-length}$, FP$_{11}$(x1), or FP$_{11}$(x4)] and 400 ng of FP$_{1-10}$ plasmid

using 293fectin (Thermo Fisher Scientific, Waltham, MA). Cells were seeded onto collagen

type I-coated (Nitta Gelatin, Osaka, Japan) glass-based 24-well plates (AGC Inc., Tokyo,

Japan) and cultured for 48 hours. Prior to imaging, cells were equilibrated for at least 1 h in

DMEM/F12 (Thermo Fisher Scientific) supplemented with 10% fetal bovine serum (Sigma-

Aldrich), and penicillin/streptomycin (Nacalai Tesque).

For FP$_{11}$ labeling, pPBbsr_EF1a-ΔSUMOstar-CFP2$_{11}$(xa)-mNeonGreen3Asp$_{11}$(xb)-sfCherry3Csp$_{11}$(xc) was cotransfected with pCMV-mPBase into HeLa$^{FP1-10x3}$ cells by using 293fectin or MDCK$^{FP1-10x3}$ cells using electroporation. Transfected cells were selected with 10 µg mL^{-1} blasticidin S (Calbiochem, San Diego, CA) and seeded onto collagen type I-coated glass-based 24-well plates.

Cells were observed with a Leica TCS SP8 FALCON confocal microscope (Leica-Microsystems, Wetzlar, Germany) equipped with an HC PL APO 20x/0.75 dry CS2 objective, an HC PL APO 40x/1.30 OIL CS2 objective, Leica HyD SMD detectors, a white light laser of 80 MHz pulse frequency, a Diode 405 (VLK 0550 T01; LASOS, Jena, Germany), a 440 nm diode laser (PDL 800-D; PicoQuant, Berlin, Germany), and a stage top incubator (Tokai Hit, Fujinomiya, Japan) to maintain 37 °C and 5% CO2. The following excitation wavelengths and emission band paths were used for the imaging: for EBFP2 imaging, 405 nm excitation, 410–450 nm emission; Cerulean or CFP2 imaging, 440 nm excitation, 460–490 nm emission; for sfGFP imaging, 488 nm excitation, 500–550 nm emission; for YFP1 imaging, 514 nm excitation, 520–550 nm emission; for mNG3A, mNG3K, or mNG3Asp imaging, 506 nm excitation, 510–550 nm emission; for sfCherry3C or sfCherry3Csp imaging, 594 nm excitation, 600–645 nm emission; for mRuby4 imaging, 561 nm excitation, 580–645 nm emission; for iRFP670 imaging, 650 nm excitation, 660–760 nm emission. To eliminate the background signal, the time gate for fluorescence detection was set from 1.0 ns to 6.0 ns.

In the multiplex imaging, the following excitation wavelengths and emission band paths were used: for CFP2 imaging, 440 nm excitation, 450–500 nm emission; for mNG3Asp imaging, 506 nm excitation, 515–580 nm emission; for sfCherry3Csp imaging, 594 nm excitation, 605–645 nm emission. To eliminate the background signal, the time gate for mNG3Asp and sfCherry3Csp fluorescence detection was set from 0.2 ns to 6.0 ns.

Images were processed and analyzed with ImageJ (*44*). For quantification of fluorescent intensity, images were segmented using Cellpose algorithm (*21*). Fluorescent intensity calculation was performed using custom MATLAB (MathWorks) and Python scripts.

Fluorescence spectral analysis

Fluorescence spectra were measured using a Leica TCS SP8 FALCON inverted microscope. Images were acquired with an HC PL APO 20x/0.75 dry CS2 objective using excitation at either 470 nm for split mNG3A or 546 nm for split sfCherry3C.

Flow cytometry analysis

Cells were suspended in PBS containing 3% FBS and analyzed or sorted with a FACS Aria IIu cell sorter (Becton Dickinson, Franklin Lakes, NJ). The following laser and emission filter combinations were used for fluorescence detection: CFP2, a 407 nm laser and an ET470/24m filter (Chroma Technology Corp., Bellows Falls, VT); mNeonGreen3Asp, a 488 nm laser, and a DF530/30 filter; sfCherry3Csp, a 561 nm laser and a DF582/15 filter (Omega Optical); iRFP670, a 633 nm laser and a DF660/20 filter (Omega Optical). Cell debris and aggregates

549   were excluded by gating for size and granularity. Data analysis was performed using FlowJo

550   software (Tree Star, Ashland, OR).

Lentivirus infection

For lentivirus production, the EGF ligand or receptor expressing plasmid, psPAX2 (Addgene

no. 12260), and pCMV-VSV-G-RSV-Rev (*45*) were co-transfected into Lenti-X 293T cells

using polyethylenimine (no. 24765-1; Polyscience Inc.). The infected cells were selected with

media containing the following antibiotics, depending on the drug resistance genes carried by

the EGF ligand or receptor expressing plasmids: 200 μg ml^{-1} hygromycin (no. 31282-04-9;

Wako).

Cell growth

For quantifying cell growth, MDCK cells labeled by Caterpie were seeded on collagen-coated

24-well glass-bottom plates (AGC Inc., Tokyo, Japan) at a density of 5×10^4 cells/ml. After 1

hour of incubation, the medium was replaced with DMEM/F12 supplemented with 100

units/ml penicillin, 100 μg/ml streptomycin, and 10% FBS. Following cell seeding, cells were

observed using a confocal microscope (Leica SP8) every 24 hours for 3 days. For observation

conditions, please refer to the "Fluorescent imaging" section. Cell numbers were counted

using the Cellpose segmentation algorithm.

Computational design for enhanced β-barrel stability and split-strand complementation

The tertiary structures of split mNG3Asp and split sfCherry3Csp were modeled using

ColabFold (*13*) implementation of AlphaFold Multimer (*14*) or AlphaFold2 (*17*) and

visualized with PyMOL (http://www.pymol.org/). The highest-confidence model of the split

sfCherry3Csp was then adopted as the starting coordinate for downstream Rosetta-based design.

To enhance global β-barrel stability of split sfCherry3Csp, *in silico* site-saturation

mutagenesis (SSM) of all 224 positions in sfCherry3C$_{1-10}$ and sfCherry3C$_{11}$ was performed

with a Rosetta-based workflow adapted from Thieker et al (*16*). For each variant, side-chain

and backbone sampling were confined to a 10 Å sphere around the mutated residue, while the

remainder of the protein was held fixed under coordinate constraints (sd = 1 outside the design

sphere; sd = 2 within a surrounding "soft sphere"). Each mutant underwent three cycles of

Cartesian FastRelax (MonomerRelax2019), and the change in Rosetta energy (ΔE) was

calculated as the difference in total_score between each mutant and the wild-type model using

the ref2015 score function.

To reinforce β$_{1-10}$/β$_{11}$ interface, two design workflows, Point-Mutation (PM) and Mutation

Cluster (MC), were adapted from the workflow of Thieker et al. and modified for interface

optimization. (1) In the PM workflow, SSM of the β$_{11}$ fragment (residues 1–16) was carried out,

with the addition of an InterfaceAnalyzer to compute binding energy (dG_separated) and

packing statistics. Residues within 10 Å of each mutation site were sampled under the same

constraint scheme used for stability design, while non-neighbor residues were held fixed; each

point mutant was relaxed by Cartesian FastRelax prior to interface analysis, and final

dG_separated values were calculated using the unconstrained ref2015 Rosetta score function.

(2) In the MC workflow, the seed mutations from the PM step were proceeded into a FastDesign

protocol using the InterfaceDesign2019 relax script to generate clustered mutations. The

neighbor-selection radius was extended to 12 Å, and an inner design shell was defined by

residues both in direct atomic contact ($\leq$ 5–7 Å, via a CloseContact selector) and satisfying the

InterfaceByVector geometric filter. A ResidueTypeConstraintGenerator was applied to inner-

shell, biasing retention of wild-type identities. Critical chromophore-interacting residues were

protected by marking them non-designable. Multi-residue combinations sampled via

FastDesign, and each cluster variant was evaluated with InterfaceAnalyzer for dG_separated

and packing quality. The scores were obtained under the unconstrained ref2015 function.

Calculation of angular data from individual cells

Three-dimensional fluorescence intensity data were acquired from individual cells by

measuring CFP2, mNG3Asp, and sfCherry3Csp signals with background subtraction. These

measurements were represented in a 3D coordinate system, with CFP2 intensities plotted on

the x-axis, mNG3Asp on the y-axis, and sfCherry3Csp on the z-axis. For each cell, the Cartesian

coordinates (x, y, z) were transformed into spherical coordinates (r, $\text{Angle}_{\text{CvsG}}$, $\text{Angle}_{\text{RvsCG}}$) using

a custom-made Python script. The coordinate transformation was performed using standard

spherical coordinate conversion equations.

$r = (x^2 + y^2 + z^2)^{1/2}$

$\text{Angle}_{\text{CvsG}} = \tan^{-1}(y/x)$

$\text{Angle}_{\text{RvsCG}} = \cos^{-1}(z/r)$

Two-component Gaussian mixture model (GMM) for FP$_{11}$ tag screening

Fluorescence intensity measurements from individual cells across 96 distinct cell populations

were transformed into angular coordinates ($\text{Angle}_{\text{CvsG}}$ and $\text{Angle}_{\text{RvsCG}}$). To enhance clustering

robustness, outliers were removed by excluding the top and bottom 10% of data points that

exhibited the greatest deviation from their respective cluster centers in both angular

measurements. An unsupervised two-component GMM was implemented using scikit-learn

(version 0.24.2; Pedregosa et al., 2011 (*46*)) to perform pairwise population comparisons. For

random sampling, a predetermined number of cell populations were selected from a total of

96 populations using simple random sampling without replacement. All computational

analyses were performed using custom Python scripts.

Cell classification using Gaussian mixture model (GMM)

Prior to angular calculations, we subtracted background and normalized the fluorescence

intensities of CFP2, mNG3Asp, and sfCherry3Csp by dividing each value by the median

intensity across all 20 cell types. Using these normalized fluorescence values, we then

calculated $Angle_{CvsG}$ and $Angle_{RvsCG}$ measurements for subsequent analysis.

The classification model was trained using GMMs implemented in the scikit-learn

library. we first fitted a single-Gaussian distribution to the training subset of each class

individually; the resulting class-specific means and covariance matrices were then

concatenated to obtain the GMM. The training dataset comprised 80% of the total data

($\sim 2.9 \times 10^4$ cells in total, $n > 10^3$ cells per cell population), consisting of two-dimensional

features ($Angle_{CvsG}$ and $Angle_{RvsCG}$ measurements) from 20 distinct cell populations. The

trained GMMs were then applied to angular data from pooled cells, and the predicted cell

labels were visualized. For each class k, we calculated the probability of each input data point

belonging to the corresponding GMM model, representing the likelihood of class

membership:

$P(x|\theta_k) = N(x|\mu_k, \Sigma_k)$

where x is the input data point, θ_k represents the parameters of the k-th GMM model, and

$N(x|\mu_k, \Sigma_k)$ denotes the normal distribution with mean μ_k and covariance Σ_k. Subsequently, the

probabilities were normalized such that the sum of probabilities across all classes equals 1 for

each data point:

$P(k|x) = P(x|\theta_k) / \Sigma_j P(x|\theta_j)$

We visualized the membership probabilities of the predicted classes. All computational

analyses were performed using custom-made Python scripts.

Cell classification using K-means

We used $Angle_{CvsG}$ and $Angle_{RvsCG}$ as the feature space coordinates for clustering. K-

means clustering was performed on the resulting dataset using scikit-learn's KMeans

algorithm with 21 clusters (random_state = 0). As shown in Figure 6B, cells labeled with tag

(BGR)=(0,0,4) corresponding to ID #4 exhibited a wide distribution of values in the

$Angle_{CvsG}$-$Angle_{RvsCG}$ space, resulting in their identification as two distinct clusters that were

subsequently merged. This accounts for the selection of 21 clusters in our analysis. The

predicted cluster labels were then remapped to match the original cell population labels for

validation. Classification accuracy was assessed by comparing the remapped cluster

assignments with ground truth labels and visualized using a confusion matrix. The overall

classification accuracy was calculated as the percentage of correctly classified cells. All

calculations were performed using Python.

Clark-Evans Index

We analyzed the spatial distribution patterns of five cell types. Individual cells were first

segmented using Cellpose, with centroids calculated and classified into five cell types via the

Caterpie method. For spatial analysis, we extracted the coordinates of each cell type,
calculated nearest neighbor distances using a k-dimensional tree algorithm, and computed the
Clark-Evans index (R). This index compares observed mean distances to those expected under
complete spatial randomness, where R = 1 indicates random distribution, R < 1 indicates
clustering, and R > 1 indicates regularity. All calculations were performed using Python.

Cell count of the nearest neighbor

We analyzed the spatial relationships between different cell types using a nearest-neighbor
approach. Cells were classified into five distinct populations based on fluorescent labels. For
each cell of the target population (label 1), we calculated the distance to every cell of other
populations (labels 2-5) and identified the closest neighboring cell using a k-dimensional tree
algorithm. To assess the statistical significance of the observed patterns, we generated 100
randomized distributions for each dataset by shuffling cell positions while preserving the total
number of each cell type. To determine whether the observed spatial associations between cell
types differed significantly from random expectations, we calculated the ratio of the
frequency with which each non-target cell type appeared as the nearest neighbor to target cells
in the original distribution divided by the mean frequency observed across 100 randomized
distributions. All calculations were performed using Python.

Statistics

679 All statistical analyses were carried out using Python. No statistical analysis was used to
680 predetermine the sample size. An unpaired Welch's t-test was used to evaluate statistically
681 significant differences. Data are expressed as the mean ± s.d. *P*-values of less than 0.05 were
682 considered to be statistically significant in two-tailed tests and were classified into four
683 categories: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and n.s. (not significant, i.e., $p \geq 0.05$).

Figure Legend

Figure 1. Caterpie: A cell identification system based on split fluorescent protein arrays

(A) Schematic overview of the Caterpie approach, illustrating cell identification through modular arrays of split fluorescent proteins. **(B)** Evolutionary relationships among split fluorescent proteins utilized in this study, presented as a phylogenetic tree. **(C)** Molecular architecture of expression constructs: Histone H2B fusions containing either full-length FP (FP$_{Full-length}$), single FP$_{11}$ fragment, or tetrameric FP$_{11}$ arrays, shown alongside the complementary FP$_{1-10}$ fragment. **(D)** Quantitative analysis of relative fluorescence intensity in HeLa cells expressing FP$_{Full-length}$, FP$_{11}$(x1) or FP$_{11}$(x4), with FP$_{1-10}$ fragments (1:4 transfection ratio). Data presented as bee swarm plots with median values indicated by black lines (n > 130 cells per condition). Measurements obtained through confocal microscopy.

Figure 2. Signal amplification and cell classification using split CFP2 arrays

(A) Molecular strategy for generating tandem CFP2$_{11}$ repeats, including detailed amino acid and nucleotide sequences at the CFP2$_{11}$(x1) insert-vector junction. **(B)** Construct architecture and expression analysis: Upper panel shows schematics CFP2$_{1-10}$ and Histone H2B fusions containing either full-length CFP2 (CFP2$_{Full-length}$) or varying copy numbers of CFP2$_{11}$ [CFP2$_{11}$(x1), (x2), (x4), and (x8)]. Lower panel presents representative confocal micrographs of HeLa cells stably co-expressing CFP2$_{1-10}$ with either CFP2$_{Full-length}$ or CFP2$_{11}$ variants. Scale bar, 20 µm. **(C)** Quantitative analysis of normalized CFP2 fluorescence in HeLa cells expressing Histone H2B-tagged CFP2$_{11}$ [(x1), (x2), (x4), or (x8)]. Data presented as bee swarm plots with median values (red lines); >1300 cells analyzed across three independent experiments. **(D)** Workflow schematic for Gaussian mixture model (GMM) and implementation. **(E)** Classification performance matrix showing prediction accuracy against true labels (rows). Data represented as mean ± SD of prediction accuracy from three independent experiments, with color intensity indicating classification accuracy. **(F)** Representative ratio image of pooled cells expressing different CFP2$_{11}$ copy number variants [(x1), (x2), (x4), or (x8)]; scale bar: 100 µm. **(G)** Cell classification map derived from panel (F), showing GMM-based assignment of individual cells to specific copy number variants [(x1), (x2), (x4), or (x8)]. **(H)** Visualization of GMM classification confidence through membership probability mapping of cells shown in panel (G).

Figure 3. Computational design of split sfCherry3C

(A) In-silico site-saturation mutagenesis of sfCherry3C$_{11}$. Left: schematic of the β$_{11}$ residues targeted for saturation mutagenesis (black spheres). Right: heatmap of mean dG_separated values in Rosetta energy unit (REU) across three independent models for each amino acid substitution at each position. (B) Cluster-mutation design of sfCherry3C$_{11}$. Left: schematic showing a representative seed point mutation (yellow dot) and the shell of neighboring residues (yellow spheres) targeted for sequential cluster mutagenesis. Right: heatmap of the

mean dG_separated across ten cluster-mutated models derived from each seed mutation at
every β_{11} site; dot size indicates the median number of cluster-mutated residues across these
ten models. (C) Schematics of mutant variants of sfCherry3C$_{1-10}$ (sfCherry3C$_{1-10M}$) and
Histone H2B-tagged mutant variant of sfCherry3C$_{11}$ (sfCherry3C$_{11M}$). (D) List of mutant
variants of sfCherry3C$_{1-10M}$ and sfCherry3C$_{11M}$. Bee swarm plot showing normalized
sfCherry3C fluorescence intensity of HeLa cells stably co-expressing Histone H2B-tagged
sfCherry3C$_{11M}$(x1) with sfCherry3C$_{1-10M}$. Red lines represent the median. (E) Rosetta models
of split sfCherry3Csp. The overall β-barrel is shown as a cartoon colored by fragment (gray,
β_{1-10}; magenta, β_{11}). Each inset zooms on an engineered site, showing the parental
conformation on the left and the corresponding design on the right, with key residues
rendered as sticks representation. All models were visualized in PyMOL. (F) Emission spectra
of HeLa cells expressing split sfCherry3C or split sfCherry3Csp with 546 nm laser excitation.

Figure 4. Optimization of split mNG3A through structure-guided C-terminal
engineering

(A) Predicted structural model of native mNG3A11(x8) generated using ColabFold
(AlphaFold2) and visualized with PyMOL. Colors represent hydrophobicity according to the
Eisenberg hydrophobicity scale. (B) Predicted structural model of parental split mNG3A
generated using ColabFold (AlphaFold Multimer) and visualized with PyMOL. Color-coded
domains: mNG3A$_{1-10}$ (green) and mNG3A$_{11}$ (light green). (C) Comparative sequence analysis
of original mNG3A$_{11}$ and engineered mutant variants (mNG3A$_{11M}$). (D) Schematic
representation of expression constructs: mNG3A$_{1-10}$ and Histone H2B fusion with the
mNG3A$_{11M}$(x8) array. Quantitative analysis of normalized mNG3A fluorescence in HeLa
cells expressing Histone H2B-tagged mNG3A$_{11}$ [(x1), (x2), (x4), or (x8)]. Data presented as
bee swarm plots with median values (red lines). (E) Quantitative analysis of normalized
mNG3A fluorescence intensity in HeLa cells stably co-expressing Histone H2B-tagged
mNG3A$_{11M}$ (x8) with mNG3A$_{1-10}$. Median values indicated by red lines. Values normalized to
set the median value of mNG3A$_{11M1}$(x8) to 1. (F) Emission spectra of HeLa cells expressing
split mNG3A or split mNG3Asp with 470 nm laser excitation. (G) Comparative analysis of
relative fluorescence intensity of HeLa cells expressing FP$_{Full-length}$, FP$_{11}$(x1) or FP$_{11}$(x4), with
cognate FP$_{1-10}$ fragments (1:4 transfection ratio). Data presented as bee swarm plots with
median values (red lines); n > 130 cells per condition. Measurements obtained through
confocal microscopy.

Figure 5. Systematic selection and validation of FP$_{11}$ tags for Caterpie implementation
(A) Schematic representation of FP$_{11}$ tags: ΔSUMO-tagged FP$_{11}$ arrays comprising paired
repeats of CFP2$_{11}$, mNG3Asp$_{11}$, and sfCherry3Csp$_{11}$ [(x0), (x1), (x2), (x4), (x8) each].
Comprehensive catalog of the FP$_{11}$ tag library comprising 96 distinct combinations, detailing
copy numbers of CFP2$_{11}$ (labeled as "C"), mNG3Asp$_{11}$ (labeled as "G"), and sfCherry3Csp$_{11}$
(labeled as "R") fragments. The marks indicate the FP$_{11}$ tags selected during primary and

secondary screening processes, and those finally chosen. **(B)** Schematic representation of

expression constructs: $FP_{1-10}x3$. **(C)** Analysis scheme for classification of cell populations

expressing various FP_{11} tags. **(D)** Matrix analysis of pairwise classification accuracy across 96

cell populations expressing various FP_{11} tags, arranged by FP_{11} tag numbers shown in panel

(A). (n > 230 cells per population). **(E)** Analysis scheme for determining the highest

achievable minimum pairwise classification accuracy among randomly selected FP_{11}-tagged

subpopulations. **(F)** Graph showing the highest achievable minimum pairwise classification

accuracy among randomly selected FP_{11}-tagged subpopulations. For each subpopulation size,

minimum accuracy between all possible pairs was determined from 10^7 random selections,

with maximum values plotted. **(G)** Comparative performance analysis of 30 candidate

populations (detailed in fig. S4) using both confocal microscopy (blue) and flow cytometry

(orange). Analysis methodology follows that of panel (F). Data presented as mean ± SD from

three independent experiments.

Figure 6. High-fidelity discrimination of twenty distinct cell populations using Caterpie

(A) Representative multicolor fluorescence micrographs of 20 distinct cell populations

expressing unique FP_{11} tag combinations. Fluorescence channels: CFP2 (blue), mNG3Asp

(green), and sfCherry3Csp (red). Copy numbers of each FP_{11} variant ($CFP2_{11}$, $mNG3Asp_{11}$,

and $sfCherry3Csp_{11}$) are indicated in the upper left corner of each panel. Scale bars: 30 μm.

(B) Left: Scatter plot showing $Angle_{CvsG}$ versus $Angle_{RvsCG}$ distributions for 20 distinct cell

populations from panel (A). (n > 1000 cells per population) Center: Distribution of label

assignments in $Angle_{CvsG}$ and $Angle_{RvsCG}$ based on Gaussian mixture model (GMM)

classification. Right: Distribution of label membership probabilities in $Angle_{CvsG}$ and

$Angle_{RvsCG}$. **(C)** Classification performance matrix showing prediction accuracy against true

population identities (rows). Color intensity indicates classification accuracy. Overall average

accuracy, 97 %. **(D)** Large-field composite image of pooled cell populations from panel (A),

displaying CFP2 (blue), mNG3Asp (green), and sfCherry3Csp (red) fluorescence channels.

Scale bar: 500 μm. **(E)** Detailed analysis of region indicated by dashed box in panel (D). Left:

Higher magnification of selected region. Center: Population assignment map following

GMM-based classification into 20 distinct categories. Right: Visualization of GMM

classification reliability through membership probability mapping.

Figure 7. Interaction dynamics in multi-population cell cultures of MDCK cells

expression EGF-family ligands and their receptors

(A) Schematic diagram showing the association pairs between EGF ligands (HBEGF, EREG)

and EGF receptors (EGFR, ErbB2). **(B)** Representative pseudocolored image of five distinct

cell types cultured together and identified using the Caterpie method. Yellow: WT, Pink:

EREG, Cyan: HBEGF, Green: EGFR, Orange: ErbB4; Copy numbers of each FP11 variant

(CFP211, mNG3Asp11, and sfCherry3Csp11) are (1,2,4), (2,0,4), (4,0,1), (2,4,0), and (8,2,1),

respectively. Scale bar: 1 mm. **(C)** Individual display of the five cell types shown in panel E.

The upper left corner of each panel indicates the re-expressed EGF signaling molecules and
the copy numbers of FP_{11} tags. **(D)** Graph showing the Clark-Evans Index calculated for each
of the five cell types. Data from three independent experiments. Black lines represent the
mean values across the three experiments. **(E)** Normalized nearest neighbor cell analysis for
evaluating spatial relationships between cells. This figure shows the ratio of heterotypic
nearest neighbor counts observed in the actual data to those counted after randomly
reassigning the labels of five cell types in the same dataset. Data from three independent
experiments. Black lines represent the mean values across the three experiments.
805

Supplementary Figure S1. Comprehensive mutational scanning of split sfCherry3C
Heatmap displaying the ΔEnergy values for saturating mutations in both $sfCherry_{1-10}$ and
$sfCherry_{11}$.

Supplementary Figure S2. Optimization of signal amplification using truncated
SUMOstar fusion strategy
(A) Schematic representation of expression constructs: $CFP2_{Full-length}$-NES and $CFP2_{11}$
variants [(x1) or (x8)]-NES with or without GB1 or ΔSUMOstar (truncated SUMOstar) tags.
(B) Amino acid sequences of SUMOstar, with ΔSUMOstar region highlighted in bold. **(C)**
Bee swarm plots showing quantitative analysis of normalized CFP2 fluorescence in HeLa
cells co-expressing $CFP2_{1-10}$ with either $CFP2_{Full-length}$-NES or $CFP2_{11}$(x1)/$CFP2_{11}$(x8)-NES
variants, with or without GB1 or ΔSUMOstar tags (constructs shown in panel A). Median
values are indicated by red lines. **(D)** Construct architecture and expression analysis: Upper
panel shows schematics of ΔSUMOstar fusions containing varying copy numbers of $CFP2_{11}$
[$CFP2_{11}$(x1), (x2), (x4), and (x8)]. Lower panel presents representative confocal micrographs
of HeLa cells stably co-expressing $CFP2_{1-10}$ with either $CFP2_{Full-length}$-NES or ΔSUMOstar-
tagged $CFP2_{11}$ variant. Scale bar: 20 μm. **(E)** Comparative analysis of normalized CFP2
fluorescence in HeLa cells expressing ΔSUMOstar-tagged $CFP2_{11}$ variants [(x1), (x2), (x4),
or (x8)] shown in panel D. Data presented as bee swarm plots with median values indicated
by red lines. **(F)** Classification performance matrix showing predictive accuracy against true
population identities (rows), with color intensity indicating classification accuracy.

Supplementary Figure S3. Influence of FP_{11} fragment order on fluorescence signal
distribution
(A) Schematic representation of expression constructs: ΔSUMO-tagged FP_{11} arrays
comprising paired repeats of $CFP2_{11}$, $mNG3Asp_{11}$, and $sfCherry3Csp_{11}$ [(x2) each, total x6].
(B) Quantitative analysis presented as bee swarm plots showing normalized fluorescence
intensities (CFP2, mNG3Asp, and sfCherry3Csp) in HeLa cells expressing six different
permutations of FP_{11} repeat arrangements, with concurrent FP_{1-10}x3 expression. Data obtained
by confocal microscopy, with median values indicated by black lines.

Supplementary Figure S4. Comparative analysis of binary classification performance across 30 selected cell populations

(A) Catalog of 30 selected FP$_{11}$ tag variants (detailed in Fig. 5A), detailing identification numbers and copy numbers of CFP2$_{11}$, mNG3Asp$_{11}$, and sfCherry3Csp$_{11}$ fragments. **(B)** Matrix analysis of pairwise classification accuracies for 30 cell populations expressing distinct FP$_{11}$ tag combinations (listed in panel A), evaluated by both microscopy (upper panel) and flow cytometry (lower panel). Data represent results from three independent experiments.

Supplementary Figure S5. Binary classification performance analysis of final 20 cell population panel

(A) Catalog of optimized FP$_{11}$ tag set comprising 20 selected variants (detailed in Fig. 5A), detailing identification numbers and copy numbers of CFP2$_{11}$, mNG3Asp$_{11}$, and sfCherry3Csp$_{11}$ fragments. **(B)** Matrix analysis of pairwise classification accuracy for 20 cell populations expressing distinct FP$_{11}$ tag combinations (defined in panel A). Comparative evaluation performed using both microscopy (left panel) and flow cytometry (right panel). Data represent mean values from three independent experiments in Supplementary Figure S3.

Supplementary Figure S6. Aggregation character of sfCherry3Csp$_{11}$(x8)

Representative images of sfCherry3Csp fluorescence in HeLa cells expressing ΔSUMOstar-tagged sfCherry3Csp$_{11}$ variants [(x4), or (x8)], with concurrent FP$_{1-10}$x3 expression.

Supplementary Figure S7. Unsupervised k-means clustering of 20 cell populations

(A) Left: Scatter plot showing the distribution of Angle$_{CvsG}$ versus Angle$_{RvsCG}$ for 20 distinct cell populations (n > 1,000 cells per population). Right: Scatter plot showing the distribution of Angle$_{CvsG}$ versus Angle$_{RvsCG}$ for individual cells grouped into 20 distinct populations using unsupervised k-means clustering. **(B)** Classification performance matrix of unsupervised k-means classification in panel (A). Color intensity indicates classification accuracy. Overall average accuracy, 96 %.

Supplementary Figure S8. High-fidelity discrimination of twenty distinct cell populations in MDCK cells using Caterpie

(A) Representative multicolor fluorescence micrographs of 20 distinct cell populations expressing FP$_{11}$ tags. Fluorescence channels: CFP2 (blue), mNG3Asp (green), and sfCherry3Csp (red). Copy numbers of each FP$_{11}$ variant (CFP2$_{11}$, mNG3Asp$_{11}$, and sfCherry3Csp$_{11}$) are indicated in the top of each panel. Scale bars: 30 μm. **(B)** Classification performance matrix showing prediction accuracy against true population identities (rows). Color intensity indicates classification accuracy. Overall average accuracy, 96%. **(C)** Top: Large-field composite image of pooled cell populations from panel (A), displaying CFP2 (blue), mNG3Asp (green), and sfCherry3Csp (red) fluorescence channels. Scale bar: 500 μm. Middle: Population assignment map following GMM-based classification into 20 distinct categories. Bottom: Visualization of GMM classification confidence through membership

871   probability mapping.

872   **Supplementary Figure S9. Cell growth of Caterpie labeled MDCK cells**

873   Growth curves showing cell density of 20 different MDCK cell types fluorescently labeled with

874   the Caterpie method measured every 24 hours. The lower panels display individual growth

875   curves for each cell type.

References

1. J. Livet, T. A. Weissman, H. Kang, R. W. Draft, J. Lu, R. A. Bennis, J. R. Sanes, J. W. Lichtman, Transgenic strategies for combinatorial expression of fluorescent proteins in the nervous system. Nature 450, 56-62 (2007).

2. M. W. Veling, Y. Li, M. T. Veling, C. Litts, N. Michki, H. Liu, B. Ye, D. Cai, Identification of neuronal lineages in the Drosophila peripheral nervous system with a "digital" multi-spectral lineage tracing system. Cell Rep. 29, 3303-3312.e3 (2019).

3. S. Cabantous, T. C. Terwilliger, G. S. Waldo, Protein tagging and detection with engineered self-assembling fragments of green fluorescent protein. Nat. Biotechnol. 23, 102-107 (2005).

4. M. D. Leonetti, S. Sekine, D. Kamiyama, J. S. Weissman, B. Huang, A scalable strategy for high-throughput GFP tagging of endogenous human proteins. Proc. Natl. Acad. Sci. U.S.A. 113, E3501-E3508 (2016).

5. N. H. Cho, K. C. Cheveralls, A. D. Brunner, K. Kim, A. C. Michaelis, P. Raghavan, H. Kobayashi, L. Savy, J. Y. Li, H. Canaj, J. Y. S. Kim, E. M. Stewart, C. Gnann, F. McCarthy, J. P. Cabrera, R. M. Brunetti, B. B. Chhun, G. Dingle, M. Y. Hein, B. Huang, S. B. Mehta, J. S. Weissman, R. Gómez-Sjöberg, D. N. Itzhak, L. A. Royer, M. Mann, M. D. Leonetti, OpenCell: Endogenous tagging for the cartography of human cellular organization. Science 375, eabi6983 (2022).

6. D. Kamiyama, S. Sekine, B. Barsi-Rhyne, J. Hu, B. Chen, L. A. Gilbert, H. Ishikawa, M. D. Leonetti, W. F. Marshall, J. S. Weissman, B. Huang, Versatile protein tagging in cells with split fluorescent protein. Nat. Commun. 7, 11046 (2016).

7. S. He, A. Cuentas-Condori, D. M. Miller, Strategy for bright, tissue-specific GFP labeling of endogenous proteins in C. elegans. Genetics 212, 387-395 (2019).

8. J. Goudeau, C. S. Sharp, J. Paw, L. Savy, M. D. Leonetti, A. G. York, D. L. Updike, C. Kenyon, M. Ingaramo, Split-wrmScarlet and split-sfGFP: tools for faster, easier fluorescent labeling of endogenous proteins in Caenorhabditis elegans. Genetics 217, iyab014 (2021).

9. R. Kamiyama, K. Banzai, P. Liu, A. Marar, R. Tamura, F. Jiang, M. A. Fitch, J. Xie, D. Kamiyama, Cell-type-specific, multicolor labeling of endogenous proteins with split fluorescent protein tags in Drosophila. Proc. Natl. Acad. Sci. U.S.A. 118, e2024690118 (2021).

10. R. Tamura, F. Jiang, J. Xie, D. Kamiyama, Multiplexed labeling of cellular proteins with split fluorescent protein tags. Commun. Biol. 4, 257 (2021).

11. S. Zhou, S. Feng, D. Brown, B. Huang, Improved yellow-green split fluorescent proteins for protein labeling and signal amplification. PLoS One 15, e0242592 (2020).

12. S. Feng, A. Varshney, D. Coto Villa, C. Modavi, J. Kohler, F. Farah, S. Zhou, N. Ali, J. D. Müller, M. K. Van Hoven, B. Huang, Bright split red fluorescent proteins for the

visualization of endogenous proteins and synapses. Commun. Biol. 2, 344 (2019).

13. G. Kim, S. Lee, E. Levy Karin, H. Kim, Y. Moriwaki, S. Ovchinnikov, M. Steinegger,
M. Mirdita, Easy and accurate protein structure prediction using ColabFold. Nat. Protoc.
20, 620–642 (2025).

14. R. Evans, M. O'Neill, A. Pritzel, N. Antropova, A. Senior, T. Green, A. Žídek, R. Bates,
S. Blackwell, J. Yim, O. Ronneberger, S. Bodenstein, M. Zielinski, A. Bridgland, A.
Potapenko, A. Cowie, K. Tunyasuvunakool, R. Jain, E. Clancy, P. Kohli, J. Jumper, D.
Hassabis, Protein complex prediction with AlphaFold-Multimer. bioRxiv
2021.10.04.463034 [Preprint] (2022); https://doi.org/10.1101/2021.10.04.463034.

15. A. Leaver-Fay, M. Tyka, S. M. Lewis, O. F. Lange, J. Thompson, R. Jacak, K. Kaufman,
P. D. Renfrew, C. A. Smith, W. Sheffler, I. W. Davis, S. Cooper, A. Treuille, D. J.
Mandell, F. Richter, Y.-E. A. Ban, S. J. Fleishman, J. E. Corn, D. E. Kim, S. Lyskov, M.
Berrondo, S. Mentzer, Z. Popović, J. J. Havranek, J. Karanicolas, R. Das, J. Meiler, T.
Kortemme, J. J. Gray, B. Kuhlman, D. Baker, P. Bradley, ROSETTA3: An object-
oriented software suite for the simulation and design of macromolecules. *Methods*
Enzymol. **487**, 545–574 (2011).

16. D. F. Thieker, J. B. Maguire, S. T. Kudlacek, A. Leaver-Fay, S. Lyskov, B. Kuhlman,
Stabilizing proteins, simplified: A Rosetta-based webtool for predicting favorable
mutations. Protein Sci. 31, e4428 (2022).

17. J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K.
Tunyasuvunakool, R. Bates, A. Žídek, A. Potapenko, A. Bridgland, C. Meyer, S. A. A.
Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back,
S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T.
Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K. Kavukcuoglu, P.
Kohli, D. Hassabis, Highly accurate protein structure prediction with AlphaFold. Nature
596, 583-589 (2021).

18. P. Zhou, A. A. Lugovskoy, G. Wagner, A solubility-enhancement tag (SET) for NMR
studies of poorly behaving proteins. J. Biomol. NMR 20, 11-14 (2001).

19. L. Liu, J. Spurrier, T. R. Butt, J. E. Strickler, Enhanced protein expression in the
baculovirus/insect cell system using engineered SUMO fusions. Protein Expr. Purif. 62,
21-28 (2008).

20. N. Milech, B. A. Longville, P. T. Cunningham, M. N. Scobie, H. M. Bogdawa, S.
Winslow, M. Anastasas, T. Connor, F. Ong, S. R. Stone, M. Kerfoot, T. Heinrich, K. M.
Kroeger, Y. F. Tan, K. Hoffmann, W. R. Thomas, P. M. Watt, R. M. Hopkins, GFP-
complementation assay to detect functional CPP and protein delivery into living cells.
Sci. Rep. 5, 18329 (2015).

21. C. Stringer, T. Wang, M. Michaelos, M. Pachitariu, Cellpose: a generalist algorithm for
cellular segmentation. Nat. Methods 18, 100-106 (2021).

22. E. Deguchi, S. Lin, D. Hirayama, K. Matsuda, A. Tanave, K. Sumiyama, S. Tsukiji, T.

Otani, M. Furuse, A. Sorkin, M. Matsuda, K. Terai, Low-affinity ligands of the epidermal
growth factor receptor are long-range signal transmitters in collective cell migration of
epithelial cells. Cell Rep. 43, 114986 (2024).

23. H. J. Snippert, L. G. van der Flier, T. Sato, J. H. van Es, M. van den Born, C. Kroon-
Veenboer, N. Barker, A. M. Klein, J. van Rheenen, B. D. Simons, H. Clevers, Intestinal
crypt homeostasis results from neutral competition between symmetrically dividing Lgr5
stem cells. Cell 143, 134-144 (2010).

24. I. Tabansky, A. Lenarcic, R. W. Draft, K. Loulier, D. B. Keskin, J. Rosains, J. Rivera-
Feliciano, J. W. Lichtman, J. Livet, J. N. H. Stern, J. R. Sanes, K. Eggan, Developmental
bias in cleavage-stage mouse blastomeres. Curr. Biol. 23, 21-31 (2013).

25. C. Ghigo, I. Mondor, A. Jorquera, N. J. Nowak, S. Wienert, S. P. Zahner, B. E. Clausen,
H. Luche, B. Malissen, F. Klauschen, M. Bajénoff, Multicolor fate mapping of
Langerhans cell homeostasis. J. Exp. Med. 210, 1657-1664 (2013).

26. D. Cai, K. B. Cohen, T. Luo, J. W. Lichtman, J. R. Sanes, Improved tools for the
Brainbow toolbox. Nat. Methods 10, 540-547 (2013).

27. K. Loulier, R. Barry, P. Mahou, Y. Le Franc, W. Supatto, K. S. Matho, S. Ieng, S.
Fouquet, E. Dupin, R. Benosman, A. Chédotal, E. Beaurepaire, X. Morin, J. Livet,
Multiplex cell and lineage tracking with combinatorial labels. Neuron 81, 505-520
(2014).

28. K. Weber, U. Bartsch, C. Stocking, B. Fehse, A multicolor panel of novel lentiviral "gene
ontology" (LeGO) vectors for functional gene analysis. Mol. Ther. 16, 698-706 (2008).

29. J. García-Marqués, L. López-Mascaraque, Clonal identity determines astrocyte cortical
heterogeneity. Cereb. Cortex 23, 1463-1472 (2013).

30. F. García-Moreno, N. A. Vasistha, J. Begbie, Z. Molnár, CLoNe is a new method to
target single progenitors and study their progeny in mouse and chick. Development 141,
1589-1598 (2014).

31. R. Sakaguchi, M. N. Leiwe, T. Imai, Bright multicolor labeling of neuronal circuits with
fluorescent proteins and chemical tags. eLife 7, e40350 (2018).

32. T. Kumamoto, F. Maurinot, R. Barry-Martinet, C. Vaslin, S. Vandormael-Pournin, M.
Le, M. Lerat, D. Niculescu, M. Cohen-Tannoudji, A. Rebsam, K. Loulier, S. Nedelec, S.
Tozer, J. Livet, Direct readout of neural stem cell transgenesis with an integration-
coupled gene expression switch. Neuron 107, 617-630.e6 (2020).

33. S. Feng, S. Sekine, V. Pessino, H. Li, M. D. Leonetti, B. Huang, Improved split
fluorescent proteins for endogenous protein labeling. Nat. Commun. 8, 370 (2017).

34. K. A. Matreyek, J. J. Stephany, M. A. Chiasson, N. Hasle, D. M. Fowler, An improved
platform for functional assessment of large protein libraries in mammalian cells. Nucleic
Acids Res. 48, e1 (2020).

35. T. Köker, A. Fernandez, F. Pinaud, Characterization of split fluorescent protein variants
and quantitative analyses of their self-assembly process. Sci. Rep. 8, 5344 (2018).

993 36. D. M. Shcherbakova, V. V. Verkhusha, Near-infrared fluorescent proteins for multicolor
994   in vivo imaging. Nat. Methods 10, 751-754 (2013).

995 37. T. Akagi, K. Sasai, H. Hanafusa, Refractory nature of normal human diploid fibroblasts
996   with respect to oncogene-mediated transformation. Proc. Natl. Acad. Sci. U.S.A. 100,
997   13567-13572 (2003).

998 38. K. Kawakami, H. Takeda, N. Kawakami, M. Kobayashi, N. Matsuda, M. Mishina, A
999   transposon-mediated gene trap approach identifies developmentally regulated genes in
1000   zebrafish. Dev. Cell 7, 133-144 (2004).

1001 39. N. Komatsu, K. Terai, A. Imanishi, Y. Kamioka, K. Sumiyama, T. Jin, Y. Okada, T.
1002   Nagai, M. Matsuda, A platform of BRET-FRET hybrid biosensors for optogenetics,
1003   chemical screening, and in vivo imaging. Sci. Rep. 8, 8984 (2018).

1004 40. J. H. Kim, S. R. Lee, L. H. Li, H. J. Park, J. H. Park, K. Y. Lee, M. K. Kim, B. A. Shin,
1005   S. Y. Choi, High cleavage efficiency of a 2A peptide derived from porcine teschovirus-1
1006   in human cell lines, zebrafish and mice. PLoS One 6, e18556 (2011).

1007 41. D. Kalderon, B. L. Roberts, W. D. Richardson, A. E. Smith, A short amino acid sequence
1008   able to specify nuclear location. Cell 39, 499-509 (1984).

1009 42. U. Fischer, J. Huber, W. C. Boelens, L. W. Mattaj, R. Lührmann, The HIV-1 Rev
1010   activation domain is a nuclear export signal that accesses an export pathway used by
1011   specific cellular RNAs. Cell 82, 475-483 (1995).

1012 43. K. Yusa, R. Rad, J. Takeda, A. Bradley, Generation of transgene-free induced pluripotent
1013   mouse stem cells by the piggyBac transposon. Nat. Methods 6, 363-369 (2009).

1014 44. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image
1015   analysis. Nat. Methods 9, 671-675 (2012).

1016 45. Miyoshi H., Blomer U., Takahashi M., Gage F. H., Verma I. M., *J. Virol.* **72**, 8150
1017   (1998).

1018 46. D. K. Barupal, O. Fiehn, Generating the blood exposome database using a
1019   comprehensive text mining and database fusion approach. Environ. Health Perspect. 127,
1020   97008 (2019).

1021

Fig.1

A

FP₁₁ tag   CFP₁₋₁₀   RFP₁₋₁₀   GFP₁₋₁₀

CFP₁₁ repeats   RFP₁₁ repeats   GFP₁₁ repeats

↓ Self-complementation

1. Labeling

HeLa^Triple1-10 Cell

20 FP₁₁ tags

2. Observation and model training

20 cell populations

Model training with a Gaussian mixture model (GMM)

Angle_RvsCG

Angle_CvsG

3. Classification (20 cells, 97%)

Pooled cells

Angle_RvsCG

Angle_CvsG

Applying the trained GMM

B

sfGFP ──── Cerulean
      ──── CFP2
      ──── YFP1
mNG2 ──── mNG3A
      ──── mNG3K
sfCherry2 ──── sfCherry3C
mRuby4

C

FP_Full   Histone H2B FP_Full-length EBFP2-NLS
                                         P2A

FP₁₁(x1)   Histone H2B EBFP2-NLS
                FP₁₁

FP₁₁(x4)   Histone H2B EBFP2-NLS

FP₁₋₁₀   FP₁₋₁₀ IRES iRFP670

D

Relative fluorescence intensity

FP_Full-length + FP₁₋₁₀
FP₁₁(x1) + FP₁₋₁₀
FP₁₁(x4) + FP₁₋₁₀

Cerulean CFP2 sfGFP sYFP1 mNG3A mNG3K sfCherry3C mRuby4

Fig.2

A

▼*Bgl*II   ▼*Bam*HI   ▼*Not*I

CFP2₁₁(x1)
Histone H2B

Insert
(*Bgl*II/*Not*I)
Vector
(*Bam*HI/*Not*I)

(x2)   (x4)   (x8)

CFP2₁₁   Linker   CFP2₁₁

Amino acid sequences   S G G R   T G G S G G R   T G G S

DNA sequences   AGATCTGGTGGAAGG....ACAGGAGGATCTGGTGGAAGG....ACAGGAGGATCC....GCGGCCGC
                TCTAGACCACCTTCC....TGTCCTCCTAGACCACCTTCC....TGTCCTCCTAGGCGCCGGCGG

*Bgl*II   *Bam*HI   *Not*I

B

CFP2₁₋₁₀ IRES iRFP670

Histone H2B CFP2_Full-length   mCherry-NLS
                              P2A

Histone H2B   ▪▪▪   mCherry-NLS
CFP2₁₁(x1), (x2), (x4), (x8)

Full-length (x1) (x2) (x4) (x8)
CFP2
mCherry

C

Normalized CFP2 intensity

Full-length (x1) (x2) (x4) (x8)

D

GMM training

CFP2 → mCherry   ⟹   CFP2 → mCherry

True   Trained GMM   Predicted
(x8)   (x8) Correct
(x4)
(x2)
(x1)   (x1) Incorrect

E

Predicted repeat numbers

	(x1)	(x2)	(x4)	(x8)
(x1)	98.5 ±1.6	0.8 ±0.6	0.7 ±0.5	0.0 ±0.0
(x2)	1.5 ±1.6	97.4 ±0.3	0.0 ±0.0	0.0 ±0.0
(x4)	0.0 ±0.0	1.8 ±0.8	96.3 ±2.6	6.1 ±1.8
(x8)	0.0 ±0.0	0.0 ±0.0	3.0 ±2.2	93.9 ±1.8

True repeat numbers

H

Trained GMM

98 %   2 %   0 %   0 %

F

Ratio image

2.8

CFP2/mCherry

0.4

G

GMM label map

(x8)
(x4)
(x2)
(x1)

GMM label reliability map

Membership probabilities (%)

Fig.3

A Point mutations

B Cluster mutations

C sfCherry3C$_{1-10M}$ Histone H2B | | mTagBFP2-NLS
                                   sfCherry3C$_{11M}$

D

	Mutation		Normalized sfCherry intensity
	1-10	11	
split sfCherry3C	-	-	
M1	A193P	S15T	
M2	A107Y	-	
M3	K48V	Y1T	
M4	E207G	-	
M5	A78K	-	
M6	H205D	T2R	
M7	-	Y1E	
M8	K48E, T50K	Y1R	
M9	H18R	-	
M10	-	E5D	
M11	D53Q	Y1D	
M12	D208S	T2E	
M13	T42Q	-	
M14	-	R9D	
M15	H76Y	-	
M16	R150V	S15Q	
M17	K48E, T50K	I3R	
M18	-	A10M	
M19	K48R	Y1T, T2I, I3D	
M20	-	T2I	
M21	H205D	T2K	
M22	-	A12V	
split sfCherry3Csp	K48V, A107Y, A193P	Y1T, S15T	

E

	WT	Design
M1	A193 S15	A193P S15T
M2	A107	A107Y
M3	Y1 K48	Y1T K48V

F

— sfCherry3C$_{11}$(x1)
— — sfCherry3Csp$_{11}$(x1)

Fig.4

A mNG3A11(x8)

F5
W8
F12
M15
M16

High
Hydrophobicity
Low

B

M16
M15
D14 F12
T13

C

                    5   10   15
mNG3A11   TELNFKEWQKAFTDMM
M1   TELNFKEWQKAFT
M2   TELNFKEWQKAFTD
mNG3Asp11   TELNFKEWQKAFTD**D**
M4   TELNFKEWQKA**Y**T
M5   TELNFKEWQKA**Y**TD
M6   TELNFKEWQKA**Y**TD**D**

D

mNG3A1–10

H2B [   ] mTagBFP2-NLS
mNG3A11M1-6 (x8)

E

Normalized mNG intensity

mNG3A11 M1 M2 mNG3Asp11 M4 M5 M6

F

Relative intensity

— mNG3A11(x1)
— mNG3Asp11(x1)

500   540   580   620
Wavelength (nm)

G

Relative fluorescence intensity

CFP2   mNG3Asp   sfCherry3Csp

Full-length (x1) (x4)   Full-length (x1) (x4)   Full-length (x1) (x4)

+FP1-10

Fig5

A FP₁₁ tag library: ΔSUMOstar ... CFP2₁₁ sfCherry3Csp₁₁ mNG3Asp₁₁

Each FP₁₁ repeat number: (x0), (x1), (x2), (x4), (x8)

B FP₁₋₁₀x3_iRFP670: sfCherry3Csp₁₋₁₀ mNG3A₁₋₁₀ P2A CFP2₁₋₁₀ IRES iRFP670

C Choosing 2 populations → Two-component Gaussian Mixture Model

D

E Randomly selected cell populations → 10⁷ trials → Minimum pairwise accuracy

F 1st screening

G 2nd screening
   Microscopy
   Flow Cytometry

Fig.6

A

HeLa cells
ID: (The repeat number of CFP2₁₁, mNG3Asp₁₁, sfCherry3Csp₁₁)

B

C

D

E

Fig.7

A

HBEGF, EREG

Caterpi labeled MDCK cells
(1,2,4) (2,0,4) (4,0,1) (2,4,0) (8,2,1)

WT   EREG   HBEGF   EGFR   ErbB4

EGFR (ErbB1)   ErbB4

Over-expression

Co-culture for 8 day (3% FBS)

WT cell

Ligand over-expressing cell   Receptor over-expressing cell

B

WT   HBEGF   EGFR
     EREG   ErbB4

1 mm

C

WT (1,2,4)   HBEGF (4,0,1)

EREG (2,0,4)   EGFR (2,4,0)

ErbB4 (8,2,1)

Splitting each cell types

1 mm

D

Clark-Evans Index

n.s.

WT EREG HBEGF EGFR ErbB4

E

Normalized cell count of nearest neighbor

n.s.
**

EREG

WT EGFR ErbB4

n.s.
***

HBEGF

WT EGFR ErbB4

**
n.s.

ErbB1

WT EREG HBEGF

*
n.s.

ErbB4

WT EREG HBEGF

Sup2

B

ΔSUMOstar

(MGHHHHHHGSLQDSEVNQEAKPEVK
PEVKPE)**THINLKVSDGSSEIFFKIK
KTTPLRRLMEAFAKRQGKEMDSLTF
LYDGIEIQADQTPEDLDMEDNDIIE
AHREQ**(IGGRDDLPSL)

A

B

| | CFP2
Relative intensity | mNG3Asp
Relative intensity | sfCherry3Csp
Relative intensity |

Sup. 4

A

ID: (The repeat number of CFP2_{11}, mNG3Asp_{11}, sfCherry3Csp_{11})

1: (4,0,0)	6: (1,8,0)	11: (2,0,4)	16: (0,8,2)	21: (1,8,1)	26: (4,1,1)
2: (0,2,0)	7: (2,2,0)	12: (4,0,1)	17: (0,8,4)	22: (1,8,2)	27: (4,1,4)
3: (0,8,0)	8: (2,4,0)	13: (0,2,8)	18: (1,1,2)	23: (2,1,1)	28: (4,2,2)
4: (0,0,2)	9: (8,1,0)	14: (0,4,8)	19: (1,2,4)	24: (2,1,2)	29: (4,4,4)
5: (0,0,8)	10: (1,0,8)	15: (0,8,1)	20: (1,4,4)	25: (2,2,8)	30: (8,2,1)

B

Sup. 5

A

ID: (The repeat number of CFP2₁₁, mNG3Asp₁₁, sfCherry3Csp₁₁)

1: (4,0,0)	5: (2,2,0)	9: (4,0,1)	13: (1,2,4)	17: (2,2,8)
2: (0,2,0)	6: (2,4,0)	10: (0,2,8)	14: (1,4,4)	18: (4,1,1)
3: (0,0,8)	7: (8,1,0)	11: (0,4,8)	15: (1,8,2)	19: (4,4,4)
4: (1,8,0)	8: (2,0,4)	12: (0,8,2)	16: (2,1,2)	20: (8,2,1)

B

Microscopy

Flow Cytometry

Sup.6

sfCherry3Csp$_{11}$(x4) sfCherry3Csp$_{11}$(x8)

50 um

A

B

A

MDCK cells

ID: (CFP2$_{11}$, mNG3Asp$_{11}$, sfCherry3Csp$_{11}$)

30 µm

B

Predicted label

True label

Accuracy (%)

C

500 µm

Identified label

Membership probabilities (%)

Sup.9