Intensity Modulation of Trichromatic Split Fluorescent Proteins for Live Cell Mapping

Mamoru Ishii[1,2*], Tomoaki Kinjo[3], Yohei Kondo[4,5], Kenta Terai[6], Kazuhiro Aoki[1], Brian Kuhlman[3,7], Michiyuki Matsuda[1,8,9*]

[1]Graduate School of Biostudies, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan.

[2]Institute of Industrial Science, The University of Tokyo, Tokyo, Japan

[3]Department of Biochemistry and Biophysics, University of North Carolina School of Medicine, Chapel Hill, NC, USA

[4]Center for One Medicine Innovative Translational Research (COMIT), Nagoya University, Nagoya, Aichi, Japan

[5]Graduate School of Medicine, Nagoya University, Tsurumai-cho, Nagoya, Aichi 466-8550, Japan

[6]Graduate School of Medicine, Tokushima University, Shinkura-cho, Tokushima 770-8501, Japan

[7]Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[8]Affiliate Graduate School, Graduate School of Medicine, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan.

[9]Integrated Graduate School of Medicine, Engineering, and Agricultural Sciences, University of Yamanashi, Chuo-shi, Yamanashi 409-3898 Japan.

*Corresponding authors. Center for Living Systems Information Science, Graduate School of Biostudies, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan
Lead correspondence: matsuda.michiyuki.2c@kyoto-u.ac.jp (M.M.), mishii@iis.u-tokyo.ac.jp (M.I.)

Running title
Multiplexed cell labeling with split FPs

Keywords
Split fluorescent proteins, Multiplexed spectral labeling, Cell labeling, Fluorescence microscopy, Protein tagging

Abstract

Multiplexed live cell-labeling technologies are essential tools for investigating complex biological systems, yet existing approaches face limitations in achieving reliable discrimination of multiple cell populations. Here, we present color and intensity modulation with split fluorescent proteins for cell labeling (Caterpie), a live cell labeling system that enables high-fidelity identification of multiple cell populations through rational design of split fluorescent proteins. Drawing inspiration from human trichromatic vision, we engineered optimized split fluorescent protein variants—split CFP2, split mNG3A, and split sfCherry3C—and developed a comprehensive library of the 11th β-strand (FP_{11}) tags comprising variable numbers of tandem repeats. Structure-guided protein engineering enhanced the performance of split mNG3A and split sfCherry3C through targeted modifications of their C-terminal sequences and binding interfaces, respectively. By systematically varying the composition and arrangement of FP_{11} repeats, we selected 20 distinct FP_{11} tags that enable robust cell population discrimination with 97% accuracy using conventional fluorescence microscopy. In these FP_{11} tag-expressing cells, we expressed multiple EGF ligands and their receptors and cultured them in bulk to examine the effects of EGF signaling on chemotaxis within the cell population. The results showed that chemotaxis changes according to the intensity of EGFR signaling. The Caterpie system holds significant potential for applications requiring precise identification and tracking of multiple cell populations in complex biological systems.

Introduction

Multiplexed spectral labeling techniques represent essential tools across diverse biological disciplines, including neuroscience, developmental biology, and immunology. These techniques enable critical applications such as discrimination of adjacent cells, specific cell populations labeling, and lineage tracing. The Brainbow system exemplifies this approach, utilizing multiple gene cassettes designed to stochastically express one of three fluorescent proteins (FPs), thereby generating distinct spectral signatures (*1*). While this system has proven particularly valuable for morphological analyses including connectomics, it presents limitations for specific cell labeling and lineage tracing due to potential variations in the relative FP expression levels across cellular generations. The development of Bitbow addressed certain limitations of earlier systems by enabling the generation of up to 31 (2^5-1) distinct color variations through random toggling of five FP genes within a single gene cassette (*2*). However, this approach faces challenges due to biased gene selection partners, resulting in fewer achievable color discrimination than theoretically predicted. These limitations underscore the advantages of rationally designed color palettes. However, conventional approaches to expanding color diversity through serial concatenation of multiple FPs face practical constraints due to increasing gene sizes. Split fluorescent proteins emerge as an elegant solution to these technical challenges.

Split fluorescent proteins (split FPs) are generated through strategic dissection of β-barrel fluorescent proteins (FPs) into two components: the 11th β-strand (FP_{11}) and a complementary segment comprising the first ten β-strands (FP_{1-10}) (*3*). While neither fragment exhibits fluorescence independently, their co-expression facilitates spontaneous reassembly, enabling chromophore maturation and subsequent fluorescence emission. This complementation system has been successfully employed for protein labeling applications, wherein target proteins are tagged with FP_{11} in cells expressing FP_{1-10} (*4*, *5*). A particularly advantageous feature of this system emerges under conditions of abundant FP_{1-10} expression: enhanced fluorescence intensity can be achieved by increasing the number of FP_{11} tags conjugated to the protein of interest, while maintaining relatively modest gene size requirements (*6–9*).

Drawing inspiration from human color perception, which discriminates diverse spectral signatures through the integration of three primary color intensities: blue, green, and red, we developed a novel multiplexed cellular labeling approach. This system exploits engineered arrays of FP_{11} fragments derived from cyan, green, and red split fluorescent proteins, each designed to complement with their cognate FP_{1-10} fragments. This method, named color and intensity modulation with split fluorescent proteins for cell labeling (Caterpie), achieves robust discrimination of 20 distinct cell populations with 97% accuracy using conventional epi-fluorescence microscopy.

Results

Split fluorescent proteins for the trichromatic color palette

We aimed to establish a trichromatic color palette through the systematic engineering of tandem FP_{11} repeats derived from cyan, green, and red split fluorescent proteins, which undergo fluorescence complementation when co-expressed with their cognate FP_{1-10} fragments (Fig. 1A). For this, we sought to select split fluorescent proteins (split FPs) that achieve brightness comparable to their full-length FP ($FP_{Full-length}$) and exhibit fluorescence signal amplification through FP_{11} repeats. A critical consideration in designing this system is the requirement for sequence divergence among the trichromatic split FPs to prevent cross-complementation between fragments from different fluorescent protein pairs (*10*). To address this constraint, we evaluated split FPs derived from distinct evolutionary lineages: sfGFP, mNG2, sfCherry2, and mRuby4 (Fig. 1B). To quantitatively assess the relative brightness of $FP_{Full-length}$, $FP_{11}(x1)$, and $FP_{11}(x4)$ of reported split FPs, we developed a dual-reporter system. We engineered expression constructs encoding Histone H2B-tagged $FP_{Full-length}$, $FP_{11}(x1)$, or $FP_{11}(x4)$ fused to EBFP2-nls (nuclear localization signal) through a self-cleaving P2A site, enabling normalization of expression levels through nuclear EBFP2 fluorescence. These constructs were co-transfected into HeLa cells with a secondary plasmid encoding FP_{1-10} linked to iRFP670 via an internal ribosome entry site (IRES) at a DNA ratio of 1:4 (Fig. 1C).

After 48 hours, cells were observed under a confocal microscope, and split FP fluorescence intensities were quantified in iRFP670 positive cells (Fig. 1D). For each fluorescent protein variant, relative fluorescence intensity values were normalized to the full-length FP. In most cases, (x1) shows decreased fluorescence brightness compared to full-length, but sfGFP, mNG3A, and mNG3K were almost equivalent fluorescence brightness. Comparative analysis of (x1) versus (x4) constructs revealed that all sfGFP-derived split FPs exhibited more than two-fold enhancement in median brightness, with CFP2 demonstrating the most substantial increase of more than three-fold. Consistent with previous findings (*11*), split mNG3A demonstrated superior amplification from (x1) to (x4) compared to split mNG3K. Split $sfCherry3C_{11}(x1)$ exhibited diminished fluorescence intensity relative to $sfCherry3C_{Full-length}$, attributable to reduced association efficiency, corroborating earlier observations (*12*). The truncated $mRuby4_{11}$ showed markedly decreased fluorescence intensity, highlighting the critical role of its C-terminal unstructured polypeptide chain (*10*). These preliminary findings indicated that while the combination of CFP2, mNG3A, and sfCherry3C provides a promising foundation for the trichromatic system, although further engineering optimization of mNG3A and sfCherry3C is necessary to achieve optimal performance across all spectral channels.

Fluorescence intensity-based cell classification using tandem CFP2 β-strand repeats

Before the engineering of mNG3A and sfCherry3C, we established a platform for cell

identification through fluorescence intensity modulation using CFP2$_{11}$ repeats. To this end, we

constructed plasmids encoding repeated sequences of CFP2$_{11}$ (Fig. 2A). Initially, we

engineered a plasmid encoding a Histone H2B-CFP2$_{11}$(x1) fusion construct, in which the

CFP2$_{11}$(x1) sequence was flanked by a BglII restriction site at its 5' terminus and BamHI and

NotI restriction sites at its 3' terminus. Through sequential ligation of BglII/NotI-digested

fragments (insert) with BamHI/NotI-digested fragments (vector), we successfully amplified

CFP2$_{11}$ to create (x2), (x4), and (x8) variants. Based on previous research (6), we

incorporated GGSGG linker sequences between CFP2$_{11}$ fragments. At the BglII-BamHI

junction, the six nucleotides encode glycine and serine residues, thereby serving as an integral

part of the linker sequence. The CFP2$_{11}$ repeats [(x1), (x2), (x4), or (x8)] fused with Histone

H2B were introduced into HeLa cells stably expressing CFP2$_{1-10}$ (HeLa$^{CFP2\ 1-10}$) (Fig. 2B). As

a control, we expressed Histone H2B-tagged CFP2$_{Full-length}$ in the same HeLa$^{CFP2\ 1-10}$ cell line.

When normalized to nuclear mCherry fluorescence, the CFP2 fluorescence intensity increased

across consecutive constructs with a factor of approximately 1.7-fold, which was slightly

lower than the theoretically expected 2-fold increment (Fig. 2C).

To identify cells expressing different numbers of CFP2$_{11}$ repeats, we implemented a

Gaussian mixture model (GMM) for cell classification (Fig. 2D). The fluorescence intensities

of mCherry and CFP2 were transformed into polar coordinates, where the Angle$_{mCherry\ vs\ CFP2}$

(ranging from 0° to 90°) represented the CFP2/mCherry intensity ratio. The angular data were

divided into training and test datasets. We trained a GMM on the training dataset to establish

classification parameters, determining the decision boundary of the Angle$_{mCherry\ vs\ CFP2}$. The

model performance was validated using the test dataset, and the predicted repeat numbers

were compared with the actual repeat numbers. The model achieved an overall accuracy of

96% (Fig. 2E). Among all constructs, CFP2$_{11}$(x8) exhibited the lowest accuracy (93.9%), with

6.1% of (x8)-expressing cells being misclassified as (x4).

To validate the system's discriminative capacity in a heterogeneous context, we analyzed

mixed populations of cells expressing four different CFP2$_{11}$ repeat variants using fluorescence

microscopy and generated CFP2/mCherry ratio images (Fig. 2F). The trained GMM

demonstrated robust performance in classifying individual cells into (x1), (x2), (x4), or (x8)

populations with 96% accuracy, which was validated through visual inspection (Fig. 2G).

Furthermore, the GMM provided membership probabilities for each cell, enabling

quantitative assessment of classification reliability (Fig. 2H). These results established that

four-level intensity modulation of a single fluorescent protein is sufficient for reliable

population discrimination.

Computational design of split sfCherry3C for enhanced β-barrel stability and

split-strand complementation

To optimize split sfCherry3C toward fluorescence intensities close to those of the

full-length protein, we modelled the sfCherry3C$_{1-10}$/sfCherry3C$_{11}$ complex with ColabFold

(*13*) (based on AlphaFold-Multimer) (*14*), and used the structure model for subsequent design

with Rosetta modeling suite (*15*). Because both global β-barrel stability and β_{1-10}/β_{11} interface

affinity are crucial for fluorescence of split FPs, we pursued two design strategies in parallel

(*16*). First, to enhance global β-barrel stability, we employed a Rosetta script adapted from the

previously described "Point-Mutation" (PM) workflow (*16*) and performed *in-silico*

site-saturation mutagenesis (SSM) across all 224 residues, ranking each variant by the change

in calculated energy (ΔE) calculated from the Rosetta *total_score* term (fig. S1). Next, to

refine the β_{1-10}/β_{11} interface, we repurposed the PM script for interface design; with the

revised script, we performed SSM at every β_{11} position and calculated the binding energy of

each single mutant from *dG_separated* score obtained with InterfaceAnalyzer protocol (Fig.

3A). Each single-mutant model was subsequently processed by the "Mutation Cluster (MC)"

(*16*) workflow modified for interface design, in which FastDesign protocol with

InterfaceDesign relax script stochastically samples sequence and rotamer space at residues

within 5–7 Å of the "seed mutation" through Monte-Carlo design/minimization cycles,

yielding multi-residue mutants that are re-evaluated with InterfaceAnalyzer (Fig. 3B). After

filtering by the scores such as *total_score*, *dG_separated*, and mutation number, we conducted

visual inspection to retain the variants that reinforce interface contacts while excluding ones

that disrupted the chromophore-forming triad M67–Y68–G69 or introduced additional

aromatic residues into β_{11} which could promote aggregation of multimeric β_{11} repeat.

Following these criteria, we selected twenty-two candidate designs for subsequent in-cell

validation.

Upon stable co-expression of β_{1-10} and β_{11} in HeLa cells, three variants (M1, M2, M3)

outperformed the parental split sfCherry3C in fluorescent intensity (Fig. 3C, D). We then

combined the mutations from the three top performing variants to generate an optimized

variant, named split sfCherry3Csp, which achieved a 2.5-fold enhancement in fluorescent

intensity compared to the parental split sfCherry3C. The M1 and M3 mutations locate at the

β_{1-10}/β_{11} interface, where the models suggest the improved association of β_{1-10} and β_{11} through

the formation of reinforced side-chain contacts. The M2 mutation lies on the opposite face of

the interface, implying an effect on β-barrel stability (Fig. 3E). The fluorescence spectra of

split sfCherry3Csp were indistinguishable to those of the parental split sfCherry3C, indicating

that the incorporated mutations enhanced the β-barrel stability and β_{1-10}/β_{11} association

without affecting the chromophore maturation (Fig. 3F).

Enhancement of complementation efficiency through structure-guided engineering of

split mNG3A

As shown in Fig. 1D, split mNG3A demonstrated only modest signal amplification, with a

1.3-fold increase in fluorescence intensity from (x1) to (x4) variants. To gain insight into the

structural factors underlying this limitation, we modelled the octameric mNG3A$_{11}$ peptide

using ColabFold (*13*) (based on AlphaFold2) (*17*). The predicted structures suggested that

mNG3A$_{11}$ repeat adopts a loosely helical conformation, with hydrophobic residues (F5, W8,
F12, M15, M16) converging to form a continuous hydrophobic interaction, despite only
moderate model confidence. These features implied that multimeric mNG3A$_{11}$ may
self-aggregate via its hydrophobic surfaces (Fig. 4A) (*17*). Additionally, inspection of the
parental mNG crystal structure (PDB: 5LTP) and the structure model of split mNG3A
suggested that the C-terminal residues D14–M15–M16 contribute minimally to the β-barrel
fold (Fig. 4B). We therefore reasoned that removing these C-terminal residues could mitigate
the aggregation propensity without perturbing β-barrel and chromophore maturation. Guided
by these insights, we implemented three C-terminal modification approaches: (1) deletion of
the C-terminal D14-M15-M16, (2) addition of one or two aspartic acid residue to preserve
charge that could electrostatically repel neighboring fragments, and (3) substitution of F12
with tyrosine to introduce polarity (Fig. 4C). To evaluate these modifications, we expressed 8
repeats of the mNG3A$_{11}$ fragments fused to Histone H2B in HeLa cells expressing
mNG3A$_{1-10}$ (Fig. 4D). Nuclear mNG3A fluorescence was quantified through microscopy,
with expression levels normalized to nuclear mTagBFP2 fluorescence intensity (Fig. 4E).
Among the modifications tested, we chose the mNG3Asp$_{11}$ variant for further analysis
because of its highest fluorescence intensity, which was 9.1-fold brighter than the parental
mNG3A$_{11}$. The fluorescence spectra obtained using the mNG3Asp$_{11}$ fragment were identical
to those observed with the mNG3A$_{11}$ fragment (Fig. 4F).
Performance evaluation of split mNG3Asp and split sfCherry3Csp was conducted in
parallel with split CFP2, following our previously established experimental framework (Fig.
4G). Quantitative analysis revealed that mNG3Asp$_{11}$(x4) achieved a 2.3-fold enhancement in
fluorescence intensity compared to its monomeric counterpart. The engineered sfCherry3Csp
demonstrated robust performance, with its monomeric variant exhibiting 80% of the
fluorescence intensity of the full-length protein, while the tetrameric construct showed a
3-fold signal amplification relative to the monomer. These significant improvements in signal
amplification efficiency indicate that both engineered variants, split mNG3Asp and split
sfCherry3Csp, now achieve performance metrics comparable to split CFP2, providing the
basis for trichromatic imaging.

Optimization of the fusion partner for the FP$_{11}$ tag
Prior to establishing the FP$_{11}$ tag library, we aimed to reduce construct size by identifying
alternatives to the relatively large Histone H2B fusion tag. We evaluated multiple fusion
protein configurations, comparing Histone H2B (126 amino acids) with two smaller
alternatives: GB1 (56 amino acids) (*18*) and ΔSUMOstar (74 amino acids), a truncated
version of SUMOstar (*19*), as well as constructs lacking fusion tags (fig. S2A). ΔSUMOstar
was engineered by removing the unstructured polypeptide regions from both termini of
SUMOstar; the resulting sequence is detailed in Figure S2B. In agreement with previous
studies (*20*), constructs lacking fusion proteins demonstrated markedly reduced fluorescence

intensity, an effect that persisted even in the octameric (x8) variant (fig. S2C). Comparative

analysis of fusion partners revealed that ΔSUMOstar-tagged constructs achieved substantially

higher fluorescence intensity than GB1-tagged variants in octameric configurations,

demonstrating the importance of fusion partner selection for optimal signal amplification. The

ΔSUMOstar-tagged constructs exhibited consistent signal enhancement, with fluorescence

intensity increasing 1.6-fold for each doubling of $CFP2_{11}$ repeats, ultimately enabling 93%

discrimination accuracy between variants (fig. S2, D to F).

To optimize the modular arrangement of the three FP_{11} variants ($CFP2_{11}$, $mNG3Asp_{11}$,

and $sfCherry3Csp_{11}$), we conducted a systematic analysis of possible configurations. In initial

experiments, we generated ΔSUMOstar fusion constructs containing six total repeats, wherein

each FP_{11} fragment was represented in duplicate (fig. S3A). The complete set of six possible

permutations was generated and stably expressed in HeLa cells constitutively expressing all

three FP_{1-10} fragments. Quantitative analysis of relative fluorescence intensities was

performed using EBFP2 fluorescence for normalization of expression levels (fig. S3B).

Among all configurations evaluated, the $CFP2_{11}(x2)$-$sfCherry3Csp_{11}(x2)$-$mNG3Asp_{11}(x2)$

arrangement demonstrated marginally superior fluorescence intensity across all spectral

channels, leading to its selection for subsequent studies.

Selection of 20 optimal FP_{11} tags

Based on these preliminary findings, we established a comprehensive library of fusion

constructs combining ΔSUMOstar with $CFP2_{11}$, $sfCherry3Csp_{11}$, and $mNG3Asp_{11}$ fragments

in this order (Fig. 5A). Each fragment was represented in varying copy numbers [(x0), (x1),

(x2), (x4), or (x8)] for each. In the first screening, we limited the FP_{11} tag combinations to 96

variants, with the total number of repeats not exceeding (x12). The integrity of all FP_{11} repeat

sequences in the 96 plasmid constructs was verified by Sanger sequencing, thereby

establishing a comprehensive FP_{11} tag library.

To generate a stable cellular platform for evaluating the FP_{11} tag library, we engineered an

expression vector encoding all three FP_{1-10} fragments (CFP_{1-10}, $sfCherry3Csp_{1-10}$, and

$mNG3A_{1-10}$) separated by self-cleaving P2A peptides, followed by an IRES and iRFP670 (Fig.

5B). Through fluorescence-activated cell sorting (FACS) based on iRFP670 expression levels,

we established a HeLa cell clone ($HeLa^{FP1-10x3\_iRFP670}$) expressing high levels of three FP_{1-10}

fragments.

The complete library of 96 FP_{11} tags was stably expressed in $HeLa^{FP1-10x3\_iRFP670}$ cells,

with resultant populations analyzed by confocal microscopy. To quantitatively assess the

fluorescence distribution patterns, we implemented spherical coordinate transformation to

calculate two angular parameters: $Angle_{CvsG}$ and $Angle_{RvsCG}$, each ranging from 0° to 90° (Fig.

5C, left). To evaluate the discriminative power of these angular parameters, we performed

pairwise population analysis using selected cell populations from the library. Cell

classification was achieved using an unsupervised two-component Gaussian mixture model

281 without prior training data (Fig. 5C, right). Classification accuracy was determined by

282 comparing model-predicted clusters against known population identities. Superior

283 classification accuracy correlated with greater separation of populations in the Angle_{CvsG} −

284 Angle_{RvsCG} parameter space. This analytical framework was systematically applied to all

285 possible pairwise combinations within the 96 FP_{11}-tagged cell library (Fig. 5D).

286  To determine the number of distinguishable cell populations, we implemented an iterative

287 sampling approach due to the vast number of possible patterns (approximately 2×10^{20}) when

288 selecting 20 FP_{11}-tagged cells from a pool of 96 (Fig. 5E). Random subsets of predetermined

289 sizes were drawn from the 96 FP_{11}-tagged cell populations, with pairwise classification

290 accuracies extracted from the comprehensive analysis presented in Fig. 5D. For each

291 FP_{11}-tagged subset, we identified the minimum classification accuracy, reflecting the subset's

292 ability to distinguish the most similar cell population pairs. This sampling process was

293 repeated 10^7 times, and the maximum achievable minimum accuracy was plotted as a function

294 of subpopulation size (Fig. 5F). The analysis revealed that a subset of 20 cell populations

295 maintained robust discrimination with a minimum pairwise classification accuracy of 90%.

296 However, expanding to 30 populations resulted in a substantial decrease in minimum

297 accuracy to 60%. Based on these quantitative insights, we selected an initial panel of 30

298 candidate FP_{11} tags (Fig. 5A, 1st) for subsequent refinement to establish the final optimized set

299 of 20 tags (Fig. 5A 2nd) that would ensure maximal discriminative power.

300  We conducted comprehensive evaluation of the 30 candidate populations using both

301 confocal microscopy and flow cytometry, with three independent experimental replicates (fig.

302 S4). The relationship between population subset size and maximum achievable minimum

303 pairwise classification accuracy was reassessed using these orthogonal detection methods (Fig.

304 5G). While fluorescence microscopy demonstrated marginally superior classification accuracy,

305 flow cytometry maintained robust discrimination capabilities, achieving >85% accuracy

306 across a set of top 20 FP_{11} tags. The final optimized set of 20 FP_{11} tags and their

307 comprehensive pairwise discrimination analysis is presented in Figure S5. Here, to optimize

308 the fluorescence signal intensity of single-color reference populations, we refined the copy

309 number composition (Fig. 5A Final): $CFP2_{11}(x8)$ and $mNG3Asp_{11}(x8)$ were implemented at

310 maximum repeat numbers, while $sfCherry3Csp_{11}$ was limited to (x4) to circumvent

311 aggregation phenomena observed at higher copy numbers (fig. S6).

High-fidelity discrimination of twenty cell populations using optimized FP_{1-10} expression

platform

314 The FP_{1-10} expression system was also refined by replacing the iRFP670 reporter with a

315 puromycin resistance cassette. Through stringent puromycin selection, we established a HeLa

316 clone cell (HeLa$^{FP1-10x3}$) expressing elevated levels of $CFP2_{1-10}$, $mNG3Asp_{1-10}$, and

317 $sfCherry3Csp_{1-10}$. The optimized set of 20 FP_{11} tags was stably expressed in HeLa$^{FP1-10x3}$ cells,

318 with individual cell populations analyzed by confocal microscopy. Fluorescence was

visualized with distinct spectral channels: CFP2 (blue), mNG3Asp (green), and sfCherry3Csp (red). Merged images revealed unique signatures based on both color composition and intensity distribution (Fig. 6A).

Cell segmentation was performed on maximum intensity projections of composite fluorescence signals using Cellpose (*21*). Mean fluorescence intensities were quantified across all three channels for each segmented cell. Following the methodology established in Figure 5C, we implemented spherical coordinate transformation to calculate $Angle_{CvsG}$ and $Angle_{RvsCG}$, generating a two-dimensional angular representation of relative fluorescence intensities (Fig. 6B, left). These parameters were used to train a Gaussian mixture model (GMM), which established both label assignment distributions and label membership probability distributions in $Angle_{CvsG}$ and $Angle_{RvsCG}$ (Fig. 6B, center and right). Validation of the trained GMM on independent test data demonstrated exceptional classification performance, achieving 97% accuracy in discriminating all 20 cell populations (Fig. 6C). Using unsupervised k-means clustering on the 20 components achieved 96% accuracy in distinguishing between the 20 cell populations, even without requiring pre-training data for individual fluorescence profiles (fig. S7).

To evaluate system performance under practical conditions, we used pooled samples containing all 20 cell populations and acquired tiled images (Fig. 6D). Application of the trained GMM enabled robust classification of individual cells, with 78% of cells being classified with more than 95% probability (Fig. 6E).

Interaction dynamics in multi-population cell cultures of MDCK cells expressing EGF-family ligands and their receptors

Finally, as an application of this system—named color and intensity modulation with split fluorescent proteins for cell labeling (Caterpie)—we studied the contribution of heterotypic cell interactions mediated by EGF-family ligands and their cognate receptors. For this, we used Madin-Darby Canine Kidney (MDCK) cells, which are widely used to study collective cell migration and cell competition regulated by the EGF signaling pathway. MDCK cells were labeled with the aforementioned FP_{1-10} and 20 FP_{11} tags with 96% accuracy (fig. S8A, B). Mixed cells were classified by applying a trained Gaussian mixture model (fig. S8C), 75% of cells being classified with more than 95% probability. The cell densities of twenty different labeled cell types at 72 hours post-labeling ranged within a 1.4-fold difference between the minimum and maximum values (fig. S9).

Among the four EGF-family ligands expressed in MDCK cells, we focused on HBEGF and EREG. HBEGF, a high-affinity ligand, binds to heparan sulfate proteoglycans to provide strong signals restricted to short distances, promoting localized cell migration. In contrast, EREG, a low-affinity ligand, diffuses quickly and remotely, propagating signals approximately four times faster than high-affinity ligands and more efficiently to distant cells (*22*). Both HBEGF and EREG bind to EGFR (ErbB1) and ErbB4. Using lentiviral

transduction, we introduced EREG, HBEGF, EGFR (ErbB1), or ErbB4 in fluorescently
labeled MDCK cells (Fig. 7A). To investigate how EGF ligands and receptors with different
affinities interact between heterotypic cells and generate spontaneous patterns, these four cell
types, along with wild-type cells, were co-cultured for 8 days under 3% low-serum conditions.
The resulting cell population was analyzed by fluorescence microscopy to identify each cell
type (Fig. 7B).
Before analyzing the positional relationships between heterotypic cells, we first examined
the distribution patterns of homotypic cells, as cell types with limited motility and dispersal
capability affect intercellular interactions with heterotypic cells. To examine homotypic cell
clustering, each cell type is individually displayed (Fig. 7C). Dispersion of each cell type was
analyzed by the Clark-Evans index; values less than 1 indicate cellular clustering. We did not
find significant homotypic cell clustering in any cell types (Fig. 7D). Then, we investigated
the frequency of adjacent heterotypic cells. When examining receptor-expressing cells
adjacent to EREG ligand or HBEGF-expressing cells (Fig. 7E), we found EGFR cells
displayed significantly higher adjacency to EREG cells than wild-type cells. Conversely,
when examining ligand-expressing cells adjacent to EGFR receptor or ErbB4-expressing cells,
HBEGF cells exhibited higher adjacency to EGFR cells and to a lesser extent to ErbB4 in
comparison to wild-type cells. These results suggest that EREG and EGF function as
chemoattractants, with EREG weakly attracting distant cells, while HBEGF strongly attracts
nearby cells.
In conclusion, the Caterpie method provides a versatile platform for analyzing cell
populations that may exhibit homotypic and/or heterotypic clustering.
379

Discussion

We have developed color and intensity modulation with split fluorescent proteins for cell labeling (Caterpie), a novel cell identification platform that achieves robust identification of 20 distinct cell populations with 96-97% accuracy. This system leverages fluorescence intensity modulation through engineered arrays of split fluorescent protein (split FP) fragments. The technological foundation of Caterpie comprises three complementary split FPs: split CFP2 and two newly engineered variants - split mNG3Asp and split sfCherry3Csp. To implement this system, we established a comprehensive toolkit consisting of 20 distinct FP_{11} tags, each constructed by fusing truncated SUMOstar protein to the 11th β-strand of the three split fluorescent proteins in varying tandem repeat configurations. These engineered FP_{11} tags demonstrate highly specific labeling of FP_{1-10}-expressing cells, enabling population discrimination with 96-97% accuracy through Gaussian mixture model (GMM) analysis of the resulting multidimensional fluorescence signatures.

Current multicolor fluorescent labeling approaches can be broadly categorized into two distinct methodological frameworks. The first relies on stochastic recombination events mediated by site-specific recombinase systems such as Cre/loxP and Flp/FRT (*1*, *2*, *23–27*). The second approach exploits the inherent randomness of transfection efficiencies or genomic integration sites (*28–32*). While both established approaches depend on stochastic processes to generate fluorescent protein expression patterns, Caterpie represents a paradigm shift through its implementation of rationally designed color palettes. This deterministic approach enables precise targeting of specific cell populations and lineages. The system holds particular promise for applications requiring high-fidelity identification and longitudinal tracking of distinct cell types within complex biological contexts. However, the full realization of Caterpie's potential necessitates further optimization of delivery methodologies. Specifically, development of refined strategies is required to achieve selective incorporation of distinct FP_{11} tag variants into target cell populations while maintaining variant segregation and preventing cross-contamination.

The theoretical prediction of an 8-fold signal enhancement for FP_{11}(x8) constructs relative to their monomeric counterparts was not fully realized in experimental measurements. Our studies with split CFP2 demonstrated approximately 5-fold enhancement in median fluorescence intensity (Fig. 2C). Further investigations revealed that the choice of fusion protein significantly impacts the signal amplification efficiency of octameric constructs (fig. S2C). In research using sfGFP, when x3, x4, or x7 $sfGFP_{11}$ was fused to β-tubulin or Lamin A/C, length and brightness correlated well, nearly achieving theoretical limits (*4*, *6*). However, when Teneurin-m was fused to x7 $sfGFP_{11}$, only a 4-fold increase in brightness was achieved (*9*). With the goal of reducing gene size, which is an advantage of our approach, we changed from the initially used Histone H2B to ΔSUMOstar and achieved almost the same amplification rate as Histone H2B, x1.6 by each repetition (fig. S2). This collective evidence

suggests that achieving linear brightness amplification through multimerization of the 11th

β-strand requires careful optimization of the fusion partner.

The application of computational protein structure prediction proved instrumental in

enhancing split fluorescent protein performance (Fig. 3). This approach represents a

significant departure from conventional optimization strategies, which typically rely on

random mutagenesis (*33*). Simultaneous introduction of mutations in both $\beta_{1\text{-}10}$ and β_{11}

regions that enhance their mutual interactions proved challenging through random

mutagenesis, because the statistical probability of obtaining cooperative mutations is

exceedingly low. Our successful implementation of structure prediction-guided optimization

suggests a broader applicability of this methodology across other split fluorescent protein

systems.

The discriminative capacity of the Caterpie system holds significant potential for

expansion through two primary avenues: the development of novel split fluorescent protein

variants and the integration of subcellular targeting sequences. Furthermore, compatibility

with the Landing Pad system (*34*), which enables precise single-copy genomic integration,

presents opportunities for multiplexed cellular labeling. These technological developments

collectively position Caterpie as a versatile platform for investigating complex biological

processes requiring simultaneous tracking of multiple distinct cell populations.

Materials and Methods

Plasmids

The following cDNAs were synthesized with optimized codons by GeneArt (Thermo Fisher Scientific, Waltham, MA): sfGFP$_{1-10}$ (*3*), CFP2$_{1-10}$ (*35*), mNeonGreen3A$_{1-10}$ (*11*), sfCherry3Csp$_{1-10}$, mRuby4$_{1-10}$ (*10*), GB1 (*18*), ΔSUMOstar (*19*). Additional split fluorescent protein variants were generated through site-directed mutagenesis of existing templates: Cerulean$_{1-10}$ (*10*) and YFP1$_{1-10}$ (*35*) from sfGFP$_{1-10}$, mNeonGreen3K$_{1-10}$ (*11*) from mNeonGreen3A$_{1-10}$, and sfCherry3C$_{1-10}$ (*12*) from sfCherry2$_{1-10}$ (Addgene: #82603). The Histone H2B coding sequence was subcloned from Addgene (plasmid # 11680 for Histone H2B; Cambridge, MA).

Vector Construction for Stable Cell Line Generation

A base Tol2 transposon vector (pT2A-IRES-iRFP670) was constructed by subcloning iRFP670 cDNA (*36*) with an internal ribosome entry site (IRES) (*37*) into the pT2A vector (*38*). CFP2$_{1-10}$ cDNA was subsequently PCR-amplified and inserted into this base vector using In-Fusion assembly (Clontech, Mountain View, CA) to generate pT2A_CFP2$_{1-10}$-IRES-iRFP670. Two distinct vector backbones, pT2A_IRES-iRFP670 and pT2ADW (containing IRES-puro cassette) (*39*), were used for multicistronic construct assembly. The following elements were PCR-amplified and assembled into both vectors using In-Fusion: sfCherry3Csp$_{1-10}$, mNeonGreen3Asp$_{1-10}$ with self-cleaving P2A peptide (*40*), and CFP2$_{1-10}$ with P2A peptide. Stable cell lines were generated through co-transfection of the constructed pT2A vectors with pCS-TP (obtained from Kawakami et al., 2004 (*38*)).

To generate pPBpuro_EF1a constructs containing CFP2$_{11}$(x1), CFP2$_{11}$(x2), CFP2$_{11}$(x4), CFP2$_{11}$(x8) -mCherry-NLS, cDNAs encoding Histone H2B, self-cleaving peptide P2A sequences, mCherry, and the nuclear localization signal (NLS) of the SV40 large T antigen (PKKKRKV) (*41*) were PCR-amplified and assembled using In-Fusion into pPBpuro_EF1a vectors (a kind gift from K. Yusa), yielding pPBpuro_EF1a_Histone H2B-P2A-mCherry-NLS. Either synthesized CFP2$_{11}$ or CFP2$_{\text{Full-length}}$ cDNA (created by PCR amplification of CFP2$_{1-10}$ cDNA and CFP2$_{11}$) was then assembled into pPBpuro_EF1a_Histone H2B-P2A-mCherry-NLS, resulting in pPBpuro_EF1a_CFP2$_{\text{Full-length}}$/CFP2$_{11}$(x1)-mCherry-NLS. Restriction enzyme sites (BglII before FP$_{11}$, BamHI and NotI after FP$_{11}$) were introduced to facilitate the construction of tandem repeats. To generate CFP2$_{11}$(x2), the CFP2$_{11}$(x1) insert and vector were digested with BglII/NotI and BamHI/NotI, respectively, and then ligated. Subsequently, CFP2$_{11}$(x4) and CFP2$_{11}$(x8) were generated using the same approach. By substituting the CFP2$_{11}$ insert with sfCherry3Csp$_{11}$ or mNeonGreen3Asp$_{11}$ inserts, multiple tandem repeats of these fragments were constructed using the same strategy. Various modifications were made to customize the constructs for specific experimental requirements: the puromycin-resistance gene (puro) was replaced with the blasticidin S-resistance gene (bsr); Histone H2B was substituted with GB1,

ΔSUMOstar; mCherry was replaced with EBFP2 or mTagBFP2; and the NLS was substituted

with the nuclear export signal (NES) of the HIV-1 rev protein (LPPLERLTLD) (*42*).

Mutant variants of sfCherry3C$_{1-10}$ were generated using overlap extension PCR and

assembled into pPB-based vectors (*43*) containing IRES-bsr (blasticidin S-resistance gene)

using In-Fusion. Additional mutant variants of mNeonGreen3A$_{11}$ and sfCherry3C$_{11}$ were

synthesized and inserted into pPB-based vectors (*43*) containing IRES-puro

(puromycin-resistance gene) using Ligation High Ver. 2 (TOYOBO). Stable cell lines were

established through co-transfection of these constructs with pCMV-mPBase (obtained from

the Wellcome Trust Sanger Institute).

Cell culture and establishment of cell lines

HeLa cells and Lenti-X 293T cells were obtained from the Human Science Research

Resources Bank and Clontech, respectively. Both cell lines were maintained in DMEM (Wako

Pure Chemical Industries) supplemented with 10% fetal bovine serum (Sigma-Aldrich) and

1% penicillin/streptomycin (Nacalai Tesque). MDCK (ECACC 84121903) cells were

purchased from the European Collection of Authenticated Cell Cultures (ECACC) through the

RIKEN BioResource Center (no. RCB0995) and maintained in DMEM (Wako Pure Chemical

Industries) supplemented with 10% fetal bovine serum (Sigma-Aldrich), 1%

penicillin/streptomycin (Nacalai Tesque).

For transposon-mediated gene transfer, pT2A_CFP2$_{1-10}$-IRES-iRFP670, or

pT2A_FP$_{1-10}$x3-IRES-iRFP670 was cotransfected with pCS-TP into HeLa cells by using

293fectin (Thermo Fisher Scientific, Waltham, MA). These obtained HeLa cells were sorted

using an FACS Aria IIu cell sorter (Becton Dickinson, Franklin Lakes, NJ) based on iRFP670

fluorescence to achieve a high expression level of the CFP2$_{1-10}$ or FP$_{1-10}$x3. Single-cell

cloning of these sorted populations to yield HeLa$^{CFP2\ 1-10\_iRFP670}$ cells and HeLa$^{FP1-10x3\_iRFP670}$

cells.

pT2A_FP$_{1-10}$x3-IRES-iRFP670 was cotransfected with pCS-TP using either 293fectin

(Thermo Fisher Scientific, Waltham, MA) for HeLa cells or electroporation with an Amaxa

nucleofector (Lonza, Basel, Switzerland) for MDCK cells. The transfected cells were selected

with 5 µg mL^{-1} puromycin (no. P-8833; Sigma-Aldrich), followed by single-cell cloning to

yield HeLa$^{FP1-10x3}$ cells or MDCK$^{FP1-10x3}$ cell lines.

Fluorescence imaging

For evaluation of split fluorescent proteins, HeLa cells in 24-well plates were co-transfected

with 100 ng of FP$_{11}$ plasmid [FP$_{Full-length}$, FP$_{11}$(x1), or FP$_{11}$(x4)] and 400 ng of FP$_{1-10}$ plasmid

using 293fectin (Thermo Fisher Scientific, Waltham, MA). Cells were seeded onto collagen

type I-coated (Nitta Gelatin, Osaka, Japan) glass-based 24-well plates (AGC Inc., Tokyo,

Japan) and cultured for 48 hours. Prior to imaging, cells were equilibrated for at least 1 h in

DMEM/F12 (Thermo Fisher Scientific) supplemented with 10% fetal bovine serum

(Sigma-Aldrich), and penicillin/streptomycin (Nacalai Tesque).

For FP$_{11}$ labeling,

pPBbsr_EF1a-ΔSUMOstar-CFP2$_{11}$(xa)-mNeonGreen3Asp$_{11}$(xb)-sfCherry3Csp$_{11}$(xc) was

cotransfected with pCMV-mPBase into HeLa$^{FP1-10x3}$ cells by using 293fectin or MDCK$^{FP1-10x3}$

cells using electroporation. Transfected cells were selected with 10 μg mL^{-1} blasticidin S

(Calbiochem, San Diego, CA) and seeded onto collagen type I-coated glass-based 24-well

plates.

Cells were observed with a Leica TCS SP8 FALCON confocal microscope

(Leica-Microsystems, Wetzlar, Germany) equipped with an HC PL APO 20x/0.75 dry CS2

objective, an HC PL APO 40x/1.30 OIL CS2 objective, Leica HyD SMD detectors, a white

light laser of 80 MHz pulse frequency, a Diode 405 (VLK 0550 T01; LASOS, Jena,

Germany), a 440 nm diode laser (PDL 800-D; PicoQuant, Berlin, Germany), and a stage top

incubator (Tokai Hit, Fujinomiya, Japan) to maintain 37 °C and 5% CO2. The following

excitation wavelengths and emission band paths were used for the imaging: for EBFP2

imaging, 405 nm excitation, 410–450 nm emission; Cerulean or CFP2 imaging, 440 nm

excitation, 460–490 nm emission; for sfGFP imaging, 488 nm excitation, 500–550 nm

emission; for YFP1 imaging, 514 nm excitation, 520–550 nm emission; for mNG3A, mNG3K,

or mNG3Asp imaging, 506 nm excitation, 510–550 nm emission; for sfCherry3C or

sfCherry3Csp imaging, 594 nm excitation, 600–645 nm emission; for mRuby4 imaging, 561

nm excitation, 580–645 nm emission; for iRFP670 imaging, 650 nm excitation, 660–760 nm

emission. To eliminate the background signal, the time gate for fluorescence detection was set

from 1.0 ns to 6.0 ns.

In the multiplex imaging, the following excitation wavelengths and emission band paths

were used: for CFP2 imaging, 440 nm excitation, 450–500 nm emission; for mNG3Asp

imaging, 506 nm excitation, 515–580 nm emission; for sfCherry3Csp imaging, 594 nm

excitation, 605–645 nm emission. To eliminate the background signal, the time gate for

mNG3Asp and sfCherry3Csp fluorescence detection was set from 0.2 ns to 6.0 ns.

Images were processed and analyzed with ImageJ (*44*). For quantification of fluorescent

intensity, images were segmented using Cellpose algorithm (*21*). Fluorescent intensity

calculation was performed using custom MATLAB (MathWorks) and Python scripts.

Fluorescence spectral analysis

Fluorescence spectra were measured using a Leica TCS SP8 FALCON inverted microscope.

Images were acquired with an HC PL APO 20x/0.75 dry CS2 objective using excitation at

either 470 nm for split mNG3A or 546 nm for split sfCherry3C.

Flow cytometry analysis

Cells were suspended in PBS containing 3% FBS and analyzed or sorted with a FACS Aria

IIu cell sorter (Becton Dickinson, Franklin Lakes, NJ). The following laser and emission filter

combinations were used for fluorescence detection: CFP2, a 407 nm laser and an ET470/24m

filter (Chroma Technology Corp., Bellows Falls, VT); mNeonGreen3Asp, a 488 nm laser, and

a DF530/30 filter; sfCherry3Csp, a 561 nm laser and a DF582/15 filter (Omega Optical); iRFP670, a 633 nm laser and a DF660/20 filter (Omega Optical). Cell debris and aggregates were excluded by gating for size and granularity. Data analysis was performed using FlowJo software (Tree Star, Ashland, OR).

Lentivirus infection

For lentivirus production, the EGF ligand or receptor expressing plasmid, psPAX2 (Addgene no. 12260), and pCMV-VSV-G-RSV-Rev (*45*) were co-transfected into Lenti-X 293T cells using polyethylenimine (no. 24765-1; Polyscience Inc.). The infected cells were selected with media containing the following antibiotics, depending on the drug resistance genes carried by the EGF ligand or receptor expressing plasmids: 200 μg ml^{-1} hygromycin (no. 31282-04-9; Wako).

Cell growth

For quantifying cell growth, MDCK cells labeled by Caterpie were seeded on collagen-coated 24-well glass-bottom plates (AGC Inc., Tokyo, Japan) at a density of 5×10^4 cells/ml. After 1 hour of incubation, the medium was replaced with DMEM/F12 supplemented with 100 units/ml penicillin, 100 μg/ml streptomycin, and 10% FBS. Following cell seeding, cells were observed using a confocal microscope (Leica SP8) every 24 hours for 3 days. For observation conditions, please refer to the "Fluorescent imaging" section. Cell numbers were counted using the Cellpose segmentation algorithm.

Computational design for enhanced β-barrel stability and split-strand complementation

The tertiary structures of split mNG3Asp and split sfCherry3Csp were modeled using ColabFold (*13*) implementation of AlphaFold Multimer (*14*) or AlphaFold2 (*17*) and visualized with PyMOL (http://www.pymol.org/). The highest-confidence model of the split sfCherry3Csp was then adopted as the starting coordinate for downstream Rosetta-based design.

To enhance global β-barrel stability of split sfCherry3Csp, *in silico* site-saturation mutagenesis (SSM) of all 224 positions in sfCherry3C□–□□ and sfCherry3C□□ was performed with a Rosetta-based workflow adapted from Thieker et al (*16*). For each variant, side-chain and backbone sampling were confined to a 10 Å sphere around the mutated residue, while the remainder of the protein was held fixed under coordinate constraints (sd = 1 outside the design sphere; sd = 2 within a surrounding "soft sphere"). Each mutant underwent three cycles of Cartesian FastRelax (MonomerRelax2019), and the change in Rosetta energy (ΔE) was calculated as the difference in total_score between each mutant and the wild-type model using the ref2015 score function.

To reinforce β□–□□/β□□ interface, two design workflows, Point-Mutation (PM) and Mutation Cluster (MC), were adapted from the workflow of Thieker et al. and modified for interface optimization. (1) In the PM workflow, SSM of the β□□ fragment (residues 1–16)

was carried out, with the addition of an InterfaceAnalyzer to compute binding energy
(dG_separated) and packing statistics. Residues within 10 Å of each mutation site were
sampled under the same constraint scheme used for stability design, while non-neighbor
residues were held fixed; each point mutant was relaxed by Cartesian FastRelax prior to
interface analysis, and final dG_separated values were calculated using the unconstrained
ref2015 Rosetta score function. (2) In the MC workflow, the seed mutations from the PM step
were proceeded into a FastDesign protocol using the InterfaceDesign2019 relax script to
generate clustered mutations. The neighbor-selection radius was extended to 12 Å, and an
inner design shell was defined by residues both in direct atomic contact ($\leq$ 5–7 Å, via a
CloseContact selector) and satisfying the InterfaceByVector geometric filter. A
ResidueTypeConstraintGenerator was applied to inner-shell, biasing retention of wild-type
identities. Critical chromophore-interacting residues were protected by marking them
non-designable. Multi-residue combinations sampled via FastDesign, and each cluster variant
was evaluated with InterfaceAnalyzer for dG_separated and packing quality. The scores were
obtained under the unconstrained ref2015 function.
603

Calculation of angular data from individual cells
Three-dimensional fluorescence intensity data were acquired from individual cells by
measuring CFP2, mNG3Asp, and sfCherry3Csp signals with background subtraction. These
measurements were represented in a 3D coordinate system, with CFP2 intensities plotted on
the x-axis, mNG3Asp on the y-axis, and sfCherry3Csp on the z-axis. For each cell, the
Cartesian coordinates (x, y, z) were transformed into spherical coordinates (r, Angle_{CvsG},
Angle_{RvsCG}) using a custom-made Python script. The coordinate transformation was
performed using standard spherical coordinate conversion equations.

$r = (x^2 + y^2 + z^2)^{1/2}$
$\text{Angle}_{CvsG} = \tan^{-1}(y/x)$
$\text{Angle}_{RvsCG} = \cos^{-1}(z/r)$

Two-component Gaussian mixture model (GMM) for FP$_{11}$ tag screening
Fluorescence intensity measurements from individual cells across 96 distinct cell populations
were transformed into angular coordinates (Angle_{CvsG} and Angle_{RvsCG}). To enhance clustering
robustness, outliers were removed by excluding the top and bottom 10% of data points that
exhibited the greatest deviation from their respective cluster centers in both angular
measurements. An unsupervised two-component GMM was implemented using scikit-learn
(version 0.24.2; Pedregosa et al., 2011 (*46*)) to perform pairwise population comparisons. For
random sampling, a predetermined number of cell populations were selected from a total of
96 populations using simple random sampling without replacement. All computational
analyses were performed using custom Python scripts.

Cell classification using Gaussian mixture model (GMM)

Prior to angular calculations, we subtracted background and normalized the fluorescence intensities of CFP2, mNG3Asp, and sfCherry3Csp by dividing each value by the median intensity across all 20 cell types. Using these normalized fluorescence values, we then calculated $Angle_{CvsG}$ and $Angle_{RvsCG}$ measurements for subsequent analysis.

The classification model was trained using GMMs implemented in the scikit-learn library. we first fitted a single-Gaussian distribution to the training subset of each class individually; the resulting class-specific means and covariance matrices were then concatenated to obtain the GMM. The training dataset comprised 80% of the total data ($\sim 2.9 \times 10^4$ cells in total, n > 10^3 cells per cell population), consisting of two-dimensional features ($Angle_{CvsG}$ and $Angle_{RvsCG}$ measurements) from 20 distinct cell populations. The trained GMMs were then applied to angular data from pooled cells, and the predicted cell labels were visualized. For each class k, we calculated the probability of each input data point belonging to the corresponding GMM model, representing the likelihood of class membership:

$$P(x|\theta_k) = N(x|\mu_k, \Sigma_k)$$

where x is the input data point, θ_k represents the parameters of the k-th GMM model, and $N(x|\mu_k, \Sigma_k)$ denotes the normal distribution with mean μ_k and covariance Σ_k. Subsequently, the probabilities were normalized such that the sum of probabilities across all classes equals 1 for each data point:

$$P(k|x) = P(x|\theta_k) / \Sigma_k P(x|\theta_k)$$

We visualized the membership probabilities of the predicted classes. All computational analyses were performed using custom-made Python scripts.

Cell classification using K-means

We used $Angle_{CvsG}$ and $Angle_{RvsCG}$ as the feature space coordinates for clustering. K-means clustering was performed on the resulting dataset using scikit-learn's KMeans algorithm with 21 clusters (random_state = 0). As shown in Figure 6B, cells labeled with tag (BGR)=(0,0,4) corresponding to ID #4 exhibited a wide distribution of values in the $Angle_{CvsG}$-$Angle_{RvsCG}$ space, resulting in their identification as two distinct clusters that were subsequently merged. This accounts for the selection of 21 clusters in our analysis. The predicted cluster labels were then remapped to match the original cell population labels for validation. Classification accuracy was assessed by comparing the remapped cluster assignments with ground truth labels and visualized using a confusion matrix. The overall classification accuracy was calculated as the percentage of correctly classified cells. All calculations were performed using Python.

Clark-Evans Index

We analyzed the spatial distribution patterns of five cell types. Individual cells were first segmented using Cellpose, with centroids calculated and classified into five cell types via the Caterpie method. For spatial analysis, we extracted the coordinates of each cell type, calculated nearest neighbor distances using a k-dimensional tree algorithm, and computed the Clark-Evans index (R). This index compares observed mean distances to those expected under complete spatial randomness, where $R = 1$ indicates random distribution, $R < 1$ indicates clustering, and $R > 1$ indicates regularity. All calculations were performed using Python.

Cell count of the nearest neighbor

We analyzed the spatial relationships between different cell types using a nearest-neighbor approach. Cells were classified into five distinct populations based on fluorescent labels. For each cell of the target population (label 1), we calculated the distance to every cell of other populations (labels 2-5) and identified the closest neighboring cell using a k-dimensional tree algorithm. To assess the statistical significance of the observed patterns, we generated 100 randomized distributions for each dataset by shuffling cell positions while preserving the total number of each cell type. To determine whether the observed spatial associations between cell types differed significantly from random expectations, we calculated the ratio of the frequency with which each non-target cell type appeared as the nearest neighbor to target cells in the original distribution divided by the mean frequency observed across 100 randomized distributions. All calculations were performed using Python.

Statistics

All statistical analyses were carried out using Python. No statistical analysis was used to predetermine the sample size. An unpaired Welch's t-test was used to evaluate statistically significant differences. Data are expressed as the mean $\pm$ s.d. *P*-values of less than 0.05 were considered to be statistically significant in two-tailed tests and were classified into four categories: $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and n.s. (not significant, i.e., $p \geq 0.05$).

Figure Legend

Figure 1. Caterpie: A cell identification system based on split fluorescent protein arrays

(A) Schematic overview of the Caterpie approach, illustrating cell identification through modular arrays of split fluorescent proteins. **(B)** Evolutionary relationships among split fluorescent proteins utilized in this study, presented as a phylogenetic tree. **(C)** Molecular architecture of expression constructs: Histone H2B fusions containing either full-length FP ($FP_{Full-length}$), single FP_{11} fragment, or tetrameric FP_{11} arrays, shown alongside the complementary FP_{1-10} fragment. **(D)** Quantitative analysis of relative fluorescence intensity in HeLa cells expressing $FP_{Full-length}$, $FP_{11}(x1)$ or $FP_{11}(x4)$, with FP_{1-10} fragments (1:4 transfection ratio). Data presented as bee swarm plots with median values indicated by black lines (n > 130 cells per condition). Measurements obtained through confocal microscopy.

Figure 2. Signal amplification and cell classification using split CFP2 arrays

(A) Molecular strategy for generating tandem $CFP2_{11}$ repeats, including detailed amino acid and nucleotide sequences at the $CFP2_{11}(x1)$ insert-vector junction. **(B)** Construct architecture and expression analysis: Upper panel shows schematics $CFP2_{1-10}$ and Histone H2B fusions containing either full-length CFP2 ($CFP2_{Full-length}$) or varying copy numbers of $CFP2_{11}$ [$CFP2_{11}(x1)$, (x2), (x4), and (x8)]. Lower panel presents representative confocal micrographs of HeLa cells stably co-expressing $CFP2_{1-10}$ with either $CFP2_{Full-length}$ or $CFP2_{11}$ variants. Scale bar, 20 µm. **(C)** Quantitative analysis of normalized CFP2 fluorescence in HeLa cells expressing Histone H2B-tagged $CFP2_{11}$ [(x1), (x2), (x4), or (x8)]. Data presented as bee swarm plots with median values (red lines); >1300 cells analyzed across three independent experiments. **(D)** Workflow schematic for Gaussian mixture model (GMM) and implementation. **(E)** Classification performance matrix showing prediction accuracy against true labels (rows). Data represented as mean ± SD of prediction accuracy from three independent experiments, with color intensity indicating classification accuracy. **(F)** Representative ratio image of pooled cells expressing different $CFP2_{11}$ copy number variants [(x1), (x2), (x4), or (x8)]; scale bar: 100 µm. **(G)** Cell classification map derived from panel (F), showing GMM-based assignment of individual cells to specific copy number variants [(x1), (x2), (x4), or (x8)]. **(H)** Visualization of GMM classification confidence through membership probability mapping of cells shown in panel (G).

Figure 3. Computational design of split sfCherry3C

(A) In-silico site-saturation mutagenesis of sfCherry3C□□. Left: schematic of the β□□ residues targeted for saturation mutagenesis (black spheres). Right: heatmap of mean dG_separated values in Rosetta energy unit (REU) across three independent models for each amino acid substitution at each position. (B) Cluster-mutation design of sfCherry3C□□. Left: schematic showing a representative seed point mutation (yellow dot) and the shell of neighboring residues (yellow spheres) targeted for sequential cluster mutagenesis. Right:

heatmap of the mean dG_separated across ten cluster-mutated models derived from each seed

mutation at every β□□ site; dot size indicates the median number of cluster-mutated residues

across these ten models. (C) Schematics of mutant variants of sfCherry3C$_{1-10}$

(sfCherry3C$_{1-10M}$) and Histone H2B-tagged mutant variant of sfCherry3C$_{11}$ (sfCherry3C$_{11M}$).

(D) List of mutant variants of sfCherry3C$_{1-10M}$ and sfCherry3C$_{11M}$. Bee swarm plot showing

normalized sfCherry3C fluorescence intensity of HeLa cells stably co-expressing Histone

H2B-tagged sfCherry3C$_{11M}$(x1) with sfCherry3C$_{1-10M}$. Red lines represent the median. (E)

Rosetta models of split sfCherry3Csp. The overall β-barrel is shown as a cartoon colored by

fragment (gray, β□–□□; magenta, β□□). Each inset zooms on an engineered site, showing

the parental conformation on the left and the corresponding design on the right, with key

residues rendered as sticks representation. All models were visualized in PyMOL. (F)

Emission spectra of HeLa cells expressing split sfCherry3C or split sfCherry3Csp with 546

nm laser excitation.

Figure 4. Optimization of split mNG3A through structure-guided C-terminal

engineering

(A) Predicted structural model of native mNG3A11(x8) generated using ColabFold

(AlphaFold2) and visualized with PyMOL. Colors represent hydrophobicity according to the

Eisenberg hydrophobicity scale. **(B)** Predicted structural model of parental split mNG3A

generated using ColabFold (AlphaFold Multimer) and visualized with PyMOL. Color-coded

domains: mNG3A$_{1-10}$ (green) and mNG3A$_{11}$ (light green). **(C)** Comparative sequence analysis

of original mNG3A$_{11}$ and engineered mutant variants (mNG3A$_{11M}$). **(D)** Schematic

representation of expression constructs: mNG3A$_{1-10}$ and Histone H2B fusion with the

mNG3A$_{11M}$(x8) array. Quantitative analysis of normalized mNG3A fluorescence in HeLa

cells expressing Histone H2B-tagged mNG3A$_{11}$ [(x1), (x2), (x4), or (x8)]. Data presented as

bee swarm plots with median values (red lines). **(E)** Quantitative analysis of normalized

mNG3A fluorescence intensity in HeLa cells stably co-expressing Histone H2B-tagged

mNG3A$_{11M}$ (x8) with mNG3A$_{1-10}$. Median values indicated by red lines. Values normalized to

set the median value of mNG3A$_{11M1}$(x8) to 1. **(F)** Emission spectra of HeLa cells expressing

split mNG3A or split mNG3Asp with 470 nm laser excitation. **(G)** Comparative analysis of

relative fluorescence intensity of HeLa cells expressing FP$_{Full-length}$, FP$_{11}$(x1) or FP$_{11}$(x4), with

cognate FP$_{1-10}$ fragments (1:4 transfection ratio). Data presented as bee swarm plots with

median values (red lines); n > 130 cells per condition. Measurements obtained through

confocal microscopy.

Figure 5. Systematic selection and validation of FP$_{11}$ tags for Caterpie implementation

(A) Schematic representation of FP$_{11}$ tags: ΔSUMO-tagged FP$_{11}$ arrays comprising paired

repeats of CFP2$_{11}$, mNG3Asp$_{11}$, and sfCherry3Csp$_{11}$ [(x0), (x1), (x2), (x4), (x8) each].

Comprehensive catalog of the FP$_{11}$ tag library comprising 96 distinct combinations, detailing

copy numbers of CFP2$_{11}$ (labeled as "C"), mNG3Asp$_{11}$ (labeled as "G"), and sfCherry3Csp$_{11}$

(labeled as "R") fragments. The marks indicate the FP_{11} tags selected during primary and
secondary screening processes, and those finally chosen. **(B)** Schematic representation of
expression constructs: $FP_{1-10}x3$. **(C)** Analysis scheme for classification of cell populations
expressing various FP_{11} tags. **(D)** Matrix analysis of pairwise classification accuracy across 96
cell populations expressing various FP_{11} tags, arranged by FP_{11} tag numbers shown in panel
(A). (n > 230 cells per population). **(E)** Analysis scheme for determining the highest
achievable minimum pairwise classification accuracy among randomly selected FP_{11}-tagged
subpopulations. **(F)** Graph showing the highest achievable minimum pairwise classification
accuracy among randomly selected FP_{11}-tagged subpopulations. For each subpopulation size,
minimum accuracy between all possible pairs was determined from 10^7 random selections,
with maximum values plotted. **(G)** Comparative performance analysis of 30 candidate
populations (detailed in fig. S4) using both confocal microscopy (blue) and flow cytometry
(orange). Analysis methodology follows that of panel (F). Data presented as mean ± SD from
three independent experiments.

Figure 6. High-fidelity discrimination of twenty distinct cell populations using Caterpie
(A) Representative multicolor fluorescence micrographs of 20 distinct cell populations
expressing unique FP_{11} tag combinations. Fluorescence channels: CFP2 (blue), mNG3Asp
(green), and sfCherry3Csp (red). Copy numbers of each FP_{11} variant (CFP2$_{11}$, mNG3Asp$_{11}$,
and sfCherry3Csp$_{11}$) are indicated in the upper left corner of each panel. Scale bars: 30 μm.
(B) Left: Scatter plot showing Angle$_{CvsG}$ versus Angle$_{RvsCG}$ distributions for 20 distinct cell
populations from panel (A). (n > 1000 cells per population) Center: Distribution of label
assignments in Angle$_{CvsG}$ and Angle$_{RvsCG}$ based on Gaussian mixture model (GMM)
classification. Right: Distribution of label membership probabilities in Angle$_{CvsG}$ and
Angle$_{RvsCG}$. **(C)** Classification performance matrix showing prediction accuracy against true
population identities (rows). Color intensity indicates classification accuracy. Overall average
accuracy, 97 %. **(D)** Large-field composite image of pooled cell populations from panel (A),
displaying CFP2 (blue), mNG3Asp (green), and sfCherry3Csp (red) fluorescence channels.
Scale bar: 500 μm. **(E)** Detailed analysis of region indicated by dashed box in panel (D). Left:
Higher magnification of selected region. Center: Population assignment map following
GMM-based classification into 20 distinct categories. Right: Visualization of GMM
classification reliability through membership probability mapping.

Figure 7. Interaction dynamics in multi-population cell cultures of MDCK cells
expression EGF-family ligands and their receptors
(A) Schematic diagram showing the association pairs between EGF ligands (HBEGF, EREG)
and EGF receptors (EGFR, ErbB2). **(B)** Representative pseudocolored image of five distinct
cell types cultured together and identified using the Caterpie method. Yellow: WT, Pink:
EREG, Cyan: HBEGF, Green: EGFR, Orange: ErbB4; Copy numbers of each FP11 variant
(CFP211, mNG3Asp11, and sfCherry3Csp11) are (1,2,4), (2,0,4), (4,0,1), (2,4,0), and (8,2,1),

respectively. Scale bar: 1 mm. **(C)** Individual display of the five cell types shown in panel E.

The upper left corner of each panel indicates the re-expressed EGF signaling molecules and

the copy numbers of FP_{11} tags. **(D)** Graph showing the Clark-Evans Index calculated for each

of the five cell types. Data from three independent experiments. Black lines represent the

mean values across the three experiments. **(E)** Normalized nearest neighbor cell analysis for

evaluating spatial relationships between cells. This figure shows the ratio of heterotypic

nearest neighbor counts observed in the actual data to those counted after randomly

reassigning the labels of five cell types in the same dataset. Data from three independent

experiments. Black lines represent the mean values across the three experiments.

Supplementary Figure S1. Comprehensive mutational scanning of split sfCherry3C

Heatmap displaying the ΔEnergy values for saturating mutations in both $sfCherry_{1-10}$ and

$sfCherry_{11}$.

Supplementary Figure S2. Optimization of signal amplification using truncated

SUMOstar fusion strategy

(A) Schematic representation of expression constructs: $CFP2_{Full-length}$-NES and $CFP2_{11}$

variants [(x1) or (x8)]-NES with or without GB1 or ΔSUMOstar (truncated SUMOstar) tags.

(B) Amino acid sequences of SUMOstar, with ΔSUMOstar region highlighted in bold. **(C)**

Bee swarm plots showing quantitative analysis of normalized CFP2 fluorescence in HeLa

cells co-expressing $CFP2_{1-10}$ with either $CFP2_{Full-length}$-NES or $CFP2_{11}(x1)/CFP2_{11}(x8)$-NES

variants, with or without GB1 or ΔSUMOstar tags (constructs shown in panel A). Median

values are indicated by red lines. **(D)** Construct architecture and expression analysis: Upper

panel shows schematics of ΔSUMOstar fusions containing varying copy numbers of $CFP2_{11}$

[$CFP2_{11}(x1)$, (x2), (x4), and (x8)]. Lower panel presents representative confocal micrographs

of HeLa cells stably co-expressing $CFP2_{1-10}$ with either $CFP2_{Full-length}$-NES or

ΔSUMOstar-tagged $CFP2_{11}$ variant. Scale bar: 20 □µm. **(E)** Comparative analysis of

normalized CFP2 fluorescence in HeLa cells expressing ΔSUMOstar-tagged $CFP2_{11}$ variants

[(x1), (x2), (x4), or (x8)] shown in panel D. Data presented as bee swarm plots with median

values indicated by red lines. **(F)** Classification performance matrix showing predictive

accuracy against true population identities (rows), with color intensity indicating

classification accuracy.

Supplementary Figure S3. Influence of FP_{11} fragment order on fluorescence signal

distribution

(A) Schematic representation of expression constructs: ΔSUMO-tagged FP_{11} arrays

comprising paired repeats of $CFP2_{11}$, $mNG3Asp_{11}$, and $sfCherry3Csp_{11}$ [(x2) each, total x6].

(B) Quantitative analysis presented as bee swarm plots showing normalized fluorescence

intensities (CFP2, mNG3Asp, and sfCherry3Csp) in HeLa cells expressing six different

permutations of FP_{11} repeat arrangements, with concurrent FP_{1-10}x3 expression. Data obtained

by confocal microscopy, with median values indicated by black lines.

Supplementary Figure S4. Comparative analysis of binary classification performance
across 30 selected cell populations
(A) Catalog of 30 selected FP_{11} tag variants (detailed in Fig. 5A), detailing identification
numbers and copy numbers of $CFP2_{11}$, $mNG3Asp_{11}$, and $sfCherry3Csp_{11}$ fragments. **(B)**
Matrix analysis of pairwise classification accuracies for 30 cell populations expressing
distinct FP_{11} tag combinations (listed in panel A), evaluated by both microscopy (upper panel)
and flow cytometry (lower panel). Data represent results from three independent experiments.

Supplementary Figure S5. Binary classification performance analysis of final 20 cell
population panel
(A) Catalog of optimized FP_{11} tag set comprising 20 selected variants (detailed in Fig. 5A),
detailing identification numbers and copy numbers of $CFP2_{11}$, $mNG3Asp_{11}$, and
$sfCherry3Csp_{11}$ fragments. **(B)** Matrix analysis of pairwise classification accuracy for 20 cell
populations expressing distinct FP_{11} tag combinations (defined in panel A). Comparative
evaluation performed using both microscopy (left panel) and flow cytometry (right panel).
Data represent mean values from three independent experiments in Supplementary Figure S3.

Supplementary Figure S6. Aggregation character of $sfCherry3Csp_{11}$(x8)
Representative images of sfCherry3Csp fluorescence in HeLa cells expressing
ΔSUMOstar-tagged $sfCherry3Csp_{11}$ variants [(x4), or (x8)], with concurrent FP_{1-10}x3
expression.

Supplementary Figure S7. Unsupervised k-means clustering of 20 cell populations
(A) Left: Scatter plot showing the distribution of $Angle_{CvsG}$ versus $Angle_{RvsCG}$ for 20 distinct
cell populations (n > 1,000 cells per population). Right: Scatter plot showing the distribution
of $Angle_{CvsG}$ versus $Angle_{RvsCG}$ for individual cells grouped into 20 distinct populations using
unsupervised k-means clustering. **(B)** Classification performance matrix of unsupervised
k-means classification in panel (A). Color intensity indicates classification accuracy. Overall
average accuracy, 96 %.

Supplementary Figure S8. High-fidelity discrimination of twenty distinct cell populations
in MDCK cells using Caterpie
(A) Representative multicolor fluorescence micrographs of 20 distinct cell populations
expressing FP_{11} tags. Fluorescence channels: CFP2 (blue), mNG3Asp (green), and
sfCherry3Csp (red). Copy numbers of each FP_{11} variant ($CFP2_{11}$, $mNG3Asp_{11}$, and
$sfCherry3Csp_{11}$) are indicated in the top of each panel. Scale bars: 30 μm. **(B)** Classification
performance matrix showing prediction accuracy against true population identities (rows).
Color intensity indicates classification accuracy. Overall average accuracy, 96%. **(C)** Top:
Large-field composite image of pooled cell populations from panel (A), displaying CFP2

(blue), mNG3Asp (green), and sfCherry3Csp (red) fluorescence channels. Scale bar: 500 µm.

Middle: Population assignment map following GMM-based classification into 20 distinct

categories. Bottom: Visualization of GMM classification confidence through membership

probability mapping.

Supplementary Figure S9. Cell growth of Caterpie labeled MDCK cells

Growth curves showing cell density of 20 different MDCK cell types fluorescently labeled

with the Caterpie method measured every 24 hours. The lower panels display individual

growth curves for each cell type.

References

1. J. Livet, T. A. Weissman, H. Kang, R. W. Draft, J. Lu, R. A. Bennis, J. R. Sanes, J. W. Lichtman, Transgenic strategies for combinatorial expression of fluorescent proteins in the nervous system. Nature 450, 56-62 (2007).

2. M. W. Veling, Y. Li, M. T. Veling, C. Litts, N. Michki, H. Liu, B. Ye, D. Cai, Identification of neuronal lineages in the Drosophila peripheral nervous system with a "digital" multi-spectral lineage tracing system. Cell Rep. 29, 3303-3312.e3 (2019).

3. S. Cabantous, T. C. Terwilliger, G. S. Waldo, Protein tagging and detection with engineered self-assembling fragments of green fluorescent protein. Nat. Biotechnol. 23, 102-107 (2005).

4. M. D. Leonetti, S. Sekine, D. Kamiyama, J. S. Weissman, B. Huang, A scalable strategy for high-throughput GFP tagging of endogenous human proteins. Proc. Natl. Acad. Sci. U.S.A. 113, E3501-E3508 (2016).

5. N. H. Cho, K. C. Cheveralls, A. D. Brunner, K. Kim, A. C. Michaelis, P. Raghavan, H. Kobayashi, L. Savy, J. Y. Li, H. Canaj, J. Y. S. Kim, E. M. Stewart, C. Gnann, F. McCarthy, J. P. Cabrera, R. M. Brunetti, B. B. Chhun, G. Dingle, M. Y. Hein, B. Huang, S. B. Mehta, J. S. Weissman, R. Gómez-Sjöberg, D. N. Itzhak, L. A. Royer, M. Mann, M. D. Leonetti, OpenCell: Endogenous tagging for the cartography of human cellular organization. Science 375, eabi6983 (2022).

6. D. Kamiyama, S. Sekine, B. Barsi-Rhyne, J. Hu, B. Chen, L. A. Gilbert, H. Ishikawa, M. D. Leonetti, W. F. Marshall, J. S. Weissman, B. Huang, Versatile protein tagging in cells with split fluorescent protein. Nat. Commun. 7, 11046 (2016).

7. S. He, A. Cuentas-Condori, D. M. Miller, Strategy for bright, tissue-specific GFP labeling of endogenous proteins in C. elegans. Genetics 212, 387-395 (2019).

8. J. Goudeau, C. S. Sharp, J. Paw, L. Savy, M. D. Leonetti, A. G. York, D. L. Updike, C. Kenyon, M. Ingaramo, Split-wrmScarlet and split-sfGFP: tools for faster, easier fluorescent labeling of endogenous proteins in Caenorhabditis elegans. Genetics 217, iyab014 (2021).

9. R. Kamiyama, K. Banzai, P. Liu, A. Marar, R. Tamura, F. Jiang, M. A. Fitch, J. Xie, D. Kamiyama, Cell-type-specific, multicolor labeling of endogenous proteins with split fluorescent protein tags in Drosophila. Proc. Natl. Acad. Sci. U.S.A. 118, e2024690118 (2021).

10. R. Tamura, F. Jiang, J. Xie, D. Kamiyama, Multiplexed labeling of cellular proteins with split fluorescent protein tags. Commun. Biol. 4, 257 (2021).

11. S. Zhou, S. Feng, D. Brown, B. Huang, Improved yellow-green split fluorescent proteins for protein labeling and signal amplification. PLoS One 15, e0242592 (2020).

12. S. Feng, A. Varshney, D. Coto Villa, C. Modavi, J. Kohler, F. Farah, S. Zhou, N. Ali, J. D. Müller, M. K. Van Hoven, B. Huang, Bright split red fluorescent proteins for the

visualization of endogenous proteins and synapses. Commun. Biol. 2, 344 (2019).

13. G. Kim, S. Lee, E. Levy Karin, H. Kim, Y. Moriwaki, S. Ovchinnikov, M. Steinegger, M.
Mirdita, Easy and accurate protein structure prediction using ColabFold. Nat. Protoc. 20,
620–642 (2025).

14. R. Evans, M. O'Neill, A. Pritzel, N. Antropova, A. Senior, T. Green, A. Žídek, R. Bates,
S. Blackwell, J. Yim, O. Ronneberger, S. Bodenstein, M. Zielinski, A. Bridgland, A.
Potapenko, A. Cowie, K. Tunyasuvunakool, R. Jain, E. Clancy, P. Kohli, J. Jumper, D.
Hassabis, Protein complex prediction with AlphaFold-Multimer. bioRxiv
2021.10.04.463034 [Preprint] (2022); https://doi.org/10.1101/2021.10.04.463034.

15. A. Leaver-Fay, M. Tyka, S. M. Lewis, O. F. Lange, J. Thompson, R. Jacak, K. Kaufman,
P. D. Renfrew, C. A. Smith, W. Sheffler, I. W. Davis, S. Cooper, A. Treuille, D. J.
Mandell, F. Richter, Y.-E. A. Ban, S. J. Fleishman, J. E. Corn, D. E. Kim, S. Lyskov, M.
Berrondo, S. Mentzer, Z. Popović, J. J. Havranek, J. Karanicolas, R. Das, J. Meiler, T.
Kortemme, J. J. Gray, B. Kuhlman, D. Baker, P. Bradley, ROSETTA3: An
object-oriented software suite for the simulation and design of macromolecules. *Methods*
Enzymol. **487**, 545–574 (2011).

16. D. F. Thieker, J. B. Maguire, S. T. Kudlacek, A. Leaver-Fay, S. Lyskov, B. Kuhlman,
Stabilizing proteins, simplified: A Rosetta-based webtool for predicting favorable
mutations. Protein Sci. 31, e4428 (2022).

17. J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K.
Tunyasuvunakool, R. Bates, A. Žídek, A. Potapenko, A. Bridgland, C. Meyer, S. A. A.
Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back,
S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T.
Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K. Kavukcuoglu, P.
Kohli, D. Hassabis, Highly accurate protein structure prediction with AlphaFold. Nature
596, 583-589 (2021).

18. P. Zhou, A. A. Lugovskoy, G. Wagner, A solubility-enhancement tag (SET) for NMR
studies of poorly behaving proteins. J. Biomol. NMR 20, 11-14 (2001).

19. L. Liu, J. Spurrier, T. R. Butt, J. E. Strickler, Enhanced protein expression in the
baculovirus/insect cell system using engineered SUMO fusions. Protein Expr. Purif. 62,
21-28 (2008).

20. N. Milech, B. A. Longville, P. T. Cunningham, M. N. Scobie, H. M. Bogdawa, S.
Winslow, M. Anastasas, T. Connor, F. Ong, S. R. Stone, M. Kerfoot, T. Heinrich, K. M.
Kroeger, Y. F. Tan, K. Hoffmann, W. R. Thomas, P. M. Watt, R. M. Hopkins,
GFP-complementation assay to detect functional CPP and protein delivery into living
cells. Sci. Rep. 5, 18329 (2015).

21. C. Stringer, T. Wang, M. Michaelos, M. Pachitariu, Cellpose: a generalist algorithm for
cellular segmentation. Nat. Methods 18, 100-106 (2021).

22. E. Deguchi, S. Lin, D. Hirayama, K. Matsuda, A. Tanave, K. Sumiyama, S. Tsukiji, T.

Otani, M. Furuse, A. Sorkin, M. Matsuda, K. Terai, Low-affinity ligands of the epidermal
growth factor receptor are long-range signal transmitters in collective cell migration of
epithelial cells. Cell Rep. 43, 114986 (2024).

23. H. J. Snippert, L. G. van der Flier, T. Sato, J. H. van Es, M. van den Born, C.
Kroon-Veenboer, N. Barker, A. M. Klein, J. van Rheenen, B. D. Simons, H. Clevers,
Intestinal crypt homeostasis results from neutral competition between symmetrically
dividing Lgr5 stem cells. Cell 143, 134-144 (2010).

24. I. Tabansky, A. Lenarcic, R. W. Draft, K. Loulier, D. B. Keskin, J. Rosains, J.
Rivera-Feliciano, J. W. Lichtman, J. Livet, J. N. H. Stern, J. R. Sanes, K. Eggan,
Developmental bias in cleavage-stage mouse blastomeres. Curr. Biol. 23, 21-31 (2013).

25. C. Ghigo, I. Mondor, A. Jorquera, N. J. Nowak, S. Wienert, S. P. Zahner, B. E. Clausen,
H. Luche, B. Malissen, F. Klauschen, M. Bajénoff, Multicolor fate mapping of
Langerhans cell homeostasis. J. Exp. Med. 210, 1657-1664 (2013).

26. D. Cai, K. B. Cohen, T. Luo, J. W. Lichtman, J. R. Sanes, Improved tools for the
Brainbow toolbox. Nat. Methods 10, 540-547 (2013).

27. K. Loulier, R. Barry, P. Mahou, Y. Le Franc, W. Supatto, K. S. Matho, S. Ieng, S.
Fouquet, E. Dupin, R. Benosman, A. Chédotal, E. Beaurepaire, X. Morin, J. Livet,
Multiplex cell and lineage tracking with combinatorial labels. Neuron 81, 505-520
(2014).

28. K. Weber, U. Bartsch, C. Stocking, B. Fehse, A multicolor panel of novel lentiviral "gene
ontology" (LeGO) vectors for functional gene analysis. Mol. Ther. 16, 698-706 (2008).

29. J. García-Marqués, L. López-Mascaraque, Clonal identity determines astrocyte cortical
heterogeneity. Cereb. Cortex 23, 1463-1472 (2013).

30. F. García-Moreno, N. A. Vasistha, J. Begbie, Z. Molnár, CLoNe is a new method to
target single progenitors and study their progeny in mouse and chick. Development 141,
1589-1598 (2014).

31. R. Sakaguchi, M. N. Leiwe, T. Imai, Bright multicolor labeling of neuronal circuits with
fluorescent proteins and chemical tags. eLife 7, e40350 (2018).

32. T. Kumamoto, F. Maurinot, R. Barry-Martinet, C. Vaslin, S. Vandormael-Pournin, M. Le,
M. Lerat, D. Niculescu, M. Cohen-Tannoudji, A. Rebsam, K. Loulier, S. Nedelec, S.
Tozer, J. Livet, Direct readout of neural stem cell transgenesis with an
integration-coupled gene expression switch. Neuron 107, 617-630.e6 (2020).

33. S. Feng, S. Sekine, V. Pessino, H. Li, M. D. Leonetti, B. Huang, Improved split
fluorescent proteins for endogenous protein labeling. Nat. Commun. 8, 370 (2017).

34. K. A. Matreyek, J. J. Stephany, M. A. Chiasson, N. Hasle, D. M. Fowler, An improved
platform for functional assessment of large protein libraries in mammalian cells. Nucleic
Acids Res. 48, e1 (2020).

35. T. Köker, A. Fernandez, F. Pinaud, Characterization of split fluorescent protein variants
and quantitative analyses of their self-assembly process. Sci. Rep. 8, 5344 (2018).

36. D. M. Shcherbakova, V. V. Verkhusha, Near-infrared fluorescent proteins for multicolor
in vivo imaging. Nat. Methods 10, 751-754 (2013).

37. T. Akagi, K. Sasai, H. Hanafusa, Refractory nature of normal human diploid fibroblasts
with respect to oncogene-mediated transformation. Proc. Natl. Acad. Sci. U.S.A. 100,
13567-13572 (2003).

38. K. Kawakami, H. Takeda, N. Kawakami, M. Kobayashi, N. Matsuda, M. Mishina, A
transposon-mediated gene trap approach identifies developmentally regulated genes in
zebrafish. Dev. Cell 7, 133-144 (2004).

39. N. Komatsu, K. Terai, A. Imanishi, Y. Kamioka, K. Sumiyama, T. Jin, Y. Okada, T.
Nagai, M. Matsuda, A platform of BRET-FRET hybrid biosensors for optogenetics,
chemical screening, and in vivo imaging. Sci. Rep. 8, 8984 (2018).

40. J. H. Kim, S. R. Lee, L. H. Li, H. J. Park, J. H. Park, K. Y. Lee, M. K. Kim, B. A. Shin, S.
Y. Choi, High cleavage efficiency of a 2A peptide derived from porcine teschovirus-1 in
human cell lines, zebrafish and mice. PLoS One 6, e18556 (2011).

41. D. Kalderon, B. L. Roberts, W. D. Richardson, A. E. Smith, A short amino acid sequence
able to specify nuclear location. Cell 39, 499-509 (1984).

42. U. Fischer, J. Huber, W. C. Boelens, L. W. Mattaj, R. Lührmann, The HIV-1 Rev
activation domain is a nuclear export signal that accesses an export pathway used by
specific cellular RNAs. Cell 82, 475-483 (1995).

43. K. Yusa, R. Rad, J. Takeda, A. Bradley, Generation of transgene-free induced pluripotent
mouse stem cells by the piggyBac transposon. Nat. Methods 6, 363-369 (2009).

44. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image
analysis. Nat. Methods 9, 671-675 (2012).

45. Miyoshi H., Blomer U., Takahashi M., Gage F. H., Verma I. M., *J. Virol.* **72**, 8150
(1998).

46. D. K. Barupal, O. Fiehn, Generating the blood exposome database using a
comprehensive text mining and database fusion approach. Environ. Health Perspect. 127,
97008 (2019).

1029

Fig.1

A

CFP$_{1-10}$ RFP$_{1-10}$ GFP$_{1-10}$

FP$_{11}$ tag

CFP$_{11}$ repeats   RFP$_{11}$ repeats   GFP$_{11}$ repeats

Self-complementation

1. Labeling

HeLa$^{Triple1-10}$ Cell

20 FP$_{11}$ tags

2. Observation and model training

20 cell populations

Model training with
a Gaussian mixture
model (GMM)

Angle$_{RvsCG}$

Angle$_{CvsG}$

3. Classification (20 cells, 97%)

Pooled cells

Applying the trained GMM

Angle$_{RvsCG}$

Angle$_{CvsG}$

B

sfGFP — Cerulean
         CFP2
         YFP1

mNG2 — mNG3A
        mNG3K

sfCherry2 — sfCherry3C

mRuby4

C

FP$_{Full}$ Histone H2B FP$_{Full-length}$ EBFP2-NLS
                                    V2A

FP$_{11}$(x1) Histone H2B   EBFP2-NLS
                     FP$_{11}$

FP$_{11}$(x4) Histone H2B   EBFP2-NLS

FP$_{1-10}$ FP$_{1-10}$ IRES iRFP670

D

Relative fluorescence intensity

- FP$_{Full-length}$ + FP$_{1-10}$
- FP$_{11}$(x1) + FP$_{1-10}$
- FP$_{11}$(x4) + FP$_{1-10}$

Cerulean CFP2 sfGFP sYFP1 mNG3A mNG3K sfCherry3C mRuby4

Fig.2

Fig.3

Fig.4

A mNG3A₁₁(x8)

F5
W8
F12
M15
M16

High
Hydrophobicity
Low

B

M16
M15
D14 F12
T13

C

		5	10	15

mNG3A₁₁   TELNFKEWQKAFTDMM
M1   TELNFKEWQKAFT
M2   TELNFKEWQKAFTD
mNG3Asp₁₁   TELNFKEWQKAFTD**D**
M4   TELNFKEWQKA**Y**T
M5   TELNFKEWQKA**Y**TD
M6   TELNFKEWQKA**Y**TD**D**

D

mNG3A₁₋₁₀

H2B [   ] mTagBFP2-NLS

mNG3A₁₁M₁₋₆ (x8)

E

Normalized mNG intensity

mNG3A₁₁ M1 M2 mNG3Asp₁₁ M4 M5 M6

F

Relative intensity

— mNG3A₁₁(x1)
— mNG3Asp₁₁(x1)

500   540   580   620
Wavelength (nm)

G

CFP2   mNG3Asp   sfCherry3Csp

Relative fluorescence intensity

Full-length (x1) (x4)   Full-length (x1) (x4)   Full-length (x1) (x4)

+FP₁₋₁₀

Fig5

A FP₁₁ tag library:

B FP₁₋₁₀x3_iRFP670: sfCherry3Csp₁₋₁₀ mNG3A₁₋₁₀ CFP2₁₋₁₀ IRES iRFP670

Fig.6

A

HeLa cells
ID: (The repeat number of CFP2$_{11}$, mNG3Asp$_{11}$, sfCherry3Csp$_{11}$)

1: (1,4,4) 2: (0,4,8) 3: (0,2,8) 4: (0,0,4) 5: (2,0,4) 6: (1,2,4) 7: (2,2,8) 8: (4,4,4) 9: (2,1,2) 10: (4,0,1)

11: (4,1,1) 12: (8,2,1) 13: (8,1,0) 14: (8,0,0) 15: (2,2,0) 16: (2,4,0) 17: (1,8,2) 18: (1,8,0) 19: (0,8,0) 20: (0,8,2)

B

Raw data
Angle$_{RvsCG}$ (degrees)
Angle$_{CvsG}$ (degrees)

GMM label assignment
Angle$_{RvsCG}$ (degrees)
Angle$_{CvsG}$ (degrees)

GMM label probability
Angle$_{RvsCG}$ (degrees)
Angle$_{CvsG}$ (degrees)
Membership probabilities (%)

C

Predicted label
True label
Accuracy (%)

D

500 μm

E

RGB image

GMM label map
Identified label

GMM label reliability map
Membership probability (%)

Fig.7

A

HBEGF, EREG

EGFR (ErbB1)   ErbB4

Caterpi labeled MDCK cells
(1,2,4) (2,0,4) (4,0,1) (2,4,0) (8,2,1)

WT EREG HBEGF EGFR ErbB4

Over-expression

Co-culture for 8 day (3% FBS)

Ligand over-expressing cell

WT cell

Receptor over-expressing cell

B

HBEGF   EGFR
WT   EREG   ErbB4

1 mm

C

WT (1,2,4)

HBEGF (4,0,1)

EREG (2,0,4)

EGFR (2,4,0)

ErbB4 (8,2,1)

1 mm

Splitting each cell types

D

Clark-Evans Index

n.s.

WT EREG HBEGF EGFR ErbB4

E

Normalized cell count of nearest neighbor

n.s.   **

EREG
WT EGFR ErbB4

n.s.   ***

HBEGF
WT EGFR ErbB4

**   n.s.

ErbB1
WT EREG HBEGF

*   n.s.

ErbB4
WT EREG HBEGF

Sup1

A

Full CFP2_Full mCherry-NLS
NES P2A

CFP2_11(x1), (x8)
x1
x8 mCherry-NLS
NES
→ GB1, ΔSUMOstar

B

ΔSUMOstar

(MGHHHHHGSLQDSEVNQEAKPEVK
PEVKPE)**THINLKVSDGSSEIFFKIK
KTTPLRRLMEAFAKRQGKEMDSLTF
LYDGIEIQADQTPEDLDMEDNDIIE
AHREQ**(IGGRDDLPSL)

C

Normalized CFP2 intensity

Full
(x1)
(x8)
***

Full-length None GB1 ΔSUMOstar

D

CFP2_11(x1), (x2), (x4), (x8)
ΔSUMOstar mCherry-NLS
NES

Full-length (x1) (x2) (x4) (x8)

CFP2

mCherry

E

Normalized CFP2 intensity

Full-length x1 x2 x4 x8

F

Predicted labels

	x1	x2	x4	x8
x1	87.8	2.3	0.0	0.0
x2	11.0	95.4	6.2	0.0
x4	1.2	1.1	91.2	2.5
x8	0.0	1.1	2.5	97.5

True labels

Sup. 3

A

Shuffling

B

Sup. 4

A

ID: (The repeat number of CFP2$_{11}$, mNG3Asp$_{11}$, sfCherry3Csp$_{11}$)

1: (4,0,0)   6: (1,8,0)   11: (2,0,4)   16: (0,8,2)   21: (1,8,1)   26: (4,1,1)
2: (0,2,0)   7: (2,2,0)   12: (4,0,1)   17: (0,8,4)   22: (1,8,2)   27: (4,1,4)
3: (0,8,0)   8: (2,4,0)   13: (0,2,8)   18: (1,1,2)   23: (2,1,1)   28: (4,2,2)
4: (0,0,2)   9: (8,1,0)   14: (0,4,8)   19: (1,2,4)   24: (2,1,2)   29: (4,4,4)
5: (0,0,8)   10: (1,0,8)   15: (0,8,1)   20: (1,4,4)   25: (2,2,8)   30: (8,2,1)

B

Sup. 5

A

ID: (The repeat number of CFP2₁₁, mNG3Asp₁₁, sfCherry3Csp₁₁)

$\text{ID: (The repeat number of } \boxed{\text{CFP2}_{11}}, \boxed{\text{mNG3Asp}_{11}}, \boxed{\text{sfCherry3Csp}_{11}})$

1: (4,0,0)   5: (2,2,0)   9: (4,0,1)   13: (1,2,4)   17: (2,2,8)
2: (0,2,0)   6: (2,4,0)   10: (0,2,8)   14: (1,4,4)   18: (4,1,1)
3: (0,0,8)   7: (8,1,0)   11: (0,4,8)   15: (1,8,2)   19: (4,4,4)
4: (1,8,0)   8: (2,0,4)   12: (0,8,2)   16: (2,1,2)   20: (8,2,1)

B

Microscopy   Flow Cytometry

Classification accuracy (%)

Sup.6

sfCherry3Csp$_{11}$(x4)   sfCherry3Csp$_{11}$(x8)

50 um

Sup.7

A

B

Sup.8

A

MDCK cells

ID: (CFP2₁₁ , mNG3Asp₁₁ , sfCherry3Csp₁₁)

30 µm

B

C

Sup.9