Dominant spinal muscular atrophy linked mutations in the cargo binding domain of
BICD2 result in altered interactomes and dynein hyperactivity.

Hannah Neiswender[1], Jessica E. Pride[1], Rajalakshmi Veeranan-Karmegam[1], Phylicia Allen[1],
Grace Neiswender[1], Avneesh Prabakar[1], Caili Hao[1], Xingjun Fan[1], and Graydon B. Gonsalvez[1]

[1]Cellular Biology and Anatomy
Medical College of Georgia, Augusta University
1460 Laney Walker Blvd
Augusta, GA, 30912, USA
11

12
[*]Address correspondence to
Graydon Gonsalvez (ggonsalvez@augusta.edu)
Cellular Biology and Anatomy
Medical College of Georgia, Augusta University
1460 Laney Walker Blvd. CB2917
Augusta, GA 30912, USA
Tel: (706) 721-1756

ABSTRACT

Cytoplasmic dynein-1 (dynein) is responsible for the transport of most cellular cargo towards the
minus end of microtubules. Dynein activation requires the multi-subunit dynactin complex and
an activating cargo adaptor. The adaptors serve to link dynein with cargo and to fully activate
the motor. Mutations in one of these activating adaptors, Bicaudal-D2 (BICD2), are associated
with a neurodegenerative disease called Spinal Muscular Atrophy with Lower Extremity
Predominance (SMALED2). The molecular defect that underlies SMALED2 is largely unknown.
In addition to interacting with dynein, BICD2 has also been shown to associate with KIF5B, a
plus-end directed microtubule motor. We hypothesized that interactome changes associated
with mutant versions of BICD2, and the resulting differences in cargo transport, might underlie
the etiology of SMALED2. To test our hypothesis, we first defined the interactome of wild-type
BICD2. This led to the identification of known BICD2 interacting proteins in addition to
potentially novel cargo such as components of the HOPS complex, a six-subunit complex
involved in endo-lysosomal trafficking. We next determined the interactome of three SMALED2
linked mutants in BICD2, two of which reside in the cargo binding domain. Interestingly, all three
mutations resulted in BICD2-mediated dynein hyper-activation. Furthermore, all three mutants
were associated with interactome changes. One of these mutants, BICD2_R747C, was deficient
in binding to HOPS complex components and the nucleoporin RANBP2. In addition, this mutant
also resulted in a gain of function interaction with GRAMD1A, a protein localized to the
endoplasmic reticulum. This gain of function interaction resulted in mis-localization of
GRAMD1A in BICD2_R747C expressing cells. Collectively, our results suggest that dynein
hyperactivity, interactome changes, and cargo transport defects might contribute to the
symptoms associated with SMALED2.

INTRODUCTION

The intracellular transport of mRNAs, proteins, vesicles, and organelles is facilitated by microtubule motors. These motors operate along microtubules, which are polarized cytoskeletal filaments. Typically, kinesin family motors carry cargo toward the microtubule plus end (Yildiz, 2025), whereas transport toward the minus end relies primarily on a single motor protein, cytoplasmic dynein-1 (dynein) (Canty and Yildiz, 2020; Reck-Peterson et al., 2018). While motor-driven transport is vital across various cell types, it holds particular importance in large, polarized cells like neurons. For instance, the axon of some neurons can extend over a meter in length, making them highly reliant on efficient transport systems. Even minor disruptions in motor-based transport can impair neuronal function and lead to disease (Franker and Hoogenraad, 2013). As a result, mutations affecting motor proteins, their activators, or adaptors are linked to a variety of neurological disorders (Sleigh et al., 2019).

The isolated dynein motor is present in an inhibited conformation and has a limited capacity to traverse the microtubule cytoskeleton (Torisawa et al., 2014; Zhang et al., 2017). Processive movement by dynein requires the multi-subunit dynactin complex and an activating cargo adaptor (McKenney et al., 2014; Schlager et al., 2014; Splinter et al., 2012). Bicaudal-D (BicD), initially identified in *Drosophila*, is one of the best characterized activating cargo adaptors (Hoogenraad and Akhmanova, 2016) (Mohler and Wieschaus, 1986). Mammals encode four orthologous proteins, BICD1, BICD2, BICDR1 and BICDR2. The closest mammalian ortholog of *Drosophila* BicD is BICD2. BICD2 contains three coiled-coil regions designated CC1, CC2, and CC3 (Fig. 1A) (Olenick and Holzbaur, 2019). The first coiled-coil, CC1, is responsible for binding to dynein and dynactin (Chowdhury et al., 2015; Urnavicius et al., 2015), whereas the third coiled-coil, CC3, mediates cargo binding (Hoogenraad and Akhmanova, 2016). In the absence of cargo, an intramolecular interaction between the CC1 and CC3 regions of BICD2 results in the protein folding into an inhibited conformation (Liu et al., 2013; Terawaki et al., 2015; Wharton and Struhl, 1989). In this state, BICD2 is unable to bind dynein (Fig. 1B). Cargo binding to CC3 competes with the intramolecular interaction, effectively opening up BICD2 and enabling the cargo-bound adaptor to interact with and activate dynein for motility (Goldman et al., 2019; Huynh and Vale, 2017; Liu et al., 2013; McClintock et al., 2018; Sladewski et al., 2018). This type of regulation ensures that the motility of dynein is tightly coupled to cargo binding.

The importance of BICD2 in dynein-mediated transport is highlighted by the fact that mutations in this gene are associated with a type of spinal muscular atrophy (SMA). SMA refers to a group of disorders characterized by progressive muscle weakness and atrophy caused by loss of motor neurons. Most often, SMA is autosomal recessive, and in the majority of these cases, causative mutations are linked to the SMN (survival of motor neurons) gene (Angilletta et al., 2023). Autosomal dominant SMA, caused predominantly by mutations in BICD2, is associated with weakness and atrophy of muscles in the feet and legs (Koboldt et al., 2020; Neveling et al., 2013; Oates et al., 2013; Peeters et al., 2013). Thus, this type of SMA is referred to as Spinal Muscular Atrophy with Lower Extremity Predominance (SMALED2). In some, the disease is relatively mild, whereas in others it is more severe, presenting with contractures, hip dysplasia, brain abnormalities, cognitive impairment, and even death (Koboldt et al., 2020).

Although the link between BICD2 and SMALED2 has been recognized, we have a limited
understanding of the molecular basis of the disease. Three SMALED2 associated mutations
within the dynein binding region of BICD2 have been characterized. These mutations have been
shown to increase the interaction between BICD2 and dynein/dynactin (Huynh and Vale, 2017).
Consequently, dynein displayed hyperactivity in motility assays (Huynh and Vale, 2017).
However, numerous additional SMALED2 mutations have also been identified within the cargo
binding domain of BICD2 (Koboldt et al., 2020; Martinez-Carrera and Wirth, 2015). The
mechanism by which these mutations result in SMALED2 is unknown. Given that BICD2 is a
cargo adaptor for dynein, we hypothesized that mutations within the cargo binding domain likely
result in interactome changes. As such, cargo transport defects caused by altered BICD2
interactomes, might underlie the etiology of SMALED2. The goal of this study was to define the
interactome of wild-type and mutant alleles of BICD2 and to identify potential interactome
changes.

RESULTS

Defining the wild-type BICD2 interactome.

In order to test our hypothesis that SMALED2 associated mutations in the cargo binding domain of BICD2 are associated with interactome changes, it was first critical for us to define the interactome of wild-type BICD2. Although BICD2 has been known to function as a dynein cargo adaptor, only a few proteins have been shown to interact with BICD2 (Hoogenraad and Akhmanova, 2016; Olenick and Holzbaur, 2019). A more comprehensive interactome of BICD2 was defined by Redwine and colleagues using proximity biotin ligation (Redwine et al., 2017). However, the goal of that study was to more broadly define the dynein interactome and potential novel BICD2 interacting proteins were not validated by follow up studies.

To begin our analysis, we generated stable cell lines expressing either RFP tagged mini-TurboID (RFP-mTrbo) or full length BICD2 tagged on the C-terminus with mini-TurboID (BICD2-mTrbo). TurboID is a proximity biotin ligase that was generated by selecting mutants of BioID that displayed much higher levels of activity in comparison to the original enzyme (Branon et al., 2018). mini-TurboID is a smaller version of TurboID, the biotinylation activity of which is more tightly coupled to the addition of exogenous biotin (Branon et al., 2018). HEK293 FLP-In T-Rex cells were used for this experiment because this enabled us to integrate the constructs at the same genomic locus across all cell lines. In addition, expression of the fusion proteins was inducible upon the addition of tetracycline. This prevents any toxic effects that might arise from constitutive expression of SMALED2 alleles of BICD2.

Cells expressing either RFP-mTrbo or BICD2_WT-mTrbo were grown to scale and proximal proteins were labeled using biotin. Biotinylated proteins were purified using streptavidin magnetic beads and identified using mass spectrometry, with the entire experiment being conducted in triplicate. Proteins enriched in the BICD2_WT-mTrbo pellet at least two-fold in comparison to the RFP-mTrbo pellet and having a p value of at least 0.05 were considered potential BICD2 interacting partners (Fig. 1C, C', Supplemental table 1). This list, which consists of 66 proteins, included several dynein components and known BICD2 interacting proteins such as RANBP2, RANGAP1, and PLK1 (Fig. 1D) (Gallisa-Sune et al., 2023; Splinter et al., 2010). In addition, four components of the HOPS complex were also specifically enriched in the BICD2_WT-mTrbo pellet (Fig. 1D). The HOPS complex consists of six subunits and is involved in the fusion of late endosomes with lysosomes (van der Beek et al., 2019). Consistent with known BICD2 functions, a cellular component GO analysis indicated an enrichment of terms such as "cytoplasmic dynein complex", "nuclear pore" and "vesicle tethering complex" (Fig. 1E).

Surprisingly, we failed to identify RAB6A in our interactome analysis. RAB6A was the first cargo identified for BICD2 using a yeast-two hybrid screen (Matanis et al., 2002). Previous studies have shown that GTP bound RAB6A is the preferred binding partner for BICD2 (Matanis et al., 2002). Thus, in order to determine whether RAB6A is capable of associating with full length BICD2-mTrbo, we transfected cells expressing either RFP-mTrbo, full length BICD2-mTrbo, or a version of BICD2-mTrbo lacking the CC3 domain along with a GTP locked mutant of RAB6A (GFP-RAB6A Q72L). Consistent with published results, GFP-RAB6A Q72L specifically associated with full length BICD2-mTrbo (Supplemental Fig. 1A). Based on this, we conclude that although a GTP locked version of RAB6A is capable of interacting with BICD2-mTrbo, the

139 interaction between endogenous RAB6A and BICD2 might not be that prevalent under
140 physiological conditions.

Subunits of the HOPS complex associate with BICD2 in vivo.
Four components of the six member HOPS complex were found to associate with BICD2 in our
interactome analysis with VPS41 being the fifth most enriched protein (Fig. 1C', D,
Supplemental table 1). We therefore chose to validate these interactions using co-
immunoprecipitation. Strains expressing either RFP-mTrbo, full length BICD2-mTrbo, or the
delta CC3 domain construct (BICD2_delCC3-mTrbo) were used for this analysis. The TurboID
constructs contain a V5 tag. Thus, V5 trap beads were used to immunoprecipitate the tagged
proteins. Bound proteins were eluted and analyzed by western blotting using antibodies against
VPS41 (Fig. 2A), VPS18 (Fig. 2B) or VPS16 (Fig. 2B). Consistent with our interactome analysis,
all three HOPS components specifically co-immunoprecipitated with full length BICD2-mTrbo
(Fig. 2A, B). A similar interaction between BICD2-mTrbo and VPS41 was also detected in cells
depleted of endogenous BICD2 (Supplemental Fig. 1B). Thus, we conclude that BICD2
associates with VPS41, VPS18, and VPS16 in vivo and that these interactions require the
BICD2 cargo binding domain.

Previously identified BICD2 cargo such as RANBP2 and RAB6A have been shown to interact
with the isolated CC3 domain of BICD2 (Matanis et al., 2002; Splinter et al., 2010). In order to
determine whether the same conditions apply for the HOPS complex components, we repeated
the experiment using the same three strains mentioned above in addition to a strain expressing
the isolated CC3 domain fused to mTrbo (BICD2_CC3-mTrbo) (Fig. 2C). Unexpectedly,
although RANBP2 was able to interact with both full length BICD2 and the isolated CC3 domain,
VPS41, VPS18 and VPS16 were only capable of associating with full length BICD2 (Fig. 2D,
Supplemental Fig. 1C). Thus, the minimal cargo binding domain of BICD2 is not sufficient for
interacting with the HOPS complex components. We also attempted the binding experiment
using an extended BICD2 C-terminal construct (residues 442-824), but even this construct failed
to efficiently bind the HOPS complex components (data now shown). Finally, we generated a
BICD2 construct that lacked the first coiled coil domain (BICD2_delCC1, residues 288-824).
Although this construct was able to bind VPS41 above background levels, the binding was
reduced in comparison to the full-length protein (Supplemental Fig. 1D). RANBP2 binding was
also reduced with this deletion construct (Supplemental Fig. 1D). It is possible that this deletion
affects the normal folding of BICD2 or that it results in the protein folding into a locked
conformation that inhibits cargo binding. Collectively, our results suggest that although the CC3
domain of BICD2 is required for interaction with the HOPS complex, it is not sufficient.

Among the HOPS components, VPS41 was by far the most enriched in the BICD2 interactome
(Fig. 1C, C'). We therefore hypothesized that the interaction between BICD2 and the HOPS
complex is mediated by VPS41. Contrary to our hypothesis, however, siRNA mediated depletion
of VPS41 had little effect on the ability of BICD2 to interact with VPS16 and VPS18
(Supplemental Fig. 1E). This suggests that BICD2 is able to independently associate with
multiple HOPS components, potentially interacting with a common motif or domain present
within these proteins. Our inability to map the minimal domain of BICD2 required for interaction
with the HOPS complex components precludes our ability to test for their direct interaction.

Thus, although our results indicate that BICD2 associates with these proteins in vivo, we cannot
conclude whether the interaction is direct or indirect.

The proper localization of VPS41 and LAMP1 vesicles requires BICD2.
We next examined the localization of the HOPS complex in control versus BICD2 depleted
HeLa cells. Although antibodies are available that can detect endogenous VPS41, VPS18, and
VPS16 by western blot, these antibodies were not suitable for detecting the native protein by
immunofluorescence. Thus, we used a GFP-tagged VPS41 construct for this analysis. In HeLa
cells, microtubule minus ends are present at the perinuclear centrosome and plus ends extend
towards the cortex. As expected, given the role of the HOPS complex in late endosome-
lysosome fusion, GFP-VPS41 positive vesicles were enriched in a perinuclear area. Dispersed
vesicles and vesicles present closer to the cortex were also observed in cells transfected with a
control siRNA (Fig. 3A, D, E). As expected, depletion of Dynein heavy chain (DHC, DYNC1H1)
resulted in a reduction of perinuclear clustering and an accumulation of vesicles close to the cell
cortex (Fig. 3B, D, E, Supplemental Fig. 2A). Because BICD2 is a cargo adaptor for dynein, we
expected that depletion of BICD2 would produce a similar phenotype. Surprisingly however,
BICD2 depletion resulted in perinuclear clustering of GFP-VPS41 vesicles (Fig. 3C, D, E,
Supplemental Fig. 2A, B). A similar phenotype was also observed using a different siRNA
targeting BICD2 (Supplemental Fig. 2B, 2D). Previous studies have shown that disruption of
BICD2-mediated dynein cargo trafficking results in fragmentation of the Golgi and defective
centrosome to nucleus tethering in cells in G2 phase of the cell cycle (Hoogenraad et al., 2001;
Splinter et al., 2010). Consistent with these earlier findings, we also found that the depletion of
BICD2 using the above siRNA resulted in Golgi fragmentation and centrosome localization
defects in HeLa cells (Supplemental 2E-H).

In addition to the HOPS complex, late endosomes and lysosomes are also positive for LAMP1
and previous studies have shown partial co-localization between V5-tagged VPS41 and LAMP1
vesicles (van der Welle et al., 2021). We therefore used a validated antibody to detect LAMP1
vesicles under these same experimental conditions. Consistent with what was observed for
GFP-VPS41, LAMP1 was present in more peripheral vesicles upon DHC depletion and was
clustered closer to the nucleus in BICD2 depleted cells (Fig. 3F-H, Supplemental Fig. 2I-K).

Although we were able to visualize GFP-VPS41 for immunofluorescence experiments using a
GFP antibody to boost signal intensity, the native fluorescence of GFP-VPS41 was consistently
very low, precluding our ability to examine the motility of these vesicles in living cells. We
therefore examined lysosome motility using a SiR lysosome kit. This reagent detects cathepsin
D present in mature lysosomes. As expected, although lysosomes were present throughout the
cell, they were enriched in a perinuclear area and a large fraction of these particles were motile
(Video 1). Depletion of BICD2 caused an even more pronounced perinuclear enrichment of
lysosomes, consistent with the localization of GFP-VPS41 and LAMP1. However, in contrast to
cells treated with the control siRNA, very few motile particles were detected in BICD2 depleted
cells (Video 2; Supplemental Fig. 2L). Thus, depletion of BICD2 results in perinuclear enriched
lysosomes that are largely immotile.

The localization of GFP-VPS41 and LAMP1 vesicles in BICD2 depleted cells is similar to the
phenotype obtained upon depletion of KIF5B, a plus-end directed Kinesin-1 motor (Guardia et

al., 2016). Although BICD2 is primarily regarded as a cargo adaptor for dynein, it has also been
shown to interact with KIF5B (Grigoriev et al., 2007). In addition, *Drosophila* BicD was recently
shown to activate the motility of Kinesin-1, the fly ortholog of KIF5B (Ali et al., 2025). In order to
more closely examine the role of BICD2 in LAMP1 vesicle localization, we over-expressed
either GFP or BICD2 in cells. In contrast to cells over-expressing GFP, LAMP1 vesicles were
peripherally scattered in BICD2 over-expressing cells. This phenotype was observed in HeLa
and Cos7 cells (Fig. 3I, J, L, M and Supplemental Fig. 2M, M'). Depletion of KIF5B in BICD2
over-expressing cells reverted this phenotype and restored the perinuclear localization of
LAMP1 vesicles (Fig. 3K-M, and Supplemental Fig. 2C). These results suggest that BICD2 likely
functions to link VPS41 and LAMP1 vesicles with KIF5B for plus-end directed transport. Our
studies do not reveal the mechanism of this linkage, however, and further studies will be needed
to fully understand the role of BICD2 in this process.

SMALED2 mutations in the BICD2 cargo binding domain result in dynein hyperactivation.
Several SMALED2 associated mutations have been identified in the BICD2 cargo binding
domain (Koboldt et al., 2020). For this initial analysis, we chose to examine BICD2_R694C and
BICD2_R747C. These mutants, which result in the substitution of arginine for cystine at the
indicated residues, were particularly intriguing because the R747C mutation was predicted to
disrupt the interaction between BICD2 and RANBP2 (Terawaki et al., 2015). By contrast,
previous studies suggest that the R694C mutation results in a higher level of interaction
between BICD2 and RANBP2 (Yi et al., 2023). In addition to these mutations in the cargo
binding domain, we also chose to analyze an additional mutant, BICD2_N188T (substitution of
asparagine for threonine). This mutation is present within the first coiled coil domain and has
been shown to result in dynein hyperactivation (Huynh and Vale, 2017). No studies have thus
far addressed the global interactome of these mutants and how they might differ from the wild-
type protein.

Stable cell lines capable of expressing BICD2_N188T, R694C and R747C with a C-terminal
mTrbo tag were generated in HEK293 FLP In T-Rex cells. We first determined whether the
cargo binding domain mutants retained their ability to interact with dynein. Wild-type and mutant
versions of BICD2-mTrbo, containing a V5 tag, were immunoprecipitated using V5 trap beads.
The co-precipitated proteins were analyzed by western blotting using antibodies against
Dynactin1/p150 Glued (DCTN1) and Dynein intermediate chain (DYNC1I1). Interestingly, not
only did BICD2_R694C and BICD2_R747C retain their interaction with dynein, both mutants
generally interacted with dynein and dynactin at a higher level than the wild-type protein (Fig.
4A). Due to experimental variability in the co-immunoprecipitation experiment, the increased
rate of binding was only statistically significant for the BICD2_R747C mutant (Supplemental Fig.
3A, B). In this regard, they were similar to the BICD2_N188T mutant which had previously been
shown to interact with dynein at a higher level and to hyperactivate the motor (Fig. 4A) (Huynh
and Vale, 2017).

We next examined the localization of wild-type and mutant BICD2 in HeLa cells. Wild-type
BICD2 was localized throughout the cell with a fraction co-localizing with Golgi and centrosomal
markers (Fig. 4B and F, Supplemental Fig. 3C, C', C''). By contrast all three mutants were
enriched at varying degrees within the centrosome (Fig. 4C-F). Notably, the centrosomal

enrichment was highly significant for BICD2_R694C and BICD2_R747C (Fig. 4F). These results
suggest that the cargo binding domain mutants might hyperactivate dynein. To test this more
directly, we monitored the distribution of peroxisomes using a well-characterized tethering assay
(Kapitein et al., 2010; Passmore et al., 2021). Peroxisomes are relatively immotile but can be
transported towards microtubule minus ends in a dynein dependent manner if they are tethered
to an activating adaptor (Fig. 4G). Thus, centrosomal clustering of peroxisomes is an indicator
of dynein activity. Peroxisomes were tethered to wild-type or mutant alleles of BICD2.
Consistent with the higher level of dynein interaction, peroxisomal clustering was increased in
cells expressing all three SMALED2 mutants (Fig. 4H and Supplemental Fig.3D-G). Collectively,
these results suggest that SMALED2 mutations within the cargo binding domain of BICD2 are
also capable of hyperactivating dynein.

As noted previously, BICD2 has also been shown to interact with the microtubule plus-end
directed motor, KIF5B (Grigoriev et al., 2007). We therefore determined whether this interaction
was maintained with the mutant alleles of BICD2. Wild-type full-length BICD2 associated with
KIF5B at slightly higher levels than the binding control, consistent with published results
(Grigoriev et al., 2007). Interestingly, all three mutants displayed a reduced interaction with
KIF5B, with the BICD2_R747C mutant being the most severe (Supplemental Fig. 3H, I). Thus,
the phenotypic outcome of dynein hyperactivity likely results from the combined effect of
increased binding between BICD2 and dynein, and reduced binding between BICD2 and KIF5B.

In contrast to HeLa cells, microtubules are organized differently in neurons. In these cells,
microtubule minus ends are localized within the cell body and plus ends extend towards the
axon (van Beuningen and Hoogenraad, 2016). Thus, anterograde movement towards the axon
tip is driven by a plus end motor, whereas retrograde transport towards the cell body involves
dynein. Primary hippocampal neurons were established from E18 rat brains and after 3 days in
vitro, plasmids containing either wild-type or mutant alleles of BICD2 were transfected into these
cells. Consistent with BICD2 being a cargo adaptor for dynein, most of the signal for the wild-
type protein was detected within the cell body (Fig. 5A). However, wild-type BICD2 could also
be detected within the axon. A similar pattern was noted for BICD2_N188T (Fig. 5B). By
contrast, axonal signal for BICD2_R694C and BICD2_R747C was reduced, reflected as a
relative increase in cell body enrichment (Fig. 5C, D). The difference in localization was
statistically significant for the R747C mutant (Fig. 5E). These results are consistent with the
cargo binding domain mutants resulting in dynein hyperactivation. Huynh and Vale found that
expression of certain SMALED2 associated BICD2 mutants, including BICD2_N188T, in
hippocampal neurons results in reduced neurite growth (Huynh and Vale, 2017). We obtained
similar results and found that this phenotype was also shared by the cargo binding domain
mutations, BICD2_R694C and BICD2_R747C (Fig. 5F).

SMALED2 mutations result in altered BICD2 interactomes.
We next determined the interactomes of the three SMALED2 mutations and compared these
interactomes to that of wild-type BICD2. As before, the entire experiment was done in triplicate.
Proteins that displayed a two-fold increased or decreased interaction with the mutant allele in
comparison to the wild-type and had a p value of at least 0.05 were considered significant (Fig.
6A-C, Supplemental tables 2, 3, 4). All three mutations, including N188T, which is present in the

CC1 domain of BICD2, displayed an altered interactome. The most dramatic difference,
however, was observed for BICD2_R747C (Fig. 6C).

In order to validate the proteomics results, we repeated the experiment in triplicate and
analyzed the pellets by western blotting. Consistent with the proteomics data, and with
published results using a mutation in mouse BicD1 (Terawaki et al., 2015), BICD2_R747C
displayed a reduced interaction with RANBP2 (Fig, 6D, E, Supplemental table 4). BICD2 has
been shown to associate indirectly with RANGAP via its direct interaction with RANBP2
(Splinter et al., 2010). Thus, as expected, a reduced level of RANGAP was detected in pellets
from BICD2_R747C in comparison to the wild-type (Supplemental table 4). A further analysis of
the proteomics results indicated a reduced association of BICD2_R747C with nuclear import
receptors such as importin beta (Supplemental table 4). This result was confirmed by repeating
the binding experiment and analyzing the eluate by western blotting (Supplemental Fig. 4A). We
hypothesized that the association between BICD2 and import receptors is likely indirect and
mediated via the direct interaction between BICD2 and RANBP2. Consistent with this notion,
the interaction of wild-type BICD2 with importin beta was greatly reduced in cells depleted of
RANBP2 (Supplemental Fig. 4C). By contrast, knock-down of RANBP2 had no effect on the
BICD2-VPS41 interaction (Supplemental Fig. 4C). Based on these results, we conclude that
BICD2_R747C is disrupted for interaction with RANBP2, and consequently also displays a
reduced association with additional nuclear import factors.

An unexpected finding was that the BICD2_R747C mutant also displayed a reduced interaction
with HOPS complex components. The proteomics result indicated a reduced interaction with
VPS41, VPS18 and VPS33 (Supplemental table 4). However, the mass spectrometry result
indicated that the association of BICD2_R747C with VPS16 was unaffected. In order to validate
these results, the experiment was repeated and analyzed by western blotting. Indeed, the
association of BICD2_R747C with VPS41 and VPS18 was greatly reduced in comparison to the
wild-type protein (Fig. 6D, F and Supplemental Fig. 4C). However, using this approach we
consistently observed a significant reduction in the association between BICD2_R747C and
VPS16 (Supplemental Fig. 4D). Based on these results, we conclude that this SMALED2
mutant is most likely compromised for interacting with the entire HOPS complex.

In addition to interactions that were lost or reduced, all three mutants were also associated with
changes in the positive direction. For instance, the mass spectrometry results revealed that the
mutants showed a stronger interaction with the centrosomal protein CSPP1 compared to wild-
type BICD2 (Supplemental Tables 2, 3, 4). Validation experiments confirmed this finding,
demonstrating that the cargo-binding domain mutants had significantly elevated interactions
with CSPP1, exceeding that observed with either wild-type BICD2 or BICD2_N188T (Fig. 6D,
G). Since BICD2_R694C and BICD2_R747C are enriched at the centrosome (Fig. 4D-F), their
altered localization most likely accounts for their increased association with centrosomal
proteins.

BICD2_R747C exhibits a gain of function interaction with GRAMD1A.
Among the three mutants analyzed, BICD2_R747C exhibited the most distinct interactome. One
particularly notable interaction was with GRAMD1A, a protein absent from the wild-type BICD2
interactome but highly enriched in the BICD2_R747C pull-down, indicating a gain-of-function

interaction. GRAMD1 encodes three isoforms: A, B, and C, with GRAMD1A being highly
expressed in the central nervous system. To validate this observation, the experiment was
repeated in triplicate and analyzed by western blotting. Consistent with the mass spectrometry
result, GRAMD1A showed only background binding to the control, wild-type BICD2,
BICD2_N188T, or BICD2_R694C. In contrast, GRAMD1A strongly associates with
BICD2_R747C (Fig. 7A, B). The same binding profile was observed in cells depleted of
endogenous BICD2 (Supplemental Fig. 5A).

We next examined the localization of GRAMD1A in Cos7 cells co-expressing GRAMD1A-
mScarlet3 with wild-type or mutant alleles of BICD2 tagged with mNeon. GRAMD1A localizes to
the endoplasmic reticulum (ER) and concentrates at sites of ER-plasma membrane contact
(Besprozvannaya et al., 2018). Cos7 cells are typically used for examining the localization of
GRAMD1A due to their large and flattened appearance (Besprozvannaya et al., 2018; Naito et
al., 2019). As expected, GRAMD1A displayed a typical ER localization pattern in cells
expressing wild-type BICD2 or BicD2_N188T (Fig. 7C, D). Much like what was observed in
HeLa cells, BICD2_R694C was often enriched in the centrosomal region in Cos7 cells (Fig. 7E,
Fig. 4D). The localization of GRAMD1A was essentially unchanged in these cells, although a
small amount of the protein co-localized with BICD2_R694C in the centrosomal region (Fig. 7E,
G). By contrast, consistent with a gain-of-function interaction, a significant fraction of GRAMD1A
localized adjacent to Pericentrin, a centrosomal marker, in cell expressing BICD2_R747C (Fig.
7F, G, Supplemental Fig. 5B-F). This effect was specific for GRAMD1A because a general ER
maker remained correctly localized in cells expressing BICD2_R747C (Supplemental Fig. 5G,
G', H, H').

Collectively, our results suggest that SMALED2 associated mutations in BICD2 hyperactive
dynein and result in gain of function and loss of function interactions with cargo. It is therefore
possible that cargo trafficking defects that arise because of these phenotypes contribute to the
symptoms associated with SMALED2.

DISCUSSION

Mutations in the dynein cargo adaptor BICD2 have been linked to spinal muscular atrophy with lower extremity predominance (SMALED2) (Koboldt et al., 2020). Mutations in the heavy chain of the dynein motor have also been implicated in a version of this disorder (Chan et al., 2018; Das et al., 2018), suggesting that defects in dynein-mediated transport contribute to its etiology. However, the molecular and cellular mechanisms underlying SMALED2 pathogenesis remain poorly understood. Previous studies have characterized mutations within the first coiled-coil domain of BICD2, a region responsible for interactions with dynein and dynactin. These analyses elegantly demonstrated that mutants such as BICD2_N188T result in dynein hyperactivity (Huynh and Vale, 2017). In addition to these mutants, however, recent studies have identified several SMALED2-associated alleles within the C-terminal cargo-binding domain of BICD2 (Ravenscroft et al., 2016; Synofzik et al., 2014). Given BICD2's role as a dynein cargo adaptor, these findings raise two important questions: (1) Is dynein hyperactivity a common feature of SMALED2-associated BICD2 mutations? and (2) Do these mutations alter the interactome of BICD2 relative to the wild-type protein? The goal of this study was to address these questions and elucidate potential molecular consequences of SMALED2 associated BICD2 mutations.

BICD2 is one of the best characterized dynein cargo adaptors. However, most studies involving BICD2 have focused on the mechanism by which this adaptor activates dynein for processive motility. Relatively little is known regarding the cargo that is linked to dynein by BICD2. In *Drosophila*, BicD links the RNA binding protein Egalitarian (Egl) with dynein for transport of specific mRNAs in the oocyte and embryo (Dienstbier et al., 2009; Goldman et al., 2021; Goldman et al., 2019; Mach and Lehmann, 1997; McClintock et al., 2018). Loss of either BicD or Egl compromises transport of these mRNAs and consequently results in defective oogenesis or embryogenesis. The first definitive cargo identified for mammalian BICD2 was the small GTP binding protein, RAB6A (Matanis et al., 2002). Despite the ability of BICD2 to directly bind RAB6A, most vesicles containing RAB6A move towards the plus end of microtubules, suggesting that their transport is primary driven by the Kinesin-1 motor, KIF5B (Grigoriev et al., 2007). Other cargos that have been shown to directly bind BICD2 are RANBP2, a nucleoporin, and Nesprin-2 (SYNE2), a LINC complex component involved in linking dynein and kinesin to the nuclear envelope (Goncalves et al., 2020; Splinter et al., 2010).

In order to determine whether SMALED2 alleles of BICD2 are associated with interactome changes, it was therefore critical for us to determine the interactome of wild-type BICD2. This was done using the promiscuous biotin ligase miniTurboID (mTrbo). In comparison to an RFP-mTrbo control, BICD2-mTrbo resulted in the biotinylation and purification of numerous known interacting partners including RANBP2, as well as several components of the dynein motor. One interesting group of potentially novel interacting proteins were components of the HOPS complex, a six-subunit complex of proteins involved in endocytic trafficking (Spang, 2016). Four of the six HOPS components were identified in the wild-type BICD2 interactome, with VPS41 being the fifth most enriched protein. However, unlike RANBP2, RAB6A, and NESPRIN-2, all of which are able to bind the isolated BICD2 cargo binding domain (Goncalves et al., 2020; Matanis et al., 2002; Splinter et al., 2010), the HOPS complex components were only able to bind full-length BICD2. The BICD2 cargo binding domain was therefore necessary but not

sufficient for interaction with HOPS components. In addition, contrary to our initial hypothesis
that VPS41 was the direct binding partner between BICD2 and the HOPS complex, BICD2
retained its interaction with VPS16 and VPS18 in cells depleted of VPS41. This suggests that
BICD2 likely recognizes a domain or motif present in several HOPS proteins. We attempted to
use Alphafold2 multimer to predict the relevant domain within HOPS proteins that interact with
BICD2. Although Alphafold2 was able to generate a high confidence prediction of the interaction
site between BICD2 and RAB6A, consistent with published results (Zhao et al., 2024), it failed to
produce a high confidence prediction for the BICD2-HOPS complex interaction (data not
shown). Thus, although we were able to validate the in vivo association between BICD2 and
VPS41, VPS16 and VPS18, we are not able to conclude whether BICD2 is capable of directly
interacting with these proteins. To the best of our knowledge, this is the first example of a BICD2
interacting proteins that display this binding characteristic. The ScaC protein from the
intracellular pathogen *O. tsutsugamushi* was recently also shown to interact with BICD2 and
although the binding site of ScaC was different from that used by RANBP2 or RAB6A, it was still
able to interact with the isolated cargo binding domain of BICD2 (Manigrasso et al., 2025).

Another unusual aspect of the BICD2-HOPS complex interaction is that it does not appear to be
linked to dynein-mediated trafficking. Depletion of dynein heavy chain resulted in the peripheral
distribution of GFP-VPS41 and LAMP1 vesicles, indicative of a reduction in minus end
transport, and a net gain in plus end directed transport. By contrast, depletion of BICD2 resulted
in the perinuclear accumulation of lysosomal vesicles that were mostly immotile. Interestingly,
however, over-expression of BICD2 caused the outward spreading of LAMP1 vesicles, a
process that depends on KIF5B (Guardia et al., 2016). Previous studies have shown that BICD2
is also able to interact with KIF5B via a central coiled coil domain (Grigoriev et al., 2007;
Hoogenraad and Akhmanova, 2016). A recent report suggests that *Drosophila* BicD is capable
of interacting with and activating the motility of Kinesin-1, the fly homolog of KIF5B (Ali et al.,
2025). Consistent with the notion that BICD2 might link late endosomal vesicles with KIF5B,
depletion of KIF5B in BICD2 over-expressing cells restored the normal localization of LAMP1
vesicles. Additional studies will be required to determine whether BICD2 is capable of directly
interacting with these vesicles and whether these vesicles are directly linked to KIF5B by
BICD2.

The motility of LAMP1 vesicles has some similarity to the transport of RAB6A exocytic vesicles.
RAB6A vesicles are transported from the area of the Golgi towards the cell periphery in a KIF5B
dependent manner and loss of either kinesin-1 or dynein results in a sharp reduction in the
number of motile particles (Grigoriev et al., 2007). In addition, mutations in BICD2 that
compromise binding to RAB6A also result in vesicles that are largely immotile (Zhao et al.,
2024). Thus, in the case of LAMP1 and RAB6A vesicles, instead of resulting in an increased
rate of minus end transport, loss of BICD2 results is compromised vesicle motility, indicating that
coordination between opposite polarity motors is critical for their motility.

As noted earlier, mutations in the CC1 region of BICD2 hyperactivate dynein (Huynh and Vale,
2017). Our findings indicate that this property is also shared by BICD2_R694C and
BICD2_R747C, mutations present within the C-terminal cargo binding domain. In the absence
of cargo, BICD2 is thought to exist in an inhibited conformation due to intramolecular

interactions between the N and C termini of the protein (Fig. 1B) (Liu et al., 2013; Terawaki et
al., 2015; Wharton and Struhl, 1989). Cargo binding to the C-terminus of BICD2 counteracts the
intramolecular interaction, enabling N-terminal residues within BICD2 to bind the
dynein/dynactin complex (Goldman et al., 2019; Huynh and Vale, 2017; Liu et al., 2013;
McClintock et al., 2018; Sladewski et al., 2018). How might mutations in BICD2 result in dynein
hyperactivation? One possibility is that these mutations disrupt the autoinhibited state of BICD2,
effectively causing BICD2 to be present in a more open and uninhibited conformation that
promotes dynein/dynactin binding. Molecular dynamics simulations suggest that the R747C
substitution causes a registry shift in the coiled coil, likely destabilizing this domain and thus
disrupting the intramolecular interaction between the N and C termini of BICD2 (Cui et al.,
2020). Another possibility is that the hyperactivation of dynein results for reduced binding
between BICD2 and KIF5B. Our results are consistent with this scenario and suggest that the
net effect of dynein hyperactivity results for three molecular changes; reduced intramolecular
BICD2 interaction, increased interaction between BICD2 and dynein, and reduced interaction
between BICD2 and KIF5B.

In addition to hyperactivating dynein, all three mutations, including BICD2_N188T, alter the
BICD2 interactome. This finding was unexpected for BICD2_N188T because this mutation is not
within the cargo binding domain. One possible explanation for this phenotype is that
BICD2_N188T is present in a more open conformation, and this change affects its binding
properties. Another possibility that is not mutually exclusive is that the different binding profile
results from the altered localization of BICD2_N188T within the cell. In comparison to wild-type
BICD2, we generally observed greater centrosomal enrichment of BICD2_N188T. In comparing
the three mutants, the general trend was that more proteins displayed a reduced interaction with
the SMALED2 mutants in comparison to wild-type BICD2. Among the three mutants analyzed,
BICD2_R747C displayed the most drastically altered interactome. This mutant displayed
reduced association with RANBP2, importin beta, and HOPS complex components.
Interestingly, this mutant also displayed numerous gain-of-function interactions. For instance,
although minimal binding was observed between wild-type BICD2 and GRAMD1A, this protein
abundantly interacted with BICD2_R747C. GRAMD1A is involved in non-vesicular transport of
accessible cholesterol from the plasma membrane to the ER and is often concentrated at sites
of plasma membrane-ER contact (Besprozvannaya et al., 2018; Sandhu et al., 2018). However,
in cells expressing BICD2_R747C, this localization pattern was disrupted and GRAMD1A co-
localized with BICD2_R747C adjacent to the centrosome.

The GRAMD1 family consists of three isoforms: GRAMD1A, GRAMD1B, and GRAMD1C.
Interestingly our interactome analysis only identified GRAMD1A as a gain-of-function interaction
partner with BICD2_R747C. It is unclear whether GRAMD1B and GRAMD1C also interact with
BICD2_R747C. However, given that GRAMD1 proteins can form hetero oligomers (Naito et al.,
2019), the BICD2_R747C-induced mislocalization of GRAMD1A could potentially affect the
distribution of other GRAMD1 isoforms as well. The GRAMD1 proteins function to sense excess
accessible cholesterol in the plasma membrane and to mediate the transport of this cholesterol
to the ER. This reduces the rate of new cholesterol synthesis by the ER, enabling the cell to
maintain cholesterol homeostasis (Sandhu et al., 2018). It will be interesting to determine
whether endogenous GRAMD1A is mislocalized in motor neurons of SMALED2 patients with

the BICD2_R747C mutation, and if this results in an expanded accessible pool of cholesterol at
the plasma membrane.

A recent study by Yi and colleagues examined the effect of the BICD2_R694C mutation on
cargo binding (Yi et al., 2023). Using in vitro experiments, they found that this mutation
enhanced RANBP2 binding while having no effect on NESPRIN-2 binding (Yi et al., 2023). Our
results using full-length BICD2 are consistent with this finding. We also observed slightly higher
binding of BICD2_R694C to RANBP2. However, due to experimental variability, the increase
was not statistically significant. The authors also examined cargo binding using a BICD2 double
mutant (F743I/R747C). Consistent with our results, this mutant displayed greatly reduced
binding to RANBP2, but bound NESPRIN-2 at a much higher level than the wild-type protein (Yi
et al., 2023). NESPRIN-2 was not identified as an interacting partner in our study for the wild-
type protein or the BICD2_R747C mutant, possibly due to its low expression level in HEK293
cells. Nevertheless, these findings, along with our interactome analysis, indicate that mutations
in the cargo binding domain of BICD2 can result in loss and gain-of-function interactions.

In conclusion, our study is the first to comprehensively examine the interactome of wild-type
BICD2 and to identify changes that occur in SMALED2 linked mutant alleles of BICD2. We find
that not only are mutations within the cargo binding domain associated with interactome
changes, but that these mutations are additionally capable of hyperactivating dynein. Some
limitations of this study are worth noting. In the current study, we chose to determine the BICD2
interactome in HEK FLP-In cells (embryonic kidney cells). These cells were chosen because it
enabled us to precisely integrate wild-type and mutant alleles of BICD2 at a specific locus. It
also enabled us to expand cultures of these cells to levels that were sufficient for proteomic
analysis. However, the main cell type affected in patients with SMALED2 are motor neurons.
Primary motor neurons are harder to culture to scale and to genetically manipulate to express
the desired wild-type or mutant BICD2 transgenes. Thus, although motor neurons were not
used in our study, the next significant challenge will be to perform these types of experiments
using motor neurons. In addition, although our study identified interactome changes between
wild-type and mutant alleles of BICD2, we cannot conclude whether these changes are
causative for the symptoms associated with SMALED2. Patients diagnosed with this disorder
display a range of symptoms, from relatively mild to more severe (Frasquet et al., 2020; Koboldt
et al., 2020). Even patients with the same genetic mutation can display a range of phenotypes
(Storbeck et al., 2017). Furthermore, disease symptoms can result from one or two interactome
changes that are critical for the health of motor neurons. Alternatively, symptoms might also be
caused from many small changes in the interactome that cumulatively affect the health of motor
neurons. Lastly, because SMALED2 is an autosomal dominant disorder, patients express wild-
type and mutant version of BICD2 in the same cell. Thus, to accurately model this disorder
studies will need to be conducted in motor neurons that are genetically edited to express
disease associated mutations in a heterozygous state.

ACKNOWLEDGEMENTS
We are grateful to Drs. Erika Holzbaur, Adam Fenton, Juan Bonifacino, Addanki Tirumala, and Raffaella de Pace for advice on transfection and culturing of hippocampal neurons. We are also grateful to Pritha Bagchi for her assistance with the mass-spectrometry analysis. This work was supported by grants from the National Institutes of Health to XF (NEI, EY032488) and G.B.G (NIGMS, R35GM145340). This work was supported in part by the Emory Integrated Proteomics Core (RRID:SCR_023530) and the Augusta University Cell Imaging Core (RRID:SCR_026799).

MATERIALS AND METHODS

Reagents used
The full list of DNA constructs, antibodies and reagents used in this work is found in Supplemental table 5.

DNA constructs
Synthesized DNA fragments were generated by either Twist biosciences or Genewiz/Azenta. All final plasmids used in this work were sequenced by Plasmidsaurus. Plasmid sequences as well as detailed cloning strategies will be provided upon request.

DNA encoding wild-type BICD2, generated by gene synthesis, was cloned into the Kpn1 and Xho1 sites of the pCDNA-FRT-TO vector (Thermo Fisher Scientific). Next, a fragment encoding mTrbo was cloned downstream of BICD2_wild-type using High fidelity assembly (NEB). A similar strategy using a synthesized fragment was used to clone mRFP-mTrbo into the pCDNA-FRT-TO vector. BICD2 mutants fused to mTrbo were generated by swapping in a synthesized fragment containing the desired mutation into the appropriate site of the BICD2_wild-type-mTrbo construct. The cDNA for wild-type BICD2 as well as the SMALED2 mutants contained silent mutation that made them resistant to bicd2 siRNA-1. The BICD2 truncations were produced by cloning either PCR generated, or gene synthesized fragments into the appropriate vector. Stable cell lines were created using these FRT constructs and the pOG44 vector (Thermo Fisher Scientific). For localization studies, PCR sub-cloning was used to move the appropriate wild-type or BICD2 mutans into the pLVX-Neo vector (Takara Bio). A fragment encoding mNeon-green along with the V5 epitope tag was cloned downstream of wild-type or mutant BICD2. The GFP-VPS41 construct was generated by cloning a synthesized fragment encoding human VPS41 into the Kpn1 and Not1 sites of the GFP-BICD2 plasmid (Addgene plasmid #161626; (Bonet-Ponce et al., 2020)). This removed the BICD2 cDNA from this vector and inserted the cDNA for VPS41 in its place. For the dynein activity experiment described in Fig. 4, a construct containing the PEX3 peroxisome targeting sequenced fused to mGreenLantern and the FRB dimerization sequence was cloned into the pLVX neo vector (Takara). The BICD2_wild type and mutant construct described above were modified by replacing the mTrbo sequence with the FKBP dimerization domain. The GRAMD1A-mScarlet3 construct was generated by cloning fragments corresponding to human GRAMD1A and mScarlet3 into the pLVX neo vector.

Cell culture
HeLa and Cos7 cells were obtained from ATCC and were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin (Thermo Fisher Scientific). The HEK Flp-In T-REx 293 Cell Line was obtained from Thermo Fisher Scientific and was cultured according to the instructions provided by the manufacturer. Stable cells were generated by culturing the transfected HEK Flp-In T-REx 293 in 100ug/ml Hygromycin B (Gibco) for 12 to 14 days, with fresh media changes every 3 days. The E18 rat hippocampus culturing kit was obtained from Transnetyx. Primary neurons were prepared according to the instructions

598 provided by the manufacturer. 40,000 cells were seeded onto poly Lysine coated
599 coverslips (Neuvitro Corporation) in each well of a 12-well plate. The cells were
600 transfected after 3 to 4 days in culture using Lipofectomine 2000 (Thermo Fisher
601 scientific).
602
Stable HEK Flp-In T-REx 293 expressing mGreenLantern tagged peroxisomes were
generated by infection with lentivirus. Lentiviral particles were produced in HEK293T
cells following a previously published protocol (Wei et al., 2024). Culture medium
containing the viral particles were passed through a 0.2 micron filter to remove cell
debris. Next, the HEK Flp-In T-REx 293 cells were infected with freshly prepared viral
particles for 48 hours. After infection, puromycin (1ug/mL) was added to the culture
medium to select for stable cell lines. Uniformly expressing mGreenLantern clones were
selected and used to integrate BICD2_wt or mutant containing the FKBP dimerization
motif at the FRT locus. Stables cells containing the integrated BICD2 constructs were
selected using Hygromycin B as described above.
613
DNA and siRNA transfections
The Qiagen Effectene reagent was used for transfecting DNA into HeLa, HEK Flp-In T-
REx 293, and Cos7 cells using the directions provided by the manufacturer. For
transfecting cells in 6 well dishes, 0.4ug of DNA was used along with 10ul of Effectene.
Primary neurons grown on glass coverslips in a 12 well dish were transfected using
Lipofectamine 2000 (Thermo Fisher Scientific).   0.5-0.8ug of DNA and 2ul of
Lipofectamine 2000 was used in each transfection. Expression of the transgenes was
induced using 0.3 ug/ml of Doxycycline 24 hours after transfection. The cells were fixed
and processed for immunofluorescence the following day. Lipofectamine RNAimax
(Thermo Fisher Scientific) was used for the transfection of siRNA according to the
instructions provided by the manufacturer.
625
Purification and analysis of biotinylated proteins
For small scale experiments (Figure 2, 6, 7, Supplemental figures 1 and 4), HEK Flp-In
T-REx 293 cells expressing the desired constructs by Tet-based induction (1ug/ml, 24
hours), were incubated with biotin (500uM) for 40 minutes. The cells were then
harvested, and lysates were prepared by resuspending the cells in RIPA buffer (50 mM
Tris-Cl [pH 7.5], 150 mM NaCl, 1% NP-40, 1 mM EDTA) containing a Halt Protease
inhibitor cocktail (Thermo Fisher Scientific). 1mg of total protein was used in the binding
experiment with 15ul of High-Capacity Streptavidin Agarose beads (Thermo Fisher
Scientific) in RIPA buffer. The binding was performed overnight at 4^0C. The samples
were washed four times using RIPA buffer, bound proteins were eluted in Laemmli
buffer and analyzed by western blotting. All western blot images were acquired on a Bio
Rad ChemiDoc MP.
638
For proteomics experiments, the same cells were grown in 10cm dishes and 5mg of
total protein was used in each purification. Biotionylated proteins were purified using
70ul of Streptavidin magnetic beads (Thermo Fisher Scientific), incubated at 4^0C
overnight. The following day, the samples were extensively washed using 1ml of the
following: 3 times with RIPA buffer, 3 times with high salt RIPA buffer (50 mM Tris-Cl

[pH 7.5], 1M NaCl, 1% NP-40, 1 mM EDTA), 3 times with RIPA buffer and 4 times with
PBS. The entire experiment was done in triplicate. After the final wash, the beads were
resuspended in 70ul of PBS and shipped to the Emory Integrated Proteomics Core on
dry ice.
648
649 **Mass spectrometry**
650 The mass spectrometry was performed at the Emory Integrated Proteomics Core
651 (RRID:SCR_023530).
652
On bead digestion: A published protocol was followed for on bead digestion of proteins
(Soucek et al., 2016). A digestion buffer containing 50 mM NH_4HCO_3 was added to the
beads. The mixture was then incubated with 1 mM dithiothreitol (DTT) at room
temperature for 30 minutes, followed by the addition of 5 mM iodoacetimide (IAA). This
mixture was incubated at room temperature for an additional 30 minutes in the dark.
Proteins were digested with 1 µg of lysyl endopeptidase (Wako) at room temperature
overnight and were further digested overnight at room temperature with 1 µg trypsin
(Promega). The resulting peptides were desalted using an HLB column (Waters) and
were dried under vacuum.
LC-MS/MS: The data acquisition by LC-MS/MS was adapted from a published
procedure (Seyfried et al., 2017). Derived peptides were resuspended in the loading
buffer (0.1% trifluoroacetic acid, TFA) and were separated on a Water's Charged
Surface Hybrid (CSH) column (150 µm internal diameter (ID) x 15 cm; particle size: 1.7
µm). The samples were run on an EVOSEP liquid chromatography system using the 15
samples per day preset gradient and were monitored on a Q-Exactive Plus Hybrid
Quadrupole-Orbitrap Mass Spectrometer (ThermoFisher Scientific). The mass
spectrometer cycle was programmed to collect one full MS scan followed by 20 data
dependent MS/MS scans. The MS scans (400-1600 m/z range, 3×10^6 AGC target, 100
ms maximum ion time) were collected at a resolution of 70,000 at m/z 200 in profile
mode. The HCD MS/MS spectra (1.6 m/z isolation width, 28% collision energy, 1×10^5
AGC target, 100 ms maximum ion time) were acquired at a resolution of 17,500 at m/z
200. Dynamic exclusion was set to exclude previously sequenced precursor ions for 30
seconds. Precursor ions with +1, and +7, +8 or higher charge states were excluded
from sequencing.
MaxQuant: Label-free quantification analysis was adapted from a published procedure
(Seyfried, Dammer et al. 2017). Spectra were searched using the search engine
Andromeda, integrated into MaxQuant, against 2022 human UniProtKB/Swiss-Prot
database (20,387 target sequences). Methionine oxidation (+15.9949 Da), asparagine
and glutamine deamidation (+0.9840 Da), and protein N-terminal acetylation (+42.0106
Da) were variable modifications (up to 5 allowed per peptide); cysteine was assigned as
a fixed carbamidomethyl modification (+57.0215 Da). Only fully tryptic peptides were
considered with up to 2 missed cleavages in the database search. A precursor mass
tolerance of ±20 ppm was applied prior to mass accuracy calibration and ±4.5 ppm after
internal MaxQuant calibration. Other search settings included a maximum peptide mass
of 6,000 Da, a minimum peptide length of 6 residues, 0.05 Da tolerance for orbitrap and

0.6 Da tolerance for ion trap MS/MS scans. The false discovery rate (FDR) for peptide
spectral matches, proteins, and site decoy fraction were all set to 1 percent.

Quantification settings were as follows: re-quantify with a second peak finding attempt
after protein identification has completed; match MS1 peaks between runs; a 0.7 min
retention time match window was used after an alignment function was found with a 20-
minute RT search space. Quantitation of proteins was performed using summed peptide
intensities given by MaxQuant. The quantitation method only considered razor plus
unique peptides for protein level quantitation.
696
Co-Immunoprecipitation
HEK293 FLP-in T-Rex cells expressing the desired constructs by Tet-based induction
(1ug/ml, 24 hours), were harvested and lysates were prepared using RIPA buffer. 1mg
of total protein was used in the binding experiment with 15ul of V5-Trap agarose beads
(ChromoTek, ProteinTech) in RIPA buffer. The binding was performed at 4^0C for 2
hours. The samples were then washed four times using RIPA buffer, bound proteins
were eluted in Laemmli buffer and analyzed by western blotting using the indicated
antibodies.
705
Immunofluorescence
Cells (HeLa, HEK293 FLP In T-Rex, Cos7, and primary rat neurons) adhered to glass
coverslips were fixed using either 4% formaldehyde for 5 minutes at room temperature
(Figures 5, 7, Supplemental figure 3, 4) or with methanol at -20^0C for 10 minutes
(Figures 3, 4, Supplemental figure 2). After fixation, cells were permeabilized by
incubating with PBST (PBS containing 0.1% TritonX-100) for 5 minutes. Next the
samples were blocked for 1 hour at room temperature using 5% normal goat serum
(Thermo Fisher Scientific). Cells were incubated overnight at 4^0C with primary antibody
in blocking solution. The next day, the coverslips were washed three times with PBS.
The cells were then incubated with secondary antibody diluted in blocking solution for 1
hour at room temperature. Following this, the cells were washed three times with PBS,
stained with DAPI (Thermo Fisher Scientific), and mounted onto slides using Prolong
Diamond antifade reagent (Thermo Fisher Scientific).
719
Microscopy
All imaging experiments were performed at the Augusta University Cell Imaging Core
(RRID:SCR_026799). Fixed images were captured on either an inverted Leica Stellaris
confocal microscope or an inverted Zeiss LSM780 equipped with Airyscan. Live imaging
of SiR labeled lysosomes was performed on an inverted Nikon AXR confocal
microscope equipped with the NSPARC detector. Images were processed for
presentation using Fiji, Adobe Photoshop, and Adobe Illustrator.
727
Quantifications
The clustering phenotype of GFP-VPS41 and LAMP1 vesicles (Fig. 3) was quantified
using Imaris 10.2. The nucleus was defined using the "surface" feature of Imaris and the
vesicles were defined using the "spots" feature. The percentage of vesicles that were
less than or equal to 10 microns from the nucleus and the percentage of vesicles that
were present a distance greater than 10 microns from the nucleus was determined

using object to object classification and the filtering feature in Imaris. The quantification
of lysosome motility in live cells (Supplemental figure 2D) was also performed in Imaris.
Vesicles were defined using the "spots" feature and were tracked over time. Motile
vesicles were defined as those that displayed movement over at least a 3 second
interval and had at least 0.1microns of displacement over the course of the imaging
experiment. This enabled us to quantify the percent of motile particles per cell and the
velocity of these vesicles. The centrosome enrichment phenotype shown in Fig. 4B-E
was quantified by determining the ratio of the mean BICD2 signal in the entire cell vs
the mean intensity in the centrosomal area demarcated by Pericentrin. This was done
using Imaris. The peroxisome clustering phenotype (Fig. 4H) was quantified using the
"cells" module of Imaris. Within this module, the nucleus was defined using the DAPI
channel and the cell boundaries were determined using phalloidin staining of F-actin.
Next, the peroxisomes were defined using the "spots" feature built into this module. This
enabled us to calculate the average distance of the peroxisomes to the nucleus on a
cell-by-cell basis. The cell body enrichment of BICD2 signal (Fig. 5E) was determined
using the "filaments" module in Imaris. This was used to define the cell body and the
axon. The mean intensity of signal in the cell body was compared to the mean intensity
of signal in the axon. The co-localization between GRAMD1A and BICD2 (Fig.7) was
determined using the "co-localization" module in Imaris. Axon length (Fig. 5F) was
quantified using the Simple neurite tracer plugin for FIJI. Densitometry for western blot
images captured on the BioRad Chemidoc MP were analyzed using the BioRad Image
Lab software. The binding level was compared to the amount of the respective protein
detected in the BICD2_wt lane. Graphing of data and statistical analysis was performed
using Graphpad Prism10.
758

 FIGURE LEGENDS
Figure 1: Wild-type BICD2 interactome
(A). Schematic of human BICD2, indicating the three coiled-coil domains and sites of interaction
with dynein/dynactin, kinesin-1, and cargo.

(B). Model of BICD2 binding to dynein/dynactin in a cargo dependent manner.

(C, C'). A volcano plot indicating proteins that were enriched with BICD2-mTrbo in comparison to
RFP-mTrbo. The dashed line along the x-axis indicates a fold enrichment of 2, whereas the
dashed line along the y-axis indicates a p value of 0.05. Specific interacting proteins were
considered those that were enriched at least two-fold with BICD2-mTrbo and had a p value of at
least 0.05. These candidates are shown in the zoomed in image in C'.

(D). Known BICD2 interacting proteins, components of the dynein motor, and components of the
HOPS complex that were specifically enriched with BICD2-mTrbo are indicated.

(E). A cellular component GO analysis of the BICD2 interactome.

Figure 2: BICD2 interacts with components of the HOPS complex in vivo.
(A). A co-immunoprecipitation experiment was performed with HEK cells expressing the
indicated constructs. The lysates were incubated with V5 trap beads to precipitate the tagged
proteins. The co-precipitating proteins were analyzed using western blotting with the indicated
antibodies. VPS41 specifically co-precipitates with BICD2_wt. Minimal binding was observed
with RFP-mTrbo or a BICD2 construct lacking the cargo binding domain.

(B). A similar co-precipitation experiment was set up as in panel A. The co-precipitating proteins
were analyzed by blotting using antibodies against VPS16, VPS18 and the V5 epitope. VPS16
and VPS18 co-precipitate specifically with BICD2_wt-mTrbo.

(C). A schematic of the BICD2 constructs used in the binding experiment.

(D). Cells expressing either RFP-mTrbo, BICD2_delCC3-mTrbo (lacking the cargo binding
domain), BICD2_wt-mTrbo (full length BICD2) or BICD2_CC3-mTrbo (just the cargo binding
domain) were used to examine the interaction with VPS41. Lysates were incubated with
streptavidin beads to precipitate biotinylated proteins. The precipitated proteins were analyzed
by blotting using the indicated antibodies. Although RANBP2 interacts with the isolated cargo
binding domain of BICD2, VPS41 does not.

Figure 3: Role of BICD2 in localization of GFP-VPS41 and LAMP1 vesicles.
(A-C). HeLa cells were transfected with either a control-siRNA (A), an siRNA targeting dynein
heavy chain (B), or an siRNA targeting BICD2 (C). Two days after the siRNA transfection, the
cells were transfected with a plasmid encoding GFP-VPS41. The next day, the cells were fixed
and processed for immunofluorescence using an antibody against GFP. The cells were
counterstained with DAPI (cyan) and Phalloidin (grey). Depletion of DHC results in an outward
spreading of GFP-VPS41 vesicles, whereas depletion of BICD2 results in more perinuclear
clustered vesicles.

(D, E). The distance of GPF-VPS41 vesicles relative to the nucleus was determined and plotted.
Vesicles present within 10 microns of the nucleus are shown in D, and those present at a
distance greater than 10 microns are shown in panel E.

(F-H). HeLa cells were transfected with the indicated siRNAs. Three days later, the cells were
fixed and processed for immunofluorescence using an antibody against LAMP1. As with GFP-
VPS41 vesicles, depletion of DHC resulted in peripheral vesicles, whereas depletion of BICD2
resulted in perinuclear clustering of LAMP1 vesicles.

(I-K). HeLa cells were transfected with either a control siRNA (I, J) or an siRNA targeting KIF5B
(K). Two days later the cells were transfected with a plasmid encoding either GFP (I) or BICD2-
mNeon (J, K). The cells were fixed on day 4 and processed for immunofluorescence using an
antibody against LAMP1. The cells were counterstained with DAPI. Over-expression of BICD2
results in the peripheral spreading of LAMP1 vesicles. This phenotype was reversed upon
knocking down KIF5B.

(L-M). The distance of LAMP1 vesicles relative to the nucleus was determined and plotted.
Vesicles present within 10 microns of the nucleus are shown in L, and those present at a
distance greater than 10 microns are shown in panel M.

The signal for GFP-VPS41 and LAMP1 is displayed using the "red hot" LUT in FIJI. The scale
bar is 20 microns. A one-way ANOVA was used for the quantifications shown in panel D, E, L
and M with the values compared to the mean of BICD2_wt. ns = not significant, *, p ≤ 0.05, ***,
p<0.001.

Figure 4: BICD2 cargo binding domain mutants hyperactivate dynein.
(A). A co-immunoprecipitation experiment was performed using HEK cells expressing the
indicated constructs. The tagged proteins were purified using V5 trap beads and the co-
precipitating proteins were analyzed by western blotting with the indicated antibodies. A greater
amount of DIC and DCTN1 co-purified with mutant BICD2 compared to the wild-type protein.

(B-E). HeLa cells expressing BICD2_wt (B), BICD2_N188T (C), BICD2_R694C (D) or
BICD2_R747C (E) were fixed and processed for immunofluorescence using antibodies against
V5 (cyan) and pericentrin (magenta). Merged images are also shown. All three mutants
displayed a centrosomal localization pattern to varying degrees. The scale bar is 20 microns.

(F). The centrosomal enrichment of BICD2_wt or mutant was quantified.

(G). Schematic of the peroxisome tethering assay.

(H). The average distance of peroxisomes to the nucleus was calculated on a cell-by-cell basis.
In comparison to wild-type BICD2, all three mutants showed increased clustering of
peroxisomes close to the nucleus. A one-way ANOVA was used for the quantifications shown in
panels F and H with the values compared to the mean of BICD2_wt. n = not significant, **,
p<0.01, ***, p<0.001, ****, p<0001.

Figure 5: Localization of BICD2 wild-type and mutants in neurons.

(A-D). E18 rat hippocampal neurons were transfected with the indicated constructs. Two days
after transfection, the cells were fixed and processed for immunofluorescence using a V5
antibody. The axon outline for cells expressing BICD2_R694C and BICD2_R747C are indicated.
Signal for wild-type BICD2 could be detected in the cell body and axon. A similar phenotype was
noted for BICD2_N188T. By contrast, BICD2_R694C and BICD2_R747C displayed reduced
axonal signal. The scale bar is 100 microns. The signal for BICD2 is displayed using the "red
hot" LUT in FIJI.

(E). Quantification of the cell body enrichment of BICD2 wild-type and mutant.

(F). The axon length of neurons expressing either wild-type or mutant alleles of BICD2 was
quantified. Expression of BICD2 mutants correlated with shorter axonal lengths. A one-way
ANOVA was used for the quantifications shown in panels E and F and the values were
compared to the mean of BICD2_wt. n = not significant, **, $p<0.01$, ***, $p<0.001$, ****, $p<0.0001$.

Figure 6: BICD2 mutations are associated with altered interactomes.
(A-C). Volcano plots comparing the interactome of BICD2_wt vs BICD2_N188T (A), vs
BICD2_R694C (B) and vs BICD2_R747C (C). Interacting proteins that show at least a twofold
change in comparison to BICD2_wt and have a p value of at least 0.05 are indicated in the
shaded boxes. Red boxes indicate proteins that display a greater interaction with BICD2
mutants vs the wild-type, whereas blue boxes indicate proteins that display a lower interaction
vs the wild-type protein.

(D) The proteomics results were validated by repeating the experiment and analyzing the bound
fractions using the indicated antibodies. Streptavidin beads were used to purify the biotinylated
proteins.

(E-G). Quantification of binding of BICD2_wt and mutants with RANBP2 (E), VPS41 (F), and
CSPP1 (G). The level of binding for the mutants was compared to BICD2_wt. Consistent with
the proteomics results, BICD2_R747C displayed reduced binding to RANBP2 and VPS41. All
three mutants bound CSPP1 at a greater level than the wild-type. A one-way ANOVA was used
for this analysis. ns = not significant, *, p ≤ 0.05, **, $p<0.01$, ***, $p<0.001$, ****, $p<0.0001$.

Figure 7: BICD2_R747C is associated with a gain of function interaction with GRAMD1A.
(A). Lysates from cells expressing BICD2_wt and mutants were incubated with streptavidin
beads to purify biotinylated proteins. Bound proteins were eluted and analyzed by blotting using
the indicated antibodies. BICD2_R747C interacted with substantially more GRAMD1A than
either the control, BICD2_wt, or the other mutants.

(B). Quantification of binding of BICD2_wt and mutants with GRAMD1A. The level of binding for
the mutants was normalized to BICD2_wt. A one-way ANOVA was used for this analysis. ns =
not significant, ***, $p<0.001$.

(C). Cos7 cells were co-transfected with constructs expressing either BICD2_wt or mutant
(magenta) along with a plasmid expressing GRAMD1A-mScarlet3 (cyan). Except for cells
expressing BICD2_R747C, GRAMD1A was localized to the ER. By contrast, GRAMD1A was
highly enriched at the centrosome in cells expressing BICD2_R747C.

(D). Quantification of the co-localization between BICD2_wt and mutants with GRAMD1A. A one-way ANOVA was used for this analysis and the values were compared to the mean of BICD2_wt. ns = not significant, ****, p<0.0001.

Supplemental Figure1

(A). HEK cells expressing the indicated constructs were transfected with a plasmid expressing GTP locked RAB6A (GFP-RAB6A Q72L). Lysates were prepared and a co-immunoprecipitation experiment was carried out using V5 trap beads. The co-precipitating proteins were analyzed using the indicated antibodies. GFP-RAB6A Q72L specifically co-precipitated with full-length BICD2.

(B). HEK cells expressing the indicated constructs were transfected with either a control siRNA or an siRNA targeting endogenous BICD2. Three days after the transfections, lysates were prepared, and a co-immunoprecipitation experiment was carried out using V5 trap beads. The co-precipitated proteins were analyzed by western blotting. RANBP2 and VPS41 co-precipitate with BICD2-wt_TurboID in cells that are depleted of endogenous BICD2.

(C). Lysates were prepared from HEK cells expressing the indicated constructs and biotinylated proteins were purified using streptavidin beads. The precipitated proteins were analyzed by blotting using antibodies against VPS16 and VSP18. Both proteins were able to interact with full-length BICD2 but not the isolated cargo binding domain.

(D). Lysates were prepared from HEK cells expressing the indicated constructs and biotinylated proteins were purified using streptavidin beads. The precipitated proteins were analyzed by blotting using antibodies against RANBP2 and VPS41. Both proteins interact with full-length BICD2 but show a reduced interaction with BICD2_delCC1.

(E). HEK cells expressing the indicated constructs were transfected with either a control siRNA or an siRNA targeting endogenous VPS41. Three days after transfection, the biotinylated proteins were purified using streptavidin beads and analyzed by western blotting using antibodies against VPS16 and VSP18. Both proteins retain their interaction with BICD2 in cells depleted of endogenous VPS41.

Supplemental Figure2

(A). Lysates were prepared from cells transfected with the indicated siRNA. The lysates were analyzed by blotting using antibodies against DHC, BICD2, and alpha-Tubulin. The siRNAs were capable of depleting the target protein.

(B). Lysates were prepared from cells transfected with two different siRNAs targeting endogenous BICD2. Both siRNAs were capable of depleting BICD2.

(C). Lysates were prepared from cells transfected with either a control siRNA or an siRNA targeting KIF5B. The kif5b siRNA was capable of efficiently depleting KIF5B protein.

(D). Representative image of HeLa cells showing the localization of GFP-VPS41 vesicles in cells transfected with bicd2 siRNA-2.

(E-F). Representative images of HeLa cells transfected with either a control siRNA (E, E') or
bicd2 siRNA-1 (F, F'). The cells were processed for immunofluorescence using an antibody
against GM130, a Golgi marker (red to white LUT). The cells were also counterstained with
DAPI (cyan). Depletion of BICD2 results in fragmentation of the Golgi.

(G-H). Representative images of HeLa cells transfected with either a control siRNA (G, G') or
bicd2 siRNA-1 (H, H'). The cells were processed for immunofluorescence using antibodies
against CyclinB (a G2 cell cycle marker, green) and Pericentrin (a centrosome marker, red). The
cells were also counterstained with DAPI (cyan). Depletion of BICD2 results mis-localization of
the centrosome away from the perinuclear area.

(I-K) Representative images of HeLa cells transfected with either a control siRNA (I), bicd2
siRNA-1 (J) or bicd2 siRNA-2 (K). The cells were processed for immunofluorescence using an
antibody against LAMP1. Depletion of BICD2 resuls in perinuclear clustering of lysosomes.

(L). The number of motile lysosome particles were quantified in HeLa cells transfected with
either a control siRNA or an siRNA targeting BICD2. The number of motile lysosomes were
greatly reduced in BICD2 depleted cells.

(M, M'). Cos7 cells were transfected with a plasmid encoding BICD2-mNeongreen. The cells
were fixed and immunostained using an antibody against LAMP1. Over-expression of BICD2
results in the outward spreading of LAMP1 vesicles. The outline of the transfected cell is
indicated in E'. The scale bar in the microscopy images is 20 microns.

Supplemental Figure3
(A-B). Quantification of binding of BICD2_wt and mutants with DIC (A) and DCTN1 (B). The
level of binding for the mutants was compared to BICD2_wt. A one-way ANOVA was used for
this analysis. ns = not significant, *, p ≤ 0.05.

(C). A representative image of HeLa cells expressing BICD2-wt_TurboID. The cells were fixed
and processed using antibodies against V5 (which recognizes TurboID, green, C) and GM130
(a Golgi marker, red, C'). The cells were also counterstained with DAPI (cyan, merged image in
C''). Wild-type BICD2_mTurboID partially co-localized with the centrosome.

(D-G). Representative images of HEK cells expressing mGreenLantern positive peroxisomes
and co-expressing BICD2_wt (A), BICD2_N188T (B), BICD2_R694C (C), and BICD2_R747C
(D) fused to the FKBP dimerization domain. The addition of rapamycin causes the tethering of
peroxisomes to the indicated BICD2 construct. BICD2 mutants result in greater perinuclear
clustering of peroxisome in comparison to the wild-type protein.

(H) Lysates were prepared from cells expressing the indicated constructs and biotinylated
proteins were purified using streptavidin beads. The purified proteins were analyzed by western
blotting using an antibody against KIF5B.

(I) The binding results in panel H was quantified. The level of binding for the mutants was
compared to BICD2_wt. A one-way ANOVA was used for this analysis. ns = not significant, *, p
≤ 0.05, ***, p<0.001. BICD2_R694C and BICD2_R747C displayed reduced binding to KIF5B.

Supplemental Figure4

(A, B). HEK cells expressing the indicated constructs were used in a binding experiment. Biotinylated proteins were purified using streptavidin beads. Bound proteins were eluted and analyzed by blotting using the indicated antibodies. Importin beta is biotinylated by wild-type BICD2 and BICD2_R694C. However, the binding between the two proteins is reduced in the BICD2_R747C background. Two replicate experiments are shown.

(C). HEK cells expressing BICD2_wt were transfected with the indicated siRNA. Three days after the siRNA transfection, lysates were prepared and biotinylated proteins were purified using streptavidin beads. Bound proteins were analyzed by blotting using the indicated antibodies. The ability of Importin beta to be biotinylated by BICD2 depends on RANBP2.

(D). Lysates were prepared from HEK cells expressing the indicated constructs. Binding was performed using streptavidin beads and the bound proteins were analyzed by blotting. VPS16 and VPS18 displayed reduced binding to BICD2_R747C.

Supplemental Figure5

(A). HEK cells expressing the indicated constructs were transfected with an siRNA targeting endogenous BICD2. Three days after transfection, lysates were prepared, and biotinylated proteins were purified using streptavidin beads. The precipitated proteins were analyzed by blotting using antibodies against RANBP2 and GRAMD1A. Whereas the interaction between BICD2_R747C and RANBP2 is reduced in cells depleted of endogenous BICD2, the interaction of this mutant with GRAMD1A is increased in these same cells.

(B-F). Cos7 cells were co-transfected with plasmids encoding BICD2_R747C-mNeon (green, B, C) and GRAMD1A-mScarlet3 (cyan, D). The cells were fixed and processed using an antibody against Pericentrin (a centrosome marker, magenta, E). The merged image is shown in F.

(G, H). Cos7 cells were co-transfected with plasmids encoding an ER marker (Addgene plasmid #137805, (Chertkova et al., 2020)) and BICD2_wt (G) or BICD2_R747C (H). In contrast to what was observed with GRAMD1A, expression of BICD2_R747C does not alter the localization of the ER marker.

Supplemental table 1
The wild-type BICD2 interactome.

Supplemental table 2
The interactome of BICD2_N188T vs BICD2_wt.

Supplemental table 3
The interactome of BICD2_R694Cvs BICD2_wt.

Supplemental table 4
The interactome of BICD2_R747Cvs BICD2_wt.

Supplemental table 5
The full list of reagents used in this manuscript.

Supplemental video 1

985 Live imaging of SiR lysosome labeled vesicles in HeLa cell transfected with a control siRNA.

986 **Supplemental video 2**
987 Live imaging of SiR lysosome labeled vesicles in HeLa cell transfected with an siRNA targeting
988 BICD2.

References

Ali, M.Y., H. Lu, P.M. Fagnant, J.E. Macfarlane, and K.M. Trybus. 2025. BicD and MAP7 collaborate to activate homodimeric Drosophila kinesin-1 by complementary mechanisms. *bioRxiv*.

Angilletta, I., R. Ferrante, R. Giansante, L. Lombardi, A. Babore, A. Dell'Elice, E. Alessandrelli, S. Notarangelo, M. Ranaudo, C. Palmarini, V. De Laurenzi, L. Stuppia, and C. Rossi. 2023. Spinal Muscular Atrophy: An Evolving Scenario through New Perspectives in Diagnosis and Advances in Therapies. *Int J Mol Sci*. 24.

Besprozvannaya, M., E. Dickson, H. Li, K.S. Ginburg, D.M. Bers, J. Auwerx, and J. Nunnari. 2018. GRAM domain proteins specialize functionally distinct ER-PM contact sites in human cells. *Elife*. 7.

Bonet-Ponce, L., A. Beilina, C.D. Williamson, E. Lindberg, J.H. Kluss, S. Saez-Atienzar, N. Landeck, R. Kumaran, A. Mamais, C.K.E. Bleck, Y. Li, and M.R. Cookson. 2020. LRRK2 mediates tubulation and vesicle sorting from lysosomes. *Sci Adv*. 6.

Branon, T.C., J.A. Bosch, A.D. Sanchez, N.D. Udeshi, T. Svinkina, S.A. Carr, J.L. Feldman, N. Perrimon, and A.Y. Ting. 2018. Efficient proximity labeling in living cells and organisms with TurboID. *Nat Biotechnol*. 36:880–887.

Canty, J.T., and A. Yildiz. 2020. Activation and Regulation of Cytoplasmic Dynein. *Trends in biochemical sciences*. 45:440–453.

Chan, S.H.S., N. van Alfen, I.J. Thuestad, J. Ip, A.O. Chan, C. Mak, B.H. Chung, A. Verrips, and E.J. Kamsteeg. 2018. A recurrent de novo DYNC1H1 tail domain mutation causes spinal muscular atrophy with lower extremity predominance, learning difficulties and mild brain abnormality. *Neuromuscul Disord*. 28:750–756.

Chertkova, A.O., M. Mastop, M. Postma, N. van Bommel, S. van der Niet, K.L. Batenburg, L. Joosen, T.W.J. Gadella, Y. Okada, and J. Goedhart. 2020. Robust and Bright Genetically Encoded Fluorescent Markers for Highlighting Structures and Compartments in Mammalian Cells. *bioRxiv*:160374.

Chowdhury, S., S.A. Ketcham, T.A. Schroer, and G.C. Lander. 2015. Structural organization of the dynein-dynactin complex bound to microtubules. *Nat Struct Mol Biol*. 22:345–347.

Cui, H., M.Y. Ali, P. Goyal, K. Zhang, J.Y. Loh, K.M. Trybus, and S.R. Solmaz. 2020. Coiled-coil registry shifts in the F684I mutant of Bicaudal D result in cargo-independent activation of dynein motility. *Traffic*. 21:463–478.

Das, J., J.B. Lilleker, K. Jabbal, and J. Ealing. 2018. A missense mutation in DYNC1H1 gene causing spinal muscular atrophy - Lower extremity, dominant. *Neurol Neurochir Pol*. 52:293–297.

Dienstbier, M., F. Boehl, X. Li, and S.L. Bullock. 2009. Egalitarian is a selective RNA-binding protein linking mRNA localization signals to the dynein motor. *Genes & development*. 23:1546–1558.

Franker, M.A., and C.C. Hoogenraad. 2013. Microtubule-based transport - basic mechanisms, traffic rules and role in neurological pathogenesis. *J Cell Sci*. 126:2319–2329.

Frasquet, M., A. Camacho, R. Vilchez, H. Argente-Escrig, E. Millet, J.F. Vazquez-Costa, R. Silla, A. Sanchez-Monteagudo, J.J. Vilchez, C. Espinos, V. Lupo, and T. Sevilla. 2020. Clinical spectrum of BICD2 mutations. *Eur J Neurol*. 27:1327–1335.

Gallisa-Sune, N., P. Sanchez-Fernandez-de-Landa, F. Zimmermann, M. Serna, L. Regue, J. Paz, O. Llorca, J. Luders, and J. Roig. 2023. BICD2 phosphorylation regulates dynein function and centrosome separation in G2 and M. *Nat Commun*. 14:2434.

Goldman, C.H., H. Neiswender, F. Baker, R. Veeranan-Karmegam, S. Misra, and G.B. Gonsalvez. 2021. Optimal RNA binding by Egalitarian, a Dynein cargo adaptor, is critical for maintaining oocyte fate in Drosophila. *RNA Biol*:1–14.

Goldman, C.H., H. Neiswender, R. Veeranan-Karmegam, and G.B. Gonsalvez. 2019. The Egalitarian binding partners Dynein light chain and Bicaudal-D act sequentially to link mRNA to the Dynein motor. *Development*. 146.

Goncalves, J.C., S. Quintremil, J. Yi, and R.B. Vallee. 2020. Nesprin-2 Recruitment of BicD2 to the Nuclear Envelope Controls Dynein/Kinesin-Mediated Neuronal Migration In Vivo. *Current biology : CB*. 30:3116–3129 e3114.

Grigoriev, I., D. Splinter, N. Keijzer, P.S. Wulf, J. Demmers, T. Ohtsuka, M. Modesti, I.V. Maly, F. Grosveld, C.C. Hoogenraad, and A. Akhmanova. 2007. Rab6 regulates transport and targeting of exocytotic carriers. *Dev Cell*. 13:305–314.

Guardia, C.M., G.G. Farias, R. Jia, J. Pu, and J.S. Bonifacino. 2016. BORC Functions Upstream of Kinesins 1 and 3 to Coordinate Regional Movement of Lysosomes along Different Microtubule Tracks. *Cell Rep*. 17:1950–1961.

Hoogenraad, C.C., and A. Akhmanova. 2016. Bicaudal D Family of Motor Adaptors: Linking Dynein Motility to Cargo Binding. *Trends in cell biology*. 26:327–340.

Hoogenraad, C.C., A. Akhmanova, S.A. Howell, B.R. Dortland, C.I. De Zeeuw, R. Willemsen, P. Visser, F. Grosveld, and N. Galjart. 2001. Mammalian Golgi-associated Bicaudal-D2 functions in the dynein-dynactin pathway by interacting with these complexes. *The EMBO journal*. 20:4041–4054.

Huynh, W., and R.D. Vale. 2017. Disease-associated mutations in human BICD2 hyperactivate motility of dynein-dynactin. *The Journal of cell biology*. 216:3051–3060.

Kapitein, L.C., M.A. Schlager, W.A. van der Zwan, P.S. Wulf, N. Keijzer, and C.C. Hoogenraad. 2010. Probing intracellular motor protein activity using an inducible cargo trafficking assay. *Biophys J*. 99:2143–2152.

Koboldt, D.C., M.A. Waldrop, R.K. Wilson, and K.M. Flanigan. 2020. The Genotypic and Phenotypic Spectrum of BICD2 Variants in Spinal Muscular Atrophy. *Ann Neurol*. 87:487–496.

Liu, Y., H.K. Salter, A.N. Holding, C.M. Johnson, E. Stephens, P.J. Lukavsky, J. Walshaw, and S.L. Bullock. 2013. Bicaudal-D uses a parallel, homodimeric coiled coil with heterotypic registry to coordinate recruitment of cargos to dynein. *Genes & development*. 27:1233–1246.

Mach, J.M., and R. Lehmann. 1997. An Egalitarian-BicaudalD complex is essential for oocyte specification and axis determination in Drosophila. *Genes & development*. 11:423–435.

Manigrasso, G., K. Saharat, P. Chaichana, C. Kullapanich, S. Atwal, J. Boulanger, T.E. Morgan, H. Kramer, J. Salje, and A.P. Carter. 2025. The intracellular bacterium Orientia tsutsugamushi uses the autotransporter ScaC to activate BICD adaptors for dynein-based motility. *Nat Commun*. 16:6122.

Martinez-Carrera, L.A., and B. Wirth. 2015. Dominant spinal muscular atrophy is caused by mutations in BICD2, an important golgin protein. *Front Neurosci*. 9:401.

Matanis, T., A. Akhmanova, P. Wulf, E. Del Nery, T. Weide, T. Stepanova, N. Galjart, F. Grosveld, B. Goud, C.I. De Zeeuw, A. Barnekow, and C.C. Hoogenraad. 2002. Bicaudal-D regulates COPI-independent Golgi-ER transport by recruiting the dynein-dynactin motor complex. *Nature cell biology*. 4:986–992.

McClintock, M.A., C.I. Dix, C.M. Johnson, S.H. McLaughlin, R.J. Maizels, H.T. Hoang, and S.L. Bullock. 2018. RNA-directed activation of cytoplasmic dynein-1 in reconstituted transport RNPs. *Elife*. 7.

McKenney, R.J., W. Huynh, M.E. Tanenbaum, G. Bhabha, and R.D. Vale. 2014. Activation of cytoplasmic dynein motility by dynactin-cargo adapter complexes. *Science*. 345:337–341.

Mohler, J., and E.F. Wieschaus. 1986. Dominant maternal-effect mutations of Drosophila melanogaster causing the production of double-abdomen embryos. *Genetics*. 112:803–822.

Naito, T., B. Ercan, L. Krshnan, A. Triebl, D.H.Z. Koh, F.Y. Wei, K. Tomizawa, F.T. Torta, M.R.
Wenk, and Y. Saheki. 2019. Movement of accessible plasma membrane cholesterol by
the GRAMD1 lipid transfer protein complex. *eLife*. 8.
Neveling, K., L.A. Martinez-Carrera, I. Holker, A. Heister, A. Verrips, S.M. Hosseini-Barkooie, C.
Gilissen, S. Vermeer, M. Pennings, R. Meijer, M. te Riele, C.J. Frijns, O. Suchowersky, L.
MacLaren, S. Rudnik-Schoneborn, R.J. Sinke, K. Zerres, R.B. Lowry, H.H. Lemmink, L.
Garbes, J.A. Veltman, H.J. Schelhaas, H. Scheffer, and B. Wirth. 2013. Mutations in
BICD2, which encodes a golgin and important motor adaptor, cause congenital
autosomal-dominant spinal muscular atrophy. *American journal of human genetics*.
92:946–954.
Oates, E.C., A.M. Rossor, M. Hafezparast, M. Gonzalez, F. Speziani, D.G. MacArthur, M. Lek,
E. Cottenie, M. Scoto, A.R. Foley, M. Hurles, H. Houlden, L. Greensmith, M. Auer-
Grumbach, T.R. Pieber, T.M. Strom, R. Schule, D.N. Herrmann, J.E. Sowden, G. Acsadi,
M.P. Menezes, N.F. Clarke, S. Zuchner, Uk10K, F. Muntoni, K.N. North, and M.M. Reilly.
2013. Mutations in BICD2 cause dominant congenital spinal muscular atrophy and
hereditary spastic paraplegia. *American journal of human genetics*. 92:965–973.
Olenick, M.A., and E.L.F. Holzbaur. 2019. Dynein activators and adaptors at a glance. *J Cell Sci*.
132.
Passmore, J.B., W. Nijenhuis, and L.C. Kapitein. 2021. From observing to controlling: Inducible
control of organelle dynamics and interactions. *Curr Opin Cell Biol*. 71:69–76.
Peeters, K., I. Litvinenko, B. Asselbergh, L. Almeida-Souza, T. Chamova, T. Geuens, E. Ydens,
M. Zimon, J. Irobi, E. De Vriendt, V. De Winter, T. Ooms, V. Timmerman, I. Tournev, and
A. Jordanova. 2013. Molecular defects in the motor adaptor BICD2 cause proximal
spinal muscular atrophy with autosomal-dominant inheritance. *American journal of
human genetics*. 92:955–964.
Ravenscroft, G., N. Di Donato, G. Hahn, M.R. Davis, P.D. Craven, G. Poke, K.R. Neas, T.M.
Neuhann, W.B. Dobyns, and N.G. Laing. 2016. Recurrent de novo BICD2 mutation
associated with arthrogryposis multiplex congenita and bilateral perisylvian
polymicrogyria. *Neuromuscul Disord*. 26:744–748.
Reck-Peterson, S.L., W.B. Redwine, R.D. Vale, and A.P. Carter. 2018. The cytoplasmic dynein
transport machinery and its many cargoes. *Nature reviews. Molecular cell biology*.
19:382–398.
Redwine, W.B., M.E. DeSantis, I. Hollyer, Z.M. Htet, P.T. Tran, S.K. Swanson, L. Florens, M.P.
Washburn, and S.L. Reck-Peterson. 2017. The human cytoplasmic dynein interactome
reveals novel activators of motility. *eLife*. 6.
Sandhu, J., S. Li, L. Fairall, S.G. Pfisterer, J.E. Gurnett, X. Xiao, T.A. Weston, D. Vashi, A.
Ferrari, J.L. Orozco, C.L. Hartman, D. Strugatsky, S.D. Lee, C. He, C. Hong, H. Jiang,
L.A. Bentolila, A.T. Gatta, T.P. Levine, A. Ferng, R. Lee, D.A. Ford, S.G. Young, E.
Ikonen, J.W.R. Schwabe, and P. Tontonoz. 2018. Aster Proteins Facilitate Nonvesicular
Plasma Membrane to ER Cholesterol Transport in Mammalian Cells. *Cell*. 175:514–529
e520.
Schlager, M.A., H.T. Hoang, L. Urnavicius, S.L. Bullock, and A.P. Carter. 2014. In vitro
reconstitution of a highly processive recombinant human dynein complex. *The EMBO
journal*. 33:1855–1868.
Seyfried, N.T., E.B. Dammer, V. Swarup, D. Nandakumar, D.M. Duong, L. Yin, Q. Deng, T.
Nguyen, C.M. Hales, T. Wingo, J. Glass, M. Gearing, M. Thambisetty, J.C. Troncoso,
D.H. Geschwind, J.J. Lah, and A.I. Levey. 2017. A Multi-network Approach Identifies
Protein-Specific Co-expression in Asymptomatic and Symptomatic Alzheimer's Disease.
Cell Syst. 4:60–72 e64.

Sladewski, T.E., N. Billington, M.Y. Ali, C.S. Bookwalter, H. Lu, E.B. Krementsova, T.A. Schroer, and K.M. Trybus. 2018. Recruitment of two dyneins to an mRNA-dependent Bicaudal D transport complex. *Elife*. 7.

Sleigh, J.N., A.M. Rossor, A.D. Fellows, A.P. Tosolini, and G. Schiavo. 2019. Axonal transport and neurological disease. *Nat Rev Neurol*. 15:691–703.

Soucek, S., Y. Zeng, D.L. Bellur, M. Bergkessel, K.J. Morris, Q. Deng, D. Duong, N.T. Seyfried, C. Guthrie, J.P. Staley, M.B. Fasken, and A.H. Corbett. 2016. The Evolutionarily-conserved Polyadenosine RNA Binding Protein, Nab2, Cooperates with Splicing Machinery to Regulate the Fate of pre-mRNA. *Mol Cell Biol*. 36:2697–2714.

Spang, A. 2016. Membrane Tethering Complexes in the Endosomal System. *Front Cell Dev Biol*. 4:35.

Splinter, D., D.S. Razafsky, M.A. Schlager, A. Serra-Marques, I. Grigoriev, J. Demmers, N. Keijzer, K. Jiang, I. Poser, A.A. Hyman, C.C. Hoogenraad, S.J. King, and A. Akhmanova. 2012. BICD2, dynactin, and LIS1 cooperate in regulating dynein recruitment to cellular structures. *Mol Biol Cell*. 23:4226–4241.

Splinter, D., M.E. Tanenbaum, A. Lindqvist, D. Jaarsma, A. Flotho, K.L. Yu, I. Grigoriev, D. Engelsma, E.D. Haasdijk, N. Keijzer, J. Demmers, M. Fornerod, F. Melchior, C.C. Hoogenraad, R.H. Medema, and A. Akhmanova. 2010. Bicaudal D2, dynein, and kinesin-1 associate with nuclear pore complexes and regulate centrosome and nuclear positioning during mitotic entry. *PLoS Biol*. 8:e1000350.

Storbeck, M., B. Horsberg Eriksen, A. Unger, I. Holker, I. Aukrust, L.A. Martinez-Carrera, W.A. Linke, A. Ferbert, R. Heller, M. Vorgerd, G. Houge, and B. Wirth. 2017. Phenotypic extremes of BICD2-opathies: from lethal, congenital muscular atrophy with arthrogryposis to asymptomatic with subclinical features. *Eur J Hum Genet*. 25:1040–1048.

Synofzik, M., L.A. Martinez-Carrera, T. Lindig, L. Schols, and B. Wirth. 2014. Dominant spinal muscular atrophy due to BICD2: a novel mutation refines the phenotype. *J Neurol Neurosurg Psychiatry*. 85:590–592.

Terawaki, S., A. Yoshikane, Y. Higuchi, and K. Wakamatsu. 2015. Structural basis for cargo binding and autoinhibition of Bicaudal-D1 by a parallel coiled-coil with homotypic registry. *Biochem Biophys Res Commun*. 460:451–456.

Torisawa, T., M. Ichikawa, A. Furuta, K. Saito, K. Oiwa, H. Kojima, Y.Y. Toyoshima, and K. Furuta. 2014. Autoinhibition and cooperative activation mechanisms of cytoplasmic dynein. *Nat Cell Biol*. 16:1118–1124.

Urnavicius, L., K. Zhang, A.G. Diamant, C. Motz, M.A. Schlager, M. Yu, N.A. Patel, C.V. Robinson, and A.P. Carter. 2015. The structure of the dynactin complex and its interaction with dynein. *Science*. 347:1441–1446.

van Beuningen, S.F., and C.C. Hoogenraad. 2016. Neuronal polarity: remodeling microtubule organization. *Curr Opin Neurobiol*. 39:1–7.

van der Beek, J., C. Jonker, R. van der Welle, N. Liv, and J. Klumperman. 2019. CORVET, CHEVI and HOPS - multisubunit tethers of the endo-lysosomal system in health and disease. *J Cell Sci*. 132.

van der Welle, R.E.N., R. Jobling, C. Burns, P. Sanza, J.A. van der Beek, A. Fasano, L. Chen, F.J. Zwartkruis, S. Zwakenberg, E.F. Griffin, C. Ten Brink, T. Veenendaal, N. Liv, C.M.A. van Ravenswaaij-Arts, H.H. Lemmink, R. Pfundt, S. Blaser, C. Sepulveda, A.M. Lozano, G. Yoon, T. Santiago-Sim, C.S. Asensio, G.A. Caldwell, K.A. Caldwell, D. Chitayat, and J. Klumperman. 2021. Neurodegenerative VPS41 variants inhibit HOPS function and mTORC1-dependent TFEB/TFE3 regulation. *EMBO Mol Med*. 13:e13258.

Wei, Z., C. Hao, K.R. Radeen, R. Srinivasagan, J.K. Chen, S. Sharma, M.E. McGee-Lawrence, M.W. Hamrick, V.M. Monnier, and X. Fan. 2024. Prevention of age-related truncation of

1189   gamma-glutamylcysteine ligase catalytic subunit (GCLC) delays cataract formation. *Sci*
1190   *Adv*. 10:eadl1088.
1191   Wharton, R.P., and G. Struhl. 1989. Structure of the Drosophila BicaudalD protein and its role in
1192   localizing the posterior determinant nanos. *Cell*. 59:881–892.
1193   Yi, J., X. Zhao, C.R. Noell, P. Helmer, S.R. Solmaz, and R.B. Vallee. 2023. Role of Nesprin-2
1194   and RanBP2 in BICD2-associated brain developmental disorders. *PLoS Genet*.
1195   19:e1010642.
1196   Yildiz, A. 2025. Mechanism and regulation of kinesin motors. *Nat Rev Mol Cell Biol*. 26:86–103.
1197   Zhang, K., H.E. Foster, A. Rondelet, S.E. Lacey, N. Bahi-Buisson, A.W. Bird, and A.P. Carter.
1198   2017. Cryo-EM Reveals How Human Cytoplasmic Dynein Is Auto-inhibited and
1199   Activated. *Cell*. 169:1303–1314 e1318.
1200   Zhao, X., S. Quintremil, E.D. Rodriguez Castro, H. Cui, D. Moraga, T. Wang, R.B. Vallee, and
1201   S.R. Solmaz. 2024. Molecular mechanism for recognition of the cargo adapter
1202   Rab6(GTP) by the dynein adapter BicD2. *Life Sci Alliance*. 7.

1203

Figure 1

Figure 2

Figure 3

Figure 4

BICD2_wt

BICD2_N188T

BICD2_R694C

BICD2_R747C

Figure 5

Figure 6

Figure 7