Emergent loss of microbial biodiversity with increasing resource diversity

Or Shalev[1*], Xiaozhou Ye[1], Joachim Kilian[2], Mark Stahl[2], Christoph Ratzke[1*]

[1]Interfaculty Institute for Microbiology and Infection Medicine Tübingen (IMIT), Cluster of Excellence EXC 2124 "Controlling Microbes to Fight Infections" (CMFI), University of Tübingen, Calwerstrasse 7/1, 72076 Tübingen

[2]Zentrum für Molekularbiologie der Pflanzen (ZMBP), University of Tübingen, Auf der Morgenstelle 32, 72076 Tübingen

*Correspondence to: or.shalev@uni-tuebingen.de, christoph.ratzke@uni-tuebingen.de

Abstract

The origin of biodiversity is arguably the most fundamental question in ecology — especially how countless microbial species coexist within a single community. A prevailing hypothesis holds that microbial species coexist by specializing on different resources, thereby reducing competition. Accordingly, increasing resource diversity is expected to promote species coexistence and boost biodiversity. Surprisingly, we observe the opposite: microbial biodiversity often declines as resource diversity increases. This unexpected result emerges from a widespread physiological response across diverse microbial taxa, in which more complex environments trigger higher overall resource uptake. This intensifies competition and drives biodiversity loss. Updating a central ecological model to include this physiological trait accurately reproduces the observed decline. Our findings demonstrate that physiological changes at the individual level can overturn classical ecological principles and scale up to restructure entire ecosystems.

Introduction

Microbial life flourishes in astonishing biodiversity. A single gram of soil can harbor over ten thousand bacterial species[1]. Yet how so many species coexist has remained a central question in ecology for nearly a century[2–4]. Classic theory holds that when species compete for the same resource, the most efficient consumer prevails, driving others to extinction through competitive exclusion[2]. Coexistence, therefore, requires niche differentiation: species avoid direct competition by exploiting different

resources[5]. By this logic, greater resource diversity should support more coexisting
species[6,7] (**Fig. 1A**).

This principle is formalized in the Consumer–Resource Model[7], which tracks
microbial growth and resource dynamics. When species are assigned random
metabolic traits and uptake efficiencies, the model predicts that increasing resource
diversity allows more species to persist—as naively expected (**Fig. 1B;**
Supplementary Fig. 1A-B). Although additional mechanisms such as stochastic
processes[8], viral infections[9], metabolic interdependencies[10], and environmental
fluctuations[11,12] have also been proposed to sustain microbial biodiversity, the concept
that resource diversity drives species biodiversity remains foundational. Yet despite
decades of theoretical support, empirical evidence for this link remains scarce[13,14],
raising the question of whether this long-standing theory truly holds in microbial
ecosystems?

As a first step toward addressing this question, we re-analyzed multi-omics data
from the Earth Microbiome Project[15] and found that resource diversity does not
consistently correlate with biodiversity (**Fig. 1C**; **Supplementary Fig. 1C-D**): in some
ecosystems biodiversity increases with higher resource diversity, while in others, it
decreases. However, natural systems are uncontrolled, making it difficult to isolate
causal effects. Yet only a few controlled experiments have directly addressed this
question[13,14] . Among them, just one—by Pacheco et al.[14]— examined how increasing
resource complexity affects diversity in a defined microbial community, finding only a
weak biodiversity increase. However, the use of a single community limits the
generalizability of these findings.

Here, we systematically tested hundreds of microbial communities and thousands
of interactions, and found that—contrary to expectations—microbial biodiversity tends
to decline as resource diversity increases.

Results

Greater resource diversity often reduces microbial biodiversity
To test the fundamental relationship between resource diversity and microbial
biodiversity, we assembled random microbial communities across a range of resource
diversities (**Fig. 1D**; see **Methods** for details). Communities were cultivated for nine

days under a daily dilution regime to allow them to stabilize (**Supplementary Fig. 2**),
and final community composition was assessed using 16S rRNA amplicon
sequencing. This yielded 701 sequenced samples across different experimental
conditions (**Supplementary Data 1**).
Contrary to predictions from the Consumer–Resource Model, microbial biodiversity
declined in a large fraction of communities as resource diversity increased, while
increases in biodiversity were rare and typically modest (**Fig. 1E**, compared with Fig.
1B; **Supplementary Fig. 1E–F**; **Supplementary Fig. 3**). These trends mirrored
patterns observed in natural ecosystems, though the declines were more consistent
and pronounced under controlled conditions (compared with Fig. 1C). Together, these
findings challenge the ecological expectation that higher resource diversity promotes
coexistence and raise the question of what drives this widespread biodiversity loss.

Figure 1: Microbial biodiversity often declines with increasing resource diversity across natural and experimental communities. A. When microbial species prefer different resources, higher resource diversity is expected to promote coexistence by reducing competition. **B.** This expectation is supported by simulations using a Consumer–Resource Model (**Supplementary Text 1**; 25 simulated communities). (**Top**) Example communities showing the weakest and strongest biodiversity responses to increasing resource number. (**Bottom**) For each community, biodiversity response was quantified as the slope (β_1) of a linear model (Shannon index = β_0 + β_1 × number of resources + ϵ). The mean slope across all simulated communities is shown in **Supplementary Fig. 1A** (Shannon index) and **Supplementary Fig. 1B** (richness). **C.** In natural microbial communities, biodiversity does not

consistently increase with resource diversity. Data from the Earth Microbiome Project[15] (462 samples
across 13 ecosystems) combine 16S-based diversity estimates with mass spectrometry-derived
resource profiles (Methods). (**Top**) Example ecosystems showing positive and negative relationships
between resource diversity and microbial biodiversity (full results in Supplementary Fig. 1C). A similar
pattern is observed using richness instead of Shannon diversity (Supplementary Fig. 1D). Notably, the
most data-supported—'animal distal gut (non-saline)'—shows a significant negative correlation.
(**Bottom**) Extending this analysis to all 13 ecosystems, we derive the slope (β_1) of Shannon diversity
index vs. resource number as described in Fig. 1B. **D.** To experimentally test the effect of resource
complexity on microbial diversity, we used 14 bacterial strains (Supplementary Data 2) to assemble 14
random synthetic communities—seven with 6 members and seven with 8 members. Each community
was cultured across five levels of resource diversity (1, 2, 4, 8, or 16 carbon sources), totaling 54
conditions per community. **E.** In our controlled experiments, microbial biodiversity generally decreased
with increasing resource diversity, while the magnitude of this effect varied across communities. (**Top**)
Communities with the strongest biodiversity decrease and increase in response to resource diversity
are shown, along with relative abundances of their member strains. Shading indicates mean ± SEM.
Results using richness are shown in Supplementary Fig. 1F. (**Bottom**) Extending this analysis to all 14
communities, we derive the slope (β_1) of Shannon diversity index vs. resource number as described in
Fig 1B.

Resource diversity fuels competition, which reduces coexistence

Microbial communities are strongly shaped by interactions between their constituent
strains. We hypothesized that the observed loss of biodiversity might result from a
fundamental shift in these interactions triggered by increasing resource diversity. To
test this, we systematically co-incubated all pairwise combinations of the 14 strains
used in our synthetic communities under a wide range of resource conditions, grouped
into three levels of resource diversity: 1, 8, and 16 resources. For each pairwise co-
culture, we measured total biomass (OD600) and relative abundances after five daily
dilutions to infer interaction outcomes. We also measured how each strain grew in
isolation in each environment, allowing us to disentangle intrinsic growth from
interaction effects. We then inferred interaction strengths between all strains by fitting
individual and pairwise data, along with community data (**Fig. 1E**), to a generalized
Lotka–Volterra (gLV) model using Bayesian regression (**Supplementary Text 2**). This
yielded interaction matrices for the three levels of resource diversity (illustrated in **Fig.
2A**; **Supplementary Fig. 4**), which, when used to simulate community assembly,
reproduced the observed decline in biodiversity (**Supplementary Fig. 5**), supporting
the validity of the inferred interactions. In these matrices, diagonal terms represent
self-inhibition (how much a strain limits its own growth; α_{self}), and off-diagonal terms

capture how strongly a strain inhibits others (α_{others}). The α values represent *per-capita* interaction strengths, meaning they are independent of population densities. According to theory, when interspecies inhibition exceeds self-inhibition ($\alpha_{others} > \alpha_{self}$), competitive exclusion will occur, where one strain will outcompete the other[16]. Therefore, if biodiversity loss is driven by pairwise interactions, we would expect increasing resource diversity to shift this balance in favor of exclusion. Interestingly, both self- and interspecies inhibition became weaker as resource diversity increased (**Fig. 2B**, top; **Supplementary Fig. 6, 7**). However, self-inhibition (α_{self}) decreased more rapidly than interspecies inhibition (α_{others}; **Fig. 2B**, middle), leading to a greater number of predicted competitive exclusions as resource complexity increased (**Fig. 2B**, bottom; **Supplementary Fig. 6C**). Counterintuitively, it is therefore not stronger per-capita inhibition of competitors, but greater individual growth—driven by reduced self-inhibition—that drives competitive exclusion and biodiversity loss. Increased growth leads to larger population sizes, which exert stronger total inhibitory effects on competitors, even as per-capita interspecies inhibition becomes weaker with increasing resource diversity.

The above findings explain the average biodiversity loss with increasing resource diversity, but the strength of this effect varied across communities (Fig. 1E). This suggests that community composition influenced the outcome, with some strains playing a larger role than others. To identify which strains drive biodiversity loss, we defined an 'impact factor'—a summary metric combining all four interaction coefficients from a pairwise interaction ($2\times\alpha_{self} + 2\times\alpha_{others}$) into a single value (**Fig. 2A**; **Methods**). This metric reflects how strongly one strain affects the growth of its interaction partner: a value of 1 indicates neutrality, values below 1 indicate suppression, and values above 1 indicate facilitation.

We hypothesized that strains whose impact factor decreased—i.e., became more suppressive—as resource diversity increased would be key drivers of biodiversity loss. To test this, we extensively simulated communities using synthetic strain sets generated by adding random noise to the experimentally derived interaction matrices (**Methods**). This approach allowed us to generalize beyond the limited statistical power of the experimental dataset. For each simulated community, we calculated biodiversity under low and high resource diversity and matched this to strain-specific changes in impact factor. Strains that became more suppressive with increasing

resource diversity consistently drove greater biodiversity loss, validating our
hypothesis (**Fig. 2C**; **Supplementary Fig. 9**).

To confirm these predictions experimentally, we examined our experimental dataset
and found that strains becoming more suppressive with increasing resource diversity
contributed more strongly to biodiversity loss (**Supplementary Fig. 10**). One strain in
particular, '79', stood out—showing a pronounced decrease in impact factor (i.e.,
increased suppression) as resource diversity rose (**Fig. 2D**; **Supplementary Fig. 11**).
Comparing communities with and without strain '79' revealed a striking pattern: those
containing it exhibited sharp biodiversity declines with increasing resource diversity,
while those without it showed slight biodiversity gains (**Fig. 2E**). In communities with
strain '79', its relative abundance increased steadily with resource diversity, eventually
dominating at 16 resources as the total abundance of other strains declined (**Fig. 2F**;
Supplementary Fig. 12). Shannon diversity mirrored this trend, decreasing in parallel
with the collapse of the remaining community members. Taken together, these
findings show that increasing resource diversity can amplify the suppressive potential
of specific strains, which in turn drives biodiversity loss.

This raises two key questions: how common is it for strains to become more
competitive as resource diversity increases, and what physiological mechanisms
underlie this shift?

Figure 2: Increasing pairwise competitiveness reduces community biodiversity as resource diversity increases. **A.** Overview of the experimental and modeling workflow. We studied pairwise interactions among 14 bacterial strains across 1, 8, and 16 carbon source conditions. Each pair was tested in an average of 31 conditions, and individual strains were profiled in all their respective conditions, totaling 619 measurements (Supplementary Data 3). OD_{600} and 16S rRNA sequencing were used to quantify growth and composition. Interaction matrices were inferred by applying Bayesian regression to a generalized Lotka–Volterra model (Methods), using observed data to estimate intra-species (α_{self}) and inter-species (α_{others}) interaction strengths (Supplementary Text 2; results in panel B). Interaction matrices were inferred by a Bayesian regression on a generalized Lotka–Volterra model (Supplementary Fig. 4), allowing comparison of intra-species (α_{self}) and inter-species (α_{others}) interactions

across resource levels (panel B). To assess strain-level contributions to diversity loss, we defined an
impact factor based on the change in a strain's partner's abundance in co-culture versus monoculture.
As examples, a drop from 6 to 2 yields an impact factor of 0.33. **B.** Resource complexity shifted
communities toward competitive exclusion. On average, α_{self} decreased more than α_{others}, reducing
coexistence potential (top and middle panels). The proportion of interactions where $\alpha_{others} > \alpha_{self}$
increased with resource number (bottom panel). Trend lines show mean ± SEM. **C.** Simulations using
noise-perturbed experimental data showed that strains with more negative impact factor slopes (1 vs. 16
resources) contributed more to biodiversity loss. The cartoon illustrates how increased pairwise
suppression by a single strain can scale up to reduce overall community diversity. **D.** Strain '79' showed
the steepest increase in suppressiveness among experimentally tested strains. For each strain, we
fitted $\log_{10}$(impact factor) = $\beta_0 + \beta_1$ × number of resources + ε across 1-, 8-, and 16-resource conditions
(3,398 values total). Strain '79' showed the most negative β_1. The cartoon illustrates how its suppressive
effect increased consistently across interaction partners with resource diversity. **E.** Communities
containing strain '79' (n = 8) showed a sharp biodiversity decline with increasing resource diversity,
while those without it (n = 6) slightly increased. Trend lines show the mean ± SEM. **F.** In these same
communities, strain '79' increased in relative abundance while other strains declined, along with a drop
in Shannon diversity. Trend lines show the mean ± SEM. Shown are four representative examples (full
data in Supplementary Fig. 12).

Increased microbial competitiveness with resource diversity is widespread

To assess how often strains become more competitive with increasing resource
diversity, we measured our established impact factor metric (illustrated in **Fig. 2A**) for
298 strains against a sentinel strain across a wide range of resource conditions,
generating ~32,000 data points (**Fig. 3A**; **Methods**). As resource complexity
increased, most strains became more suppressive (**Fig. 3B**, top; **Supplementary Fig.**
13). However, within this trend, we identified two distinct response types: mildly
influenced strains (189) and highly influenced strains (109), the latter showing a much
sharper gain in suppression under complex conditions (**Fig. 3B**, bottom;
Supplementary Figs. 14, 15). Both types were phylogenetically conserved,
suggesting that evolutionary constraints shape competitive strategies (**Fig. 3B**, top).
Notably, these response types were already distinguishable based on how strains
behaved in individual-resource environments (**Supplementary Figs. 16–17**). The
responses of highly and mildly influenced strains were strongly quantitative, scaling
with the number of available resources. Highly influenced strains started out less
suppressive than mildly influenced strains under simple conditions, but became more
suppressive as resource complexity increased (**Fig. 3C**). Strikingly, these strains also

lost the facilitative effects that were common in individual carbon sources (**Fig. 3B**, top;
Supplementary Fig. 15).

Given the increasing dominance of more competitive strains in our experiments
(**Fig. 2F**), we hypothesized that highly influenced strains would also become more
abundant in natural ecosystems as resource diversity increased. To test this and
extend our findings beyond the lab, we mapped metagenomic data from environmental
samples in the Earth Microbiome Project[15] (**Fig. 1B**) to the genomes of the 298 strains
tested in the sentinel assay (**Fig. 3D**). Because highly influenced strains show a
phylogenetic signal (**Fig. 3B**), their close relatives are expected to share similar traits.
Indeed, only reads mapping to highly influenced strains increased significantly with
resource diversity, unlike mildly influenced strains or unmapped reads (**Fig. 3E**).
These findings suggest that highly influenced strains not only become more
competitive in the lab, but also tend to dominate natural ecosystems as resource
complexity increases—reinforcing the ecological relevance of our experimental
results.

Figure 3: Resource complexity consistently increases microbial competitiveness, particularly

among a subset of strains A. We performed a high-throughput experiment using a luminescent

sentinel strain (*Pseudomonas aeruginosa PA01*) to quantify microbial interactions. Its growth was

measured alone and in co-culture with each of 298 partner strains across 16 levels of resource diversity

(1–16 resources), including 22 individual resources and 278 random mixtures (Supplementary Data 4).

This yielded 32,216 impact factor measurements quantifying each partner's effect across conditions. **B.**

K-means clustering (Supplementary Fig. 14) classified strains into two types based on their impact

factor response: highly influenced (109 strains) and mildly influenced (189 strains). Both types became

more suppressive on average with increased carbon complexity, but highly influenced strains showed a

much stronger shift. (**Top**) Heatmap shows $\log_{10}$(impact factor) across individual resources and

selected mixtures (gray = missing data). Strains are ordered by core-genome phylogeny. Right: each
strain's deviation in resource mixtures (≥2 resources) from its mean impact across individual resources.
black points represent each strain's average deviation. In total, 6,556 impact factor values were
calculated for individual resources (all shown in the heatmap) and 26,660 for resource mixtures
(heatmap and plot on the right). (**Bottom**) Histogram shows the distribution of mean $\log_{10}$(impact factor)
differences between individual and mixed resources (2–16 resources) for highly and mildly influenced
strains. The bimodal pattern reflects the distinct groupings, with dashed lines indicating each group's
mean. **C.** Highly influenced strains show a sharp decline in impact factor, indicating greater suppression,
with increasing resource complexity; mildly influenced strains shift only slightly. Lines show mean ±
SEM, with smoothed generalized additive model fit and 95% confidence intervals. **D.** Metagenomic
reads from 462 Earth Microbiome Project samples across 18 ecosystems were mapped to the full
genomes of the 298 tested strains, grouped by their response to resource complexity (highly or mildly
influenced; see panel B and Methods). Read counts per group were calculated and converted to relative
abundances. Illustration: a metagenomic read mapping to a highly influenced strain. **E.** The relative
abundance of highly influenced strains increases with resource complexity across all samples, in
contrast to mildly influenced or unmapped reads. Pearson correlation (r) and p-values are shown.

Emergent resources uptake raises microbial competitiveness

Microbes often interact by chemically modifying their environment—for instance, by
depleting shared resources. Therefore, we examined whether microbial alterations to
the growth medium—capturing their metabolic footprint—could reveal the mechanism
underlying the observed increase in competitiveness with resource diversity.
Specifically, we grew a subset of highly and mildly influenced strains in individual- and
mixed-resource settings, filtered the cultures, and collected supernatants. We then
compared true mixed-resource supernatants to their equivalent mixtures of single-
resource supernatants (illustrated in **Fig. 4A**). If metabolism were unaffected by
resource number, both would show similar chemical profiles. However, gas
chromatography–mass spectrometry (GC–MS) showed that residual key nutrients
were lower in true mixed-resource supernatants, indicating increased carbon uptake
with greater resource diversity (**Fig. 4C**; **Supplementary Fig. 18**)—also providing a
probable explanation for the stronger microbial growth observed in mixed resources
(**Fig. 2B**). This effect was particularly pronounced in highly influenced strains and
evident across both consumed and secreted metabolites (**Supplementary Fig. 19**).
To assess whether this enhanced uptake leads to stronger competition, we
compared sentinel strain growth in true mixed-resource supernatants to its growth in
corresponding mixtures of single-resource supernatants. In line with the GC–MS

results, the sentinel strain grew markedly less in true mixed-resource supernatants,
especially those from highly influenced strains and under higher resource diversity
(**Fig. 4C**). This reduction was not due to toxicity, as nutrient replenishment restored
growth in nearly all cases (**Supplementary Fig. 20**). Together, these results show that
highly influenced strains increase resource uptake as resource diversity rises, leaving
fewer nutrients available and creating a more competitive environment for other
microbes (illustrated in **Fig. 4D**).

To test whether this mechanism is sufficient to explain the observed biodiversity
loss (**Fig. 1D**), we incorporated increased resource uptake of highly influenced strains
into the classical Consumer–Resource Model (**Supplementary Text 3**; **Methods**).
Across simulations varying resource complexity and the proportion of highly influenced
strains, the modified model frequently predicted a decline in biodiversity with
increasing resource complexity—unlike the original model—with stronger effects as
the fraction of highly influenced strains increased (**Fig. 4E**; **Supplementary Figs. 21–**
22). This trend held across a wide range of uptake increases (**Supplementary Fig.**
23). Notably, even a small proportion of highly influenced strains was enough to limit
the expected biodiversity increase with resource diversity (compared to communities
without them), resulting in the dominance of these strains (**Supplementary Fig. 22**).
This outcome closely aligns with our experimental results, where the decrease in
diversity under high-resource conditions was largely driven by a single strain ('79'; **Fig.**
2A–C).

Collectively, our results show that a distinct subset of strains with synergistic carbon
uptake becomes increasingly competitive as resource complexity rises—ultimately
reducing microbial diversity. An emergent physiological trait at the individual level thus
drives the unexpected biodiversity loss at the community scale.

Figure 4: Emergent resource uptake with increasing resource complexity enhances microbial competitiveness. A. We tested whether highly influenced strains increase their resource uptake in mixtures beyond what is expected from independent uptake. Strains were grown in individual resources and in mixtures. Supernatants were then extracted, and individual supernatants were combined to create supernatant mixtures (see main text and Methods for details). Metabolic depletion was analyzed by GC-MS, and interaction effects were assessed via sentinel strain luminescence after 24 hours. **B.** Highly influenced strains show greater metabolite depletion in resource mixtures than in their corresponding supernatant mixtures, and this effect intensifies with increasing resource complexity. Shown is the relative change in metabolite peak area (normalized to an internal standard) for 96

identified metabolites, comparing each resource mixture to its matched supernatant mixture. Mann-
Whitney U tests assessed whether the median relative change deviated from 1. For mildly influenced
strains in two-resource conditions, values >1 indicate less depletion (i.e., lower uptake) in the resource
mixture. Significance: ns ($p > 0.05$), * ($p \leq 0.05$), ** ($p \leq 0.01$), *** ($p \leq 0.001$), **** ($p \leq 0.0001$). **C.** Highly
influenced strains suppress the sentinel strain more strongly in resource mixtures than in supernatant
mixtures, and this effect increases with resource complexity. Mildly influenced strains show the same
trend, but less pronounced. Shown is the change in sentinel strain luminescence after 24 h. Significance
levels as in panel B. **D.** Our results suggest that highly influenced strains increase carbon uptake as
resource diversity increases. For example, when a focal resource (A) is supplied alone, uptake is similar
between strain types. But in a mixture, only highly influenced strains show increased uptake of A. **E.**
Highly influenced strain behavior reduces microbial diversity as resource complexity increases in a
modified Consumer-Resource Model. We introduced a strain-specific parameter γ that increases
uptake affinity (i.e., reduces K_m) as resource diversity increases (equations in Supplementary Text 3).
We simulated communities of randomly generated strains across resource complexities using 21 highly
influenced strain fractions (0–100% in 5% steps; 20 replicates each). Shown is the slope (β_1) from a
linear model of Shannon diversity vs. resource number, plotted against the fraction of highly influenced
strains. The 0% case corresponds to the default Consumer-Resource Model (Fig. 1A). The trend line
shows the mean, ±SD. Results are robust for γ > 0.03 (γ = 0.05 shown; Supplementary Fig. 23), where γ
controls increased uptake in highly influenced strains (Supplementary Text 3).

Discussion

Our findings show that, contrary to ecological intuition and widely used models,
increasing resource diversity in microbial communities often reduces biodiversity.
Since biodiversity is closely linked to ecosystem resilience[17], this result has important
implications. One particularly striking example is in the context of dietary guidelines,
which often associate food variety with greater gut microbial richness[18]—yet our
results suggest that this strategy may backfire. More broadly, biodiversity loss due to
nutrient enrichment (e.g., fertilization) is frequently observed in forest[19,20], agricultural[21],
and aquatic ecosystems[22,23], though these cases typically involve increases in nutrient
quantity rather than diversity. These outcomes are typically explained by limiting-factor
trade-offs, such as shifts in competition toward light[23,24]. Our study isolated resource
diversity alone, keeping total nutrient concentration constant, and still observed
systematic diversity loss. This demonstrates that resource complexity itself can be
sufficient to drive competitive exclusion.

Conventional ecological models often ignore microbial physiology[4,25]. We show that
accounting for it can radically alter predicted outcomes and improve model accuracy,
as physiological shifts can cascade to reshape entire ecosystems. This finding is
surprising within an ecological framework, yet the underlying physiological pattern—
emergent changes in microbial resource uptake—is consistent with mounting
evidence: microbes exhibit markedly different behavior in mixed-resource
environments, including diauxic shifts and lag-phase dynamics[26], and distinct,
unexpected gene expression profiles[27]. While microbial physiology is undeniably
complex, we find that viewing it at a coarse-grained level reveals a surprisingly simple
and measurable link to competitive behavior—one that is sufficient to explain
community-level patterns. This echoes previous work showing that even rough
measures of cellular regulation can strongly predict bacterial growth rates[28]. Together,
this argues for tighter integration between microbial physiology and ecological
modeling. Doing so offers a more realistic view of how microbial ecosystems behave—
and why they sometimes defy expectations.

Finally, our finding reopens a fundamental question: how is microbial coexistence
maintained despite competition for the same resources? While recent theories offer
new perspectives on this paradox[8–12], it remains unclear whether existing frameworks
—alone or in combination—are sufficient to explain the diversity patterns observed in
nature, or whether additional, undiscovered mechanisms shape microbial ecosystems
in ways we have yet to uncover.

Methods

Media and Bacterial Culture

Bacteria were pre-cultured from frozen stocks in nutrient medium (NM) composed of
1% yeast extract and 1% soytone (both Sigma-Aldrich) for 24 h at 30 °C in shaking
(225 rpm). Following pre-culturing, cultures were diluted 1:5 in PBS and inoculated into
M9 medium supplemented with the relevant carbon sources (see **Supplementary**
Table 1 for full list and vendor details) and 1:100 diluted NM. Inoculation was
performed using an acoustic liquid handler (Echo 525, Beckman Coulter) by
transferring 25 nL of 1:5 diluted culture into 40 µL of media per well. For both
community and pairwise experiments (Figs. 1 and 2), cultures were passaged daily by
diluting 1:27 into fresh media (1.5 µL into 39 µL). Communities were sampled on days

9 and 10 (after 8 and 9 passages, respectively), and pairwise interactions were
sampled on day 5 (after 4 passages). All experiments were conducted in 384-well
plates. Community and pairwise experiments were carried out in transparent plates
with lids (VWR), while sentinel strain experiments were performed in black plates with
lids (Sarstedt, Lumox model) optimized for luminescence readouts.
403

Resources and their Mixtures

404
A total of 22 individual resources were used (**Supplementary Table 1**). Resources
were dispensed into 384-well plates containing M9 media without carbon sources
using an acoustic liquid handler (Echo 525, Beckman) prior to microbial inoculation.
For all resource conditions (individual or mixtures), a final concentration of 0.2%
(mass/volume) was maintained. Mixture conditions ranged from 2 to 16 resources and
were assigned randomly.
411

Strain library and whole-genome sequencing

412
Community and pairwise experiments
For all community and pairwise experiments, we used 14 bacterial strains with distinct
16S rRNA gene sequences, enabling unambiguous taxonomic identification (strain
names and 16S sequences listed in **Supplementary Data 2**). These strains are part of
a *Caenorhabditis elegans*-associated isolate collection of the Ratzke lab and
represent a subset of the 16 strains previously used in a similar context[29].
419

420 *Sentinel strain experiments*
421 For the sentinel strain experiments, we used 298 bacterial strains isolated from the gut
422 of *Caenorhabditis elegans* (**Supplementary Data 5**). Strains were provided by Dr.
423 Christoph Ratzke and Prof. Hinrich Schulenburg.
424

Bacterial whole-genome sequencing
Of the 298 strains used in this study, 92 had been previously sequenced and
assembled (**Supplementary Data 5**; courtesy of Prof. Hinrich Schulenburg). The
remaining 206 strains were sequenced as part of this work. Genomic DNA was
extracted using the peqGOLD Bacteria DNA Kit (VWR). Whole-genome libraries were
prepared with the NEBNext Ultra II FS DNA Library Prep Kit for Illumina (New England
Biolabs), using dual index oligos for barcoding. To reduce reagent use and cost, the

protocol was miniaturized to 1:7 of the standard volume. This scaled-down protocol
was benchmarked against the standard 1:1 protocol and produced comparable
results. Libraries were sequenced on an Illumina NovaSeq platform to a mean depth of
~120×. Raw reads were trimmed using fastp (v0.23.2)[30] and assembled de novo with
SPAdes (v3.15.5)[30] using the --isolate flag and default error correction. Assemblies
were generated using 24 threads per sample. Assembly quality was assessed with
QUAST (v5.2)[32], and metrics were summarized across strains. Contamination was
evaluated at the contig level using DIAMOND[33] followed by MEGAN[34] with GTDB
taxonomy. Genomes were excluded if ≥10% of contigs were assigned to taxa
inconsistent with the expected lineage, based on contradictions at or above the genus
level. A core genome phylogeny of the strain collection was constructed using panX[35]
(default bacterial parameters, 95% core gene threshold). Sequencing statistics in
Supplementary Data 6.
445

Luciferase tagging of the sentinel strain
Pseudomonas aeruginosa PA01 ('PA01') was genomically transformed with the
luxCDABE operon using the mini-Tn7 insertion system as previously described106.
Briefly, the plasmids pUC18T-mini-Tn7T-Gm-lux (Addgene #64953) and pTNS2
(Addgene #64968) were curated from E.coli DH5a using a plasmid extraction kit
(peqGOLD Plasmid MiniPrep Kit II; VWR). PA01 cells were made competent for
transformation by treatment with a sucrose 300 mM solution PA01, and were then
electroporated with 50 ng of the curated pUC18T-mini-Tn7T-Gm-lux and pTNS2
plasmids. After recovery in 1 ml of LB at 30 °C, the electroporated cells were plated on
an LB-agar+Gm30 plate. Plates were incubated for 24 hours at 30 °C, and colonies
were tested for their bioluminescence activity using a plate reader. Positive colonies
were then further grown in a selective medium (LB+Gm30), and were stocked in 25%
glycerol at -80 °C.
459

460 **Community composition based on 16S rRNA gene sequencing**
461 Amplicon-based 16S rRNA gene sequencing was used to profile bacterial community
462 composition in pairwise and multispecies experiments.
463

DNA extraction

Genomic DNA was extracted using a miniaturized protocol in 384-well plates. Pellets
were resuspended in TES buffer, and 10 µL was transferred to fresh PCR plates. Lysis
was initiated by adding metapolyzyme (in PBS) and ReadyLyse (Lucigen, 1:5 in TES),
dispensed via an ECHO acoustic liquid handler, followed by overnight incubation at
30 °C with shaking (1,350 rpm). The next day, Proteinase K (New England Biolabs, 1:2
dilution of 20 mg/mL stock) and 20% SDS (1% final) were added, followed by
incubation at 55 °C for 30 min. RNase A (Omega Bio-Tek, 1:10 dilution) was added
and incubated for 5 min at room temperature. Proteinase K was inactivated at 95 °C
for 5 min. Samples were centrifuged (4,000 rpm, 10 min), and supernatants were
collected. Lysates were diluted 1:6 and then 1:3 in nuclease-free water. DNA
concentrations were measured using the PicoGreen assay, and 0.1–10 ng was used
as template for 16S amplification.

16S library preparation and sequencing
Full-length 16S rRNA genes were amplified using a barcoded PCR protocol with
primers 27F and 1492R (barcode sequences listed in **Supplementary Data 7**),
followed by sequencing on an Oxford Nanopore platform. PCR-grade water and 2×
Hot-Start PCR Master Mix (Biotechrabbit) were combined in ECHO-qualified plates,
and 4.4 µL of reaction mix was dispensed into 384-well PCR plates using an ECHO
acoustic liquid handler (Beckman Coulter). After centrifugation, 50 nL of barcode-
specific primers and 500 nL of genomic DNA were added to each well using the same
system. Thermocycling was performed as follows: 94 °C for 5 min; 29 cycles of 94 °C
for 60 s, 55 °C for 60 s, and 72 °C for 2.5 min; followed by 72 °C for 10 min and a final
hold at 4 °C. Barcoded amplicons were pooled per tube, and libraries were prepared
using Oxford Nanopore's short-read amplicon workflow according to the
manufacturer's protocol, with halved reaction and bead-cleanup volumes. Sequencing
was performed on a MinION device (Oxford Nanopore Technologies).

Read mapping and relative abundance estimation
Raw Nanopore reads were split into 100 parts using seqkit[36] and filtered by quality
(Q ≥ 15) and length (1,500–1,700 bp) using NanoFilt (v2.8.0)[37]. Demultiplexing was
performed with a custom barcode set using minibar (https://github.com/calacademy-
research/minibar), allowing edit distances up to 6 bases. Reverse complementation,
where needed, was performed using a custom Python script

(rev_cmplt_minibar_output.py). Strain-specific 16S consensus sequences were
generated by pooling reads across replicates, filtering (Q ≥ 18), and aligning the top
1,000 reads per strain with MAFFT (v7.508)[38]. Consensus sequences were computed
using a custom script applying a 90% agreement threshold and IUPAC ambiguity
codes, and then combined into a FASTA file to serve as a custom reference database.
Chimeric reads were removed using vsearch (v2.21.1)[39] with --uchime_ref against this
reference. Non-chimeric reads were mapped to the database using minimap2
(v2.26)[40] with -ax map-ont. Alignment files were sorted and indexed using samtools
(v1.17)[41], and per-sample read counts were generated using samtools idxstats. The
final count matrix was compiled with a custom script (make_count_table_updated.py).
All custom scripts and workflow code are available at
https://github.com/orshalevsk/Emergent_Biodiv_loss.
511

512 **OD600 and bioluminescence measurement**
513 *OD600*
514 A 384-well plate containing bacterial cultures was mixed for 15 s at 2,000 rpm using a
515 Mixmate plate shaker (Eppendorf), then transferred to a FLUOstar Omega plate
516 reader (BMG Labtech) to measure absorbance at 600 nm.
517

Bioluminescence
Bacterial cultures were transferred to a Lumox 384-well plate (Sarstedt) for
bioluminescence measurement. First, 2.5 µL of NM solution was dispensed into each
well. Then, 20 µL of bacterial culture was reverse-dispensed into the same wells using
a VIAFLO 384 liquid handler (Integra) and mixed with NM for 30 s at 1,500 rpm using a
Mixmate plate shaker (Eppendorf). The Lumox plate was incubated at 30 °C in a
FLUOstar Omega plate reader (BMG Labtech) for 2 min to allow NM to stimulate
bacterial metabolism (notably in *Pseudomonas aeruginosa*), enhancing
bioluminescence signal consistency. Bioluminescence was then recorded using the
plate reader's luminescence emission filter, with a gain setting of 3,600 and an
exposure time of 1 s.
529

Experimental design
Communities, pairwise and individual strains experiments

Pre-cultures were prepared as described above (Media and Bacterial Culture), then
dispensed into 384-well plates containing M9 media with the appropriate carbon
sources and 1:100 NM. Metadata for all conditions and strain combinations used in the
community experiment are provided in **Supplementary Data 1**; metadata for pairwise
and single-strain experiments are provided in **Supplementary Data 3**. All
experimental conditions were fully randomized across wells and plates. At the end of
each experiment, bacterial density (OD_{600}) was measured for all wells, including
individual strains, pairwise combinations, and communities. Only pairwise and
community samples were further processed for 16S sequencing: samples were
centrifuged at 4,000 rpm for 5 min, supernatants were removed, and pellets were
stored at –80 °C. DNA extraction and 16S rRNA gene sequencing procedures are
described below (DNA extraction and 16S sequencing). Individual strain growth was
measured after 72 hours (carrying capacity, OD_{600}), pairwise cultures were serially
passaged for five days, followed by a final 24-hour incubation before OD_{600}
measurement and 16S profiling and communities were serially passaged for 9 days,
followed by a final 24-hour incubation before OD_{600} measurement.

Sentinel strain experiments
The full experimental setup for sentinel strain co-cultures is described above (see
Media and Bacterial Culture and Resources). Briefly, the sentinel strain and 298 co-
cultured strains were grown overnight in NM medium and diluted 1:5 in PBS (without
washing). Cultures were then dispensed into 384-well plates containing the
appropriate media and resource conditions. Six experimental batches were
performed: three using individual resources (biological replicates), and three using
distinct mixed-resource conditions (no replicates). In the individual-resource
experiments, all 298 strains were co-incubated with the sentinel strain across 22
single-resource conditions. The sentinel strain was also grown alone with 16 replicates
per condition. In the mixed-resource experiments, one batch included all 298 strains
co-incubated with the sentinel strain in all conditions. In the other two batches, 80
strains were randomly selected per condition to maximize the number of unique
resource combinations tested. In these batches, the sentinel strain alone was included
with 8 replicates per condition. All conditions were fully randomized across wells and
plates, with each experiment distributed over 20–24 separate 384-well plates per
batch. Plates were incubated at 30 °C for 24 h before luminescence was measured.

The average luminescence of the sentinel strain grown alone in each condition was
used to calculate the impact factor of co-cultured strains, as shown in Figs. 2A and 3A.
Impact factor values for individual resources represent the average across three
replicate experiments ('5', '6', and '7'). Full experimental conditions and metadata are
provided in **Supplementary Data 4**.

Supernatant experiments

The full setup for supernatant experiments is described above (see Media and
Bacterial Culture and Resources). Briefly, selected highly influenced and mildly
influenced strains were grown overnight in NM medium and diluted 1:5 in PBS (without
washing). Cultures were dispensed into 384-well plates containing M9 medium
supplemented with either individual or mixed carbon sources, with each strain
inoculated alone into four adjacent wells per condition (e.g., A1, A2, B1, B2) to ensure
sufficient volume for supernatant extraction. Plates were incubated at 30 °C for 24 h.
To extract supernatants, technical replicates (four wells per condition) were pooled into
96-well plates using a VIAFLO 384 liquid handler (Integra). Each set of four wells
(40 µL per well) yielded a combined volume of 160 µL. The pooled cultures were
filtered through 0.2 µm filter plates (PALL) into a new sterile 96-well plate, removing
cells and leaving only the cell-free supernatant. This design enabled combinatorial
mixing of supernatants to reconstruct mixed-carbon environments. Mixing was
performed using a VIAFLO 96 liquid handler by combining supernatants from wells on
different plates (e.g., A1 from plate 1 with carbon A and A1 from plate 2 with carbon B
were combined 1:1 to generate A+B mixtures). For the sentinel growth assay (Fig. 4C),
eight strains were randomly selected—four highly influenced and four mildly
influenced—were tested across 66 mixed-resource conditions. For the mass
spectrometry experiment (Fig. 4B), a subset of four of these strains (two highly
influenced, two mildly influenced) was tested across nine mixed-resource conditions.
Full strain lists and conditions are provided in **Supplementary Data 8**.

GC−MS sample preparation and analysis

A total of 72 samples (20 µL each) were thawed on ice and mixed with 50 µL of internal
standard solution containing 60 µM $13C_6$-glucose and 60 µM 3-hydroxybenzoic acid
(equivalent to 3,000 pmol each). Samples were dried overnight using a vacuum
concentrator (Eppendorf Concentrator) to remove water, then placed in a desiccator

over phosphorus pentoxide for 1 h. Derivatization was performed following a protocol adapted from Fiehn et al[42]. Briefly, 50 μL of methoxylamine hydrochloride in pyridine (20 mg/mL, freshly prepared) was added to each sample. Samples were incubated in an ultrasonic bath for 10 min, followed by shaking at 30 °C for 90 min at 1,400 rpm. After a brief centrifugation, 70 μL of MSTFA (N-methyl-N-trimethylsilyltrifluoroacetamide; Sigma-Aldrich) was added. Samples were then incubated at 40 °C for 60 min at 1,200 rpm and left at room temperature for an additional 2 h. Final centrifugation was performed at 13,000 rpm for 10 min at 4 °C (Hettich tabletop centrifuge, swing-out rotor), and 60 μL of supernatant was transferred to GC vials for injection. GC-MS analysis was carried out on a Shimadzu TQ 8040 triple quadrupole mass spectrometer coupled to a high-performance liquid chromatography system. Chromatographic separation was achieved on a Restek Rxi-5Sil MS column. Detailed GC and MS parameters are provided in **Supplementary Data 9**. Data acquisition and peak integration were performed using LabSolutions Insight GCMS software, with manual curation of peak boundaries.

Earth Microbiome Project data handling

Alpha diversity vs. resource complexity analysis

Data were retrieved from the Earth Microbiome Project (EMP) study by Shaffer et al.[15] via Qiita (study ID 13114: https://qiita.ucsd.edu/study/description/13114). We used both 16S-based alpha diversity and predicted microbially derived metabolite richness, as previously computed in Shaffer et al.[15] (Fig. 2C; Supplementary Figs. 1C–D). Alpha diversity was taken from the column alpha_16s_deblur_nosingletons_rar5k_shannon, which reports Shannon diversity after singleton removal and rarefaction to 5,000 sequences per sample. This depth was identified by the authors as optimal for cross-sample comparison. Resource diversity was taken from alpha_lcms_fbmn_microbial_richness, defined as the number of HPLC–MS peaks >0 in each sample following Feature-Based Molecular Networking (FBMN), and interpreted as microbially associated metabolite richness. Samples were filtered to include only those with non-missing alpha diversity values (477 of 618 total). To ensure statistical robustness, we retained only environments (as defined by EMP level "empo_4") with ≥6 samples, resulting in 462 samples across 13 distinct ecosystems. We used the "empo_4" level of environmental classification, which provides the most specific ecosystem labels available in the EMP dataset and avoids confounding that

634 can arise at broader levels (e.g., "empo_1"), where ecologically distinct environments
635 —such as animal guts and plants—may be grouped together.
636
Mapping metagenomic reads to highly and mildly influenced strains and abundance
estimation
To assess the environmental representation of highly and mildly influenced strains, we
mapped metagenomic shotgun reads from the Earth Microbiome Project[15] to our full
genome collection of 298 experimentally characterized strains. Genome assemblies
were merged into a single reference database, and a minimap2 index (v2.26)[40] was
created. Reads were aligned using minimap2 with the -ax sr --secondary=no
parameters to exclude secondary alignments (pipeline:
minimap2_pipline_no_multi_match.sh). Alignment files (SAM) were processed using a
custom script (make_count_table_updated_with_mistmatch_filter_db.py) to filter for
high-quality matches (--min_mapq 50) and compute per-strain read counts, along with
unmapped reads (**Supplementary Data 10**). To ensure sufficient coverage and
minimize noise, only samples with a total mapped read count ≥10,000 were retained.
This filtering yielded 453 metagenomes spanning 18 distinct ecosystems. For each
sample, raw counts were normalized to relative abundances. Strains were categorized
as highly or mildly influenced based on their experimentally measured impact on
biodiversity, and reads were binned accordingly into highly influenced, mildly
influenced, or unmapped groups. Relative abundances for each category were then
calculated (**Supplementary Data 11**). These values were integrated with matched
metabolomic profiles from the Earth Microbiome Project[15] for downstream ecological
analyses. All custom scripts used in this workflow are available at
https://github.com/orshalevsk/Emergent_Biodiv_loss.
659
Consumer-Resource model
For the classic Consumer-Resource model see **Supplementary Text 1**. For the highly
influenced strains modified version see **Supplementary Text 3**. For the highly
influenced strains version, we simulated 420 communities of 50 randomly generated
strains across seven resource complexities (1–64; **Fig. 4E**; **Supplementary Figs. 21-**
23).
666

667 **Inferring interaction matrices from experimental data using the generalized**
668 **Lotka–Volterra model**
669 See **Supplementary Text 2**. Note that error bars for α_{self} in Figure 2B (middle) may be
670 overestimated as the interaction values are not independent (see Supplementary Fig.
671 6A).
672
673
Estimating single-species Shapley contributions to biodiversity loss
We used the interaction matrices obtained from experimental data for 1 and 16 carbon
sources (methodology described in **Supplementary Text 2**; results in
Supplementary Fig. 4) and added Gaussian noise (mean = 0, variance = 0.7) to each
entry to generate a synthetic set of 14 species resembling the 14 experimentally
measured strains. We produced in total 800 such sets of 14 species and took from
each set randomly six or eight species respectively (400x six species and 400x eight
species). We then simulated all possible subcommunities of these 6 or 8 species using
the generalized Lotka–Volterra model under both 1- and 16-resource conditions. Since
the interaction matrices were obtained from steady state data, we could not estimate
the single species' *per capita* growth rates and set them to be one instead. Since we
are only interested in steady state outcomes which are not impacted by the *per capita*
growth rates only ensuring that the systems reached steady state was sufficient for
further analysis (**Supplementary Fig. 8**). For each simulated subcommunity, we
computed the steady-state Shannon diversity index under both conditions and
calculated the change in diversity as: $\Delta Shannon = Shannon_{16} - Shannon_1$. We then
computed Shapley values, representing the contribution of each strain in a six- or eight-
species community to the overall change in Shannon diversity between 1 and 16
resources. Next, using the same interaction matrices (**Supplementary Fig. 4**), we
calculated each species' mean impact factor on the other members of its community
under both resource conditions—analogous to the analysis shown in Fig. 2D. Finally,
we computed the change in mean impact factor with increasing resource diversity and
plotted these values against the corresponding Shapley values for each species (**Fig.**
2C).
698
699 **Code availability**

700 The code used in this study is available at
701 https://github.com/orshalevsk/Emergent_Biodiv_loss.
702
703 **Data availability**
704 All sequencing data will be deposited in the European Nucleotide Archive (ENA) prior
705 to publication and will be made available upon request during peer review.
706

References

1. Gans, J., Wolinsky, M. & Dunbar, J. Computational Improvements Reveal Great Bacterial

Diversity and High Metal Toxicity in Soil. *Science* **309**, 1387–1390 (2005).

2. Georgy, G. & Georgii, F. *The Struggle For Existence*. (Baltimore: Williams & Wilkins, 1934).

3. Volterra, V. Variations and Fluctuations of the Number of Individuals in Animal Species living

together. *ICES J. Mar. Sci.* **3**, 3–51 (1928).

4. Lotka, A. J. Elements of Physical Biology. *Sci. Prog. Twent. Century 1919-1933* **21**, 341–343

(1926).

5. Futuyma, D. J. & Moreno, G. THE EVOLUTION OF ECOLOGICAL SPECIALIZATION. *Annu. Rev. Ecol.*

Evol. Syst. **19**, 207–233 (1988).

6. Hutchinson, G. E. Homage to Santa Rosalia or Why Are There So Many Kinds of Animals? *Am.*

Nat. **93**, 145–159 (1959).

7. Macarthur, R. & Levins, R. The Limiting Similarity, Convergence, and Divergence of Coexisting

Species. *Am. Nat.* **101**, 377–385 (1967).

8. Jetschke, G. The Unified Neutral Theory of Biodiversity and Biogeography. *Ecology* **83**, 1771–

1772 (2002).

9. Thingstad, T. F. Elements of a theory for the mechanisms controlling abundance, diversity, and

biogeochemical role of lytic bacterial viruses in aquatic systems. *Limnol. Oceanogr.* **45**, 1320–

1328 (2000).

10. Morris, J. J., Lenski, R. E. & Zinser, E. R. The Black Queen Hypothesis: Evolution of Dependencies

through Adaptive Gene Loss. *mBio* **3**, e00036-12 (2012).

11. Chesson, P. L. & Warner, R. R. Environmental Variability Promotes Coexistence in Lottery

Competitive Systems. *Am. Nat.* **117**, 923–943 (1981).

12. Chesson, P. Multispecies Competition in Variable Environments. *Theor. Popul. Biol.* **45**, 227–276

(1994).

13. Dal Bello, M., Lee, H., Goyal, A. & Gore, J. Resource–diversity relationships in bacterial

communities reflect the network structure of microbial metabolism. *Nat. Ecol. Evol.* **5**, 1424–

1434 (2021).

14. Pacheco, A. R., Osborne, M. L. & Segrè, D. Non-additive microbial community responses to

environmental complexity. *Nat. Commun.* **12**, 2365 (2021).

15. Shaffer, J. P. *et al.* Standardized multi-omics of Earth's microbiomes reveals microbial and

metabolite diversity. *Nat. Microbiol.* **7**, 2128–2150 (2022).

16. Hofbauer, J. & Sigmund, K. *Evolutionary Games and Population Dynamics*. (Cambridge University

Press, Cambridge, 1998). doi:10.1017/CBO9781139173179.

17. Tilman, D., Reich, P. B. & Knops, J. M. H. Biodiversity and ecosystem stability in a decade-long

grassland experiment. *Nature* **441**, 629–632 (2006).

18. Xu, Z. & Knight, R. Dietary effects on human gut microbiome diversity. *Br. J. Nutr.* **113 Suppl**, S1-

5 (2015).

19. Meunier, C. L., Gundale, M. J., Sánchez, I. S. & Liess, A. Impact of nitrogen deposition on forest

and lake food webs in nitrogen-limited environments. *Glob. Change Biol.* **22**, 164–179 (2016).

20. Ceulemans, T., Merckx, R., Hens, M. & Honnay, O. Plant species loss from European semi-natural

grasslands following nutrient enrichment – is it nitrogen or is it phosphorus? *Glob. Ecol.*

Biogeogr. **22**, 73–82 (2013).

21. Piggott, J. J., Lange, K., Townsend, C. R. & Matthaei, C. D. Multiple Stressors in Agricultural

Streams: A Mesocosm Study of Interactions among Raised Water Temperature, Sediment

Addition and Nutrient Enrichment. *PLOS ONE* **7**, e49873 (2012).

22. *Eutrophication: Causes, Consequences and Control: Volume 2*. (Springer Netherlands, Dordrecht,

754   2014). doi:10.1007/978-94-007-7814-6.

755   23. Hautier, Y., Niklaus, P. A. & Hector, A. Competition for Light Causes Plant Biodiversity Loss After

756   Eutrophication. *Science* **324**, 636–638 (2009).

757   24. Borer, E. T. *et al*. Herbivores and nutrients control grassland plant diversity via light limitation.

758   *Nature* **508**, 517–520 (2014).

759   25. MacArthur, R. Species packing and competitive equilibrium for many species. *Theor. Popul. Biol*.

760   **1**, 1–11 (1970).

761   26. Basan, M. *et al*. A universal trade-off between growth and lag in fluctuating environments.

762   *Nature* **584**, 470–474 (2020).

763   27. New, A. M. *et al*. Different Levels of Catabolite Repression Optimize Growth in Stable and

764   Variable Environments. *PLOS Biol*. **12**, e1001764 (2014).

765   28. Wu, C. *et al*. Cellular perception of growth rate and the mechanistic origin of bacterial growth

766   law. *Proc. Natl. Acad. Sci*. **119**, e2201585119 (2022).

767   29. Ye, X., Shalev, O. & Ratzke, C. Biotic resistance predictably shifts microbial invasion regimes. *Nat.*

768   *Commun*. **16**, 3952 (2025).

769   30. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

770   *Bioinformatics* **34**, i884–i890 (2018).

771   31. Bankevich, A. *et al*. SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-

772   Cell Sequencing. *J. Comput. Biol*. **19**, 455–477 (2012).

773   32. Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool for genome

774   assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

775   33. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND. *Nat.*

776   *Methods* **12**, 59–60 (2015).

777   34. Huson, D. H., Auch, A. F., Qi, J. & Schuster, S. C. MEGAN analysis of metagenomic data. *Genome*

778   *Res*. **17**, 377–386 (2007).

35. Ding, W., Baumdicker, F. & Neher, R. A. panX: pan-genome analysis and exploration. *Nucleic*

Acids Res. **46**, e5 (2018).

36. Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File

Manipulation. *PLOS ONE* **11**, e0163962 (2016).

37. De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M. & Van Broeckhoven, C. NanoPack: visualizing

and processing long-read sequencing data. *Bioinformatics* **34**, 2666–2669 (2018).

38. Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid multiple sequence

alignment based on fast Fourier transform. *Nucleic Acids Res.* **30**, 3059–3066 (2002).

39. Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. VSEARCH: a versatile open source tool

for metagenomics. *PeerJ* **4**, e2584 (2016).

40. Li, H. New strategies to improve minimap2 alignment accuracy. *Bioinformatics* **37**, 4572–4574

(2021).

41. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079

(2009).

42. Fiehn, O. Metabolomics by Gas Chromatography-Mass Spectrometry: the combination of

targeted and untargeted profiling. *Curr. Protoc. Mol. Biol. Ed. Frederick M Ausubel Al* **114**,

30.4.1-30.4.32 (2016).

Acknowledgements

C.R. received funding from the European Research Council (ERC) under the

European Union's Horizon 2020 research and innovation programme (grant

agreement No. 948753), the Deutsche Forschungsgemeinschaft (DFG, German

Research Foundation) grants 468972576 and 540605007, and the Cluster of

Excellence EXC 2124 "Controlling Microbes to Fight Infections" (CMFI). O.S. received

funding from the DFG (grant 516931136). We thank Hinrich Schulenburg for

generously providing a C. elegans strain library, and Andreas Peschel for providing the

original Pseudomonas aeruginosa PA01 strain. We also thank the entire Ratzke lab for

constructive feedback on the project and the manuscript. The bacterial illustrations in
the figures were created by Bala Akaba (balaakaba@gmail.com).
809

Author Contributions Statement

O.S. Conceived the study, led the conceptualization, performed all experiments, data
analysis, and coding, generated all figures except where noted, and was responsible
for writing—original draft and review & editing. C.R. Contributed to conceptualization,
developed the generalized Lotka–Volterra modeling framework (including simulations
and Bayesian inference), and contributed to writing—review & editing. X.Y.
Contributed to conceptualization, extended the Consumer–Resource model to include
highly influenced strains, performed related simulations, and contributed to writing—
review & editing. J.K. and M.S. Conducted the GC–MS measurements and performed
metabolite data analysis.

Competing Interests Statement

The authors declare no competing interests.

825