Quantitative RNA pseudouridine landscape reveals dynamic modification patterns and

evolutionary conservation across bacterial species

Letong Xu[1,#], Shenghai Shen[2,#], Yizhou Zhang[1], Zhihao Guo[4], Beifang Lu[1], Jiadai Huang[1], Runsheng

Li[5,6], Yitong Shen[2], Li-Sheng Zhang[2,3,*] and Xin Deng[1,4,5,*]

[1] Department of Biomedical Sciences, City University of Hong Kong, Kowloon Tong, Hong

Kong SAR, China

[2] Division of Life Science, The Hong Kong University of Science and Technology, Kowloon,

Hong Kong SAR, China.

[3] Department of Chemistry, The Hong Kong University of Science and Technology, Kowloon,

Hong Kong SAR, China.

[4] Shenzhen Research Institute, City University of Hong Kong, Shenzhen, Guangdong 518057,

China

[5] Tung Biomedical Sciences Center, City University of Hong Kong, Hong Kong, China

[6] Department of Infectious Diseases and Public Health, City University of Hong Kong,

Kowloon Tong, Hong Kong SAR, China

* Correspondence: xindeng@cityu.edu.hk (Xin Deng) and zhangls@ust.hk (Li-Sheng Zhang)

[#] These authors contributed equally to the paper as the first authors.

Abstract

Pseudouridine (Ψ) modifications are the most abundant RNA modifications; however, their distribution and functional significance in bacteria remain largely unexplored compared to eukaryotic systems. In this study, we present the first transcriptome-wide and quantitative mapping of Ψ modifications across five diverse bacterial species (*Bacillus cereus*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Pseudomonas syringae*) at single-base resolution, utilizing the optimized BID-seq method for bacterial RNA. Our analysis revealed growth phase-dependent dynamics of pseudouridylation in bacterial tRNA and mRNA, particularly in genes enriched in core metabolic pathways. Comparative analysis demonstrated evolutionarily conserved features of Ψ modifications, such as dominant motif contexts, Ψ clustering within operons, etc. Functional analysis indicated Ψ modifications influence bacterial mRNA stability, translation, and interactions with specific RNA-binding proteins (RBPs) in response to changing cellular demands during growth phase transitions. The integrated computational analysis on local RNA architecture was conducted to elucidate the structure-dependent Ψ modifications in bacterial RNA. Furthermore, we developed an integrated deep learning framework, combining Transformer-GNN-based neural networks (pseU_NN) to capture both RNA sequence and structural features for effective prediction of Ψ-modified sites. Overall, our study provides valuable insights into the landscapes of bacterial RNA Ψ modifications and establishes a foundation for future mechanistic investigations into the functions of Ψ in bacterial RNA regulation.

Keywords: Bacteria, RNA modification, Microbiology, Bioinformatics, Pseudouridine.

Introduction

RNA modifications are a crucial layer of post-transcriptional regulation in biological systems, with approximately 170 distinct chemical modifications identified to date[1]. Pseudouridine (Ψ), often referred to as the "fifth nucleoside," is one of the most prevalent and evolutionarily conserved RNA modifications[2,3]. This unique modification arises from a specific isomerization process in which uridine undergoes a site-specific intramolecular rearrangement[4]. Ψ can thermodynamically stabilize RNA structures by enhancing base stacking and increasing the rigidity of the sugar-phosphate backbone, which helps maintain the structural folding of functional RNAs such as tRNA and rRNA[5,6]. For example, Ψ at the 55[th] position of tRNA, a universally conserved modification in both eukaryotes and prokaryotes, is crucial for regulating tRNA stability and aminoacylation levels[7,8]. Ψ modifications in rRNA play important roles in rRNA biogenesis and function, as well as in mRNA translation in both mammalian cells and bacteria[9–11].

The regulation of mRNA translation is highly complex, and mRNA stability significantly influences gene expression in humans. Ψ-modified mRNAs demonstrate enhanced stability due to their resistance to RNase L-mediated degradation[12]. Pre-mRNA is found to be pseudouridylated co-transcriptionally, with Ψ enriched near alternative splicing regions and RNA-binding protein (RBP) binding sites[13]. Moreover, Ψ located within exon regions can alter codon properties to modulate translation, while Ψ modifications at stop codons promote ribosomal readthrough[14–16]. Ψ also facilitates the low-level synthesis of peptide products from individual mRNA sequences in human cells and increases the rate at which near-cognate tRNA^{Val} interacts with ΨUU codons[17], suggesting a more complex regulatory role for Ψ in translation.

Until recently, comprehensive investigations of bacterial RNA pseudouridylation have been limited due to technical challenges in precisely mapping and quantifying Ψ modifications at single-nucleotide resolution. While eukaryotic mRNA can be easily isolated and assessed using poly-A selection methods, bacterial transcripts lack poly-A tails and are predominantly

composed of ribosomal RNA, which accounts for over 95% of total RNA[18]. This methodological gap has hindered the functional exploration of Ψ roles in bacterial RNA. A previously reported CMC-based method can selectively label Ψ sites and produce truncation signatures during reverse transcription. However, this method often exhibits several drawbacks, including relatively low sensitivity and limitations in quantifying Ψ modification levels. Recently, a new technique named Bisulfite-induced deletion sequencing (BID-seq) has emerged, utilizing unique deletion signatures induced at Ψ-modifed sites to achieve base-resolution and quantitative characterization of Ψ sites across the transcriptome[19,20].

To address these challenges and conduct an in-depth study of bacterial Ψ modifications, we developed an optimized BID-seq method for bacterial RNA, termed baBID-seq. By combining efficient rRNA depletion in baBID-seq, we expanded our quantitative Ψ analysis to four representative bacterial species: *K. pneumoniae*, *B. cereus*, *P. aeruginosa*, and *P. syringae*, examining both exponential and stationary growth phases. Our comprehensive analysis revealed the landscape of Ψ modifications on bacterial rRNA, tRNA, and mRNA, highlighting evolutionarily conserved Ψ features across bacterial strains. We investigated the sequence and structural properties of local mRNA regions that influence Ψ deposition. Dynamic Ψ modifications at specific sites in tRNA and mRNA were observed, showing distinct accumulation patterns during the stationary growth phase. In *P. syringae*, we explored the roles of Ψ under nutrient-deficient conditions and found a positive correlation between mRNA translation efficiency (TE) and Ψ intensity[21]. In *P. aeruginosa*, we provided evidence for the potential regulatory functions of Ψ in promoting mRNA interactions with the Hfq chaperone [22]. Furthermore, we employed an attention-based graph neural network (GNN) classification approach, integrating RNA sequence and local structural features to predict potential Ψ modification sites. Collectively, our systematic analysis revealed a novel and dynamic landscape of Ψ modifications, uncovering evolutionarily conserved modification features and key motifs and structural elements within mRNA regions that impact Ψ modification.

Results

Optimized BID-seq quantitatively maps Ψ modification in bacterial rRNA and tRNA

To investigate Ψ modifications in bacteria, we primarily applied the standard BID-seq

protocol[20] to total RNA isolated from *E. coli* and *P. aeruginosa* during exponential and

stationary growth phases. Ψ sites on rRNA were identified through deletion signatures at

single-base resolution, and the observed deletion ratios were utilized to assess Ψ modification

fraction. By characterizing Ψ sites with significantly higher deletion ratios in the 'BID-seq

treated' samples compared to the 'input' samples, we detected 9 out of 10 known Ψ sites, as

well as one additional Ψ site, that were significantly modified in 23S and 16S rRNA,

respectively (**Fig. 1a,b**). For example, Ψ781 in 16S rRNA of *P. aeruginosa* exhibited a

distinct deletion signature in the 'BID-seq treated' samples (**Supplementary Fig. 1c**).

Although Ψ is known to play a critical role in regulating rRNA local structure and

biogenesis[11], bacterial RNA BID-seq (baBID-seq) pointed out that the number of Ψ-modified

sites is noticeably lower than the typical >100 Ψ sites found in mammalian rRNA[19]. We then

compared the rRNA Ψ fraction in *E. coli* and *P. aeruginosa* across the two growth stages, and

almost all Ψ sites are highly conserved, with only slight variations in Ψ site deletion ratio

(**Fig. 1a,b**).

To study Ψ modifications on bacterial RNA species beyond rRNA, we employed probe-based

rRNA depletion[23] as a critical step in our optimized protocol for baBID-seq. We carefully

established RNA fragmentation conditions to generate fragmented RNA of ~60–70 bp in

length, ensuring efficient adaptor ligation; meanwhile, size selection of amplified library

DNA by native PAGE gel effectively minimized contamination from adaptor dimers or

dsDNA of unexpected sizes (**Supplementary Fig. 1a**). We then applied baBID-seq protocol

to four bacterial species*: K. pneumoniae*, *B. cereus*, *P. aeruginosa*, and *P. syringae*. Despite

the rRNA depletion in baBID-seq, the Ψ sites in the remaining 16S rRNA of these four

strains exhibited conserved modification fraction, serving as benchmarks for assessing

baBID-seq library quality when stimulating deletion signatures (**Supplementary Fig. 1b**).

baBID-seq successfully captured various RNA species, including rRNA, tRNA, and mRNA.

We characterized hundreds of Ψ sites on rRNA-depleted RNA isolated from four bacterial

strains across exponential and stationary growth phases. The results from baBID-seq

quantitatively demonstrated an excellent correlation between biological replicates (**Fig. 1c**

and **Supplementary Fig. 2a**). While Ψ sites on rRNA and tRNA consistently showed stable

modification levels across biological replicates, certain mRNA Ψ sites displayed greater

variability.

Given the conservation of tRNA Ψ sites across biological replicates, we used the average Ψ

fraction at each specific tRNA Ψ site for downstream analysis. baBID-seq quantitatively

maps Ψ modifications at various positions within tRNA, including the stem and loop of the

T-arm, the anticodon arm, and the D-arm (**Fig. 1d** and **Supplementary Fig. 2b,c**). To

investigate tRNA Ψ dynamics during exponential versus stationary growth phases, we

quantified Ψ fraction differences at each specific site across four strains. Our analysis

revealed that most Ψ sites on bacterial tRNA consistently exhibited higher modification

fractions in the stationary phase across all examined strains (**Fig. 1e**). In *K. pneumoniae*, the

Ψ sites within the T-arm, D-arm, and anticodon arm collectively showed a reduced

modification fraction in the exponential phase compared to the stationary phase (**Fig. 1d**).

Similarly, in *B. cereus* and *P. syringae*, most tRNA Ψ sites within the T-arm displayed lower

Ψ fractions during the exponential phase (**Supplementary Fig. 2b,c**). Previous research has

shown that the T-arm of tRNA can globally influence tRNA maturation and regulate

translation in *E. coli*[8]. Thus, this growth phase-dependent pattern of tRNA pseudouridylation

suggests a coordinated regulatory mechanism that may affect mRNA translation as bacteria

adapt to changing environmental conditions.

In addition to rRNA and tRNA, baBID-seq also identified highly conserved Ψ sites on

mRNA between biological replicates (**Fig. 1f** and **Supplementary Fig. 2d**), providing strong

confidence in the identification of genuine Ψ modifications. For subsequent analysis, we

focused exclusively on Ψ sites that were consistently detected across biological replicates.

BID-seq uncovers abundant Ψ modification in bacterial mRNA

With the identification of highly conserved Ψ modifications in bacterial mRNA enabled by baBID-seq, we proceeded to analyze their distribution patterns and quantitative features across the transcriptome. In total, we detected over 3,000 Ψ sites in the mRNA of four bacterial strains. Notably, the metagene plot revealed that most Ψ sites were enriched within the coding sequences (CDS) of bacterial mRNAs, and this distribution pattern shows remarkable consistency across the four strains (**Fig. 2a,b**), which was also similar to the observations in human cell lines[19]. Overall, the average Ψ modification in bacterial mRNA (mean Ψ fraction: 15%) was lower than that in 16S and 23S rRNA (mean Ψ fraction: 40%). Most mRNA Ψ sites were primarily distributed below a 25% Ψ fraction, while a smaller proportion reached modification fraction above 50% as highly modified ones (**Fig. 2c** and **Supplementary Fig. 3a**).

Since Ψ modifications in mRNA untranslated regions (UTRs) can influence mRNA processing in eukaryotic cells[2,13], and the UTRs play a crucial role in post-translational regulation in bacteria[24], we conducted a detailed investigation of Ψ modifications in the upstream and downstream UTRs. We found that UTRs pseudouridylation accounted for 7% to 16.4% of total Ψ sites across the transcriptome in both stationary and exponential phases (**Supplementary Fig. 3b,c**). Notably, in *P. aeruginosa* and *B. cereus*, we observed a significantly higher modification fraction for Ψ sites within downstream regions compared to coding sequences (CDS) in both growth phases (**Fig. 2d** and **Supplementary Fig. 3d**). In contrast, *P. aeruginosa* exhibited a higher Ψ modification level in the upstream regions of mRNA during the exponential phase (**Fig. 2d**).

We defined a Ψ-modified gene as any mRNA containing one or more Ψ sites in any growth phase and identified both phase-shared and phase-unique genes across the four strains (**Fig.**

2e). We also noted that individual genes frequently harbored multiple Ψ sites, with the

number of sites varying dynamically across bacterial growth phases (**Fig. 2f**).

Motif analysis of Ψ modifications in bacterial mRNA, rRNA and tRNA

Previous research has shown that different RNA species exhibit unique Ψ modification

patterns, with Ψ synthases targeting specific sequence motifs in tRNA and rRNA (such as the

RluA motif ΨURAA)[5,25]. To identify the sequence determinants of Ψ modifications and

uncover RNA-type-specific Ψ motif contexts across bacterial transcriptomes, we conducted a

comprehensive motif analysis of Ψ-modified sites with a fraction above 2%, which could be

reliably identified in both growth phases. The sequence context analysis focused on 5-

nucleotide motifs centered on each Ψ site. We calculated the percentages of Ψ motifs by

dividing the count of each unique Ψ motif by the total number of Ψ motifs detected in the

mRNA of a single bacterial strain (**Fig. 3a**).

Our analysis revealed a diverse array of Ψ motifs within bacterial mRNA based on baBID-

seq data. Notably, GCΨCG, GGΨCG, and CCΨCG were the most abundant motifs observed

in the three Gram-negative bacteria, while GUΨGU and GGΨGU were the dominant motifs

in *B. cereus* (**Fig. 3a**). We then analyzed the quantitative features of Ψ sites within these

diverse motif contexts and found that the average Ψ fractions for different motifs ranged from

3.4% to 96.6%, indicating varying Ψ installation efficiencies regulated by the properties of

different Ψ synthases (**Fig. 3b**). Overall, we summarized the top 10 frequent Ψ motifs for

bacterial mRNA of *P. aeruginosa*, and *P. syringae*: GUΨCG, (CC/CU/GC/GG/UC)ΨCG,

(CU/GC/UC)ΨCC and GCΨGG (**Fig. 3c**).

baBID-seq also reveals Ψ motif contexts in bacterial rRNA and tRNA. While a variety of Ψ

motifs were identified in bacterial rRNA, the GUΨCG motif stands out as the predominant Ψ

motif in tRNA across the four bacterial strains (**Supplementary Fig. 4a–d**). Notably, the

GUΨCG motif is well-characterized within the T-arm of tRNAs (at position 55) and is

specifically modified by TruB family[5,8,19,26–28]. According to our baBID-seq data, GUΨCG

has been confirmed as the predominant motif in both tRNA and rRNA (mean fraction: 68%),

as well as in mRNA (mean fraction: 19%) (**Fig. 3c** and **Supplementary Fig. 4c,d**),

suggesting that TruB may also play a role in Ψ installation in bacterial mRNA. We also

identified several other key motifs, including UUGC, UUGA, and UUAAA, which

correspond to the previously characterized RluA motif in *E. coli*[25]. Overall, no distinct

sequence motifs were universally enriched in the mRNA of the four bacterial strains

(**Supplementary Fig. 4e–h**), likely due to the complex interactions among multiple Ψ

synthases involved in U-to-Ψ conversion on mRNA.

To determine whether sequence preferences for Ψ modifications vary under different growth

conditions, we calculated both motif frequency and average Ψ fraction for each Ψ motif in *P.*

aeruginosa. Notably, we observed a decrease in the frequency of the GCΨCG motif (**Fig. 3d**),

as the most abundant Ψ motif among the three Gram-negative bacteria. For other top Ψ

motifs, the associated modification levels showed only slight variations between the

exponential and stationary phases. To gain insights into the sequence features related to Ψ

modifications, we analyzed the nucleotide composition within 10 base pairs flanking Ψ sites

in bacterial mRNA. While most strains exhibited non-unique differences in GC content, *B.*

cereus displayed a notably higher GC ratio (normalized to the genomic background GC

content) compared to the other three bacterial strains (**Supplementary Fig. 4i–k**). To explore

the relatively distinct motif pattern in *B. cereus*, we conducted comparative orthology-based

analyses to identify pseudouridine synthase sequences among the five bacterial strains

(**Supplementary Fig. 4l**). A unique interactive regulation of pseudouridine synthases in *B.*

cereus may explain its divergent Ψ sequence patterns compared to other bacterial strains.

Evolutionary conservation of clustered Ψ modifications in bacterial orthologous genes

To investigate the evolutionary conservation of mRNA Ψ modifications among bacterial

strains, we analyzed orthologous genes, focusing on both the preservation of modification

sites and the functional characteristics of Ψ-modified genes. Based on clustered Ortho groups

and genome annotations, our results revealed 225 homologous genes carrying Ψ

modifications in at least two bacterial strains (**Fig. 4a**). We systematically characterized the

biological functions of these homologous Ψ-modified genes among the four bacterial strains

through pathway enrichment analysis, using annotated gene sets from the Kyoto

Encyclopedia of Genes and Genomes (KEGG). This analysis uncovered three distinct but

interconnected metabolic clusters. The primary cluster showed highly significant enrichment

in central carbon metabolism, including glycolysis, TCA cycle, oxidative phosphorylation,

and amino acid biosynthesis (**Fig. 4a** and **Supplementary Fig. 5a**). Notably, gene clusters

essential for ATP production, such as *atpA* and *atpD*, were also enriched. Among the four

bacterial strains, the functional enrichment of Ψ-modified genes in core metabolic pathways,

combined with the conservation of Ψ motif contexts (**Fig. 3a**), provides evidence for the

functional importance and evolutionary conservation of a specific group of Ψ-modified

transcripts across bacterial systems. In addition to the homologous Ψ-modified genes, we

identified 633 strain-specific Ψ-modified mRNAs, highlighting the dynamic nature of Ψ

modifications and suggesting potential strain-specific regulatory mechanisms.

We then conducted a functional investigation of Ψ-modified genes in the two growth phases

of *P. aeruginosa*. The Gene Ontology (GO) enrichment analysis revealed that Ψ-modified

genes were significantly enriched in GO terms related to energy metabolism during the

exponential phase compared to the stationary phase, highlighting distinct state-specific GO

patterns (**Supplementary Fig. 5b**). The exponential phase is known to exhibit elevated levels

of metabolism-related genes, reinforcing our findings regarding the biological relevance of Ψ

modifications. Notably, Ψ-modified genes showed a preferential enrichment in type IV

swarming motility during the stationary phase. For instance, the key transcription factor *algR*,

which regulates multiple virulence factors and promotes *P. aeruginosa* twitching motility,

serves as a representative example among the Ψ-modified genes[29]. This finding suggests a

specific role for Ψ modifications in coordinating motility behaviors during the stationary

growth phase, potentially linking RNA modification to adaptive virulence mechanisms when

nutrient availability is limited. Besides, we observed that homologous genes exhibited more

stable Ψ fraction patterns, likely due to their enriched functions in fundamental metabolic

processes (**Supplementary Fig. 5c**).

To explore the regional distribution of Ψ modifications in greater detail, we conducted a 500

bp sliding window analysis across all detected mRNA transcripts in four bacterial strains,

identifying specific regions with multiple Ψ sites using baBID-seq (**Supplementary Fig. 5d**).

Further analysis revealed that most Ψ-enriched regions corresponded to operons or gene

clusters. We identified Ψ clusters containing two or more Ψ sites in operons such as the

evolutionarily conserved *atp* operon[30] and *eno* operon, across all strains[31,32] (**Fig. 4b**). This

phenomenon was also observed in other homologous gene clusters or operons, such as *rpoA*,

fusA, *groE*, and *rpc* operons[33] (**Supplementary Fig. 5e**), suggesting that Ψ modification may

play a role in regulating post-transcriptional processes[2]. Upon examining the Ψ distribution

within operons and gene clusters in more detail, we observed distinct regional patterns of Ψ

modifications. Some Ψ modifications accumulated within the 3' UTR regions of *atp* operon,

for instance, in *atpD* of *P. aeruginosa* (**Fig. 4b**). In *P. aeruginosa*, Ψ sites in the *atpD* gene

were predominantly located near stop codon, whereas in *B. cereus*, most Ψ modifications

were concentrated at the translation initiation region (**Fig. 4b**). This detailed regional analysis

of Ψ modifications within gene clusters revealed strain-specific distribution patterns in the

conserved operons. The flexible regional Ψ modification patterns observed in operons such as

atp may regulate gene expression to align with bacteria-specific metabolic needs.

Growth state-dependent dynamic Ψ modification in bacterial mRNA

Ψ modification levels on mRNA have been reported to fluctuate under stress conditions in

human cells[34]. Our study also observed alterations in Ψ modifications across the four

bacterial strains under different growth conditions. We conducted a detailed comparative

analysis of Ψ sites between the two growth phases, identifying both newly emerged and

diminished Ψ modification events, as well as alteration in modification fractions at conserved

sites. The quantitative baBID-seq approach allowed us to pinpoint dynamic Ψ modifications

in response to bacterial metabolic shifts and changes in growth states. We initially compared

the distribution of modification fractions for all mRNA Ψ sites in the exponential phase

versus the stationary phase. *K. pneumoniae* and *B. cereus* exhibited significantly higher

global Ψ levels during the stationary phase (**Supplementary Fig. 6a,b**). In contrast, no matter

in nutrient-enriched or minimal media (MM) conditions, *P. aeruginosa* and *P. syringae* did

not show significant changes in global Ψ fractions between the two phases (**Supplementary**

Fig. 6c,d).

We then focused on changes in Ψ fractions at specific sites, setting a cutoff of 10% variation

between the two growth phases across the four bacterial strains. In *P. aeruginosa* and *P.*

syringae, we identified numerous phase-specific Ψ sites that either emerged or disappeared,

as well as many conserved Ψ sites exhibiting changes in Ψ fractions above 10% (**Fig. 5a,b**).

For instance, in *P. aeruginosa*, key genes linked to energy metabolism, such as *sucB, sucC*,

and *gltA*, displayed Ψ sites of increased modification fraction during the exponential phase of

heightened metabolic activity. The stop codon region of *secY*, a protein essential for type II

secretion systems in *P. aeruginosa*, contained a highly modified Ψ site that was only detected

during the exponential growth phase; similar Ψ fraction dynamics were observed for *secY* in

K. pneumoniae and *B. cereus* (**Supplementary Fig. 6a–c**).

In addition to analyzing Ψ sites, we also calculated Ψ intensity for each gene, defined as the

sum of modification fractions for all Ψ sites within a single mRNA molecule. During the

transition from the exponential to the stationary phase, the Ψ intensity of specific genes

shifted significantly (**Fig. 5c–e** and **Supplementary Fig. 6e**). In the exponential phase, many

genes involved in metabolism, amino acid biosynthesis, and protein synthesis exhibited

higher Ψ intensity, such as *rpsD*, *rpsT*, *tufB*, *lon*, *nuoD*, etc., reflecting their critical role in

supporting rapid growth; meanwhile, for these genes, their decreased Ψ intensity observed in

the stationary phase may suggest a coordinated Ψ reprogramming that helps bacteria adapt to

reduced nutrient availability and increased cell density by downregulating energy-intensive

processes. Overall, the dynamic nature of Ψ modifications—including newly emerged and

disappeared Ψ sites, as well as altered Ψ fractions across the two growth conditions—

indicates their potential roles as responsive epi-transcriptomic switches that facilitate bacteria

to be adapted to varying environmental challenges and metabolic demands.

Ψ correlates with bacterial mRNA metabolism and function

It is well established that Ψ can enhance mRNA stability and translation in mammals[35],

parasites[36,37], and plants[37]. However, the effects of Ψ on bacterial mRNA have yet to be

investigated. We normalized gene expression levels by calculating transcripts per kilobase

million (TPM) for the two growth phases and grouped the adequately expressed genes into

Ψ-modified mRNA (Ψ-mRNA) and non-Ψ-modified mRNA (non-Ψ-mRNA). The analysis

revealed that the expression level of Ψ-mRNA was significantly higher than that of non-Ψ-

mRNA in *P. aeruginosa* and *P. syringae* when cultured in a nutrient-sufficient medium (**Fig.**

5f,g and **Supplementary Fig. 6f**). However, *P. syringae* exhibited more moderate changes in

mRNA expression between the different growth phases under minimal medium (MM)

conditions (**Fig. 5h**). Our findings suggest that Ψ may stabilize mRNA in a growth phase-

dependent manner in bacteria.

We then conducted comprehensive analyses of TE in *P. syringae* with alterations in Ψ

modifications under two distinct conditions (King's B medium, KB and minimal medium,

MM) to examine whether Ψ impacts bacterial mRNA translation. Although we did not

observe a universal global correlation between changes in Ψ modifications and TE for all

genes under both conditions, a larger proportion of genes did show a positive correlation

between TE and Ψ modifications (**Fig. 5i**). Our findings partially align with previous studies

in mammals, which suggested that Ψ modifications could enhance mRNA translation

efficiency in bacteria and may play more complex roles in translation regulation.

Hfq is a major bacterial post-transcriptional regulator that functions as a pivotal RNA-binding

protein (RBP), orchestrating various cellular processes[22]. Its regulatory mechanisms have

been extensively characterized, including the alteration of RNA structure and the facilitation

of sRNA-mRNA interactions[38], highlighting its fundamental role in coordinating gene

expression networks [39,40]. To investigate the potential association between Hfq and Ψ

modifications, we performed an integrative analysis combining our data with previously

published Hfq RIP-seq data[22] in *P. aeruginosa*. We examined both exponential and stationary

growth phases to evaluate whether Hfq targets Ψ-modified regions or whether Ψ

modifications influence Hfq-RNA interactions. Our results indicated that the distance

between Ψ sites and Hfq peak centers significantly decreased during the stationary phase

(**Supplementary Fig. 6g**); meanwhile, Hfq-bound genes accounted for a substantial

proportion of mRNA Ψ sites, with 12.5% during the exponential phase and 27.3% during the

stationary phase (**Fig. 5j**). By defining the Hfq binding score as the sum of Hfq peaks binding

strength per gene, we found that, in the stationary phase, Hfq exhibited a stronger binding

affinity for genes carrying more Ψ modifications (**Fig. 5k**). These results suggest that Ψ

modifications may facilitate mRNA-Hfq interactions to some extent. Overall, dynamic Ψ

modifications have been shown to impact bacterial mRNA stability, translation, and RBP

interactions in response to changing cellular demands during growth phase transitions.

358 Integrated computational analysis reveals structure-dependent Ψ modifications in

359 bacterial RNA

Although some Ψ modifications shared common sequence motifs, we observed an array of

diverse motif contexts at Ψ sites in bacterial mRNA (**Fig. 3a,b**). This observation aligns with

previous studies demonstrating that pseudouridine synthases, such as PUS1 and TruB,

preferentially recognize RNA local structures beyond primary sequence motifs for Ψ

installation[5,41–43]. To model and study the widespread Ψ modifications on bacterial mRNA,

which are hypothesized to be structure-dependent, we incorporate Ψ modification

determinants through comprehensive clustering analyses of local RNA sequences and

structural elements to test our hypothesis. We firstly calculated the predicted secondary

structure 41nt centered GUΨC motif, with the representative Ψ sites of 50%~96%

modification fraction (**Fig. 6a**). Interestingly, all GUΨC motifs with varying Ψ fractions are

predicted to occur within RNA loop structures. To determine whether RNA structural factors

influence Ψ deposition and modification fractions, we compared the structural and fractional

characteristics of the two highly prevalent motifs, GUΨC and GCΨCG, by clustering

predicted RNA structures across all RNA species in *P. aeruginosa*. The results indicated that,

aside from tRNA and rRNA, which clustered together due to their distinct structural features,

we observed small clusters within certain mRNAs, such as *guaB*, *recA*, and *PA4943*. Overall,

we did not find characteristic structural signatures that could completely distinguish GUΨC

from the GCΨCG motif (**Fig. 6b**). To gain deeper insights, we conducted structural clustering

analyses of all RNA species across different strains (**Supplementary Fig. 7a-d**). Given that

some pseudouridine synthases target specific structural features, we anticipated highly

distinguishable clustering results. However, contrary to our expectations, such distinct

clustering patterns were not observed. This may suggest that certain pseudouridine synthases,

such as RluA[25], do not solely rely on structural features for target recognition. Notably, we

identified clusters of Ψ sites with similar structures that exhibited higher modification

fractions, including *sucC* and *sucB* in *P. aeruginosa* (**Supplementary Fig. 7a**), as well as

sdhA, *PSPPH_RS14750*, and *atpB* in *P. syringae* (**Supplementary Fig. 7c**). Integrating these

findings with the distinctive Ψ fraction patterns exhibited by various motifs (**Fig. 3b**), we

conclude that both local sequence and structural features of RNA may play important roles in

Ψ installation.

Transformer-GNN-based neural networks for integrated prediction of Ψ

In next-generation sequencing (NGS) data, variable read coverage dictated by gene

expression patterns or limited sequencing depth can lead to missed potential Ψ sites. To

address this limitation, we implemented a strategy that integrates RNA sequence and

structural features for a transcriptome-wide scan of Ψ sites, resulting in a more

comprehensive inventory of potential Ψ sites. Previous studies have shown that the sequence

context surrounding Ψ sites can serve as a reliable predictor for RNA modifications[28,44].

Building on this, we developed a deep learning model that accurately captures both sequence

and structural features of known Ψ sites across various RNA species, allowing us to predict

potential modification sites that may be condition-dependent or below the detection

thresholds of BID-seq. We extracted sequence segments of 41, 61, and 81 nucleotides (± 20,

± 30, and ± 40 bp) centered on each Ψ site, applying window shifts of ± 5, ± 10, and ± 15 bp

respectively. The input sequences were then embedded using one-hot encoding and processed

through a multi-head transformer module. Simultaneously, we utilized adjacency matrices

representing local RNA structures (predicted using MXfold2[45] as input for a graph neural

network (GNN) module. Features extracted from both modules were combined through

weighted concatenation and subsequently processed using a residual block. We employed

binary cross-entropy loss for predicting the likelihood of Ψ modifications. This hybrid

transformer-GNN architecture effectively integrated both sequence and structural

characteristics across various RNA sequences (**Fig. 6c**), termed as pseU_NN.

We used 3,377 high-confidence Ψ sites (with fraction values >2%) as positive samples. The

negative samples consisted of 3,400 randomly selected sites that contained the unique Ψ

motif but showed no evidence of Ψ deposition in our experimental data. The dataset was

divided into 4,744 training samples, 1,016 test samples, and 1,017 validation samples. The

model consistently performed well across different input dimensions (41, 61, and 81

nucleotides), with all variants achieving AU-ROC scores exceeding 0.8 after convergence

(**Fig. 6d**). Using 41-nucleotide inputs, our methodology achieved impressive validation

metrics, including an area under the precision-recall curve (AU-PRC) of 0.878 and an area

under the receiver operating characteristic curve (AU-ROC) of 0.84 (**Fig. 6e**). Models trained

with alternative input sequence lengths also demonstrated strong performance metrics

(**Supplementary Fig. 7e,f**). These results set the basis for further development of effective

deep learning tools for transcriptome-wide Ψ prediction in both bacterial and mammalian

RNAs.

Discussion

RNA modifications in bacteria, particularly Ψ, are less characterized than their well-studied eukaryotic counterparts. Leveraging recent advances in quantitative sequencing methods such as BID-seq[19,20], we present comprehensive, single-nucleotide resolution maps of Ψ modifications, complete with stoichiometric information, across four bacterial strains under different growth phases. Our findings confirm the widespread occurrence of Ψ modifications in bacterial RNA and provide a functional analysis of these modifications. This extensive dataset serves as a valuable resource for understanding the evolutionary and functional significance of RNA Ψ modifications in bacteria.

The Ψ modification in tRNA plays critical regulatory roles in tRNA aminoacylation, stability, and the formation of functional structures[8,26,46,47]. Our analysis revealed that tRNA Ψ modification is present in varying fractions, with a stronger modification level observed in the TΨC loop compared to the anticodon-arm and D-arm loops. Previous studies indicate that the tRNA T-arm is highly modified, not only by Ψ but also by other uridine modifications like 5-methyluridine (m^5U)[48]. Both Ψ and m^5U modifications globally enhance tRNA aminoacylation, and also independently influence specific tRNA modifications, such as 3-(3-amino-3-carboxypropyl)uridine at position 47[8]. The TΨC loop is crucial for the interaction between tRNA and the ribosome, facilitating the formation of the tRNA-ribosome complex[48] Interestingly, we observed a novel phenomenon where Ψ modifications on the tRNA T-arm increase in the stationary growth phase compared to the exponential phase. Given that codon composition and mRNA expression are closely correlated[49], the dynamic Ψ modification in the TΨC loop may regulate bacterial mRNA translation by modulating T-arm interactions with the ribosome. Besides, other RNA modifications on tRNA may be influenced by the dynamic Ψ pattern during growth phase transitions, potentially creating a feedback loop where existing modifications affect the biogenesis of subsequent modifications. Overall, this condition-dependent Ψ modification in tRNA may represent a new mechanism by which bacteria adapt to their environment, anticipating further investigation in future research.

Previous studies have demonstrated that both a deficiency and an excess of pseudouridine can

severely impair ribosomal translation and proper assembly in *E. coli*[11,50]. The stable fraction

of Ψ modification observed in *E. coli* rRNA across two different growth phases suggests that

rRNA Ψ may be tightly regulated to maintain essential rRNA function. The role of Ψ in

mRNA remains unclear across the three domains of life. Our results reveal a comprehensive

mRNA Ψ landscape in four bacterial strains. We found that the overall fraction of mRNA Ψ

modifications was significantly lower than that of rRNA and tRNA, consistent with findings

in plants and mammals[19,37]. The distribution and stoichiometric patterns of mRNA Ψ between

Gram-positive and Gram-negative bacteria exhibited similarities. Notably, we identified

evolutionarily conserved Ψ modifications in mRNAs encoding proteins involved in energy

generation, ATP binding, amino acid synthesis, and protein translation, mirroring

observations in mammals and plants[19,37]. We also discovered clusters of multiple Ψ sites

enriched in specific operons related to conserved functions, which were detected across

multiple strains.

To date, limited research has focused on Ψ modifications and their alterations in bacteria. In

this study, we profiled and uncovered dynamic changes in Ψ modifications during growth

phase transitions among four bacterial strains. During the metabolically active phase of *P.*

aeruginosa, we observed increased Ψ modifications in many metabolism-related genes

(**Supplementary Fig. 5b**). In the stationary phase, our analysis revealed reduced

pseudouridylation within the coding regions of *fimV*, a gene that encodes an inner membrane

protein in *P. aeruginosa* responsible for regulating intracellular cyclic AMP levels, type IV-

mediated twitching motility, and type II secretion system genes[51,52]. Ψ modifications in

coding regions are known to alter codon properties on mRNA, leading to reconstituted

translation and promoting the low-level synthesis of multiple peptides[17]. These findings

suggest a potential new mechanism for regulating bacterial metabolism and quorum sensing

in *P. aeruginosa*. Previous studies have demonstrated that pseudouridylation enhances

mRNA stability in mammals and plants[19,20,37], and our research confirms and extends these

observations to bacterial systems.

It has been reported that methionine aminoacyl tRNA[Met] synthetase can target Ψ1074 in yeast

[53], and several Ψ sites overlap with RNA-binding protein (RBP) binding regions[13]. To

systematically investigate dynamic Ψ modifications in bacterial RNA and their potential

impact on RBP binding, we conducted an integrative analysis that combined our BID-seq

data with Hfq RIP-seq data from P. aeruginosa. Hfq is an RNA chaperone that recognizes 5-

repeat AAN motifs in P. aeruginosa and plays a crucial role in regulating various post-

transcriptional processes, including mediating sRNA-mRNA interactions[38]. Hfq can trigger

mRNA structural reprogramming, and RNA structural switches may facilitate or hinder the

process of pseudouridylation[21,41]. We observed an increased binding affinity of Hfq to Ψ-

modified mRNA during the stationary phase compared to the exponential growth phase. This

discovery opens new avenues for understanding the regulatory interactions between Ψ

modifications and Hfq, which may closely correlate with the restructuring of mRNA through

the direct or indirect effects of Ψ modifications.

Ψ modification is specifically recognized in RNA partial structures or local motif contexts by

the TruB pseudouridine synthase[54]. Our comprehensive analysis of Ψ-containing sequences

in mRNA revealed conserved motif signatures that closely resemble those found in tRNA and

rRNA. This conservation pattern suggests that tRNA and rRNA Ψ synthases may directly

recognize similar structural and sequence elements in mRNA, as evidenced by PUS1 and

PUS6, which can both add Ψ to tRNA and mRNA[41,53]. Through an integrated approach

combining RNA sequence and structural clustering analysis, we demonstrated that both the

selection of Ψ sites and the modification fraction are strongly associated with the local

secondary structure of RNA molecules[41]. From a structural perspective, this specificity likely

explains why certain potential modification sites with appropriate motifs remain unmodified

or exhibit dynamic Ψ levels under different growth conditions.

Deep learning methods are increasingly utilized for RNA modification prediction. Building

on previous concepts, such as attention-based multi-label neural networks that predict

multiple RNA modifications using sequence context[44], we developed a hybrid transformer-

GNN architecture (pseU_NN) that integrates RNA structural features with multi-head

attention mechanisms to predict potential Ψ modification sites, particularly those that may have eluded detection due to low or absent gene expression. The robustness of our model was rigorously validated using high-confidence sites of various lengths and predicted RNA structures as input. pseU_NN enables reliable prediction of potential Ψ sites across different bacterial contexts, providing a comprehensive map of potential Ψ sites within bacterial transcriptomes.

Overall, this study presents the first comprehensive landscape of Ψ modifications across multiple diverse bacterial strains using optimized BID-seq. We systematically revealed Ψ fractions in tRNA, rRNA, and mRNA under exponential and stationary growth conditions in four strains, as well as under nutrient-deficient conditions in P. syringae. Our findings indicated that Ψ modifications on rRNA were stable under both growth conditions in *E. coli* and *P. aeruginosa*, while significant variations in Ψ fractions were observed in tRNA and mRNA across different growth phases. The motif conservation analysis provides insights into the activity of pseudouridine synthases on bacterial mRNA. We further identified evolutionarily conserved patterns of Ψ-enriched modifications in conserved operons involved in fundamental metabolic pathways across bacterial strains. In summary, our comprehensive study enhances the understanding of the regulatory roles of Ψ modifications across the bacterial kingdom, paving the way for future mechanistic investigations into the unknown functions of Ψ in bacterial RNA regulation.

Materials and Methods

Bacteria strains and growth conditions

The wild-type *Pseudomonas syringae pv. phaseolicola* 1448A strain was cultured in King's B (KB) medium[55] (20 g/L proteose peptone, 1.5 g/L K_2HPO_4, 1.5 g/L $MgSO_4 \cdot 7H_2O$, and 10 mL/L glycerol) at 28°C for 12 hours (overnight) until reaching an optical density at 600 nm (OD_{600}) of 1–2, corresponding to stationary phase. *Bacillus cereus* ATCC 14579, *Pseudomonas aeruginosa*, *Escherichia coli* K-12 MG1655, and *Klebsiella pneumoniae* CR-HvKP4 were grown in Luria-Bertani (LB) broth at 37°C until achieving an OD_{600} of 1–2 for

stationary phase samples. For exponential phase samples, all bacterial strains were first

cultured to stationary phase as described, then subcultured into fresh medium and incubated

under the same conditions until reaching an OD_{600} of 0.5–0.6. For *P. syringae* in MM,

exponential phase cells were harvested, washed three times with freshly prepared minimal

medium[56] (MM), resuspended in MM at an OD_{600} of 0.1–0.2, and cultured for an additional 6

hours.

baBID-seq library construction

RNA was extracted using RNA isolation Kit V2(Vazyme, #RC112-01). The extracted RNA

processed using DNase I (RNase-free, NEB # M0303S) and collected using RNA Clean &

Concentrator™-5 (Zymo #R1014). The RiboRID technique was used for rRNA depletion[23].

For strains B.cereus and K. pneumoniae, rRNA was depleted with NEBNext® rRNA

Depletion Kit (Bacteria) (#E7850L). RNA concentration in each step was tested using Qubit

RNA Assays.

One hundred nanograms of RNA generated from the ribosome removal process from each

biological replicate was used for the library construction. The fragmentation was optimized to

4 min under 70 °C with fragmentation buffer used (Invitrogen). For the following steps, we

strictly followed the BIDseq protocal[20]. The final amplified cDNA are collected and then

optimal library fragments are selected using native PAGE gels. 40% Acrylamide/Bis (29:1)

10% native polyacrylamide gel used. The optimal band (175-200bp) used collected. 400 μL

of 1x TE buffer and soak the gel thoroughly at 37 °C for 1 h on the thermal shaker at 600 rpm.

The gel was crushed and snap-freezed using liquid nitrogen and incubated on a thermal

shaker at 37 °C at 600rpm for 12 hours. The supernatant collected using Spin-X followed the

DNA precipitation. The constructed libraries were sequenced on the Illumina NovaSeq

sequencing platform in paired reads mode and single-end reads used for following BID-seq

data processing.

Sequencing data processing and analysis

Sequencing data were subjected to a refined bioinformatic workflow adapted from previously

established protocols. Raw sequencing reads underwent adapter trimming via Cutadapt (v.3.5)

and PCR duplicate elimination using BBMap tools (v.38.73). The filtered reads were initially

aligned to a curated repository of non-coding RNA sequences (including rRNAs, tRNAs, and

other small RNAs). Subsequently, unmapped reads were subjected to genome alignment with

optimized mapping parameters tailored for Ψ detection. The resultant alignment files were

processed with Samtools (v.1.13) to generate strand-specific BAM files, which were then

interrogated using bam-readcount (v.1.0.1) to quantify nucleotide deletion events and

calculate coverage metrics. Ψ modification sites were identified by integrating deletion rate

profiles, site coverage, and background signal correction derived from "input" libraries.

Bacterial samples from distinct temporal phases were analyzed as discrete entities to

minimize batch effects. The quantitative assessment of Ψ levels was achieved by

transforming raw deletion ratios according to previously reported calibration curves, yielding

high-confidence Ψ fractions at single-nucleotide resolution.

TPM calculation

The TPM value for a specific transcript *i* is calculated as:

$$TPM_i = \left(\frac{q_i/l_i}{\Sigma_j\left(q_j/l_j\right)}\right) \times 10^6$$

q_i was the number of reads mapped to transcript i. l_i is the length of transcript I (in

nucleotides). 10^6 was the scaling factor to express the result as transcripts per million[57].

Evolutionary analysis

Orthologous gene analysis was performed using OrthoFinder[58] software to identify

homologous gene clusters across the five bacterial strains. The analysis incorporated

complete genome sequences and their corresponding GFF (General Feature Format)

annotation files with default parameters. The resulting orthologous gene clusters were

subsequently utilized for KEGG (Kyoto Encyclopedia of Genes and Genomes) pathway

analysis. Genes that were not clustered in the orthology analysis were excluded from the

KEGG pathway mapping to ensure reliable results. Functional annotations were performed

using eggNOG-mapper v2[59].

RNA structure analysis

Pseudouridine modification sites were filtered with a fraction of larger than 2%. RNA

sequences were generated by extracting 20nt upstream and downstream of each modification

site, yielding 41 nucleotide sequences. These sequences were subsequently used for RNA

secondary structure prediction with MXfold2[45] and sequence clustering analysis with

RNAclust[60]. Results analysis and visualization were performed using customized scripts

written in R.

pseU_GNN

Training data

The negative control dataset was constructed by scanning bacterial genomes for

pseudouridylation-compatible sequence motifs absent from the experimentally verified

modification sites in baBID-seq data, then generating sequence segments of 41, 61, and 81

nucleotides with these unmodified motifs at their centers. MXfold2[45] is used to predict the

secondary structure of all sequence as input for training and validation process. 3,377 high-

confidence Ψ sites (with fraction values >2%) as positive samples. The negative samples

consisted of 3,400 randomly selected sites that contained the unique Ψ motif but showed no

evidence of Ψ deposition in our experimental data. The dataset was divided into 4,744

training samples, 1,016 test samples, and 1,017 validation samples for each sequence length.

Model Architecture

The model architecture consisted of three main components: two dense graph encoder

extracting RNA secondary structure features generated from MXfold2 and one-hot embedded

sequence fed into 4 layers multi-head transformer to extract sequence features. Features

extracted from both modules were combined through weighted concatenation and

subsequently processed using a residual block (fully connected layers) followed by sigmoid

activation function to predict the possibility of Ψ modification.

Training and Evaluation

The pseU_NN was trained using binary cross-entropy loss with the Adam optimizer. We

implemented a dynamic learning rate scheduler that adjusted the rate based on validation

AUC-ROC performance.

For the final evaluation, we utilized an independent validation dataset, with F1 scores and

precision-recall curves guiding prediction threshold selection. This balanced approach

ensured optimal sensitivity and specificity - critical in biological classification systems where

both false positives and false negatives carry significant consequences.

The F1 score is the harmonic mean of precision and recall. The definition of TP, FP and FN

with shifting was adopted in previous studies[61,62].

$$\text{F1 } score \ = \frac{2 \times precision \times recall}{precision + recall}$$

Acknowledgments

This study was funded by the Guangdong Major Project of Basic and Applied Basic Research (2020B0301030005), National Natural Science Foundation of China (32172358), General Research Funds of Hong Kong (21103018, 11101619, and 11102720) and Early Career Scheme of Hong Kong (26103623). The funders had no role in study design, data collection, interpretation, or the decision to submit the work for publication.

Data availability

All sequencing data files have been submitted to the National Center for Biotechnology Information (NCBI) Gene Expression Omnibus (GEO) database with the reference code of GSE292335.

Code availability

The code for pseU_NN used for this paper are available at https://github.com/Dylan-LT/pseU_NN.git

Author contributions

X.D. and L.X. conceived the original idea. X.D. and L.-S.Z. supervised the project. L.X. and Y.Z. designed and performed experiments. L.X., S.S., and Y.S. performed bioinformatic analysis and prepared figures. L.X. and Z.G. designed pseU_NN. B.L., Z.G., and R.L. assisted with experiments. L.X., L.-S.Z., and X.D. wrote the manuscript. All the authors read the paper and agreed with the final version.

Conflicts of interest

The authors declare that they have no competing interests.

Reference

1.   Roundtree, I. A., Evans, M. E., Pan, T. & He, C. Dynamic RNA Modifications in Gene

Expression Regulation. *Cell* vol. 169 Preprint at

https://doi.org/10.1016/j.cell.2017.05.045 (2017).

2.   Rodell, R., Robalin, N. & Martinez, N. M. Why U matters: detection and functions of

pseudouridine modifications in mRNAs. *Trends in Biochemical Sciences* vol. 49

Preprint at https://doi.org/10.1016/j.tibs.2023.10.008 (2024).

3.   Cerneckis, J., Cui, Q., He, C., Yi, C. & Shi, Y. Decoding pseudouridine: an emerging

target for therapeutic development. *Trends in Pharmacological Sciences* vol. 43

Preprint at https://doi.org/10.1016/j.tips.2022.03.008 (2022).

4.   Yu, C. T. & Allen, F. W. Studies of an isomer of uridine isolated from ribonucleic

acids. *BBA - Biochimica et Biophysica Acta* **32**, (1959).

5.   Pan, H., Agarwalla, S., Moustakas, D. T., Finer-Moore, J. & Stroud, R. M. Structure of

tRNA pseudouridine synthase TruB and its RNA complex: RNA recognition through a

combination of rigid docking and induced fit. *Proc Natl Acad Sci U S A* **100**, (2003).

6.   Roovers, M. *et al.* Formation of the conserved pseudouridine at position 55 in archaeal

tRNA. *Nucleic Acids Res* **34**, (2006).

7.   Ishida, K. *et al.* Pseudouridine at position 55 in tRNA controls the contents of other

modified nucleotides for low-temperature adaptation in the extreme-thermophilic

eubacterium Thermus thermophilus. *Nucleic Acids Res* **39**, (2011).

8.   Schultz, S. K. *et al.* Modifications in the T arm of tRNA globally determine tRNA

maturation, function, and cellular fitness. *Proceedings of the National Academy of*

Sciences **121**, e2401154121 (2024).

9.   Sloan, K. E. *et al.* Tuning the ribosome: The influence of rRNA modification on

eukaryotic ribosome biogenesis and function. *RNA Biology* vol. 14 Preprint at

https://doi.org/10.1080/15476286.2016.1259781 (2017).

10.   Zhao, Y., Rai, J. & Li, H. Regulation of translation by ribosomal RNA

pseudouridylation. *Sci Adv* **9**, (2023).

11.   Leppik, M., Liiv, A. & Remme, J. Random pseuoduridylation in vivo reveals critical
region of Escherichia coli 23S rRNA for ribosome assembly. *Nucleic Acids Res* **45**,
(2017).

12.   Anderson, B. R. *et al.* Nucleoside modifications in RNA limit activation of 2′-5′-
oligoadenylate synthetase and increase resistance to cleavage by RNase L. *Nucleic*
Acids Res **39**, (2011).

13.   Martinez, N. M. *et al.* Pseudouridine synthases modify human pre-mRNA co-
transcriptionally and affect pre-mRNA processing. *Mol Cell* **82**, (2022).

14.   Karijolich, J. & Yu, Y.-T. Modifying the genetic code: Converting nonsense codons
into sense codons by targeted pseudouridylation. *Nature* **474**, (2012).

15.   Hoernes, T. P. *et al.* Nucleotide modifications within bacterial messenger RNAs
regulate their translation and are able to rewire the genetic code. *Nucleic Acids Res* **44**,
(2016).

16.   Karijolich, J. & Yu, Y. T. Converting nonsense codons into sense codons by targeted
pseudouridylation. *Nature* **474**, 395–399 (2011).

17.   Eyler, D. E. *et al.* Pseudouridinylation of mRNA coding sequences alters translation.
Proc Natl Acad Sci U S A **116**, (2019).

18.   Liang, S. T., Xu, Y. C., Dennis, P. & Bremer, H. mRNA composition and control of
bacterial gene expression. *J Bacteriol* **182**, (2000).

19.   Dai, Q. *et al.* Quantitative sequencing using BID-seq uncovers abundant
pseudouridines in mammalian mRNA at base resolution. *Nat Biotechnol* **41**, 344–354
(2023).

20.   Zhang, L. S. *et al.* BID-seq for transcriptome-wide quantitative sequencing of mRNA
pseudouridine at base resolution. *Nat Protoc* **19**, (2024).

21.   Hua, C. *et al.* Hfq mediates transcriptome-wide RNA structurome reprogramming
under virulence-inducing conditions in a phytopathogen. *Cell Rep* **43**, (2024).

22.   Trouillon, J., Han, K., Attrée, I. & Lory, S. The core and accessory Hfq interactomes
across Pseudomonas aeruginosa lineages. *Nat Commun* **13**, (2022).

23.   Choe, D. *et al.* RiboRid: A low cost, advanced, and ultra-efficient method to remove
ribosomal RNA for bacterial transcriptomics. *PLoS Genet* **17**, (2021).

24.   Adams, P. P. *et al.* Regulatory roles of escherichia coli 5' utr and orf-internal rnas
detected by 3' end mapping. *Elife* **10**, (2021).

25.   Schaening-Burgos, C., LeBlanc, H., Fagre, C., Li, G.-W. & Gilbert, W. V. RluA is the
major mRNA pseudouridine synthase in Escherichia coli. *PLoS Genet* **20**, e1011100-
(2024).

26.   De Crécy-Lagard, V. *et al.* Survey and summary: Matching tRNA modifications in
humans to their known and predicted enzymes. *Nucleic Acids Research* vol. 47
Preprint at https://doi.org/10.1093/nar/gkz011 (2019).

27.   Veerareddygari, G. R., Singh, S. K. & Mueller, E. G. The pseudouridine synthases
proceed through a glycal intermediate. *J Am Chem Soc* **138**, (2016).

28.   Hoang, C. & Ferré-D'Amaré, A. R. Cocrystal structure of a tRNA Ψ55 pseudouridine
synthase: Nucleotide flipping by an RNA-modifying enzyme. *Cell* **107**, (2001).

29.   Kong, W. *et al.* ChIP-seq reveals the global regulator AlgR mediating cyclic di-GMP
synthesis in Pseudomonas aeruginosa. *Nucleic Acids Res* **43**, (2015).

30.   Ventura, M., Canchaya, C., Van Sinderen, D., Fitzgerald, G. F. & Zink, R.
Bifidobacterium lactis DSM 10140: Identificafion of the atp (atpBEFHAGDC) operon
and analysis of its genetic structure, characteristics, and phylogeny. *Appl Environ*
Microbiol **70**, (2004).

31.   Salgado, H., Moreno-Hagelsieb, G., Smith, T. F. & Collado-Vides, J. Operons in
Escherichia coli: Genomic analyses and predictions. *Proc Natl Acad Sci U S A* **97**,
(2000).

32.   Commichau, F. M. *et al.* Novel activities of glycolytic enzymes in Bacillus subtilis:
Interactions with essential proteins involved in mRNA processing. *Molecular and*
Cellular Proteomics **8**, (2009).

731 33. Salgado, H. *et al.* RegulonDB (version 5.0): Escherichia coli K-12 transcriptional

732   regulatory network, operon organization, and growth conditions. *Nucleic Acids Res* **34**,

733   (2006).

734 34. Li, X. *et al.* Chemical pulldown reveals dynamic pseudouridylation of the mammalian

735   transcriptome. *Nat Chem Biol* **11**, (2015).

736 35. Karikó, K. *et al.* Incorporation of pseudouridine into mRNA yields superior

737   nonimmunogenic vector with increased translational capacity and biological stability.

738   *Molecular Therapy* **16**, (2008).

739 36. Nakamoto, M. A., Lovejoy, A. F., Cygan, A. M. & Boothroyd, J. C. MRNA

740   pseudouridylation affects RNA metabolism in the parasite Toxoplasma gondii. *RNA* **23**,

741   (2017).

742 37. Li, H. *et al.* Quantitative RNA pseudouridine maps reveal multilayered translation

743   control through plant rRNA, tRNA and mRNA pseudouridylation. *Nat Plants* **11**, 234–

744   247 (2025).

745 38. Chihara, K. *et al.* Conditional Hfq Association with Small Noncoding RNAs in

746   Pseudomonas aeruginosa Revealed through Comparative UV Cross-Linking

747   Immunoprecipitation Followed by High-Throughput Sequencing. *mSystems* **4**, (2019).

748 39. Sobrero, P. & Valverde, C. The bacterial protein Hfq: Much more than a mere RNA-

749   binding factor. *Critical Reviews in Microbiology* vol. 38 Preprint at

750   https://doi.org/10.3109/1040841X.2012.664540 (2012).

751 40. dos Santos, R. F., Arraiano, C. M. & Andrade, J. M. New molecular interactions

752   broaden the functions of the RNA chaperone Hfq. *Current Genetics* vol. 65 Preprint at

753   https://doi.org/10.1007/s00294-019-00990-y (2019).

754 41. Carlile, T. M. *et al.* mRNA structure determines modification by pseudouridine

755   synthase 1. *Nat Chem Biol* **15**, (2019).

756 42. Lange, S. J. *et al.* Global or local? Predicting secondary structure and accessibility in

757   mRNAs. *Nucleic Acids Res* **40**, (2012).

758   43.   Safra, M., Nir, R., Farouq, D., Slutzkin, I. V. & Schwartz, S. TRUB1 is the
759   predominant pseudouridine synthase acting on mammalian mRNA via a predictable
760   and conserved code. *Genome Res* **27**, (2017).

761   44.   Song, Z. *et al.* Attention-based multi-label neural networks for integrated prediction
762   and interpretation of twelve widely occurring RNA modifications. *Nat Commun* **12**,
763   (2021).

764   45.   Sato, K., Akiyama, M. & Sakakibara, Y. RNA secondary structure prediction using
765   deep learning with thermodynamic integration. *Nat Commun* **12**, (2021).

766   46.   Krutyhołowa, R., Zakrzewski, K. & Glatt, S. Charging the code — tRNA modification
767   complexes. *Current Opinion in Structural Biology* vol. 55 Preprint at
768   https://doi.org/10.1016/j.sbi.2019.03.014 (2019).

769   47.   de Crécy-Lagard, V. & Jaroch, M. Functions of Bacterial tRNA Modifications: From
770   Ubiquity to Diversity. *Trends in Microbiology* vol. 29 Preprint at
771   https://doi.org/10.1016/j.tim.2020.06.010 (2021).

772   48.   Chou, H. J., Donnard, E., Gustafsson, H. T., Garber, M. & Rando, O. J.
773   Transcriptome-wide Analysis of Roles for tRNA Modifications in Translational
774   Regulation. *Mol Cell* **68**, (2017).

775   49.   Gouy, M. & Gautier, C. Codon usage in bacteria: Correlation with gene expressivity.
776   *Nucleic Acids Res* **10**, (1982).

777   50.   O'Connor, M., Leppik, M. & Remme, J. Pseudouridine-free Escherichia coli
778   ribosomes. *J Bacteriol* **200**, (2018).

779   51.   Semmler, A. B. T., Whitchurch, C. B., Leech, A. J. & Mattick, J. S. Identification of a
780   novel gene, fimV, involved in twitching motility in Pseudomonas aeruginosa.
781   *Microbiology (N Y)* **146**, (2000).

782   52.   Buensuceso, R. N. C. *et al.* The conserved tetratricopeptide repeat-containing C-
783   Terminal domain of pseudomonas aeruginosa fimv is required for its cyclic AMP-
784   dependent and -independent functions. *J Bacteriol* **198**, (2016).

53. Levi, O. & Arava, Y. S. Pseudouridine-mediated translation control of mRNA by

methionine aminoacyl tRNA synthetase. *Nucleic Acids Res* **49**, (2021).

54. Machnicka, M. A., Olchowik, A., Grosjean, H. & Bujnicki, J. M. Distribution and

frequencies of post-transcriptional modifications in tRNAs. *RNA Biol* **11**, (2014).

55. King, E. O., Ward, M. K. & Raney, D. E. Two simple media for the demonstration of

pyocyanin and fluorescin. *J Lab Clin Med* **44**, (1954).

56. Huynh, T. V., Dahlbeck, D. & Staskawicz, B. J. Bacterial blight of soybean:

Regulation of a pathogen gene determining host cultivar specificity. *Science (1979)*

245, (1989).

57. Zhao, Y. *et al.* TPM, FPKM, or Normalized Counts? A Comparative Study of

Quantification Measures for the Analysis of RNA-seq Data from the NCI Patient-

Derived Models Repository. *J Transl Med* **19**, (2021).

58. Emms, D. M. & Kelly, S. OrthoFinder: Phylogenetic orthology inference for

comparative genomics. *Genome Biol* **20**, (2019).

59. Cantalapiedra, C. P., Herṇandez-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J.

eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain

Prediction at the Metagenomic Scale. *Mol Biol Evol* **38**, (2021).

60. Engelhardt, J., Heyne, S., Will, S. & Reiche, K. RNAclust Documentation.

Bioinformatics Group, Department of Computer Science, University of Leipzig (2010).

61. Yu, M. *et al.* SnapHiC: a computational pipeline to identify chromatin loops from

single-cell Hi-C data. *Nature Methods 2021 18:9* **18**, 1056–1059 (2021).

62. Wang, F. *et al.* A Lightweight Framework For Chromatin Loop Detection at the

Single-Cell Level. *Advanced Science* **10**, 2303502 (2023).

Figure 1

Figure 1. BID-seq identifies precise pseudouridine (Ψ) modification sites in ribosomal RNA and reveals dynamic Ψ modification patterns in transfer RNA. a,b Ψ modifications detected on 16S and 23S ribosomal RNA with baBID-seq of *E. coli* (**a**) and *P. aeruginosa* (**b**) total RNA during exponential and stationary growth phases. All Ψ sites in panels **a** and **b** were identified using filtration criteria of deletion ratio > 0.02 and *p*-value <= 1×10^{-4} **c** Pearson correlation analysis of Ψ modification fractions at individual sites between biological replicates. **d** Ψ modification fraction alteration pattern observed on specific sites in *K. pneumoniae* different tRNA regions. The left panel depicts the general tRNA secondary structure in *K. pneumoniae*. Structural regions are indicated in the legend and highlighted with corresponding colors. The right scatter plot's y-axis depicts the Ψ fraction difference, calculated as (exponential phase Ψ fraction) - (stationary phase Ψ fraction). **e** Heatmap displaying the tRNA Ψ fraction alteration features detected across different conditions. Color represents the average Ψ fraction values (range from 2% to 100%) at specific sites within tRNA isoacceptors for each strain. The tRNA tags (labeled in y-axis) comprise the amino acids transferred by each tRNA, the corresponding anticodons, and the Ψ position on the tRNA molecules. **f** Venn plot shows the overlap of detected Ψ sites between biological replicates for *P. aeruginosa* and *P. syringae* during each growth phase.

Figure 2

Figure 2. baBID-seq uncovers Ψ modification in bacterial mRNA CDS and UTRs. a

Density plot depicting the distribution of Ψ modifications in mRNA across different growth

phases and conditions. **b** Distribution of mRNA Ψ fraction showing strain and growth phase-

specific patterns of Ψ distribution (right Ψ density plot across each strain's mRNA). **c** mRNA

Ψ faction and counts under different conditions. **d** Right pie charts show the proportion of Ψ

sites in intergenic versus coding regions, and violine plots compare Ψ fraction values

between untranslated regions (upstream and downstream UTRs) and coding regions.

Statistical significance was determined using the Wilcoxon Signed-Rank test; ns, *p*-

value ≥ 0.05; *p-value < 0.05; **p-value < 0.01; ***p-value < 0.001 and ****p-value < 0.0001.

e The pie charts show the Ψ-modified gene overlap in two growth phases across four strains.

f Density plot shows the Ψ site numbers per transcript.

Figure 3

a

b

c

d

Figure 3. Comparative analysis of Ψ modification motif across strains. a Comparison of

overall mRNA 5-mer Ψ motif ratios across four strains. Motif ratios are calculated by

dividing the count of each specific 5-mer motif centered on Ψ by the total number of motifs

detected in each individual strain mRNA. Ψ-modified sites with a fraction above 2% are used

here. **b** Distribution of Ψ fraction (ranging from 2% to 100%) for each motif detected. **c**

Scatter pie chart shows the proportional distribution of top 10 (ranked by motif abundance)

Ψ-containing motif counts categorized by RNA types and bacterial strains. **d** The scatter plot

illustrates the relationship between the average modification fraction and abundance of motifs

in *P. aeruginosa* in exponential (*P. aeruginosa*_exp) and stationary (*P. aeruginosa*_stat)

growth phases. The average Ψ fraction was calculated as the sum of Ψ fractions for each

individual motif divided by its frequency.

Figure 4

Figure 4. Evolutionary conservation of clustered Ψ modifications in orthologous genes.
a The bar plot depicts the Ψ modified homologous genes among bacterial species (green box)
and strain-specific genes (orange box). Functional networks generated by clustering KEGG
pathway enrichment results of Ψ-modified homologous genes present across two or more
strains. The dot size in the network indicates the gene number contained in specific KEGG
pathways. The *p*-value for each pathway was calculated with Fisher's Exact test. **b** Regions

868   with Ψ enrichment across the *atp* and *eno* operons in four strains. Dark red dots represent

869   highly modified sites with Ψ fraction values exceeding 50%.

Figure 5

Figure 5. Growth state-dependent dynamics of Ψ modification fraction. a,b Heatmap

showing the mRNA Ψ fractions alteration of each site in *P. aeruginosa* (a) and *P. syringae*

(b). The color intensity reflects Ψ fraction at each site. Only sites with >10% absolute difference in Ψ fraction between exponential and stationary phases are displayed. Blank boxes signify either unmodified sites or those with Ψ fractions below 2%. The annotation combining position label and gene name indicates the precise location of Ψ modification within genes. **c** Scatter plot shows Ψ intensity alteration between two growth phases in *P. aeruginosa*. For each mRNA, Ψ intensity is calculated as the sum of all Ψ fractions throughout the transcript. **d** Similar to **c**, the scatter plot shows Ψ intensity alteration in *P. syringae* under two growth phases. **e** scatter plot shows Ψ intensity alteration in *P. syringae* under two growth phases in MM medium condition. **f-h** Box plot shows the modified and unmodified mRNA transcripts per kilobase million (TPM) changing between exponential and stationary growth phases under different conditions: *P. aeruginosa* cultured in LB medium (**f**), *P. syringae* cultured in KB medium (**g**) and *P. syringae* cultured in MM medium (**h**). Y-axis shows $\log_2$(TPM at exponential phase / TPM at stationary phase) of each mRNA. The red color presents Ψ-mRNA, and the blue color indicates no-Ψ-mRNA. Wilcoxon Signed-Rank test; ns, p-value ≥ 0.05; *p-value < 0.05; **p-value < 0.01; ***p-value < 0.001 and ****p-value < 0.0001. **i** The scatter plot illustrates the correlation between translation efficiency (TE) alteration ($\log_2$Foldchange of (TE_MM/TE_KB)) and Ψ intensity difference (Ψ intensity of MM-Ψ intensity of KB) in *P. syringae* cultured under MM medium (TE_MM) versus KB conditions (TE_KB). **j** The proportion of Hfq-bound Ψ-mRNA (red color for exponential growth phase mRNA and blue color for stationary phase mRNA) versus those non-Ψ-mRNA (grey color) across two growth conditions in *P. aeruginosa*. **k** The scatter plot shows a correlation between mRNA Ψ intensity and Hfq binding score, where the Hfq binding score is calculated as the sum of each mRNA peak binding strength ($\log_2$FoldChange value of each peak).

Figure 6

Figure 6. Structure-dependent patterns of Ψ modifications and transformer-GNN-based deep learning network for Ψ prediction. a Predicted RNA secondary structures containing the GUΨCG motif with corresponding Ψ fraction values and gene identifiers annotated. MXfold2 is employed to model these structures using 20nt flanking sequences extending from each modification site. **b** Sequence and structure clustering of 41-nucleotide RNA segments centered by Ψ sites with fraction values greater than 0.1 and containing either GCΨCG (black branch color) or GUΨC (red branch color) motifs. The circular visualization features three concentric layers: the inner layer displays the Ψ fraction value, the middle layer indicates RNA type, and the outer layer represents the GC ratio (%) of each 41nt RNA segment. The red dot around the circle marked the position of RNA displayed in (**a**). **c** Architecture of pseU_NN. The model integrates sequence and structural information through two parallel pathways: 1) A sequence analysis branch with one-hot embedding followed by a multi-head transformer module, and 2) A structure analysis branch that processes RNA secondary structure adjacent matrices through a graph convolution module to extract structural features. The features extracted by the two modules are further weighted and merged as input for residual blocks (fully connected layers). **d** Receiver operating characteristic (ROC) curve depicting the performance of the pseudouridine prediction model using different input sequence lengths of 81nt, 61nt and 41nt. **e** Multi-metric assessment showing precision-recall curve (AUC 0.878), F1 curve, and ROC curve (AUC 0.843) of the pseU_NN model on 41nt validation datasets, achieving a peak F1 score of 0.764. **f** Distribution of pseU_NN prediction scores on 41nt validation datasets.